Actomyosin Contractility is a Potent Suppressor of Mesoderm Induction by Human Pluripotent Stem Cells

Loic Fort[1,3], Wenjun Wang[2], and Ian Macara[1,3]

[1] Dept of Cell and Developmental Biology, Vanderbilt University School of Medicine Nashville TN 37240 USA

[2] Dept of Biomedical Engineering, Vanderbilt University Nashville TN 37240 USA

[3] Authors for correspondence

Abstract

The activation of WNT signaling in human pluripotent stem cells drives efficient conversion to lateral mesoderm, which can be further differentiated into cardiomyocytes. Stabilization of the WNT effector β-catenin promotes expression of mesoderm-specifying genes such as Brachyury (encoded by *TBXT*) and Eomesodermin and drives an epithelial-mesenchymal transition (EMT) that occurs at 49-52 hrs post WNT activation. Interestingly, WNT activation also triggers an early contraction of stem cell colonies, with increased myosin light chain (MLC) phosphorylation, resulting in cellular morphological changes. Small molecule inhibitors and genetic approaches were applied to determine if contractility is related to mesoderm differentiation. Contrary to expectation, suppressing contractility accelerated differentiation; Brachyury induction was enhanced, and EMT initiated as early as 24 hrs post WNT activation. Conversely, expression of constitutively active Rho kinase or myosin light chain kinase, to increase contractility, efficiently blocked any induction of mesoderm, and cells fail to undergo EMT. Mechanistically, reduced contractility promoted active β-catenin levels in the cytoplasm and nucleus. We propose that the early contraction of human pluripotent stem cell colonies in response to WNT activation initially delays mesoderm induction by restraining accumulation of active β-catenin, suggesting a role for contractility as a temporal gate.

Introduction

Mechanical forces play major roles at many stages of animal development: Starting as early as embryo compaction at the 8-cell stage[1], forces are crucial through gastrulation and later during tissue morphogenesis. Internal forces are largely generated by acto-myosin contractility, and in epithelia are transmitted across cell boundaries by intercellular junctions, enabling collective organization. Tension is generated by these junctions and by adhesion to the extracellular matrix (ECM). Early embryogenesis is marked by waves of cell fate specification to create the three major lineages, followed by further differentiation, and these events are likely coordinated with changes in physical forces created by the cells, to which they then respond. Mechanistically, however, the connections between cell fate and mechanical force remain unclear.

Human embryonic stem cell (hESC) and induced pluripotent stem cell (hiPSC) colonies can be induced to select the mesoderm lineage and eventually to differentiate into functional cardiomyocytes. Pluripotent stem cells are strongly epithelial and lineage specification relies on an epithelial-to-mesenchymal transition (EMT) that is reminiscent of some aspects of gastrulation. When treated with the morphogen BMP4, hESCs can self-organize into 2D "gastruloids" [2], and if plated on compliant hydrogel surfaces they will form "gastrulation-like" nodes that express Brachyury, a mesoderm marker. Micropatterning and traction stress mapping demonstrated that regions of high tension promote Brachyury expression, through release of β-catenin from the cell junctions and consequent activation of WNT signaling, which is known to be required for mesoderm induction [3].

Mesoderm can also be generated from human pluripotent stem cells by direct WNT pathway activation using the small molecule GSK3β inhibitor Chiron 99021 (hereafter CHIR), which protects β-catenin from degradation. Whether mechanical forces play any role in this pathway to mesoderm induction is unknown. We observed that hiPSCs induced by CHIR to differentiate into mesoderm show colony contraction, increased phospho-MLC (myosin light chain) and increased optical retardance, which is a non-invasive measure of actomyosin contractility. From this correlation between contractility and mesoderm induction, plus prior work on BMP4-driven mesoderm in hESCs, we expected that inhibiting actomyosin activity would block induction. Surprisingly, however, we obtained the opposite result: suppressing MLC phosphorylation using small molecule inhibitors or genetic tools substantially promoted mesoderm induction, while

further increasing actomyosin contractility was sufficient to block differentiation and EMT

completely. Mechanistically, we found that promoting cell relaxation led to an increase in active

β-catenin in the cytoplasm and nucleus.

Results

<u>WNT-mediated mesoderm specification increases actomyosin contractility</u>

To explore how pluripotent stem cells sense, integrate and respond to physical cues, we

first investigated whether differentiating cells experience differential forces along their

conversion from hiPSC to mesoderm. hiPSCs were treated with CHIR to stabilize β-catenin and

thereby activate the WNT pathway, triggering pluripotency exit and commitment to primitive

streak and mesoderm lineages. We noticed that hiPSC colonies showed rapid retraction of their

edge following CHIR addition **(Supp. Fig 1A)**, consistent with observations from other groups [4].

Having previously reported an essential and permissive role for apoptosis during mesoderm

specification [5], we tested whether colony retraction was related to the wave of cell death that

occurs following CHIR addition. However, co-treatment with a pan-Caspase inhibitor (Q-VD-

OPH) did not affect retraction of colony edges **(Supp. Fig 1A)**, ruling out an effect of cell density

and cell death.

Commitment to the mesoderm lineage also resulted in thicker epithelial layers,

containing taller cells **(Supp. Fig 1B-D)**, suggesting profound morphological changes that are

highly dependent on the cellular contractile state. Consistent with the role of nuclei as a

mechanosensor [6,7], differentiating cells exhibit significantly smaller nuclear area after 48h of

CHIR treatment **(Supp. Fig 1E-F)**. To address the variation of cell density, we correlated each of

our nuclear area measurements with the actual cell number **(Supp. Fig 1G)**. As expected,

nuclear area negatively correlates with cell density before differentiation. However, we did not

find any correlation following CHIR treatment, together suggesting that the smaller nuclear size

at 48h of differentiation is unlikely to be related to cell density. While these observations were

indirect, they point towards a differential contractile status early in mesoderm commitment,

switching from low contractile undifferentiated cells to high contractility. To directly test for

higher contractility, differentiating cells were stained for phospho-T18/S19 Myosin Light Chain 2

(ppMLC), a key regulator of contractility. Immunofluorescences reveal a gradual increase of

ppMLC2 intensity over time, with the formation of supracellular fibers extending across several

cells **(Supp. Fig 1H-I)**. Finally, we took advantage of the birefringence property of actin fibers to

measure light retardance using quantitative polarization microscopy. Birefringence of a material

changes proportionally to the applied strain, providing an orthogonal way to probe for

contractile status during mesoderm commitment. Confirming our previous data, retardance

values increased during hiPSC-to-mesoderm conversion **(Supp. Fig 1J)**. Specificity of the

retardance value as a readout for contractility was tested by incubating undifferentiated hiPSCs

with a ROCK inhibitor (Y-27632) to suppress MLC phosphorylation. Together, these data suggest

that hiPSC differentiating along the cardiac mesoderm trajectory experience increasing

contractile forces mediated by MLC2 activity.

<u>Inhibition of actomyosin contractility promotes hiPSc and hESC conversion to the mesoderm</u>

<u>lineage</u>

We hypothesized initially that increased contractility might be required for pluripotent

cell to acquire their mesoderm identity. To test this hypothesis, hiPSCs were differentiated in the

presence of ROCK inhibitor, Y-27632. We first confirmed that MLC2 phosphorylation was

reduced by ROCK inhibition **(Supp. Fig 2A-B)**. Contrary to expectations, however, blocking

contractility strongly promoted Brachyury expression, a mesoderm marker (encoded by the

TBXT gene, also known as *T* gene) **(Fig 1A-B)**. Expression of other key mesoderm (*MESP1* and

TBX6) and EMT (*SNAI1* and *SNAI2*) genes was also upregulated following Y-27632 treatment **(Fig**

1C-F). To rule out an effect for ROCK inhibition on global transcription, we examined expression

levels of genes unrelated to mesodermal fate and did not detect any response to decreased

contractility, suggesting a specific role of actomyosin contractility in cell fate specification **(Supp.**

Fig 2C).

Normal hiPSCs synchronously undergo EMT at 49-52hrs post-CHIR addition. Brachyury

expression is initially restricted to the colony edges at 24hrs, before broader expression across

the colony at later timepoints (**Fig. 1G-H**). However, Y-27632 treatment resulted in markedly

accelerated EMT, as shown by loss of the epithelial marker ZO-1 from tight junctions (**Fig. 1G, I**)

and increased expression of the mesenchymal marker Slug **(Supp. Fig 2D)**, plus more

widespread Brachyury expression **(Fig. 1G-H)**. To validate these data, we used a more specific

ROCK inhibitor, H-1152, which led to similar decreases in ppMLC2 **(Supp. Fig 2E-F)** and

recapitulated the early EMT and higher induction of mesoderm markers **(Supp. Fig 2G-M)**.

Importantly, these effects of ROCK inhibition were conserved in hESCs expressing a knock-in T-

mNeonGreen fusion [3] **(Fig. 1J-K)** and in wild-type H9 hESCs treated with CHIR. Importantly,

there was no effect of reduced contractility on exit from the pluripotency state **(Fig 1L and**

Supp. Fig 2N, P).

Because apoptosis-mediated ATP release is crucial to enable mesoderm fate

specification [5], we probed for PARP cleavage as an apoptosis marker but did not detect any

difference following addition of Y-27632 or H-1152 **(Supp. Fig 2L and 2N-O)**. MLC2

phosphorylation was previously shown to be spatially regulated by ROCK and Myosin Light

Chain Kinase (MLCK) in polarized and non-polarized cells [8,9]. Therefore, we tested the

involvement of MLCK by treating cells with ML-7, a potent MLCK inhibitor [10]. ML-7 led to

efficient reduction in ppMLC2 **(Fig. 1N-O)** and strongly enhanced Brachyury expression at 48h

(Fig. 1P-Q), without affecting cell death. Together these data unexpectedly show that

pharmacological blockade of MLC2-mediated force generation enhances mesoderm

commitment quantitatively and accelerates the EMT required for conversion to cardiac

mesoderm.

<u>Genetic suppression of actomyosin contractility promotes stem cell conversion to the</u>

<u>mesoderm lineage</u>

To rule out off-target effects of the small molecule inhibitors, we designed a genetic

approach to suppress contractility in human cells. Dephosphorylation of ppMLC is mediated by

Myosin Light Chain Phosphatase, an heterotrimer composed of a catalytic subunit PP1cβ, a

Myosin phosphatase-targeting regulatory subunit (MYPT1) and a protein of unknown function

M20 [11,12] **(Fig 2A-B)**. Based on previous work describing an optogenetic construct to control

actomyosin contractility [13], we created a truncated MYPT1, fused with a nuclear export

sequence and a mNG reporter (**Fig 2B**), that is expected to act as a constitutively active mutant

and promote MLC2 dephosphorylation. To validate the activity of this new tool, we cloned this

construct (referred to as MYPT1^{CA}-NES-mNG) under a doxycycline-inducible promoter and

established a stable hiPSC line. Global reduction of ppMLC2 was observed by immunoblot and

immunofluorescence after doxycycline addition (**Fig 2C-E**). Strikingly, activating MYPT1^{CA}

recapitulated the effects of pharmaceutical inhibition of contractility: differentiating MYPT1^{CA}

cells displayed enhanced Brachyury expression (**Fig 2F-J**) and earlier induction of the EMT genes

Snail and Slug (**Fig 2F**) following Doxycycline induction, compared to controls. Apoptosis

induction was not affected in this system (**Fig 2H**). We conclude that reducing MLC2-driven

contractility, using either pharmaceutical or genetic strategies, accelerates mesoderm

specification.

<u>Contractility is a potent inhibitor of mesoderm identity</u>

Next, we sought to test if constitutively increased contractility impacts hiPSC-to-mesoderm

conversion. We stably expressed constitutively active ROCK2 or MLCK (ROCK2^{CA} and MLCK^{CA})

under a Doxycycline inducible promoter with ubiquitous expression of a mVenus reporter.

Green transduced cells were mixed with unlabeled WT hiPSCs, pre-treated or not with

Doxycycline to turn on ROCK2^{CA} or MLCK^{CA} before starting the differentiation protocol for 72h

(**Fig 3A**). We first confirmed that mVenus-positive cells treated with Doxycycline showed signs of

higher contractility. In the co-culture set up with ROCK2^{CA}/WT or MLCK^{CA}/WT cells treated with

Doxycycline, mVenus-positive clusters showed strong cortical F-actin and higher ppMLC2

staining organized as fibers compared to WT mVenus-negative cells, which show more diffused

staining (**Supp. Fig 3A, C**). These data support the idea that mVenus-positive cells experience

higher contractility, allowing us to test whether cell fate was affected. ROCK2^{CA}/WT co-cultures

were treated with water (+ Veh) or Doxycycline (+Dox) and stained for mesoderm (Brachuyry)

(**Fig. 3B**), primitive streak (EOMES) (**Fig 3D**) and EMT markers (Slug and ZO-1) (**Fig 3F**). mVenus-

positive and negative area were drawn based on the mVenus coverage across each cluster

(**Supp. Fig 3B-D**) and percentages of cells expressing each marker were reported. Following

Doxycycline induction, mVenus-positive clusters showed a dramatic decrease in mesoderm,

primitive streak and mesenchymal markers compared to mVenus-negative cells or Vehicle-

treated cells (**Fig. 3C-E-G**). Because we previously showed that MLCK inhibition affects

mesoderm commitment the same way as ROCK inhibition (**Fig. 1P-Q**), we sought to address the

effect of MLCK-driven contractility on cell identity. Importantly, MLCK^{CA}-positive clusters failed

to commit to the primitive streak and mesoderm lineage and to trigger the EMT program

required for mesoderm identity (**Supp. Fig 3E-G**). These data support the surprising conclusion

that constant and pre-mature activation of ROCK/MLCK-mediated contractility prevents

mesoderm conversion and EMT and is sufficient to counteract the effect of CHIR.

<u>Reduced cell contractility promotes nuclear accumulation of stable β-Catenin</u>

Mechanistically, we first hypothesized that Hippo signaling, one of the main mechanosensitive

pathways in epithelial tissues, drives mesoderm specification acceleration upon cell relaxation.

YAP, a downstream effector of the Hippo pathway, has been previously associated with cell fate

patterning in 2D gastruloids [14]. Activation of the Hippo pathway leads to phosphorylation of YAP

at Ser127, restricting its localization to the cytosol and promoting degradation. In contrast, non-

phosphorylated YAP translocates to the nucleus and forms a complex with co-transcriptional

activator TEAD1-4, together regulating gene expression. Surprisingly, however, phosphorylated

YAP was not affected during differentiation or by cell relaxation (**Supp. Fig 4 A-B**). We next

directly probed for *CTGF* and *CYR61* expression, two well-described YAP target genes. Gene

expression strongly decreased during differentiation but was not affected by cellular contractile

status (**Supp. Fig 4C**). Together, these data rule out a significant role for the Hippo pathway in

the acceleration of mesodermal differentiation following cell relaxation.

Next, we investigated the positive feedback loop reported during BMP4-driven

mesoderm commitment, which relies on mechanical tension-dependent secretion of canonical

WNT ligands [3]. We treated our MYPT1[CA]-NES-mNG cells with Doxycycline to trigger cell

relaxation and probed for canonical and non-canonical WNT ligands. As a positive control,

expression of the mesoderm marker *TBX6* was increased in low contractile cells compared to

Vehicle-treated control cells. Interestingly, canonical WNT ligand expression (*WNT3A* and

WNT8A) was strongly upregulated by suppression of contractility, while non-canonical *WNT4*

expression was unaffected neither by CHIR treatment nor by the reduction of contractility

(**Supp. Fig 4E**). To test whether increased WNT3A/8A expression was responsible for increased

mesoderm commitment in low-contractility cells, we blocked WNT ligand processing and

secretion using IWP-2 (Porcupine inhibitor). Addition of IWP-2 did not delay mesoderm gene

expression in Y-27632-treated cells, suggesting that this pathway does not drive accelerated

mesoderm specification (**Supp. Fig 4F**).

We then turned our attention to β-catenin, which exists in epithelial cells both as a

cytosolic molecule and bound to E-Cadherin at adherens junctions and plays key

mechanosensitive functions during development and tissue homeostasis [3,15]. Whether each

pool shares common regulatory mechanisms is still debated [16]. In the absence of WNT signaling,

cytosolic β-catenin is associated with a destruction complex, where it is phosphorylated at

Ser45 by CK1, which primes the protein for subsequent phosphorylation by GSK3β, driving its

ubiquitination and degradation. WNT activation disrupts the destruction complex, permitting

the accumulation of stable, non-phosphorylated β-catenin. Therefore, non-phospho β-catenin

(hereafter "active" β-catenin) signaling can be assessed from the level of non-phosphorylated

Ser45. We evaluated active β-catenin shuttling to the nucleus by isolating cytosolic and nuclear

fractions from MYPT1^{CA} cells (**Fig 4A**). Purity of each fraction was assessed from enrichment of a

nuclear and cytosolic marker, Lamin A/C and GAPDH respectively (**Supp. Fig 4D**). Strikingly,

active β-catenin was strongly upregulated in both the cytosolic and the nuclear fractions in low

contractile differentiating cells (24h + Dox) (**Fig. 4B, C**). Immunostaining confirmed the

enrichment of active β-catenin in the nucleus 24h post differentiation (**Fig. 4D-F**). We validated

this observation by imaging endogenous mEGFP-β-catenin in a knock-in line treated with CHIR

in either control or low contractile (Y-27632) conditions. First, we noticed cytosolic

accumulation of mEGFP-β-catenin within 5h of CHIR treatment, regardless of the presence of Y-

27632. However, consistent with the immunostaining data, mEGFP-β-catenin later became

enriched in the nuclei of cells under low contractile stress (**Supp. Fig 4G-H**). Collectively, these

findings suggest that the promotion of β-catenin nuclear accumulation by cytoskeletal

relaxation accelerates mesoderm program. However, this model can only hold true if the

concentration of CHIR used in our study (7.5 μM) does not saturate the β-catenin signaling

response. To test this premise, we doubled the concentration of CHIR and probed for expression

of WNT target genes as a read-out for WNT signaling (**Fig. 4G**). *LEF1* and *AXIN2* expression

responded in a dose-dependent manner to CHIR. Interestingly, however, at any given CHIR

concentration, inhibition of cell contractility by addition of Y-27632 further enhanced WNT-

target gene responses. Together, these results demonstrate that cell relaxation amplifies active

β-catenin translocation to the nucleus synergistically with CHIR to enhance mesoderm

commitment.

Discussion

Synchronization of developmental processes is key to development. Germ layer specification

relies on tight spatio-temporal control of genetic programs, which is concomitant to profound

morphological changes during gastrulation. Here, we investigated how differentiating cells sense

and integrate physical cues. Our data revealed an unexpected role for acto-myosin contractility

as a potent inhibitor of the mesoderm lineage. We discovered that low contractility promotes

nuclear localization of β-catenin, which correlates with higher and more rapid mesoderm

marker expression (**Fig. 4H**). β-catenin exists as two pools, one bound to the adherens junctions

through E-cadherin and one cytosolic pool that is central to WNT signaling. The regulation and

exchange of β-catenin molecules between these 2 pools remains poorly understood. E-cadherin

binding to β-catenin prevents its nuclear translocation and transcriptional co-activator function

[17]. SRC-mediated phosphorylation of β-catenin at residue Y654 reduces binding to E-cadherin [18],

so one possibility might be that SRC activity is inversely dependent on contractility in

pluripotent stem cells. Alternatively, however, Gayrard et al [19] reported a mechanism through

which, in migrating cells, SRC-dependent phosphorylation of FAK causes actin remodeling and E-

cadherin tension relaxation, resulting in the release and nuclear translocation of β-catenin. In

our system, reduced tension leads to β-catenin accumulation in the cytosolic compartment and

translocation into the nucleus as observed by cell fractionation and confocal imaging (**Fig. 4B, D**

and Supp. Fig 4G).

We note that the response of pluripotent stem cells to changes in contractility and cell-adhesion

tension depends strongly on the environment. For example, contrary to our study, mechanical

tension has been reported to promote Brachyury expression in Bilateria[20] while in gastrulation-

like models of hESC, Muncie et al showed that high cell adhesion tension can promote

mesoderm induction[3]. In this study, authors proposed a dual function for mechanical tension:

First, BMP4-driven Brachyury expression relies on mechanical stretch of cell junction that

triggers a β-catenin-mechanosensitive response by exposing the conserved Y654 residue.

Phosphorylation by SRC-family kinase destabilizes junctional β-catenin and initiates its nuclear

translocation. A positive feedback loop reinforces mesoderm commitment through increased

canonical WNT ligand secretion. These discrepancies may be explained by differences in the

differentiation protocol (BMP4 versus CHIR) and possibly by the substrate/micropattern used [21],

providing differential biochemical and mechanical inputs. Indeed, our protocol directly activates

WNT pathway by inhibition of the destruction complex, in contrast with BMP4 activation.

Our pharmacological and genetic data confirm and expand on a previous study using Y-27632 to

enhance mesendodermal differentiation [22], while ruling out potential off target effects of the

drug. Our study also suggests that cell fate can be modulated by actomyosin contractility. In

Xenopus, decreased ROCK-dependent actomyosin contractility promotes an ectoderm-to-

mesoderm transition, with ectodermal cells adopting mesoderm-like properties [23]. While we

have not tested for alternative cell fate following CHIR addition, our differentiation protocol was

adapted from Lian et al [24,25] and was not reported to lead to ectodermal identity. Additionally,

we note that differentiating cells lose their pluripotency marks, regardless of their contractile

status, suggesting that the rate of pluripotency exit is not affected (Supp. Fig 2N, P).

Alternative mechanisms might also complement this pathway. For instance, a previous study

shows that hiPSC epithelial cohesiveness is a crucial barrier for BMP4-signaling activation, by

blocking BMP4 diffusion to the basolaterally-localized BMP receptors and restricting Brachyury

expression to the colony edge [26]. However, in our system, the use small molecule inhibitor as a

differentiating cue eliminates this diffusion barrier problem. Additionally, while we noticed

increased canonical WNT ligand expression following cell relaxation, we ruled out a feed-

forward loop by blocking WNT ligand secretion (Supp. Fig 4D-E). Finally, we investigated the

activity of the Hippo pathway, as one of the main mechanosensitive pathways. Surprisingly

however, we did not detect signature of differential activity during mesoderm specification.

Unlike other study on several epithelial and non-epithelial cell lines[27], ROCK inhibition did not

affect YAP phosphorylation or expression of YAP target genes. One explanation might be that

pluripotent stem cells are insensitive to Hippo activation because of their inability to perceive

cell-cell interactions as previously reported [28].

In several in vivo and in vitro models, including ours, initiation of mesoderm commitment

correlates with tissue deformation and cell contraction [3,29–32]. In *Drosophila,* contraction of the

posterior end of the embryo is mediated by supercellular active myosin II cables reminiscent

with our observation in 2D [32]. Higher contractility also leads to deformation of nuclear

morphology, as reported by others [30,33]. Therefore, how can we reconciliate this response with

the fact that reducing contractility promotes mesoderm identity? We propose a model in which

increasing contractility following the first 24h of differentiation dampens the CHIR response so

as to temporally synchronize mesoderm identity. Indeed, in the developing embryo,

specification of lineages required deep morphological changes and cell compaction, mostly

mediated by actomyosin contractility [34–36]. It is widely accepted that high density cells, similar to

ingressing cells during gastrulation, have low contractility compared to cells at low density, both

in vitro and in vivo [37]. Our work suggests that this mechanical gate prevents mesoderm

specification until mechanical tension is released. Inducing low contractility prior to CHIR

treatment, disrupts this mechanical gate, resulting to accelerated mesoderm identity and EMT.

Limitations of the study

In this study, we aimed to understand if and how mechanical tension is important for mesoderm

lineage specification. To this end, we used a simple 2D system, allowing us to control most

parameters while only perturbing actomyosin contractility. However, this 2D system is limited in

its physiological parallels. Translation of these data into 2.5D and 3D models of gastrulation (2D

and 3D gastruloid respectively), and alternatively into early mouse development will test the

generalization these findings. Moreover, while we consistently observed higher β-catenin in the

nucleus upon cell relaxation, further work will be required to address whether it reflects an

increase of import or a decrease of export from the nucleus.

Finally, it remains to be addressed whether increased nuclear β-catenin results from increased

binding to its target gene promoters, such as the Brachyury promoter, which contains two

TCF/LEF-1/β-catenin binding sites within the first 500pb [38,39]. Further work on chromatin

rearrangement will be required to better understand whether enhancer/repressor are spatially

modulated by mechanical tension.

Acknowledgements

We thank Cynthia Reinhart-King for giving us access to the QPOL facility. We thank the Weaver

lab (UCSF) and the Gardel lab (University of Chicago) for sharing the T-mNG knock-in hESC line.

We also thank all members of the Macara lab, especially Vaishna Vamadevan and Christian de

Caestecker for valuable suggestions and advice. This work was funded by an American Heart

Association postdoctoral fellowship (Award ID 23POST1018347 to L.F.), a Vanderbilt

Mechanobiology fellowship (to L.F.) and a grant from the National Institutes of Health

(GM070902 to I.G.M.).

Author contributions

L.F. conceived the study, performed and quantified all experiments except the QPOL, and wrote

the manuscript. W.W. imaged and quantified the QPOL data. I.G.M. supervised the project, and

edited the manuscript

Declaration of interest

The authors declare no competing interests.

Material & Methods

Research compliance and ethical regulation

All experiments using human ESCs were performed using the WA09 (H9) cell line under the

supervision of the Vanderbilt Institutional Human Pluripotent Cell Research Oversight

Committee (protocol IRB no. 160146). No cells were sourced directly from human embryos.

Reagents

Common laboratory reagents used in this study are listed in Supplementary Table 1

Cell culture and maintenance

Human GM25256 iPSCs were obtained from the Coriell Institute. The mEGFP–CTNNB1 knock-in

GM25256 iPSC cell line was obtained from the Allen Cell Collection, Coriell Institute (cell ID

AICS-0058-067). The human ESC line H9 (WA09) was obtained from the WiCell Research

Institute. The T-mNG knock-in hESC H9 was made by Valerie Weaver lab (UCSF) and kindly

provided by Margaret Gardel (University of Chicago). The iPSC and ESC H9 lines were cultured

on Matrigel-coated six-well plates (Matrigel diluted at 42 μg ml−1 in DMEM/F12 medium) and

cultured in mTeSR1 medium. The medium was changed daily until the cells reached 70%

confluency. Cells were passaged using Gentle cell dissociation reagent for 4 min, resuspended in

mTseR1 medium as small clusters and replated at 1:7.

HEK293T cells were obtained from the ATCC. Cells were grown in 10% FBS DMEM medium and

passaged at 1:10 every 2-3 days.

All cell lines used in this study were maintained at 37 °C under 5% CO_2.

Cell freezing and thawing

Human iPSCs and ESCs were harvested from culture dishes using Gentle cell dissociation

reagent and centrifuged at 120g for 3 min. The pellets were resuspended in mTeSR1 medium

supplemented with 10% DMSO and aliquoted into cryovials. Cells were first transferred to

−80 °C for 24 h before long-term storage in liquid nitrogen. iPSCs and ESCs were slowly thawed

using mTeSR1, centrifuged and resuspended in mTeSR1 supplemented with 10 µM Y-27632 for

24 h.

Mesoderm differentiation protocol

Differentiation was initiated when cells reach 70-80% confluency. Cells were treated with 7.5

µM CHIR99021 (hereafter CHIR) diluted in RPMI media supplemented with 1X B27 minus insulin

and 100U/mL Penicillin and 100 µg/mL streptomycin (1X) for up to 72 hours.

For pharmaceutical inhibition of contractility, cells were pre-treated overnight with respective

drug or vehicle (water) diluted in mTeSR1. Differentiation was initiated by switching to CHIR

media as described above, supplemented with respective drug or vehicle (water). Similarly, cells

were pre-induced overnight with Doxycycline or vehicle (water) diluted in mTeSR1 before

initiating the differentiation protocol in the presence of Doxycycline or water. For concentration

of each drug, please refer to Supplementary Table 1.

Lentiviral preparation

HEK193T were grown at 40% confluency in 10 cm plate and transfected using Calcium-

Phosphate. Briefly, 20 µg of lentiviral plasmid, 15 µg of pSPAX2 (Addgene #8454) and 6 µg

pMD2G (Addgene #12260) were mixed with 450 µL sterile water and 50 µL of 2.5M $CaCl_2$

solution, previously filter sterilized. DNA mixture was added dropwise into 500 µL of 2X HeBS

(50 mM HEPES, 10 mM KCl, 12 mM dextrose, 280 mM NaCl and 1.5 mM Na2PO4, pH 7.04) with

constant vortexing. Solution was added dropwise on HEK 293T cells for 6h. Medium was

replaced with 8mL of fresh 10% FBS DMEM medium for 48h. Supernatant was filtered through a

0.22 µm membrane and concentrated using a 100kDa cut-off Amicon centrigal unit. Aliquots

were frozen at -80 °C.

SDS-PAGE and Western blotting

Cells were washed with ice cold 1X PBS and lysed with 70 µL RIPA buffer (150 mM NaCl, 10 mM

Tris–HCl pH 7.5, 1 mM EDTA, 1% Triton X-100, 0.1% SDS) supplemented with and 1X protease

and phosphatase inhibitors. Mechanical disruption was performed by scrapping the cells off the

plate, transferring the lysate into a 1.5mL Eppendorff tube and vortexing for 5 seconds followed

by 5 min incubation on ice. Soluble proteins were collected by centrifugation at 16000 g (13200

rpm) for 10min at 4°C. Protein concentration was measured using Precision Red following the

manufacturer's instructions. Lysates in sample buffer were boiled for 5min.

For ppMLC blotting, cells were washed with 1X PBS and directly lysed using 250 µL of boiling 1X

Sample buffer. Lysates were sonicated 3 times for 10 seconds with 60 seconds incubation on ice

between each blast.

30 µg of proteins were resolved on 1.5mm thick bis-tris acrylamide gels and transferred onto

0.2 µm nitrocellulose membrane using BioRad wet transfer system (50 V for 2hours).

Membranes were blocked for > 30min at room temperature (RT) with 5% BSA in 1X TBS-T

(10 mM Tris pH 8.0, 150 mM NaCl and 0.5% Tween 20) and incubated overnight at 4°C and

gentle rocking with primary antibodies diluted in blocking buffer (Supplementary Table 2).

Membranes were washed 3 times with 1X TBS-T and incubated for 1h at RT with AlexaFluor-

conjugated secondary antibodies (Supplementary Table 2). After 3 additional washes in 1X TBS-

T, membranes were scanned using Li-Cor Odyssey CLx system, analyzed and processed using

Image StudioLite v. 5.2.

Immunofluorescence

Cells were cultured on Matrigel-coated Ø 12mm coverglass, fixed with 4% paraformaldehyde for

10min, permeabilized for 5min at RT (20mM Glycine, 0.05% Triton X-100 in 1X PBS), and

incubated with blocking buffer for > 30min (5% BSA in 1X PBS). Coverslips were transferred into

a dark humidified chamber and incubated with primary and AlexaFluor-conjugated secondary

antibodies (Supplementary Table 2) diluted in blocking buffer for 2 hours. Coverslips were

washed in 1X PBS and mounted on glass slide using Fluoromount-G mounting solution.

For ppMLC staining, samples were incubated overnight with primary antibody.

Image acquisition, analysis and processing

Micrographs were obtained using an inverted Nikon A1-R confocal microscope equipped with a

x20 objective (Numerical aperture NA 0.75), a x40 oil objective (NA 1.20) and a x100 oil

objective (NA 1.40). Z-stacks covering the entire cell height were acquired. Maximum Intensity

Projections were obtained and images were analyzed and processed using Fiji (version

2.1.0/1.54f).

MYPT1[CA]-NES-mNeonGreen cloning strategy

gBlock corresponding to truncated human MYPT1 (aa 1-300) was built in fusion with a PKI super

NES and mNeonGreen sequence. gBlock was amplified using Fusion polymerase and the

following primers: Fwd 5'-CTGCTGACCGGTACCATGGCGGACGCGAAGC-3' and Rev 5'-

CATCATACGCGTCTACGATCCGCCACCGC-3'. PCR product and recipient backbone (pInducer10b-

HA-KRas G12V Addgene #164928) were digested with AgeI-HF and MluI-HF for 1h at 37°C, gel

purified and ligated overnight at 16°C using T4 ligase. Ligation reactions were transformed into

chemically competent E.Coli Stbl3 strain by heatshock (45 sec at 42°C) and selected on

Ampicillin plate. Clones were isolated and set up for MiniPrep (QIAprep Spin MiniPrep kit). The

presence of the insert was confirmed by restriction digest check. Positive clones were sent for

full plasmid sequencing provided by Genewiz Plasmid-EZ service before MidiPrep using Takara

Bio NucleoBond XtraMidi kit.

Flow cytometry analysis

T-mNeonGreen knock-in hESC were transduced with a lentivector expressing Histone 2B-

mScarlet (pWPI-H2B-mScarlet). Cells were pre-treated overnight with Vehicle (water) or 10 µM

Y-27632. Differentiation was initiated for 24 hours as described above and cells were harvested

as single cells. Briefly, cells were washed and gentle cell dissociation buffer was added for 8 min

at 37°C. Cell suspension was centrifuged and resuspended in 500 µL PBS. Cells were passed

through a Ø 40 µm strainer directly into 500 µL of 8% PFA (final concentration 4%) and fixed for

10 min on a wheel before sending them for cytometry using a 5-laser Fortessa analyzer (70 µm

nozzle). Singlets were selected based of forward and side scatter. mScarlet (PE-Texas-RedA) and

mNeonGreen (GFP-A) population were selected by drawing a gate on double negative WT hESC

shifted by one order of magnitude. Percentage of mNeonGreen/mScarlet double positive was

reported.

RNA isolation and RT-qPCR

Cells were washed and RNAs were isolated using RNeasy mini kit following manufacturer's instructions. 1 μg of RNA was reverse transcribed to cDNA using the SuperScript III first-strand synthesis kit. cDNA was diluted 1:5 in water and mixed with Maxima SYBR Green/1mM of each primer (Supplementary Table 3). Quantitative PCR was performed on a BioRad CFX96 thermocycler. C_t values from technical triplicates were averaged and normalized to GAPDH using the $\Delta\Delta C_t$ formula.

Cell Fractionation

Following appropriate treatment, cells were washed with ice cold 1X PBS and collected as single cell suspension using gentle cell dissociation as previously described. Cell suspension was washed with cold 1X PBS and resuspended in 500 μL of cold PBS. 100 μL of the cell suspension was set aside and mixed with 30 μL of 4X sample buffer. Sample was sonicated for 15 sec at 20% power 3 times, boiled 5min and stored at -20 °C (Whole Cell Lysate fraction or WCL). Cell fractionation was performed using Cell signaling kit (#9038) and their protocol. Briefly, the remaining 400 μL of cell suspension was centrifuged at 500 x g at 4°C for 5min. Supernatant was removed and pellet was resuspended in 250 μL of CIB buffer, vortexed for 5 sec and incubated on ice for 5 min. Samples were centrifuged at 500 x g at 4°C for 5min and supernatant was stored in pre-chilled tube (= Cytoplasmic fraction). Pellet was resuspended in 250 μL of MIB buffer, vortexed for 15 sec and incubated on ice for 5 min. Samples were centrifuged at 8000 x g at 4°C for 5min and supernatant was discarded (= Membrane and organelle fraction). Pellet was resupended in 125 μL of CyNIB buffer, sonicated for 5 sec at 20% power 3 times (cytoskeletal and nuclear fraction). Each fraction was mixed with the appropriate volume of 4X sample buffer to obtain a final concentration of 1X. Samples were boiled for 5min and centrifuged before loading 15 μL.

Reproducibility and statistics

All datasets were analyzed using GraphPad Prism10 (version 10.0.3). Individual measurements (n) were averaged for each biological repeat (N). Datasets were tested for normality (Shapiro-Wilk test) before applying the appropriate statistical test on the biological repeat (N), except if

mentioned differently in the figure legend. For dataset displayed as superplots, individual

measurements and biological repeats are represented by small and large dots respectively. Each

dataset for a biological repeat is color coded. Error bar represent S.D. Significance levels are

given as follows and exact P values are indicated in each figure: NS, not significant ($P \geq 0.05$);

*$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; and ****$P < 0.0001$.

All experiments, including representative MIPs and western blots, were performed at least three

times independently as biological repeats, unless stated otherwise in the legends. No statistical

method was used to pre-determine the sample size, the experiments were not randomized and

the investigators were not blinded to allocation during experiments and outcome assessment.

Figure legends

<u>Figure 1: Pharmaceutical inhibition of actomyosin contractility promotes hiPSc and hESC</u>

<u>conversion to the mesoderm lineage</u>

A, B – Immunoblot of Brachyury during hiPSC-to-mesoderm commitment (0-48h) in the

presence (+ Y-27632) or absence (+ Veh) of ROCK inhibitor. Molecular weights (M.W.) are

displayed on the right side (A). Brachyury expression was quantified by densitometry and

normalized to α-Tubulin as loading control across N=3 independent biological repeats (B). Mean

and S.D. are displayed. Two-tailed unpaired t test was performed.

C-F – Relative expression of mesoderm markers (*MESP1, TBX6*) (C, D) and EMT markers (*SNAI1,*

SNAI2) (E, F) 48h post CHIR treatment in the presence (+Y-27) or absence (+Veh) of ROCK

inhibitor. N=3 independent biological repeats except for *MESP1* N=4 independent biological

repeats. Mean and S.D. are displayed. Two-tailed unpaired t test was performed.

G-I – Representative MaxIP immunofluorescences during hiPSC-to-mesoderm commitment (0-

48h) in the presence (+ Y-27632) or absence (+ Veh) of ROCK inhibitor. Cells were stained with a

nuclear marker (DNA), EMT marker (ZO-1 – Inverted LUT) and Mesoderm marker (T/Bra–

Inverted LUT). Scale bar = 50 μm (G). Quantification of Brachyury expression (H) and ZO-1

intactness (I) is represented as violin plots with individual measurements (small dots) averaged

for each biological repeat (large dots). Median (Plain red line) and quartiles (Dotted black lines).

n=10-15 technical repeats across N=3 independent biological repeats. One-way ANOVA with

Tukey's multiple comparisons post-test was performed on biological repeats.

J, K – Analysis of Brachyury expression by flow cytometry following 24h CHIR treatment in the

presence (+Y-27632) or absence (+Veh) of ROCK inhibitor using Bra-mNeonGreen knock-in H9

hESC line (J). Quantification of the percentage of cells mNeonGreen-positive from N=4

independent biological repeats. Mean and S.D. are displayed. Two-tailed unpaired t test was

performed (K).

L, M – Immunoblot of Brachyury (mesoderm) and Nanog (Pluripotency marker) using hESC H9 in

the presence (+ Y-27632) or absence (+ Veh) of ROCK inhibitor. Molecular weights (M.W.) are

displayed on the right side (L). Brachyury expression was quantified by densitometry and

normalized to α-Tubulin as loading control across N=3 independent biological repeats (M).

Mean and S.D. are displayed. Two-tailed unpaired t test was performed.

N, O – Representative MaxIP immunofluorescence of hiPSCs treated with a MLCK inhibitor (ML-

7) or Vehicle (DMSO) and stained for nuclei (DNA) and phospho T18/S19 Myosin Light Chain 2

(ppMLC2 – Inverted LUT). Scale bar = 50 μm. Magnified views of the yellow dotted ROI are

shown for ppMLC2. Scale bar = 20 μm (N). Percentage of cellular area positive for ppMLC2

following Vehicle or ML-7 treatment is reported as violin pots. Median (Plain red line) and

quartiles (Dotted black lines). n=15 technical repeats across N=3 independent biological repeats.

Two-tailed unpaired t test was performed on the biological repeats.

P, Q - Immunoblot of Brachyury (mesoderm) and PARP (Cell death marker) using hiPSCs in the

presence (+ ML-7) or absence (+ Veh) of MLCK inhibitor at basal state (0h) and during CHIR

treatment (24h, 48h). Molecular weights (M.W.) are displayed on the right side (P). Brachyury

expression was quantified by densitometry and normalized to α-Tubulin as loading control

across N=3 independent biological repeats. Mean and S.D. are displayed. Mann-Whitney test

was performed for the 24h timepoints and two-tailed unpaired t test was performed for the 48h

timepoints (Q).

<u>Figure 2: Genetic inhibition of actomyosin contractility promotes hiPSc conversion to the</u>

<u>mesoderm lineage</u>

A – Summary of the main regulators of MLC phosphorylation.

B – Schematic representation of full length MYPT1 (1030 aa), interacting with PP1c phosphatase

and protein of unknown function M20. Truncation of MYPT1 Cterm Δ301-1030 leads to a

constitutive recruitment and activation of PP1C. Truncated MYPT1 was fused with NES-

mNeonGreen and is referred to as MYPT1^{CA}-NES-mNG.

C – Immunoblot of phospho T18/S19 MLC2 (ppMLC2) and total MLC2 in WT hiPSCs treated with

Doxycycline or hiPSC expressing MYPT1^{CA}-NES-mNG in the presence or absence of Doxycycline.

α-Tubulin was used as loading control. Molecular weights (M.W.) are displayed on the right side

D, E – Representative MaxIP immunofluorescence of hiPSC expressing MYPT1^{CA}-NES-mNG and

treated or not with Doxycycline. Cells were fixed and stained for phospho T18/S19 MLC2

(ppML2 - Magenta), DNA (blue) and tight junction marker ZO-1 (Orange). Scale bar = 50 μm (D).

Fraction of the cellular area positive for ppMLC2 following Doxycycline induction of MYPT1^{CA}-

NES-mNG hiPSC and reported as violin plots. Median (Plain red line) and quartiles (Dotted black

lines). n=32 technical repeats across N=3 independent biological repeats. Two-tailed unpaired t

test was performed on the biological repeats (E).

F, G – Representative immunoblot for Brachyury (mesoderm marker), Snail and Slug (EMT

markers) using WT and MYPT1^{CA}-NES-mNG hiPSCs, following CHIR treatment (0h-48h) in the

presence or absence of Doxycycline. Molecular weights (M.W.) are displayed on the right side

(F). Brachyury expression was quantified by densitometry and normalized to α-Tubulin as

loading control across N=3-4 independent biological repeats. Mean and S.D. are displayed. A

one-way ANOVA with Tukey's multiple comparisons post-test was performed (G).

H – Immunoblot of PARP (cell death marker) and Brachyury (Mesoderm marker) during

mesoderm commitment (0h-48h CHIR) using MYPT1^{CA}-NES-mNG hiPSCs induced with

Doxycycline. Molecular weights (M.W.) are displayed on the right side.

I, J – Representative MaxIP immunofluorescences of WT and MYPT1^{CA}-NES-mNG hiPSCs treated

with CHIR for 24h +/- Doxycycline. Cells were stained for nuclei (DNA), EMT marker (ZO-1 –

Inverted LUT) and Mesoderm marker (T/Bra – Inverted LUT). Scale bar = 100 μm (I).

Quantification of Brachyury expression is presented as violin plots. Median (Plain red line) and

quartiles (Dotted black lines). n=15-16 technical repeats across N=3 independent biological

repeats. One-way ANOVA with Šidák's multiple comparisons post-test was performed on

biological repeats (J).

Figure 3: Increased actomyosin contractility is sufficient to block mesoderm commitment and

EMT

A – Experimental design for co-culture experiment. hiPSC were transduced with lentivector

expressing ROCK2^{CA} or MLCK^{CA}. Transduced cells are mVenus positive and are mixed with WT

mVenus-negative hiPSCs. ROCK2^{CA}/MLCK^{CA} expression is induced by addition of Doxycycline 24h

prior to initiating mesoderm differentiation. Doxycycline induction was maintained during the 3

days of differentiation using CHIR before fixing and staining.

B-H – Representative MaxIP immunofluorescence for Brachyury (B), EOMES (D), Slug (F) and ZO-

1 (H) in Vehicle (H2O) and Doxycycline-induced ROCK2^{CA} co-culture. mVenus-positive cell

clusters are highlighted by an orange dotted line. Individual channels are inverted LUT. Scale bar

= 50 μm. For (H), magnified mVenus-positive and mVenus-negative areas are shown as insets.

Scale bar = 20 μm. Quantification of Brachyury (C), EOMES (E) and Slug-positive cells (G) is

reported as violin plot, comparing mVenus-positive vs mVenus-negative cluster in the presence

or absence of Doxycycline. For Brachyury, n = 7 (+ Veh) and n=15 (+ Dox) across N=3

independent biological repeats. For EOMES, n=9 across N=3 independent biological repeats. For

Slug, n=9 (+ Veh) and n=10 (+ Dox) across N=3 independent biological repeats. One-way ANOVA

with Tukey's multiple comparisons post-test was performed (C, E, G).

<u>Figure 4: Reduced cell contractility promotes nuclear accumulation of active β-Catenin</u>

A – Experiment design for treatments and cell fractionation.

B-C – Representative immunoblot as explained in (A). MYPT1^{CA}-NES-mNG hiPSCs were induced

with CHIR for 24h in the presence or absence of Doxycycline. Whole cell lysate (WCL) was

collected and loaded as input while cytosolic and nuclear fractions were extracted. Fraction

purity was probed using cytosolic (GAPDH) and nuclear (Lamin A/C) markers. Active β-catenin

(non-phospho S45) enrichment was probed for each fraction and α-Tubulin was used as loading

control. Molecular weights (M.W.) are displayed on the right side (B). Quantification of active β-

catenin enrichment in the cytosolic (left) and nuclear (right) fractions. N=3 independent

biological repeats. Mean and S.D. are displayed. One-way ANOVA with Šidák's multiple

comparisons post-test was performed (C).

D-F – Representative confocal slice and MaxIP for active non-phospho S45 β-catenin (Inverted

LUT) and mNeonGreen respectively. MYPT1^{CA}-NES-mNG hiPSCs were fixed at 0h and 24h post

CHIR treatment complemented or not with Doxycycline. Scale bar = 50 μm. Insets represent a

magnified view of the active non-phospho S45 β-catenin with an example of the nuclei binary

mask overlaid (pink). Scale bar = 10 μm (D). Schematic representation of the quantification

pipeline. (E). Quantification of active non-phospho S45 β-catenin mean intensity as depicted in

(E). n = 10-12 technical repeats across N=3 independent biological repeats. Mean and S.D. are

displayed. One-way ANOVA with Šidák's multiple comparisons post-test was performed (F).

G – Relative expression of WNT target genes (*LEF1* and *AXIN1*) with increasing concentration of

CHIR in the presence (Magenta) or absence (Cyan) of ROCK inhibitor Y-27632. N=7 independent

biological repeats. Mean and S.E.M. are displayed. One-way ANOVA with Šidák's multiple

comparisons post-test was performed.

H – Graphical abstract. Bottom right panel: Upon constitutively high contractile forces,

mesoderm specification is strongly inhibited. Bottom right panel: Cell relaxation promotes

mesoderm conversion by driving accumulation of nuclear β-catenin and licensing of WNT

pathway signaling.

Supplementary Figure 1: hiPSC differentiation along the cardiac mesoderm increases

actomyosin-driven contractility (Related to Figure 1)

A – Bright field imaging of WT hiPSCs treated with CHIR in the presence or absence of a pan-

Caspase inhibitor Q-VD-OPH. Retraction area is highlighted in yellow. Time is presented as

hh:mm. Scale bar = 20 μm.

B, C – Confocal XZ orthogonal views (B) and Maximum Intensity Projections (MaxIP) (C) of

hiPSCs at basal state (0h CHIR) or post-CHIR treatment (24h and 48h CHIR), stained for DNA

(Cyan) and F-Actin (Grey). Scale bar = 10 μm.

D – Quantification of the monolayer height at 0h, 24h and 48h post CHIR treatment from (B).

Violin plots represent individual measurements (small dots) averaged for each biological repeat

(large dots). Median (Plain red line) and quartiles (Dotted black lines). n=32 technical repeats

across N=7 independent biological repeats. A Kruskal-Wallis test with Dunn's multiple

comparison post-test was performed on biological repeats.

E, F – Representative MaxIP immunofluorescence of WT hiPSC during CHIR treatment. Fixed

cells were stained for DNA marker (Hoechst). Scale bar = 20 μm (E). Quantification of average

nuclear size from n=15 fields of view across N=3 independent biological repeats. Median (Plain

red line) and quartiles (Dotted black lines) are displayed. Two-tailed unpaired t test was

performed (F).

G – Correlation of cell area over cell density for each field of view during CHIR treatment.

Biological repeats are color-coded. Pearson correlation coefficient is displayed.

H – Representative MaxIP immunofluorescence of hiPSCs at basal state (0h) or post-CHIR

treatment (24 and 48h) stained for phospho T18/S19 MLC2 (ppMLC2). Pixels are color-coded by

intensity using the fire LUT.

I – Percentage of total imaged area that is ppMLC2-positive at 0h, 24h, 48h post CHIR is

reported as violin plots. Median (Plain red line) and quartiles (Dotted black lines). n=15

technical repeats across N=3 independent biological repeats. A Kruskal-Wallis test with Dunn's

multiple comparison post-test was performed on the technical repeats.

J – Retardance measurements obtained from hiPSCs at basal state (0h), post-CHIR treatment

(24h, 48h, 72h) or treated with Y-27632 at 0h (Y-27) using Quantitative Polarization Microscopy

imaging. Median (Plain red line) and quartiles (Dotted black lines). n=240 technical repeats for

0h, 24h, 48h, 72h and n=120 technical repeats for Y-27 across N=3 independent biological

repeats. A one-way ANOVA with Šidák's multiple comparisons post-test was performed on the

biological repeats.

<u>Supplementary Figure 2: Pharmaceutical inhibition of actomyosin contractility promotes hiPSc</u>

<u>and hESC conversion to the mesoderm lineage (Related to Figure 2)</u>

A, B - Representative MaxIP immunofluorescences of hiPSCs treated with a ROCK inhibitor (Y-

27632) or Vehicle (H_2O) and stained for nuclei marker (DNA) and phospho T18/S19 Myosin Light

Chain 2 (ppMLC2 – Inverted LUT). Scale bar = 50 μm. Magnified views of the yellow dotted ROI

are shown for ppMLC2. Scale bar = 20 μm (A). Fraction of the cellular area positive for ppMLC2

following Y-27632 treatment and reported as violin pots. Median (Plain red line) and quartiles

(Dotted black lines). n=30 technical repeats across N=3 independent biological repeats. Two-

tailed unpaired t test was performed on the biological repeats (B).

C – Relative expression of unrelated genes 48h post CHIR treatment in the presence (+Y-27) or

absence (+Veh) of ROCK inhibitor. N=4 independent biological repeats. Mean and S.D. are

displayed. Two-tailed unpaired t test was performed.

D - Representative MaxIP immunofluorescences of hiPSC treated 48h with CHIR in the presence

(+ Y-27632) or absence (+ Veh) of ROCK inhibitor. Cells were stained for nuclei marker (DNA),

EMT markers (ZO-1 and Slug – Inverted LUT). Scale bar = 50 μm.

E, F - Representative MaxIP immunofluorescences of hiPSCs treated with a ROCK inhibitor (H-

1152) or Vehicle (H_2O) and stained for nuclei marker (DNA) and phospho T18/S19 Myosin Light

Chain 2 (ppMLC2 – Inverted LUT). Scale bar = 50 μm. Magnified views of the yellow dotted ROI

are shown for ppMLC2. Scale bar = 20 μm (E). Fraction of the cellular area positive for ppMLC2

following H-1152 treatment and reported as violin pots. Median (Plain red line) and quartiles

(Dotted black lines). n=15 technical repeats across N=3 independent biological repeats. Two-

tailed unpaired t test was performed on the biological repeats (F).

G-K - Relative expression of mesoderm markers (*TBXT*, *MESP1, TBX6*) (G-I) and EMT markers

(*SNAI1, SNAI2*) (J, K) 48h post CHIR treatment in the presence (+H-1152) or absence (+Veh) of

ROCK inhibitor. N=3 independent biological repeats. Mean and S.D. are displayed. Two-tailed

unpaired t test was performed.

L – Immunoblot probing for mesoderm marker expression (T/Bra) and cell death marker (PARP

cleavage) during hiPSC-to-mesoderm commitment (0h-48h CHIR) in the presence of ROCK

inhibitor (+H-1152). Addition of the pan-caspases inhibitor Q-VD-OPH (+ Q-VD) to the

differentiation medium totally blocks mesoderm commitment and was used as negative control.

α-Tubulin was used as loading control. Molecular weights (M.W.) are displayed on the right side.

M – Representative MaxIP immunofluorescences for mesoderm marker Brachyury (T/Bra –

Inverted LUT) and nuclei. Cells were fixed 24h post-CHIR treatment in the presence or absence

of ROCK inhibitor H-1152.

N-P – Representative immunoblot of PARP (cell death), T/Bra (mesoderm), Oct4 and Nanog

(Pluripotency markers) using hESC H9 in the presence (+ Y-27632) or absence (+ Veh) of ROCK

inhibitor. Molecular weights (M.W.) are displayed on the right side (N). PARP cleavage (O) and

Nanog expression (P) was quantified by densitometry and normalized to α-Tubulin as loading

control across N=4 (PARP) and N=3 (Nanog) independent biological repeats. Mean and S.D. are

displayed. Two-tailed unpaired t test was performed.

Supplementary Figure 3: Increased actomyosin contractility is sufficient to block mesoderm

commitment and EMT (Related to Figure 3)

A – Representative MaxIP immunofluorescence of Doxycycline-induced ROCK2^{CA} hiPSCs

(mVenus-positive) mixed with WT hiPSCs (mVenus-negative). Cells were stained for DNA

(Nuclei), F-actin (top – Inverted LUT) and phospho T18/S19 Myosin Light Chain 2 (Bottom -

ppMLC2 – Inverted LUT). Scale bar = 50 µm. Magnified views comparing mVenus-positive

(Green dotted ROI) and mVenus-negative (Black dotted ROI) areas are shown for F-Actin and

ppMLC2. Scale bar = 10 µm.

B – Percentage of the ROI area positive for mVenus based on cluster considered as mVenus-

Negative and Positive using the ROCK2^{CA} line. On average, mVenus negative clusters have <15%

of their area positive for mVenus. Mean and S.E.M are displayed. n=7 (no Dox) and n=15 fields

of view (+Dox) across N=3 independent biological experiments.

C – Representative MaxIP immunofluorescence of Doxycycline-induced MLCK^{CA} hiPSCs (mVenus-

positive) mixed with WT hiPSCs (mVenus-negative). Cells were stained for DNA (Nuclei), F-actin

(top – Inverted LUT) and phospho T18/S19 Myosin Light Chain 2 (Bottom - ppMLC2 – Inverted

LUT). Scale bar = 50 µm. Magnified views comparing mVenus-positive (Green dotted ROI) and

mVenus-negative (Black dotted ROI) areas are shown for F-Actin and ppMLC2. Scale bar = 10

µm.

D – Percentage of the ROI area positive for mVenus based on cluster considered as mVenus-

Negative and Positive using the MLCK^{CA} line. On average, mVenus negative clusters have <15%

of their area positive for mVenus. Mean and S.E.M are displayed. n=7 (no Dox) and n=18 fields

of view (+Dox) across N=3 independent biological experiments.

E, F - Representative MaxIP immunofluorescence for Brachyury in Vehicle (+Veh) and

Doxycycline-induced MLCK^{CA} co-culture. mVenus-positive cell clusters are highlighted by an

orange dotted line. Individual channels are inverted LUT. Scale bar = 50 µm (E). Percentage of

Brachyury-positive nuclei is reported as violin plot, comparing mVenus-positive vs mVenus-

negative cluster in the presence or absence of Doxycycline. Median (Plain red line) and quartiles

(Dotted black lines). n=7 (+Veh) and n=18 (+ Dox) fields of view (small dots) across N=3

independent biological repeats (large dots). One-way ANOVA with Tukey's multiple comparisons

post-test was performed (F).

G - Representative MaxIP immunofluorescence for EOMES, Slug and ZO-1 in Doxycycline-

induced MLCK^{CA} co-culture. mVenus-positive cell clusters are highlighted by an orange dotted

line. Individual channels are inverted LUT. Scale bar = 50 µm.

<u>Supplementary Figure 4: Reduced cell contractility promotes nuclear accumulation of active β-</u>

<u>catenin (Related to Fig. 4)</u>

A, B – Immunoblot for phospho S127 YAP (pYAP)and total YAP following differentiation (0-48h CHIR) in the presence or absence of ROCK inhibitor Y-27632. Brachyury expression was also probed as an internal positive control and α-Tubulin was used as loading control. Molecular weights (M.W.) are displayed on the right side (A). Ratio of phospho/total YAP was quantified across N=4 independent biological repeats. Mean and S.D. are displayed. One-way ANOVA with Šidák's multiple comparisons post-test was performed.

C – Relative expression of YAP-target genes *CTGF* and *CYR61* during differentiation in the presence or absence of ROCK inhibitor Y-27632. N=4 independent biological repeats. Mean and S.D. are displayed. One-way ANOVA with Šidák's multiple comparisons post-test was performed.

D - Relative expression of canonical WNT (*WNT3A, WNT8A*), non-canonical WNT (*WNT4*) and mesoderm marker (*TBX6*) in MYPT1^{CA}-NES-mNG cells treated with CHIR (0-48h) in the presence of Vehicle or Doxycycline. N=3 independent biological repeats. Mean and S.D. are displayed. One-way ANOVA with Šidák's multiple comparisons post-test was performed.

E – Relative expression of *TBX6* and *MESP1* (mesoderm markers) following CHIR treatment in the presence of Y-27632 complemented or not with 7.5 μM Porcupine inhibitor (IWP-2 or DMSO respectively). N=3 independent biological repeats. Mean and S.D. are displayed. One-way ANOVA with Tukey's multiple comparisons post-test was performed.

F – Compartment purity following fractionation was assessed by quantifying Lamin A/C and GAPDH enrichment in both, cytosolic and nuclear fractions across N=4 independent biological repeats.

G, H – Still pictures from confocal timelapse imaging of mEGFP-β-catenin knock-in hiPSCs treated with CHIR +/- Y-26732. Time is shown in hours. Scale bar = 10 μm (G). mEGFP-β-catenin intensity is reported as a scan line plot across a cell at 0h and 20h (H).

References

1. Firmin, J. *et al*. Mechanics of human embryo compaction. *Nature* **629**, 646–651 (2024).
2. Deglincerti, A. *et al*. Self-organization of human embryonic stem cells on micropatterns. *Nat Protoc* **11**, 2223–2232 (2016).
3. Muncie, J. M. *et al*. Mechanical Tension Promotes Formation of Gastrulation-like Nodes and Patterns Mesoderm Specification in Human Embryonic Stem Cells. *Dev Cell* **55**, 679-694.e11 (2020).

768   4.   Hookway, C. *et al.* A human induced pluripotent stem (hiPS) cell model for the
769   holistic study of epithelial to mesenchymal transitions (EMTs). *BioRxiv* (2024).
770   5.   Fort, L., Gama, V. & Macara, I. G. Stem cell conversion to the cardiac lineage requires
771   nucleotide signalling from apoptosing cells. *Nat Cell Biol* **24**, 434–447 (2022).
772   6.   Lomakin, A. J. *et al.* The nucleus acts as a ruler tailoring cell responses to spatial
773   constraints. *Science (1979)* **370**, (2020).
774   7.   de Leeuw, N. F., Budhathoki, R., Russell, L. J., Loerke, D. & Blankenship, J. T. Nuclei as
775   mechanical bumpers during epithelial remodeling. *Journal of Cell Biology* **223**,
776   (2024).
777   8.   Totsukawa, G. *et al.* Distinct roles of ROCK (Rho-kinase) and MLCK in spatial
778   regulation of MLC phosphorylation for assembly of stress fibers and focal adhesions
779   in 3T3 fibroblasts. *J Cell Biol* **150**, 797–806 (2000).
780   9.   Shi, Y., Ohler, L. S., de Jesus, B. N. & Dickinson, D. J. Opposing actomyosin pools
781   generate cortical flows that establish epithelial polarity. *Biorxiv* (2024).
782   10.   Makishima, M. *et al.* Induction of differentiation of human leukemia cells by
783   inhibitors of myosin light chain kinase. *FEBS Lett* **287**, 175–177 (1991).
784   11.   Matsumura, F., Yamakita, Y. & Yamashiro, S. Myosin Phosphatase-targeting Subunit 1
785   Controls Chromatid Segregation. *Journal of Biological Chemistry* **286**, 10825–10833
786   (2011).
787   12.   Kiss, A., Erdődi, F. & Lontay, B. Myosin phosphatase: Unexpected functions of a long-
788   known enzyme. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* **1866**,
789   2–15 (2019).
790   13.   Yamamoto, K. *et al.* Optogenetic relaxation of actomyosin contractility uncovers
791   mechanistic roles of cortical tension during cytokinesis. *Nat Commun* **12**, 7145
792   (2021).
793   14.   Stronati, E. *et al.* YAP1 regulates the self-organized fate patterning of hESC-derived
794   gastruloids. *Stem Cell Reports* **17**, 211–220 (2022).
795   15.   Röper, J.-C. *et al.* The major β-catenin/E-cadherin junctional binding site is a primary
796   molecular mechano-transductor of differentiation in vivo. *Elife* **7**, (2018).
797   16.   van der Wal, T. & van Amerongen, R. Walking the tight wire between cell adhesion
798   and WNT signalling: a balancing act for β-catenin. *Open Biol* **10**, (2020).
799   17.   Orsulic, S., Huber, O., Aberle, H., Arnold, S. & Kemler, R. E-cadherin binding prevents
800   β-catenin nuclear localization and β-catenin/LEF-1-mediated transactivation. *J Cell
801   Sci* **112**, 1237–1245 (1999).
802   18.   Roura, S., Miravet, S., Piedra, J., de Herreros, A. G. & Duñach, M. Regulation of E-
803   cadherin/Catenin Association by Tyrosine Phosphorylation. *Journal of Biological
804   Chemistry* **274**, 36734–36740 (1999).
805   19.   Gayrard, C., Bernaudin, C., Déjardin, T., Seiler, C. & Borghi, N. Src- and confinement-
806   dependent FAK activation causes E-cadherin relaxation and β-catenin activity.
807   *Journal of Cell Biology* **217**, 1063–1077 (2018).
808   20.   Brunet, T. *et al.* Evolutionary conservation of early mesoderm specification by
809   mechanotransduction in Bilateria. *Nat Commun* **4**, 2821 (2013).
810   21.   Llewellyn, J., Charrier, A., Cuciniello, R., Helfer, E. & Dono, R. Substrate stiffness
811   alters layer architecture and biophysics of human induced pluripotent stem cells to
812   modulate their differentiation potential. *iScience* **27**, 110557 (2024).
813   22.   Maldonado, M., Luu, R. J., Ramos, M. E. P. & Nam, J. ROCK inhibitor primes human
814   induced pluripotent stem cells to selectively differentiate towards mesendodermal

lineage via epithelial-mesenchymal transition-like modulation. *Stem Cell Res* **17**,
222–227 (2016).

23. Kashkooli, L., Rozema, D., Espejo-Ramirez, L., Lasko, P. & Fagotto, F. Ectoderm to
mesoderm transition by down-regulation of actomyosin contractility. *PLoS Biol* **19**,
e3001060 (2021).

24. Lian, X. *et al.* Robust cardiomyocyte differentiation from human pluripotent stem
cells via temporal modulation of canonical Wnt signaling. *Proceedings of the*
National Academy of Sciences **109**, (2012).

25. Lian, X. *et al.* Directed cardiomyocyte differentiation from human pluripotent stem
cells by modulating Wnt/β-catenin signaling under fully defined conditions. *Nat*
Protoc **8**, 162–175 (2013).

26. Vasic, I. *et al.* Loss of TJP1 disrupts gastrulation patterning and increases
differentiation toward the germ cell lineage in human pluripotent stem cells. *Dev Cell*
58, 1477-1488.e5 (2023).

27. Dupont, S. *et al.* Role of YAP/TAZ in mechanotransduction. *Nature* **474**, 179–183
(2011).

28. Pagliari, S. *et al.* YAP–TEAD1 control of cytoskeleton dynamics and intracellular
tension guides human pluripotent stem cell mesoderm specification. *Cell Death*
Differ **28**, 1193–1207 (2021).

29. Leptin, M. & Grunewald, B. Cell shape changes during gastrulation in *Drosophila*.
Development **110**, 73–84 (1990).

30. Skory, R. M. *et al.* The nuclear lamina couples mechanical forces to cell fate in the
preimplantation embryo via actin organization. *Nat Commun* **14**, 3101 (2023).

31. Caldarelli, P. *et al.* Self-organized tissue mechanics underlie embryonic regulation.
Nature **633**, 887–894 (2024).

32. Rozbicki, E. *et al.* Myosin-II-mediated cell shape changes and cell intercalation
contribute to primitive streak formation. *Nat Cell Biol* **17**, 397–408 (2015).

33. Dahl, K. N., Kahn, S. M., Wilson, K. L. & Discher, D. E. The nuclear envelope lamina
network has elasticity and a compressibility limit suggestive of a molecular shock
absorber. *J Cell Sci* **117**, 4779–4786 (2004).

34. Maître, J.-L. *et al.* Adhesion Functions in Cell Sorting by Mechanically Coupling the
Cortices of Adhering Cells. *Science (1979)* **338**, 253–256 (2012).

35. Krieg, M. *et al.* Tensile forces govern germ-layer organization in zebrafish. *Nat Cell*
Biol **10**, 429–436 (2008).

36. Turlier, H. & Maître, J.-L. Mechanics of tissue compaction. *Semin Cell Dev Biol* **47–48**,
110–117 (2015).

37. Loza, A. J. *et al.* Cell density and actomyosin contractility control the organization of
migrating collectives within an epithelium. *Mol Biol Cell* **27**, 3459–3470 (2016).

38. Yamaguchi, T. P., Takada, S., Yoshikawa, Y., Wu, N. & McMahon, A. P. T (Brachyury) is
a direct target of Wnt3a during paraxial mesoderm specification. *Genes Dev* **13**,
3185–3190 (1999).

39. Arnold, S. J. *et al.* Brachyury is a target gene of the Wnt/β-catenin signaling pathway.
Mech Dev **91**, 249–258 (2000).

Figure 1

Figure 2

Figure 3

Figure 4

A MYPT1^CA-NES-mNG

B

	WCL				Cytosolic				Nuclear			
	+Veh		+Dox		+Veh		+Dox		+Veh		+Dox	
CHIR	0h	24h	0h	24h	0h	24h	0h	24h	0h	24h	0h	24h

M.W. (kDa)

- GAPDH — 37
- Lamin A / Lamin C — 75
- T/Bra — 50
- Active β-catenin — 100
- α-Tubulin — 50

C

Cytosolic Active β-catenin — Normalized enrichment (A.U.)
** p=0.0037
*** p=0.0008

Nuclear Active β-catenin — Normalized enrichment (A.U.)
** p=0.0028
** p=0.0038

CHIR 0h 24h | 0h 24h
+ Veh | + Dox

D

Columns: Nuclei | Active β-catenin | mNeonGreen* | Magnification + Nuclear outline

Rows: 0h CHIR (+Veh, +Dox); 24h CHIR (+Veh, +Dox)

Scale bars: 50 µm, 10 µm

Confocal slice | Confocal slice | * MaxIP

E Quantification pipeline

Cell A, Cell B
Hoechst
Active non-phospho S45 β-catenin

Nucleus A, Nucleus B
Create binary mask using Hoechst channel → Apply mask to β-catenin channel and measure nuclear mean intensity

F

Nuclear active β-catenin (Mean Intensity)
* p=0.0107
** p=0.0036
n.s. p=0.9433

CHIR 0h 24h | 0h 24h
+ Veh | + Dox

G

LEF1 — Normalized mRNA expression (ΔΔCt)
n.s. p=0.4906
*** p=0.0006
* p=0.0234
*** p=0.0002
**** p<0.0001
**** p<0.0001

AXIN2 — Normalized mRNA expression (ΔΔCt)
* p=0.0496
*** p=0.0002
* p=0.0331
**** p<0.0001
**** p<0.0001
*** p=0.0002

Veh Y-27 | Veh Y-27 | Veh Y-27
0h | 7.5 µM | 15 µM
24h CHIR

H Working model

Contractility ⇄ Mesoderm identity

Contractility ⊣ Mesoderm identity

Contractility → Nuclear β-catenin → WNT signaling → Mesoderm identity

Supplementary Figure 1

Supplementary Figure 1 (Related to Figure 2)

Supplementary Figure 3 (Related to Figure 3)

Supplementary Figure 4 (Related to Figure 4)