Actomyosin Contractility is a Potent Suppressor of Mesoderm Induction by Human Pluripotent Stem Cells

Loic Fort[1,3], Vaishna Vamadevan[1], Wenjun Wang[2], and Ian G. Macara[1,3]

[1] Dept of Cell and Developmental Biology, Vanderbilt University School of Medicine, Nashville TN 37240 USA

[2] Dept of Biomedical Engineering, Vanderbilt University Nashville TN 37240 USA

[3] Authors for correspondence

Abstract

The activation of WNT signaling in human pluripotent stem cells (hPSCs) drives efficient conversion to lateral mesoderm, which can be further differentiated into cardiomyocytes. Stabilization of the WNT effector β-catenin promotes expression of mesoderm-specifying genes such as *TBXT* (which encodes Brachyury) and drives an epithelial-mesenchymal transition (EMT). Mechanical forces are essential for the self-organization and development of vertebrate embryos but the role of forces, especially actomyosin contractility, in mesoderm specification has remained controversial. We discovered that, unexpectedly, increasing actomyosin contractility by expression of constitutively active Rho kinase or MLC kinase, efficiently blocked any induction of mesoderm by WNT signaling, and cells failed to undergo EMT. Conversely, the suppression of contractility by inhibitors and genetic approaches significantly accelerated differentiation: Brachyury induction was enhanced, and EMT initiated 24hrs earlier than in control settings. These data were initially puzzling because we observed that WNT signaling was sufficient by itself to promote contractility of hiPSC colonies. Notably, however, we showed that contractility must be inhibited prior to WNT activation to efficiently promote mesoderm specification, suggesting that reduced tension primes the pluripotent state, not the induced state, to accelerate differentiation along this trajectory. Mechanistically, we found that loss of contractility decreased junctional β-catenin and promoted active β-catenin levels in the cytoplasm and nucleus. Increased contractility had opposite effects, highlighting actomyosin contractility at the pluripotent state as a key regulator of WNT signaling responsiveness through effects on adherens junctions.

Introduction

Mechanical forces play major roles at many stages of animal development: Starting as early as compaction at the 8-cell stage [1], mechanics plays a central role in self-organization to shape the embryo [2–5]. In vitro, the differentiation of pluripotent stem cells (PSCs) into germ layers is governed by dynamic interplay between mechanical forces, signaling cues, and transcriptional regulators. Internal forces are largely generated by actomyosin contractility, and in epithelia are transmitted across cell boundaries by intercellular junctions, enabling collective organization [6–11]. But how actomyosin contractility is coupled to cell fate specification through target gene expression has remained unclear.

Human embryonic stem cell (hESC) and induced pluripotent stem cell (hiPSC) colonies can be induced to select the mesoderm lineage and eventually to differentiate into functional cardiomyocytes [12–14] Both types of PSCs are strongly epithelial, and lineage specification relies on an epithelial-to-mesenchymal transition (EMT) that is reminiscent of some aspects of gastrulation [15–17].

Mesoderm can also be generated from hPSCs by direct WNT pathway activation using the small molecule GSK3β inhibitor Chiron 99021 (hereafter CHIR), which protects cytosolic β-catenin from degradation through a similar mechanism to WNT ligand signaling. The stabilized β-catenin then enters the nucleus and binds to TCF transcription factors at target gene promoters. A separate pool of β-catenin in epithelial cells is tightly associated with E-cadherin at the adherens junctions, but whether this pool contributes to WNT signaling and mesoderm induction by hPSCs, or participates in WNT responses in other situations, remains controversial [18]. For example, increased junctional β-catenin has been reported to correlate with increased nuclear β-catenin, though the mechanism for this phenomenon was not explained [17]. To the contrary, in *Drosophila* imaginal disks, actomyosin contractility can promote E-cadherin accumulation at adherens junctions[11], which reduces β-catenin-mediated gene expression [11], while deletion of E-cadherin promotes WNT-responsive gene expression [19]. These studies in *Drosophila* suggest that cytoplasmic β-catenin and β-catenin at the adherens junctions are dynamically linked, but whether a similar titration system applies in human cells and, particularly in hPSCs, remains unknown. Although there are multiple studies on the impact of extracellular matrix stiffness [16,17], tissue confinement [20], and EMT [21] on the E-cadherin/β-catenin

interaction and its downstream consequences, there has been to our knowledge no direct

interrogation of how actomyosin contractility impacts mesoderm commitment by hPSCs.

From prior work on BMP4-driven mesoderm in hESCs [16,17], we expected that inhibiting

actomyosin activity would block induction. Surprisingly, however, we obtained the opposite

result: increasing contractility was sufficient to completely block differentiation and EMT, while

suppressing MLC phosphorylation using small molecule inhibitors or genetic tools substantially

promoted mesoderm induction. These data were also puzzling because induction of

differentiation alone promotes contractility of stem cell colonies. However, we discovered that

the effects of elevated or suppressed actomyosin contractility have a temporal dependence,

since their effects on differentiation depend on priming of the pluripotent state, not on events

post-induction. Mechanistically, we found that promoting cell relaxation led to loss of β-catenin

from cell junctions and an increase in active β-catenin in the cytoplasm and nucleus, suggesting

that the β-catenin pools are dynamically coupled through actomyosin contractility to control

mesoderm specification.

Results

<u>Contractility is a potent inhibitor of mesoderm identity</u>

We initially hypothesized that actomyosin contractility might support mesoderm

specification, given prior reports implicating mechanical forces in lineage commitment [5,16,17].

The broad-spectrum phosphatase inhibitor Calyculin A has been used in several studies of the

impact of enhanced contractility on embryonic development [5,9] but it blocks dephosphorylation

of scores of proteins, not just ppMLC. Therefore, we treated WT hiPSCs before and during

differentiation with RhoA activator II (CN03), a highly specific bacterial toxin derivative that

converts the RhoA Glutamine 63 to Glutamate, locking RhoA in a constitutively active form [22,23],

which will then specifically activate ROCK to phosphorylate MLC (**Supplementary Figure 1A**).

Staining for phospho-T18/S19 Myosin Light Chain 2 (ppMLC2) confirmed that CN03 treatment

led to higher levels of contractility by 24 hrs post-CHIR addition (**Supplementary Figure 1B**).

Strikingly, CN03-treated cells failed to express any detectable Brachyury (encoded by *TBXT* gene,

also known as *T* gene), a mesoderm marker. (**Supplementary Figure 1C-D**). However, CN03

treatment caused cell toxicity past 48h, preventing further analysis.

For this reason, we turned to genetic approaches by stably expressing constitutively

active ROCK2 or MLCK (ROCK2^{CA} and MLCK^{CA}) under a Doxycycline inducible promoter, with

constitutive expression of a Venus reporter. We first confirmed that Venus^{Pos} cells treated with

Doxycycline showed signs of higher contractility. In co-cultures containing ROCK2^{CA}/WT or

MLCK^{CA}/WT cells treated with Doxycycline, Venus^{Pos} clusters showed strong cortical F-actin and

higher ppMLC2 staining organized as fibers, compared to WT Venus^{Neg} cells, which show more

diffuse staining (**Supplementary Figure 1E, G**). These data support the idea that Venus^{Pos} cells

experience higher contractility, allowing us to test the impact on cell fate. As an in situ control

unlabeled WT hiPSCs were mixed 1:1 with green (Venus^{Pos}) transduced cells, and the cultures

were pre-treated with Doxycycline (or vehicle) to turn on ROCK2^{CA} or MLCK^{CA} before starting the

differentiation protocol. We prolonged the differentiation to 72 hrs in this co-culture system to

better assess the impact of enhanced contractility (**Figure 1A**). Venus-positive and negative

areas were outlined based on Venus expression across clusters (**Supplementary Figure 1F, H**)

and percentages of cells expressing mesoderm and EMT markers were measured. Strikingly,

following 72h of differentiation in the presence of doxycycline, Venus^{Pos} clusters of ROCK2^{CA}

cells showed a dramatic decrease in mesoderm (**Figure 1B, C**), primitive streak (**Figure 1D, E**)

compared to Venus^{Neg} cells or Vehicle-treated cells. Venus^{Pos} cells also failed to undergo EMT,

which usually occurs 50-52h post differentiation, as observed by the absence of Slug expression

and the persistence of epithelial marker ZO-1 (**Figure 1F-H**).

In addition to ROCK2, MLC2 phosphorylation was previously shown to be spatially

regulated by Myosin Light Chain Kinase (MLCK) in polarized and non-polarized cells [24,25] .

Therefore, we also probed the effect of MLCK-driven contractility on cell identity by expressing a

constitutively active mutant. Consistent with the effects of ROCK2^{CA}, MLCK^{CA}-positive clusters

failed to commit to the primitive streak and mesoderm lineage and to trigger the EMT program

required for mesoderm identity (**Supplementary Figure 1I-K**). These data support the surprising

conclusion that activation of ROCK/MLCK-mediated contractility is sufficient to counteract the

effect of WNT signaling activated by CHIR and completely prevents mesoderm conversion and

EMT.

<u>Genetic suppression of actomyosin contractility promotes stem cell conversion to the</u>

<u>mesoderm lineage</u>

This unexpected finding prompted us to investigate whether inhibition of contractility might

conversely promote mesoderm differentiation. We designed a genetic approach to suppress

contractility in human cells. Dephosphorylation of ppMLC is mediated by MLC Phosphatase, a

heterotrimer composed of a catalytic subunit PP1cβ, a Myosin phosphatase-targeting regulatory

subunit (MYPT1) and a protein of unknown function M20 [26,27] (**Figure 2A-B**). Based on previous

work describing an optogenetic construct to control actomyosin contractility [28], we created a

truncated MYPT1, fused with a nuclear export sequence and a mNeonGreen (mNG) reporter

(**Figure 2B**), which is expected to act as a constitutively active mutant and promote MLC2

dephosphorylation.

To validate the activity of this new tool, we cloned the construct (referred to as

MYPT1^{CA}-NES-mNG) under a doxycycline-inducible promoter and established a stable hiPSC line.

Global reduction of ppMLC2 was observed by immunoblot and immunofluorescence after

doxycycline addition (**Figure 2C-E**). Strikingly, MYPT1^{CA} cells pre-treated overnight with

doxycycline and followed by differentiation (**Figure 2F**), displayed enhanced Brachyury

expression (**Fig 2G-K**) and earlier induction of the EMT genes Snail and Slug (**Figure 2G**)

following Doxycycline induction, compared to controls. Because apoptosis is crucial to enable

mesoderm fate specification [15] we probed for PARP cleavage as an apoptosis marker but did not

detect any difference (**Figure 2I**). We conclude that reducing MLC2-driven contractility

significantly accelerates mesoderm specification.

Promotion of hiPSC and hESC conversion to the mesoderm lineage by actomyosin relaxation is

time-dependent

We next used small molecule inhibitors as an orthogonal method to validate our

previous findings. This approach allowed us to expand our investigation to hESCs, while also

providing better control over the timing of inhibition.

First, hiPSCs were pre-treated overnight with the ROCK inhibitor, Y-27632 to suppress

actomyosin contractility at the pluripotent state then differentiated by addition of CHIR in the

continued presence Y-27632 (**Figure 3A**). We confirmed that MLC2 phosphorylation was

reduced by ROCK inhibition (**Supplementary Figure 2A-B**). Similar to the genetic approach,

pharmaceutical blockage of contractility strongly promoted Brachyury expression (**Figure 3B-C**).

Expression of other key mesoderm (*MESP1* and *TBX6*) and EMT (*SNAI1* and *SNAI2*) genes was

also upregulated following Y-27632 treatment **(Figure 3D-G)**. To rule out an effect of ROCK

inhibition on global transcription, we examined expression levels of genes unrelated to

mesodermal fate and did not detect any response to decreased contractility, suggesting a

specific role for actomyosin contractility in mesoderm specification **(Supplementary Figure 2C)**.

At the cellular level, Brachyury expression is initially restricted to the colony edges at

24hrs, before broader expression across the colony at later timepoints **(Figure 3H-I)**. However, Y-

27632 treatment resulted in more widespread Brachyury expression **(Figure 1H-I)**, markedly

accelerated EMT as shown by loss of the epithelial marker ZO-1 from tight junctions **(Figure 1H,**

J), and increased expression of the mesenchymal marker Slug **(Supplementary Figure 2D)**. To

validate these data, we used a more specific ROCK inhibitor, H-1152, which led to similar

decreases in ppMLC2 **(Supplementary Figure 2E-F)** and recapitulated the early EMT and higher

induction of mesoderm markers **(Supplementary Figure 2G-M)**. Apoptosis was not affected

following Y-27632 or H-1152 treatment **(Supplementary Figure 2L)**. Moreover, these effects of

ROCK inhibition were conserved in hESCs expressing a knock-in T-mNeonGreen fusion [16] **(Figure**

3K-L) and in wild-type H9 hESCs treated with CHIR **(Supplementary Figure 2N-O)**. Importantly,

similar to hiPSCs, there was no effect of reduced contractility on apoptosis initiation

(Supplementary Figure 2N and 2P) or on exit from the pluripotency state **(Supplementary**

Figure 2N and 2Q).

Because MLCK activation was sufficient to block cell commitment **(Supplementary**

Figure 1I-K), we tested the effect of ML-7, a potent MLCK inhibitor [29]. ML-7 led to efficient

reduction in ppMLC2 **(Figure 3M-N)** and strongly enhanced Brachyury expression at 48hrs**,**

without affecting cell death **(Figure 3O-P)**. Together these data show, unexpectedly, that

pharmacological blockade of MLC2-mediated force generation enhances mesoderm

commitment quantitatively and accelerates the EMT required for conversion to cardiac

mesoderm.

These results were particularly surprising, because hiPSC colonies show rapid retraction

of their edges following CHIR addition **(Supplementary Figure 3A and Supplementary Video 1)**,

consistent with observations from other groups [30]. Having previously reported an essential and

permissive role for apoptosis during mesoderm specification [15], we tested whether colony

retraction was related to the wave of cell death that follows CHIR addition. However, co-

treatment with a pan-Caspase inhibitor (Q-VD-OPH) did not prevent retraction of colony edges

(Supplementary Figure 3A), ruling out an effect of cell density and cell death.

Commitment to the mesoderm lineage also resulted in thicker epithelial layers,

containing taller cells **(Supplementary Figure 3B-D)**, suggesting morphological changes that are

highly dependent on the cellular contractile state. Consistent with the role of nuclei as a

mechanosensor [31,32], differentiating cells exhibit significantly smaller nuclear area after 48h of

CHIR treatment **(Supplementary Figure 3E-F)**. To address the variation of cell density, we

correlated each of our nuclear area measurements with the actual cell number **(Supplementary**

Figure 3G). As expected, nuclear area negatively correlates with cell density before

differentiation. However, we did not find any correlation following CHIR treatment, together

suggesting that the smaller nuclear size at 48h of differentiation is unlikely to be related to cell

density.

While these observations were indirect, they point towards a change in contractile

status during mesoderm commitment. To directly test for higher contractility, differentiating

cells were stained ppMLC2. Immunofluorescence revealed a gradual increase of ppMLC2

intensity over time **(Supplementary Figure 3H-I)**. Finally, we took advantage of the birefringence

property of actin fibers to measure optical retardance using quantitative polarization

microscopy [33]. Birefringence of a material changes proportionally to the applied strain,

providing an orthogonal way to probe for contractile status during mesoderm commitment.

Confirming our previous data, retardance values increased during hiPSC-to-mesoderm

conversion **(Supplementary Figure 3J)**. Specificity of the retardance value as a readout for

contractility was tested by incubating undifferentiated hiPSCs with Y-27632, which reduced

retardance. Together, these data suggest that hiPSC differentiating along the cardiac mesoderm

trajectory, in the absence of any external perturbation, experience increasing contractile forces

mediated by MLC2 phosphorylation.

How can we reconcile the surprising observation that differentiating cells intrinsically

become more contractile, with the inhibitory function of actomyosin contractility on

mesodermal specification? We wondered if the timing of inhibitor treatment was important,

and therefore staggered the timing such that Y-27632 was added either 12 hrs prior

differentiation (-12h), at the time of differentiation (0h) or 12 hrs following CHIR treatment

(+12h), and analyzed Brachyury expression after 24hrs post-CHIR addition (**Figure 3Q**). Pre-

treatment of pluripotent stem cells with ROCK inhibitor (-12h) strongly promoted mesoderm

differentiation. Co-treatment with ROCK inhibitor and CHIR reduced the effect, and addition of

ROCK inhibitor at +12hrs had no effect on differentiation (**Figure 3R-S**). We conclude that

reduced contractility primes pluripotent stem cells for specification but has no effect once

specification has been initiated by WNT activation.

<u>Hippo pathway and force-dependent WNT ligand secretion do not contribute to enhanced</u>

<u>mesoderm specification</u>

Mechanistically, how do changes in contractility impact cell fate specification? A major

target of mechanical force in epithelia is the Hippo pathway. YAP, a downstream effector of

Hippo signaling, has been previously associated with cell fate patterning in 2D gastruloids [34], and

substrate stiffness-driven mesoderm specification [35]. Low tension activates this pathway,

resulting in YAP phosphorylation and degradation. Conversely, high tension suppresses Hippo

signaling, blocking phosphorylation of YAP. The non-phosphorylated YAP translocates to the

nucleus to regulate gene expression. Surprisingly, however, phosphorylated YAP levels were not

affected during hiPSC differentiation or by cell relaxation (**Supplementary Figure 4A, B**). We

next directly probed for *CTGF* and *CYR61* expression, two well-described YAP target genes. Gene

expression strongly decreased during differentiation, as expected given the reported repressive

functions of YAP during mesendoderm specification [36], but expression level was not affected by

the cellular contractile status (**Supplementary Figure 4C**). Together, these data rule out a

significant role for the Hippo pathway in the acceleration of mesodermal differentiation

following cell relaxation.

Next, we investigated the positive feedback loop reported for BMP4-driven mesoderm

commitment, which relies on tension-dependent secretion of canonical WNT ligands [16]. We

treated our MYPT1^{CA}-NES-mNG cells with Doxycycline to trigger cell relaxation and probed for

canonical and non-canonical WNT ligands. As expected, expression of *TBX6* (a mesoderm

marker) increased in low contractile cells. Interestingly, canonical WNT ligand expression

(*WNT3A* and *WNT8A*) was strongly upregulated by suppression of contractility, while non-

canonical *WNT4* expression was unaffected by CHIR treatment or by relaxation (**Supplementary**

Figure 4D). To test whether increased *WNT3A/8A* expression was responsible for increased

mesoderm commitment in low-contractility cells, we blocked WNT ligand processing and

secretion using IWP-2, a Porcupine inhibitor. Addition of IWP-2 did not reverse mesoderm gene

expression in Y-27632-treated cells, suggesting that this pathway does not drive accelerated

mesoderm specification in response to cell relaxation (**Supplementary Figure 4E**). Together,

these data ruled out the involvement of two major pathways, previously thought to link cell

tension to mesoderm commitment [16,35] .

<u>Intercellular adhesion and Adherens Junctions mediate cell contractility</u>

Seeking a potential mechanism, we turned our attention to intercellular adherens

junctions (AJs). These junctions are under tension, generated by actomyosin interactions with

vinculin/α-catenin, which in turn are coupled to β–catenin and E-cadherin [37]. In addition,

tension across AJs is mediated by homotypic and calcium-dependent interactions between E-

cadherin molecules on adjacent cells [38]. As a major signaling hub, we reasoned that AJs might

transduce mechanical forces to impact cell fate specification. We first tested whether disrupting

AJs using EGTA (a calcium chelator) would affect cell specification. EGTA treatment caused

colony decompaction (**Figure 4A and Supplementary Fig 5A**), consistent with previous literature

linking AJ-disengagement with reduced force transmission [39,40]. Strikingly, EGTA treatment

triggered a strong enhancement of mesodermal and EMT marker expression (**Figure 4B-E and**

Supplementary Figure 5B-C). Knowing that reduced contractility mimics this phenotype, we

tested whether EGTA treatment would alter contractility. Staining for ppMLC2 confirmed that

EGTA treated cells are in a low contractile state (**Supplementary Figure 5D-E**).

As an orthogonal approach, we sought to uncouple force transmission from the

actomyosin network to E-cadherin via vinculin/α-catenin. To this end, we created α-catenin

knockdown (*CTNNA1* KD) hiPSCs, with a 50% decrease in protein expression (**Supplementary**

Figure 5F-G). First, because AJs are required for stemness acquisition during fibroblast

reprograming to iPSCs [41], we checked that the *CTNNA1* knockdown cells had not lost

pluripotency (**Supplementary Figure 5F, H-I**). Notably, ppMLC2 levels were 50% lower in the KD

cells (**Figure 4F-G**), confirming that α-catenin is a major contributor to cell actomyosin activity.

The *CTNNA1* KD hiPSCs exhibited higher and earlier expression of Brachyury. Strikingly, however,

these cells failed to further respond to Y-27632 treatment, while control cells responded as

expected by increasing Brachyury expression (**Figure 4H-I**). Together these data demonstrate

that AJ disruption reduces actomyosin contractility and that α-catenin-mediated force

transmission is a major determinant of mesoderm specification kinetics.

<u>Junctional β-catenin localization scales with intrinsic actomyosin contractility in undifferentiated</u>

<u>hiPSCs</u>

Our discovery of a key role for AJ mechanics in cell fate specification suggested that β-

catenin, which in epithelial cells is mostly bound to E-Cadherin at AJs, might respond to changes

in actomyosin contractility by modulating WNT signaling responsiveness. Release of β-catenin

from AJs, for example, might increase the cytoplasmic and nuclear pools to amplify expression

of *TBXT* and other mesoderm genes. Mechanosensitive functions for β-catenin during

development and tissue homeostasis have been described [16,42] but whether the junction-

associated and cytoplasmic β-catenin pools both contribute to WNT signaling is still

controversial [18], and other studies have suggested a connection in which increased junctional β-

catenin correlates with increased nuclear β-catenin [17] .

We hypothesized that loss of tension at AJs would release β-catenin to promote WNT-

responsive gene expression, and conversely that increased tension would promote binding at

AJs, thereby titrating out the free cytoplasmic/nuclear β-catenin. To test this mechanism,

undifferentiated hiPSCs were treated with either the Rho activator CN03 or ROCK inhibitor Y-

27632, causing an increase or decrease in ppMLC2, respectively (**Figure 5A-B and**

Supplementary Figure 6A). Strikingly, under each condition, junctional β-catenin levels

correlated closely with the level of active ppMLC2 in the cells (**Figure 5B-C and Supplementary**

Figure 6B). To rule out staining artifacts, we performed comparable experiments using our

EGFP-β-catenin knock-in hiPSC line (**Figure 5D**), with a similar conclusion (**Figure 5E-F,**

Supplementary Figure 6C and Supplementary Video 2). Importantly, overall β-catenin

expression is not affected by contractile state, suggesting that contractility does not regulate β-

catenin level but rather its localization (**Supplementary Figure 6D-F**). Together, these data

demonstrate that β-catenin association with AJs scales directly with actomyosin contractility.

<u>Cell relaxation promotes nuclear β-catenin localization and occupancy at *TBXT* promoter</u>

In the absence of WNT signaling, cytosolic β-catenin is maintained at a low level by

association with a destruction complex, where it is phosphorylated at Ser45 by CK1, which

primes the protein for subsequent phosphorylation by GSK3β, driving its ubiquitination and

degradation. WNT activation disrupts the destruction complex, permitting the accumulation of

stable, non-phosphorylated β-catenin, which can enter the nucleus to drive gene expression.

This "active" state can be assessed from the detection of non-phosphorylated Ser45 β-catenin.

To test for differential accumulation of active β-catenin, we isolated cytosolic and nuclear

fractions from cells that contain the Dox-inducible MYPT1^{CA}-NES-mNG construct

(**Supplementary Figure 7A**). Purity of each fraction was assessed for enrichment of a nuclear

and cytosolic marker, Lamin A/C and GAPDH, respectively (**Supplementary Figure 7B**).

Fractionation showed that active β-catenin was strongly enriched in both the cytosolic and the

nuclear fractions of cells treated with CHIR + Dox (24h) versus cells treated only with CHIR

(**Figure 6A-B**). Immunostaining confirmed the nuclear enrichment of active β-catenin (**Figure 6C**

and Supplementary Figure 7C-D) after 24h of differentiation in low contractile conditions.

We validated this observation by imaging endogenous mEGFP-β-catenin knock-in hiPSCs

treated with CHIR in either control or low contractile (+Y-27632) conditions. First, we noticed

nuclear accumulation of mEGFP-β-catenin within 5h of CHIR treatment, regardless of the

contractile status of the cells. However, while β-catenin signal decreased over time in vehicle-

treated cells, remaining visible after 20h in only a few nuclei, low contractile cells showed a

robust and enhanced nuclear localization (**Figure 6D-E and Supplementary Video 3**).

Finally, knowing that β-catenin is enriched in the nucleus upon low contractility, we

focused on the ability of β-catenin to interact with its binding partners and to regulate WNT-

responsive genes. Because of the timing of our phenotype, we ruled out an involvement of β-

catenin interactions with transcription factors (such as SOX family, SMAD, TBX3) [43,44], as they are

involved later during specification [45]. In addition, ChIP-seq experiments report that TCFs/β-

catenin binding is the main event during pluripotency exit and early specification [45,46].

Mammalian genomes encode four TCF proteins (TCF7, TCF7L1, TCF7L2 and LEF1), all of which

bind to similar DNA sequences [47], show redundant functions [46,48] and co-occupy similar DNA

regions [49,50]. TCF3 (encoded by *TCF7L1*) is the most highly expressed member in hiPSC

(**Supplementary Figure 7E**) but is widely reported to be repressive [51–54]. Therefore, we focused

on the second most highly expressed factor, TCF4 (encoded by *TCF4L2*). Nuclei isolated from Y-

27632-treated EGFP- β-catenin knock-in hiPSCs showed higher binding between β-catenin and

TCF4 compared to vehicle-treated cells (**Figure 6F-H**). Finally, CUT & RUN assays showed that

low contractility significantly increased β-catenin recruitment to *TBXT* promoter during

differentiation (**Figure 6I and Supplementary Figure 7F**), supporting a priming model. Together,

these findings suggest that the promotion of nuclear β-catenin accumulation by cytoskeletal

relaxation accelerates mesoderm program by binding to its TCF co-mediator. However, this

model can only hold true if the concentration of CHIR used in our study (7.5 µM) does not

saturate the β-catenin signaling response. To test this premise, we doubled the concentration of

CHIR and probed for expression of WNT target genes as a read-out for WNT signaling

(**Supplementary Figure 7G**). *LEF1* and *AXIN2* expression responded in a dose-dependent

manner to CHIR. Interestingly, however, at any given CHIR concentration, inhibition of cell

contractility further enhanced WNT-target gene responses.

Collectively, we demonstrate that actomyosin contractility regulates β-catenin

availability and that cell relaxation amplifies active β-catenin translocation to the nucleus

synergistically with CHIR to enhance mesoderm commitment (**Figure 7**).

Discussion

Here, we have identified an inverse relationship between actomyosin contractility and

mesoderm specification in hPSCs. Mechanistically, our data support a model in which adherens

junctions serve as dynamic reservoirs that sequester β-catenin away from the cytoplasmic

signaling pool. We propose that the inter-dependence of actomyosin contractility and AJs

creates a titration mechanism for β-catenin availability, thereby tuning mesoderm specification.

Previous papers have reported that, contrary to our observations, increased mechanical tension

promotes β-catenin-dependent mesoderm induction. For example, tension has been reported

to promote Brachyury expression in hESC [17], in Bilateria [55] and Cnidaria [20] , while in gastrulation-

like models of hESC, Muncie et al argued that high cell adhesion tension on compliant substrate

can promote mesoderm induction [16]. This study proposed a dual function for mechanical

tension. First, BMP4-driven Brachyury expression relies on mechanical stretch of cell junctions,

allowing destabilization of β-catenin from AJs and its nuclear translocation. Following this

initiation stage, a positive feedback loop reinforces mesoderm commitment through increased

WNT ligand secretion. Strikingly, on Matrigel-coated plates, we observe the opposite outcome.

Importantly, our unconstrained system directly focuses on contractility rather than tension

geometry or substrate properties. Our findings suggest that intrinsic contractility, even in

unpatterned settings, is sufficient to guide cell fate decisions. We provide causal and cell-intrinsic evidence that actomyosin-generated forces act upstream of mesodermal fate, in the pluripotent state, to inhibit cell specification. Second, our protocol directly activates WNT signaling by inhibition of the destruction complex, as do WNT ligands, in contrast with the indirect effects of BMP4 activation. Importantly, a previous study showed that epithelial cohesiveness and tight junctions block access by BMP4 to its receptor on the lateral plasma membrane [56]. Therefore, BMP4 signaling can only activate Brachyury expression at colony edges and corners [16]. However, our protocol uses a cell-permeable small molecule inhibitor as the differentiating cue, circumventing this diffusion barrier. Third, our findings distinguish our model from alternative force-mediated mechanisms such as Hippo/YAP signaling [35] or tension driven WNT ligand secretion [16], neither of which accounted for the enhanced mesoderm specification. One explanation might be that pluripotent stem cells are insensitive to Hippo activation because of their inability to perceive cell-cell interactions, as previously reported [35]. Finally, our system relies on activating or reducing contractility at the pluripotent state by pre-treating cells prior to differentiation and maintaining treatment during cell commitment. We observed a timing-dependent response during which enhanced mesoderm identity was only achieved following ROCK inhibition pre-treatment of cells in the pluripotent state. Therefore, our study suggests that low contractility primes pluripotent cells to enter the differentiation trajectory. Such a licensing mechanism would likely involve cis-regulatory element-mediated chromatin priming that can potentiate gene activity prior to specification events [57]. Alternatively, we note that actomyosin-dependent epigenome regulation was previously associated with RNA-polymerase II activity and lineage commitment in adult stem cells [58].

We here describe a titration mechanism by actomyosin contractility that allows a synchronization with developmental processes during cell commitment. β-catenin exists in two pools, one bound to the adherens junctions through E-cadherin and one cytosolic pool that is central to WNT signaling. The regulation and exchange of β-catenin molecules between these two pools remains poorly understood. E-cadherin binding to β-catenin prevents its nuclear translocation and transcriptional co-activator function [19]. Src-mediated phosphorylation of β-catenin at residue Y654 has been reported to reduce binding to E-cadherin [59], so one possibility might be that Src tyrosine kinase activity is anti-dependent on contractility in pluripotent stem cells. Alternatively, however, Gayrard et al [21] reported an inverse mechanism through which, in

migrating cells, Src-dependent phosphorylation of FAK causes actin remodeling and E-cadherin tension relaxation, resulting in the release and nuclear translocation of β-catenin. In our system, reduced tension causes loss of β-catenin from AJs, accumulation in the cytosolic compartment, and translocation into the nucleus. Finally, in *Xenopus*, decreased ROCK-dependent actomyosin contractility promotes an ectoderm-to-mesoderm transition, suggesting force-dependent lineage plasticity is a conserved mechanism [60]. While we have not tested for alternative cell fate following CHIR addition, our differentiation protocol was adapted from Lian et al [12,13] and was not reported to lead to ectodermal identity.

Acknowledgements

We thank Cynthia Reinhart-King for giving us access to the QPOL facility. We thank the Weaver lab (UCSF) and the Gardel lab (University of Chicago) for sharing the T-mNG knock-in hESC line. We thank Dr. Vivian Gama lab (Vanderbilt University) for providing the NSC lysate, Pax6 and Otx1/2 antibodies. We also thank all members of the Macara lab, especially Dr. Christian de Caestecker for valuable suggestions and advice. This work was funded by an American Heart Association postdoctoral fellowship (Award ID 23POST1018347 to L.F.), a 2024 Vanderbilt Mechanobiology fellowship (to L.F.), a 2024 and 2025 Vanderbilt Mechanobiology fellowship (to L.F.) and a grant from the National Institutes of Health (GM070902 to I.G.M.).

Author contributions

L.F. conceived the study, performed and quantified all experiments except the ones stated below, and wrote the manuscript. V.V. performed the CUT & RUN assay. W.W. imaged and quantified the QPOL data. I.G.M. supervised the project, and edited the manuscript.

Data availability statement

The data are available from the corresponding author upon reasonable request.

Declaration of interest

The authors declare no competing interests.

Material & Methods

Research compliance and ethical regulation

All experiments using human ESCs were performed using the WA09 (H9) cell line under the supervision of the Vanderbilt Institutional Human Pluripotent Cell Research Oversight Committee (protocol IRB no. 160146). No cells were sourced directly from human embryos.

Reagents

Common laboratory reagents used in this study are listed in Supplementary Table 1

Cell culture and maintenance

Human GM25256 iPSCs were obtained from the Coriell Institute. The mEGFP–CTNNB1 knock-in GM25256 iPSC cell line was obtained from the Allen Cell Collection, Coriell Institute (cell ID AICS-0058-067). The human ESC line H9 (WA09) was obtained from the WiCell Research Institute. The T-mNG knock-in hESC H9 was made by Valerie Weaver lab (UCSF) and kindly provided by Margaret Gardel (University of Chicago). The iPSC and ESC H9 lines were cultured on Matrigel-coated six-well plates (Matrigel diluted at 42 µg ml−1 in DMEM/F12 medium) and cultured in mTeSR1 medium. The medium was changed daily until the cells reached 70% confluency. Cells were passaged using Gentle cell dissociation reagent for 4 min, resuspended in mTSeR1 medium as small clusters and replated at 1:7. HEK293T cells were obtained from the ATCC. Cells were grown in 10% FBS DMEM medium and passaged at 1:10 every 2-3 days. All cell lines used in this study were maintained at 37 °C under 5% CO2.

Cell freezing and thawing

Human iPSCs and ESCs were harvested from culture dishes using Gentle cell dissociation reagent and centrifuged at 120g for 3 min. The pellets were resuspended in mTeSR1 medium supplemented with 10% DMSO and aliquoted into cryovials. Cells were first transferred to −80 °C for 24 h before long-term storage in liquid nitrogen. iPSCs and ESCs were slowly thawed using mTeSR1, centrifuged and resuspended in mTeSR1 supplemented with 10 µM Y-27632 for 24 h.

Mesoderm differentiation protocol and drug treatments

Differentiation was initiated when cells reach 70-80% confluency. Cells were treated with 7.5

µM CHIR99021 (hereafter CHIR) diluted in RPMI media supplemented with 1X B27 minus insulin

and 100U/mL Penicillin and 100 µg/mL streptomycin (1X).

For pharmaceutical inhibition of contractility, cells (50% confluent) were pre-treated overnight

or 3h with drug or vehicle diluted in mTeSR1. Differentiation was initiated the day after (cells

reached 70-80% confluent) by switching to CHIR media as described above, supplemented with

the drug or vehicle. For MYPT1^{CA}, cells were pre-induced overnight with Doxycycline or vehicle

(water) diluted in mTeSR1 before initiating the differentiation protocol in the presence of

Doxycycline or water.

For pharmaceutical stimulation of cell contractility, 70-80% confluent cells were pre-

treated for 3h with 4 µg/mL CN03 or water in mTeSR1, followed by CHIR treatment

supplemented with CN03 or water. For EGTA treatment, 70-80% confluent cells were pre-

treated for 30min with 1mM of EGTA or water in mTeSR1, followed by CHIR treatment

supplemented with 10 µM EGTA or water. For concentration of each drug, please refer to

Supplementary Table 1.

Lentiviral preparation

HEK293T were grown at 40% confluency in 10 cm plate and transfected using Calcium-

Phosphate. Briefly, 20 µg of lentiviral plasmid, 15 µg of pSPAX2 (Addgene #8454) and 6 µg

pMD2G (Addgene #12260) were mixed with 450 µL sterile water and 50 µL of 2.5M CaCl$_2$

solution, previously filter sterilized. DNA mixture was added dropwise into 500 µL of 2X HeBS

(50 mM HEPES, 10 mM KCl, 12 mM dextrose, 280 mM NaCl and 1.5 mM Na2PO4, pH 7.04) with

constant vortexing. Solution was added dropwise on HEK 293T cells for 6h. Medium was

replaced with 8mL of fresh 10% FBS DMEM medium for 48h. Supernatant was filtered through a

0.22 µm membrane and concentrated using a 100kDa cut-off Amicon centrigal unit. Aliquots

were frozen at -80 °C.

For generation of CRISPR cells, single-guide RNA was selected using Benchling design

tool, and are listed in Supplementary Table 3. Annealed oligonucleotides were cloned into

pLentiCrispRv2-Puro as described by Sanjana and colleagues

(https://doi.org/10.1038/nmeth.3047). Lentiviral particles were prepared as described above.

hiPSCs were transduced in suspension and selected 48h post-transduction using 1 μg/mL of

Puromycin.

<u>SDS-PAGE and Western blotting</u>

Cells were washed with ice cold 1X PBS and lysed with 70 μL RIPA buffer (150 mM NaCl, 10 mM

Tris–HCl pH 7.5, 1 mM EDTA, 1% Triton X-100, 0.1% SDS) supplemented with and 1X protease

and phosphatase inhibitors. Mechanical disruption was performed by scrapping the cells off the

plate, transferring the lysate into a 1.5mL Eppendorff tube and vortexing for 5 seconds followed

by 5 min incubation on ice. Soluble proteins were collected by centrifugation at 16000 g (13200

rpm) for 10min at 4°C. Protein concentration was measured using Precision Red following the

manufacturer's instructions. Lysates in sample buffer were boiled for 5min.

For ppMLC blotting, cells were washed with 1X PBS and directly lysed using 250 μL of

boiling 1X sample buffer. Lysates were sonicated 3 times for 10 seconds with 60 seconds

incubation on ice between each blast. 30 μg of proteins were resolved on 1.5mm thick bis-tris

acrylamide gels and transferred onto 0.2 μm nitrocellulose membrane using BioRad wet transfer

system (50 V for 2hours). Membranes were blocked for > 30min at room temperature (RT) with

5% BSA in 1X TBS-T (10 mM Tris pH 8.0, 150 mM NaCl and 0.5% Tween 20) and incubated

overnight at 4°C and gentle rocking with primary antibodies diluted in blocking buffer

(Supplementary Table 2). Membranes were washed 3 times with 1X TBS-T and incubated for 1h

at RT with AlexaFluor-conjugated secondary antibodies (Supplementary Table 2). After 3

additional washes in 1X TBS-T, membranes were scanned using Li-Cor Odyssey CLx system,

analyzed and processed using Image StudioLite v. 5.2.

<u>Immunofluorescence</u>

Cells were cultured on Matrigel-coated Ø 12mm coverglass, fixed with 4% paraformaldehyde for

10min, permeabilized for 5min at RT (20mM Glycine, 0.05% Triton X-100 in 1X PBS), and

incubated with blocking buffer for > 30min (5% BSA in 1X PBS). Coverslips were transferred into

a dark humidified chamber and incubated with primary and AlexaFluor-conjugated secondary

antibodies (Supplementary Table 2) diluted in blocking buffer for 2 hours. Coverslips were

washed in 1X PBS and mounted on glass slide using Fluoromount-G mounting solution.

For ppMLC staining, samples were incubated overnight with primary antibody.

<u>Image acquisition, analysis and processing</u>

Micrographs were obtained using an inverted Nikon A1-R confocal microscope equipped with a

x20 objective (Numerical aperture NA 0.75), a x40 oil objective (NA 1.20) and a x100 oil

objective (NA 1.40). Z-stacks covering the entire cell height were acquired. Maximum Intensity

(MaxIP) Projections were obtained and images were analyzed and processed using Fiji (version

2.1.0/1.54f). Images processing did not include a denoising step except for Supplementary

Video 3.

<u>MYPT1^{CA}-NES-mNeonGreen cloning strategy and cell line generation</u>

A gBlock was obtained from IDT, corresponding to truncated human MYPT1 (aa 1-300) in fusion

with a PKI super Nuclear Export Signal and mNeonGreen sequence. The gBlock was amplified

using Fusion polymerase and the following primers: Fwd 5'-

CTGCTGACCGGTACCATGGCGGACGCGAAGC-3' and Rev 5'-CATCATACGCGTCTACGATCCGCCACCGC-

3'. PCR product and recipient backbone (pInducer10b-HA-KRas G12V Addgene #164928) were

digested with AgeI-HF and MluI-HF for 1h at 37°C, gel purified, and ligated overnight at 16°C

using T4 ligase. Ligation reactions were transformed into chemically competent E.Coli Stbl3

strain and selected on Ampicillin plates. Plasmids were isolated from single colonies (QIAprep

Spin MiniPrep kit). The presence of the insert was confirmed by restriction digestion. Positive

clones were sent for full plasmid sequencing provided by Genewiz Plasmid-EZ service before

plasmid purification using Takara Bio NucleoBond XtraMidi kit. WT hiPSCs were transduced as

described above and cells were selected 48h later with 1 µg/mL of Puromycin.

<u>Quantitative Polariztion Microscopy (QPOL)</u>

The contractility of cells was measured using quantitative polarization microscopy (QPOL) as

described previously [33]. Briefly, QPOL was built on an inverted Axiovert microscope equipped

with an Axiocam 506 color camera. A motorized linear polarizer (Thorlabs, Newton, NJ) and a

circular polarizer were positioned in the illumination plane above the condenser and in the

imaging plane, respectively. Images were captured using a 20x/0.5 N.A. polarization objective.

For each field of view, image sequences were acquired with 10° intervals of the rotating

polarizer over the range of 0–180° using Zen software. The polarized image sequences were

then processed using a custom MATLAB code to generate pixel-by-pixel retardance maps. The

obtained retardance images then underwent background subtraction and subsequent analysis

with ImageJ. The retardance signal proportional to cell contractility was quantified as the

average value of the background-subtracted retardance over the cell area.

Flow cytometry analysis

T-mNeonGreen knock-in hESC were transduced with a lentivector expressing Histone 2B-

mScarlet (pWPI-H2B-mScarlet). Cells were pre-treated overnight with Vehicle (water) or 10 µM

Y-27632. Differentiation was initiated for 24 hrs as described above and cells were harvested as

single cells. Briefly, cells were washed and gentle cell dissociation buffer was added for 8 min at

37°C. Cell suspension was centrifuged and resuspended in 500 µL PBS. Cells were passed

through a Ø 40 µm strainer directly into 500 µL of 8% PFA (final concentration 4%) and fixed for

10 min on a wheel before sending them for cytometry using a 5-laser Fortessa analyzer (70 µm

nozzle). Singlets were selected based of forward and side scatter. mScarlet (PE-Texas-RedA) and

mNeonGreen (GFP-A) population were selected by drawing a gate on double negative WT hESC

shifted by one order of magnitude. Percentage of mNeonGreen/mScarlet double positive was

reported using the online flow cytometry analysis resource https://floreada.io.

RNA isolation and RT-qPCR

Cells were washed and RNAs were isolated using RNeasy mini kit following manufacturer's

instructions. 1 µg of RNA was reverse transcribed to cDNA using the SuperScript III first-strand

synthesis kit. cDNA was diluted 1:5 in water and mixed with Maxima SYBR Green/1mM of each

primer (Supplementary Table 3). Quantitative PCR was performed on a BioRad CFX96

thermocycler with thermal cycling conditions as followed: Initial denaturation: 95°C for 10 min

followed by 40 cycles of denaturation at 95°C for 15 sec, annealing at 58°C for 30 sec, extension

at 72°C. Final step was performed for 10 sec at 95°C. C_t values from technical triplicates were

averaged and normalized to GAPDH using the $\Delta\Delta C_t$ formula.

TCF expression levels (Supplementary Figure 7E) were curated from the transcriptomic data

available on the Allen Institute website (https://www.allencell.org/genomics.html).

Cell Fractionation

Following appropriate treatment, cells were washed with ice cold 1X PBS and collected as single

cell suspension using gentle cell dissociation as previously described. Cell suspension was

washed with cold 1X PBS and resuspended in 500 μL of cold PBS. 100 μL of the cell suspension

was set aside and mixed with 30 μL of 4X sample buffer. Samples were sonicated for 15 sec at

20% power 3 times, boiled 5min and stored at -20 °C (Whole Cell Lysate fraction or WCL).

Cell fractionation was performed using Cell signaling kit (#9038) and their protocol. Briefly, the

remaining 400 μL of cell suspension was centrifuged at 500 x g at 4°C for 5min. Supernatant was

removed and pellet was resuspended in 250 μL of CIB buffer, vortexed for 5 sec and incubated

on ice for 5 min. Samples were centrifuged at 500 x g at 4°C for 5min and supernatant was

stored in pre-chilled tube (= Cytoplasmic fraction). Pellet was resuspended in 250 μL of MIB

buffer, vortexed for 15 sec and incubated on ice for 5 min. Samples were centrifuged at 8000 x g

at 4°C for 5min and supernatant was discarded (= Membrane and organelle fraction). Pellet was

resuepended in 125 μL of CyNIB buffer, sonicated for 5 sec at 20% power 3 times (cytoskeletal

and nuclear fraction). Each fraction was mixed with the appropriate volume of 4X sample buffer

to obtain a final concentration of 1X. Samples were boiled for 5min and centrifuged before

loading 15 μL.

Nuclear extraction and CUT & RUN Assay

Nuclei extraction: 500,000 cells (per sample) were harvested using TrypLE, washed with

1X PBS, and centrifuged at 600 × g for 3 min at RT. The pellet was resuspended in Nuclear

Extraction (NE) buffer (20 mM HEPES, pH 7.9, 10 mM KCl, 0.1% Triton X-100, 20% glycerol, 1 mM

MnCl$_2$, 0.5 mM spermidine, 1X Roche Complete Protease Inhibitor EDTA-Free) and incubated for

10 min at 4 °C. After incubation, samples were centrifuged at 600 × g, and the isolated nuclei

were resuspended in 100 μL of NE buffer per sample. Nuclei were incubated with 10 μL of

magnetic Concanavalin A beads for 10 min at RT to facilitate binding.

DNA purification: Bead-bound nuclei were incubated with 3 μL of anti-β-catenin

antibody in 50 μL of digitonin buffer (20 mM HEPES, pH 7.5, 150 mM NaCl, 0.5 mM spermidine,

0.01% digitonin, 1X Protease Inhibitor Cocktail) containing 2 mM EDTA overnight at 4 °C.

Following incubation, samples were washed twice with digitonin buffer and resuspended in 50

μL of the same buffer supplemented with 1.5 μL of pAG-MNase. Samples were incubated for 10

min at RT, followed by two additional washes, and subjected to chromatin digestion at 4 °C for 2

hrs with 2 mM CaCl$_2$. To quench the reaction, 33 µL of STOP buffer (340 mM NaCl, 20 mM EDTA,

4 mM EGTA, 50 µg/mL RNase A, 50 µg/mL glycogen) was added, followed by incubation at 37 °C

for 10 min. Beads were collected using a magnetic rack, and the supernatant containing

released DNA was transferred to a new tube. DNA was purified using DNA purification buffers

and spin columns following the manufacturer's instructions and eluted in 30 µL of buffer

<u>qPCR reaction & Data analysis</u>: Reactions were prepared using 1 µL of purified DNA from

the CUT and RUN, with 2 µL Forward and reverse primer mix (5 µM each), 5 µL SYBR Green

master mix (Applied Biosystems), and 2 µL of nuclease-free water. Thermal cycling condition

were set up on a BioRad CFX96 thermocycler as followed: Initial denaturation: 95°C for 10 min

followed by 40 cycles of denaturation at 95°C for 15 sec, annealing at 60°C for 30 sec, extension

at 72°C. Final step was performed for 10 sec at 95°C. Ct values from technical triplicates were

averaged, and ΔCt was calculated by subtracting Ct IgG control from target. Fold enrichment

was calculated using the ΔΔCt method: Fold enrichment=$2^{-\Delta\Delta Ct}$.

<u>GFP-Trap immunoprecipitation</u>

EGFP-CTNNB1 knock-in hiPSCs were harvested using TripLE and nuclei were isolated as

described above. For each reaction, two wells were collected. Both 0h and 24h nuclei were

cryopreserved by slowly freezing them in NE buffer using an isopropanol-filled chiller at -80°C.

Nuclei were quickly thawed using a 37°C water bath, centrifuged at 600 x g and resuspended

using 100 µL of lysis buffer (10 mM Tris/HCl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.5 % NP40).

Nuclear lysates were incubated 5 min on ice and cleared by centrifugation (16000 x g for 10 min

at 4°C. Soluble protein concentration was measured using Precision Red following the

manufacturer's instructions and pellets were mixed with 1X sample buffer (insoluble fraction).

Meanwhile, GFP-Trap beads (20 µL/reaction) were equilibrated three times with 500 µL of ice-

cold dilution buffer (10 mM Tris/HCl pH 7.5, 150 mM NaCl, 0.5 mM EDTA). 150-200 µg of soluble

proteins were prepared in 500 µL of ice-cold dilution buffer and mixed with equilibrated GFP-

Trap beads. Reactions were incubated for 2 hrs at 4°C on a wheel. Beads were then pelleted

(2500 x g, 3min at 4°C) and washed three times with 500 µL of ice-cold wash buffer (10 mM

Tris/HCl pH 7.5, 150 mM NaCl, 0.05 % NP40, 0.5 mM EDTA). Proteins were eluted by adding 50

µL of 2X sample buffer and boiled for 5 min. Eluted proteins were resolved by western blot as

described above. TCF4 binding was measured by densitometry of the TCF4 band (prey)

normalized to the GFP band (bait).

<u>Reproducibility and statistics</u>

All datasets were analyzed using GraphPad Prism10 (version 10.0.3). Individual measurements

(n) were averaged for each biological repeat (N). Datasets were tested for normality (Shapiro-

Wilk test for normality of biological replicates) before applying the appropriate statistical test on

the biological repeats (N), except if mentioned differently in the figure legend. Error bars

represent S.D. except where stated otherwise. For datasets displayed as superplots, individual

measurements and biological repeats are represented by small and large dots respectively.

Violin plots display median (thick dotted line) and quartiles (thin dotted lines). Each dataset for

a biological repeat is color coded. Significance levels are given as follows and exact p-values are

indicated in each figure: n.s., not significant ($p \geq 0.05$); $*p < 0.05$; $**p < 0.01$; $***p < 0.001$; and

$****p < 0.0001$.

All experiments, including representative MaxIPs and western blots, were performed at

least three times independently as biological repeats, unless stated otherwise in the legends.

No statistical method was used to pre-determine the sample size, the experiments were not

randomized and the investigators were not blinded to allocation during experiments and

outcome assessment.

Figure legends

<u>Figure 1: Increased actomyosin contractility is sufficient to block mesoderm commitment and</u>

<u>EMT</u>

(A) Experimental design for co-culture experiment. hiPSC were transduced with lentivector

expressing ROCK2^{CA} or MLCK^{CA}. mVenus positive-transduced cells were mixed with mVenus-

negative WT hiPSCs. ROCK2^{CA}/MLCK^{CA} expression is induced by addition of Doxycycline 24h

prior to initiating mesoderm differentiation. Doxycycline induction was maintained during the 3

days of differentiation using CHIR before fixing and staining.

(B-H) Representative MaxIP immunofluorescence for Brachyury (B), EOMES (D), Slug (F) and ZO-

1 (H) in Vehicle and Doxycycline-induced ROCK2^{CA} co-culture. mVenus-positive cell clusters are

highlighted by an orange dotted line. Individual channels are presented as inverted LUT. Scale

bar = 50 µm. For (H), magnified mVenus-positive and mVenus-negative areas are shown as

insets. Scale bar = 20 µm. Quantification of Brachyury (C), EOMES (E) and Slug-positive cells (G)

is reported as violin plot, comparing mVenus-positive *vs* mVenus-negative clusters in the

presence or absence of Doxycycline. Median (Plain red line) and quartiles (Dotted black lines).

For Brachyury, n = 7 (+ Veh) and n=15 (+ Dox) across N=3 independent biological repeats. For

EOMES, n=9 across N=3 independent biological repeats. For Slug, n=9 (+ Veh) and n=10 (+ Dox)

across N=3 independent biological repeats. One-way ANOVA with Tukey's multiple comparisons

post-test was performed.

Figure 2: Genetic inhibition of actomyosin contractility promotes hiPSC conversion to the

mesoderm lineage

(A) Summary of the main regulators of MLC phosphorylation.

(B) Schematic representation of full length human MYPT1 (1030 aa), interacting with PP1c

phosphatase and a protein of unknown function M20. Truncation of MYPT1 Cterm Δ301-1030

leads to a constitutive recruitment and activation of PP1C. Truncated MYPT1 was fused with

NES-mNeonGreen and is referred to as MYPT1^{CA}-NES-mNG.

(C) Immunoblot of phospho T18/S19 MLC2 (ppMLC2) and total MLC2 in WT hiPSCs treated with

Doxycycline or hiPSC expressing MYPT1^{CA}-NES-mNG in the presence or absence of Doxycycline.

α-Tubulin was used as loading control. Molecular weights (M.W.) are displayed on the right side.

(D, E) – Representative MaxIP immunofluorescence of hiPSC expressing MYPT1^{CA}-NES-mNG and

treated or not with Doxycycline. Cells were fixed and stained for phospho T18/S19 MLC2

(ppML2 - Magenta), DNA (blue) and tight junction marker ZO-1 (Orange). Scale bar = 50 µm (D).

Fraction of the cellular area positive for ppMLC2 following Doxycycline induction of MYPT1^{CA}-

NES-mNG hiPSC and reported as violin plots. Median (Plain red line) and quartiles (Dotted black

lines). n=32 technical repeats across N=3 independent biological repeats. Two-tailed unpaired t

test was performed on the biological repeats (E).

(F) Experimental design. hiPSC cells were transduced with pInducer20-MYPT1^{CA}-NES-mNG and

stable population was selected with puromycin. MYPT1^{CA} cells were treated or not with

doxycycline for 16h and treated with CHIR supplemented or not with doxycycline. Parental WT

line was differentiated with Doxycycline.

(G, H) Representative immunoblot for Brachyury (mesoderm marker), Snail and Slug (EMT

markers) using WT and MYPT1^{CA}-NES-mNG hiPSCs, following CHIR treatment (0h-48h) in the

presence or absence of Doxycycline, as shown in (F). Molecular weights (M.W.) are displayed on

the right side (G). Brachyury expression was quantified by densitometry and normalized to α-

Tubulin as loading control across N=3-4 independent biological repeats. Mean and S.D. are

displayed. A one-way ANOVA with Tukey's multiple comparisons post-test was performed (H).

(I) Immunoblot of PARP (cell death marker) and Brachyury (Mesoderm marker) during

mesoderm commitment (0hr-48hrs CHIR) using MYPT1^{CA}-NES-mNG hiPSCs induced +/

Doxycycline, as shown in (F). Molecular weights (M.W.) are displayed on the right side.

(J, K) Representative MaxIP immunofluorescences of WT and MYPT1^{CA}-NES-mNG hiPSCs treated

with CHIR for 24h +/- Doxycycline as shown in (F). Cells were stained for nuclei (DNA), EMT

marker (ZO-1 – Inverted LUT) and Mesoderm marker (T/Bra – Inverted LUT). Scale bar = 100 μm

(J). Quantification of Brachyury expression is presented as violin plots. Median (Plain red line)

and quartiles (Dotted black lines). n=15-16 technical repeats across N=3 independent biological

repeats. One-way ANOVA with Šidák's multiple comparisons post-test was performed on

biological repeats (K).

<u>Figure 3: Pharmaceutical inhibition of actomyosin contractility promotes hiPSC and hESC</u>

<u>conversion to the mesoderm lineage</u>

(A) Experimental design. WT hiPSCs or hESCs were pre-treated overnight with small molecules

to lower contractility, followed by differentiation.

(B) Immunoblot of Brachyury during hiPSC-to-mesoderm commitment (0-48h) in the presence

(+ Y-27632) or absence (+ Veh) of ROCK inhibitor, as shown in (A). Molecular weights (M.W.) are

displayed on the right side.

(C) Brachyury expression was quantified by densitometry and normalized to α-Tubulin as

loading control across N=3 independent biological repeats. Mean and S.D. are displayed. Two-

tailed unpaired t test was performed.

(D - G) Relative expression of mesoderm markers (*MESP1, TBX6*) (D, E) and EMT markers

(*SNAI1, SNAI2*) (F, G) 48h post CHIR treatment in the presence (+Y-27) or absence (+Veh) of

ROCK inhibitor as shown in (A). N=3 independent biological repeats except for *MESP1* N=4

independent biological repeats. Mean and S.D. are displayed. Two-tailed unpaired t test was

performed.

(H) – Representative MaxIP immunofluorescences during hiPSC-to-mesoderm commitment (0-

48h) in the presence (+ Y-27632) or absence (+ Veh) of ROCK inhibitor as shown in (A). Cells

were stained with a nuclear marker (DNA), EMT marker (ZO-1 – Inverted LUT) and Mesoderm

marker (T/Bra– Inverted LUT). Scale bar = 50 µm.

(I, J) Quantification of Brachyury expression (I) and ZO-1 intactness (J) is represented as violin

plots with individual measurements (small dots) averaged for each biological repeat (large dots).

Median (Plain red line) and quartiles (Dotted black lines). n=10-15 technical repeats across N=3

independent biological repeats. One-way ANOVA with Tukey's multiple comparisons post-test

was performed on biological repeats.

(K, L) Analysis of Brachyury expression by flow cytometry following 24h CHIR treatment in the

presence (+Y-27632) or absence (+Veh) of ROCK inhibitor, as shown in (A), using Bra-

mNeonGreen knock-in H9 hESC line (K). Quantification of the percentage of mNeonGreen-

positive cells from N=4 independent biological repeats. Mean and S.D. are displayed. Two-tailed

unpaired t test was performed (L).

(M, N) Representative MaxIP immunofluorescence of hiPSCs treated with a MLCK inhibitor (ML-

7) or Vehicle (DMSO), as shown in (A). Cells were stained for nuclei (DNA) and phospho T18/S19

Myosin Light Chain 2 (ppMLC2 – Inverted LUT). Scale bar = 50 µm. Magnified views of the yellow

dotted ROI are shown for ppMLC2. Scale bar = 20 µm (M). Percentage of cellular area positive

for ppMLC2 following Vehicle or ML-7 treatment is reported as violin pots. Median (Plain red

line) and quartiles (Dotted black lines). n=15 technical repeats across N=3 independent

biological repeats. Two-tailed unpaired t test was performed on the biological repeats (N).

(O, P) Immunoblot of Brachyury (mesoderm) and PARP (cell death marker) using hiPSCs in the

presence (+ ML-7) or absence (+ Veh) of MLCK inhibitor at basal state (0h) and during CHIR

treatment (24hrs, 48hrs), as shown in (A). Cells treated with CHIR and Q-VD (Caspase inhibitor)

for 48hrs were used as control for PARP immunoblot. Molecular weights (M.W.) are displayed on

the right side (O). Brachyury expression was quantified by densitometry and normalized to α-

Tubulin as loading control across N=3 independent biological repeats. Mean and S.D. are

displayed. Mann-Whitney test was performed for the 24hrs timepoints and two-tailed unpaired

t test was performed for the 48hrs timepoints (P).

(Q) Experimental design. Effects of Y-27632 was tested at different time along the mesoderm

commitment. ROCK inhibitor was added 12h before differentiation (-12h), at the time of

differentiation (0h) or 12h after differentiation (+12h). All conditions were collected 24h post

differentiation.

(R, S) Representative immunoblot for Brachyury following addition of Vehicle or Y-27632 as

shown in (Q). Molecular weights (M.W.) are displayed on the right side (R). Brachyury expression

was quantified by densitometry and normalized to α-Tubulin as loading control across N=3

independent biological repeats. Mean and S.D. are displayed. Two-way ANOVA with Šidák's

multiple comparisons post-test was performed (S).

Figure 4: Adherens junctions disengagement promotes mesoderm fate

(A) Experimental design. hiPSCs were pre-treated with an acute concentration of EGTA to

disrupt calcium-dependent E-Cadherin junctions. Following pre-treatment, hiPSCs were

differentiated in the presence of a lower concentration of EGTA.

(B-E) Representative immunoblot of hiPSCs treated as shown in (A) and probed for Brachyury

(mesoderm marker), Snail and Slug (EMT markers) and Pax6 and Otx1/2 (Neuroectoderm

markers as negative controls). Neural Stem Cell (NSC) lysate was used as positive control for

Pax6 and Otx1/2. Non-specific band is designated by an asterisk. Molecular weights (M.W.) are

displayed on the right side (B). Expression of Brachyury (C), Snail (D) and Slug (E) was quantified

by densitometry and normalized to α-Tubulin across N=3 independent biological repeats. Mean

and S.D. are displayed. One-Way ANOVA with Šidák's multiple comparisons post-test was

performed.

(F, G) Representative immunoblot of control (Ctr) and α-catenin knockdown hiPSC lines at basal

state and probed for total and phospho-MLC2. Molecular weights (M.W.) are displayed on the

right side (F). Quantification of active MLC2 (ppMLC2/MLC2) was measured by densitometry

across N=3 independent biological repeats. Mean and S.D. are displayed. One-Way ANOVA with

Dunnett's multiple comparison post-test was performed (G).

(H, I) Representative immunoblot of control (Ctr) and α-catenin knockdown (*CTNNA1* KD #1 and

#2) hiPSC lines, differentiated for up to 48hrs in the presence or absence of ROCK inhibitor Y-

27632 and probed for α-catenin, Brachyury and α-Tubulin as loading control. Molecular weights

(M.W.) are displayed on the right side (H). Normalized Brachyury expression was measured by

densitometry across N=3 independent biological repeats. Mean and S.D. are displayed. One-

Way ANOVA with Šidák's multiple comparisons post-test was performed (I).

Figure 5: Contractility differentially regulates β-catenin localization at adherens junctions

(A) Experimental design

(B) Representative immunofluorescence of fixed hiPSCs treated with Vehicle, 4µg/mL CN03 or

10 µM Y-27632 for 3h and stained for ppMLC2 and β-catenin (inverted LUT), as shown in (A).

Scale bare = 50 µm.

(C) Linear regression analysis of junctional β-catenin and cellular ppMLC2 was quantified across

N=3 independent biological repeats (representing n=16 for Vehicle, n=19 for CN03 and n=17

technical repeats for Y-27632). R^2 value is reported.

(D) Experimental design

(E, F) Representative still pictures from mEGFP-β-catenin knock-in (KI) hiPSCs treated with

Vehicle, 4µg/mL CN03 or 10 µM Y-27632 for 3h, as shown in (D). Scale bare = 50 µm (E).

Quantification of junctional β-catenin is reported for the different treatment across N=4

independent biological repeats (representing n=25 technical repeats). Violin plots representing

median and quartiles. One-Way ANOVA with Dunnett's multiple comparison post-test was

performed on biological repeats (F).

Figure 6: Reduced cell contractility promotes β-catenin nuclear activity and enhances binding to

mesodermal gene

(A) Representative immunoblot as explained in Supplementary Figure 6A. MYPT1[CA]-NES-mNG

hiPSCs were induced with CHIR for 24hrs in the presence or absence of Doxycycline. Whole cell

lysate (WCL) was collected and loaded as input while cytosolic and nuclear fractions were

extracted. Fraction purity was probed using cytosolic (GAPDH) and nuclear (Lamin A/C) markers.

Active β-catenin (non-phospho S45) enrichment was probed for each fraction and α-Tubulin was

used as loading control. Molecular weights (M.W.) are displayed on the right side.

(B) Quantification of active β-catenin enrichment in the cytosolic (left) and nuclear (right)

fractions. N=4 independent biological repeats. Mean and S.D. are displayed. One-way ANOVA

with Šidák's multiple comparisons post-test was performed.

(C) Representative immunofluorescence for active non-phospho S45 β-catenin (Inverted LUT –

confocal slice) and mNeonGreen (MaxIP). MYPT1^{CA}-NES-mNG hiPSCs were fixed at 0h and 24h

post CHIR treatment +/- Doxycycline. Scale bar = 50 μm. Insets represent a magnified view of

the active non-phospho S45 β-catenin with an example of the nuclei binary mask overlaid

(pink). Scale bar = 10 μm.

(D, E) Stills confocal timelapse imaging of mEGFP-β-catenin knock-in hiPSCs treated with CHIR

+/- Y-26732 (Related to Supplementary movie 3). Time is shown in hours. Scale bar = 10 μm (D).

mEGFP-β-catenin intensity is reported as a scan line plot across a cell at 0h and 20h (E).

(F) Experiment design for GFP-Trap immunoprecipitation. mEGFP-β-catenin knock-in hiPSCs

were pre-treated with Vehicle or Y-27632 and differentiated for 24h. Nuclei were isolated and

nuclear lysates were mixed with GFP-Trap beads.

(G, H) Representative GFP-Trap immunoprecipitation of mEGFP-β-catenin knock-in hiPSCs,

probed for TCF7L2/TCF4 and α-Tubulin as loading control. Molecular weights (M.W.) are

displayed on the right side (G). Quantification of TCF4 (Prey) binding to mEGFP-β-catenin (Bait)

was measured by densitometry across N=3 independent biological repeats. Mean and S.D. are

displayed. One-way ANOVA with Šidák's multiple comparisons post-test was performed.

(I) CUT & RUN assay for IgG or β-catenin binding to *TBXT* promoter in WT hiPSC pre-treated with

Veh or Y-27632 (0h CHIR) or following 24h of CHIR treatment in the presence of absence of

ROCK inhibitor. N=3 independent biological repeats. Mean and S.D. are displayed. One-way

ANOVA with Šidák's multiple comparisons post-test was performed.

Figure 7: Working model

Effect of actomyosin contractility on mesoderm lineage commitment. Each panel represents a

specific contractile status at basal state (0h - top) and after 48h (bottom) of CHIR treatment. At

basal state, β-catenin is degraded and localized at AJs. The amount of junctional β-catenin

correlates with the level of contractility experienced by hiPSCs. During differentiation, β-catenin

accumulates in the nucleus and activates a mesoderm program through its binding with TCF4.

Supplementary Figure 1: Increased actomyosin contractility is sufficient to block mesoderm

commitment and EMT (Related to Figure 1)

(A) Experimental design. hiPSCs were pre-treated with 4 µg/mL CN03 or vehicle for 3 hours.

Differentiation was initiated with CHIR supplemented with Vehicle or 4 µg/mL CN03 for 24h

before fixation and staining.

(B) Representative MaxIP immunofluorescence of hiPSC treated as explained in (A) and stained

for phospho-MLC2 (Inverted LUT). Scale bar = 50 µm

(C, D) Representative MaxIP immunofluorescence of hiPSC treated as explained in (A) and

stained for Brachyury (Inverted LUT), nuclei and ZO-1. Scale bar = 50 µm (C). Violin plot (median,

quartiles) reporting the percentage of Brachyury-positive nuclei after 24h of differentiation and

quantified across N=4 independent biological repeats (representing n=20 technical repeats).

Unpaired t test was performed on biological replicates (D).

(E) Representative MaxIP immunofluorescence of Doxycycline-induced ROCK2^{CA} hiPSCs

(mVenus-positive) mixed with WT hiPSCs (mVenus-negative). Cells were stained for DNA

(Nuclei), F-actin (top – Inverted LUT) and phospho T18/S19 Myosin Light Chain 2 (Bottom -

ppMLC2 – Inverted LUT). Scale bar = 50 µm. Magnified views comparing mVenus-positive

(Green dotted ROI) and mVenus-negative (Black dotted ROI) areas are shown for F-Actin and

ppMLC2. Scale bar = 10 µm.

(F) Percentage of the ROI area positive for mVenus based on cluster considered as mVenus-

Negative and Positive using the ROCK2^{CA} line. On average, mVenus negative clusters have <15%

of their area positive for mVenus. Mean and S.E.M are displayed. n=7 (no Dox) and n=15 fields

of view (+Dox) across N=3 independent biological experiments.

(G) Representative MaxIP immunofluorescence of Doxycycline-induced MLCK^{CA} hiPSCs (mVenus-

positive) mixed with WT hiPSCs (mVenus-negative). Cells were stained for DNA (Nuclei), F-actin

(top – Inverted LUT) and phospho T18/S19 Myosin Light Chain 2 (Bottom - ppMLC2 – Inverted

LUT). Scale bar = 50 µm. Magnified views comparing mVenus-positive (Green dotted ROI) and

mVenus-negative (Black dotted ROI) areas are shown for F-Actin and ppMLC2. Scale bar = 10

µm.

(H) Percentage of the ROI area positive for mVenus based on cluster considered as mVenus-

Negative and Positive using the MLCK^{CA} line. On average, mVenus negative clusters have <15%

of their area positive for mVenus. Mean and S.E.M are displayed. n=7 (no Dox) and n=18 fields

of view (+Dox) across N=3 independent biological experiments.

(I, J) Representative MaxIP immunofluorescence for Brachyury in Vehicle (+Veh) and

Doxycycline-induced MLCK^{CA} co-culture. mVenus-positive cell clusters are highlighted by an

orange dotted line. Individual channels are inverted LUT. Scale bar = 50 µm (I). Percentage of

Brachyury-positive nuclei is reported as violin plot, comparing mVenus-positive *vs* mVenus-

negative cluster in the presence or absence of Doxycycline. Median (Plain red line) and quartiles

(Dotted black lines). n=7 (+Veh) and n=18 (+ Dox) fields of view (small dots) across N=3

independent biological repeats (large dots). One-way ANOVA with Tukey's multiple comparisons

post-test was performed (J).

(K) Representative MaxIP immunofluorescence for EOMES, Slug and ZO-1 in Doxycycline-

induced MLCK^{CA} co-culture. mVenus-positive cell clusters are highlighted by an orange dotted

line. Individual channels are inverted LUT. Scale bar = 50 µm.

<u>Supplementary Figure 2: Pharmaceutical inhibition of actomyosin contractility promotes hiPSC</u>

<u>and hESC conversion to the mesoderm lineage (Related to Figure 3)</u>

(A, B) Representative MaxIP immunofluorescences of hiPSCs treated with a ROCK inhibitor (Y-

27632) or Vehicle (H$_2$O) and stained for nuclei marker (DNA) and phospho T18/S19 Myosin Light

Chain 2 (ppMLC2 – Inverted LUT). Scale bar = 50 µm. Magnified views of the yellow dotted ROI

are shown for ppMLC2. Scale bar = 20 µm (A). Fraction of the cellular area positive for ppMLC2

following Y-27632 treatment and reported as violin pots. Median (Plain red line) and quartiles

(Dotted black lines). n=30 technical repeats across N=3 independent biological repeats. Two-

tailed unpaired t test was performed on the biological repeat (B).

(C) Relative expression of genes not directly related to mesoderm commitment at 0h and 48h

post CHIR treatment in the presence (+Y-27) or absence (+Veh) of ROCK inhibitor. N=4

independent biological repeats. Mean and S.D. are displayed. Two-tailed unpaired t test was

performed.

(D) Representative MaxIP immunofluorescences of hiPSC treated 48h with CHIR in the presence

(+ Y-27632) or absence (+ Veh) of ROCK inhibitor. Cells were stained for nuclei marker (DNA),

EMT markers (ZO-1 and Slug – Inverted LUT). Scale bar = 50 µm.

(E, F) Representative MaxIP immunofluorescences of hiPSCs treated with a ROCK inhibitor (H-

1152) or Vehicle (H$_2$O) and stained for nuclei marker (DNA) and phospho T18/S19 Myosin Light

Chain 2 (ppMLC2 – Inverted LUT). Scale bar = 50 µm. Magnified views of the yellow dotted ROI

are shown for ppMLC2. Scale bar = 20 μm (E). Fraction of the cellular area positive for ppMLC2

following H-1152 treatment and reported as violin pots. Median (Plain red line) and quartiles

(Dotted black lines). n=15 technical repeats across N=3 independent biological repeats. Two-

tailed unpaired t test was performed on the biological repeats (F).

(G-K) Relative expression of mesoderm markers (*TBXT*, *MESP1, TBX6*) (G-I) and EMT markers

(*SNAI1, SNAI2*) (J, K) 48h post CHIR treatment in the presence (+H-1152) or absence (+Veh) of

ROCK inhibitor. N=3 independent biological repeats. Mean and S.D. are displayed. Two-tailed

unpaired t test was performed.

(L) Immunoblot probing for mesoderm marker expression (T/Bra) and cell death marker (PARP

cleavage) during hiPSC-to-mesoderm commitment (0h-48h CHIR) in the presence of ROCK

inhibitor (+H-1152). Addition of the pan-caspase inhibitor Q-VD-OPH (+ Q-VD) to the

differentiation medium totally blocks mesoderm commitment and was used as control. α-

Tubulin was used as loading control. Molecular weights (M.W.) are displayed on the right side.

(M) Representative MaxIP immunofluorescences for mesoderm marker Brachyury (T/Bra –

Inverted LUT) and nuclei. Cells were fixed 24h post-CHIR treatment in the presence or absence

of ROCK inhibitor H-1152. Scale bar = 50 μm

(N) Representative immunoblot of PARP (cell death), T/Bra (mesoderm), Oct4 and Nanog

(Pluripotency markers) using hESC H9 in the presence (+ Y-27632) or absence (+ Veh) of ROCK

inhibitor, as shown in Figure 3A. Molecular weights (M.W.) are displayed on the right side.

(O-Q) Brachyury expression (O), PARP cleavage (P) and Nanog expression (Q) was quantified by

densitometry and normalized to α-Tubulin as loading control across N=4 (PARP) and N=3

(Brachyury and Nanog) independent biological repeats. Mean and S.D. are displayed. Two-tailed

unpaired t test was performed.

<u>Supplementary Figure 3: hiPSC differentiation along the cardiac mesoderm increases</u>

<u>actomyosin-driven contractility</u>

(A) Bright field imaging of WT hiPSCs treated with CHIR in the presence or absence of a pan-

Caspase inhibitor Q-VD-OPH. Retraction area is highlighted in yellow. Time is presented as

hh:mm. Scale bar = 20 μm. (Related to Supplementary movie 1).

(B, C) Confocal XZ orthogonal views (B) and Maximum Intensity Projections (MaxIP) (C) of hiPSCs

at basal state (0h CHIR) or post-CHIR treatment (24h and 48h CHIR), stained for DNA (Cyan) and

F-Actin (Grey). Scale bar = 10 μm.

(D) Quantification of the monolayer height at 0h, 24h and 48h post CHIR treatment from (B).

Violin plots represent individual measurements (small dots) averaged for each biological repeat

(large dots). Median (Plain red line) and quartiles (Dotted black lines). n=32 technical repeats

across N=7 independent biological repeats. A Kruskal-Wallis test with Dunn's multiple

comparison post-test was performed on biological repeats.

(E-G) Representative MaxIP immunofluorescence of WT hiPSC during CHIR treatment. Fixed cells

were stained for DNA marker (Hoechst). Scale bar = 20 μm (E). Average nuclear size from n=15

fields of view across N=3 independent biological repeats. Median (Plain red line) and quartiles

(Dotted black lines) are displayed. Two-tailed unpaired t test was performed (F). Correlation of

nuclear area over cell number for each field of view during CHIR treatment. Biological repeats

are color-coded. Pearson correlation coefficient is displayed for each treatment (G).

(H) Representative MaxIP immunofluorescence of hiPSCs at basal state (0h) or post-CHIR

treatment (24 and 48h) stained for phospho T18/S19 MLC2 (ppMLC2). Pixels are color-coded by

intensity using the fire LUT. Scale bar = 50 μm

(I) Percentage of ppMLC2-positive area at 0hr, 24hrs, 48hrs post CHIR is reported as violin plots.

Median (Plain red line) and quartiles (Dotted black lines). n=15 technical repeats across N=3

independent biological repeats. A Kruskal-Wallis test with Dunn's multiple comparison post-test

was performed on the technical repeats.

(J) Retardance measurements obtained from hiPSCs at basal state (0h), post-CHIR treatment

(24h, 48h, 72h) or treated with Y-27632 at 0h (Y-27) using Quantitative Polarization Microscopy

imaging. Median (Plain red line) and quartiles (Dotted black lines). n=240 technical repeats for

0hr, 24hrs, 48hrs, 72hrs and n=120 technical repeats for Y-27 across N=3 independent biological

repeats. A one-way ANOVA with Šidák's multiple comparisons post-test was performed on the

biological repeats.

Supplementary Figure 4: Key mechanosensitive pathways are not involved in mesoderm

specification enhancement following decreased contractility

(A, B) Immunoblot for phospho S127 YAP (pYAP) and total YAP following differentiation (0-48h

CHIR) in the presence or absence of ROCK inhibitor Y-27632. Brachyury expression was also

probed as an internal positive control and α-Tubulin was used as loading control. Molecular

weights (M.W.) are displayed on the right side (A). Ratio of phospho/total YAP was quantified

across N=4 independent biological repeats. Mean and S.D. are displayed. One-way ANOVA with

Šidák's multiple comparisons post-test was performed (B)

(C) Relative expression of YAP-target genes *CTGF* and *CYR61* during differentiation +/- Y-27632.

N=4 independent biological repeats. Mean and S.D. are displayed. One-way ANOVA with Šidák's

multiple comparisons post-test was performed.

(D) Relative expression of canonical WNT (*WNT3A, WNT8A*), non-canonical WNT (*WNT4*) and

mesoderm marker (*TBX6*) in MYPT1^{CA}-NES-mNG cells treated with CHIR (0-48h) in the presence

of Vehicle or Doxycycline. N=3 independent biological repeats. Mean and S.D. are displayed.

One-way ANOVA with Šidák's multiple comparisons post-test was performed.

(E) Relative expression of *TBX6* and *MESP1* (mesoderm markers) following CHIR treatment +/- Y-

27632 complemented or not with 7.5 µM Porcupine inhibitor (IWP-2 or DMSO respectively).

N=3 independent biological repeats. Mean and S.D. are displayed. One-way ANOVA with Tukey's

multiple comparisons post-test was performed.

<u>Supplementary Figure 5: EGTA treatment and validation *CTNNA1* knockdown hiPSCs (related to</u>

<u>Figure 4)</u>

(A) Bright field picture showing the colony morphology following EGTA pre-treatment. Colony

appears less compact and individual cell boundaries can be distinguished. Scale bar = 20 µm.

(B, C) Representative immunofluorescence of hiPSCs treated as shown in Figure 4A and stained

for Brachyury. Scale bar = 50 µm (B). Quantification of Brachyury-positive nuclei is shown

following EGTA treatment. Mean and S.D. are displayed. n=18 (Veh) and n=20 fields of view

across N=3 independent biological repeats. Two-tailed unpaired t test was performed on

biological repeats (C).

(D, E) Representative immunofluorescence of hiPSC treated as shown in Figure 4A and stained

for ppMLC2. Scale bar = 50 µm (D). Quantification of ppMLC2-positive area is shown following

EGTA treatment. Mean and S.D. are displayed. n=20 (Veh) and n=25 fields of view across N=4

independent biological repeats. Two-tailed unpaired t test was performed on biological repeats

(E).

(F-I) Representative immunoblot of control (Ctr) or *CTNNA1* knockdown hiPSCs, probed for α-

catenin, Nanog and Oct-4 (pluripotency) and α-Tubulin as loading control. Molecular weights

(M.W.) are displayed on the right side (F). Quantification of α-catenin (G), Nanog (H) and Oct-4

(I) expression was obtained by densitometry across N=5-6 (α-catenin), N=4 (Nanog) and N=3

(Oct-4) independent biological repeats. Mean and S.D. are displayed. One-Way ANOVA with

Dunnett's multiple comparisons post-test was performed.

1018

<u>Supplementary Figure 6: Contractility affects β-catenin localization at adherens junctions</u>

<u>(related to Figure 5)</u>

(A) Immunoblot assessing the effects of CN03 treatment at 4h and 16h. Membrane was probed

for total and phospho-ML2 (MLC2 and ppMLC2 respectively), Nanog and Oct-4 and α-Tubulin as

loading control. Molecular weights (M.W.) are displayed on the right side.

(B) Independent technical repeats from Figure 5 A-B. Repeats across the 3 biological repeats

were averaged and shown as a dot with error bars (S.D.).

(C) Still picture from overnight confocal imaging of mEGFP-β-catenin knock-in hiPSC treated with

Vehicle, CN03 and Y-27632 at basal state. Pixels are color-coded by intensity using the Red Fire

LUT. Scale bar = 20 μm. (Related to Supplementary video 2).

(D) Experimental design. Cells were pre-treated overnight with Vehicle or Y-27632 or CN03 in

mTeSR1 and differentiated using CHIR media complemented with Vehicle or Y-27632 or CN03

for 24h before collecting protein lysates

(E, F) Representative immunoblot for total β-catenin. Membrane was also probed for Brachyury

to show effect of each drug and α-Tubulin as loading control. Molecular weights (M.W.) are

displayed on the right side (E). β-catenin expression was quantified across N=6 (Vehicle and Y-

27632) and N=4 (CN03) independent biological repeats. Mean and S.D. are displayed. One-way

ANOVA with Šidák's multiple comparisons post-test was performed (F).

1037

<u>Supplementary Figure 7: Reduced cell contractility promotes nuclear accumulation of β-catenin</u>

<u>(related to Figure 6)</u>

(A) Experiment design for treatments and cell fractionation.

(B) Compartment purity following fractionation was assessed by quantifying Lamin A/C and

GAPDH enrichment in both, cytosolic and nuclear fractions across N=4 independent biological

repeats.

(C) Schematic representation of the quantification pipeline for nuclear β-catenin.

(D) Quantification of active non-phospho S45 β-catenin mean intensity as depicted in (C). n =

10-12 technical repeats across N=3 independent biological repeats. Mean and S.D. are

displayed. One-way ANOVA with Šidák's multiple comparisons post-test was performed.

(E) Expression of TCF genes (transcripts per million – tpm) from parental WTC hiPSC obtained

from the Allen Institute transcriptomic data. N=6 independent biological repeats.

(F) Positive control for CUT & RUN assay using IgG and β-catenin binding to WNT target genes

(*AXIN2* and *SP5*) between 0h and 24h of CHIR treatment. Mean and S.D. are displayed. N=3

independent biological repeats. One-Way ANOVA with Dunnett's multiple comparisons post-test

was performed.

(G) Relative expression of WNT target genes (*LEF1* and *AXIN2*) with increasing concentration of

CHIR in the presence (Magenta) or absence (Cyan) of ROCK inhibitor Y-27632. N=7 independent

biological repeats. Mean and S.E.M. are displayed. One-way ANOVA with Šidák's multiple

comparisons post-test was performed.

1058

1059   <u>Supplementary Video 1</u>

1060   Bright field movie of hiPSC colony treated with CHIR supplemented with Vehicle (DMSO) or Q-

1061   VD-OPH (cell death inhibitor). Time in hh:mm. Scale bar = 20 μm.

1062

1063   <u>Supplementary Video 2</u>

1064   Confocal imaging of mEGFP-β-catenin knock-in hiPSC treated with Vehicle (water), 4μg/mL

1065   CN03 or 10 μM Y-27632 in mTeSR1. Red Fire LUT reflects the pixel intensity. Time in hh:mm:ss.

1066   Scale bar = 20 μm.

1067

1068   <u>Supplementary Video 3</u>

1069   Confocal imaging of mEGFP-β-catenin knock-in hiPSC treated with CHIR supplemented with

1070   Vehicle (water) or 10 μM Y-27632. Time in hh:mm. Scale bar = 10 μm.

1071

References

1.   Firmin, J. *et al.* Mechanics of human embryo compaction. *Nature* **629**, 646–651 (2024).

2.   Mundhe, G. *et al.* Spatial patterning of contractility by a mechanogen gradient underlies *Drosophila* gastrulation. Preprint at https://doi.org/10.1101/2025.04.11.648359 (2025).

3.   Lecuit, T. & Lenne, P.-F. Cell surface mechanics and the control of cell shape, tissue patterns and morphogenesis. *Nat Rev Mol Cell Biol* **8**, 633–644 (2007).

4.   Goodwin, K. & Nelson, C. M. Mechanics of Development. *Dev Cell* **56**, 240–250 (2021).

5.   Caldarelli, P. *et al.* Self-organized tissue mechanics underlie embryonic regulation. *Nature* **633**, 887–894 (2024).

6.   De Belly, H., Paluch, E. K. & Chalut, K. J. Interplay between mechanics and signalling in regulating cell fate. *Nat Rev Mol Cell Biol* **23**, 465–480 (2022).

7.   Khalilgharibi, N. *et al.* Stress relaxation in epithelial monolayers is controlled by the actomyosin cortex. *Nat Phys* **15**, 839–847 (2019).

8.   Yang, Y.-A. *et al.* Local contractions regulate E-cadherin rigidity sensing. *Sci Adv* **8**, (2022).

9.   Shyer, A. E. *et al.* Emergent cellular self-organization and mechanosensation initiate follicle pattern in the avian skin. *Science (1979)* **357**, 811–815 (2017).

10.   Behrndt, M. *et al.* Forces Driving Epithelial Spreading in Zebrafish Gastrulation. *Science (1979)* **338**, 257–260 (2012).

11.   Hall, E. T., Hoesing, E., Sinkovics, E. & Verheyen, E. M. Actomyosin contractility modulates Wnt signaling through adherens junction stability. *Mol Biol Cell* **30**, 411–426 (2019).

12.   Lian, X. *et al.* Robust cardiomyocyte differentiation from human pluripotent stem cells via temporal modulation of canonical Wnt signaling. *Proceedings of the National Academy of Sciences* **109**, (2012).

13.   Lian, X. *et al.* Directed cardiomyocyte differentiation from human pluripotent stem cells by modulating Wnt/β-catenin signaling under fully defined conditions. *Nat Protoc* **8**, 162–175 (2013).

14.   Zhao, M., Tang, Y., Zhou, Y. & Zhang, J. Deciphering Role of Wnt Signalling in Cardiac Mesoderm and Cardiomyocyte Differentiation from Human iPSCs: Four-dimensional control of Wnt pathway for hiPSC-CMs differentiation. *Sci Rep* **9**, 19389 (2019).

15.   Fort, L., Gama, V. & Macara, I. G. Stem cell conversion to the cardiac lineage requires nucleotide signalling from apoptosing cells. *Nat Cell Biol* **24**, 434–447 (2022).

16.   Muncie, J. M. *et al.* Mechanical Tension Promotes Formation of Gastrulation-like Nodes and Patterns Mesoderm Specification in Human Embryonic Stem Cells. *Dev Cell* **55**, 679-694.e11 (2020).

17.   Przybyla, L., Lakins, J. N. & Weaver, V. M. Tissue Mechanics Orchestrate Wnt-Dependent Human Embryonic Stem Cell Differentiation. *Cell Stem Cell* **19**, 462–475 (2016).

18.   van der Wal, T. & van Amerongen, R. Walking the tight wire between cell adhesion and WNT signalling: a balancing act for β-catenin. *Open Biol* **10**, (2020).

19. Orsulic, S., Huber, O., Aberle, H., Arnold, S. & Kemler, R. E-cadherin binding prevents
β-catenin nuclear localization and β-catenin/LEF-1-mediated transactivation. *J Cell*
Sci **112**, 1237–1245 (1999).
20. Pukhlyakova, E., Aman, A. J., Elsayad, K. & Technau, U. β-Catenin–dependent
mechanotransduction dates back to the common ancestor of Cnidaria and Bilateria.
Proceedings of the National Academy of Sciences **115**, 6231–6236 (2018).
21. Gayrard, C., Bernaudin, C., Déjardin, T., Seiler, C. & Borghi, N. Src- and confinement-
dependent FAK activation causes E-cadherin relaxation and β-catenin activity.
Journal of Cell Biology **217**, 1063–1077 (2018).
22. Schmidt, G. *et al*. Gln 63 of Rho is deamidated by Escherichia coli cytotoxic
necrotizing factor-1. *Nature* **387**, 725–729 (1997).
23. Flatau, G. *et al*. Toxin-induced activation of the G protein p21 Rho by deamidation of
glutamine. *Nature* **387**, 729–733 (1997).
24. Totsukawa, G. *et al*. Distinct roles of ROCK (Rho-kinase) and MLCK in spatial
regulation of MLC phosphorylation for assembly of stress fibers and focal adhesions
in 3T3 fibroblasts. *J Cell Biol* **150**, 797–806 (2000).
25. Shi, Y., Ohler, L. S., de Jesus, B. N. & Dickinson, D. J. Opposing actomyosin pools
generate cortical flows that establish epithelial polarity. *Biorxiv* (2024).
26. Matsumura, F., Yamakita, Y. & Yamashiro, S. Myosin Phosphatase-targeting Subunit 1
Controls Chromatid Segregation. *Journal of Biological Chemistry* **286**, 10825–10833
(2011).
27. Kiss, A., Erdődi, F. & Lontay, B. Myosin phosphatase: Unexpected functions of a long-
known enzyme. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* **1866**,
2–15 (2019).
28. Yamamoto, K. *et al*. Optogenetic relaxation of actomyosin contractility uncovers
mechanistic roles of cortical tension during cytokinesis. *Nat Commun* **12**, 7145
(2021).
29. Makishima, M. *et al*. Induction of differentiation of human leukemia cells by
inhibitors of myosin light chain kinase. *FEBS Lett* **287**, 175–177 (1991).
30. Hookway, C. *et al*. A human induced pluripotent stem (hiPS) cell model for the
holistic study of epithelial to mesenchymal transitions (EMTs). *BioRxiv* (2024).
31. Lomakin, A. J. *et al*. The nucleus acts as a ruler tailoring cell responses to spatial
constraints. *Science (1979)* **370**, (2020).
32. de Leeuw, N. F., Budhathoki, R., Russell, L. J., Loerke, D. & Blankenship, J. T. Nuclei as
mechanical bumpers during epithelial remodeling. *Journal of Cell Biology* **223**,
(2024).
33. Wang, W., Miller, J. P., Pannullo, S. C., Reinhart-King, C. A. & Bordeleau, F.
Quantitative assessment of cell contractility using polarized light microscopy. *J*
Biophotonics **11**, (2018).
34. Stronati, E. *et al*. YAP1 regulates the self-organized fate patterning of hESC-derived
gastruloids. *Stem Cell Reports* **17**, 211–220 (2022).
35. Pagliari, S. *et al*. YAP–TEAD1 control of cytoskeleton dynamics and intracellular
tension guides human pluripotent stem cell mesoderm specification. *Cell Death*
Differ **28**, 1193–1207 (2021).
36. Meyer, K., Lammers, N. C., Bugaj, L. J., Garcia, H. G. & Weiner, O. D. Optogenetic
control of YAP reveals a dynamic communication code for stem cell fate and
proliferation. *Nat Commun* **14**, 6929 (2023).

37. Pinheiro, D. & Bellaïche, Y. Mechanical Force-Driven Adherens Junction Remodeling
and Epithelial Dynamics. *Dev Cell* **47**, 3–19 (2018).
38. Koch, A. W., Pokutta, S., Lustig, A. & Engel, J. Calcium Binding and Homoassociation
of E-cadherin Domains. *Biochemistry* **36**, 7697–7705 (1997).
39. Smutny, M. *et al*. Myosin II isoforms identify distinct functional modules that support
integrity of the epithelial zonula adherens. *Nat Cell Biol* **12**, 696–702 (2010).
40. le Duc, Q. *et al*. Vinculin potentiates E-cadherin mechanosensing and is recruited to
actin-anchored sites within adherens junctions in a myosin II–dependent manner.
Journal of Cell Biology **189**, 1107–1115 (2010).
41. Bedzhov, I. *et al*. Adhesion, but not a specific cadherin code, is indispensable for ES
cell and induced pluripotency. *Stem Cell Res* **11**, 1250–1263 (2013).
42. Röper, J.-C. *et al*. The major β-catenin/E-cadherin junctional binding site is a primary
molecular mechano-transductor of differentiation in vivo. *Elife* **7**, (2018).
43. Funa, N. S. *et al*. β-Catenin Regulates Primitive Streak Induction through
Collaborative Interactions with SMAD2/SMAD3 and OCT4. *Cell Stem Cell* **16**, 639–
652 (2015).
44. Zimmerli, D. *et al*. TBX3 acts as tissue-specific component of the Wnt/β-catenin
transcriptional complex. *Elife* **9**, (2020).
45. Mukherjee, S., Luedeke, D. M., McCoy, L., Iwafuchi, M. & Zorn, A. M. SOX
transcription factors direct TCF-independent WNT/β-catenin responsive
transcription to govern cell fate in human pluripotent stem cells. *Cell Rep* **40**, 111247
(2022).
46. Moreira, S. *et al*. A Single TCF Transcription Factor, Regardless of Its Activation
Capacity, Is Sufficient for Effective Trilineage Differentiation of ESCs. *Cell Rep* **20**,
2424–2438 (2017).
47. Cadigan, K. M. & Waterman, M. L. TCF/LEFs and Wnt Signaling in the Nucleus. *Cold
Spring Harb Perspect Biol* **4**, a007906–a007906 (2012).
48. Gerner-Mauro, K. N., Akiyama, H. & Chen, J. Redundant and additive functions of the
four Lef/Tcf transcription factors in lung epithelial progenitors. *Proceedings of the
National Academy of Sciences* **117**, 12182–12191 (2020).
49. Guo, Q. *et al*. A β-catenin-driven switch in TCF/LEF transcription factor binding to
DNA target sites promotes commitment of mammalian nephron progenitor cells.
Elife **10**, (2021).
50. Blassberg, R. *et al*. Sox2 levels regulate the chromatin occupancy of WNT mediators
in epiblast progenitors responsible for vertebrate body formation. *Nat Cell Biol* **24**,
633–644 (2022).
51. Yi, F., Pereira, L. & Merrill, B. J. Tcf3 Functions as a Steady-State Limiter of
Transcriptional Programs of Mouse Embryonic Stem Cell Self-Renewal. *Stem Cells*
26, 1951–1960 (2008).
52. Pereira, L., Yi, F. & Merrill, B. J. Repression of Nanog Gene Transcription by Tcf3 Limits
Embryonic Stem Cell Self-Renewal. *Mol Cell Biol* **26**, 7479–7491 (2006).
53. Wray, J. *et al*. Inhibition of glycogen synthase kinase-3 alleviates Tcf3 repression of
the pluripotency network and increases embryonic stem cell resistance to
differentiation. *Nat Cell Biol* **13**, 838–845 (2011).
54. Cole, M. F., Johnstone, S. E., Newman, J. J., Kagey, M. H. & Young, R. A. Tcf3 is an
integral component of the core regulatory circuitry of embryonic stem cells. *Genes
Dev* **22**, 746–755 (2008).

55. Brunet, T. *et al*. Evolutionary conservation of early mesoderm specification by
mechanotransduction in Bilateria. *Nat Commun* **4**, 2821 (2013).
56. Vasic, I. *et al*. Loss of TJP1 disrupts gastrulation patterning and increases
differentiation toward the germ cell lineage in human pluripotent stem cells. *Dev Cell*
58, 1477-1488.e5 (2023).
57. Maytum, A. *et al*. Chromatin priming elements direct tissue-specific gene activity
before hematopoietic specification. *Life Sci Alliance* **7**, e202302363 (2024).
58. Le, H. Q. *et al*. Mechanical regulation of transcription controls Polycomb-mediated
gene silencing during lineage commitment. *Nat Cell Biol* **18**, 864–875 (2016).
59. Roura, S., Miravet, S., Piedra, J., de Herreros, A. G. & Duñach, M. Regulation of E-
cadherin/Catenin Association by Tyrosine Phosphorylation. *Journal of Biological*
Chemistry **274**, 36734–36740 (1999).
60. Kashkooli, L., Rozema, D., Espejo-Ramirez, L., Lasko, P. & Fagotto, F. Ectoderm to
mesoderm transition by down-regulation of actomyosin contractility. *PLoS Biol* **19**,
e3001060 (2021).
1226

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A **Experimental design**

WT → 3h mTeSR1 + Veh or Y-27632 or CN03 → Fix & Stain Imaging

B mTeSR1

+ Veh   + CN03   + Y27

ppMLC2

β-catenin

50 μm

C

Junctional β-catenin (% Area) vs ppMLC2 (% Area)

+ Veh, + CN03, + Y27; ● Repeat 1, ■ Repeat 2, ▲ Repeat 3

R² = 0.9592

D **Experimental design**

EGFP-CTNNB1 knock-in → 3h or O/N mTeSR1 + Veh or Y-27632 or CN03 → Imaging

E mTeSR1

+ Veh   + CN03   + Y-27

EGFP-β-catenin KI

50um

F

Junctional β-catenin (% Area)

** p=0.0072
* p=0.0129

Veh, CN03, Y-27

Figure 6

Figure 7

Supplementary Figure 1 - Related to Figure 1

Supplementary Figure 2 - Related to Figure 3

Supplementary Figure 3

A

CHIR + DMSO

| 00:00 | 08:15 | 14:35 |

CHIR + Q-VD-OPH

| 00:00 | 07:25 | 14:35 |

Brightfield

20 μm

20 μm

B

F-Actin / Nuclei

0h CHIR

24h CHIR

48h CHIR

10 μm

10 μm

10 μm

C

Nuclei   F-Actin

| 0h CHIR | 24h CHIR | 48h CHIR |

10 μm   10 μm   10 μm

D

Monolayer height (μm)

**
p=0.0014
*
p=0.0133

0h   24h   48h
CHIR

E

Nuclei

| 0h CHIR | 48h CHIR |

20 μm   20 μm

F

Average nuclear size (μm²)

*
p=0.0258

0h   48h
CHIR

G

Area (μm²)

0h CHIR R²=0.6876
48h CHIR

R²=0.04714

Count

H

CHIR treatment

| 0h | 24h | 48h |

ppMLC2

50 μm

Pixel intensity

I

ppMLC⁺ area (%)

**
p=0.0054
**
p=0.0026

0h   24h   48h
CHIR

J

Retardance (nm)

*
p=0.0447
****
p<0.0001
****
p<0.0001
***
p=0.0009

0h   24h   48h   72h   Y-27
CHIR

Supplementary Figure 4

Supplementary Figure 5 - Related to Figure 4

Supplementary Figure 6 - Related to Figure 5

A

4 hrs | 16 hrs
CN03 | CN03
H20 | 1 µg/mL | 4 µg/mL | H20 | 1 µg/mL | 4 µg/mL | M.W. (kDa)

ppMLC2 — 20
MLC2 — 20
Nanog — 37
Oct4 — 37
α-Tubulin — 50

B

+ Veh   + CN03   + Y27

Junctional β-catenin (% Area) vs ppMLC2 (% Area)

C

mTeSR1

+ Veh | + CN03 | + Y-27

EGFP-β-catenin KI

0h

14h

20 µm

2000 / 1500 / 1000 / 500 / 0

D <u>Experimental design</u>

Pre-treatment | Differentiation

Veh or Y-27 or CN03 | CHIR + Veh or Y-27 or CN03

-16h | 0h | 24h

E

0h CHIR | 24h CHIR
Veh | Y-27 | CN03 | Veh | Y-27 | CN03 | M.W. (kDa)

Total β-catenin — 100
T/Bra — 50
Tubulin — 50

F

Normalized βcatenin expression

n.s. p=0.8017
n.s. p=0.7314

CHIR 0h 24h | 0h 24h | 0h 24h
Veh | Y-27632 | CN03