Methylomes reveal recent evolutionary changes in populations of two plant species

Kevin Korfmann[ab1], Andreas Zauchner[a1], Bing Huo[a], Corinna Grünke[a], Yitong Wang[a], Aurélien Tellier[at2], Ramesh Arunkumar [at2]

[a]Section of Population Genetics, School of Life Sciences, Technical University of Munich, Liesel-Beckmann Strasse 2, 85354 Freising, Germany

[b]Present address: Department of Biology 77 Klamath Hall 1210 University of Oregon Eugene, OR 97403

[1] These authors contributed equally

[2] These authors contributed equally

[†] Authors for correspondence (Aurélien Tellier - aurelien.tellier@tum.de, Ramesh Arunkumar – arunkumar.ramesh@tum.de)

Abstract

Plant DNA methylation changes occur hundreds to thousands of times faster than DNA mutations and can be transmitted transgenerationally, making them useful for studying population-scale patterns in clonal or selfing species. However, a state-of-the-art approach to use them for inferring population genetic processes and demographic histories is lacking. To address this, we compare evolutionary signatures extracted from CG methylomes and genomes in *Arabidopsis thaliana* and *Brachypodium distachyon*. While methylation variants (SMPs) are less effective than genetic variants (SNPs) for identifying population differentiation in *A. thaliana*, they can classify phenotypically divergent *B. distachyon* subgroups that are otherwise genetically undistinguishable. The site frequency spectra generated using methylation sites from varied genomic locations and evolutionary conservation exhibit similar shapes, indicating minimal noise when all CG sites are analyzed jointly. Nucleotide diversity is three orders of magnitude higher for methylation variants than for genetic variants in both species, driven by the higher epimutation rate. Correlations between SNPs and SMPs in nucleotide diversity and allele frequencies at gene exons are weak or absent in *A. thaliana*, possibly because the two sources of variation reflect evolutionary forces acting at different timescales. Linkage disequilibrium quickly decays within 250 bp for methylation variants in both plant species. Finally, we developed a deep learning-based demographic inference approach. We identified recent population expansions in *A. thaliana* and *B. distachyon* using methylation variants that were not identified when using genetic variants. Our study demonstrates the unique evolutionary insights methylomes provide that genetic variation alone cannot reveal.

Keywords: Methylation site frequency spectra, demographic inference, deep learning, Tajima's *D*, linkage disequilibrium, nucleotide diversity, clocklike regions, gene body methylation, *Arabidopsis thaliana*, *Brachypodium distachyon*

Background

Understanding how variation is partitioned within and between populations is a central goal in evolutionary biology. Increasing evidence suggests that epigenetic modifications can vary within and between populations [1]. Nevertheless, our knowledge of how epigenomes evolve through neutral and selective forces and, consequently, their capacity to reveal population demographic histories is limited. DNA cytosine methylation, a conserved epigenetic modification in eukaryotes [2], occurs in plants within three sequence contexts: CG, CHG, and CHH (H represents A, T, or C) [3]. DNA methylation is known to be involved in regulating gene expression, suppressing transposable element activity, and stabilizing the genome [3]. Long-term changes in the methylome are mainly shaped by genomic alterations such as the proliferation of transposable elements and mutations affecting DNA methylation pathways, whereas heritable stochastic epimutations that occur independent of genomic background drive short-term changes [4].

The unique characteristics of epimutations make them appealing markers for evolutionary investigations. In plants, CG methylation is maintained during cell division by METHYLTRANSFERASE 1 (MET1) [3]. Spontaneous stochastic loss or gain of methylation at CG sites can be transmitted stably across multiple generations [5]. Van der Graaf *et al*. have

estimated the forward (gain of DNA methylation) and backward (loss of methylation) epimutation rates per CG site in the model plant *Arabidopsis thaliana* to be approximately 2.56×10^{-4} and 6.3×10^{-4}, respectively, per generation per haploid methylome [6]. These rates are several orders of magnitude higher than the mutation rate of 7×10^{-9} base substitutions per site per generation in the same population [7]. The increased rates of change at the epigenetic level may facilitate evolutionary inference over recent timescales, where the utility of slower-evolving genetic markers is limited [8, 9].

As transiently induced epimutations can be mistaken for stably inherited markers, a recent study exposed genetically identical *A. thaliana* to various environmental conditions to identify genomic regions—termed clocklike regions—where CG divergence rates are independent of environmental stimuli [10]. While these regions are valuable for evolutionary studies, they are challenging to identify in most species. Therefore, the same study observed that gene body methylated regions (gbM) exhibit a pattern similar to that of CG divergence as the clocklike regions [10]. gbM refers to a subclass of genes that display elevated CG methylation. They are mostly found in moderately and constitutively expressed housekeeping genes [11]. As they can be identified based on the distribution of methylation in the gene bodies, their use may facilitate the investigation of evolutionary processes in nonmodel plant species.

Previous attempts to explore how methylation variants evolve have either neglected demographic (neutral) models, mixed environmentally sensitive and insensitive methylation sites, or did not adequately assess the interdependence of genetic and methylation variants. Charlesworth and Jain's analytic framework for hypermutable sites [12] has been applied to create a methylation site frequency spectrum (mSFS) [4]. Previous studies generated mSFS for genic CG sites and the methylation states of gbM genes, finding either no selection or only weak selection acting on the methylome in *A. thaliana* [4, 13] and maize [14]. However, these studies do not incorporate an explicit demographic model while assessing selection. Another study found that positive selection acted on methylation variants linked to metabolite regulation in *A. thaliana*, even after accounting for demographic histories [15]. Other studies in *Brachypodium distachyon* [16], *Fragaria vesca* [17, 18], lettuce [19], and grapevines [20] have compared evolutionary inferences using genetic and methylation variants by analyzing all CG sites as the environmental sensitivity of individual sites is unknown in most plant species. Recently, Sellinger *et al*. utilized methylation as a secondary marker alongside genetic variants to enhance demographic inference over short and long timescales [8]. The conceptual novelty was to consider that the underlying genealogy at a locus (*i.e.* the coalescent tree tracing back the ancestry of the sample back in time) is common for SNPs and SMPs (under a neutral model of evolution). This represents the first population genomics inference model combining DNA mutations, epimutations, and recombination along the genome, and the basis for developing a Sequential Markovian Coalescent method (SMCm). However, that study exhibits a fair amount of noise in the inference of past evolutionary history when it was applied to methylome data from *A. thaliana* [8]. These inaccuracies are due to a lack of knowledge about the optimal SMP markers to use, the relevant genomic context, the relationship between SNPs and SMPs, and the SMCm method being very sensitive and not accounting well for methylated region effects (so-called differentially methylated regions or DMRs). A related question is whether DMRs can be used as well for evolutionary inferences.

Given these limitations, our goal was to assess how well methylation variants reveal known evolutionary patterns inferred using genetic variants. We chose *A. thaliana* and *B. distachyon*

for our analyses because their evolutionary histories are well-described [16, 21]. We found that the capacity of methylation variants to differentiate populations is contingent on when populations diverge. We also show that CG sites across the methylome capture within- and between-population patterns comparable to environmentally insensitive sites (clocklike regions and gbM) [10] in samples grown in a common garden. We found weak or no local correlations between genetic and methylation variants, indicating they reflect different aspects of population evolutionary histories. Additionally, the linkage between neighboring methylation variants was minimal. Conventional genome-based demographic inference methods, such as the Sequential Markovian Coalescent [22], show limitations in using methylation variants due to high sensitivity to violations of the Markovian properties of the underlying coalescent model along the genome [8]. Moreover, since such methods require knowing the variant accumulation rates and distributions—often unknown for methylation in many species—we developed a deep learning method that uses methylation as a neutral marker to estimate past demography. Unlike conventional convolutional networks, we use (to our knowledge) the first Encoder-Only Transformer architecture [23-25] method, focusing on the placement of mutations on a hidden ancestral recombination graph. We track estimates of coalescent events in a time-discretized transition matrix by using heterogeneous sites per genomic window as input, thereby applying a coarse-grained approach, similar to tsABC [22]. By aligning training and real data distributions, we show that rapidly changing methylation variants can reveal recent demographic shifts in selfing and clonal species.

Results

CG methylation variants have variable performance in identifying population structure

We analyzed genetic and methylation diversity using 63 samples each from the Central European (CEU) and the Iberian non-relicts (IBnr) *A. thaliana* populations (Table S1). These populations were used by Strütt *et al*. [22]. Across the two populations, 31-33% of CG methylation sites in clocklike regions were polymorphic (Figure S1A), which is similar to a previous estimate for CG sites in *A. thaliana* mutation accumulation lines [26]. In contrast, 1.6-1.7% of coding sites and 2.7-2.8% of non-coding sites were polymorphic (Figure S1A). We compared our results from *A. thaliana* with those from 29 samples from the clonally propagating grass *B. distachyon*. Wilson *et al*. used SNPs to identify clonal groups among Turkish *B. distachyon*, with samples within each group showing high levels of genetic similarity [27]. Eichten *et al*. identified high levels of consistency between the known genetic relationships for the clonal groups and the extent of CG divergence [16]. Here, we restrict our focus to SMPs and DMRs occurring within gbM regions and two clonal groups of *B. distachyon* with average bisulfite sequencing depths greater than 10 [16] (Group 1: 22 samples, Group 6: 7 samples) (Table S1). Between 4-7% of CG methylation sites within gbM regions were polymorphic in *B. distachyon* (Figure S1B). The median length of DMRs in *B. distachyon* was 107 bp (1st percentile: 44 bp, 99% percentile: 717 bp) which is smaller than the median length in *A. thaliana* DMRs, which was 181 bp (1st percentile: 78 bp, 99% percentile: 1125 bp) (Figure S2).

Figure 1. Methylomes outperform genetic variants in detecting population subdivision in *Brachypodium distachyon* but not *Arabidopsis thaliana*. (**A**) Neighbor-joining trees were generated using p-distance matrices for non-coding single nucleotide polymorphisms (SNP), coding SNPs, clocklike region single methylation polymorphisms (SMP), and clocklike region differentially methylated regions (DMR) for Central European (CEU) and the Iberian non-relicts (IBnr) *A. thaliana*. (**B**) Neighbor-joining trees generated using p-distance matrices estimated for SMP and DMR in gbM regions of Turkish *Brachypodium distachyon* clonal groups.

We estimated the proportion of sites at which two samples were different (p-distance) and clustering to uncover how the net total of genetic and methylation variation differed between samples from the CEU and IBnr populations. In this approach, a clear separation between the CEU and IBnr populations was apparent with both coding and non-coding single nucleotide polymorphisms (SNP) (Figure 1A). Although single methylation polymorphisms (SMP) and differentially methylated regions (DMR) largely separated the two populations, some samples were separated by only minimal distances (Figure 1A). Clustering indicated nine or more

208 groups within the SNP data (Figure S3A-B), whereas only two groups were apparent in the
209 SMP and DMR datasets (Figure S3C-D).
210
Furthermore, we performed discriminant analysis of principal components (DAPC) analyses
to identify how correlated changes in subsets of genetic and methylation variants can be used
to discriminate the two populations. DAPC indicated greater discriminatory power with coding
and non-coding SNPs compared to SMPs and DMRs (Figure S3E). Although p-distance
matrix-based comparisons could have been influenced by the greater number of SNPs (Figure
1A), marker numbers were comparable for the DAPC, and yet SMPs and DMRs showed poor
discriminatory capability (Figure S3E). SMPs and DMRs that occur within gbM genes have
comparable performance to those occurring within clocklike regions in discriminating the two
populations (Figure S4). High methylation rates in *A. thaliana* may have caused multiple
changes at the same site (so-called homoplasy accounted for as a finite site model [4, 12,
28]), resulting in the underestimation of the true levels of divergence. Indeed, the divergence
between IBnr and CEU populations is estimated to have occurred after the last glacial
maximum ca. 10,000-20,000 years ago [29-31].
224
We then performed a similar analysis using the *B. distachyon* accessions. In contrast to the
clonal grouping based on the genetic profiles [27], SMPs and DMRs were capable of
additionally distinguishing the two subgroups of Group 1 (Figures 1B, S5). The trees estimated
using the p-distance matrices were shorter for SMPs compared to DMRs (Figure 1B), possibly
due to increased error in this dataset because of the shallow bisulfite sequencing depths.
However, both trees were longer compared to those in *A. thaliana* (Figure S4A) indicating
greater methylome differentiation in *B. distachyon*. Both our results using gbM regions and the
results from Eichten *et al*. that use the entire collection of CG sites [16] reveal a difference in
methylation profiles between the two subgroups of Group 1. We obtained phenotype data for
the two subgroups [32] and found significant differences in the longitude (Figure S6), elevation
of collection site, seed lengths, widths, yield production, and germination percentages (Figure
S7). Our findings suggest that methylation variants can reveal subpopulation structure in
recent timescales, where there has been little time for genetic differences and multiple
methylation changes to accumulate.
239
240
241 **Genomic contexts have little influence on the shape of SMP distributions**

Figure 2. Methylation site frequency spectra generated using single methylation polymorphisms were similar regardless of genomic contexts. Methylation site frequency spectra (mSFS) generated using differentially methylated regions (DMR) exhibited greater variability. (**A**) mSFS for the Central European (CEU, Salmon) and the Iberian non-relicts (IBnr, Cyan) *Arabidopsis thaliana* populations using CG sites that occur in clocklike, gene body methylated (gbM), non-gbM, and low and high Sorting Intolerant from Tolerant (SIFT) score regions. Only sites with minor allele frequencies (MAF) >0.02 were used to generate the mSFS. (**B**) mSFS for *Brachypodium distachyon* clonal groups 1.1, 1.2, and 6 for single methylation polymorphisms (SMP) and DMRs occurring within gbM and non-gbM regions. Only sites with minor allele frequencies (MAF) >0.04 were used to generate the mSFS. For (A) and (B), the numbers in each panel indicate the number of sites used for the analyses.

We sought to understand how genomic contexts influenced the methylation site frequency
spectra (mSFS) in *A. thaliana*. The mSFS reveals how methylation rates, demographic
changes, and natural selection influence changes in the methylome over time [4, 12]. We
found a greater frequency of unmethylated alleles at most polymorphic methylation sites for
the CEU and IBnr populations, but CEU consistently had higher frequencies of unmethylated
alleles (Figure 2A). The shapes of the mSFS for SMPs are consistent with previous findings
that the backward methylation rate promoting the change from methylated to unmethylated is
higher than the forward rate [6] and are consistent with previous mSFS generated by
combining *A. thaliana* samples from extended geographic regions [4, 13]. The shape of the
mSFS for SMPs in clocklike and gbM regions were similar, supporting their interchangeability.
We also studied the mSFS for intervals containing genes with low and high Sorting Intolerant
From Tolerant (SIFT) scores, which are positively correlated with the level of evolutionary
constraint that genes are under [33]. Our results indicate that the level of gene constraint has
little impact on the mSFS shape (Figure 2A).

When compared to SMPs, the DMR mSFS for *A. thaliana* were more dependent on the
underlying set of intervals used to generate them. For DMRs that occur in intervals containing
non-gbM genes, and high and low SIFT-score genes, more samples have the unmethylated
state reflecting the underlying SMP distributions (Figure 2A). This is also true when DMRs
across the entire genome are analyzed (Figure 2A). In contrast, for DMRs in clocklike regions
and gbM genes, more samples exhibit the methylated state (Figure 2A). The mSFS shapes
for DMRs show greater variability compared to those for SMPs, which may reflect underlying
differences in their molecular basis.

In contrast to the mSFS for SMPs in *A. thaliana* (Figure 2A), the mSFS for all three clonal
groups of *B. distachyon* indicated an excess of methylated alleles within gbM and non-gbM
SMPs, although it was less pronounced for the latter (Figure 2B). There was less consistency
in the shapes of the mSFS for DMRs in gbM and non-gbM regions (Figure 2B), similar to our
observation when *A. thaliana* DMRs were analyzed (Figure 2A).

Methylation and genetic variants show unique evolutionary signatures in *A. thaliana*

Figure 3. Unique evolutionary signatures in genomes and methylomes. (**A**) Estimates of
pairwise nucleotide diversity (π) scaled by the number of cytosines per region for single
methylation polymorphisms (π_{SMP}) and the number of bases for single nucleotide
polymorphisms (π_{SNP}) for the Central European (CEU) and the Iberian non-relicts (IBnr)
Arabidopsis thaliana populations. (**B**) Estimates of Tajima's *D* per region for CEU and IBnr.
Comparisons of (**C**) π_{SNP} versus π_{SMP} and (**D**) Tajima's *D* for SNPs (D_{SNP}) and SMPs (D_{SMP})
per gene for CEU and IBnr. The numbers in the titles of each panel indicate the number of
sites used for the analyses. Correlation tests were conducted to assess the associations
between π and Tajima's *D*. (**E**) π_{SMP} scaled by the number of cytosines per region for
Brachypodium distachyon clonal groups 1.1, 1.2, and 6. (**F**) D_{SMP} per region for *B. distachyon*.
For (A), (B), (E), and (F), the numbers above the boxplots indicate the number of intervals.

300 Letters indicate group-wise differences in means following Tukey's honest significant
301 difference tests. For (F), the color scheme and number of sites are the same as in (E).
302
303
We next sought to understand how genetic and methylation polymorphisms are distributed
within *A. thaliana* populations. We first calculated the pairwise nucleotide diversity (π) for
SMPs (π_{SMP}) and SNPs (π_{SNP}) from noncoding, clocklike, gbM, non-gbM, and high and low
SIFT genic regions. We scaled estimates by the total number of sites (polymorphic and
monomorphic) to ensure that the results were comparable (that is per site estimates). π_{SMP} is
three orders of magnitude higher than π_{SNP} (Figure 3A), likely driven by the high rates of
change in methylomes [6]. π_{SNP} in the CEU population and π_{SMP} in CEU and IBnr populations
for clocklike regions were significantly higher than in other regions (Figure 3A), possibly
indicating better detection of environmental insensitivity with more polymorphic sites. π_{SMP} was
higher for gbM and high-SIFT genes than at non-gbM and low-SIFT, respectively, but no such
difference was seen for π_{SNP} (Figure 3A), indicating that constraint at the genic level has a
minor effect on the levels of local methylation diversity.
316
We then compared how selective and demographic processes influence genomes and
methylomes by estimating Tajima's D for CEU and IBnr. For methylomes, we used a modified
Tajima's D using the D^m [28] that accounts for the hypermutability of the methylation sites.
D_{SNP} was slightly negative for noncoding sites in the CEU and IBnr populations (Figure 3B),
reflecting the known history of population expansions [22]. The higher D_{SNP} for IBnr (Figure
3B) is consistent with larger effective population sizes over recent timescales [22]. In contrast,
IBnr had significantly lower D_{SMP} compared to CEU (Figure 3B). This could be due to variation
in methylation rates, purifying selection, or demographic and selective (purifying selection)
histories. Therefore, we used SMCm [8], which employs fixed mutation rates and compares
the distributions of mutation and methylation markers, to estimate the forward and backward
methylation rates for the IBnr and CEU populations. Methylation rates were only slightly higher
in CEU but the ratio of backward to forward rates was similar in both populations (Figure S8).
Hypermutability should result in greater frequencies of intermediate variants [12], but we did
not observe such a pattern (Figure 2A). Tajima's D is based on a comparison of the number
of segregating sites versus π. The disparity in D_{SMP} appears to be driven primarily by the
variation in the number of segregating sites per region (Figure S9). This difference suggests
that the demographic histories of the two populations may be slightly different.
334
We then estimated Tajima's D in different genomic contexts for CEU and IBnr. Clocklike
regions had significantly higher D_{SNP} and D_{SMP} compared to other regions (Figure 3B). The D^m
[28] method assumes that forward and backward methylation rates are equivalent. Although
the ratio of the backward to forward methylation rates in clocklike regions is 1.36, the ratios
are 2.11 and 2.84 for gbM and non-clocklike regions, respectively [10], indicating the
assumption of D^m is less strongly violated for clocklike regions. D_{SMP} for non-gbM regions was
slightly but significantly higher compared to the estimates for gbM regions (Figure 3B),
suggesting that ignoring the differences between regions can introduce minimal noise when
inferring evolutionary signatures from methylomes. The differences in D_{SMP} between high-SIFT
vs low-SIFT were not consistent in both populations (Figure 3B). There was only a weak
positive correlation for π (Figure 3C) generated for SNPs vs SMPs that occur in coding regions
and no significant correlation for Tajima's D (Figure 3D).
347

We compared π and Tajima's *D* for *B. distachyon* SMPs generated using SNPs for the
Anatolian and Turkish lineage (B_East) [34, 35] to which all three clonal groups belong. Scaled
π for SMPs was larger for the *B. distachyon* clonal groups compared to the *A. thaliana*
populations, but there were no consistent differences between gbM and non-gbM regions or
among clonal groups (Figure 3A, 3E). Similar to *A. thaliana* (Figure 3A), scaled π for SMPs in
all three clonal groups was three orders of magnitude larger than the median SNP-based
scaled π (~$1.8×10^{-6}$) for the B_East lineage [34, 35] (Figure 3E). Similar to *A. thaliana* (Figure
3B), D_{SMP} in the three clonal groups was negative. D_{SMP} was slightly but significantly larger in
non-gbM regions compared to gbM regions (Figure 3F). These negative values contrast with
the positive D_{SNP} for genetic variants in the B_East lineage [34]. Unlike in *A. thaliana* (Figure
S9), this contrast cannot be attributed primarily to the differences in the number of segregating
sites (Figure S10), meaning that it cannot be explained solely by the extent of genotyping.
D_{SMP} was lower for Group 1.1 compared to the other two groups (Figure 3F), which is
consistent with the higher frequency of rare alleles in this group (Figure 2B).
362
363
364   **Linkage disequilibrium quickly erodes for methylation variants**
365

366
Figure 4. Weak linkage disequilibrium (LD) between neighboring single methylation
polymorphisms (SMP). (**A**) Pairwise r^2 estimated using coding single nucleotide
polymorphisms (SNP), non-coding SNPs, and SMPs for Central European (CEU) and Iberian
non-relicts (IBnr) *Arabidopsis thaliana* populations. The number of sites used for comparisons
is indicated within the plots. (**B**) Pairwise r^2 was estimated using SMPs for the clonal groups.
For (A) and (B), the number of sites used for comparisons is indicated within the plots.
373
374
An important consideration for population genetic analysis is the presence of sites with
independent evolutionary histories [8]. Therefore, we assessed the levels of pairwise linkage
disequilibrium (LD) to identify the number of independent polymorphic SNPs and SMPs.
Pairwise LD decay was stronger for noncoding SNPs compared to coding SNPs in *A. thaliana*
(Figure 4A). However, LD decay was much stronger for SMPs than for SNPs with r^2 dropping
below 0.1 within the first 250 bp (Figure 4A). LD decays to 0.1-0.25 for SMPs sites within gbM
regions for *B. distachyon* within the first 100 bp but remains constant thereafter (Figure 4B)
and is stronger compared to LD in *A. thaliana* over large distances (Figure 4A).
383
384
385   **Demographic inference using genetic and methylation variants**

A)

1. Count number of heterozygous markers per window

Compute for pairwise comparisons of samples

2. Multiply scaling factor (s) and number of markers to approximate average TMRCA estimate per window

3. Counting transitions from window to window to approximate transition matrix

Scaled number of makers (w_1, w_2, w_3, …)

4. Predict demography from approximated transition matrix

Model: Encoder-only Transformer
Input: Approximated Transition Matrix
Output: Population size at n time points

387
Figure 5. Demographic inference using mutation or methylation markers. **(A)** Schematic of
the approximated transition matrix inference framework. The genome is first divided into 20
Kbp bins, and heterozygous markers are counted for all pairwise comparisons in a sample of

10 individuals. These counts are then scaled by an empirically chosen factor to ensure
sufficient values in the approximated transition matrix by aligning the number of markers per
window to the TMRCA in that window. The framework constructs this matrix by concatenating
all pairwise comparisons into a long sequence and applying a sliding window of size 2 and
step size 1. Each pair of consecutive values in this window represents coordinates in the
transition matrix, which is incremented at those positions. This process results in an
unnormalized transition matrix that captures the genetic relationships between samples.
Finally, an encoder-only transformer neural network takes this noisy matrix as input, implicitly
learning the fixed scaling factor, and predicts population sizes at various time points. Inference
using over 42 time steps using (**B**) single nucleotide polymorphisms (SNP) and (**C**) single
methylation polymorphisms (SMP) for the central European (CEU) and the Iberian non-relicts
(IBnr) *Arabidopsis thaliana* populations. (**D**) Demographic inference using SMP for Groups 1.1
and 1.2 of *Brachypodium distachyon*. SMP inference was done using sequences from all five
chromosomes (light lines, average in bold), while SNP inference is based on chromosome
one. (**E**) Inference of simulated sawtooth demography using five replicates, 4Mbp segments,
and 10 samples.
407
408
We developed a novel inference method that focuses on large genome blocks (20 Kbp
windows) likely spanning multiple coalescent genealogies (Figure 5A), to estimate the times
to the most recent common ancestor (TMRCA). Our method assumes that the number of
heterozygous markers per window is correlated with the average TMRCA. Essentially, longer
coalescent times allow more mutations and epimutations to accumulate. To mitigate the
possibility that older genealogies (long TMRCA) can be unexpectedly short due to immediate
recombination, our approach uses 20 Kbp windows to average out genealogies. We apply a
scaling to the number of markers to estimate the TMRCA (Figure S11). Although the chosen
scaling factor will not perfectly offset all variable demographic samples—given that we trained
using a fixed recombination rate and a fixed window size—offsets are minimal and addressed
during training. An encoder-only transformer neural network processes this data, learns the
scaling factor, and predicts population sizes over time. This is done by summarizing the
TMRCA of two consecutive windows in the empirical transition matrix. The differences
between each pair of samples are used to infer their TMRCA. The TMRCA values for all pairs
across multiple intervals (windows) in the genome are then represented as a transition matrix.
This method combines raw genotype data restructuring with deep learning to infer
demographic history.
426
For deep learning inference, the distributions of heterozygous markers in training and real data
must overlap to ensure in-distribution inference. For the training data, we collapsed each
training transition matrix into a single scalar by summing the flattened matrix and plotted the
distribution of unnormalized transition matrices (Figure S12A). To ensure the real data *from A.*
thaliana and *B. distachyon* matched the training data, we analyzed 20 randomly sampled
genomic regions. For the SNP data, we selected start positions uniformly between 0-28 Mbp,
whereas for the methylation data, the start positions were chosen uniformly between 0-0.5
Mbp. For both data types, we determined each region's length by sampling from a uniform
distribution between 1-3 Mbp. Summing the resulting transition matrices for the IBnr, CEU,
Group 1.1, and Group 1.2 regions (Figure S12B-D) produced values within the training data
range (Figure S12A). Despite the differential properties of SMPs and SNPs, the distribution of

heterozygous markers per 20 Kbp window was similar across populations (Figure S12B-C),
supporting the use of a common training dataset for deep learning.
440
We applied our coarse−grained inference approach to *A. thaliana* and *B. distachyon*. We
generated approximated transition matrices from 10 samples and 4 Mbp sequence lengths.
The 45 pairwise comparisons totaling 180 Mbp with 20 Kbp windows yielded 9,000 pairs of
TMRCA values per matrix. We detected a population bottleneck ~100,000 years ago (100 kya)
in CEU and IBnr using SNPs (Figure 5B), which was consistent with the findings of a previous
study [22]. This signal was not clearly detected with SMPs (Figure 5C), indicating their limited
use for ancient events. Both SNPs and SMPs indicated another bottleneck between 1-10 kya
(Figure 5B-C), but SMPs also revealed a subsequent population expansion (Figure 5C)
undetected by SNPs (Figure 5B). The population contraction began earlier in IBnr, followed
by a slightly stronger expansion compared to CEU (Figure 5C), which may explain its more
negative Tajima's *D* estimate for methylation variants (Figure 3B). Using SMPs, we recovered
two bottlenecks (~100 kya and between 1-10 kya) in *B. distachyon* (Figure 5D) that were
previously reported using SNPs for samples combined from multiple clonal groups in this
species [34]. The steeper expansion of IBnr compared to CEU (*A. thaliana*) and Group 1.1
compared to Group 1.2 (*B. distachyon*) in the last 10,000 years likely led to the observed
excess of rare alleles (Figures 2B, 3F).
457
We validated our demographic inference with two approaches. First, we simulated the
sawtooth demography observed in both species, confirming that our deep learning method
accurately detected multiple bottleneck and expansion events (Figure 5E). Second, we
simulated methylation markers with high rates of change (10^{-4}, 10^{-5}, or 10^{-6} per site per
generation) and backward rates 2.5 times larger than forward rates, based on the estimates
provided in [6], to test whether deviations from the infinite sites model affect inference. We
found that our method accurately recovered demographic histories beyond 0.6 kya and before
5, 22, and 442 kya for methylation rates of 10^{-4}, 10^{-5}, and 10^{-6} per site per generation,
respectively (Figure S13). Conservatively, this suggests that inferences 20 kya and beyond
are unlikely to be accurate for CEU and IBnr due to their estimated rates of methylation change
being around 10^{-5} per site per generation (Figure S8). Note that the demographic estimates
with SNPs are likely accurate for older events (>20kya). Nonetheless, we are confidently
estimating recent population expansion with SMPs (Figure 5C), highlighting the potential of
methylation markers to reveal insights beyond what genetic variants alone can provide.
472
473
474 **Discussion**
475
We observed contrasting patterns in the capacity of methylation variants to distinguish
population subdivisions in *A. thaliana* and *B. distachyon*. In *A. thaliana*, the infinite sites model
is likely violated for methylation, resulting in the erosion of methylome differentiation likely due
to the older divergence between populations. In *B. distachyon*, samples from different altitudes
with varying seed morphologies exhibited distinct methylation profiles in a common garden,
despite their genetic similarity. While SMP distributions are similar across different genomic
contexts, DMRs are more context dependent, making the former preferable for population-
scale analyses. We found weak negative correlations between π and no local correlations for
Tajima's *D* between genetic variants and methylation variants in *A. thaliana*, underscoring their
distinct molecular and evolutionary properties, similar to the conclusions reached by Schmitz

et al. (2013) [36]. We found slightly stronger linkage between neighboring SMPs compared to a previous estimate in wild *A. thaliana* sampled across the Northern Hemisphere [36], likely due to our focus on individual populations with less genetic structure [22]. SMPs and SNPs recover recent and older demographic changes, respectively, providing a comprehensive reconstruction of population evolutionary histories.

The differing ability to detect population subdivisions in *A. thaliana* and *B. distachyon* using methylation variants could lead to differences in divergence times, methylation rates, or local adaptation. As the post-glacial re-colonization of the Iberian Peninsula and the spread northward started ~16 kya [29], multiple changes at the same methylation site (homoplasy) during this period may have caused an underestimation of the true levels of divergence in *A. thaliana*. The greater capacity to detect group subdivision is *B. distachyon* could reflect a recent separation by altitude contributing to methylome divergence. We indeed rule out the role of adaptation such as found in *F. vesca* [17], because we generated inferences using gbM regions that are highly conserved in angiosperms [11]. The observed divergence could be due to low rates of methylation change, but we could not estimate methylation rates for *B. distachyon* as our rate estimate relies on comparing the distributions of genetic and methylation variants. The most parsimonious explanation is that the two subgroups diverged recently. As the B_East lineage split from the western Mediterranean lineage 23 kya [34, 35], the split between the closely related subgroups is likely to have been far more recent, ensuring that the infinite sites model was not violated for methylation variants. Our findings underscore the necessity for methods distinguishing neutral demographic changes from selective pressures on the methylome, elucidating why SMPs behave differently across species.

Our results support the use of individual CG sites from all genomic contexts for population-scale analyses. Our between-population comparisons using CG sites in gbM regions successfully identified the known population structure within *B. distachyon*, similar to a previous study using genic CG sites [16]. Further supporting this conclusion is the consistent similarity in the mSFS shape when using SMPs, regardless of their genomic context. The *A. thaliana* [21] and *B. distachyon* [16] methylomes, generated under similar environmental conditions, ensure that sites affected solely by transient environmental effects would not be recognized as SMPs. Another study using all CG sites from lettuce methylomes, cultivated under uniform environmental conditions, also demonstrated the capacity to distinguish multiple species [19]. Some noise can be introduced by analyzing the whole methylome as non-gbM regions show a slight skew towards variants with intermediate frequencies, but this is likely to be offset by the increased number of sites for inference. The distribution of DMR states is tied to the genomic context. gbM results from the combined activity of MET1, which is enhanced by nearby existing methylation, as well as the histone variant H2A.Z and the DNA demethylase ROS1 [37]. The molecular basis underlying the clock-like, non-gbM regions, as well as regions with high and low SIFT scores, remains unknown. However, given that our mSFS for these regions differ, this suggests potential for differences in their molecular basis. Studying DMRs can nevertheless be valuable for imputing methylation states when sequencing depths are low [38], while the impact of such imputation on population genetics inference needs to be evaluated. Overall, using a restricted set of CG sites may be more appropriate when analyzing accessions grown under variable environments together or for tracking long-term methylome divergence [10].

The weak local correlations between genetic and methylation diversity and the association
between gene constraint and methylation diversity in *A. thaliana* can partly be attributed to
methylation affecting the mutability of neighboring sites [39-41] and transposable element
polymorphisms influencing methylation variation [42, 43]. However, only about 12% of
methylation quantitative trait loci (QTLs) in *A. thaliana* are within 10Kbp of the epiallelic target
[36]. In maize, approximately 50% of methylation QTLs occur in *cis* [44], indicating a potentially
stronger trend in this species. The higher rate of methylation changes compared to mutation
accumulation [6, 7] may also explain why methylation differences persist despite genetic
homogeneity. Our simulations also show that coalescent histories of markers with vastly
different rates of change diverge, which can explain the general lack of correlation between π
and Tajima's *D* for local genetic and methylation diversity. Our results are similar to those of
previous studies in lettuce [19] and cotton [45], which also found higher levels of polymorphism
in the methylome compared to the genome. The lack of correlation or similarity between the
Tajima's *D* estimates from the genome and methylome in *A. thaliana* and *B. distachyon*
supports the notion that the SNPs and SMPs likely reflect different parts of the common
underlying coalescent genealogy at a locus: SNPs reflect the older part of the tree, whereas
SMPs would reflect the topology and branch lengths in the more recent past. It is then not
surprising that for recent positive selection events (selective sweeps), the signatures are
similar for both types of markers, and SMPs may even be more accurate in detecting them
[15]. Our study mostly focuses on local correlations in diversity, rather than genome-wide
correlations, where life history traits and genetic differentiation levels also play a significant
role [46-52].

555
Our study demonstrates that methylation variants can independently detect and allow
inference of historical demographic changes. This represents a significant improvement over
a previous approach [8], where methylation was used only as a secondary marker to
supplement genetic variant-based inferences. Furthermore, our improved deep learning
method does not require precise methylation rate parameters and is less sensitive to the
presence of invariant methylation sites (compared to the SMCm in [8]). Instead, it trains on
diverse scenarios to match observed variant distributions. For *B. distachyon*, methylation
variants revealed expansions also detected by genetic variants [34]. In *A. thaliana*, genetic
variant-based demographic inferences fail to detect recent population expansions (this study
and [53]), which can be uncovered through methylation variants instead. Our simulations
confirmed the utility of rapidly mutating markers for inferring demographic changes. Overall,
our study highlights the value of methylation markers for understanding recent shifts in
population size.

569
570
571   **Conclusions**

572
Our findings demonstrate that CG methylation can provide new insights into population
structure in clonal and selfing species, similar to its previously demonstrated utility in *A.*
thaliana from America and the clonal seagrass *Zostera marina* [10]. We demonstrate the utility
of methylation variants for uncovering very recent demographic events, where SNP markers
are limited due to their slower rate of change. We also illustrate its usefulness in clonal groups,
where the paucity of SNPs would render such inferences either impossible or necessitate very
deep and costly levels of DNA sequencing. Conversely, at older timescales, methylation
markers are unlikely to be informative because the high rates of change would have resulted

in the violation of the infinite sites model. We show that SNPs and SMPs are complementary
for generating demographic inferences over short and long timescales, highlighting the value
of a recently developed method that leverages both sources of variation for inferring
demography [8]. Our work demonstrates that methylomes contain unique evolutionary insights
unattainable through studying genomic variation in isolation.
586
587
588 **Methods**
589
590 *Genetic and methylation variant calling in Arabidopsis thaliana*
591
We obtained the *A. thaliana* 1001 genomes [30] SNP calls from
https://1001genomes.org/data/GMI-MPI/releases/v3.1/. We only retained biallelic SNPs for
our analyses. As C/T variant calls can become confounded with methylation calls for data
generated using bisulfite sequencing, we generated sample-specific genome references for
mapping bisulfite sequencing reads. We substituted the genetic variant calls for each sample
into the TAIR10 genome reference [54] using SAMtools [55], BCFtools consensus [55], and
fastq split [56]. We obtained bisulfite sequencing data for the 1001 *A. thaliana* from [21]. We
first removed poor-quality reads using Trimmomatic [57] with the following options:
ILLUMINACLIP:TruSeq3-SE.fa:2:30:10 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15
MINLEN:36 for single-end reads and ILLUMINACLIP:TruSeq3-PE.fa:2:30:10:2:True
LEADING:3 TRAILING:3 SLIDINGWINDOW:4:20 MINLEN:36 for paired-end reads. We
subsequently mapped reads to the sample-specific reference genomes using Bismark with
the aid of the HISAT2 aligner [58]. We ran Bismark deduplicate_bismark to remove optical
duplicates and ran bismark_methylation_extractor to extract methylation calls. We studied
cytosine sites with a coverage of at least five reads similar to what was done in [21]. To identify
the methylation state of a site, we performed a binomial test with the non-conversion rate being
used as the hypothesized probability of success following the method in [59]. The non-
conversion rate was calculated as the proportion of methylation in the chloroplast as done in
[59]. Sites were assigned as methylated if the false discovery rate corrected *P*-values following
the binomial test was less than 0.01. Otherwise, they were designated as unmethylated. We
used jDMR [60] to identify DMRs in the CG contexts as this method accounts for coverage
when calculating likelihood scores for methylation states. For this, we called the methylation
status per site using METHimpute (function: callMethylationSeparate) [38], extracted cytosine
clusters in the CG context ensuring that there was a minimum of eight cytosines in each cluster
(function: CfromFASTAv4 and makeReg with default options makeRegnull = c(FALSE), N.boot
= 10^5, N.sim.C = "all", fp.rate = 0.01, set.tol = 0.01), and identified the methylation states of
the regions in each sample (function: runMethimputeRegions), ensuring that we used a
minimum of eight cytosines and a coverage of three.
620
63 IBnr and 67 CEU samples overlapped between the SNP and methylation variant call files.
We randomly subsampled 63 samples from the CEU population to ensure the numbers were
comparable to those of the IBnr population. For our analyses, we only used sites where at
least 80% of the samples within each population were genotyped. The proportions of variant
SNP and SMP sites were calculated by dividing the number of sites with minor allele
frequencies greater than 0.02 by the total number of variant and invariant sites.
627
Population (epi)genetic metric estimation for Arabidopsis thaliana

To estimate π and Tajima's *D*, we created fasta files for each genomic region containing the
sequences for each sample. We substituted the variants from the SNP vcf into the TAIR10
reference genome using the approach described previously for each sample, extracted the
region of interest, and converted it to a fasta format. The regions could be the clocklike regions
described in [10], genic regions comprising all exons, or non-coding regions comprising of all
regions other than exons. Genic regions were further split into gbM genes using the list for *A.*
thaliana provided by [61] and high SIFT-score (>0.75) and low SIFT-score (<0.3) genes using
the list provided by [62].

We estimated π_{SNP} and D_{SMP} using pegas [63] and estimated π_{SMP} and D_{SMP} using the D^m [28].
We only analyzed intervals that had at least three polymorphic sites and sequences with
lengths greater than 10bp. We calculated the number of cytosines in each interval using
METHimpute [38]. To compare π and Tajima's *D* among the different genomic contexts, we
built a linear model where these metrics were the dependent variable, and the concatenated
population and genomic context groupings were the predictor variables. We then applied a
Tukey honest significant differences test using the agricolae R package
(https://myaseen208.com/agricolae//index.html) to identify significant differences between
individual predictor levels. We performed a correlation test to compare the π and Tajima's *D*
for SMPs and SNPs per exon.

We used VCFtools [64] to estimate the allele frequencies for SMPs and DMRs. We used
PopLDdecay [65] to estimate pairwise LD using the r^2 metric analyzing all sites with a minor
allele frequency greater than 0.005.

Between population comparisons for Arabidopsis thaliana

We combined the CEU and IBnr vcfs and used VCF2Dis (https://github.com/BGI-
shenzhen/VCF2Dis) to estimate the p-distance matrix. We used this matrix to generate
neighbor-joining trees using FastME [66] and plotted it using ggtree [67]. We also performed
DAPC to discriminate samples from the CEU and IBnr populations. We thinned the dataset so
that only sites that were separated by 1Kbp were sampled to reduce the computational time.
We converted vcf files into a genlight format using the *genomic_converter* function in the R
package radiator (https://thierrygosselin.github.io/radiator/). We used the find.clusters function
in adegenet [68] to identify the number of clusters in each dataset allowing for a maximum of
20 clusters for SMPs/DMRs and 30 clusters for SNPs. As DAPC regions require fully typed
data, we imputed missing values using the mean value. Then, we performed cross-validation
using adegenet [68] to identify the number of informative principal components. We retained
a maximum of 15 principal components, used 90% of the data for training, and calculated
group-wise assignment success. We centered but did not scale variables. We used 30
replicates for each principal component. Finally, we performed DAPC using adegenet [68].

Methylation rate estimation *for Arabidopsis thaliana*

We generated the multihetsep files for SNPs for the same 5Mbp regions that were used for
SMPs (described in the section immediately above). We combined the SNP and SMP
multihetsep files. We randomly selected 20 samples from each population to ensure
reasonable runtimes. We used the *Methylation_rate_estimation* function in SMCm to estimate

forward and backward rates [8]. We set the mutation rate as 7×10^{-9} per site per generation
and the recombination rate as 3.5×10^{-8} per site per generation. We ignored regional influences
on methylation rates and estimated rates under a site-only model. We used the known
estimates of forward (2.56×10^{-4} per generation per haploid methylome) and backward
methylation rates (6.3×10^{-4} per generation per haploid methylome) for *A. thaliana* [6] as
priors.
683
684 *Genetic and methylation variant calling in Brachypodium distachyon*
685
We obtained whole genome sequencing data (29 samples) for *B. distachyon* from [27] and
whole genome bisulfite sequencing data from [16]. We used Bd21-3 v1.2 genome reference
and the GenBank annotation from the Joint Genome Institute (http://phytozome.jgi.doe.gov/)
for variant calling. We used Trimmomatic [57] to trim reads using the paired-end options as
stated above for the whole genome sequencing data and the single-end option for the whole
genome bisulfite sequencing data. We mapped whole genome sequencing data reads to the
reference using bwa-mem [69]. We used Picard Toolkit (https://broadinstitute.github.io/picard/)
to add read groups and mark duplicate reads. We used SAMtools [55] to remove unpaired
reads, remove reads with mapping quality of less than 20, and sort reads. We called genetic
variants using GATK HaplotypeCaller [70]. We only retained SNPs with depths greater than
five and quality scores greater than 30 using BCFtools [55]. Using this SNP variant call file,
we perform methylation variant calling in the same way as was done for *A. thaliana*. We
obtained the sample localities from [32] and created a map showing their occurrence with Map
Maker (https://maps.co/).
700
701 *Within and between population analyses in Brachypodium distachyon*
702
We used a minor allele frequency threshold of 10% to identify polymorphic sites in each clonal
group because fewer samples than *A. thaliana* were analyzed. For the analysis of SMPs and
DMRs, we only used variants that occurred within CG-gbM regions identified in [71]. We
estimated mSFS, LD, p-distance matrices, π, and Tajima's *D* in a similar way as we did for *A.*
thaliana. We thinned the SMP dataset so that only sites that were separated by 3Kbp were
sampled to reduce the computational time for the DAPC analyses.
709
710 *Generating the approximated transition matrix for demographic inference*
711
712 We created a training and testing dataset consisting of a combined 60×10^3 simulations
713 (80%/20%, respectively). Of these 10×10^3 were constant demographies (*i.e.*, no demographic
714 change) with a random population size drawn between 10×10^3 and 200×10^3, and the other
715 50×10^3 simulations were random walk demographies, with time steps being defined for i $\in$
716 {0,1,...,K − 1} using:
717

718
$$(\text{population\_time})_i = \frac{\exp\left(\frac{i}{K-1} \cdot \ln(1 + \delta t_{\max})\right) - 1}{\delta}$$

719
720 where *K*=42 is the number of time points, $\delta = 0.001$ is the growth rate, and $t_{\max} = 2.1 \times 10^6$ is the
721 maximum time.
722

The population size was drawn for each of the *K*=42 time points in the following way: an initial
population size was randomly selected between a minimum (N_min = 10×10^3) and maximum
(N_max = 1.2×10^6) value. For each subsequent time point, a new population size was
proposed by multiplying the previous size by a random factor between 0.1 and 10. If this
proposed size fell outside the allowed range (N_min to N_max), new proposals were
generated until an acceptable value was found. This process was repeated for all 42 time
points, resulting in a sequence of population sizes that exhibit both gradual and abrupt
changes while remaining within the specified bounds. Nevertheless, a sliding window average
with a size of 6 and a step size of 1 was applied to smooth extreme changes. This approach
for sampling demographies was adapted from Sanchez *et al*. [72].

The obtained population sizes and times were passed onto *msprime* [73] to simulate the
corresponding ancestral recombination graphs using a recombination rate of 3.4×10^{-8} per
base pair and generation. For SNPs, branches were mutated at a rate of 7×10^{-9} per base pair
and generation, leading to a ρ over θ ratio of 4.86.

Simulated and real data are summarized as a genotype matrix (*GM*) with dimensions $n \times m$,
where $n,m \in N$ represent the number of samples and mutations, respectively. Each element
g_{ij} of *GM*, where i $\in$ {1,...,n} and j $\in$ {1,...,m}, denotes the presence (g_{ij} = 1) or absence (g_{ij} = 0)
of mutation *j* in sample *i*. We generated a corresponding vector of positions of size *m* for each
segregating site containing floating point numbers of the location in the genome.

For each random population size parameter, we used a genome of length 4Mbp,10 samples
(ploidy=1), and the *GM* and positions vector to construct a density estimate of mutations along
the genome (in theory, corresponding to the respective TMRCA at that region). To ensure the
presence of enough markers, we segmented the genome into 20Kbp pieces and counted the
number of heterozygous markers between all pairwise $\binom{10}{2}$ combinations. The 4Mbp
sequences of all the comparisons were concatenated and log-transformed. Likewise, the
population time steps were also log-scaled. To make sure that the number of markers per
window fell approximately into the time windows, we scaled the markers by an arbitrarily

chosen scaling factor ($=e^{8.2}$) (Figure S13). The importance of the given scaling factor is
negligible, due to all the transformation steps occurring in the pre-training, and therefore any
noisy representation presented to the learning algorithm will be part of the training.
757
Each *msprime* simulation (under a random demographic history) yielded one *approximated*
transition matrix, which was obtained by counting the transitions (scaled number of
heterozygous sites in window t → window t+1) from one 20 Kbp window to the next using the
following algorithm: a sliding window of size 2 and step size 1 was applied along the sequence
and each pair of consecutive values in this window represents coordinates in the transition
matrix, which was incremented at those positions leading to a filled matrix which remained
unnormalized. Lastly, the entire transition matrix was incremented by one at each entry and
scaled by the natural log to squeeze all values approximately into the same range to avoid
training instability.
767
768 *Model description, training, and inference*
769
The inference approach can be described as a supervised learning approach using the
approximated transition matrices x with defined dimensions $R^{L=42 \times K=42}$ as the input and the
population size at each time step as the predicted variable. The input is a square matrix where
both dimensions (K and L) are 42. The model architecture is an encoder-only transformer [74]
(https://github.com/lucidrains/x-transformers). It consists of three main components: an input
projection layer, a transformer encoder, and an output projection layer. It can be formalized
as:

1. Input Projection: $x_{proj} = W_{in}x + b_{in}$, where $W_{in} \in R^{D_{enc} \times K_{in}}, b_{in} \in R^{D_{enc}}, x \in R^{L \times K_{in}}$
778
779 2. Transformer Encoder: $z = \text{Encoder}(x_{proj})$, where $z \in R^{L \times D_{enc}}$
780
781 The Encoder employs a configuration with dimension D_{enc} = 512, depth N_{enc} = 12, number
782 of attention heads H_{enc} = 10 heads, Gated Linear Unit activation in feed-forward layers [75],
783 residual attention connections [76], and rotary positional embeddings [77].
784
785 3. Dimension Reduction: $z_{reduced} = \sum_{i=1}^{L} z_i$, where $z_{reduced} \in R^{D_{enc}}$
786
787 4. Output Projection: $y = W_{out}z_{reduced} + b_{out}$, where $W_{out} \in R^{D_{out} \times D_{enc}}, b_{out} \in R^{D_{out}}$
788
789 In this formalization, sequence length *L* = the input dimension of the transition matrix K_{in} =
790 42, which represents the square input matrix of 42 x 42 dimensions, and the output
791 dimension D_{out} = 42.
792
The neural network model underwent training for 50 epochs, employing mean squared error
as the loss function. The training process leverages the Accelerate framework from Hugging
Face to enhance performance [78]. An Accelerator object was configured with mixed precision
using bfloat16 and gradient accumulation over 12 steps (with batch size 32). Throughout the
training regiment, both the training and test losses were continuously monitored to evaluate
model performance and assess convergence.
799

Generating data for demographic inference

We first generated unique variant call files for each sample from the population variant call files using BCFtools [55]. We then used the generate_multihetsep.py from MSMC-tools (https://github.com/stschiff/msmc-tools/tree/master) to combine all the samples per chromosome. For *A. thaliana* SNPs, we used the entire chromosome 1 as we needed a long contiguous fragment to generate the approximated transition matrix. For SMPs, we used 5Mbp segments in each of the five chromosomes (Table S2) as the SMP density was far higher. We used the list of genotyped cytosine positions as a genomic location mask for the SMPs. For *B. distachyon*, we included singletons to support the demographic inference given the small sample sizes in this species. The data were polarized by identifying the minor (less frequent) alleles and encoding them as derived, whereas the major alleles were encoded as ancestral. The resulting genotype matrix represents this polarized, unphased data, where each row corresponds to an individual and each column corresponds to a genomic position.

Methylation simulation model

We explored the boundaries of our course-grained demographic inference by adding a new mutation model to msprime [73]. We simulated a sawtooth demographic model with cyclical population size changes over time and a baseline effective population size of 2×10^5 individuals. Simulations were conducted on a 5Mbp sequence with a recombination rate of 3.4×10^{-8} per base pair per generation (ploidy=1), to isolate the influence of methylation dynamics.

A custom matrix mutation simulation was implemented to model methylation and demethylation processes, employing 2 allelic states: 0 (unmethylated) and 1 (methylated). The model incorporates 3 key rates: the site introduction rate (μ), methylation rate (β), and demethylation rate (δ). The methylation and demethylation rates are measured as factors of the site entry rate, where β = the methylation factor$\cdot\mu$ and δ = the demethylation factor$\cdot\mu$. The transition probabilities between the unmethylated and methylated states are determined by the ratio $\beta/(\beta+\delta)$ for methylation and $\delta/(\beta+\delta)$ for demethylation. The resulting 2×2 transition matrix encapsulates these probabilities:

$$\begin{bmatrix} 1 - [\beta/(\beta + \delta)] & \beta/(\beta + \delta) \\ \delta/(\beta + \delta) & 1 - [\delta/(\beta + \delta)] \end{bmatrix}$$

This framework allowed us to explore various methylation scenarios by adjusting β and δ relative to μ. We investigated cases where $\beta = \mu$ and $\delta = 2.5\mu$, $\beta = 10\mu$ and $\delta = 25\mu$, and $\beta = 100\mu$ and $\delta = 250\mu$, representing low, medium, and high methylation dynamics respectively, with the demethylation rate 2.5 times higher than the methylation rate. By varying the epimutation rate compared to the mutation rate, we test the robustness of our approach in correlating the number of markers in one Kbp windows with the average coalescent time. This proportionality, and the robustness of our method, is expected to decrease for higher epimutation rates, as these would violate the infinite site model for older coalescent times more strongly (due to homoplasy, [4]).

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The scripts for analyzing and plotting all the results, except for the demographic inference, have been published on GitHub (arunkumarramesh/methevolve) and archived using Zenodo (https://doi.org/10.5281/zenodo.13991812). The demographic inference method is available on GitHub (kevinkorfmann/aTMi) and has been archived using Zenodo (https://doi.org/10.5281/zenodo.13332833).

Competing interests

The authors declare that they have no competing interests.

Funding

RA was funded by the Peter and Traudl Engelhorn Foundation.

Authors' contributions

RA and AT designed the research. RA, KK, AW, BH, CG, and YW performed the research. RA, AT, KK, and AW wrote the paper. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank Thibaut Paul Patrick Sellinger for helping with the estimation of methylation rates. This work was supported by the de.NBI Cloud within the German Network for Bioinformatics Infrastructure (de.NBI) and ELIXIR-DE (Forschungszentrum Jülich and W-de.NBI-001, W-de.NBI-004, W-de.NBI-008, W-de.NBI-010, W-de.NBI-013, W-de.NBI-014, W-de.NBI-016, W-de.NBI-022).

References

1.   Richards CL, Alonso C, Becker C, Bossdorf O, Bucher E, Colome-Tatche M, Durka
W, Engelhardt J, Gaspar B, Gogol-Doring A, et al: **Ecological plant epigenetics:**
Evidence from model and non-model species, and the way forward. *Ecol Lett*
2017, **20:**1576-1590.
2.   Feng S, Cokus SJ, Zhang X, Chen PY, Bostick M, Goll MG, Hetzel J, Jain J, Strauss
SH, Halpern ME, et al: **Conservation and divergence of methylation patterning**
in plants and animals. *Proc Natl Acad Sci U S A* 2010, **107:**8689-8694.
3.   Zhang H, Lang Z, Zhu JK: **Dynamics and function of DNA methylation in plants.**
Nat Rev Mol Cell Biol 2018, **19:**489-506.
4.   Vidalis A, Zivkovic D, Wardenaar R, Roquis D, Tellier A, Johannes F: **Methylome**
evolution in plants. *Genome Biol* 2016, **17:**264.
5.   Cortijo S, Wardenaar R, Colome-Tatche M, Gilly A, Etcheverry M, Labadie K,
Caillieux E, Hospital F, Aury JM, Wincker P, et al: **Mapping the epigenetic basis**
of complex traits. *Science* 2014, **343:**1145-1148.
6.   van der Graaf A, Wardenaar R, Neumann DA, Taudt A, Shaw RG, Jansen RC,
Schmitz RJ, Colome-Tatche M, Johannes F: **Rate, spectrum, and evolutionary**
dynamics of spontaneous epimutations. *Proc Natl Acad Sci U S A* 2015,
112:6676-6681.
7.   Ossowski S, Schneeberger K, Lucas-Lledo JI, Warthmann N, Clark RM, Shaw RG,
Weigel D, Lynch M: **The rate and molecular spectrum of spontaneous**
mutations in Arabidopsis thaliana. *Science* 2010, **327:**92-94.
8.   Sellinger T, Johannes F, Tellier A: **Improved inference of population histories by**
integrating genomic and epigenomic data. *eLife* 2023, **12:**RP89470.
9.   Yao N, Schmitz RJ, Johannes F: **Epimutations Define a Fast-Ticking Molecular**
Clock in Plants. *Trends Genet* 2021, **37:**699-710.
10. Yao N, Zhang Z, Yu L, Hazarika R, Yu C, Jang H, Smith LM, Ton J, Liu L, Stachowicz
JJ, et al: **An evolutionary epigenetic clock in plants.** *Science* 2023, **381:**1440-
1445.
11. Muyle AM, Seymour DK, Lv Y, Huettel B, Gaut BS: **Gene Body Methylation in**
Plants: Mechanisms, Functions, and Important Implications for
Understanding Evolutionary Processes. *Genome Biol Evol* 2022, **14:**evac038.
12. Charlesworth B, Jain K: **Purifying selection, drift, and reversible mutation with**
arbitrarily high mutation rates. *Genetics* 2014, **198:**1587-1602.
13. Muyle A, Ross-Ibarra J, Seymour DK, Gaut BS: **Gene body methylation is under**
selection in Arabidopsis thaliana. *Genetics* 2021, **218:**iyab061.
14. Xu G, Lyu J, Li Q, Liu H, Wang D, Zhang M, Springer NM, Ross-Ibarra J, Yang J:
Evolutionary and functional genomics of DNA methylation in maize
domestication and improvement. *Nat Commun* 2020, **11:**5539.
15. Shirai K, Sato MP, Nishi R, Seki M, Suzuki Y, Hanada K: **Positive selective**
sweeps of epigenetic mutations regulating specialized metabolites in
plants. *Genome Res* 2021, **31:**1060-1068.
16. Eichten SR, Srivastava A, Reddiex AJ, Ganguly DR, Heussler A, Streich JC, Wilson
PB, Borevitz JO: **Extending the Genotype in Brachypodium by Including DNA**
Methylation Reveals a Joint Contribution with Genetics on Adaptive Traits.
G3 (Bethesda) 2020, **10:**1629-1637.

17. Le Vève A, Sammarco I, Latzel V, Lafon Placette C: **DNA methylation shows**
footprints of altitude selection in the clonal plant species Fragaria vesca.
bioRxiv 2024, **2024.03.19.585697**.

18. Sammarco I, Diez Rodriguez B, Galanti D, Nunn A, Becker C, Bossdorf O,
Munzbergova Z, Latzel V: **DNA methylation in the wild: epigenetic**
transgenerational inheritance can mediate adaptation in clones of wild
strawberry (Fragaria vesca). *New Phytol* 2024, **241:**1621-1635.

19. Cao S, Sawettalake N, Li P, Fan S, Shen L: **DNA methylation variations underlie**
lettuce domestication and divergence. *Genome Biol* 2024, **25:**158.

20. Rodriguez-Izquierdo A, Carrasco D, Anand L, Magnani R, Catarecha P, Arroyo-
Garcia R, Rodriguez Lopez CM: **Epigenetic differences between wild and**
cultivated grapevines highlight the contribution of DNA methylation during
crop domestication. *BMC Plant Biol* 2024, **24:**504.

21. Kawakatsu T, Huang SC, Jupe F, Sasaki E, Schmitz RJ, Urich MA, Castanon R,
Nery JR, Barragan C, He Y, et al: **Epigenomic Diversity in a Global Collection of**
Arabidopsis thaliana Accessions. *Cell* 2016, **166:**492-505.

22. Strütt S, Sellinger T, Glemin S, Tellier A, Laurent S: **Joint inference of**
evolutionary transitions to self-fertilization and demographic history using
whole-genome sequences. *Elife* 2023, **12:**e82384.

23. Vaswani A, Shazeer N, Parmar N, Uszkoreit J, Jones L, Gomez AN, Kaiser L,
Polosukhin I: **Attention Is All You Need.** *arXiv* 2017, **1706.03762**.

24. Devlin J, Chang M-W, Lee K, Toutanova K: **BERT: Pre-training of Deep**
Bidirectional Transformers for Language Understanding. *arXiv* 2018,
1810.04805.

25. Korfmann K, Gaggiotti OE, Fumagalli M: **Deep Learning in Population Genetics.**
Genome Biol Evol 2023, **15**.

26. Becker C, Hagmann J, Muller J, Koenig D, Stegle O, Borgwardt K, Weigel D:
Spontaneous epigenetic variation in the Arabidopsis thaliana methylome.
Nature 2011, **480:**245-249.

27. Wilson PB, Streich JC, Murray KD, Eichten SR, Cheng R, Aitken NC, Spokas K,
Warthmann N, Gordon SP, Accession C, et al: **Global Diversity of the**
Brachypodium Species Complex as a Resource for Genome-Wide
Association Studies Demonstrated for Agronomic Traits in Response to
Climate. *Genetics* 2019, **211:**317-331.

28. Wang J, Fan C: **A neutrality test for detecting selection on DNA methylation**
using single methylation polymorphism frequency spectrum. *Genome Biol*
Evol 2014, **7:**154-171.

29. Hewitt G: **Post-glacial re-colonization of European biota.** *Biological Journal of*
the Linnean Society 1999, **68:**87-112.

30. The1001GenomesConsortium: **1,135 Genomes Reveal the Global Pattern of**
Polymorphism in Arabidopsis thaliana. *Cell* 2016, **166:**481-491.

31. Sharbel TF, Haubold B, Mitchell-Olds T: **Genetic isolation by distance in**
Arabidopsis thaliana: biogeography and postglacial colonization of Europe.
Mol Ecol 2000, **9:**2109-2118.

32. Filiz E, Ozdemir BS, Budak F, Vogel JP, Tuna M, Budak H: **Molecular,**
morphological, and cytological analysis of diverse Brachypodium
distachyon inbred lines. *Genome* 2009, **52:**876-890.

33.   Ng PC, Henikoff S: **SIFT: Predicting amino acid changes that affect protein**
986   **function.** *Nucleic Acids Research* 2003, **31:**3812-3814.

987   34.   Minadakis N, Williams H, Horvath R, Caković D, Stritt C, Thieme M, Bourgeois Y,
988   Roulin AC: **The demographic history of the wild crop relative Brachypodium**
989   **distachyon is shaped by distinct past and present ecological niches.** *Peer*
990   *Community Journal* 2023, **3:**e84.

991   35.   Stritt C, Gimmi EL, Wyler M, Bakali AH, Skalska A, Hasterok R, Mur LAJ, Pecchioni
992   N, Roulin AC: **Migration without interbreeding: Evolutionary history of a highly**
993   **selfing Mediterranean grass inferred from whole genomes.** *Mol Ecol* 2022,
994   **31:**70-85.

995   36.   Schmitz RJ, Schultz MD, Urich MA, Nery JR, Pelizzola M, Libiger O, Alix A, McCosh
996   RB, Chen H, Schork NJ, Ecker JR: **Patterns of population epigenomic diversity.**
997   *Nature* 2013, **495:**193-198.

998   37.   Briffa A, Hollwey E, Shahzad Z, Moore JD, Lyons DB, Howard M, Zilberman D:
999   **Millennia-long epigenetic fluctuations generate intragenic DNA methylation**
1000   **variance in Arabidopsis populations.** *Cell Syst* 2023, **14:**953-967 e917.

1001   38.   Taudt A, Roquis D, Vidalis A, Wardenaar R, Johannes F, Colome-Tatche M:
1002   **METHimpute: imputation-guided construction of complete methylomes**
1003   **from WGBS data.** *BMC Genomics* 2018, **19:**444.

1004   39.   Xia J, Han L, Zhao Z: **Investigating the relationship of DNA methylation with**
1005   **mutation rate and allele frequency in the human genome.** *BMC Genomics*
1006   2012, **13:**S7.

1007   40.   Mugal CF, Ellegren H: **Substitution rate variation at human CpG sites**
1008   **correlates with non-CpG divergence, methylation level and GC content.**
1009   *Genome Biol* 2011, **12:**R58.

1010   41.   Kusmartsev V, Drozdz M, Schuster-Bockler B, Warnecke T: **Cytosine Methylation**
1011   **Affects the Mutability of Neighboring Nucleotides in Germline and Soma.**
1012   *Genetics* 2020, **214:**809-823.

1013   42.   Stuart T, Eichten SR, Cahn J, Karpievitch YV, Borevitz JO, Lister R: **Population**
1014   **scale mapping of transposable element diversity reveals links to gene**
1015   **regulation and epigenomic variation.** *Elife* 2016, **5:**e20777.

1016   43.   Shen Y, Zhang J, Liu Y, Liu S, Liu Z, Duan Z, Wang Z, Zhu B, Guo YL, Tian Z: **DNA**
1017   **methylation footprints during soybean domestication and improvement.**
1018   *Genome Biol* 2018, **19:**128.

1019   44.   Eichten SR, Briskine R, Song J, Li Q, Swanson-Wagner R, Hermanson PJ, Waters
1020   AJ, Starr E, West PT, Tiffin P, et al: **Epigenetic and genetic influences on DNA**
1021   **methylation variation in maize populations.** *Plant Cell* 2013, **25:**2783-2797.

1022   45.   Zhao T, Guan X, Hu Y, Zhang Z, Yang H, Shi X, Han J, Mei H, Wang L, Shao L, et al:
1023   **Population-wide DNA methylation polymorphisms at single-nucleotide**
1024   **resolution in 207 cotton accessions reveal epigenomic contributions to**
1025   **complex traits.** *Cell Res* 2024.

1026   46.   Avramidou EV, Korakaki E, Malliarou E, Boutsios S: **Studying the Genetic and**
1027   **the Epigenetic Diversity of the Endangered Species Juniperus drupacea**
1028   **Labill. towards Safeguarding Its Conservation in Greece.** *Forests* 2023,
1029   **14:**1271.

1030   47.   Avramidou EV, Moysiadis T, Ganopoulos I, Michailidis M, Kissoudis C, Valasiadis
1031   D, Kazantzis K, Tsaroucha E, Tsaftaris A, Molassiotis A, et al: **Phenotypic,**

1032    **Genetic, and Epigenetic Variation among Diverse Sweet Cherry Gene Pools.**
1033    *Agronomy* 2021, **11:**680.

1034 48. Wang MZ, Li HL, Li JM, Yu FH: **Correlations between genetic, epigenetic and**
1035    **phenotypic variation of an introduced clonal herb.** *Heredity (Edinb)* 2020,
1036    **124:**146-155.

1037 49. Langford N, Fargeot L, Blanchet S: **Spatial covariation between genetic and**
1038    **epigenetic diversity in wild plant and animal populations: a meta-analysis.** *J*
1039    *Exp Biol* 2024, **227**.

1040 50. Medrano M, Alonso C, Bazaga P, Lopez E, Herrera CM: **Comparative genetic and**
1041    **epigenetic diversity in pairs of sympatric, closely related plants with**
1042    **contrasting distribution ranges in south-eastern Iberian mountains.** *AoB*
1043    *Plants* 2020, **12:**plaa013.

1044 51. Fargeot L, Loot G, Prunier JG, Rey O, Veyssiere C, Blanchet S: **Patterns of**
1045    **Epigenetic Diversity in Two Sympatric Fish Species: Genetic vs.**
1046    **Environmental Determinants.** *Genes (Basel)* 2021, **12:**107.

1047 52. Lele L, Ning D, Cuiping P, Xiao G, Weihua G: **Genetic and epigenetic variations**
1048    **associated with adaptation to heterogeneous habitat conditions in a**
1049    **deciduous shrub.** *Ecol Evol* 2018, **8:**2594-2606.

1050 53. Durvasula A, Fulgione A, Gutaker RM, Alacakaptan SI, Flood PJ, Neto C,
1051    Tsuchimatsu T, Burbano HA, Pico FX, Alonso-Blanco C, Hancock AM: **African**
1052    **genomes illuminate the early history and transition to selfing in Arabidopsis**
1053    **thaliana.** *Proc Natl Acad Sci U S A* 2017, **114:**5213-5218.

1054 54. Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, Muller R,
1055    Dreher K, Alexander DL, Garcia-Hernandez M, et al: **The Arabidopsis**
1056    **Information Resource (TAIR): improved gene annotation and new tools.**
1057    *Nucleic Acids Res* 2012, **40:**D1202-1210.

1058 55. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A,
1059    Keane T, McCarthy SA, Davies RM, Li H: **Twelve years of SAMtools and**
1060    **BCFtools.** *Gigascience* 2021, **10:**giab008.

1061 56. Nassar LR, Barber GP, Benet-Pages A, Casper J, Clawson H, Diekhans M, Fischer
1062    C, Gonzalez JN, Hinrichs AS, Lee BT, et al: **The UCSC Genome Browser**
1063    **database: 2023 update.** *Nucleic Acids Res* 2023, **51:**D1188-D1195.

1064 57. Bolger AM, Lohse M, Usadel B: **Trimmomatic: a flexible trimmer for Illumina**
1065    **sequence data.** *Bioinformatics* 2014, **30:**2114–2120.

1066 58. Krueger F, Andrews SR: **Bismark: a flexible aligner and methylation caller for**
1067    **Bisulfite-Seq applications.** *Bioinformatics* 2011, **27:**1571-1572.

1068 59. Lister R, O'Malley RC, Tonti-Filippini J, Gregory BD, Berry CC, Millar AH, Ecker JR:
1069    **Highly integrated single-base resolution maps of the epigenome in**
1070    **Arabidopsis.** *Cell* 2008, **133:**523-536.

1071 60. Hazarika RR, Serra M, Zhang Z, Zhang Y, Schmitz RJ, Johannes F: **Molecular**
1072    **properties of epimutation hotspots.** *Nat Plants* 2022, **8:**146-156.

1073 61. Williams CJ, Dai D, Tran KA, Monroe JG, Williams BP: **Dynamic DNA methylation**
1074    **turnover in gene bodies is associated with enhanced gene expression**
1075    **plasticity in plants.** *Genome Biol* 2023, **24:**227.

1076 62. Hamala T, Tiffin P: **Biased Gene Conversion Constrains Adaptation in**
1077    **Arabidopsis thaliana.** *Genetics* 2020, **215:**831-846.

63. Paradis E: **pegas: an R package for population genetics with an integrated-**
modular approach. *Bioinformatics* 2010, **26:**419-420.

64. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker
RE, Lunter G, Marth GT, Sherry ST, et al: **The variant call format and VCFtools.**
Bioinformatics 2011, **27**.

65. Zhang C, Dong SS, Xu JY, He WM, Yang TL: **PopLDdecay: a fast and effective**
tool for linkage disequilibrium decay analysis based on variant call format
files. *Bioinformatics* 2019, **35:**1786-1788.

66. Lefort V, Desper R, Gascuel O: **FastME 2.0: A Comprehensive, Accurate, and**
Fast Distance-Based Phylogeny Inference Program. *Mol Biol Evol* 2015,
32:2798-2800.

67. Yu G, Smith DK, Zhu H, Guan Y, Lam TTY: **ggtree: an r package for visualization**
and annotation of phylogenetic trees with their covariates and other
associated data. *Methods in Ecology and Evolution* 2017, **8:**28-36.

68. Jombart T: **adegenet: a R package for the multivariate analysis of genetic**
markers. *Bioinformatics* 2008, **24:**1403–1405.

69. Li H: **Aligning sequence reads , clone sequences and assembly contigs with**
BWA-MEM. 2013, **00:**1-3.

70. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A,
Garimella K, Altshuler D, Gabriel S, Daly M, DePristo MA: **The Genome Analysis**
Toolkit: A MapReduce framework for analyzing next-generation DNA
sequencing data. *Genome Research* 2010, **20:**1297–1303.

71. Niederhuth CE, Bewick AJ, Ji L, Alabady MS, Kim KD, Li Q, Rohr NA, Rambani A,
Burke JM, Udall JA, et al: **Widespread natural variation of DNA methylation**
within angiosperms. *Genome Biol* 2016, **17:**194.

72. Sanchez T, Cury J, Charpiat G, Jay F: **Deep learning for population size history**
inference: Design, comparison and combination with approximate Bayesian
computation. *Mol Ecol Resour* 2021, **21:**2645-2660.

73. Baumdicker F, Bisschop G, Goldstein D, Gower G, Ragsdale AP, Tsambos G, Zhu
S, Eldon B, Ellerman EC, Galloway JG, et al: **Efficient ancestry and mutation**
simulation with msprime 1.0. *Genetics* 2022, **220**.

74. Paszke A, Gross S, Massa F, Lerer A, Bradbury J, Chanan G, Killeen T, Lin Z,
Gimelshein N, Antiga L, et al: *pyTorch: An Imperative Style, High-Performance*
Deep Learning Library. Curran Associates, Inc.; 2019.

75. Shazeer N: **GLU Variants Improve Transformer.** *arXiv* 2020:2002.05202.

76. He R, Ravula A, Kanagal B, Ainslie J: **RealFormer: Transformer Likes Residual**
Attention. *arXiv* 2021:2012.11747.

77. Su J, Ahmed M, Lu Y, Pan S, Bo W, Liu Y: **RoFormer: Enhanced transformer with**
Rotary Position Embedding. *Neurocomputing* 2024, **568:**127063.

78. Gugger S, Debut L, Wolf T, Schmid P, Mueller Z, Mangrulkar S, Sun M, Bossan B:
Accelerate: Training and inference at scale made simple, efficient and
adaptable. *https://githubcom/huggingface/accelerate* 2022.

1120