**Methylomes reveal recent evolutionary changes in populations of two plant species**

Kevin Korfmann[ab1], Andreas Zauchner[a1], Bing Huo[a], Corinna Grünke[a], Yitong Wang[a], Aurélien

Tellier[a†2], Ramesh Arunkumar [a†2]

[a]Section of Population Genetics, School of Life Sciences, Technical University of Munich,

Liesel-Beckmann Strasse 2, 85354 Freising, Germany

[b]Present address: Department of Biology 77 Klamath Hall 1210 University of Oregon Eugene,

OR 97403

[1] These authors contributed equally

[2] These authors contributed equally

[†] Authors for correspondence (Aurélien Tellier - aurelien.tellier@tum.de, Ramesh Arunkumar

– arunkumar.ramesh@tum.de)

**Abstract**

Plant DNA methylation changes occur hundreds to thousands of times faster than DNA mutations and can be transmitted transgenerationally, making them useful for studying population-scale patterns in clonal or selfing species. However, a state-of-the-art approach to use them for inferring population genetic processes and demographic histories is lacking. To address this, we compare evolutionary signatures extracted from CG methylomes and genomes in *Arabidopsis thaliana* and *Brachypodium distachyon*. While methylation variants (SMPs) are less effective than genetic variants (SNPs) for identifying population differentiation in *A. thaliana*, they can classify phenotypically divergent *B. distachyon* subgroups that are otherwise genetically indistinguishable. The site frequency spectra generated using methylation sites from varied genomic locations and evolutionary conservation exhibit similar shapes, indicating minimal noise when all CG sites are analyzed jointly. Nucleotide diversity is three orders of magnitude higher for methylation variants than for genetic variants in both species, driven by the higher epimutation rate. Correlations between SNPs and SMPs in nucleotide diversity and allele frequencies at gene exons are weak or absent in *A. thaliana*, possibly because the two sources of variation reflect evolutionary forces acting at different timescales. Linkage disequilibrium quickly decays within 250 bp for methylation variants in both plant species. Finally, we developed a deep learning-based demographic inference approach. We identified recent population expansions in *A. thaliana* and *B. distachyon* using methylation variants that were not identified when using genetic variants. Our study demonstrates the unique evolutionary insights methylomes provide that genetic variation alone cannot reveal.

**Keywords:** Methylation site frequency spectra, demographic inference, deep learning, Tajima's *D*, linkage disequilibrium, nucleotide diversity, clocklike regions, gene body methylation, *Arabidopsis thaliana*, *Brachypodium distachyon*

**Background**

Understanding how variation is partitioned within and between populations is a central goal in evolutionary biology. Increasing evidence suggests that epigenetic modifications can vary within and between populations [1]. Nevertheless, our knowledge of how epigenomes evolve through neutral and selective forces and, consequently, their capacity to reveal population demographic histories is limited. DNA cytosine methylation, a conserved epigenetic modification in eukaryotes [2], occurs in plants within three sequence contexts: CG, CHG, and CHH (H represents A, T, or C) [3]. DNA methylation is known to be involved in regulating gene expression, suppressing transposable element activity, and stabilizing the genome [3]. Long-term changes in the methylome are mainly shaped by genomic alterations such as the proliferation of transposable elements and mutations affecting DNA methylation pathways, whereas heritable stochastic epimutations that occur independent of genomic background drive short-term changes [4].

The unique characteristics of epimutations make them appealing markers for evolutionary investigations. In plants, CG methylation is maintained during cell division by METHYLTRANSFERASE 1 (MET1) [3]. Spontaneous stochastic loss or gain of methylation at CG sites can be transmitted stably across multiple generations [5]. Van der Graaf *et al*. have estimated the forward (gain of DNA methylation) and backward (loss of methylation) epimutation rates per CG site in the model plant *Arabidopsis thaliana* to be approximately $2.56 \times 10^{-4}$ and $6.3 \times 10^{-4}$, respectively, per generation per haploid methylome [6]. These rates are several orders of magnitude higher than the mutation rate of $7 \times 10^{-9}$ base substitutions per site per generation in the same population [7]. The increased rates of change at the epigenetic level may facilitate evolutionary inference over recent timescales, where the utility of slower-evolving genetic markers is limited [8, 9].

As transiently induced epimutations can be mistaken for stably inherited markers, a recent study exposed genetically identical *A. thaliana* to various environmental conditions to identify

genomic regions—termed clocklike regions—where CG divergence rates are independent of environmental stimuli [10]. While these regions are valuable for evolutionary studies, they are challenging to identify in most species. Therefore, the same study observed that gene body methylated regions (gbM) exhibit a pattern similar to that of CG divergence in clocklike regions [10]. gbM refers to a subclass of genes that display elevated CG methylation. They are mostly found in moderately and constitutively expressed housekeeping genes [11]. As they can be identified based on the distribution of methylation in the gene bodies, their use may facilitate the investigation of evolutionary processes in nonmodel plant species.

Previous attempts to explore how methylation variants evolve have either neglected demographic (neutral) models, mixed environmentally sensitive and insensitive methylation sites, or did not adequately assess the interdependence of genetic and methylation variants. Charlesworth and Jain's analytic framework for hypermutable sites [12] has been applied to create a methylation site frequency spectrum (mSFS) [4]. Previous studies generated mSFS for genic CG sites and the methylation states of gbM genes, finding either no selection or only weak selection acting on the methylome in *A. thaliana* [4, 13] and maize [14]. However, these studies do not incorporate an explicit demographic model while assessing selection. Another study found that positive selection acted on methylation variants linked to metabolite regulation in *A. thaliana*, even after accounting for demographic histories [15]. Other studies in *Brachypodium distachyon* [16], *Fragaria vesca* [17, 18], lettuce [19], and grapevines [20] have compared evolutionary inferences using genetic and methylation variants by analyzing all CG sites as the environmental sensitivity of individual sites is unknown in most plant species. Recently, Sellinger *et al*. utilized methylation as a secondary marker alongside genetic variants to enhance demographic inference over short and long timescales [8]. The conceptual novelty was to consider that the underlying genealogy at a locus (*i.e.* the coalescent tree tracing back the ancestry of the sample back in time) is common for SNPs and SMPs (under a neutral model of evolution). This represents the first population genomics inference model combining DNA mutations, epimutations, and recombination along the genome, and the basis for

developing a Sequential Markovian Coalescent method (SMCm). However, that study exhibits a fair amount of noise in the inference of past evolutionary history when it was applied to methylome data from *A. thaliana* [8]. These inaccuracies are due to a lack of knowledge about the optimal SMP markers to use, the relevant genomic context, the relationship between SNPs and SMPs, and the SMCm method being very sensitive and not accounting well for methylated region effects (so-called differentially methylated regions or DMRs). A related question is whether DMRs can be used for evolutionary inferences as well.

Given these limitations, our goal was to assess how well methylation variants reveal known evolutionary patterns inferred using genetic variants. We chose *A. thaliana* and *B. distachyon* for our analyses because their evolutionary histories are well-described [16, 21]. We found that the capacity of methylation variants to differentiate populations is contingent on when populations diverge. We also show that CG sites across the methylome capture within- and between-population patterns comparable to environmentally insensitive sites (clocklike regions and gbM) [10] in samples grown in a common garden. We found weak or no local correlations between genetic and methylation variants, indicating they reflect different aspects of population evolutionary histories. Additionally, the linkage between neighboring methylation variants was minimal. Conventional genome-based demographic inference methods, such as the Sequential Markovian Coalescent [22], show limitations in using methylation variants due to high sensitivity to violations of the Markovian properties of the underlying coalescent model along the genome [8]. Moreover, since such methods require knowing the variant accumulation rates and distributions—often unknown for methylation in many species—we developed a deep learning method that uses methylation as a neutral marker to estimate past demography. Unlike conventional convolutional networks, we use (to our knowledge) the first Encoder-Only Transformer architecture [23-25] method, focusing on the placement of mutations on a hidden ancestral recombination graph. We track estimates of coalescent events in a time-discretized transition matrix by using heterogeneous sites per genomic window as input, thereby applying a coarse-grained approach, similar to tsABC [22]. By

aligning training and real data distributions, we show that rapidly changing methylation variants

can reveal recent demographic shifts in selfing and clonal species.

**Results**

**CG methylation variants have variable performance in identifying population structure**

We analyzed genetic and methylation diversity using 63 samples each from the Central

European (CEU) and the Iberian non-relicts (IBnr) *A. thaliana* populations (Table S1). These

populations were used by Strütt *et al*. [22]. Across the two populations, 31-33% of CG

methylation sites in clocklike regions were polymorphic (Figure S1A), which is similar to a

previous estimate for CG sites in *A. thaliana* mutation accumulation lines [26]. In contrast, 1.6-

1.7% of coding sites and 2.7-2.8% of non-coding sites were polymorphic (Figure S1A). We

compared our results from *A. thaliana* with those from 29 samples from the clonally

propagating grass *B. distachyon*. Wilson *et al*. used SNPs to identify clonal groups among

Turkish *B. distachyon*, with samples within each group showing high levels of genetic similarity

[27]. Eichten *et al*. identified high levels of consistency between the known genetic

relationships for the clonal groups and the extent of CG divergence [16]. Here, we restrict our

focus to SMPs and DMRs occurring within gbM regions and two clonal groups of *B. distachyon*

with average bisulfite sequencing depths greater than 10 [16] (Group 1: 22 samples, Group 6:

7 samples) (Table S1). Between 4-7% of CG methylation sites within gbM regions were

polymorphic in *B. distachyon* (Figure S1B). The median length of DMRs in *B. distachyon* was

107 bp (1$^{st}$ percentile: 44 bp, 99% percentile: 717 bp) which is smaller than the median length

in *A. thaliana* DMRs, which was 181 bp (1$^{st}$ percentile: 78 bp, 99% percentile: 1125 bp) (Figure

S2).

**Figure 1**. Methylomes outperform genetic variants in detecting population subdivision in *Brachypodium distachyon* but not *Arabidopsis thaliana*. (**A**) Neighbor-joining trees were generated using p-distance matrices for non-coding single nucleotide polymorphisms (SNP), coding SNPs, clocklike region single methylation polymorphisms (SMP), and clocklike region differentially methylated regions (DMR) for Central European (CEU) and the Iberian non-relicts (IBnr) *A. thaliana*. (**B**) Neighbor-joining trees generated using p-distance matrices estimated for SMP and DMR in gbM regions of Turkish *Brachypodium distachyon* clonal groups.

We estimated the proportion of sites at which two samples were different (p-distance) and

clustering to uncover how the net total of genetic and methylation variation differed between

samples from the CEU and IBnr populations. In this approach, a clear separation between the

CEU and IBnr populations was apparent with both coding and non-coding single nucleotide

polymorphisms (SNP) (Figure 1A). Although single methylation polymorphisms (SMP) and

differentially methylated regions (DMR) largely separated the two populations, some samples

were separated by only minimal distances (Figure 1A). Clustering indicated nine or more

groups within the SNP data (Figure S3A-B), whereas only two groups were apparent in the

SMP and DMR datasets (Figure S3C-D).

Furthermore, we performed discriminant analysis of principal components (DAPC) analyses

to identify how correlated changes in subsets of genetic and methylation variants can be used

to discriminate the two populations. DAPC indicated greater discriminatory power with coding

and non-coding SNPs compared to SMPs and DMRs (Figure S3E). Although p-distance

matrix-based comparisons could have been influenced by the greater number of SNPs (Figure

1A), marker numbers were comparable for the DAPC, and yet SMPs and DMRs showed poor

discriminatory capability (Figure S3E). SMPs and DMRs that occur within gbM genes have

comparable performance to those occurring within clocklike regions in discriminating the two

populations (Figure S4). High methylation rates in *A. thaliana* may have caused multiple

changes at the same site (so-called homoplasy accounted for as a finite site model [4, 12,

28]), resulting in the underestimation of the true levels of divergence. Indeed, the divergence

between IBnr and CEU populations is estimated to have occurred after the last glacial

maximum ca. 10,000-20,000 years ago [29-31].

We then performed a similar analysis using the *B. distachyon* accessions. In contrast to the

clonal grouping based on the genetic profiles [27], SMPs and DMRs were capable of

additionally distinguishing the two subgroups of Group 1 (Figures 1B, S5). The trees estimated

using the p-distance matrices were shorter for SMPs compared to DMRs (Figure 1B), possibly

due to increased error in this dataset because of the shallow bisulfite sequencing depths.

However, both trees were longer compared to those in *A. thaliana* (Figure S4A) indicating

greater methylome differentiation in *B. distachyon*. Both our results using gbM regions and the

results from Eichten *et al*. that use the entire collection of CG sites [16] reveal a difference in

methylation profiles between the two subgroups of Group 1. We obtained phenotype data for

the two subgroups [32] and found significant differences in the longitude (Figure S6), elevation

of collection site, seed lengths, widths, yield production, and germination percentages (Figure

S7). Our findings suggest that methylation variants can reveal subpopulation structure in

recent timescales, where there has been little time for genetic differences and multiple

methylation changes to accumulate.

**Genomic contexts have little influence on the shape of SMP distributions**

**Figure 2**. Methylation site frequency spectra generated using single methylation polymorphisms were similar regardless of genomic contexts. Methylation site frequency spectra (mSFS) generated using differentially methylated regions (DMR) exhibited greater variability. (**A**) mSFS for the Central European (CEU, Salmon) and the Iberian non-relics (IBnr, Cyan) *Arabidopsis thaliana* populations using CG sites that occur in clocklike, gene body methylated (gbM), non-gbM, and low and high Sorting Intolerant from Tolerant (SIFT) score regions. Only sites with minor allele frequencies (MAF) >0.02 were used to generate the

mSFS. (**B**) mSFS for *Brachypodium distachyon* clonal groups 1.1, 1.2, and 6 for single

methylation polymorphisms (SMP) and DMRs occurring within gbM and non-gbM regions.

Only sites with minor allele frequencies (MAF) >0.04 were used to generate the mSFS. For

(A) and (B), the numbers in each panel indicate the number of sites used for the analyses.

We sought to understand how genomic contexts influenced the methylation site frequency

spectra (mSFS) in *A. thaliana*. The mSFS reveals how methylation rates, demographic

changes, and natural selection influence changes in the methylome over time [4, 12]. We

found a greater frequency of unmethylated alleles at most polymorphic methylation sites for

the CEU and IBnr populations, but CEU consistently had higher frequencies of unmethylated

alleles (Figure 2A). The shapes of the mSFS for SMPs are consistent with previous findings

that the backward methylation rate promoting the change from methylated to unmethylated is

higher than the forward rate [6] and are consistent with previous mSFS generated by

combining *A. thaliana* samples from extended geographic regions [4, 13]. The shape of the

mSFS for SMPs in clocklike and gbM regions were similar, supporting their interchangeability.

We also studied the mSFS for intervals containing genes with low and high Sorting Intolerant

From Tolerant (SIFT) scores, which are positively correlated with the level of evolutionary

constraint that genes are under [33]. Our results indicate that the level of gene constraint has

little impact on the mSFS shape (Figure 2A).

When compared to SMPs, the DMR mSFS for *A. thaliana* were more dependent on the

underlying set of intervals used to generate them. For DMRs that occur in intervals containing

non-gbM genes and high and low SIFT-score genes, more samples have the unmethylated

state reflecting the underlying SMP distributions (Figure 2A). This is also true when DMRs

across the entire genome are analyzed (Figure 2A). In contrast, for DMRs in clocklike regions

and gbM genes, more samples exhibit the methylated state (Figure 2A). The mSFS shapes

for DMRs show greater variability compared to those for SMPs, which may reflect underlying

differences in their molecular basis.

In contrast to the mSFS for SMPs in *A. thaliana* (Figure 2A), the mSFS for all three clonal

groups of *B. distachyon* indicated an excess of methylated alleles within gbM and non-gbM

SMPs, although it was less pronounced for the latter (Figure 2B). There was less consistency

in the shapes of the mSFS for DMRs in gbM and non-gbM regions (Figure 2B), similar to our

observation when *A. thaliana* DMRs were analyzed (Figure 2A).

**Methylation and genetic variants show unique evolutionary signatures in *A. thaliana***

**Figure 3**. Unique evolutionary signatures in genomes and methylomes. (**A**) Estimates of pairwise nucleotide diversity (π) scaled by the number of cytosines per region for single methylation polymorphisms ($\pi_{SMP}$) and the number of bases for single nucleotide polymorphisms ($\pi_{SNP}$) for the Central European (CEU) and the Iberian non-relicts (IBnr) *Arabidopsis thaliana* populations. (**B**) Estimates of Tajima's $D$ per region for CEU and IBnr. Comparisons of (**C**) $\pi_{SNP}$ versus $\pi_{SMP}$ and (**D**) Tajima's $D$ for SNPs ($D_{SNP}$) and SMPs ($D_{SMP}$)

per gene for CEU and IBnr. The numbers in the titles of each panel indicate the number of

sites used for the analyses. Correlation tests were conducted to assess the associations

between π and Tajima's *D*. (**E**) $\pi_{SMP}$ scaled by the number of cytosines per region for

*Brachypodium distachyon* clonal groups 1.1, 1.2, and 6. (**F**) $D_{SMP}$ per region for *B. distachyon*.

For (A), (B), (E), and (F), the numbers above the boxplots indicate the number of intervals.

Letters indicate group-wise differences in means following Tukey's honest significant

difference tests. For (F), the color scheme and number of sites are the same as in (E).

We next sought to understand how genetic and methylation polymorphisms are distributed

within *A. thaliana* populations. We first calculated the pairwise nucleotide diversity (π) for

SMPs ($\pi_{SMP}$) and SNPs ($\pi_{SNP}$) from noncoding, clocklike, gbM, non-gbM, and high and low

SIFT genic regions. We scaled estimates by the total number of sites (polymorphic and

monomorphic) to ensure that the results were comparable (that is per-site estimates). $\pi_{SMP}$ is

three orders of magnitude higher than $\pi_{SNP}$ (Figure 3A), likely driven by the high rates of

change in methylomes [6]. $\pi_{SNP}$ in the CEU population and $\pi_{SMP}$ in CEU and IBnr populations

for clocklike regions were significantly higher than in other regions (Figure 3A), possibly

indicating better detection of environmental insensitivity with more polymorphic sites. $\pi_{SMP}$ was

higher for gbM and high-SIFT genes than at non-gbM and low-SIFT, respectively, but no such

difference was seen for $\pi_{SNP}$ (Figure 3A), indicating that constraint at the genic level has a

minor effect on the levels of local methylation diversity.

We then compared how selective and demographic processes influence genomes and

methylomes by estimating Tajima's *D* for CEU and IBnr. For methylomes, we used a modified

Tajima's *D* using the $D^m$ [28] that accounts for the hypermutability of the methylation sites.

$D_{SNP}$ was slightly negative for noncoding sites in the CEU and IBnr populations (Figure 3B),

reflecting the known history of population expansions [22]. The higher $D_{SNP}$ for IBnr (Figure

3B) is consistent with larger effective population sizes over recent timescales [22]. In contrast,

IBnr had significantly lower $D_{SMP}$ compared to CEU (Figure 3B). This could be due to variation

in methylation rates, purifying selection, or demographic and selective (purifying selection)

histories. Therefore, we used SMCm [8], which employs fixed mutation rates and compares

the distributions of mutation and methylation markers, to estimate the forward and backward

methylation rates for the IBnr and CEU populations. Methylation rates were only slightly higher

in CEU but the ratio of backward to forward rates was similar in both populations (Figure S8).

Hypermutability should result in greater frequencies of intermediate variants [12], but we did

not observe such a pattern (Figure 2A). Tajima's $D$ is based on a comparison of the number

of segregating sites versus π. The disparity in $D_{SMP}$ appears to be driven primarily by the

variation in the number of segregating sites per region (Figure S9). This difference suggests

that the demographic histories of the two populations may be slightly different.

We then estimated Tajima's $D$ in different genomic contexts for CEU and IBnr. Clocklike

regions had significantly higher $D_{SNP}$ and $D_{SMP}$ compared to other regions (Figure 3B). The $D^m$

method assumes that forward and backward methylation rates are equivalent [28]. Although

the ratio of the backward to forward methylation rates in clocklike regions is 1.36, the ratios

are 2.11 and 2.84 for gbM and non-clocklike regions, respectively [10], indicating the

assumption of $D^m$ is less strongly violated for clocklike regions. $D_{SMP}$ for non-gbM regions was

slightly but significantly higher compared to the estimates for gbM regions (Figure 3B),

suggesting that ignoring the differences between regions can introduce minimal noise when

inferring evolutionary signatures from methylomes. The differences in $D_{SMP}$ between high-SIFT

vs low-SIFT were not consistent in both populations (Figure 3B). There was only a weak

positive correlation for π (Figure 3C) generated for SNPs vs SMPs that occur in coding regions

and no significant correlation for Tajima's $D$ (Figure 3D).

We compared π and Tajima's $D$ for *B. distachyon* SMPs generated using SNPs for the

Anatolian and Turkish lineage (B_East) [34, 35] to which all three clonal groups belong. Scaled

π for SMPs was larger for the *B. distachyon* clonal groups compared to the *A. thaliana*

populations, but there were no consistent differences between gbM and non-gbM regions or

among clonal groups (Figure 3A, 3E). Similar to *A. thaliana* (Figure 3A), scaled π for SMPs in

all three clonal groups was three orders of magnitude larger than the median SNP-based

scaled π (~1.8×10⁻⁶) for the B_East lineage [34, 35] (Figure 3E). Similar to *A. thaliana* (Figure

3B), $D_{SMP}$ in the three clonal groups was negative. $D_{SMP}$ was slightly but significantly larger in

non-gbM regions compared to gbM regions (Figure 3F). These negative values contrast with

the positive $D_{SNP}$ for genetic variants in the B_East lineage [34]. Unlike in *A. thaliana* (Figure

S9), this contrast cannot be attributed primarily to the differences in the number of segregating

sites (Figure S10), meaning that it cannot be explained solely by the extent of genotyping.

$D_{SMP}$ was lower for Group 1.1 compared to the other two groups (Figure 3F), which is

consistent with the higher frequency of rare alleles in this group (Figure 2B).

**Linkage disequilibrium quickly erodes for methylation variants**

**Figure 4**. Weak linkage disequilibrium (LD) between neighboring single methylation

polymorphisms (SMP). (**A**) Pairwise r² estimated using coding single nucleotide

polymorphisms (SNP), non-coding SNPs, and SMPs for Central European (CEU) and Iberian

non-relicts (IBnr) *Arabidopsis thaliana* populations. The number of sites used for comparisons

is indicated within the plots. (**B**) Pairwise r² was estimated using SMPs for the clonal groups.

For (A) and (B), the number of sites used for comparisons is indicated within the plots.

An important consideration for population genetic analysis is the presence of sites with

independent evolutionary histories [8]. Therefore, we assessed the levels of pairwise linkage

disequilibrium (LD) to identify the number of independent polymorphic SNPs and SMPs.

Pairwise LD decay was stronger for noncoding SNPs compared to coding SNPs in *A. thaliana*

(Figure 4A). However, LD decay was much stronger for SMPs than for SNPs with $r^2$ dropping

below 0.1 within the first 250 bp (Figure 4A). LD decays to 0.1-0.25 for SMPs sites within gbM

regions for *B. distachyon* within the first 100 bp but remains constant thereafter (Figure 4B)

and is stronger compared to LD in *A. thaliana* over large distances (Figure 4A).

**Demographic inference using genetic and methylation variants**

**A)**

*1. Count number of heterozygous markers per window*

Cont.

w₁  w₂  w₃

Compute for pairwise comparisons of samples

*2. Multiply scaling factor (s) and number of markers to approximate average TMRCA estimate per window*

*3. Counting transitions from window to window to approximate transition matrix*

Scaled number of makers (w₁, w₂, w₃, …)

*4. Predict demography from approximated transition matrix*

Model: Encoder-only Transformer
Input: Approximated Transition Matrix
Output: Population size at n time points

**Figure 5.** Demographic inference using mutation or methylation markers. **(A)** Schematic of

the approximated transition matrix inference framework. The genome is first divided into 20

Kbp bins, and heterozygous markers are counted for all pairwise comparisons in a sample of 10 individuals. These counts are then scaled by an empirically chosen factor to ensure sufficient values in the approximated transition matrix by aligning the number of markers per window to the TMRCA in that window. The framework constructs this matrix by concatenating all pairwise comparisons into a long sequence and applying a sliding window of size 2 and step size 1. Each pair of consecutive values in this window represents coordinates in the transition matrix, which is incremented at those positions. This process results in an unnormalized transition matrix that captures the genetic relationships between samples. Finally, an encoder-only transformer neural network takes this noisy matrix as input, implicitly learning the fixed scaling factor, and predicts population sizes at various time points. Inference using over 42 time steps using (**B**) single nucleotide polymorphisms (SNP) and (**C**) single methylation polymorphisms (SMP) for the central European (CEU) and the Iberian non-relicts (IBnr) *Arabidopsis thaliana* populations. (**D**) Demographic inference using SMP for Groups 1.1 and 1.2 of *Brachypodium distachyon*. SMP inference was done using sequences from all five chromosomes (light lines, average in bold), while SNP inference is based on chromosome one. (**E**) Inference of simulated sawtooth demography using five replicates, 4Mbp segments, and 10 samples.

We developed a novel inference method that focuses on large genome blocks (20 Kbp windows) likely spanning multiple coalescent genealogies (Figure 5A), to estimate the times to the most recent common ancestor (TMRCA). Our method assumes that the number of heterozygous markers per window is correlated with the average TMRCA. Essentially, longer coalescent times allow more mutations and epimutations to accumulate. To mitigate the possibility that older genealogies (long TMRCA) can be unexpectedly short due to immediate recombination, our approach uses 20 Kbp windows to average out genealogies. We apply a scaling to the number of markers to estimate the TMRCA (Figure S11). Although the chosen scaling factor will not perfectly offset all variable demographic samples—given that we trained

using a fixed recombination rate and a fixed window size—offsets are minimal and addressed

during training. An encoder-only transformer neural network processes this data, learns the

scaling factor, and predicts population sizes over time. This is done by summarizing the

TMRCA of two consecutive windows in the empirical transition matrix. The differences

between each pair of samples are used to infer their TMRCA. The TMRCA values for all pairs

across multiple intervals (windows) in the genome are then represented as a transition matrix.

This method combines raw genotype data restructuring with deep learning to infer

demographic history.

For deep learning inference, the distributions of heterozygous markers in training and real data

must overlap to ensure in-distribution inference. For the training data, we collapsed each

training transition matrix into a single scalar by summing the flattened matrix and plotted the

distribution of unnormalized transition matrices (Figure S12A). To ensure the real data *from A.*

*thaliana* and *B. distachyon* matched the training data, we analyzed 20 randomly sampled

genomic regions. For the SNP data, we selected start positions uniformly between 0-28 Mbp,

whereas for the methylation data, the start positions were chosen uniformly between 0-0.5

Mbp. For both data types, we determined each region's length by sampling from a uniform

distribution between 1-3 Mbp. Summing the resulting transition matrices for the IBnr, CEU,

Group 1.1, and Group 1.2 regions (Figure S12B-D) produced values within the training data

range (Figure S12A). Despite the differential properties of SMPs and SNPs, the distribution of

heterozygous markers per 20 Kbp window was similar across populations (Figure S12B-C),

supporting the use of a common training dataset for deep learning.

We applied our coarse−grained inference approach to *A. thaliana* and *B. distachyon*. We

generated approximated transition matrices from 10 samples and 4 Mbp sequence lengths.

The 45 pairwise comparisons totaling 180 Mbp with 20 Kbp windows yielded 9,000 pairs of

TMRCA values per matrix. We detected a population bottleneck ~100,000 years ago (100 kya)

in CEU and IBnr using SNPs (Figure 5B), which was consistent with the findings of a previous

study [22]. This signal was not clearly detected with SMPs (Figure 5C), indicating their limited use for ancient events. Both SNPs and SMPs indicated another bottleneck between 1-10 kya (Figure 5B-C), but SMPs also revealed a subsequent population expansion (Figure 5C) undetected by SNPs (Figure 5B). The population contraction began earlier in IBnr, followed by a slightly stronger expansion compared to CEU (Figure 5C), which may explain its more negative Tajima's *D* estimate for methylation variants (Figure 3B). Using SMPs, we recovered two bottlenecks (~100 kya and between 1-10 kya) in *B. distachyon* (Figure 5D) that were previously reported using SNPs for samples combined from multiple clonal groups in this species [34]. The steeper expansion of IBnr compared to CEU (*A. thaliana*) and Group 1.1 compared to Group 1.2 (*B. distachyon*) in the last 10,000 years likely led to the observed excess of rare alleles (Figures 2B, 3F).

We validated our demographic inference with two approaches. First, we simulated the sawtooth demography observed in both species, confirming that our deep learning method accurately detected multiple bottleneck and expansion events (Figure 5E). Second, we simulated methylation markers with high rates of change ($10^{-4}$, $10^{-5}$, or $10^{-6}$ per site per generation) and backward rates 2.5 times larger than forward rates, based on the estimates provided in [6], to test whether deviations from the infinite sites model affect inference. We found that our method accurately recovered demographic histories beyond 0.6 kya and before 5, 22, and 442 kya for methylation rates of $10^{-4}$, $10^{-5}$, and $10^{-6}$ per site per generation, respectively (Figure S13). Conservatively, this suggests that inferences 20 kya and beyond are unlikely to be accurate for CEU and IBnr due to their estimated rates of methylation change being around $10^{-5}$ per site per generation (Figure S8). Note that the demographic estimates with SNPs are likely accurate for older events (>20kya). Nonetheless, we are confidently estimating recent population expansion with SMPs (Figure 5C), highlighting the potential of methylation markers to reveal insights beyond what genetic variants alone can provide.

**Discussion**

We observed contrasting patterns in the capacity of methylation variants to distinguish population subdivisions in *A. thaliana* and *B. distachyon*. In *A. thaliana*, the infinite sites model is likely violated for methylation, resulting in the erosion of methylome differentiation likely due to the older divergence between populations. In *B. distachyon*, samples from different altitudes with varying seed morphologies exhibited distinct methylation profiles in a common garden, despite their genetic similarity. While SMP distributions are similar across different genomic contexts, DMRs are more context dependent, making the former preferable for population-scale analyses. We found weak negative correlations between π and no local correlations for Tajima's *D* between genetic variants and methylation variants in *A. thaliana*, underscoring their distinct molecular and evolutionary properties, similar to the conclusions reached by Schmitz *et al.* (2013) [36]. We found slightly stronger linkage between neighboring SMPs compared to a previous estimate in wild *A. thaliana* sampled across the Northern Hemisphere [36], likely due to our focus on individual populations with less genetic structure [22]. SMPs and SNPs recover recent and older demographic changes, respectively, providing a comprehensive reconstruction of population evolutionary histories.

The differing ability to detect population subdivisions in *A. thaliana* and *B. distachyon* using methylation variants could lead to differences in divergence times, methylation rates, or local adaptation. As the post-glacial re-colonization of the Iberian Peninsula and the spread northward started ~16 kya [29], multiple changes at the same methylation site (homoplasy) during this period may have caused an underestimation of the true levels of divergence in *A. thaliana*. The greater capacity to detect group subdivision is *B. distachyon* could reflect a recent separation by altitude contributing to methylome divergence. We indeed rule out the role of adaptation such as found in *F. vesca* [17], because we generated inferences using gbM regions that are highly conserved in angiosperms [11]. The observed divergence could be due to low rates of methylation change, but we could not estimate methylation rates for *B. distachyon* as our rate estimate relies on comparing the distributions of genetic and

methylation variants. The most parsimonious explanation is that the two subgroups diverged

recently. As the B_East lineage split from the western Mediterranean lineage 23 kya [34, 35],

the split between the closely related subgroups is likely to have been far more recent, ensuring

that the infinite sites model was not violated for methylation variants. Our findings underscore

the necessity for methods distinguishing neutral demographic changes from selective

pressures on the methylome, elucidating why SMPs behave differently across species.

Our results support the use of individual CG sites from all genomic contexts for population-

scale analyses. Our between-population comparisons using CG sites in gbM regions

successfully identified the known population structure within *B. distachyon*, similar to a

previous study using genic CG sites [16]. Further supporting this conclusion is the consistent

similarity in the mSFS shape when using SMPs, regardless of their genomic context. The *A.*

*thaliana* [21] and *B. distachyon* [16] methylomes, generated under similar environmental

conditions, ensure that sites affected solely by transient environmental effects would not be

recognized as SMPs. Another study using all CG sites from lettuce methylomes, cultivated

under uniform environmental conditions, also demonstrated the capacity to distinguish

multiple species [19]. Some noise can be introduced by analyzing the whole methylome as

non-gbM regions show a slight skew towards variants with intermediate frequencies, but this

is likely to be offset by the increased number of sites for inference. The distribution of DMR

states is tied to the genomic context. gbM results from the combined activity of MET1, which

is enhanced by nearby existing methylation, as well as the histone variant H2A.Z and the DNA

demethylase ROS1 [37]. The molecular basis underlying the clock-like, non-gbM regions, as

well as regions with high and low SIFT scores, remains unknown. However, given that our

mSFS for these regions differ, this suggests potential for differences in their molecular basis.

Studying DMRs can nevertheless be valuable for imputing methylation states when

sequencing depths are low [38], while the impact of such imputation on population genetics

inference needs to be evaluated. Overall, using a restricted set of CG sites may be more

appropriate when analyzing accessions grown under variable environments together or for

tracking long-term methylome divergence [10].

The weak local correlations between genetic and methylation diversity and the association

between gene constraint and methylation diversity in *A. thaliana* can partly be attributed to

methylation affecting the mutability of neighboring sites [39-41] and transposable element

polymorphisms influencing methylation variation [42, 43]. However, only about 12% of

methylation quantitative trait loci (QTLs) in *A. thaliana* are within 10Kbp of the epiallelic target

[36]. In maize, approximately 50% of methylation QTLs occur in *cis* [44], indicating a potentially

stronger trend in this species. The higher rate of methylation changes compared to mutation

accumulation [6, 7] may also explain why methylation differences persist despite genetic

homogeneity. Our simulations also show that coalescent histories of markers with vastly

different rates of change diverge, which can explain the general lack of correlation between π

and Tajima's *D* for local genetic and methylation diversity. Our results are similar to those of

previous studies in lettuce [19] and cotton [45], which also found higher levels of polymorphism

in the methylome compared to the genome. The lack of correlation or similarity between the

Tajima's *D* estimates from the genome and methylome in *A. thaliana* and *B. distachyon*

supports the notion that the SNPs and SMPs likely reflect different parts of the common

underlying coalescent genealogy at a locus: SNPs reflect the older part of the tree, whereas

SMPs would reflect the topology and branch lengths in the more recent past. It is then not

surprising that for recent positive selection events (selective sweeps), the signatures are

similar for both types of markers, and SMPs may even be more accurate in detecting them

[15]. Our study mostly focuses on local correlations in diversity, rather than genome-wide

correlations, where life history traits and genetic differentiation levels also play a significant

role [46-52].

Our study demonstrates that methylation variants can independently detect and allow

inference of historical demographic changes. This represents a significant improvement over

a previous approach [8], where methylation was used only as a secondary marker to

supplement genetic variant-based inferences. Furthermore, our improved deep learning

method does not require precise methylation rate parameters and is less sensitive to the

presence of invariant methylation sites (compared to the SMCm in [8]). Instead, it trains on

diverse scenarios to match observed variant distributions. For *B. distachyon*, methylation

variants revealed expansions also detected by genetic variants [34]. In *A. thaliana*, genetic

variant-based demographic inferences fail to detect recent population expansions (this study

and [53]), which can be uncovered through methylation variants instead. Our simulations

confirmed the utility of rapidly mutating markers for inferring demographic changes. Overall,

our study highlights the value of methylation markers for understanding recent shifts in

population size.

**Conclusions**

Our findings demonstrate that CG methylation can provide new insights into population

structure in clonal and selfing species, similar to its previously demonstrated utility in *A.*

*thaliana* from America and the clonal seagrass *Zostera marina* [10]. We demonstrate the utility

of methylation variants for uncovering very recent demographic events, where SNP markers

are limited due to their slower rate of change. We also illustrate its usefulness in clonal groups,

where the paucity of SNPs would render such inferences either impossible or necessitate very

deep and costly levels of DNA sequencing. Conversely, at older timescales, methylation

markers are unlikely to be informative because the high rates of change would have resulted

in the violation of the infinite sites model. We show that SNPs and SMPs are complementary

for generating demographic inferences over short and long timescales, highlighting the value

of a recently developed method that leverages both sources of variation for inferring

demography [8]. Our work demonstrates that methylomes contain unique evolutionary insights

unattainable through studying genomic variation in isolation.

**Methods**

*Genetic and methylation variant calling in Arabidopsis thaliana*

We obtained the *A. thaliana* 1001 genomes [30] SNP calls from https://1001genomes.org/data/GMI-MPI/releases/v3.1/. We only retained biallelic SNPs for our analyses. As C/T variant calls can become confounded with methylation calls for data generated using bisulfite sequencing, we generated sample-specific genome references for mapping bisulfite sequencing reads. We substituted the genetic variant calls for each sample into the TAIR10 genome reference [54] using SAMtools [55], BCFtools consensus [55], and fastq split [56]. We obtained bisulfite sequencing data for the 1001 *A. thaliana* from [21]. We first removed poor-quality reads using Trimmomatic [57] with the following options: ILLUMINACLIP:TruSeq3-SE.fa:2:30:10 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:36 for single-end reads and ILLUMINACLIP:TruSeq3-PE.fa:2:30:10:2:True LEADING:3 TRAILING:3 SLIDINGWINDOW:4:20 MINLEN:36 for paired-end reads. We subsequently mapped reads to the sample-specific reference genomes using Bismark with the aid of the HISAT2 aligner [58]. We ran Bismark deduplicate_bismark to remove optical duplicates and ran bismark_methylation_extractor to extract methylation calls. We studied cytosine sites with a coverage of at least five reads similar to what was done in [21]. To identify the methylation state of a site, we performed a binomial test with the non-conversion rate being used as the hypothesized probability of success following the method in [59]. The non-conversion rate was calculated as the proportion of methylation in the chloroplast as done in [59]. Sites were assigned as methylated if the false discovery rate corrected *P*-values following the binomial test was less than 0.01. Otherwise, they were designated as unmethylated. We used jDMR [60] to identify DMRs in the CG contexts as this method accounts for coverage when calculating likelihood scores for methylation states. For this, we called the methylation status per site using METHimpute (function: callMethylationSeparate) [38], extracted cytosine clusters in the CG context ensuring that there was a minimum of eight cytosines in each cluster (function: CfromFASTAv4 and makeReg with default options makeRegnull = c(FALSE), N.boot

= 10^5, N.sim.C = "all", fp.rate = 0.01, set.tol = 0.01), and identified the methylation states of

the regions in each sample (function: runMethimputeRegions), ensuring that we used a

minimum of eight cytosines and a coverage of three.

63 IBnr and 67 CEU samples overlapped between the SNP and methylation variant call files.

We randomly subsampled 63 samples from the CEU population to ensure the numbers were

comparable to those of the IBnr population. For our analyses, we only used sites where at

least 80% of the samples within each population were genotyped. The proportions of variant

SNP and SMP sites were calculated by dividing the number of sites with minor allele

frequencies greater than 0.02 by the total number of variant and invariant sites.

*Population (epi)genetic metric estimation for Arabidopsis thaliana*

To estimate $\pi$ and Tajima's $D$, we created fasta files for each genomic region containing the

sequences for each sample. We substituted the variants from the SNP vcf into the TAIR10

reference genome using the approach described previously for each sample, extracted the

region of interest, and converted it to a fasta format. The regions could be the clocklike regions

described in [10], genic regions comprising all exons, or non-coding regions comprising all

regions other than exons. Genic regions were further split into gbM genes using the list for *A.*

*thaliana* provided by [61] and high SIFT-score (>0.75) and low SIFT-score (<0.3) genes using

the list provided by [62].

We estimated $\pi_{SNP}$ and $D_{SMP}$ using pegas [63] and estimated $\pi_{SMP}$ and $D_{SMP}$ using the $D^m$ [28].

We only analyzed intervals that had at least three polymorphic sites and sequences with

lengths greater than 10bp. We calculated the number of cytosines in each interval using

METHimpute [38]. To compare $\pi$ and Tajima's $D$ among the different genomic contexts, we

built a linear model where these metrics were the dependent variable, and the concatenated

population and genomic context groupings were the predictor variables. We then applied a

Tukey    honest    significant    differences    test    using    the    agricolae    R    package

(https://myaseen208.com/agricolae//index.html) to identify significant differences between

individual predictor levels. We performed a correlation test to compare the π and Tajima's *D*

for SMPs and SNPs per exon.

We used VCFtools [64] to estimate the allele frequencies for SMPs and DMRs. We used

PopLDdecay [65] to estimate pairwise LD using the $r^2$ metric analyzing all sites with a minor

allele frequency greater than 0.005.

*Between population comparisons for Arabidopsis thaliana*

We combined the CEU and IBnr vcfs and used VCF2Dis (https://github.com/BGI-

shenzhen/VCF2Dis) to estimate the p-distance matrix. We used this matrix to generate

neighbor-joining trees using FastME [66] and plotted it using ggtree [67]. We also performed

DAPC to discriminate samples from the CEU and IBnr populations. We thinned the dataset so

that only sites that were separated by 1Kbp were sampled to reduce the computational time.

We converted vcf files into a genlight format using the *genomic_converter* function in the R

package radiator (https://thierrygosselin.github.io/radiator/). We used the find.clusters function

in adegenet [68] to identify the number of clusters in each dataset allowing for a maximum of

20 clusters for SMPs/DMRs and 30 clusters for SNPs. As DAPC regions require fully typed

data, we imputed missing values using the mean value. Then, we performed cross-validation

using adegenet [68] to identify the number of informative principal components. We retained

a maximum of 15 principal components, used 90% of the data for training, and calculated

group-wise assignment success. We centered but did not scale variables. We used 30

replicates for each principal component. Finally, we performed DAPC using adegenet [68].

Methylation rate estimation *for Arabidopsis thaliana*

We generated the multihetsep files for SNPs for the same 5Mbp regions that were used for

SMPs (described in the section immediately above). We combined the SNP and SMP

multihetsep files. We randomly selected 20 samples from each population to ensure

reasonable runtimes. We used the *Methylation_rate_estimation* function in SMCm to estimate

forward and backward rates [8]. We set the mutation rate as $7\times10^{-9}$ per site per generation

and the recombination rate as $3.5\times10^{-8}$ per site per generation. We ignored regional influences

on methylation rates and estimated rates under a site-only model. We used the known

estimates of forward ($2.56\times10^{-4}$ per generation per haploid methylome) and backward

methylation rates ($6.3\times10^{-4}$ per generation per haploid methylome) for *A. thaliana* [6] as

priors.

*Genetic and methylation variant calling in Brachypodium distachyon*

We obtained whole genome sequencing data (29 samples) for *B. distachyon* from [27] and

whole genome bisulfite sequencing data from [16]. We used Bd21-3 v1.2 genome reference

and the GenBank annotation from the Joint Genome Institute (http://phytozome.jgi.doe.gov/)

for variant calling. We used Trimmomatic [57] to trim reads using the paired-end options as

stated above for the whole genome sequencing data and the single-end option for the whole

genome bisulfite sequencing data. We mapped whole genome sequencing data reads to the

reference using bwa-mem [69]. We used Picard Toolkit (https://broadinstitute.github.io/picard/)

to add read groups and mark duplicate reads. We used SAMtools [55] to remove unpaired

reads, remove reads with a mapping quality of less than 20, and sort reads. We called genetic

variants using GATK HaplotypeCaller [70]. We only retained SNPs with depths greater than

five and quality scores greater than 30 using BCFtools [55]. Using this SNP variant call file,

we perform methylation variant calling in the same way as was done for *A. thaliana*. We

obtained the sample localities from [32] and created a map showing their occurrence with Map

Maker (https://maps.co/).

*Within and between population analyses in Brachypodium distachyon*

We used a minor allele frequency threshold of 10% to identity polymorphic sites in each clonal

group due to the smaller number of samples being analyzed when compared to *A. thaliana*.

For the analysis of SMPs and DMRs, we only used variants that occurred within CG-gbM

regions identified in [71]. We estimated mSFS, LD, p-distance matrices, π, and Tajima's *D* in

a similar way as we did for *A. thaliana*. We thinned the SMP dataset so that only sites that

were separated by 3Kbp were sampled to reduce the computational time for the DAPC

analyses.

*Generating the approximated transition matrix for demographic inference*

We created a training and testing dataset consisting of a combined $60 \times 10^3$ simulations

(80%/20%, respectively). Of these $10 \times 10^3$ were constant demographies (*i.e.*, no demographic

change) with a random population size drawn between $10 \times 10^3$ and $200 \times 10^3$, and the other

$50 \times 10^3$ simulations were random walk demographies, with time steps being defined for i ∈

{0,1,...,K − 1} using:

$$(\text{population\_time})_i = \frac{\exp\left(\frac{i}{K-1} \cdot \ln(1 + \delta t_{max})\right) - 1}{\delta}$$

where *K*=42 is the number of time points, $\delta = 0.001$ is the growth rate, and $t_{max} = 2.1 \times 10^6$ is the

maximum time.

The population size was drawn for each of the *K*=42 time points in the following way: an initial

population size was randomly selected between a minimum (*N_min* = $10 \times 10^3$) and maximum

(*N_max* = $1.2 \times 10^6$) value. For each subsequent time point, a new population size was

proposed by multiplying the previous size by a random factor between 0.1 and 10. If this

proposed size fell outside the allowed range (N_*min* to *N_max*), new proposals were

generated until an acceptable value was found. This process was repeated for all 42 time

points, resulting in a sequence of population sizes that exhibit both gradual and abrupt

changes while remaining within the specified bounds. Nevertheless, a sliding window average

with a size of 6 and a step size of 1 was applied to smooth extreme changes. This approach

for sampling demographies was adapted from Sanchez *et al.* [72].

The obtained population sizes and times were passed onto *msprime* [73] to simulate the

corresponding ancestral recombination graphs using a recombination rate of $3.4 \times 10^{-8}$ per

base pair and generation. For SNPs, branches were mutated at a rate of $7 \times 10^{-9}$ per base pair

and generation, leading to a $\rho$ over $\theta$ ratio of 4.86.

Simulated and real data are summarized as a genotype matrix (*GM*) with dimensions $n \times m$,

where $n,m \in \mathbb{N}$ represent the number of samples and mutations, respectively. Each element

$g_{ij}$ of *GM*, where $i \in \{1,...,n\}$ and $j \in \{1,...,m\}$, denotes the presence ($g_{ij} = 1$) or absence ($g_{ij} = 0$)

of mutation *j* in sample *i*. We generated a corresponding vector of positions of size *m* for each

segregating site containing floating point numbers of the location in the genome.

For each random population size parameter, we used a genome of length 4Mbp,10 samples

(ploidy=1), and the *GM* and positions vector to construct a density estimate of mutations along

the genome (in theory, corresponding to the respective TMRCA at that region). To ensure the

presence of enough markers, we segmented the genome into 20Kbp pieces and counted the

number of heterozygous markers between all pairwise $\binom{10}{2}$ combinations. The 4Mbp

sequences of all the comparisons were concatenated and log-transformed. Likewise, the

population time steps were also log-scaled. To make sure that the number of markers per

window fell approximately into the time windows, we scaled the markers by an arbitrarily

chosen scaling factor ($=e^{8.2}$) (Figure S13). The importance of the given scaling factor is

negligible, due to all the transformation steps occurring in the pre-training, and therefore any

noisy representation presented to the learning algorithm will be part of the training.

Each *msprime* simulation (under a random demographic history) yielded one *approximated*

transition matrix, which was obtained by counting the transitions (scaled number of

heterozygous sites in window t → window t+1) from one 20 Kbp window to the next using the

following algorithm: a sliding window of size 2 and step size 1 was applied along the sequence

and each pair of consecutive values in this window represents coordinates in the transition

matrix, which was incremented at those positions leading to a filled matrix which remained

unnormalized. Lastly, the entire transition matrix was incremented by one at each entry and

scaled by the natural log to squeeze all values approximately into the same range to avoid

training instability.

*Model description, training, and inference*

The inference approach can be described as a supervised learning approach using the

*approximated* transition matrices *x* with defined dimensions $R^{L=42 \times K=42}$ as the input and the

population size at each time step as the predicted variable. The input is a square matrix where

both dimensions (K and L) are 42. The model architecture is an encoder-only transformer [74]

(https://github.com/lucidrains/x-transformers). It consists of three main components: an input

projection layer, a transformer encoder, and an output projection layer. It can be formalized

as:

1. Input Projection: $x_{proj} = W_{in} x + b_{in}$, where $W_{in} \in R^{D_{enc} \times K_{in}}, b_{in} \in R^{D_{enc}}, x \in R^{L \times K_{in}}$

2. Transformer Encoder: $z = \text{Encoder}(x_{proj})$, where $z \in R^{L \times D_{enc}}$

The Encoder employs a configuration with dimension $D_{enc}$ = 512, depth $N_{enc}$ = 12, number

of attention heads $H_{enc}$ = 10 heads, Gated Linear Unit activation in feed-forward layers [75],

residual attention connections [76], and rotary positional embeddings [77].

3. Dimension Reduction: $z_{reduced} = \sum_{i=1}^{L} z_i$ , where $z_{reduced} \in R^{D_{enc}}$

4. Output Projection: $y = W_{out} z_{reduced} + b_{out}$, where $W_{out} \in R^{D_{out} \times D_{enc}}, b_{out} \in R^{D_{out}}$

In this formalization, sequence length $L$ = the input dimension of the transition matrix $K_{in}$ =

42, which represents the square input matrix of 42 x 42 dimensions, and the output

dimension $D_{out}$ = 42.

The neural network model underwent training for 50 epochs, employing mean squared error

as the loss function. The training process leverages the Accelerate framework from Hugging

Face to enhance performance [78]. An Accelerator object was configured with mixed precision

using bfloat16 and gradient accumulation over 12 steps (with batch size 32). Throughout the

training regiment, both the training and test losses were continuously monitored to evaluate

model performance and assess convergence.

*Generating data for demographic inference*

We first generated unique variant call files for each sample from the population variant call

files using BCFtools [55]. We then used the generate_multihetsep.py from MSMC-tools

(https://github.com/stschiff/msmc-tools/tree/master) to combine all the samples per

chromosome. For *A. thaliana* SNPs, we used the entire chromosome 1 as we needed a long

contiguous fragment to generate the approximated transition matrix. For SMPs, we used 5Mbp

segments in each of the five chromosomes (Table S2) as the SMP density was far higher. We

used the list of genotyped cytosine positions as a genomic location mask for the SMPs. For

*B. distachyon*, we included singletons to support the demographic inference given the small

sample sizes in this species. The data were polarized by identifying the minor (less frequent)

alleles and encoding them as derived, whereas the major alleles were encoded as ancestral.

The resulting genotype matrix represents this polarized, unphased data, where each row

corresponds to an individual and each column corresponds to a genomic position.

*Methylation simulation model*

We explored the boundaries of our course-grained demographic inference by adding a new

mutation model to msprime [73]. We simulated a sawtooth demographic model with cyclical

population size changes over time and a baseline effective population size of $2\times10^5$

individuals. Simulations were conducted on a 5Mbp sequence with a recombination rate of

$3.4\times10^{-8}$ per base pair per generation (ploidy=1), to isolate the influence of methylation

dynamics.

A custom matrix mutation simulation was implemented to model methylation and

demethylation processes, employing 2 allelic states: 0 (unmethylated) and 1 (methylated). The

model incorporates 3 key rates: the site introduction rate ($\mu$), methylation rate ($\beta$), and

demethylation rate ($\delta$). The methylation and demethylation rates are measured as factors of

the site entry rate, where $\beta$ = the methylation factor·$\mu$ and $\delta$ = the demethylation factor·$\mu$. The

transition probabilities between the unmethylated and methylated states are determined by

the ratio $\beta/(\beta+\delta)$ for methylation and $\delta/(\beta+\delta)$ for demethylation. The resulting 2×2 transition

matrix encapsulates these probabilities:

790
$$\begin{bmatrix} 1 - [\beta/(\beta + \delta)] & \beta/(\beta + \delta) \\ \delta/(\beta + \delta) & 1 - [\delta/(\beta + \delta)] \end{bmatrix}$$

This framework allowed us to explore various methylation scenarios by adjusting $\beta$ and $\delta$

relative to $\mu$. We investigated cases where $\beta = \mu$ and $\delta = 2.5\mu$, $\beta = 10\mu$ and $\delta = 25\mu$, and $\beta =$

$100\mu$ and $\delta = 250\mu$, representing low, medium, and high methylation dynamics respectively,

with the demethylation rate 2.5 times higher than the methylation rate. By varying the

epimutation rate compared to the mutation rate, we test the robustness of our approach in

correlating the number of markers in one Kbp windows with the average coalescent time. This

proportionality, and the robustness of our method, is expected to decrease for higher

epimutation rates, as these would violate the infinite site model for older coalescent times more

strongly (due to homoplasy, [4]).

**Declarations**

*Ethics approval and consent to participate*

Not applicable

*Consent for publication*

Not applicable

*Availability of data and materials*

The scripts for analyzing and plotting all the results, except for the demographic inference,

have been published on GitHub (arunkumarramesh/methevolve) and archived using Zenodo

(https://doi.org/10.5281/zenodo.13991812). The demographic inference method is available

on GitHub (kevinkorfmann/aTMi) and has been archived using Zenodo

(https://doi.org/10.5281/zenodo.13332833).

*Competing interests*

The authors declare that they have no competing interests.

*Funding*

RA was funded by the Peter and Traudl Engelhorn Foundation.

*Authors' contributions*

RA and AT designed the research. RA, KK, AW, BH, CG, and YW performed the research. RA,

AT, KK, and AW wrote the paper. All authors read and approved the final manuscript.

*Acknowledgements*

We would like to thank Thibaut Paul Patrick Sellinger for helping with the estimation of

methylation rates. This work was supported by the de.NBI Cloud within the German Network

for Bioinformatics Infrastructure (de.NBI) and ELIXIR-DE (Forschungszentrum Jülich and W-

de.NBI-001, W-de.NBI-004, W-de.NBI-008, W-de.NBI-010, W-de.NBI-013, W-de.NBI-014, W-

de.NBI-016, W-de.NBI-022).

**References**

1. Richards CL, Alonso C, Becker C, Bossdorf O, Bucher E, Colome-Tatche M, Durka

W, Engelhardt J, Gaspar B, Gogol-Doring A, et al: **Ecological plant epigenetics:**

**Evidence from model and non-model species, and the way forward.** *Ecol Lett*

2017, **20:**1576-1590.

2. Feng S, Cokus SJ, Zhang X, Chen PY, Bostick M, Goll MG, Hetzel J, Jain J, Strauss

SH, Halpern ME, et al: **Conservation and divergence of methylation patterning**

**in plants and animals.** *Proc Natl Acad Sci U S A* 2010, **107:**8689-8694.

3. Zhang H, Lang Z, Zhu JK: **Dynamics and function of DNA methylation in plants.**

*Nat Rev Mol Cell Biol* 2018, **19:**489-506.

4. Vidalis A, Zivkovic D, Wardenaar R, Roquis D, Tellier A, Johannes F: **Methylome**

**evolution in plants.** *Genome Biol* 2016, **17:**264.

5. Cortijo S, Wardenaar R, Colome-Tatche M, Gilly A, Etcheverry M, Labadie K,

Caillieux E, Hospital F, Aury JM, Wincker P, et al: **Mapping the epigenetic basis**

**of complex traits.** *Science* 2014, **343:**1145-1148.

6.   van der Graaf A, Wardenaar R, Neumann DA, Taudt A, Shaw RG, Jansen RC,

Schmitz RJ, Colome-Tatche M, Johannes F: **Rate, spectrum, and evolutionary**

**dynamics of spontaneous epimutations.** *Proc Natl Acad Sci U S A* 2015,

**112:**6676-6681.

7.   Ossowski S, Schneeberger K, Lucas-Lledo JI, Warthmann N, Clark RM, Shaw RG,

Weigel D, Lynch M: **The rate and molecular spectrum of spontaneous**

**mutations in Arabidopsis thaliana.** *Science* 2010, **327:**92-94.

8.   Sellinger T, Johannes F, Tellier A: **Improved inference of population histories by**

**integrating genomic and epigenomic data.** *eLife* 2023, **12:**RP89470.

9.   Yao N, Schmitz RJ, Johannes F: **Epimutations Define a Fast-Ticking Molecular**

**Clock in Plants.** *Trends Genet* 2021, **37:**699-710.

10.  Yao N, Zhang Z, Yu L, Hazarika R, Yu C, Jang H, Smith LM, Ton J, Liu L, Stachowicz

JJ, et al: **An evolutionary epigenetic clock in plants.** *Science* 2023, **381:**1440-

1445.

11.  Muyle AM, Seymour DK, Lv Y, Huettel B, Gaut BS: **Gene Body Methylation in**

**Plants: Mechanisms, Functions, and Important Implications for**

**Understanding Evolutionary Processes.** *Genome Biol Evol* 2022, **14:**evac038.

12.  Charlesworth B, Jain K: **Purifying selection, drift, and reversible mutation with**

**arbitrarily high mutation rates.** *Genetics* 2014, **198:**1587-1602.

13.  Muyle A, Ross-Ibarra J, Seymour DK, Gaut BS: **Gene body methylation is under**

**selection in Arabidopsis thaliana.** *Genetics* 2021, **218:**iyab061.

14.  Xu G, Lyu J, Li Q, Liu H, Wang D, Zhang M, Springer NM, Ross-Ibarra J, Yang J:

**Evolutionary and functional genomics of DNA methylation in maize**

**domestication and improvement.** *Nat Commun* 2020, **11:**5539.

15.   Shirai K, Sato MP, Nishi R, Seki M, Suzuki Y, Hanada K: **Positive selective**

**sweeps of epigenetic mutations regulating specialized metabolites in**

**plants.** *Genome Res* 2021, **31:**1060-1068.

16.   Eichten SR, Srivastava A, Reddiex AJ, Ganguly DR, Heussler A, Streich JC, Wilson

PB, Borevitz JO: **Extending the Genotype in Brachypodium by Including DNA**

**Methylation Reveals a Joint Contribution with Genetics on Adaptive Traits.**

*G3 (Bethesda)* 2020, **10:**1629-1637.

17.   Le Vève A, Sammarco I, Latzel V, Lafon Placette C: **DNA methylation shows**

**footprints of altitude selection in the clonal plant species Fragaria vesca.**

*bioRxiv* 2024, **2024.03.19.585697**.

18.   Sammarco I, Diez Rodriguez B, Galanti D, Nunn A, Becker C, Bossdorf O,

Munzbergova Z, Latzel V: **DNA methylation in the wild: epigenetic**

**transgenerational inheritance can mediate adaptation in clones of wild**

**strawberry (Fragaria vesca).** *New Phytol* 2024, **241:**1621-1635.

19.   Cao S, Sawettalake N, Li P, Fan S, Shen L: **DNA methylation variations underlie**

**lettuce domestication and divergence.** *Genome Biol* 2024, **25:**158.

20.   Rodriguez-Izquierdo A, Carrasco D, Anand L, Magnani R, Catarecha P, Arroyo-

Garcia R, Rodriguez Lopez CM: **Epigenetic differences between wild and**

**cultivated grapevines highlight the contribution of DNA methylation during**

**crop domestication.** *BMC Plant Biol* 2024, **24:**504.

21.   Kawakatsu T, Huang SC, Jupe F, Sasaki E, Schmitz RJ, Urich MA, Castanon R,

Nery JR, Barragan C, He Y, et al: **Epigenomic Diversity in a Global Collection of**

**Arabidopsis thaliana Accessions.** *Cell* 2016, **166:**492-505.

22. Strütt S, Sellinger T, Glemin S, Tellier A, Laurent S: **Joint inference of**

**evolutionary transitions to self-fertilization and demographic history using**

**whole-genome sequences.** *Elife* 2023, **12:**e82384.

23. Vaswani A, Shazeer N, Parmar N, Uszkoreit J, Jones L, Gomez AN, Kaiser L,

Polosukhin I: **Attention Is All You Need.** *arXiv* 2017, **1706.03762**.

24. Devlin J, Chang M-W, Lee K, Toutanova K: **BERT: Pre-training of Deep**

**Bidirectional Transformers for Language Understanding.** *arXiv* 2018,

**1810.04805**.

25. Korfmann K, Gaggiotti OE, Fumagalli M: **Deep Learning in Population Genetics.**

*Genome Biol Evol* 2023, **15**.

26. Becker C, Hagmann J, Muller J, Koenig D, Stegle O, Borgwardt K, Weigel D:

**Spontaneous epigenetic variation in the Arabidopsis thaliana methylome.**

*Nature* 2011, **480:**245-249.

27. Wilson PB, Streich JC, Murray KD, Eichten SR, Cheng R, Aitken NC, Spokas K,

Warthmann N, Gordon SP, Accession C, et al: **Global Diversity of the**

**Brachypodium Species Complex as a Resource for Genome-Wide**

**Association Studies Demonstrated for Agronomic Traits in Response to**

**Climate.** *Genetics* 2019, **211:**317-331.

28. Wang J, Fan C: **A neutrality test for detecting selection on DNA methylation**

**using single methylation polymorphism frequency spectrum.** *Genome Biol*

*Evol* 2014, **7:**154-171.

29. Hewitt G: **Post-glacial re-colonization of European biota.** *Biological Journal of*

*the Linnean Society* 1999, **68:**87-112.

30.  The1001GenomesConsortium: **1,135 Genomes Reveal the Global Pattern of**

**Polymorphism in Arabidopsis thaliana.** *Cell* 2016, **166:**481-491.

31.  Sharbel TF, Haubold B, Mitchell-Olds T: **Genetic isolation by distance in**

**Arabidopsis thaliana: biogeography and postglacial colonization of Europe.**

*Mol Ecol* 2000, **9:**2109-2118.

32.  Filiz E, Ozdemir BS, Budak F, Vogel JP, Tuna M, Budak H: **Molecular,**

**morphological, and cytological analysis of diverse Brachypodium**

**distachyon inbred lines.** *Genome* 2009, **52:**876-890.

33.  Ng PC, Henikoff S: **SIFT: Predicting amino acid changes that affect protein**

**function.** *Nucleic Acids Research* 2003, **31:**3812-3814.

34.  Minadakis N, Williams H, Horvath R, Caković D, Stritt C, Thieme M, Bourgeois Y,

Roulin AC: **The demographic history of the wild crop relative Brachypodium**

**distachyon is shaped by distinct past and present ecological niches.** *Peer*

*Community Journal* 2023, **3:**e84.

35.  Stritt C, Gimmi EL, Wyler M, Bakali AH, Skalska A, Hasterok R, Mur LAJ, Pecchioni

N, Roulin AC: **Migration without interbreeding: Evolutionary history of a highly**

**selfing Mediterranean grass inferred from whole genomes.** *Mol Ecol* 2022,

**31:**70-85.

36.  Schmitz RJ, Schultz MD, Urich MA, Nery JR, Pelizzola M, Libiger O, Alix A, McCosh

RB, Chen H, Schork NJ, Ecker JR: **Patterns of population epigenomic diversity.**

*Nature* 2013, **495:**193-198.

37.  Briffa A, Hollwey E, Shahzad Z, Moore JD, Lyons DB, Howard M, Zilberman D:

**Millennia-long epigenetic fluctuations generate intragenic DNA methylation**

**variance in Arabidopsis populations.** *Cell Syst* 2023, **14:**953-967 e917.

38.    Taudt A, Roquis D, Vidalis A, Wardenaar R, Johannes F, Colome-Tatche M:

**METHimpute: imputation-guided construction of complete methylomes**

**from WGBS data.** *BMC Genomics* 2018, **19:**444.

39.    Xia J, Han L, Zhao Z: **Investigating the relationship of DNA methylation with**

**mutation rate and allele frequency in the human genome.** *BMC Genomics*

2012, **13:**S7.

40.    Mugal CF, Ellegren H: **Substitution rate variation at human CpG sites**

**correlates with non-CpG divergence, methylation level and GC content.**

*Genome Biol* 2011, **12:**R58.

41.    Kusmartsev V, Drozdz M, Schuster-Bockler B, Warnecke T: **Cytosine Methylation**

**Affects the Mutability of Neighboring Nucleotides in Germline and Soma.**

*Genetics* 2020, **214:**809-823.

42.    Stuart T, Eichten SR, Cahn J, Karpievitch YV, Borevitz JO, Lister R: **Population**

**scale mapping of transposable element diversity reveals links to gene**

**regulation and epigenomic variation.** *Elife* 2016, **5:**e20777.

43.    Shen Y, Zhang J, Liu Y, Liu S, Liu Z, Duan Z, Wang Z, Zhu B, Guo YL, Tian Z: **DNA**

**methylation footprints during soybean domestication and improvement.**

*Genome Biol* 2018, **19:**128.

44.    Eichten SR, Briskine R, Song J, Li Q, Swanson-Wagner R, Hermanson PJ, Waters

AJ, Starr E, West PT, Tiffin P, et al: **Epigenetic and genetic influences on DNA**

**methylation variation in maize populations.** *Plant Cell* 2013, **25:**2783-2797.

45.    Zhao T, Guan X, Hu Y, Zhang Z, Yang H, Shi X, Han J, Mei H, Wang L, Shao L, et al:

**Population-wide DNA methylation polymorphisms at single-nucleotide**

resolution in 207 cotton accessions reveal epigenomic contributions to

complex traits. *Cell Res* 2024.

46.  Avramidou EV, Korakaki E, Malliarou E, Boutsios S: **Studying the Genetic and**

**the Epigenetic Diversity of the Endangered Species Juniperus drupacea**

**Labill. towards Safeguarding Its Conservation in Greece.** *Forests* 2023,

**14:**1271.

47.  Avramidou EV, Moysiadis T, Ganopoulos I, Michailidis M, Kissoudis C, Valasiadis

D, Kazantzis K, Tsaroucha E, Tsaftaris A, Molassiotis A, et al: **Phenotypic,**

**Genetic, and Epigenetic Variation among Diverse Sweet Cherry Gene Pools.**

*Agronomy* 2021, **11:**680.

48.  Wang MZ, Li HL, Li JM, Yu FH: **Correlations between genetic, epigenetic and**

**phenotypic variation of an introduced clonal herb.** *Heredity (Edinb)* 2020,

**124:**146-155.

49.  Langford N, Fargeot L, Blanchet S: **Spatial covariation between genetic and**

**epigenetic diversity in wild plant and animal populations: a meta-analysis.** *J*

*Exp Biol* 2024, **227**.

50.  Medrano M, Alonso C, Bazaga P, Lopez E, Herrera CM: **Comparative genetic and**

**epigenetic diversity in pairs of sympatric, closely related plants with**

**contrasting distribution ranges in south-eastern Iberian mountains.** *AoB*

*Plants* 2020, **12:**plaa013.

51.  Fargeot L, Loot G, Prunier JG, Rey O, Veyssiere C, Blanchet S: **Patterns of**

**Epigenetic Diversity in Two Sympatric Fish Species: Genetic vs.**

**Environmental Determinants.** *Genes (Basel)* 2021, **12:**107.

52. Lele L, Ning D, Cuiping P, Xiao G, Weihua G: **Genetic and epigenetic variations**

**associated with adaptation to heterogeneous habitat conditions in a**

**deciduous shrub.** *Ecol Evol* 2018, **8:**2594-2606.

53. Durvasula A, Fulgione A, Gutaker RM, Alacakaptan SI, Flood PJ, Neto C,

Tsuchimatsu T, Burbano HA, Pico FX, Alonso-Blanco C, Hancock AM: **African**

**genomes illuminate the early history and transition to selfing in Arabidopsis**

**thaliana.** *Proc Natl Acad Sci U S A* 2017, **114:**5213-5218.

54. Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, Muller R,

Dreher K, Alexander DL, Garcia-Hernandez M, et al: **The Arabidopsis**

**Information Resource (TAIR): improved gene annotation and new tools.**

*Nucleic Acids Res* 2012, **40:**D1202-1210.

55. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A,

Keane T, McCarthy SA, Davies RM, Li H: **Twelve years of SAMtools and**

**BCFtools.** *Gigascience* 2021, **10:**giab008.

56. Nassar LR, Barber GP, Benet-Pages A, Casper J, Clawson H, Diekhans M, Fischer

C, Gonzalez JN, Hinrichs AS, Lee BT, et al: **The UCSC Genome Browser**

**database: 2023 update.** *Nucleic Acids Res* 2023, **51:**D1188-D1195.

57. Bolger AM, Lohse M, Usadel B: **Trimmomatic: a flexible trimmer for Illumina**

**sequence data.** *Bioinformatics* 2014, **30:**2114–2120.

58. Krueger F, Andrews SR: **Bismark: a flexible aligner and methylation caller for**

**Bisulfite-Seq applications.** *Bioinformatics* 2011, **27:**1571-1572.

59. Lister R, O'Malley RC, Tonti-Filippini J, Gregory BD, Berry CC, Millar AH, Ecker JR:

**Highly integrated single-base resolution maps of the epigenome in**

**Arabidopsis.** *Cell* 2008, **133:**523-536.

60.    Hazarika RR, Serra M, Zhang Z, Zhang Y, Schmitz RJ, Johannes F: **Molecular**

1013            **properties of epimutation hotspots.** *Nat Plants* 2022, **8:**146-156.

1014    61.    Williams CJ, Dai D, Tran KA, Monroe JG, Williams BP: **Dynamic DNA methylation**

1015            **turnover in gene bodies is associated with enhanced gene expression**

1016            **plasticity in plants.** *Genome Biol* 2023, **24:**227.

1017    62.    Hamala T, Tiffin P: **Biased Gene Conversion Constrains Adaptation in**

1018            **Arabidopsis thaliana.** *Genetics* 2020, **215:**831-846.

1019    63.    Paradis E: **pegas: an R package for population genetics with an integrated-**

1020            **modular approach.** *Bioinformatics* 2010, **26:**419-420.

1021    64.    Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker

1022            RE, Lunter G, Marth GT, Sherry ST, et al: **The variant call format and VCFtools.**

1023            *Bioinformatics* 2011, **27**.

1024    65.    Zhang C, Dong SS, Xu JY, He WM, Yang TL: **PopLDdecay: a fast and effective**

1025            **tool for linkage disequilibrium decay analysis based on variant call format**

1026            **files.** *Bioinformatics* 2019, **35:**1786-1788.

1027    66.    Lefort V, Desper R, Gascuel O: **FastME 2.0: A Comprehensive, Accurate, and**

1028            **Fast Distance-Based Phylogeny Inference Program.** *Mol Biol Evol* 2015,

1029            **32:**2798-2800.

1030    67.    Yu G, Smith DK, Zhu H, Guan Y, Lam TTY: **ggtree: an r package for visualization**

1031            **and annotation of phylogenetic trees with their covariates and other**

1032            **associated data.** *Methods in Ecology and Evolution* 2017, **8:**28-36.

1033    68.    Jombart T: **adegenet: a R package for the multivariate analysis of genetic**

1034            **markers.** *Bioinformatics* 2008, **24:**1403–1405.

1035 69. Li H: **Aligning sequence reads , clone sequences and assembly contigs with**

1036   **BWA-MEM.** 2013, **00:**1-3.

1037 70. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A,

1038   Garimella K, Altshuler D, Gabriel S, Daly M, DePristo MA: **The Genome Analysis**

1039   **Toolkit: A MapReduce framework for analyzing next-generation DNA**

1040   **sequencing data.** *Genome Research* 2010, **20:**1297–1303.

1041 71. Niederhuth CE, Bewick AJ, Ji L, Alabady MS, Kim KD, Li Q, Rohr NA, Rambani A,

1042   Burke JM, Udall JA, et al: **Widespread natural variation of DNA methylation**

1043   **within angiosperms.** *Genome Biol* 2016, **17:**194.

1044 72. Sanchez T, Cury J, Charpiat G, Jay F: **Deep learning for population size history**

1045   **inference: Design, comparison and combination with approximate Bayesian**

1046   **computation.** *Mol Ecol Resour* 2021, **21:**2645-2660.

1047 73. Baumdicker F, Bisschop G, Goldstein D, Gower G, Ragsdale AP, Tsambos G, Zhu

1048   S, Eldon B, Ellerman EC, Galloway JG, et al: **Efficient ancestry and mutation**

1049   **simulation with msprime 1.0.** *Genetics* 2022, **220**.

1050 74. Paszke A, Gross S, Massa F, Lerer A, Bradbury J, Chanan G, Killeen T, Lin Z,

1051   Gimelshein N, Antiga L, et al: *pyTorch: An Imperative Style, High-Performance*

1052   *Deep Learning Library*. Curran Associates, Inc.; 2019.

1053 75. Shazeer N: **GLU Variants Improve Transformer.** *arXiv* 2020**:**2002.05202.

1054 76. He R, Ravula A, Kanagal B, Ainslie J: **RealFormer: Transformer Likes Residual**

1055   **Attention.** *arXiv* 2021**:**2012.11747.

1056 77. Su J, Ahmed M, Lu Y, Pan S, Bo W, Liu Y: **RoFormer: Enhanced transformer with**

1057   **Rotary Position Embedding.** *Neurocomputing* 2024, **568:**127063.

1058    78.    Gugger S, Debut L, Wolf T, Schmid P, Mueller Z, Mangrulkar S, Sun M, Bossan B:

1059            **Accelerate: Training and inference at scale made simple, efficient and**

1060            **adaptable.** *https://githubcom/huggingface/accelerate* 2022.

1061