LongTrack: long read metagenomics-based precise tracking of bacterial strains

and their genomic changes after fecal microbiota transplantation

Yu Fan[1], Mi Ni[1], Varun Aggarwala[1], Edward A. Mead[1], Magdalena Ksiezarek[1], Lei Cao[1], Michael
A. Kamm[2], Thomas J. Borody[3], Sudarshan Paramsothy[4], Nadeem O. Kaakoush[5], Ari Grinspan[6],
Jeremiah J. Faith[1] and Gang Fang[1#]

[1] *Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA*

[2] *Department of Gastroenterology and Medicine, St. Vincent's Hospital, Melbourne, VIC, Australia*

[3] *Centre for Digestive Diseases, Sydney, NSW, Australia*

[4] *University of Sydney, Sydney, Australia*

[5] *School of Biomedical Sciences, University of New South Wales, Sydney, Australia*

[6] *Division of Gastroenterology, Icahn School of Medicine at Mount Sinai, New York, NY, USA.*

[#] Address correspondence to: *gang.fang@mssm.edu*

Abstract

Fecal microbiota transplantation (FMT) has revolutionized the treatment of recurrent

Clostridioides difficile infection (rCDI) and is being evaluated across other diseases. Accurate

tracking of bacterial strains, the functional units of the microbiota, that stably engraft in

recipients, is critical for understanding the determinants of strain engraftment, evaluating their

correlation with clinical outcomes, and guiding the development of therapeutic bacterial

consortia. While short-read sequencing has advanced our understanding of FMT, it faces

challenges in the strain-level *de novo* metagenomic assembly of microbiome samples. In this

study, we described a novel framework, *LongTrack*, which uses long-read metagenomic

assemblies and rigorous informatics tailored for FMT strain tracking. We first highlighted

LongTrack advantage over short-read approaches especially when multiple strains co-exist in

the same sample. We then showed LongTrack uncovered hundreds of engrafted strains across

six FMT cases of rCDI and inflammatory bowel disease patients, validated by complementary

bacterial culturing. Furthermore, beyond conventional strain tracking, long reads also allowed us

to assess the genomic and epigenomic stability of successfully engrafted strains during the 5-

year follow-ups, revealing structural variations in their genomes, which may be associated with

strain adaptation in a new host environment. Combined, our study advocates the use of long

read metagenomics and LongTrack to enhance strain tracking in future FMT studies, paving the

way for the development of more effective defined biotherapeutic as an alternative to FMT.

Introduction

Fecal microbiota transplantation (FMT) has emerged as a new therapeutic strategy,

transforming the treatment paradigms for patients with recurrent *Clostridioides difficile* infection

(rCDI)[1–3]. By transferring gut microbiota from a healthy donor into a patient, FMT effectively

restores the patient's gut microbial ecology[4,5]. Potential applications of FMT extend beyond rCDI

onto treating a broader array of diseases[6,7]. Notably, the use of FMT in the treatment of

inflammatory bowel disease (IBD) is under investigation with multiple clinical trials[8–10].

Additionally, there is growing interest in the potential of FMT as an adjunctive tool in cancer

immunotherapy[11–13] .

The FMT procedure operates on the principle of transferring gut microbiota, which

constitutes a diverse array of bacterial strains, the functional impact units of the gut

microbiota[14–16], from a healthy donor into a recipient[4,17]. Therefore, the precise tracking of these

strains, from the point of transplantation to their eventual engraftment within the recipient's gut,

is of critical importance[18,19]. Specifically, reliable strain tracking can help researchers understand

factors that facilitate or hinder engraftment[5,17]. Moreover, comprehensive strain tracking can

provide vital information on how the engraftment of specific strains correlates with clinical

outcomes[18,20]. Lastly, a detailed understanding of bacterial strain engraftment can guide the

development of defined bacterial consortiums, composed of selected bacterial strains with

beneficial effects, for safer therapeutic applications[19–21].

Several previous studies, powered by high throughput metagenomic shotgun

sequencing, have provided valuable insights into strain engraftment in FMT in various

diseases[18,20–25]. These studies relied on short-read sequencing and metagenomic analyses,

which can detect single nucleotide polymorphisms (SNPs) and/or gene clusters that help

determine the source of a strain observed in the recipient after FMT[22–24,26–28]. However, as

demonstrated in a recent study that did comprehensive bacteria isolation combined with strain

tracking using high-purity and high-completeness genome sequences of bacterial isolates[20],

existing methods that use short-read sequencing alone face challenges when multiple strains

co-exist in a single microbiome sample[29–34]. Specifically, the complexity of recovering the

number of strains in a metagenome poses a critical challenge in subsequent strain tracking[29,35].

Bacterial isolation effectively addressed this challenge by converting a complex

microbiome sample into a finite set of discrete bacterial isolates[20]. Once the genomes of

individual bacterial isolates from donor and pre-FMT recipient samples are assembled (high-

purity and high-completeness), downstream tracking (post-FMT) can be reliably achieved based

on the unique k-mers of each strain in short-read metagenomic sequencing samples even with

modest-to-low sequencing depth[20]. However, standard culturing methods tend to miss certain

bacterial taxa due to the selective nature of culture media and/or conditions[36]. Additionally, it is

not feasible (labor intensive and time consuming) to perform comprehensive bacterial culture

across a large number of fecal samples, making it challenging to scale up to large FMT cohorts.

Strain-resolved *de novo* meta-genome assembly, if achievable, can greatly complement the bias

and scalability of culture-based approaches.

In this context, long-read sequencing could empower bacterial strain tracking in FMT.

Specifically, the ability of long-read sequencing to sequence DNA molecules of thousands of

kilobases in a single read is much more effective at *de novo* genome assembly, constructing

metagenome-assembled genomes (MAGs) at strain resolution[32,37,38]. Unlike short-read

sequencing, which often result in fragmented assemblies due to its limitations in spanning

repetitive and highly homologous regions, long-reads sequencing facilitates the recovery of

complete, contiguous genomes, providing more accurate strain-level resolution[38,39]. Additionally,

it enhances functional annotations, improves the detection of horizontal gene transfer, and

reveals intraspecies diversity that short-read methods fail to capture[40–42]. With this rationale, we

aimed to address two questions in this study. (Q1) How to reliably use long-read MAGs for

strain tracking? To answer this question, we developed a new framework, *LongTrack*, that uses

rigorous informatics tailored to reliably use long-read MAGs for strain tracking (**Fig. 1**). We built

upon a recent study[20] that conducted a systematic culture of bacterial strains from FMT donors

and recipients for rCDI with 5-year follow-ups (**Fig. 1A**), which serve as a reliable, complex and

realistic ground truth for rigorous evaluation of LongTrack (**Fig. 1B**). We systematically

evaluated LongTrack in comparison with short-read based alternative methods and observed

superior performance of LongTrack, especially in distinguishing co-existing bacterial strains

within a microbiome sample for reliable FMT strain tracking (**Fig. 1C**). (Q2) How stable are the

genomes of the engrafted strains (before and after FMT)? We asked this question to expand

beyond the standard scope of strain tracking building on the FMT cohort with longitudinal

sample collections across five years. In addition to *de novo* assembly, long-read sequencing

has advantages in strain-level read mapping, which facilitates the detection of complex

structural genomic variations within the same strains from microbiome data[40,42]. Its precise

quantitative analysis further reveals structural variations (SVs) within these strains as they

transfer from donors to post-FMT recipients (**Fig 1D**). In addition, we also built on the unique

ability of long-read sequencing technology for the direct detection of bacterial DNA

methylation[43–47] (which contributes to epigenetic heterogeneity and plasticity[45,48]), and highlight

epigenetic variations within the same strains during FMT and follow-ups. These in-depth

genomic and epigenomic analyses suggest the versatile capacity of individual strains during

FMT and long-term adaptation. Combined, this study introduces *LongTrack* (**Fig. 1C and 1D**), a

novel and rigorous framework that advocates the incorporation of long-read sequencing to

improve the tracking of bacterial strains and their genomic dynamics in future FMT studies.

Results

High purity, strain-level Long-read MAGs serve as a solid foundation for LongTrack.

The success of strain tracking in FMT depends on obtaining strain-resolved and high purity

genomes from both the donor and the recipient before FMT. Strain resolution is crucial as it

allows differentiation among co-existing strains of the same species, while the high purity of

MAGs avoids chimeric contigs or bins (comprising mixed strains) and supports accurate strain

tracking, forming a robust foundation for LongTrack.

Bacterial isolation is an effective method for achieving this, as sequenced bacteria

isolates derived from individual colonies yield discrete genomes with high purity and

completeness. With this rationale, we built on the comprehensive collection of 1000+ bacteria

isolates with individually assembled genomes from a recent study[20] as a ground truth that is

reliable, complex, and realistic to evaluate the long-read MAGs, and their accuracy in FMT

strain tracking. This study involved a cohort of seven healthy donors and thirteen recipients

using FMT for the treatment of rCDI. For evaluation purposes, we selected four FMT cases from

this cohort (3 donors and 4 recipients) from which, 184 unique strains (105 species) were

isolated, individually sequenced, and assembled[20]. In total, 25 fecal samples were taken from

the donors and recipients at multiple time points before and after FMT for up to 5 years (**Supp**

Fig. S1). We built on the previously generated short-read sequencing data[20], and generated

new long-read sequencing data for 15 samples (**Methods**. Additionally, we generated deeper

short-read data with two times the depth of the long-read data for two samples to enhance

detection power and enable a more comprehensive evaluation along with LongTrack (**Methods**).

To construct strain-level long-read MAGs, we employed metaFlye v2.9[49] for

metagenomic assembly from long-read Nanopore sequencing and MaxBin2[50] for species-level

binning. Furthermore, we utilized Strainberry[32] for haplotype phasing each species-level bin and

employed metaWRAP[51] for final strain-level binning. (**Supp Fig. S2, Methods**). From the three

donors and four recipients, we obtained a total of 460 long-read MAGs (**Fig. 2A**), which

collectively represent 460 unique strains (across 312 species), including 60 strains that were

also recovered from isolate culturing (**Fig. 2A**). Additional short-read Illumina sequencing data

was generated to facilitate method evaluation, resulting in 135 short-read MAGs from Donor 1

(D1) (**Fig. 2B**) and 20 short-read MAGs from Recipient 1 pre (R1) using metaSPAdes[52]. We also

compared the performance of another short-read assembly tool, MEGAHIT[53], for D1. MEGAHIT

produced 131 shared MAGs (ANI > 99.4%) with similar completeness and contamination levels

(**Supp Fig. S3A and S3B).** However, MEGAHIT[53] identified a smaller number of shared strains

with isolate genomes (**Fig 2B**, **Supp Fig. S3C).** Based on these results, we used MAGs from

metaSPAdes[52] for subsequent analyses. Contamination levels for all 460 long-read MAGs

(across 3 donors and 4 recipients pre-FMT samples) and 155 short-read MAGs (across 1 donor

and 1 recipient pre-FMT sample) were assessed using CheckM[54], revealing lower

contamination and higher contig N50 values in long-read MAGs than short-read MAGs (**Fig. 2C**

& 2D).

In our in-depth evaluation, we focused on sample D1, which had the most bacterial

isolates (136 unique strains, **Methods**), out of which 31 and 23 were shared with the 124 long-

read MAGs and 135 short-read MAGs, respectively (**Fig. 2B**). Notably, in comparison with the

isolate genomes, long-read MAGs include all 23 MAGs recovered by short-read MAGs along

with 8 additional strains represented only by long-reads MAGs (**Fig. 2B**). Both long-read and

short-read MAGs spanned the same 17 taxonomic orders. When we compared long-read MAGs

and isolate genomes, the former contained 8 unique orders, while the latter featured 5 unique

orders (**Fig. 2E**). These overlapping strains provide an opportunity to evaluate the reliability of

long-read MAGs in comparison with short-read MAGs generated from the same sample, using

genomes assembled from bacterial isolates as ground truths (**Methods**). The properties of long-

read MAGs and short-read MAGs were thoroughly evaluated regarding their strain resolution

and purity. First, we assessed their contamination levels by mapping long-read and short-read

MAGs to their corresponding isolate genomes (**Fig. 2F, Supp Fig. S4A, Methods**). Counting

the percentage of unaligned bases (MAGs to isolate genomes) as contamination, this method

builds on the valuable isolate genomes to provide more reliable insights than estimates by

CheckM[54] (**Supp Fig. S4B**). The contamination levels were lower for long-read MAGs (1.34%)

compared to short-read MAGs (4.77%), both for species with a single strain and those with co-

existing strains (**Fig. 2F and Supp Fig. S4C and S4D**). We further evaluated the similarity

between long-read and short-read MAGs and their corresponding isolate genomes using k-mer

similarity, which consistently indicated long-read MAGs exhibit a higher similarity (**Supp Fig.**

S5A). Importantly, long-read sequencing effectively recovered the same number of strains as

bacterial culture from the same donor sample (**Fig. 2G)**. In contrast, these co-existing strains

were not completely recovered in the short-read MAGs (**Fig. 2G**). It is worth noting that, despite

the comparable completeness of long-read and short-read MAGs (**Supp Fig. S5B**), the higher

purity of long-read MAGs (**Fig. 2F**) and strain resolution is expected to be critically important for

FMT strain tracking, because even a partially complete MAG that reliably represents individual

strains can provide unique k-mers valuable information for strain tracking[20,55], as we will also

evaluate in the next section. Using viralVerify (v1.1)[56] and PLASMe (v1.1)[57], we identified 11

plasmids from the 60 isolate genomes (shared with long-read MAGs). Among these, 7 plasmids,

including one circular plasmid, were recovered from long-read MAGs. The remaining 4 plasmids

that were only identified from isolate genomes may have been missed due to experimental size

selection steps that are used in standard long-read metagenomic library preparation. Using

isolate-based plasmid-to-strain assignment as the ground truth, we found that none of the 7

plasmids were correctly binned with their matching chromosomes. This analysis highlights that

plasmid binning in microbiome settings remain a challenge due to the usually inconsistent copy

numbers that could confuse coverage-based binning, and need to be assisted by more

specialized methods such as Hi-C[58]. While the shared strains demonstrate the reliability of long-

read MAGs, it's important to note that long-read MAGs and bacterial isolates capture different

additional strains and species due to their inherent methodological differences. Specifically,

long-read MAGs, based on metagenomic sequencing, favor strains with modest-to-high relative

abundance and tend to miss low-abundance ones, while bacterial culture favors species and

strains selected by culture media rather than abundance. For instances, *Akkermansia*

muciniphila and *Faecalibacterium prausnitzii* (**Fig. 2A**), two species that have been associated

with several diseases and treatment responses[59,60], have modest-to-high relative abundances

(0.63% and 3.94% respectively) in metagenomic sequencing but were missed by conventional

culture methods used in the previous study[20]. Conversely, *Escherichia coli* and *Enterococcus*

faecalis were cultured from D1 but not recovered by long-read sequencing because their

relative abundances were both below 0.01% (**Fig. 2A**). 37.1% and 65.9% of the total

metagenomic reads of D1 can be mapped to the isolate genomes and long-read MAGs

(**Methods**), respectively. When we combined the isolate genomes and long-read MAGs, >76%

of the total metagenomic reads can be mapped to the merged reference (**Supp Fig. S5C**).

Interestingly, the 31 common strains (13.5% of the 229 strains from long-read MAGs and

cultured isolates combined) represented a substantial proportion (30.7%) of the total

metagenomic reads, because of the relatively high abundance of these strains. Isolate genomes

and long-read MAGs uniquely explain 6.5% and 38.77% of the total reads, respectively. These

highlight the complementarity between long-read MAGs and bacterial culture, each providing

valuable insights into the microbial community, as we will further investigate in the context of

FMT strain tracking.

This section provides a comprehensive evaluation of long-read MAGs with a large

collection of isolate genomes from matched fecal samples. We showed that long-read MAGs

are more reliable than short-read MAGs for resolving strain-level MAGs *de novo* and at better

purity. Our analyses also highlighted the complementarity of long-read MAGs and bacterial

culture in capturing different strains and species. This section lays the groundwork for the

subsequent section, in which we will describe the rigorous design of the LongTrack method to

ensure the reliable utilization of long-read MAGs to achieve precise and scalable FMT strain

tracking.

LongTrack enables reliable use of Long-read MAGs for precise FMT strain tracking

Once individual isolate genomes from donor or pre-FMT recipient are assembled, strain tracking

can be done reliably based on unique k-mers (n=31) selected from isolate genomes and short-

read sequencing of post-FMT samples, providing a robust ground truth for evaluating

LongTrack[20]. This k-mer based strategy (called Strainer) for strain tracking has been

demonstrated in a previous work[20] with superior performance compared to several SNP-based

approaches[20] including inStrain[27], StrainFinder[23] , and ConStrains[22], particularly in resolving

strains with modest-to-low abundance and multiple co-existing strains. Our independent

evaluation of LongTrack for strain tracking confirmed these findings, showing consistent results

with the previous study[20] in that the k-mer based approach is more sensitive than the SNP-

based method (StrainFinder[23]): while the SNP-based method works well for strains with

relatively modest-to-high abundance, it tends to miss strains with modest-to-low abundance

(**Supp Fig. S6**).

However, the k-mer based strain tracking method (Strainer) is not compatible with long-

read MAGs, because while some of the long-read MAGs are nearly complete, others are not

(taxa with modest-to-low abundance). K-mers selected from incomplete long-read MAGs are not

necessarily strain-specific and can lead to false strain tracking results (**Methods**). Therefore, to

reliably use long-read MAGs for FMT strain tracking, we made important methodology

adaptations in LongTrack to ensure a more thoughtful selection of unique k-mers to avoid false

positives (**Methods**). Specifically, in LongTrack, for similar co-existing strains in one fecal

sample, or two similar strains each in the donor sample and recipient pre-FMT samples, we

selected unique k-mers only from their common genomic regions (**Fig. 1C, Methods**). The

rationale of this adaptation is to avoid the selection of unreliable k-mers that appear to be

unique to a strain but are not, due to the incompleteness of some long-read MAGs, which is the

key difference between high purity long-read MAGs and the complete genomes of bacterial

isolates[20]. More specifically, as illustrated in **Fig. 1C**, Strains A1 and A2 both belong to Species

A and represent co-existing strains derived from the same donor. If the k-mers identified for

Strain A1 in Step 1a were from genomic regions missing in the incomplete assembly of Strain

A2 (but actually existing in Strain A2), these k-mers are not truly unique to Strain A1, which

would lead to unreliable strain tracking results (not able to precisely differentiate between A1 vs

A2). In addition, we used negative controls to estimate the confidence of strain tracking, based

on which each strain is determined to be present (confidence score >95%) or absent in a post-

FMT sample (**Methods**). To evaluate the performance of LongTrack, we conducted precision

and sensitivity analyses for strain tracking using LongTrack with long-read MAGs versus

Strainer (the previous method) with short-read MAGs across 3 donors and 4 recipients, with

multiple fecal sampling time points each (326 unique strain presence/absence events).

LongTrack achieved perfect precision (1.00) and sensitivity (1.00) (**Fig. 3A**) when compared to

isolate genomes as the ground truth[20]. In contrast, strain tracking using Strainer with short-read

MAGs showed lower sensitivity (0.77) (**Fig. 3A**). Additionally, we compared the number of

unique k-mers used for strain tracking in cases involving multiple co-existing strains, using the

original Strainer method (with high completeness isolate genomes) versus LongTrack with high-

purity long-read MAGs (**Fig. 3B**). Although only k-mers from shared genomic regions were

retained, and LongTrack generally kept fewer unique k-mers compared to Strainer (based on

isolate genomes with high completeness), LongTrack still retained a sufficient number of unique

k-mers, supporting reliable strain tracking in post-FMT samples (**Fig. 3A and 3B**). Specifically,

we first built upon the two comprehensively characterized FMT cases, D1->R1, and D1-> R2,

for which strain tracking results are well validated against the ground truth[20]. For the 31 common

strains as described in the previous section, we evaluated LongTrack with long-read MAGs vs.

Strainer with isolate genomes for strain tracking and found that they had 100% consistency (**Fig.**

3C). In contrast, although Strainer with short-read MAGs allowed us to track 23 of the 31 strains

(74%), it failed to track the other 8 strains in post-FMT samples: this was largely because it was

not able to resolve the multiple strains co-existing within the same donor sample as described in

the previous section. In addition, we also observed that Strainer with short-read MAGs led to six

false positives tracking, which was due to the >5% contamination levels when compared to the

isolate genomes. For instance, the short-read MAGs *Bifidobacterium adolescentis* and

Bacteroides uniformis have contamination levels of 8.76% and 5.8%, respectively. Specifically,

28% of the contamination of the *B. adolescentis* is from a *Bacteroides vulgatus* strain within the

sample (**Supp Fig. S7A)** which led to the subsequent unreliable selection of unique k-mers and

the ultimate false positive tracking results in the Year 5 post-FMT sample from R1 (**Fig. 3C**). If

short-read MAGs with >5% contamination were marked as unreliable, the sensitivity would be

reduced from 0.77 to 0.39 (**Fig. 3A**), and accuracy of Strainer with short-read MAGs for strain

tracking decreased from 74% to 45% (**Supp Fig. S7B)**, while LongTrack strain tracking

maintained 100% consistency when compared to the Strainer with isolate genomes (**Supp Fig.**

S7B). We also applied LongTrack to two additional rCDI FMT cases: D2->R3 and D3->R4, and

also found 100% consistency with well-validated results derived from Strainer using isolate

genomes for the strains shared between long-read MAGs and ground truth i.e. isolate genomes

(**Fig. 3D and 3E**). A noteworthy observation is that completeness does not significantly affect

tracking accuracy, provided the long-read MAGs are of high-purity. For instance, the

Bacteroides caccae long-read MAG, despite having a low completeness of 17.24%, contained

sufficient unique k-mers to accurately track strains in the post-FMT samples (**Fig 3C**).

To further understand the performance of LongTrack with long-read MAGs compared to

Strainer with short-read MAGs, we designed additional evaluations with two mock FMT

scenarios with *in silico* spike-in of specific strains into D1 and R1 (**Methods**). In the first

evaluation, we spiked-in three strains of *Phocaeicola massiliensis* (Str1-3) to D1, and a fourth

strain of *P. massiliensis* (Str 4) to R1 pre-FMT (**Fig. 3F**). All four strains were added to post-FMT

recipient R1. While LongTrack were able to recover all four strains and support accurate

tracking of the four strains in post-FMT recipient R1, short-read MAGs only recovered two of the

four strains (**Fig. 3F**). In the second evaluation, we spiked-in ten pairs of strains, each with high

similarity (ANI>99.5%), from ten different species, one strain to D1 and the other to R1 pre-FMT,

similarly for all the ten pairs. All twenty strains were added to post-FMT recipient R1 (**Fig. 3H**).

LongTrack correctly tracked all the ten pairs of strains in post-FMT recipients, while Strainer with

short-read MAGs missed two of the strains (**Fig. 3H**).

In summary, the above evaluations support the reliability of LongTrack for FMT strain

tracking. Specifically, LongTrack can identify strain-specific k-mers from long-read MAGs to

differentiate highly similar strains within and between the donor and recipient pre-FMT samples

to achieve precise FMT strain tracking. This evaluation served as the foundation for the next

section, where we used LongTrack with long-read MAGs to track uncultured strains to better

understand strain engraftment in FMT for rCDI and IBD beyond culture-based approaches.

LongTrack helps better understand strain engraftment in rCDI and IBD patients

Building on the reliability of LongTrack-based FMT strain tracking as evaluated in the previous

section, we next aimed to scale up strain tracking to more comprehensively understand strain

engraftment in FMT. Across all the four cases of rCDI FMT, long-read sequencing recovered 396

MAGs with diverse taxonomy distribution (**Fig. 4A, Methods**) that were not captured by the

previous culture-based study[20].

We performed LongTrack to measure the engraftment of donor strains in the recipients,

focusing on uncultured strains from long-read. To summarize strain engraftment up to five years

after FMT, we used a metric named *proportional engraftment of donor strains* (PEDS), which is

calculated as the number of donor strains detected in a post-FMT recipient divided by the total

number of strains identified from the donor in the long-read MAGs. We found consistent and

stable engraftment of donor strains (**Fig. 4B**), with PEDS values comparable to cultured strains

(**Supp Figs. S8A**)**,** and individual trajectories of strain engraftment for donor-recipient pairs

(**Supp Figs. S9A-C)**, respectively. In these individuals, we report an average PEDS of 98% of

uncultured strains at the first week, which stabilized ~96% at 8 weeks (a commonly noted

clinical endpoint for measuring efficacy) and remained high at 63% even 5 years later. This

demonstrates that gut microbiota manipulation by FMT can lead to a fairly stable long-term

engraftment of donor 's uncultured strains in patients with rCDI.

In parallel with tracking donor strains, we also monitored the persistence of recipient

strains post-FMT using a metric called *proportional persistence of recipient strains (PPRS),*

defined similarly to PEDS but for recipient strains. We observed a major reduction (47%) in

PPRS of recipient strains post-FMT, both for cultured and uncultured strains, indicating a major

shift in the gut microbiota following FMT (**Fig. 4C, Supp Figs. S8B**). The absence of *C. difficile*

in post-FMT samples from recipient R1 is noteworthy, suggesting that the transplantation led to

a more balanced and health-promoting gut microbiota (**Supp Fig. S9A**). Some recipient strains

demonstrated resilience and were still detectable years post-FMT (**Supp Fig. S9A-C**), which

highlights the dynamic nature of the gut microbiota and underscores the potential for certain

recipient strains to withstand the microbial reshuffling induced by FMT.

330   The above analyses of PEDS and PPRS at the final time are consistent with the

previous study[20] that focused on strains resolved by bacterial cultures, but highlight the

hundreds of uncultured strains identified through long-read MAGs, providing a more complete

view of donor strain engraftment in FMTs for rCDI. When we examine the individual trajectories

of long-read MAGs recovered strains for donor-recipient pairs (**Supp Figs. S9A-C**), we found

several strains that stably engrafted in the recipient after five years include many species linked

to human health and diseases. One notable example, *A. muciniphila* (**Supp Figs. S9A, S9B**),

initially present in donor D1 and donor D2, is a mucin-degrading bacterium that has been linked

to health benefits in the context of several human diseases and treatment outcomes[60–65]. It has

been shown to play a role in preventing the overgrowth of harmful pathogens such as *C. difficile*

through the modulation of gut microbiota and metabolites[32]. Moreover, its presence is thought to

bolster gut barrier integrity and modulate the immune response, thereby creating an

environment less favorable for *C. difficile* colonization and infection[33]. *A. muciniphila* strains from

donors (D1 and D2) demonstrated stable engraftment in post-FMT recipients, with persistence

observed for up to five years (**Supp Figs. S9A, S9B**). In contrast, an *A. muciniphila* strain from

pre-FMT recipient R3 was not detected in the corresponding post-FMT recipient R3 four weeks

after the FMT (**Supp Fig. S9B**). *F. prausnitzii* (**Supp Fig. S9A)** is another species well resolved

by long-read MAGs but not the large-scale culture in the previous study[20]. Specifically, it's

known for its anti-inflammatory effects and its ability to produce short-chain fatty acids like

butyrate, which are beneficial for the health of the colon and the entire gastrointestinal tract[59,67–

71] . This bacterium is often considered a marker of a healthy gut and has been the focus of

research for its potential role in maintaining gut health and preventing IBD[67,68,72]. *F. prausnitzii*

strain from donor D1 was detected in post-FMT recipient R1 eight weeks after FMT, but not

observed five years after the FMT (**Supp Fig. S9A**). *Ruminococcus bromii* (**Supp Figs. S9A, S9C)** is another species tracked by LongTrack, known for its ability to break down resistant starches into short-chain fatty acids like butyrate[73]. These short-chain fatty acids contribute to gut health, serve as an energy source for colon cells, and have anti-inflammatory properties[73,74]. *R. bromii* strains from donors D1 and D3 showed stable engraftment in post-FMT recipients, with persistence observed up to the last collection time point (**Supp Figs. S9A, S9C**).

Next, we preformed LongTrack to expand the tracking analysis to strains represented by long-read MAGs from recipient post-FMT samples and reversely tracked them in earlier time points (**Methods**). The rationale of this design was that some strains were indeed present in donor or recipient pre-FMT but at too low abundance to be recovered by long-read MAGs, yet their abundances can be higher at recipient post-FMT samples, hence assembled by long reads and can be reversely tracked in earlier time points to have a more complete view of strain engraftment in FMT. Indeed, this design helped us identify 17, 8 and 10 additional strains from the three FMT cases D1->R1, D2->R3, and D3-R4, respectively (**Supp Fig. S10**). All these cases have clear evidence that they were from either the donor sample or recipient pre-FMT samples but just at low abundance so were not recovered by long-read MAGs from pre-FMT samples. This strategy helps make the study of strain engraftment in FMT more comprehensive. Interestingly, in this analysis, we also observed a few cases where a novel strain was recovered from post-FMT recipients, but no evidence supported the presence of these strains in any of the earlier times, neither donor nor recipient pre-FMT (**Supp Fig. S11**), which suggested that these strains may have been acquired from the sourced unrelated to the donor or recipient. For example, a strain of *Acidaminococcus intestini* from the year 5 sample of post-FMT recipient R1 was absent at every earlier time or in the donor (**Supp Fig. S11A**), was likely acquired from the environment sometime between week 8 and year 5, unless its abundance was below detections limit in all the earlier time points.

In addition to the four cases of FMT for the treatment of rCDI, we further applied the

LongTrack to two FMT cases for the treatment of IBD[8,75] (**Supp Fig. S12**), which represents

pre-FMT recipients with more complex microbiome than rCDI patients. Different from the FMT

cases for rCDI patients, the two IBD patients (R5 and R6) each received fecal samples from six

donors (D4 - D9), as multiple donors are often used in the FMT treatment of IBD considering the

more diverse and resilient pre-FMT gut microbiome in IBD patients compared to rCDI

patients[8,75]. This greater complexity of FMT provided a great opportunity to demonstrate the

power of LongTrack for strain tracking. Here, we conducted long-read sequencing on two of the

donors (D4 and D5) and the two pre-FMT recipients, recovered 188 MAGs (R5: 45, R6: 52, D4:

52, D5: 39). Out of these strains, 58 (64%) and 32 (35%) strains from the two donors

successfully engrafted the recipient 16 weeks post-FMT (8 weeks of FMT dosage then 8 weeks

after cessation) for the two recipients, respectively (**Figs. 4D, 4E)**. We also estimated PPRS

post-FMT and observed that 80% and 96% of strains from recipient pre-FMT were present after

16 weeks post-FMT (**Fig. 4D, 4E**). Compared to the FMT cases for rCDI at a similar time point

(8 weeks), we observed lower engraftment of donor strains and higher persistence of recipient

strains post-FMT in the two FMT cases for IBD, which is consistent with the expectation that the

gut microbiome of IBD patients is more diverse than those of rCDI patients, thus exhibiting a

stronger resilience upon FMT introduction of donor strains[8]. Because fecal samples of IBD

patients exhibited greater diversity than those from patients with rCDI, we observed several

coexisting strains in the post-FMT recipients, where strains from donors and recipients were

detected. Specifically, co-existing strains of the same species were observed for 19 and 20

species in the two IBD FMT cases (**Fig. 4F, 4H**). Notably, while nearly all strains from the pre-

FMT recipient samples remained stable 16 weeks post-FMT, some donor strains were not

detectable at 16 weeks post-FMT, suggesting endogenous strains in IBD patients pre-FMT are

more resilient (than donor strains) in the same host after FMT.

In summary, LongTrack empowered the scalable tracking of hundreds of additional

strains missed by conventional culture from the same FMT cases in a previous study, and

provided a more comprehensive view of strain engraftment in six FMT cases across rCDI and

IBD patients. The newly resolved strains that stably engrafted the post-FMT recipient include

species that have been implicated with healthy benefits across several human diseases, such

as *A. muciniphila* and *F. prausnitzii.* The complex co-existing strains post-FMT in the two IBD

FMT cases also demonstrated the power of LongTrack based strain tracking for more complex

diseases beyond rCDI.

Long-read sequencing of post-FMT samples uncovered genomic structural variations

within engrafted strains.

In addition to determining strain presence/absence during FMT, it's crucial to understand how

donor strains adapt to the new microbial and host environments in the recipient. The pressure to

adapt can drive the selection for genomic variations in these donor strains, a topic of interest but

not yet extensively studied in FMT research[76–78]. While short-read sequencing is powerful in

detecting single nucleotide variations (SNVs) and small indels, long-read sequencing has great

advantages in the detection of more complex structural variations (SVs, such as insertions,

deletions, and inversions). In addition, another advantage of third-generation long-read

sequencing technology, represented by the Nanopore and Pacific Biosciences (PacBio)

platforms, is that they allow the direct detection of bacterial DNA methylation[43–47,79]. Therefore,

we aimed to use long-read sequencing of donor and post-FMT samples to assess the stability

and dynamics of SVs and DNA methylation of individual strains across the longitudinal samples

collected over a five-year post-FMT.

For SV analyses, we focused on the 255 long-read MAGs and 41 isolated genomes

without co-existing strains from the rCDI donors (D1-D3), because genomic analysis of single

strain simplifies the calling and interpretation of strain-level SVs in this exploration. A systematic

analysis of long-read sequencing data of the post-FMT recipients (R1-R4) across various time

points led to the detection of 38 SVs, which included 6 deletions, 20 insertions, and 12

inversions, as determined by stringent filtering criteria to maximize reliability of genome-specific

read mapping (**Methods**). Further comparison of read alignments of SV regions— specifically, a

43 bp deletion in *Parabacteroides merdae* and a 1,250 bp insertion in *Alistipes putredinis*—

showed that both long-read sequencing methods provided clear and consistent flanking

sequence alignments (**Fig. 5A**), illustrating they were from the same strain. In contrast, short-

read sequencing plots displayed several mismatches, making it challenging to reliably attribute

the sequences to the same strain (**Fig. 5A**). Specifically, out of the 38 SVs, 12 were identified in

four strains that we were able to isolate from both donor and post-FMT recipient samples. To

validate these SVs, long-read sequencing was performed on newly cultured isolates from these

strains. Paired donor and post-FMT recipient isolates showed high genome-wide ANI (>99.9%) ,

confirming that the strains from donor and post-FMT samples are indeed the same strain.

Among the 12 SVs, 11 alternative SV genotypes (91.67%), indeed exist in the same strain as

validated by long-read sequencing of the isolates from donor and post-FMT samples,

demonstrating the high reliability of SV detection within the same strain using long-read

sequencing (**Fig. 5B, Fig. S13**). The remaining 26 SVs were identified in five strains that were

de novo assembled into near-complete long-read MAGs in both donor and post-FMT samples.

Importantly, comparison of paired donor and post-FMT samples confirmed that all the 26 SVs

originated from the same strain, with genome-wide ANI >99.70% (mean ANI = 99.85%).

Of the 38 SVs (**Fig. 5B**), 15 were located within genic regions, while 23 were found in

intergenic regions. Notably, 5 SVs were identified in TonB-dependent receptor genes or in their

upstream regions (**Fig. 5B**). These receptors, predominantly located in the outer membrane of

Gram-negative bacteria, are integral for nutrient transport, particularly iron, into bacterial cells[80].

Previous research has suggested that deletions in these TonB-dependent receptor genes might

be associated with immune-mediated selective pressures[81]. For example, an inversion in a

strain of *Bacteroides stercoris* from D1 had high frequency exclusively in late-stage post-FMT

sample, across TonB-dependent receptor susC genes, which may be associated with immune

adaptation to the new host environment **(Fig. 5C)**. In *A. putredinis* D1, the deletion shown in **Fig**

5C encompasses an IS110 family transposase gene designated ISCaa7, was observed with a

high prevalence post-FMT **(Fig. 5C)**. The ISCaa7 gene encodes a transposase, an enzyme that

mediates the movement of transposons (or jumping genes) within the genome of a single cell[82].

This alteration may affect genetic diversity, evolution, and antibiotic resistance propagation,

highlighting the impact on microbial adaptability[83]. Moreover, in *Gemmiger formicili* D2, an

insertion proximal to yesO gene (Fig. 5C), encoding an ABC transporter substrate-binding

protein, which likely plays a significant role in nutrient uptake and metabolic processes [84].

We also observed some more complex SVs that involve bacterial restriction-modification

systems (RMS), which play a vital role in safeguarding bacterial genomes by cleaving

unmethylated foreign DNA while methylating their own DNA[45,48,85–88]. These SVs can alter the

activity or specificity of DNA methyltransferases (MTases), driving genome-wide methylation

changes that regulate gene expression and enhance phenotypic diversity[89–91], promoting

bacterial adaptation to stresses such as antibiotics and host immune responses[92,93]. These

epigenetic phase variations (ePV) driven by SVs provide reversible, heritable phenotypic states

with adaptive advantages[48,90]. Our approach leverages the power of long-read sequencing to

simultaneously detect SVs and DNA methylation patterns at single-molecule resolution.

Compared to short-read methods, long-read technologies can more reliably map SVs, and link

RMS genotypes to methylation motifs, revealing the impact of SVs on epigenomic changes. For

example, we detected a complex SV spanning a Type I RMS S subunits of *P. merdae.* The

nested inverted repeats create a complex genomic shuffling among six different genotypes (**Fig.**

5D). Notably, genotype 5, predominantly observed in the donor, had a drastic reduction post-

FMT, while genotype 6 was absent in the donor but significantly increased in post-FMT recipient.

Because the specificity unit, S, of the Type I RMS system determines the sequence recognition

motif of the MTases, the genotype switches of this RMS locus can create genome-wide

methylation changes[41]. To test this hypothesis, we used PacBio sequencing data, which offers

advantages in detecting 6-methyladenine (6mA) at single molecule resolution[46,79,86], and indeed

observed a DNA methylation motif CA6mAYNNNNNNRTC exclusively in post-FMT samples

(**Fig. 5E**). This observation suggests some SVs can influence genome-wide changes in

methylation, which might empower the *P. merdae* strain to better adapt to the new gut

environment of the post-FMT recipient. Further investigation revealed that the other DNA

methylation motifs in this *P. merdae* strain were stable between donor and recipient across

different time points (**Fig. 5E**). These results suggest RMS-associated SVs may be involved in

creating clonal epigenetic heterogeneity within a single strain, which often mediate the

expression of multiple genes as a bet hedging strategy for adaptation[87,94,95]. This analysis also

illustrates the power of long-read sequencing to resolve SVs along with DNA methylation

analysis to empower more comprehensive microbiome studies.

In addition, we examined 11 unique methylation motifs across 10 single strains with

high-coverage Nanopore long-read data in Donor D1 (**Fig. 5F**). Among these, 1 motif was

identified as 5-methylcytosine (5mC), 2 as 4-methylcytosine (4mC) motifs, and 8 as 6mA motifs,

all of which conform to Type II methylation patterns. These motifs demonstrated consistency

and stability in both donors and recipients over various time points, which correlates with the

absence of SVs in their flanking genomic regions.

In summary, to assess the genomic stability of engrafted strains, we built on the

strengths of long-read sequencing data from post-FMT samples to detect both complex

genomic SVs and epigenomic variations. This analysis illustrate a new approach to study the

genomic stability and dynamics of donor strains that engrafted post-FMT recipients, which may

lead to new understandings of bacterial adaptation in new host microbiome environment.

Discussion

In this study, we present a novel approach, LongTrack, that uses long-read sequencing and

tailored informatics for precision FMT strain tracking, addressing a critical gap in the

comprehensive tracking of microbial strains.

Methodological advancements: Although long-read MAGs overcome the limitations of

short-read sequencing in the *de novo* assembly of co-existing, highly related bacterial strains,

special considerations in LongTrack were critical in the reliable use of long-read MAGs for strain

tracking. A key methodological innovation of LongTrack is the rigorous selection of strain-unique

k-mers from long-read MAGs that are compatible with the high-purity yet sometime incomplete

long-read MAGs (for low-abundance strains), which are different from separately assembled

bacterial isolate genomes[20]. To resolve this issue, LongTrack applies a critical step that retains

only those unique k-mers originating from assembled genomic regions shared across the

conspecific strains both within and between FMT donor and recipient samples. Once reliable

strain-unique k-mers are selected by LongTrack from long-read MAGs, subsequent strain

tracking can then be reliably performed using Strainer with short-read sequencing data. We also

used negative controls to assign confident scores to assess the strain tracking results for

reliable applications. This LongTrack framework was rigorously validated against ground truth

methods based on Strainer with isolate genomes from large-scale bacterial culture.

The previous isolate-based method served as valuable controls for methods evaluation,

although we would like to acknowledge that culture-based approaches have biases and

limitations as they rely on standard culturing conditions, which often miss certain bacterial taxa

due to the selective nature of culture media. For instance, strains like *A. muciniphila* and *F.*

prausnitzii were often missed under standard culture conditions, despite being culturable from

stool samples under more stringent conditions[96,97]. Nonetheless, the inclusion of isolate

genomes in our study was primarily intended to evaluate LongTrack with long-read MAGs

against Strainer with short-read MAGs, particularly in resolving co-existing strains. Importantly,

the use of isolate strains with pure and complete genome assemblies provides a solid approach

to evaluate strain tracking, as demonstrated in the previous work that Strainer provides more

reliable strain tracking than several SNP-based methods[20].

Application of LongTrack to FMT cases for both rCDI and IBD patients demonstrated the

method's applicability across diverse microbiome complexities. Notably, the application of

LongTrack allowed the identification of a large number of strains across diverse taxonomy,

including many strains not cultured by standard conditions[20], and revealed more complete

insights into their long-term engraftment in rCDI patients up to five years post-FMT. In IBD FMT

trios, LongTrack demonstrated its effectiveness in more complex microbiomes with greater

strain diversity, and the FMT from multiple donors to IBD patients. Due to the relatively small

sample size, the study of rCDI and IBD FMT samples were mostly for methods evaluation and

demonstration of broad applicability, instead of generalized findings on strain engraftment.

Future studies on large cohorts could further explore the influence of strain richness on

engraftment dynamics and its therapeutic implications[75].

Beyond identifying strains that are stably engrafted post-FMT, this study also made the

first attempt to assess the genomic dynamics of donor strains that might reflect their adaptation

to their new surroundings within the recipient's microbial and host environments. This adaptation

process, which may select for certain genomic changes in the engrafted strains, remained a

relatively unexplored area in FMT research. A number of recent studies highlighted the

importance of SVs (insertions, deletions, and inversions), in addition to SNVs, in microbiome

adaptation[66,81,82]. However, reliable detection of strain-level SVs, especially large SVs, across

the longitudinal samples poses a challenge to short-read sequencing but can be better

approached by long-read sequencing[83]. Long reads have the advantages in detecting complex

genomic structures like shufflons, dynamic DNA inversion systems known to influence bacterial

adaptation by generating structural diversity[87,94,95]. So, our study has utilized these advantages

of long-read sequencing to examine the stability and variations in SVs in individual strains

across longitudinal samples collected over five years following FMT, unveiling dynamic changes

over time may reflect strain adaptation to the new environment. For some of the SVs that

involve bacterial RMS, we further investigate their impact on genome-wide DNA methylation

patterns. This analysis capitalized on the unique capability of long-read sequencing, which

enables simultaneous detection of complex SVs and single-molecule resolution of DNA

methylation to assess both genomic and epigenomic changes in donor strains post-FMT.

In this study, we employed two long-read sequencing technologies: Oxford Nanopore

and PacBio sequencing. Oxford Nanopore technology produces ultra-long reads that are highly

effective for *de novo* assembling high-quality MAGs and detecting SVs. PacBio sequencing

offers the advantage of high-accuracy long read and quantitative detection of 6mA and 4mC at

single-molecule resolution[79], facilitating the strain deconvolution and the identification of longer

motifs such as CA6mAYNNNNNNRTC which were exclusively observed in post-FMT strains

associated with genotype 6 (**Fig 5E**). By leveraging the complementary strengths of these

technologies, we suggest a new dimension to characterize how donor strains adapt in post-FMT

recipients over time.

While LongTrack offers novel insights, it does inherently face limitations in recovering

super low abundance strains. To mitigate this, we propose two complementary strategies. First,

we would like to emphasize the complementarity and integration between long-read sequencing

and culture-based methods that can provide a more comprehensive view of the FMT strain

engraftment. This synergistic approach is especially valuable for tracking strains that are either

challenging to culture or present in low abundance, thus ensuring a more complete microbial

representation. Second, the challenges posed by low-abundance strains to metagenomic

approaches can be mitigated by targeted enrichment methods. Specifically, recent studies have

demonstrated efficient enrichment of specific taxa of interests based on flow cytometry[84] or

bacterial methylome patterns[85]. These enriched subsets can then be sequenced and analyzed

separately, enhancing the recovery and tracking of low abundance strains.

As the cost of both Nanopore and PacBio sequencing platforms continue to decrease,

we expect LongTrack to be broadly used for comprehensive strain tracking in FMT studies,

paving the way towards better understanding FMT and the development of more effective

defined biotherapeutics. More broadly, LongTrack can also help precise strain tracking in other

biomedical studies to enhance our understanding of strain transmission, genomic and

epigenomic variation, and their roles in health and diseases[101–104].

Figure legend

Figure 1. Description of FMT samples, matched bacteria isolates and the core design of LongTrack for reliable FMT strain tracking and detecting intra-strain structural variations in engrafted strains.

A, Stool samples were collected from three FMT donors and four FMT recipients at multiple time points over five years. Comprehensive bacterial culture, isolate genome assemblies, and short read-metagenomic sequencing have been performed in the previous study by Aggarwala et al, and shown along with FMT samples. Long read metagenomic sequencing data were newly generated for this study. **B**, Long-read data from donors' and recipients' samples before FMT were used to construct *de novo* metagenome-assembled genomes (long-read MAGs). The overlapping isolate genomes enable a comprehensive evaluation of the reliability of long-read MAGs, which serves for the foundation of LongTrack. **C**, Schematics of the core design of LongTrack for FMT strain tracking based on strain-level long-read (LR) MAGs. Strain-unique k-mers (k=31) were identified for each strain-level LR MAG from donor or pre-FMT recipient as the markers by eliminating common k-mers present in public database, or the other MAGs in the same FMT donor and recipient (**Methods**). To ensure specificity, strain-unique k-mers are only selected from shared regions across multiple strain-level LR MAGs within the same species. The presence of these final unique k-mers of a specific MAG (above a confidence score, **Methods**) in a post-FMT recipient's short-read metagenomic data indicates that the corresponding MAG is present (green color) in the post-FMT recipient sample. **D**, Long-read metagenomic sequencing data were also utilized for strain-specific detection of structure variations (SVs) that differ between donor and post-FMT samples. The references for putative genotypes were created and used to map and quantify specific SVs in each sample, allowing tracking of SV changes between hosts and/or along time. The long-read approach has unique advantage in strain-level read mapping that allows reliably detection of intra-strain variations,

which are challenging for short reads due to their limitations in accurately mapping reads to

specific strains.

Figure 2. High-purity, strain-level long-read MAGs serve as a solid foundation for

LongTrack.

A, The Venn diagram displays the number of strains that were recovered by long-read (LR)

MAGs (yellow color) or isolate genomes from bacterial culture (red color) from 3 donors and 4

recipients, with the left bar plot illustrating examples of species with MAGs with modest-to-high

relative abundance but were missed by culture, and with right bar plot illustrating examples of

isolates that were cultured but not recovered by long-read sequencing because of their very low

abundances. **B**, The Venn diagram displays the number of strains that can be recovered by LR

MAGs (yellow color), short-read (SR) MAGs (blue) or isolate genomes (red) in D1. LR MAGs

include all the 23 strains recovered by SR MAGs, along with 8 additional strains not assembled

by short reads. **C**, Contamination levels of 460 LR MAGs (from three donors and four recipients)

and 135 SR MAGs (from Donor 1, D1, for which we generated ultra-deep data) assessed using

CheckM, showing significantly lower contamination in LR MAGs ($p < 0.01$, **). **D**, Distributions of

relative abundance (log scale) and contig N50 (kb) for LR MAGs and SR MAGs. **E,** Order-level

taxonomic comparison of LR MAGs, SR MAGs, and isolate genomes from D1. **F**, Contamination

level for 60 LR MAGs (from Donor 1-3 and Recipient 1-4) and 23 SR MAGs (from D1),

evaluated across various species with a single strain or multiple strains. LR MAGs consistently

show lower contamination. Contamination levels were assessed by mapping long-read and

short-read MAGs to their corresponding isolate genomes and calculating the percentage of

unaligned bases as a measure of contamination. **G**, Long reads (LR) sequencing have a unique

advantage to resolve multiple co-existing strains from the same species than short reads (SR)

sequencing.

Figure 3. Comprehensive evaluation of LongTrack for FMT strain tracking.

A, Precision and sensitivity of strain tracking using LongTrack (methods tailored for long-read

MAGs) versus Strainer with short-read MAGs (SR MAGs). Across 3 donors and 4 recipients

with multiple fecal sampling time points (326 unique strain presence/absence events),

LongTrack achieved 100% consistency with isolate-based strain tracking, even under stringent

quality filtering (**Methods**). In contrast, Strainer with SR MAG-based strain tracking (n = 186)

showed lower sensitivity (0.77 overall, reduced to 0.39 under stringent filtering). **B**, Numbers of

unique k-mers used for strain tracking in cases of multiple co-existing strains by the original

Strainer method (with complete genomes assembled for isolates) and the LongTrack method

(with long-read MAGs). Although some k-mers from shared regions are filtered out by

LongTrack, enough unique k-mers remain for reliable strain tracking in post-FMT samples. **C**,

Presence (green) or absence (gray) of strains in post-FMT recipients determined by strain-

specific unique k-mers selected from the Strainer with isolate genomes (considered the ground

truth), LongTrack with LR MAGs, or Strainer with SR MAGs. Discrepancies with the ground truth

are indicated by an "X" marker. **(C-E).** The evaluation of strain tracking was performed for four

FMT cases of rCDI: D1->R1 (**C**), D1->R2 (**C**), D2->R3 (**D**), and D3->R4 (**E**). **F**, Further

evaluation of LongTrack for FMT strain tracking using *in silico* spike-in data that represent the

co-existence of multiple strains of the same species in the donor sample. Specifically, we

spiked-in three strains of *Phocaeicola massiliensis* (Str 1-3) to D1, and a fourth strain of *P.*

massiliensis (Str 4) to R1 pre-FMT. All four strains were added to post-FMT recipient R1. **H**,

Further evaluation of LongTrack for FMT strain tracking using *in silico* spike-in data that

represent cases when there are high-similarity strains present in between the donor and pre-

FMT samples using *in silico* spike-in data. Specifically, we spiked-in ten pairs of strains, each

with high similarity (ANI>99.5%), from ten different species, one strain to D1 and the other to

pre-FMT recipient R1, similarly for all the ten pairs. All twenty strains were added to post-FMT

recipient R1.

Figure 4. LongTrack uncovered hundreds of engrafted strains across six FMT cases of

rCDI and IBD patients.

A, Diverse taxonomy distribution of long-read MAGs that are not cultured in the previous study

by Aggarwala *et al*. **B**, Proportional engraftment of donor strains (PEDS) in individual post-FMT

recipients at various time points of the 5-year follow up. **C**, Proportional persistence of recipient

strains (PPRS) in individual post-FMT recipients at various time points of the 5-year follow up. **D**

& **E**, PEDS and PPRS analysis for the two FMT cases for IBD patients R5 (**D**) and R6 (**E**),

where three colors are used to represent the strains from the two Donors (D4 and D5) or the

recipient. (**F** & **H**) The presence and absence of co-existing strains in the two IBD patients (R5,

F, R6, **H**) after FMT that originated from the two Donors (D4 and D5) or the recipient.

Figure 5. Long reads uncovered complex structural variations in engrafted strains during

the 5-year follow-up after rCDI FMT.

A, IGV plots illustrating the advantage of long reads in the detection of intra-strain structural

variation (SV) detection, including a 43 bp deletion in *Parabacteroides merdae* and a 1250 bp

insertion in *Alistipes putredinis*, over short-read sequencing. Flanking sequences of SVs show

clear and consistent alignments in all long-read IGV plots, supporting that they are from the

same strain. In contrast, short-read plots display several mismatches, making it difficult to

reliably determine whether the sequences belong to the same strain. **B**, The heatmap plot

summarizing all the SVs detected in the donors and post-FMT recipients at early (E) and late (L)

time points after FMT including inversions (INV), deletions (DEL), and insertions (INS). Samples

with insufficient coverage over a specific SV region are marked by an asterisk (*). The SVs in

the strains were marked with a blue cross (+) when their SVs were detected in the isolate

genomes from the fecal samples of both donors and recipients. SVs are also categorized by

their location: intergenic regions (IR) and genic regions (GR). **C**, Three examples of SVs with

changes in the frequency of the two genotypes across donor and post-FMT recipient at different

time points. The first one illustrates an inversion (detected in *Bacteroides stercoris*), the second

one illustrates deletion (detected in *Alistipes putredinis*) changes in genotype frequency both in

the early and late stage post-FMT, while the third one shows insertion (detected in *Gemmiger*

formicilis) and change in genotype frequency only at late stage post-FMT. Pie charts

demonstrate the frequency of two genotypes shown across the donor (D1 or D2) and post-FMT

recipient (R1 or R3) at early (E) and late (L) time points after FMT. **D**, A complex inversion

(shufflon) involving six genotypes is located on restriction endonuclease subunit S of the RMS

system in *Parabacteroides merdae*. The frequency of the genotypes varies across donor and

post-FMT recipient at different time points, with major changes observed for Genotype 6 *i.e.*

absence in donor and increases after FMT. **E**, DNA adenine methylation (6mA) motifs in the *P.*

merdae strain detected across donor and post-FMT recipient with different time points;

CA6mAYNNNNNNRTC corresponds to Genotype 6 of this RMS phase variant, which is only

detectable after FMT. **F**, DNA methylation motifs detected across several single strains

assessing their epigenomic stability in the donor (D1) and post-FMT recipient (R1) over time.

Acknowledgments

This work was supported in part by the staff and resources of the Microbiome Translational Center and Department of Scientific Computing at the Icahn School of Medicine at Mount Sinai. This work was supported by grant no. R35 GM139655 (G.F.) from the National Institutes of Health. G.F. is a Hirschl Research Scholar by Irma T. Hirschl/Monique Weill-Caulier Trust and a Nash Family Research Scholar.

Competing interests

S.P. has served as a consultant / steering committee member for Vedanta Biosciences and has received speaker / advisory board fees from AbbVie, Dr Falk Pharma, Ferring, Janssen and Takeda. J.F. is a consultant for Vedanta Biosciences and Genfit. The remaining authors declare no competing interests.

Author contributions

Initiated study: G.F. and J.F. Developed methodology: Y.F. and G.F. Data analysis: Y.F., J.F. and G.F. Data interpretation: Y.F., M.N., V.A., E.A.M., M.K., L.C., J.F. and G.F. Collection and interpretation of clinical data: M.A.K., T.J.B., S.P., N.O.K. and A.G. Supervised research: G.F. Wrote first draft of paper: Y.F. and G.F. Approved paper: all authors.

Reference

1. Feuerstadt, P. *et al.* SER-109, an Oral Microbiome Therapy for Recurrent *Clostridioides difficile* Infection. *N Engl J Med* **386**, 220–229 (2022).

2. Khanna, S. *et al.* Efficacy and Safety of RBX2660 in PUNCH CD3, a Phase III, Randomized, Double-Blind, Placebo-Controlled Trial with a Bayesian Primary Analysis for the Prevention of Recurrent *Clostridioides difficile* Infection. *Drugs* **82**, 1527–1538 (2022).

3. Kelly, C. R. *et al.* Effect of Fecal Microbiota Transplantation on Recurrence in Multiply Recurrent *Clostridium difficile* Infection: A Randomized Trial. *Ann Intern Med* **165**, 609–616 (2016).

4. Ooijevaar, R. E., Terveer, E. M., Verspaget, H. W., Kuijper, E. J. & Keller, J. J. Clinical Application and Potential of Fecal Microbiota Transplantation. *Annu Rev Med* **70**, 335–351 (2019).

5. Li, S. S. *et al.* Durable coexistence of donor and recipient strains after fecal microbiota transplantation. *Science* **352**, 586–589 (2016).

6. Pamer, E. G. Fecal microbiota transplantation: effectiveness, complexities, and lingering concerns. *Mucosal Immunol* **7**, 210–214 (2014).

7. Bethlehem, L. *et al.* Microbiota therapeutics for inflammatory bowel disease: the way forward. *The Lancet Gastroenterology & Hepatology* **9**, 476–486 (2024).

8. Paramsothy, S. *et al.* Multidonor intensive faecal microbiota transplantation for active ulcerative colitis: a randomised placebo-controlled trial. *The Lancet* **389**, 1218–1228 (2017).

9. Rossen, N. G. *et al.* Findings From a Randomized Controlled Trial of Fecal Transplantation for Patients With Ulcerative Colitis. *Gastroenterology* **149**, 110-118.e4 (2015).

10. Chu, N. D. *et al.* Dynamic Colonization of Microbes and Their Functions after Fecal Microbiota Transplantation for Inflammatory Bowel Disease. *mBio* **12**, e00975-21.

11.   Sevcikova, A. *et al.* The Impact of the Microbiome on Resistance to Cancer Treatment

with Chemotherapeutic Agents and Immunotherapy. *International Journal of Molecular*

Sciences **23**, 488 (2022).

12.   Davar, D. *et al.* Fecal microbiota transplant overcomes resistance to anti-PD-1 therapy in

melanoma patients. *Science* **371**, 595–602 (2021).

13.   Routy, B. *et al.* Fecal microbiota transplantation plus anti-PD-1 immunotherapy in

advanced melanoma: a phase I trial. *Nat Med* **29**, 2121–2132 (2023).

14.   Zhao, S. *et al.* Adaptive evolution within the gut microbiomes of healthy people. *Cell Host*

Microbe **25**, 656-667.e8 (2019).

15.   Ghensi, P. *et al.* Strong oral plaque microbiome signatures for dental implant diseases

identified by strain-resolution metagenomics. *npj Biofilms Microbiomes* **6**, 1–12 (2020).

16.   Lloyd-Price, J. *et al.* Strains, functions and dynamics in the expanded Human

Microbiome Project. *Nature* **550**, 61–66 (2017).

17.   Porcari, S. *et al.* Key determinants of success in fecal microbiota transplantation: From

microbiome to clinic. *Cell Host & Microbe* **31**, 712–733 (2023).

18.   Ianiro, G. *et al.* Variability of strain engraftment and predictability of microbiome

composition after fecal microbiota transplantation across different diseases. *Nat Med* **28**,

1913–1923 (2022).

19.   Duvallet, C. *et al.* Framework for rational donor selection in fecal microbiota transplant

clinical trials. *PLOS ONE* **14**, e0222881 (2019).

20.   Aggarwala, V. *et al.* Precise quantification of bacterial strains after fecal microbiota

transplantation delineates long-term engraftment and explains outcomes. *Nat Microbiol* **6**,

1309–1318 (2021).

21.   Podlesny, D. *et al.* Identification of clinical and ecological determinants of strain

engraftment after fecal microbiota transplantation using metagenomics. *CR Med* **3**, (2022).

22.   Luo, C. *et al.* ConStrains identifies microbial strains in metagenomic datasets. *Nat*

Biotechnol **33**, 1045–1052 (2015).

23.   Smillie, C. S. *et al.* Strain-tracking reveals the determinants of bacterial engraftment in

the human gut following fecal microbiota transplantation. *Cell Host Microbe* **23**, 229-240.e5

(2018).

24.   Tamburini, F. B. *et al.* Precision Identification of Diverse Bloodstream Pathogens in the

Gut Microbiome. *Nat Med* **24**, 1809–1814 (2018).

25.   Schmidt, T. S. B. *et al.* Drivers and determinants of strain dynamics following fecal

microbiota transplantation. *Nat Med* **28**, 1902–1912 (2022).

26.   Beghini, F. *et al.* Integrating taxonomic, functional, and strain-level profiling of diverse

microbial communities with bioBakery 3. *Elife* **10**, e65088 (2021).

27.   Olm, M. R. *et al.* InStrain enables population genomic analysis from metagenomic data

and sensitive detection of shared microbial strains. *Nat Biotechnol* **39**, 727–736 (2021).

28.   Blanco-Míguez, A. *et al.* Extending and improving metagenomic taxonomic profiling with

uncharacterized species using MetaPhlAn 4. *Nat Biotechnol* **41**, 1633–1644 (2023).

29.   Van Rossum, T., Ferretti, P., Maistrenko, O. M. & Bork, P. Diversity within species:

interpreting strains in microbiomes. *Nat Rev Microbiol* **18**, 491–506 (2020).

30.   Gounot, J.-S. *et al.* Genome-centric analysis of short and long read metagenomes

reveals uncharacterized microbiome diversity in Southeast Asians. *Nat Commun* **13**, 6044

(2022).

31.   Sczyrba, A. *et al.* Critical Assessment of Metagenome Interpretation – a benchmark of

computational metagenomics software. *Nat Methods* **14**, 1063–1071 (2017).

32.   Vicedomini, R., Quince, C., Darling, A. E. & Chikhi, R. Strainberry: automated strain

separation in low-complexity metagenomes using long reads. *Nat Commun* **12**, 4485 (2021).

33. Tamburini, F. B. *et al.* Short- and long-read metagenomics of urban and rural South

African gut microbiomes reveal a transitional composition and undescribed taxa. *Nat*

Commun **13**, 926 (2022).

34. Ghazi, A. R., Münch, P. C., Chen, D., Jensen, J. & Huttenhower, C. Strain Identification

and Quantitative Analysis in Microbial Communities. *J Mol Biol* **434**, 167582 (2022).

35. Yan, Y., Nguyen, L. H., Franzosa, E. A. & Huttenhower, C. Strain-level epidemiology of

microbial communities and the human microbiome. *Genome Medicine* **12**, 71 (2020).

36. Tidjani Alou, M. *et al.* State of the Art in the Culture of the Human Microbiota: New

Interests and Strategies. *Clinical Microbiology Reviews* **34**, 10.1128/cmr.00129-19 (2020).

37. Kim, C. Y., Ma, J. & Lee, I. HiFi metagenomic sequencing enables assembly of accurate

and complete genomes from human gut microbiota. *Nat Commun* **13**, 6367 (2022).

38. Moss, E. L., Maghini, D. G. & Bhatt, A. S. Complete, closed bacterial genomes from

microbiomes using nanopore sequencing. *Nat Biotechnol* **38**, 701–707 (2020).

39. Gehrig, J. L. *et al.* Finding the right fit: evaluation of short-read and long-read

sequencing approaches to maximize the utility of clinical microbiome data. *Microbial*

Genomics **8**, 000794 (2022).

40. Agustinho, D. P. *et al.* Unveiling microbial diversity: harnessing long-read sequencing

technology. *Nat Methods* **21**, 954–966 (2024).

41. Kuleshov, V. *et al.* Synthetic long-read sequencing reveals intraspecies diversity in the

human microbiome. *Nat Biotechnol* **34**, 64–69 (2016).

42. Chen, L. *et al.* Short- and long-read metagenomics expand individualized structural

variations in gut microbiomes. *Nat Commun* **13**, 3175 (2022).

43. Tourancheau, A., Mead, E. A., Zhang, X.-S. & Fang, G. Discovering multiple types of

DNA methylation from bacteria and microbiome using nanopore sequencing. *Nat Methods* **18**,

491–498 (2021).

44.   Beaulaurier, J. *et al.* Metagenomic binning and association of plasmids with bacterial

host genomes using DNA methylation. *Nat Biotechnol* **36**, 61–69 (2018).

45.   Beaulaurier, J., Schadt, E. E. & Fang, G. Deciphering bacterial epigenomes using

modern sequencing technologies. *Nat Rev Genet* **20**, 157–172 (2019).

46.   Kong, Y. *et al.* Critical assessment of DNA adenine methylation in eukaryotes using

quantitative deconvolution. *Science* **375**, 515–522 (2022).

47.   Fang, G. *et al.* Genome-wide mapping of methylated adenine residues in pathogenic

Escherichia coli using single-molecule real-time sequencing. *Nat Biotechnol* **30**, 1232–1239

(2012).

48.   Sánchez-Romero, M. A. & Casadesús, J. The bacterial epigenome. *Nat Rev Microbiol*

18, 7–20 (2020).

49.   Kolmogorov, M. *et al.* metaFlye: scalable long-read metagenome assembly using repeat

graphs. *Nat Methods* **17**, 1103–1110 (2020).

50.   Wu, Y.-W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: an automated binning algorithm

to recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**, 605–607 (2016).

51.   Uritskiy, G. V., DiRuggiero, J. & Taylor, J. MetaWRAP—a flexible pipeline for genome-

resolved metagenomic data analysis. *Microbiome* **6**, 158 (2018).

52.   Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P. A. metaSPAdes: a new versatile

metagenomic assembler. *Genome Res* **27**, 824–834 (2017).

53.   Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-

node solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics **31**, 1674–1676 (2015).

54.   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. *Genome Res* **25**, 1043–1055 (2015).

55.   Faith, J. J. *et al.* Strain population structure varies widely across bacterial species and

predicts strain colonization in unrelated individuals. 2020.10.17.343640 Preprint at

https://doi.org/10.1101/2020.10.17.343640 (2020).

56.   Antipov, D., Raiko, M., Lapidus, A. & Pevzner, P. A. Metaviral SPAdes: assembly of

viruses from metagenomic data. *Bioinformatics* **36**, 4126–4129 (2020).

57.   Tang, X., Shang, J., Ji, Y. & Sun, Y. PLASMe: a tool to identify PLASMid contigs from

short-read assemblies using transformer. *Nucleic Acids Res* **51**, e83 (2023).

58.   Kent, A. G., Vill, A. C., Shi, Q., Satlin, M. J. & Brito, I. L. Widespread transfer of mobile

antibiotic resistance genes within individual gut microbiomes revealed through bacterial Hi-C.

Nat Commun **11**, 4379 (2020).

59.   Leylabadlo, H. E. *et al.* The critical role of *Faecalibacterium prausnitzii* in human health:

An overview. *Microbial Pathogenesis* **149**, 104344 (2020).

60.   Cani, P. D., Depommier, C., Derrien, M., Everard, A. & de Vos, W. M. *Akkermansia*

muciniphila: paradigm for next-generation beneficial microorganisms. *Nat Rev Gastroenterol*

Hepatol **19**, 625–637 (2022).

61.   Derrien, M., Belzer, C. & de Vos, W. M. *Akkermansia muciniphila* and its role in

regulating host functions. *Microbial Pathogenesis* **106**, 171–181 (2017).

62.   Bae, M. *et al. Akkermansia muciniphila* phospholipid induces homeostatic immune

responses. *Nature* **608**, 168–173 (2022).

63.   Wu, Z. *et al. Akkermansia muciniphila* Ameliorates *Clostridioides difficile* Infection in

Mice by Modulating the Intestinal Microbiome and Metabolites. *Front Microbiol* **13**, 841920

(2022).

64.   Zheng, M. *et al.* The role of *Akkermansia muciniphila* in inflammatory bowel disease:

Current knowledge and perspectives. *Front. Immunol.* **13**, (2023).

65.   Derosa, L. *et al.* Intestinal *Akkermansia muciniphila* predicts clinical response to PD-1

blockade in patients with advanced non-small-cell lung cancer. *Nat Med* **28**, 315–324 (2022).

66.   Everard, A. *et al.* Cross-talk between *Akkermansia muciniphila* and intestinal epithelium

controls diet-induced obesity. *Proc Natl Acad Sci U S A* **110**, 9066–9071 (2013).

67.   Cao, Y., Shen, J. & Ran, Z. H. Association between *Faecalibacterium prausnitzii*

Reduction and Inflammatory Bowel Disease: A Meta-Analysis and Systematic Review of the

Literature. *Gastroenterology Research and Practice* **2014**, e872725 (2014).

68.   Miquel, S. *et al. Faecalibacterium prausnitzii* and human intestinal health. *Current*

Opinion in Microbiology **16**, 255–261 (2013).

69.   Zhao, H. *et al.* Systematic review and meta-analysis of the role of *Faecalibacterium*

prausnitzii alteration in inflammatory bowel disease. *Journal of Gastroenterology and*

Hepatology **36**, 320–328 (2021).

70.   Sokol, H. *et al. Faecalibacterium prausnitzii* is an anti-inflammatory commensal

bacterium identified by gut microbiota analysis of Crohn disease patients. *Proceedings of the*

National Academy of Sciences **105**, 16731–16736 (2008).

71.   Zhang, J. *et al.* An immune-competent human gut microphysiological system enables

inflammation-modulation by *Faecalibacterium prausnitzii. npj Biofilms Microbiomes* **10**, 1–11

(2024).

72.   Quévrain, E. *et al.* Identification of an anti-inflammatory protein from *Faecalibacterium*

prausnitzii, a commensal bacterium deficient in Crohn's disease. *Gut* **65**, 415–425 (2016).

73.   Ze, X., Duncan, S. H., Louis, P. & Flint, H. J. *Ruminococcus bromii* is a keystone species

for the degradation of resistant starch in the human colon. *ISME J* **6**, 1535–1543 (2012).

74.   Abell, G. C. J., Cooke, C. M., Bennett, C. N., Conlon, M. A. & McOrist, A. L. Phylotypes

related to *Ruminococcus bromii* are abundant in the large bowel of humans and increase in

response to a diet high in resistant starch. *FEMS Microbiology Ecology* **66**, 505–515 (2008).

75.   Chen-Liaw, A. *et al.* Gut microbiota strain richness is species specific and affects

engraftment. *Nature* **637**, 422–429 (2025).

76.   Durrant, M. G. & Bhatt, A. S. Microbiome genome structure drives function. *Nat Microbiol*

4, 912–913 (2019).

77.   Behling, A. H. *et al.* Horizontal gene transfer after faecal microbiota transplantation in

adolescents with obesity. *Microbiome* **12**, 26 (2024).

78.   Moss, E. L. *et al.* Long-term taxonomic and functional divergence from donor bacterial

strains following fecal microbiota transplantation in immunocompromised patients. *PLOS*

ONE **12**, e0182585 (2017).

79.   Kong, Y., Mead, E. A. & Fang, G. Navigating the pitfalls of mapping DNA and RNA

modifications. *Nat Rev Genet* **24**, 363–381 (2023).

80.   Noinaj, N., Guillier, M., Travis J. Barnard & Buchanan, S. K. TonB-Dependent

Transporters: Regulation, Structure, and Function. *Annual Review of Microbiology* **64**, 43–60

(2010).

81.   Dingemans, J. *et al.* The deletion of TonB-dependent receptor genes is part of the

genome reduction process that occurs during adaptation of Pseudomonas aeruginosa to the

cystic fibrosis lung. *Pathog Dis* **71**, 26–38 (2014).

82.   Babakhani, S. & Oloomi, M. Transposons: the agents of antibiotic resistance in bacteria.

Journal of Basic Microbiology **58**, 905–917 (2018).

83.   Partridge, S. R., Kwong, S. M., Firth, N. & Jensen, S. O. Mobile Genetic Elements

Associated with Antimicrobial Resistance. *Clinical Microbiology Reviews* **31**,

10.1128/cmr.00088-17 (2018).

84.   Maqbool, A. *et al.* The substrate-binding protein in bacterial ABC transporters: dissecting

roles in the evolution of substrate specificity. *Biochemical Society Transactions* **43**, 1011–

1017 (2015).

85.   Vasu, K. & Nagaraja, V. Diverse Functions of Restriction-Modification Systems in

Addition to Cellular Defense. *Microbiology and Molecular Biology Reviews* **77**, 53–72 (2013).

86.   Beaulaurier, J. *et al.* Single molecule-level detection and long read-based phasing of

epigenetic variations in bacterial methylomes. *Nat Commun* **6**, 7438 (2015).

87.   Srikhanta, Y. N., Fox, K. L. & Jennings, M. P. The phasevarion: phase variation of type III

DNA methyltransferases controls coordinated switching in multiple genes. *Nat Rev Microbiol*

8, 196–206 (2010).

88.   Oliveira, P. H. & Fang, G. Conserved DNA Methyltransferases: A Window into

Fundamental Mechanisms of Epigenetic Regulation in Bacteria. *Trends in Microbiology* **29**,

28–40 (2021).

89.   Brunet, Y. R., Bernard, C. S., Gavioli, M., Lloubès, R. & Cascales, E. An epigenetic

switch involving overlapping fur and DNA methylation optimizes expression of a type VI

secretion gene cluster. *PLoS Genet* **7**, e1002205 (2011).

90.   Lim, H. N. & van Oudenaarden, A. A multistep epigenetic switch enables the stable

inheritance of DNA methylation states. *Nat Genet* **39**, 269–275 (2007).

91.   Seib, K. L., Srikhanta, Y. N., Atack, J. M. & Jennings, M. P. Epigenetic Regulation of

Virulence and Immunoevasion by Phase-Variable Restriction-Modification Systems in

Bacterial Pathogens. *Annu Rev Microbiol* **74**, 655–671 (2020).

92.   Huang, X. *et al.* Prevalence of phase variable epigenetic invertons among host-

associated bacteria. *Nucleic Acids Res* **48**, 11468–11485 (2020).

93.   Oliveira, P. H. *et al.* Epigenomic characterization of *Clostridioides difficile* finds a

conserved DNA methyltransferase that mediates sporulation and pathogenesis. *Nat Microbiol*

5, 166–180 (2020).

94.   Furuta, Y. *et al.* Methylome Diversification through Changes in DNA Methyltransferase

Sequence Specificity. *PLOS Genetics* **10**, e1004272 (2014).

95.   Manso, A. S. *et al.* A random six-phase switch regulates pneumococcal virulence via

global epigenetic changes. *Nat Commun* **5**, 5055 (2014).

96.   Lopez-Siles, M., Duncan, S. H., Garcia-Gil, L. J. & Martinez-Medina, M.

953   *Faecalibacterium prausnitzii*: from microbiology to diagnostics and prognostics. *ISME J* **11**,

954   841–852 (2017).

955   97.   Wu, H. *et al.* Strategies for high cell density cultivation of *Akkermansia muciniphila* and

956   its potential metabolism. *Microbiol Spectr* **12**, e0238623 (2024).

957   98.   Zhernakova, D. V. *et al.* Host genetic regulation of human gut microbial structural

958   variation. *Nature* **625**, 813–821 (2024).

959   99.   West, P. T., Chanin, R. B. & Bhatt, A. S. From genome structure to function: insights into

960   structural variation in microbiology. *Current Opinion in Microbiology* **69**, 102192 (2022).

961   100.   Ji, P., Zhang, Y., Wang, J. & Zhao, F. MetaSort untangles metagenome assembly by

962   reducing microbial community complexity. *Nat Commun* **8**, 14306 (2017).

963   101.   Bokulich, N. A. *et al.* Antibiotics, birth mode, and diet shape microbiome maturation

964   during early life. *Sci Transl Med* **8**, 343ra82 (2016).

965   102.   Yassour, M. *et al.* Natural history of the infant gut microbiome and impact of antibiotic

966   treatment on bacterial strain diversity and stability. *Sci Transl Med* **8**, 343ra81 (2016).

967   103.   Jiang, X. *et al.* Invertible promoters mediate bacterial phase variation, antibiotic

968   resistance, and host adaptation in the gut. *Science* **363**, 181–187 (2019).

969   104.   Sarkar, A. *et al.* Microbial transmission in the social microbiome and host health and

970   disease. *Cell* **187**, 17–43 (2024).

Materials and Methods

Fecal sample collection and DNA extraction

The adult fecal samples were from a previous study[1] with Institutional Review Board approval (rCDI: HS# 11-01669, IBD: HREC/13/SVH/69). Fecal DNA was extracted with a QIAamp PowerFecal Pro DNA Kit (Qiagen, 51804) according to the manufacturer's instructions. Briefly, 64 - 436 (average is 181) mg of fecal sample, and 800 µl of Solution CD1 were added to the Bead Tube (provided). After gentle vortexing, the sample was homogenized with the TissueLyser II machine (Qiagen, 85300) for 10 min at 25 Hz speed (5 min each side). The homogenized samples were centrifuged to collect the supernatant at 15,000 x g for 1 min. The subsequent column-based extraction steps were performed as described in the QIAamp PowerFecal Pro DNA Kit protocol.

Library preparation and sequencing

Nanopore long-read sequencing

Native libraries were prepared using the Oxford Nanopore Technology Inc. (New York, NY, USA) Native barcoding kit (EXP-NBD104 and 114 for R9.4.1 flow cells, and SQK-NBD114.24 for R10.4.1 flow cells), Ligation Sequencing Kit (SQK-LSK109 for R9, and SQK-LSK114 for R10.4.1). Prior to library preparation, genomic DNA (gDNA) was sheared using Covaris g-tubes (Covaris, Woburn, MA, USA 520079) at 6000 RPM in an Eppendorf 5424R centrifuge (Eppendorf, Enfield, CT, USA, 5406000640) and cleaned up with AMPure XP beads (Beckman Coulter, Indianapolis, IN, USA, A63882) at 0.8 x concentration. Whole genome amplification (WGA) samples were prepared using REPLI-g Mini Kits (QIAGEN, Hilden, Germany, 150023) according to the protocol with 12.5 ng of input DNA per sample and 16 h incubation at 30 °C, followed by denaturation at 65 °C for 3 minutes. Next, WGA samples were treated with T7

endonuclease I (NEB M0302S) to maximize nanopore sequencing yield according to ONT

documentation, then cleaned up with AMPure XP beads as above. Subsequently, samples were

processed the same as above native libraries. Samples were sequenced on R9.4.1 and R10.4.1

MinION flow cells and post-run basecalled using Guppy (v5.0.7 for R9.4.1 data and 6.3.4 for

R10.4.1 data)[2] with the SUP model dna_r9.4.1_450bps_sup.cfg and

dna_r10.4.1_e8.2_400bps_sup.cfg, respectively.

Illumina short-read sequencing

Library construction for Illumina sequencing was executed utilizing the seqWell Plexwell 384 kit.

DNA was systematically barcoded, and ligation products underwent purification. Size selection

and PCR amplification of the DNA fragments followed. The prepared library was then

sequenced on an Illumina HiSeq platform, generating paired-end reads of 150 base pairs (bp)

each.

PacBio long-read sequencing

DNA SMRTbell libraries were prepared according to the manufacturer's instructions for the

SMRTbell Express Template Prep Kit 2.0, then each library was annealed to sequencing primer

together with the Sequel 2.1 polymerase, followed by loading to the a 8M Sequel II sequencing

chip (for Sequel II) and 25M Revio sequencing chip.

Metagenomic assembly, strain-level separation and taxonomic annotation

Long-read metagenomic assembly

Initially, the raw long reads were subjected to a metagenomic assembly for long-read MAGs

process using the metaFlye tool (v2.9)[3] with "---nano-raw --meta --keep-haplotypes --iterations

2". The resulting metagenome-assembled genomes (MAGs) were partitioned into individual bins

corresponding to the species level using the MaxBin (v2.0) software[4]. To annotate the species-

level bins, we applied the Genome Taxonomy Database Toolkit (GTDB-tk v1.6.0)[5]. For each

species-level bin, we utilized Strainberry (v1.1)[6] to separate the contigs into the strain level

based on haplotype phasing. Subsequently, these separated contigs were binned to strain level

using the "binning" module of metaWRAP (v1.3.2)[7], which could combine the results from three

metagenomic binning software - MaxBin2[4], metaBAT2[8], and CONCOCT[9]. The quality of the

reconstructed long-read MAGs was evaluated using the "bin_refinement" module in metaWRAP

(v1.3.2)[7], which combines the results of multiple binning algorithms and selects high-quality bins

based on completeness and contamination estimates, as well as estimated by CheckM

(v.1.1.3)[10]. The strain-level long-read MAGs with completeness of more than 10% were selected

for further analysis. We used viralVerify (v1.1)[11] and PLASMe (v1.1)[12] to identify plasmids in

each assembly. For viralVerify, only contigs shorter than 500 kbp with prediction scores greater

than five were considered. Annotations labeled as 'Plasmid' or 'Uncertain - plasmid or

chromosomal' were classified as plasmids. For PLASMe, only results with high precision were

included. Overlapping plasmids identified by both methods were treated as the final results.

Short-read metagenomic assembly

We conducted a metagenomic assembly of the short reads using metaSPAdes (v3.15.2)[13] and

MEGAHIT (v1.1.3)[14] with default settings to construct short-read MAGs. Subsequently, we

utilized metaWRAP (v1.3.2)[7] to bin and refine the reconstructed short-read MAGs. The

completeness and contamination level for the short-read MAGs were estimated using CheckM

(v.1.1.3)[10], and only those with completeness of more than 10% were selected for further

analysis. Taxonomic assignment of the reconstructed short-read MAGs was performed using

GTDB-tk (v1.6.0)[5].

1041

Evaluation of long-read MAGs

The previous study identified a total of 156 unique strains, which were from 156 different

bacterial isolates previously isolated from three distinct donors[1]. Specifically, 136 bacterial

isolates were from Donor 1 (D1), 13 were from Donor 2 (D2), and 7 were from Donor 3 (D3).

The 156 bacterial isolates were either directly isolated from the donors or from post-FMT

recipients, which could be traced back to their original donors. The resulting assemblies for

each isolate serve as reference individual genomes. To detect shared strains between long-read

MAGs/short-read MAGs and genomes of bacterial isolates, we utilized fastANI (1.32)[15] for

average nucleotide identity (ANI) calculation, applying a 99.4 % cut-off[3]. We assessed the

contamination level by mapping long-read and short-read MAG to its corresponding isolate

genome, calculated as the percentage of unaligned bases. Mummer (v4.0.0b)[17] was employed

to identify the mapping regions between long-read MAG/short-read MAG and its corresponding

genome of isolate. We also assessed the genomic similarity of shared strains between the long-

read MAGs/short-read MAGs and their corresponding isolate genomes determined by k-mer

similarity as in the previous study[1], with the following formula:

$$\text{Kmer Similarity} = \frac{Number\ of\ shared\ kmer\ between\ Genome\ A\ and\ Genome\ B}{Lesser\ total\ kmer\ count\ of\ either\ Genome\ A\ and\ Genome\ B}$$

Where:

  K-mer size is set to 31.

  Shared k-mers are those that appear in both Genome A and B.

  Total k-mer count is the count of all distinct 31-mers in a genome.

  The k-mer similarity between two genomes A and B was determined by generating a

hash for genomes with k-mer size 31 and quantifying the proportion of k-mers shared in both

genomes A and B divided by the lesser total k-mer count of either

1064

1065 **The proportion of reads associated with each strain/MAG**

1066 We analyzed long-read and short-read metagenomic sequencing data from D1 by mapping to a

1067 merged set of genomes, which included genomes from bacterial isolates and uncultured long-

1068 read MAGs or short-read MAGs. For the long-read data, we used 136 genomes from bacterial

1069 isolates and 105 uncultured long-read MAGs. The proportion of reads associated with each

1070 strain/MAG was determined by calculating the percentage of metagenomic reads that were

1071 successfully mapped to each genome/MAG in this merged set. Similarly, the short-read data

1072 were aligned to the same 136 genomes of bacterial isolates, supplemented with 120 uncultured

1073 short-read MAGs, to quantify the proportion of reads associated with each strain/MAG.

1074

1075 **LongTrack framework designation for FMT strain tracking based on strain-unique k-mers**

1076 The LongTrack framework for FMT strain tracking follows a three-step algorithmic process (**Fig.**

1077 **1C**). LongTrack strain tracking relies on strain-unique k-mers, built on previous methods[1], and

1078 incorporates important methodological adaptations compared to previous studies to ensure the

1079 reliable selection of unique k-mers and to minimize false positives.

1080 *Step 1: Identification of strain-unique k-mers from each long-read MAG*

1081 a. Extract strain-unique k-mers from each long-read MAG

1082 Following established methods[1], the LongTrack framework extracts k-mers (k=31) for each

1083 strain-level long-read MAG from the donor or pre-FMT recipient using KMC (version 3.0.0)[18].

1084 Strain-unique k-mers are then identified by eliminating common k-mers shared in 100,000

1085 bacterial genomes in the NCBI database (as of 2018-11-20), 116 unrelated metagenomic

1086 samples from a previous study[1], and other long-read MAGs within the same FMT case.

b. Obtain strain-unique k-mers in co-existing strains

Relying solely on strain-unique k-mers identified in Step 1a can lead to false strain-tracking

results due to the varying completeness of long-read MAGs, especially for similar co-existing

strains. For instance, as shown in **Fig. 1C**, Strains A1 and A2 both belong to Species A and are

co-existing strains from the same donor. If the k-mers identified for Strain A1 in Step 1a

correspond to genomic regions missing in Strain A2 due to incomplete assembly, these k-mers

may also appear in Strain A2 simply because of the high genomic similarity between the two

strains. In such cases, the k-mers are not truly unique to Strain A1 but are instead masked in

Strain A2 by assembly gaps, leading to unreliable results.

To address this issue, LongTrack introduces methodological adaptations to ensure the reliable

selection of strain-unique k-mers and to minimize false positives. Specifically, for similar co-

existing strains within a single fecal sample or closely related strains found separately in the

donor and pre-FMT recipient samples, LongTrack applies a filter to retain only those unique k-

mers originating from genomic regions shared among multiple co-existing strains in same FMT

case. Mummer (v4.0.0b)[17] is used to identify these common genomic regions.

This approach prevents the erroneous selection of k-mers that might seem strain-specific but

are unreliable due to MAG incompleteness—a critical distinction between high-purity long-read

MAGs and complete genomes from bacterial isolates. The resulting retained strain-unique k-

mers serve as markers for each long-read MAG, enabling reliable strain tracking in subsequent

steps.

Step 2: Assign short reads with final strain-unique k-mers to metagenomic samples

In the second step, the final strain-unique k-mers identified in Step 1b were utilized to assign

short reads from post-FMT recipient microbiome data to their corresponding long-read MAGs.

Short reads containing strain-unique k-mers were mapped back to their respective long-read

MAGs using Bowtie 2.4.1[19], with parameters optimized for accuracy ('very-sensitive,' 'no-mixed,'

and 'no-discordant'). The successful alignment of short reads to genomic regions containing

strain-unique k-mers confirmed the presence of the corresponding MAG in the post-FMT

recipient, facilitating accurate strain tracking.

Step 3: Discard results with low confidence score

To ensure the reliability of strain tracking, we calculated a confidence score by comparing read

enrichment in five unrelated metagenomic samples (negative controls) as done in the previous

study[1]. Results with low confidence scores (< 95% confidence) were discarded to maintain the

accuracy of strain tracking.

1120

Evaluation of LongTrack for FMT strain tracking

Evaluation of LongTrack for FMT strain tracking through comparison with ground truth

For a comprehensive assessment of LongTrack for FMT strain tracking, we utilized the shared

strains between long-read MAGs/short-read MAGs and genomes of bacterial isolates in four

FMT cases of rCDI. This evaluation was conducted by comparing tracking results in the post-

FMT recipients at different time points based on Strainer with genomes of bacterial isolates

(considered the ground truth) and LongTrack/Strainer with short-read MAGs. We applied two

key metrics:

1. Precision: Precision quantifies the proportion of correct tracking results (True Positives,

TP) relative to all tracking results predicted by the LongTrack/Strainer with short-read

MAGs. It is calculated as TP divided by the sum of True Positives (TP) and False

Positives (FP).

2. Sensitivity: Sensitivity evaluates the effectiveness of LongTrack/Strainer with short-read

MAGs in accurate strain tracking. It is determined by the ratio of True Positives (TP) to

the total number of True Positives (TP) and False Negatives (FN).

The accuracy of strain tracking based on LongTrack/Strainer with short-read MAGs was

evaluated using the formula: the number of consistent tracking results between ground truth and

LongTrack/Strainer with short-read MAGs divided by the total number of tracking results,

expressed as a percentage. The tracking results were presented as a heatmap using the R

package "pheatmap v1.0.12".

Evaluation of FMT strain tracking between LongTrack and StrainFinder, which serves as a

representative methods of SNP-based tracking

To evaluate FMT strain tracking between LongTrack and StrainFinder[20], we utilized *in*

silico spike-in datasets (**Supp Fig. S6**). For donor D1, we spiked in 3% *Clostridium*

tyrobutyricum at 60x depth and 2% *Turicibacter sanguinis* at 40x depth into long-read

microbiome dataset. The simulated Nanopore long reads were generated using nanosim (v

3.0.0)[21] for metagenomic assembly. The error profile was trained using real sequencing data

from a large number of nanopore sequencing reads, and the read length distribution was

modeled to match that of the actual Donor 1 sequencing data. Similarly, we produced simulated

short-read data with identical abundances of these strains for donor D1 in the short-read

microbiome datasets. For the post-FMT recipient R1, we spiked in short reads to the short-read

metagenomic sequencing data, which was used for strain tracking. We designed two distinct

spike-in datasets: A) The first dataset A (low abundance) featured a composition of 0.25% (5x

depth) *Clostridium tyrobutyricum*, along with paired short reads, each having a length of 150 bp.

Furthermore, this dataset included 0.1% (2x depth) *Turicibacter sanguinis*, with paired short

reads, also 150 bp in length. B) The second dataset B (high abundance) presented a

composition of 2.5% (50x depth) *Clostridium tyrobutyricum*, accompanied by paired short reads,

each measuring 150 bp in length. Additionally, this dataset contained 1% (20x depth)

Turicibacter sanguinis, once again with paired short reads, each 150 bp in length. We simulated

paired short reads with a length of 150 bp using randomreads.sh from the bbmap package

(v38.86)[22]. Subsequently, LongTrack and StrainFinder[20] were performed for strain tracking

analysis.

Evaluation of LongTrack for FMT strain tracking when there are high-similarity strains present in

between the donor and pre-FMT samples.

To evaluate LongTrack/Strainer with short-read MAGs for FMT strain tracking when

there are high-similarity strains present in between the donor and pre-FMT samples, we spiked-

in ten pairs of strains, each with high similarity (ANI>99.5%), from ten different species, one

strain to D1 and the other to R1 pre-FMT, similarly for all the ten pairs **(Fig. 3F)**. All twenty

strains were added to R1 post-FMT. To facilitate metagenomic assembly, we generated

simulated long-read data/short-read data for D1 and R1 pre-FMT into their long-read/short-read

microbiome datasets. Furthermore, we simulated short-read paired reads, each with a length of

150 bp, to R1 post-FMT short-read microbiome datasets. The simulated Nanopore long reads

were generated using the nanosim (v 3.0.0)[21]. The short reads were generated using the

randomreads.sh from the bbmap package (v38.86)[22]. LongTrack/Strainer with short-read MAGs

was performed for strain tracking analysis.

Evaluation of LongTrack for FMT strain tracking when there are co-existing strains of the same

species in one donor sample.

To evaluate LongTrack/Strainer with short-read MAGs for FMT strain tracking when

there are co-existing strains of the same species in one donor sample, we spiked-in three

strains of *Phocaeicola massiliensis* (Strains 1-3) to D1 and a fourth strain of *P. massiliensis*

(Strain 4) to R1 pre-FMT **(Fig. 3H)**. For metagenomic assembly, long-read data/short-read data

were generated for these strains in D1 and R1 pre-FMT microbiome datasets. Furthermore, we

simulated short-read paired reads of 4 strains (Strains 1-4), each with a length of 150 bp, to R1

post-FMT short-read microbiome datasets. The simulated Nanopore long reads with an average

length of 5 kb, were generated using the nanosim (v 3.0.0)[21]. The short reads were generated

using the randomreads.sh from the bbmap package (v38.86)[22]. LongTrack/Strainer with short-

read MAGs was performed for strain tracking analysis.

1188

Application of LongTrack: strain tracking in rCDI and IBD cases, reverse tracking and

identification of novel strains

Application of LongTrack in strain tracking of rCDI cases

We initially retrieved long-read MAGs from 3 donors and 4 pre-FMT recipients. Using

LongTrack, we examined whether these MAGs were present in the metagenomic sequencing

data of the corresponding post-FMT recipients across multiple time points.

Application of LongTrack in strain tracking of IBD cases

In the IBD cases, the recipients (R5 and R6) had received fecal microbiota

transplantation from six donors (D4 - D9), and strain tracking was performed for long-read

MAGs from two donors (D4 and D5). To avoid the influence of common k-mers from the

remaining four donors, we further excluded the k-mers that were present in the short-read

metagenomic sequencing data of the other four donors. Subsequently, we assessed whether

these MAGs were present in the metagenomic sequencing data of the corresponding post-FMT

recipients across multiple time points using LongTrack.

LongTrack-based reverse tracking and novel strain identification

We initially retrieved 402 uncultured long-read MAGs from post-FMT recipients (R1-R4). We

then identified long-read MAGs shared between the post-FMT recipient and the respective

donor or pre-FMT recipient using an ANI exceeding 99.4% in the same FMT cases. We

confirmed that 39 long-read MAGs are unique to the post-FMT recipients and investigated

whether they were present in the metagenomic sequencing data of the corresponding donor

and pre-FMT recipient using LongTrack. If the long-read MAGs were not tracked in microbiome

data of donor, pre-FMT or any earlier time points of post-FMT, we considered them to be novel

strains.

1212

Detection of structural variation in engrafted strains using long-read sequencing

We utilized NGMLR (v 0.2.7)[23] to perform the alignment of long-read microbiome data of post-

FMT recipients (R1-R4) across various time points to the uncultured long-read MAGs and

isolated genomes from the respective rCDI donors (D1-D3). Following the alignment process,

we employed Sniffles (v1.0.12)[24] for the detection of structural variations (SVs) within the long-

read microbiome data. To ensure the reliability of our SV detection, we applied several filtering

criteria: 1) We excluded SVs involving multiple strains within the same species, as these are

more prone to mismatches and may not accurately represent structural variations. 2) For

mapping quality control, we retained aligned reads with identity greater than 98%, length more

than 1000 bp, and MAPQ (Mapping Quality) score greater than 60. 3) We kept SVs that had a

coverage depth of greater than 5 in long-read metagenomic sequencing data of donor and at

least one post-FMT, ensuring that the detected structural variations were well-supported by

sequencing data.

1226

Validation of structural variations using isolate long-read sequencing

We collected 18 isolates for SV validation, including:

○ 2 *Alistipes onderdonkii* isolates: 1 from the donor and 1 from 8 weeks post-FMT.

○ 4 *Bacteroides stercoris* isolates: 2 from the donor, 1 from 8 weeks post-FMT, and 1 from

6 months post-FMT.

○ 5 *Bifidobacterium adolescentis* isolates: 2 from the donor, 1 from 8 weeks post-FMT, and

2 from 6 months post-FMT.

○ 7 *Parabacteroides merdae* isolates: 3 from the donor, 2 from 8 weeks post-FMT, and 2

from 6 months post-FMT.

Native libraries were prepared using the Oxford Nanopore Technologies Native Barcoding Kit

(SQK-NBD114.24) for R10.4.1 flow cells and the Ligation Sequencing Kit (SQK-LSK114).

Samples were sequenced on R10.4.1 MinION flow cells, and basecalling was performed post-

run using Guppy (v6.3.4)[2] with the SUP model dna_r10.4.1_e8.2_400bps_sup.cfg.

For alignment, we utilized NGMLR (v0.2.7)[23] to map the single isolate long-read sequencing

data to the corresponding long-read MAGs. SVs were detected using Sniffles (v1.0.12)[24]. We

then cross-validated the detected SVs from metagenomic sequencing data to determine if they

could also be identified in the single isolate long-read sequencing data.

1244

Methylation analysis using long read sequencing

De novo methylation motif discovery using Nanopore long-read sequencing

We utilized Nanodisco (v1.0.2)[25] to identify *de novo* methylation motifs in long-read MAGs from

the D1 . First, raw fast5 reads from both native and WGA metagenomic sequencing data of D1

were aligned to the long-read MAGs with "nanodisco preprocess". The alignments for each

long-read MAG were then segregated. Only long-read MAGs exhibiting a coverage above 20 x

were retained to following analysis. We then computed current differences (PA) between the

native (retaining methylation) and WGA (methylation-free) data for each genomic position using

"nanodisco difference". To conclude, "nanodisco motif and refine" was employed for *de novo*

methylation motif discovery based on these current differences. For D1 5-year, R1 post-FMT 8-

week, and R1 post-FMT 5-year, the raw fast5 reads from native microbiome data were aligned

to the retained long-read MAGs from D1 with "nanodisco preprocess" and separated the

alignments into each MAG. Subsequently, we computed the current differences by comparing

the native data with the WGA data from D1 for each MAG using "nanodisco difference".

"Nanodisco refine" was employed to verify the presence of D1 detected methylation motifs in the

samples of D1 5-year, R1 post-FMT 8-week, and R1 post-FMT 5-year based on the current

differences.

De novo methylation motif discovery using PacBio long-read sequencing

For PacBio long-read data of D1 (generated by the Sequel II platform), the raw reads pre-

processing was performed with SMRTlink (v8.0). Raw multiplexed files were first de-multiplexed

with lima. We mapped the subreads to long-read MAGs using pbalign (v0.4.1) with default

parameters and separated the alignments into each MAG. The IPD ratio was calculated with

ipdSummary (v2.4.1) from the alignments for each MAG. The *de novo* methylation motifs were

detected by using motifMaker (in SMRTlink v8.0 package) with default parameters. For PacBio

long-read data of D1 5-year, R1 8-week and R1 5-year (generated by the Revio platform), the

raw reads pre-processing was performed with SMRTlink (v13.0), and methylation motif

detection was performed by our in-house scripts.

1272

Method reference

1. Aggarwala, V. *et al.* Precise quantification of bacterial strains after fecal microbiota

transplantation delineates long-term engraftment and explains outcomes. *Nat. Microbiol.* **6**,

1309–1318 (2021).

2. Wick, R. R., Judd, L. M. & Holt, K. E. Performance of neural network basecalling tools for

Oxford Nanopore sequencing. *Genome Biol.* **20**, 129 (2019).

3. Kolmogorov, M. *et al.* metaFlye: scalable long-read metagenome assembly using repeat

graphs. *Nat. Methods* **17**, 1103–1110 (2020).

4. Wu, Y.-W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: an automated binning algorithm to

recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**, 605–607 (2016).

5. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927 (2020).

6. Vicedomini, R., Quince, C., Darling, A. E. & Chikhi, R. Strainberry: automated strain

separation in low-complexity metagenomes using long reads. *Nat. Commun.* **12**, 4485 (2021).

7. Uritskiy, G. V., DiRuggiero, J. & Taylor, J. MetaWRAP—a flexible pipeline for genome-

resolved metagenomic data analysis. *Microbiome* **6**, 158 (2018).

8. Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient genome

reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

9. Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition. *Nat. Methods*

11, 1144–1146 (2014).

10.   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. *Genome Res.* **25**, 1043–1055 (2015).

11.   Antipov, D., Raiko, M., Lapidus, A. & Pevzner, P. A. Metaviral SPAdes: assembly of

viruses from metagenomic data. *Bioinforma. Oxf. Engl.* **36**, 4126–4129 (2020).

12.   Tang, X., Shang, J., Ji, Y. & Sun, Y. PLASMe: a tool to identify PLASMid contigs from

short-read assemblies using transformer. *Nucleic Acids Res.* **51**, e83 (2023).

13.   Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P. A. metaSPAdes: a new versatile

metagenomic assembler. *Genome Res.* **27**, 824–834 (2017).

14.   Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-

node solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinforma. Oxf. Engl. **31**, 1674–1676 (2015).

15.   Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High

throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat.*

Commun. **9**, 5114 (2018).

16.   Rodriguez-R, L. M. *et al.* An ANI gap within bacterial species that advances the

definitions of intra-species units. 2022.06.27.497766 Preprint at

https://doi.org/10.1101/2022.06.27.497766 (2023).

17.   Marçais, G. *et al.* MUMmer4: A fast and versatile genome alignment system. *PLOS*

Comput. Biol. **14**, e1005944 (2018).

18.   Kokot, M., Dlugosz, M. & Deorowicz, S. KMC 3: counting and manipulating k-mer

statistics. *Bioinforma. Oxf. Engl.* **33**, 2759–2761 (2017).

19.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat.*

Methods **9**, 357–359 (2012).

20.   Smillie, C. S. *et al.* Strain-tracking reveals the determinants of bacterial engraftment in

the human gut following fecal microbiota transplantation. *Cell Host Microbe* **23**, 229-240.e5

(2018).

21.   Yang, C., Chu, J., Warren, R. L. & Birol, I. NanoSim: nanopore sequence read simulator

based on statistical characterization. *GigaScience* **6**, gix010 (2017).

22.   Bushnell, B. BBMap: A Fast, Accurate, Splice-Aware Aligner. (2014).

23.   Sedlazeck, F. J. *et al.* Accurate detection of complex structural variations using single

molecule sequencing. *Nat. Methods* **15**, 461–468 (2018).

24.   Smolka, M. *et al.* Detection of mosaic and population-level structural variants with

Sniffles2. *Nat. Biotechnol.* 1–10 (2024) doi:10.1038/s41587-023-02024-y.

25.   Tourancheau, A., Mead, E. A., Zhang, X.-S. & Fang, G. Discovering multiple types of

DNA methylation from bacteria and microbiome using nanopore sequencing. *Nat. Methods*

18, 491–498 (2021).

1330

1331

Fig.1

A   **FMT samples with cultured isolates**

PreFMT (Day 0)   Week 1   Year 5

Donors

Recipients

🧫   Isolate culture for genome assembly

★   Long-read sequencing for MAGs

▲   Short-read sequencing for strain tracking

B   **High purity, strain level long-read MAGs**

Isolate genomes

Long-read MAGs

Method validation

LongTrack

C   **LongTrack enables reliable FMT strain tracking**

Species A   Species B

Strain A1   Strain A2   Strain B1   Strain B2

Step 1a. Extract strain-unique k-mers from each LR MAG

Step 1b. Obtain strain-unique k-mers in co-existing strains by retaining the k-mers from shared genome regions

Donor   PreFMT   PostFMT Week 1   Year 5

Step 2. Assign short reads with strain-unique k-mers to each metagenomic sample

	Donor	PreFMT	PostFMT Week 1	Year 5
Strain A1	🟩	⬜	🟩	🟩
Strain A2	🟩	⬜	🟩	⬜
Strain B1	🟩	⬜	🟩	⬜
Strain B2	⬜	🟩	⬜	⬜

Step 3. Discard results with low confidence score and obtain Presence/absence Table

Strain can be detected   🟩 Yes   ⬜ No

D   **LongTrack resolves intra-strain structural variations in engrafted strains**

Long reads

Genotype 1 (G1)

Genotype 2 (G2)

Donor   G1

PostFMT Week 1   G2 / G1

PostFMT Year 5   G2 / G1

Fig.2

A Long-read MAGs · Donors & Recipients · Isolate genomes

B Donor 1

C

D

E Donor 1 Order taxon — Presence / Absence

F Species with single strain · Species with multiple strains

G Strain can be recovered correctly — Yes / No

Fig.3

Fig.4

Fig.5

A

Parabacteroides merdae

Short-read Alignment

Nanopore Long-read Alignment

PacBio Long-read Alignment

43bp Deletion

Alistipes putredinis

Short-read Alignment

Nanopore Long-read Alignment

PacBio Long-read Alignment

1250bp Insertion

B

Strain	Type	Donor	Early	Late	Gene/downstream gene
Agathobacter rectalis D1	INV			*	IR: DNA/RNA helicase
	INS			*	IR: Type II toxin-antitoxin system prevent-host-death family
	DEL			*	IR: Nucleotidyltransferase domain-containing protein
	INS			*	IR: Transcriptional regulator
✚ ***Alistipes onderdonkii* D1**	INV				GR: Restriction endonuclease subunit S
	INV				IR: Transcription antitermination protein RfaH
	INS				IR: Hypothetical protein
	INV				IR: Hypothetical protein
Alistipes putredinis D1	DEL				GR: IS110 family transposase ISCaa7
	INS				IR: Hypothetical protein
	INV				GR: Type-1 restriction enzyme EcoKI specificity protein
	INS				GR: OmpA family protein
	DEL				IR: DUF4836 family protein
	INS				IR: Hypothetical protein
	INV				IR: Site-specific integrase
	INS				GR: Hypothetical protein
	INS				IR: Lysine 2,3-aminomutase
	INS				IR: Peptide chain release factor RF2
✚ ***Bacteroides stercoris* D1**	INV				GR: TonB-dependent receptor SusC
Bacteroides uniformis D1	INV				IR: UpxY family transcription antiterminator
	INS				GR: Alkaline phosphatase
✚ ***Bifidobacterium adolescentis* D1**	INS			*	GR: H(+)/Cl(-) exchange transporter ClcA
✚ ***Parabacteroides merdae* D1**	INV				GR: Restriction endonuclease subunit S
	INS			*	IR: TonB-dependent receptor
	INV			*	GR: Hypothetical protein
	INS			*	GR: TonB-dependent receptor SusC
	INV			*	IR: Hypothetical protein
	DEL				GR: TonB-dependent receptor SusC
Gemmiger formicilis D2	DEL				IR: Hypothetical protein
	INV				GR: Restriction endonuclease subunit S
	INS				IR: ABC transporter substrate-binding protein YesO
	INS				IR: Type II secretion system protein
	INS		*		IR: tRNA-Ala(tgc)
	DEL		*		IR: Protein VirD4
	INS		*		IR: Hypothetical protein
	INS		*		GR: DUF87 domain-containing protein
Phocaeicola plebeius D2	INS				GR: TonB-dependent receptor P26

SV frequency
0 — 1

* Low/No coverage; IR: Intergenic region; GR: Genic region

The SVs in ✚ strain can be detected in the isolate genomes from fecal of donor and recipient

C

Bacteroides stercoris

SusC SusC

Genotype 1

Genotype 2

D1 → R1 E → R1 L

Alistipes putredinis

Soj ISCaa7

Genotype 1

Genotype 2

D1 → R1 E → R1 L

Gemmiger formicilis

YesO

Genotype 1

Genotype 2

D2 → R3 E → R3 L

■ Genotype 1
■ Genotype 2
E: Early Stage
L: Late Stage

D

Parabacteroides merdae Type I MTases

R M S1 S2 S3

Genotype 1
Genotype 2
Genotype 3
Genotype 4
Genotype 5
Genotype 6

D1 → R1 E → R1 L

■ Genotype 1   ■ Genotype 3   ■ Genotype 5   E: Early Stage
■ Genotype 2   ■ Genotype 4   ■ **Genotype 6**   L: Late Stage

E

Species	Motifs	Donor 1		Post-FMT R1		Methylation detection
		Day 0	Year 5	Week 8	Year 5	
Parabacteroides merdae	TCCGG6mA					■ Yes
	CA6mAGNNNNNNNTTC					□ No
	GRNT6mAYNNNNNNRTTA					
	CA6mAYNNNNNNRTC					← Only in **Genotype 6**

F

		Day 0	Year 5	Week 8	Year 5
Agathobacter faecis	GGW4mCC				
Akkermansia muciniphila	RA6mATTY				
Anaerobutyricum hallii	GGNC5mC				
Anaerostipes hadrus	CTS6mAG				
Bacteroides uniformis	AC6mAGG				
	AGT6mACT				
Mediterraneibacter faecis	GGW4mCC				
Ruminococcus D bicirculans	CTGC6mAG				
Bifidobacterium bifidum	CTGC6mAG				
Dorea longicatena	GC6mATC				
Eubacterium R sp000434995	A4mCGGC				
	GCC6mAT				
	C6mATG				