Macrophages expressing chimeric cytokine receptors have an inflammatory phenotype and anti-tumoral activity upon IL-10 or TGFβ stimulation

Sabrina Traxel[1], Florian Schmidt[1], Corina Beerli[1], Dinh-Van Vuong[1], Roberto F. Speck[1*], Simon Bredl [1*]

[1] Department of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, University of Zurich, Raemistrasse 100, 8091 Zurich, Switzerland

*co-senior authors

CORRESPONDING ADDRESSES:

Dr. rer. nat. Simon Bredl

Department of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, University of Zurich, Raemistrasse 100, 8091 Zurich, Switzerland

e-mail: simon.bredl@gmail.com

tel: +41 43 253 32 78

Dr. sc. nat. Sabrina Traxel

Department of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, University of Zurich, Raemistrasse 100, 8091 Zurich, Switzerland

e-mail: sabrinatraxel@gmail.com

tel: +41 43 253 44 15

DECLARATIONS

Ethics approval and consent to participate

Written consent for the use of buffy coats for research purposes was obtained from blood donors by the Blood Donation Centre.

Consent for publication

Not applicable.

Competing interests

SB and RS hold a patent associated to the ChCR (WO2024133517). No potential competing interests were declared by the other authors.

Funding

This project was funded by Gilead Swiss Fellowship (ST), UZH Postdoc grant (CB), Innosuisse 103.487 IP-LS (SB, RS), Claudia von Schilling Stiftung (SB, RS), Lotte und Adolf Hotz-Sprenger Stiftung (SB, RS), OPO Stiftung (SB, RS), Gebauer Stiftung (SB, RS), Novartis Foundation for Medical Biological Research (#19B139) (SB, RS), USZ Foundation (SB, RS), UZH Entrepreneur Fellowship (SB), Arnold U. und Susanne Huggenberger-Bischoff Stiftung (SB, RS), Peter Bockhoff Stiftung (SB, RS), Mach-Gaensslen Stiftung (SB, RS).

Authors' contributions

Conceptualization: ST, RS, SB

Chimeric cytokine receptor conceptualization and design: SB

Experiments: ST, FS, CB, VDV

Study supervision: RS, SB

Manuscript writing: ST, RS, SB

Funding acquisition: ST, CB, RS, SB

Acknowledgments

We thank the anonymous donors for donating the buffy coats. Flow cytometry was performed with equipment of the Flow Cytometry Facility, University of Zurich. We thank for the assistance in cell sorting experiments by the Flow Cytometry Facility, University of Zurich. We thank the Functional Genomics Center Zurich (FGCZ) of University of Zurich and ETH Zurich, and in particular Timothy Sykes, Hubert Rehrauer*,* and Peter Leary, for the support on RNA sequencing analysis. We thank Jeremy Luban for providing us with pcDNA3.1 Vpx and p8.9NDSB-DPAVDLL plasmids for virus production and Patrick Salmon for providing us with pCWX Dest backbone. We thank Patrick Turko for his advice on analysis of TCGA data. We thank Sarah Brünigk for her advice with the RNAseq data analysis. The results shown here are in part based upon data generated by the TCGA Research Network: https://www.cancer.gov/tcga. ChatGPT was used to improve wording, grammar and spelling of this manuscript.

ABSTRACT

Background

Triple-negative breast cancer (TNBC) is a highly aggressive subtype of breast cancer that lacks hormone receptors and HER2 amplification, making it unresponsive to standard hormone or HER2-targeted therapies. Although immune checkpoint inhibitors (ICIs) have shown promise in tumors with high lymphocyte infiltration, their efficacy remains limited in tumors with minimal lymphocyte infiltration. To overcome this challenge, we developed a novel approach to reprogram tumor-associated macrophages (TAMs) from an immunosuppressive to an inflammatory phenotype within the tumor microenvironment (TME), aiming to enhance therapeutic outcomes in TNBC.

Methods

We designed a chimeric cytokine receptor (ChCR) that triggers STAT1 signaling upon stimulation with IL-10 or TGFβ, cytokines prevalent in the TME that typically drive immunosuppression. Human primary macrophages were transduced with a lentiviral vector expressing the ChCR, and changes in their phenotype, secretome, and transcriptome were analyzed following stimulation. The anti-tumoral activity of these reprogrammed macrophages was assessed using co-culture assays with 3D TNBC spheroids.

Results

ChCR expressing macrophages showed robust STAT1 activation in response to IL-10 or TGFβ stimulation, resulting in an inflammatory phenotype similar to IFNγ activation, as confirmed by phenotypic markers, and transcriptomic profiling. These ChCR-stimulated macrophages demonstrated significant anti-tumoral effects in 3D TNBC spheroids. Moreover, ChCR stimulation led to the upregulation of genes associated with good response to ICIs, among others *CXCL9* and *CXCL10*, chemokines essential for lymphocyte recruitment. Secretion of CXCL10 was also confirmed by ELISA.

Conclusion

We successfully engineered a ChCR that reprograms TAMs within an IL-10- and TGFβ-rich environment, inducing an inflammatory phenotype and anti-tumoral activity. The expression of *CXCL9* and *CXCL10* further supports lymphocyte recruitment, potentially facilitating greater lymphocyte infiltration and disrupting the immunosuppressive TME. This approach offers a promising strategy to improve immunotherapy outcomes in TNBC patients, particularly those with low immune cell infiltration, thereby addressing a critical unmet clinical need.

88 **LIST OF ABBREVIATIONS**

90	CAR	Chimeric antigen receptor
91	ChCR	Chimeric cytokine receptor
92	EF1a	Elongation factor 1 alpha core promoter
93	FDR	False discovery rate
94	GO:BP	Gene Ontology biological process
95	hPGK	Human phosphoglycerate kinase promoter
96	ICI	Immune checkpoint inhibitor
97	MFI	Median fluorescence intensity
98	NES	Normalized enrichment score
99	NR	No response
100	PCA	Principle component analysis
101	R	Response
102	RLU	Relative light units
103	RNAseq	RNA sequencing
104	SEAP	Secreted embryonic alkaline phosphatase
105	SFFV	Spleen focus-forming virus promoter
106	TAM	Tumor associated macrophage
107	TCGA	The Cancer Genome Atlas
108	TME	Tumor microenvironment
109	TNBC	Triple negative breast cancer
110	tNGFR	Truncated nerve growth factor receptor

INTRODUCTION

Triple negative breast cancer (TNBC) is an aggressive subtype of breast cancer that lacks both hormone receptor expression and HER2 amplification and is therefore not amenable to hormone or HER2 targeted therapies (reviewed in[1]). Current first line treatment for both high-risk early-stage TNBC and advanced stage, PD-L1 positive TNBC (combined positive score ≥10) is the immune checkpoint inhibitor (ICI) pembrolizumab in combination with chemotherapy. Early stage TNBC have a good prognosis with 86.6% overall survival at 60 months when treated with neoadjuvant pembrolizumab and chemotherapy.[2] In patients with advanced stage TNBC the pembrolizumab and chemotherapy combination improves the median overall survival from 16.1 months to 23 months, when compared to the placebo and chemotherapy cohort, which is still a dismal prognosis for the patients.[3] Thus, there is clearly an unmet need for patients with advanced TNBC.

Adoptive cell therapy using chimeric antigen receptor (CAR) T cells has achieved remarkable success in hematological malignancies[4] and there was great hope that CAR T cell therapy would also change the therapeutic landscape for solid tumors. To date, CAR T cells had only limited success in solid tumors.[5][6] Several phase I clinical trials of CAR-based immunotherapies for TNBC are currently underway.[7] While we eagerly await the first results of adoptive CAR T cell therapies in TBNC, alternative innovative treatments must be sought.

A critical determinant of disease progression of solid tumors including TNBCs, is their immunophenotype, which is based on the number and localization of CD8+ T cells in the tumor microenvironment (TME).[8-10] Three main immunophenotypes have been defined: hot (high number of intratumoral T cells), excluded (T cells restricted to the tumor margin) or cold (lack of T cells).[11] Patients with a hot TME, including patients suffering from TNBC, have a better prognosis and show better response to ICI treatment than patients with an excluded or cold TEM. [8][10][12-14] In TNBC, 42-54% of patients have a tumor with excluded or cold immunophenotype.[8][10][15] Therefore, reprogramming a cold into a hot TME is of great therapeutic interest.

Myeloid cells including tumor associated macrophages (TAMs) constitute up to 30% of the tumor mass.[16] TAMs differentiate from monocytes that are actively recruited into the TME.[17] In general, macrophages can adopt a wide range of phenotypes depending on the stimuli received and tissue environment.[18] Consequently, their activities include immunosuppression, tissue repair, phagocytosis, inflammation, and cytotoxicity. Inflammatory macrophages are induced by pro-inflammatory signals like IFNγ and/or LPS, whereas immunosuppressive macrophages are induced by immunosuppressive cytokines like IL-4, IL-13, IL-10, or TGFβ. IL-10 and TGFβ are expressed in the TME of solid tumors, such as in TNBC[19-21] and consequently push TAMs to adopt an immunosuppressive phenotype. TAMs inhibit the immune system, support tumor growth, angiogenesis, and tumor cell migration (reviewed in[22]). In contrast, inflammatory macrophages have anti-tumoral activity by attracting T cells, by increased antigen presentation and their direct tumoricidal effects.[20][23-26] In fact, tumors with high number of inflammatory macrophages are more likely to have a hot TME, a better prognosis and better response to ICIs.[8][27-29] Thus, reprogramming TAMs towards an inflammatory phenotype is a hot topic in the field of cancer immunotherapy.[30]

We believe that macrophages should be reprogrammed to switch to inflammatory and anti-tumoral macrophages locally in the TME. Systemically activated macrophages may be retained in various tissues, unable to reach their destination. Furthermore, local activation would be less likely to have off-target effects. Therefore, we developed chimeric cytokine receptor (ChCR) expressing macrophages that utilize immunosuppressive cytokines enriched in the TME to switch to

148   inflammatory macrophages. Specifically, we engineered macrophages to express a ChCR that binds either IL-10 or TGFβ

149   and induces an IFNγ signaling.

MATERIALS AND METHODS

Ethics statement

Monocytes used were obtained from buffy coats, anonymously provided by the Blood Donation Service Zurich, Swiss Red Cross, Schlieren. Written consent for the use of buffy coats for research purposes was obtained from blood donors by the Blood Donation Centre.

Cell lines

HEK293T (ATCC) were maintained in Dulbecco′s Modified Eagle′s Medium - high glucose (DMEM, Sigma-Aldrich, D6429) supplemented with 10% heat-inactivated fetal bovine serum (FBS, Gibco, 10270-106) (hereafter called D10). HEK-Blue[TM] IFN-γ cells (Invivogen, hkb-infg)) were maintained in D10 supplemented with 100μg/ml Zeocin (Invivogen, ant-zn-1) and 30μg/ml Blasticidin (Invivogen, ant-bl-1). MDA-MB-231 (ATCC) were maintained in RPMI-1640 (Sigma-Aldrich, R8758) supplemented with 10% heat-inactivated FBS (Gibco, 10270-106) (hereafter called R10). BT-549 (ATCC) were maintained in R10 supplemented with 10ug/ml insulin (Sigma- Aldrich, I9278). THP-1 (ATCC) were maintained in R10 supplemented with 2-mercaptoethanol (Gibco, 31350010). MRC-5 (ATCC-CCL-171) were maintained in Eagle's Minimum Essential Medium (EMEM, ATCC-30-2003) supplemented with 10% heat inactivated FBS. All cells were maintained at 5% CO_2 and 37°C in a humidified incubator.

Monocyte derived macrophages

Peripheral mononuclear cells were isolated from buffy coats of healthy donors (Blood Donation Service Zurich, Swiss Red Cross, Schlieren) using density gradient centrifugation on lymphoprep (Axon Lab, 1858). Monocytes were isolated using CD14 microbeads (Miltenyi Biotec, 130-050-201) and plated at 150'000 cells/cm^2 in F+ high attachment plates (Sarstedt) in RPMI-1640 supplemented with Penicillin-Streptomycin (P/S, Gibco, 15140-122). Two hours after plating, medium was replaced with RPMI-1640 supplemented with P/S and 10% human AB serum (Sigma Aldrich, H6914), hereafter called RH10. Monocytes were differentiated for 6-11 days with medium change every 2-3 days. To polarize macrophages, they were stimulated for additional 48h with IFNγ (Biolegend, 570204), IL-10 (Biolegend, 571004) or TGFβ1 (Peprotech, 100-21) at indicated concentrations.

Plasmid construction

To generate ChCR plasmids, amino acid sequences of the human IL10RA, IL10RB, TGFβR1, TGFβR2, IFNGR1 and IFNGR2 receptor were obtained from UniProt. Supplementary Table 1 shows Uniprot sequences used for the ChCR design. The ChCR were constructed as follows: The extracellular domains of IL10RA, IL10RB, TGFβR1, or TGFβR2 were fused to the transmembrane and extracellular domain of the IFNGR1 or IFNGR2, including a glycine-serine linker for flexibility.[31][32] As an N-terminal signal peptide, we used the one from the corresponding intracellular IFNGR. All variants were analyzed with DeepTMHMM to verify *in silico* signal peptide cleavage, transmembrane domains and surface expression.[33] Sequences were ordered codon-optimized at GenScript Biotech Corporation. For expression of single chains, ChCR sequences were subcloned into a pCWX Dest lentiviral vector (Gift from Patrick Salmon) under the control of an elongation factor 1 alpha core promoter (EF1a). To simultaneously express both chains, ChCR sequences were subcloned into a pCWX Dest lentiviral vector separated by a P2A sequence and under the control of a spleen focus-forming virus (SFFV) promoter. All lentiviral vectors also encode for a truncated nerve growth factor receptor (tNGFR) driven by a

187   human phosphoglycerate kinase (hPGK) promoter as a reporter for transduction. Mock lentiviral vector only encodes for

188   tNGFR.

189   To generate the Luciferase-EGFP plasmid, luciferase and EGFP sequences, separated by a T2A sequence, were ordered as

190   gene blocks and subcloned into a pCWX Dest lentiviral vector under the control of EF1a promoter.

Lentivirus production and titration

Lentiviral vectors were produced in HEK293T. Briefly, 13 Million cells were plated in T150 flasks. The day after, cells

were co-transfected using PEI (Polysciences, 23966) with pMD2.G, p8.9NDSB-DPAVDLL[34] (gift from Jeremy Luban),

pcDNA3.1 Vpx239[34] (gift from Jeremy Luban) and pCWX Dest transfer vector at a molar ratio of 251:107.5:756.1:45. 4h

after transfection, medium was changed and supplemented with 5mM sodium butyrate (Sigma Aldrich, B5887). Virus was

harvested 52h after transfection and clarified by centrifugation. Lentiviral supernatants were concentrated by ultrafiltration

using 100kDa cut-off Amicon filters (Merck Milipore, UFC910024). Titration of lentiviruses was done by transducing

HEK293T. Three days after transduction, transduced cells were stained for tNGFR and analysed by flow cytometry on a

LSR II Fortessa (BD Biosciences).

THP-1 transduction

201   THP-1 were transduced with lentiviruses at MOI 1 by spinoculation (800g, 45min, 32°C) in presence of protamine sulfate

202   (100ug/ml, Sigma-Aldrich, P402). Transduced cells were sorted based on tNGFR+ expression using MACSelect LNGFR

203   microbeads (Miltenyi Biotec, 130-091-330).

Monocyte transduction

205   To transduce monocytes, lentivirus was added 2h after plating of monocytes in RPMI + P/S + 5% human AB serum at a

206   MOI of 10 or 15 in presence of protamine sulfate (100ug/ml, Sigma-Aldrich, P4020). After overnight incubation, medium

207   was changed to RH10. Transduction efficiency was assessed between day 6 and day 12 of differentiation and was defined

208   as %tNGFR+ cells. In general, macrophages with transduction rates below 65% were excluded from experiments. In early

209   experiments assessing IL-10 ChCR macrophage polarization, transduction rates down to 55% were accepted.

IFNγ signaling reporter cells

211   HEK-Blue™ IFN-γ were transduced with lentiviruses by spinoculation in the presence of 8ug/ml polybrene (SantaCruz,

212   sc-134220). To measure secreted embryonic alkaline phosphatase (SEAP), transduced cells were detached mechanically

213   and seeded in 96-well plate. Cells were incubated with cytokines as indicated for 24h. Thereafter, 30ul of the cell

214   supernatant was incubated with 170ul Quanti-Blue solution (Invivogen, rep-qbs) and incubated for 1h at 37°C. OD was

215   measured at 620nm using Spectramax iD3 plate reader (Molecular Devices).

GFP-Luciferase expressing tumor cells

217   MDA-MB-231 cell line was transduced with lentiviral construct to express Luciferase and EGFP. Cells were sorted for

218   high EGFP expression on a BD FACSAria III (BD Biosciences).

Flow cytometry analysis

To assess transduction rate and polarization marker expression, macrophages were harvested for flow cytometry by

incubation with 5mM EDTA in PBS on ice for 45min, followed by gentle pipetting. Detached macrophages were first

treated with True-Stain FcX (Biolegend, 422301) and thereafter stained with antibodies (Table 1). Stained samples were

acquired on LSR II Fortessa (BD Life Sciences). Untransduced or mock transduced cells were used as gating control for

tNGFR and IL-10Rα/IL-10Rβ/TGFβR2, respectively. Data was analyzed using FlowJo v10.10 software (BD Life

Sciences). Median fluorescence intensity (MFI) of polarization marker was normalized to the MFI of the corresponding

isotype by subtraction. Corresponding isotype controls were used to gate for CD38+ cells.

Table 1: Flow cytometry antibodies

Epitope	Fluorochrome	Company	Category Number
anti-mouse	BV421	Biolegend	405317
anti-rat	BV421	Biolegend	405414
CD11b m/h	FITC	Biolegend	101206
CD163	BV711	Biolegend	333630
CD206	PE	BioLegend	321105
CD38	BV605	BioLegend	303532
CD86	BV785	Biolegend	305441
HLA-DR	BV510	BioLegend	307646
IgG1 Isotype	BV711	Biolegend	400167
IgG1 Isotype	PE	BD Pharmingen	555749
IgG1 Isotype	BV605	Biolegend	400161
IgG2b Isotype	BV785	Biolegend	400355
IgG2a Isotype	BV510	Biolegend	400267
IL-10Rα	unconjugated	Biolegend	308802
IL-10Rα	PE	Miltenyi Biotec	130-121-341
IL-10Rβ	unconjugated	R&D Systems	MAB874
IL-10Rβ	APC	R&D Systems	FAB874A
NGFR (CD271)	Alexa Fluor 647	BD Biosciences	560326
NGFR (CD271)	BV510	BD Biosciences	563451
NGFR (CD271)	APC	BioLegend	345108
pSTAT1 (Tyr701)	PE	Biolegend	666403
TGFβR2	PE	Biolegend	399703

STAT1 phosphoflow

THP-1 or human primary macrophages were used for STAT1 phosphoflow experiments. THP-1 were plated in round

bottom 96-well plates in R10. Macrophages were detached as described for flow cytometry analysis and plated in round

bottom ultra low attachment 96-well plates (Corning, 7007) in RH10. Cells were rested for 1h at 37°C and thereafter

stimulated for 30min as indicated. Cells were fixed by adding one volume of 4% PFA (Santa Cruz) and 15min incubation

at 37°C. Next, cell surface markers were stained before cells were permeabilized with True-Phos Perm Buffer (Biolegend).

Cells were stained for intracellular pSTAT1 (Tyr701) and analyzed on LSR II Fortessa (BD Biosciences).

IL10 and TGFB1 expression in TNBC

Gene expression data from 195 TNBC patients were obtained from The Cancer Genome Atlas (TCGA). TNBC patients in

the TCGA dataset were previously identified by Lehmann et al.[35] Raw counts were retrieved, preprocessed and normalized

using the R package TCGABiolinks (2.33.0) without filtration.[36] Percentile expression of IL10 and TGFB1 across all genes

was calculated by dividing the rank expression with the total analyzed genes.

RNA sequencing and analysis

For bulk RNA sequencing (RNAseq), monocytes were transduced and differentiated into macrophages. Macrophages were stimulated for 48h as indicated. Total RNA was extracted using Quiashredder (Qiagen, 79254) and RNeasy Mini Kit (Qiagen, 74104) with on column DNA digestion (Qiagen, 79254). RNAseq was performed at the Functional Genomics Center Zurich (FGCZ). The NovaSeq X Plus (Illumina, Inc, California, USA) was used for cluster generation and sequencing according to standard protocol. Sequencing was paired end 150 bp. Between 28 and 86 million reads were generated per sample. RNAseq analysis was performed using the SUSHI framework,[37] which encompassed the following steps: Read quality was inspected using FastQC,[38] and sequencing adaptors removed using fastp,[39] alignment of the RNA-Seq reads using the STAR aligner,[40] and with the GENCODE human genome build GRCh38.p13 (Release 42) as the reference[41] ; the counting of gene-level expression values using the featureCounts function of the R package Rsubread.[42] Differential expression analysis was performed using the generalized linear model as implemented by the DESeq2 Bioconductor R package.[43] Principle component analysis (PCA) was performed using the prcomp function from stats R package. For gene set enrichment analysis (GSEA), differentially expressed genes were ranked by log2 fold change. Gene Ontology (GO) GSEA tests were performed using the gseGO function respectively of the clusterProfiler Bioconductor R package.[44] Hallmark and custom pathways GSEA was performed using the GSEA function of the Bioconductor R package clusterProfiler (4.13.0)[44] and the hallmark gene sets (MSigDb 2023.2)[45] on R version (4.4.0). To assess the GSEA of genes upregulated or downregulated in macrophages in anti-PDL-1 responders, genes upregulated or downregulated in baseline tumor macrophages of responders[27] were assembled as custom pathways referred to as responder and non-responder, respectively. Unless stated otherwise, all R functions were executed on R version 4.3.2 (R Core Team, 2020) and Bioconductor version 3.17.

CXCL10 ELISA

To analyze CXCL10 secretion, differentiated macrophages were stimulated for 48h as indicated. Thereafter, supernatant was collected and cell debris removed by centrifugation. Supernatants were stored at -80°C. CXCL10 in supernatant was measured using the ELISA MAX™ Deluxe Set Human CXCL10 (IP-10) kit (Biolegend, 439904). OD450nm and OD570 as background were measured using Multiskan SkyHigh plate reader (Thermo Scientific™).

3D spheroid assay

Heterotypic spheroids were generated by mixing 1000 MDA-MB-231 and 1500 MRC-5 cells in Akura™ 96 Spheroid Microplates (InSphero, CS-PB15) in R10. Plates were centrifuged at 100g for 5 min to remove air bubbles and incubated at 37°C, 5% CO_2 overnight. To support the aggregation and maturation of the spheroids, the plates were placed on a tilting stand at a 30° angle. The following day, macrophages were detached using 5mM EDTA in PBS and gentle pipetting and added at a 10:1 ratio (macrophages:MDA-MB-231 at seeding) and stimulated as indicated. Medium and stimuli were replaced every 2-3 days. Fluorescent (EGFP) images were captured using the Cytation 3 (BioTek) with a 4x objective. Images were processed and analyzed with the Gen5 (BioTek) software. To determine the cell number, the mean fluorescence intensity of the full image was determined and the background of an empty well subtracted.

Luciferase assay

To assess an endpoint quantification of tumor cells in the spheroids, luciferase activity of the remaining MDA-MB-231 was determined. Spheroids were washed twice with PBS to get rid of luciferase in the supernatant and left with 50 ul

residual PBS per well. 50 ul of BrightGloTM substrate (Promega, E2610) was added per well, consequently leading to cell

lysis and release of contained luciferase. The lysate was transferred into a 96-well white plate (Pierce™, 15042). After a

minimum waiting time of 2 min, luciferase activity was determined using GloMax® Explorer luminometer (Promega,

GM3500). The background signal was determined using a medium control and was subtracted from the samples.

WST1 assay

To assess the ability of genetically engineered macrophages to affect TNBC proliferation, macrophages were stimulated

for 2d with IFNγ, IL-10 or TGFβ at indicated concentrations. Supernatants were harvested, clarified by centrifugation and

stored at -80°C. For WST1 assay, 5'000 BT-549 or 10'000 MDA-MB-231 cells were plated in 96-well plates. The next

day, medium was replaced with supernatants from stimulated macrophages or RH10 as untreated control. 72h after addition

of supernatants, viability was measured by incubating cells for 1h with cell proliferation reagent WST-1 (Roche,

5015944001). OD440nm and OD650nm as background were measured using Multiskan SkyHigh plate reader (Thermo

Scientific™).

Statistical analysis

The statistical analysis of TCGA and previously published single cell RNAseq data was performed in R (4.4.0). The

statistical analysis of bulk RNAseq data was performed using the Sushi framework of the FGCZ.[37] The remaining statistical

analysis was performed using Graphpad Prism 10.2.0 using statistical tests specified in the figure legends.

RESULTS

IL10 and *TGFB1* are expressed in TNBC

IL-10 and TGFβ1 have previously been shown to be expressed in TNBC patients.[19-21] To confirm this data, we analyzed publicly available RNAseq data from 195 TNBC patients for *IL10* and *TGFB1* expression. We found that *IL10* and *TGFB1* were expressed among the top 73% (median) and top 90% (median) genes (figure 1). Thus, TCGA gene expression analysis confirms previously published results of IL10 and TGFβ expression in TNBC patients.

IL-10 and TGFβ ChCR induce STAT1 signaling

To convert IL-10 or TGFβ signals into IFNγ signals, we designed two ChCR, which consist of the extracellular domains of IL10 receptor or TGFβ receptor, respectively, fused to the intracellular domains of the IFNγ receptor (figure 2A and B). To control for transduction efficacy, we also incorporated the truncated NGFR (tNGFR) in the lentiviral vector.

In a first instance, we verified IFNγ signaling through the IL-10 ChCR in HEK-Blue IFNγ reporter cell line. In the HEK-Blue IFNγ reporter cell line, IFNγ-induced STAT1 signaling leads to expression of secreted embryonic alkaline phosphatase (SEAP), which can be measured in the supernatant using a colorimetric assay. We stably expressed either the full IL-10 ChCR or the individual components, i.e. IL-10RA fused to IFNGR1 and IL-10RB fused to IFNGR2 in the HEK-Blue IFNγ reporter cells. (online supplemental figure S1A). All cell lines were transduced at >90% efficiency (data not shown) and ChCR expression was confirmed by flow cytometry (online supplemental figure S1B). We found that IL-10 induced IFNγ signaling in a dose-dependent manner when the full IL-10 ChCR was expressed (online supplemental figure S1C). Surprisingly, IL-10RA-IFNGR1 alone triggered moderate IFNγ signaling, but both chains of the ChCR were necessary to achieve maximum STAT1 signaling. We also verified that the expression of the full TGFβ ChCR in HEK-Blue IFNγ reporter cell line induced STAT1 signaling (data not shown). Next, we proceeded to the characterization of both ChCRs in the monocyte cell line THP-1, which more closely approximates the properties of primary human macrophages. We found a prominent upregulation of IL-10Rα and TGFβR2 and a moderate one of IL10Rβ using flow cytometry (figure 2C). We tested several TGFβR1 antibodies, all suffered from non-specificity and therefore we do not have expression data for TGFβR1. Since TGFβR1 is on the first position of the polycistronic vector and TGFβR2 expression is detectable, we assume that TGFβR1 is expressed as well.

Importantly, IL-10 and TGFβ induced a dose-dependent activation of STAT1 in IL-10 or TGFβ ChCR expressing THP-1, respectively, (figure 2D). As expected, untransduced and mock transduced THP-1 cells only showed pSTAT1 signaling upon IFNγ stimulation.

To investigate the ChCRs in primary macrophages, we first optimized the protocol for transducing monocytes. We achieved a transduction efficiency of 70.2 ± 14.5% (mean±SD) across all vectors (figure 2E). Again, we saw a very prominent upregulation of IL-10Rα and TGFβR2 and a lesser one of IL-10Rβ (figure 2F). IL-10 induced STAT1 activation to a similar level like IFNγ, whereas TGFβ induced only a little, but consistent STAT1 activation in the macrophages expressing the ChCRs (figure 2G). Thus, both IL-10 and TGFβ ChCR variants trigger the IFNγ signaling pathway.

ChCR expression induces an inflammatory phenotype upon IL-10 / TGFβ stimulation

We next assessed whether triggering the ChCRs induces an inflammatory phenotype in primary human macrophages. We focused on the expression of polarization markers typically upregulated on macrophages either upon IFNγ (i.e., HLA-DR,

CD38, CD86) or IL-10 stimulation (i.e., CD163). Importantly, both, IL-10 and TGFβ induced the expression of HLA-DR,

CD38 and CD86 in the macrophages expressing the corresponding ChCRs (figure 3A and B). Moreover, IL-10 blocked

the upregulation of CD163 in IL-10 ChCR transduced macrophages, while IL-10 massively increased its expression in the

controls (figure 3A). Notably, CD163 is upregulated in response to IL-10 but not to TGFβ and therefore we did not measure

CD163 expression in TGFβ ChCR expressing macrophages. Overall, we found that, ChCR expressing macrophages indeed

show a phenotype similar to IFNγ stimulated macrophages upon stimulation with IL10 or TGFβ.

ChCR expressing macrophages have a IFNγ-like transcriptome

Next, we performed RNAseq to extend the characterization of ChCR expressing macrophages. Principle component

analysis (PCA) showed that ChCR expressing macrophages stimulated with their corresponding ligand clustered together

with IFNγ stimulated-macrophages, indicating a IFNγ-like transcriptome profile induced by stimulation of the ChCRs

(figure 4A). As expected, differential gene expression analysis of ChCR expressing macrophages showed the upregulation

of many IFNγ stimulated genes upon IL-10 or TGFβ stimulation when compared to mock transduced macrophages

stimulated equally (figure 4B). Similarly, gene set enrichment analysis (GSEA) of hallmark gene sets showed that IFNγ

stimulated genes, among other *CXCL9/CXCL10/CIITA,* were significantly enriched in ChCR expressing macrophages

upon stimulation (figure 4C). Moreover, GSEA of gene ontology biological processes (GO:BP) showed enrichment of

genes associated with antigen presentation (MHC-II HLA genes) and several immune related pathways (figure 4D). Thus,

we could confirm that ChCR expressing macrophages have an IFNγ-like transcriptome.

ChCR expressing macrophage upregulate genes associated with anti-PD-L1 response in TNBC patients

CXCL9 and CXCL10 have been associated with better response to immunotherapy and a hot TME in TNBC and other

tumors.[8 27 28 46] Interestingly, both *CXCL9* and *CXCL10* were upregulated in ChCR expressing macrophages in RNAseq,

which we confirmed for CXCL10 by ELISA (figure 5A and B). Zhang et al have performed single cell RNAseq of TNBC

patients treated with chemotherapy and anti-PD-L1 and have identified a TAM transcriptome profile associated with anti-

PD-L1 response.[27] Thus, we assessed the expression of other genes associated with good response to ICI treatment in our

ChCR expressing macrophages upon stimulation. Indeed, 50 (IL-10 ChCR) and 58 (TGFβ ChCR) of 195 response

associated genes were also significantly upregulated in ChCR expressing macrophages upon stimulation with the respective

ligand in comparison to mock transduced macrophages (data not shown). Moreover, the heatmap analyzing response

associated genes shows a cluster of genes also upregulated in ChCR expressing macrophages (figure 5C). Genes also

upregulated in ChCR expressing macrophages upon stimulation belong i.e. to hallmark pathways of IFNγ response and

allograft rejection (i.e. *CD74, CXCL9, STAT1, SERPING1, B2M…*) and TNFα signaling via NF-κB (i.e. *TNF, NFKBIA,*

CCL4, ICAM1…). In addition, we have performed gene set enrichment analysis of responder gene set (R, upregulated in

anti-PD-L1 responders) and non-responder gene set (NR, downregulated in anti-PD-L1 responders). We found that the

responder gene set was indeed enriched in ChCR expressing macrophages upon stimulation with the respective ligand

(figure 5D). In contrast, the gene set associated with non-response was not enriched in ChCR expressing macrophages.

Thus, the transcriptome of our genetically engineered macrophages could be beneficial for the response to anti-PD-L1

therapy.

ChCR expressing macrophages have anti-tumoral activity

Inflammatory macrophages are reported to have anti-tumoral effects.[20 23-25] Therefore, we explored whether ChCR

engineered macrophages have an anti-tumoral effect in a heterotypic TNBC spheroid model. This spheroid model consisted

of the TNBC cell line MDA-MB-231 and MRC5 fibroblasts. The presence of fibroblasts was essential as MDA-MB-231

cells alone did not form spheroids. One day after formation of MDA-MB-231 and MRC-5 spheroids, we added the

genetically engineered macrophages. To monitor the viability of MDA-MB-231 cells easily, we engineered the cell line to

express both EGFP and luciferase. We used EGFP to measure MDA-MB-231 number by microscopy over time and

luciferase activity to determine the end-point cell number. Indeed, the IL-10- or TGFβ ChCR expressing macrophages

limited the expansion of MDA-MB-231 spheroid when added together with the corresponding cytokines, as assessed by

mean EGFP intensity (figure 6A-D). We observed similar results when measuring the remaining luciferase activity on day

6, even though there was only a trend for TGFβ ChCR (figure 6E and F). Next, we were wondering whether the anti-

tumoral effect of ChCR expressing macrophages was mediated by soluble factors or cell-contact. We found that supernatant

from stimulated ChCR expressing macrophages alone reduced the proliferation of MDA-MB-231 and BT-549 cells (online

supplemental figure S2), indicating secreted factors, are, at least in part, responsible for the anti-tumoral effect observed in

the spheroid model.

DISCUSSION

The cold TME is highly suppressive and makes tumors non-responsive to ICI. The presence of inflammatory macrophages
in the TME increases the responsiveness to ICIs [8 27 28] and therefore, TAMs or macrophages recruited to the TME represent
a therapeutic target. Here, we developed ChCR expressing macrophages that are designed to switch to inflammatory
macrophages in the presence of IL-10 and TGFβ in the TME. We show that ChCR expressing macrophages indeed have a
phenotype similar to IFNγ-induced inflammatory macrophages. Importantly, ChCR expressing macrophages also have a
direct anti-tumoral activity.

In some ways, the TME is a standalone area for the best tumor growth. Immunosuppressive cytokines are prevalent, among
other IL-10 and TGFβ. In TCGA RNAseq data, we confirmed that the transcripts of these cytokines are present in TNBC.
Therefore, we constructed ChCRs by combining the extracellular domains of the IL-10 and TGFβ receptors with the
intracellular signaling domains of the IFNγ receptor to activate macrophages in the TME, leveraging the local enrichment
of IL-10 and TGFβ. Please, note that IFNγ is the crucial cytokine for inducing the inflammatory activation of macrophages.
To provide flexibility to the ChCRs, we added a glycine-serine linker between the extracellular domains and the
transmembrane domains of the ChCRs.[31 32] To ensure targeting the ChCR to the cell surface, we used the signal peptides
from the corresponding IFNγR subunit. The characterization in terms of surface expression and signaling clearly showed
that we have created functional ChCRs.

Our results suggest that ChCR expressing macrophages become inflammatory in response to IL-10 or TGFβ. These ChCR
expressing macrophages exhibit several attributes that would allow them to disrupt an immunosuppressive TME. Firstly,
RNAseq showed that ChCR expressing macrophages upregulated the expression of the chemokines *CXCL9* and *CXCL10*
upon IL-10 or TGFβ stimulation, which we confirmed for CXCL10 by ELISA .Both in TNBC and beyond, there are several
studies that underline the importance of CXCL9 and CXCL10 expression in the TME for the efficacy of a checkpoint
inhibitor treatment.[8 27 28 46-49] It is noteworthy that Sloas et al. recently reported that macrophages engineered with an IL-
10/IFNλ switch receptor, similar to our IL-10 ChCR, led to an increased T cell infiltration into the tumor *in vivo* in a
syngeneic tumor mouse model.[50 51] Secondly, HLA-DR and CD86 were also upregulated as determined by flow cytometry.
These surface proteins are crucial for T-cell activation and antigen presentation. RNAseq analysis also showed enrichment
of pathways associated with antigen presentation. Therefore, we believe that ChCR expressing macrophages could lead to
attraction, infiltration and activation of T cells and thereby initiating a positive feedback loop that results in T cells secreting
IFNγ, which in turn leads to TAMs adopting an inflammatory phenotype.[29] Ultimately, we show that that ChCR expressing
macrophages have a direct anti-tumoral effect when adding to a TNBC spheroid composed of MDA-MB-231 cells and
MRC5 fibroblasts. The anti-tumoral effect is, at least partially, mediated by secreted factors, as conditioned medium
obtained from ChCR expressing macrophages reduced the viability of the TNBC cells MDA-MB-231 and BT-549. It is
well known that macrophages exhibit anti-tumoral activity upon IFNγ and LPS stimulation.[24-26 52] Here we show that IFNγ-
like signaling alone via the ChCR render macrophages anti-tumorally active.

We noticed that spheroids co-cultured with IL-10 or TGFβ ChCR expressing macrophages showed a reduced spheroid
dissemination compared to the controls (data not shown). It is highly speculative but this reduced dissemination may be
due to the anti-tumoral activity of the ChCR expressing macrophages, while TAMs commonly promote metastasis [53]

We observed differences in the performance of the two ChCR variants evaluated: the TGFβ ChCR responded to a lower
dose of TGFβ than the IL-10 ChCR to IL-10 as shown by the STAT1 activation in THP-1 and CXCL10 ELISA in primary

macrophages. These differences could be due to distinct affinities to the corresponding cytokine or distinct activation

thresholds of the ChCRs. The difference in activation threshold might be dependent on the combination of extracellular

domain with the intracellular domain of the IFNGR2, which is the limiting subunit in inducing signaling.[54] For instance,

in the case of the IL-10 ChCR, we fused the low affinity IL-10Rb to IFNGR2 (reviewed in[55]), whereas in the case of TGFβ

ChCR, we fused the high affinity TGFβR2 to the IFNGR2.[56] Importantly, in the complex setting of spheroids, macrophages

expressing IL-10 ChCR performed similar to macrophages expressing the TGFβ ChCR, when saturating concentrations

were used (figure 6). Ultimately, *in vivo* experiments will be needed to assess and compare activity and toxicity of both

ChCR variants.

As mentioned above, patients with advanced, PD-L1 positive TNBC benefit from adding pembrolizumab to

chemotherapy.[3] TAMs isolated from patients who responded to anti-PD-L1 treatment show a very unique transcriptomic

pattern.[27] The RNAseq data we obtained from ChCR expressing macrophages in response to the corresponding cytokines,

showed upregulation of key genes associated with better response to PD-L1 ICI. Therefore, we have good reasons to

assume that PD1 ICI treatment in concert with adoptively transferred ChCR expressing macrophages could result in an

additive or synergistic anti-tumoral activity.

What is the weakness of the data? The ChCR expressing macrophages present several attributes which may render them

anti-tumoral active. However, all the assays including the spheroid assays do not recapitulate the complexity of a tumor

with its rich cell content, low pH, high levels of adenosine and hypoxia. Spheroid assays at the best try to mimic it. We

will push forward and investigate the anti-tumor activity of these ChCR expressing macrophages in vivo in orthotopic

TNBCs mouse models.

LITERATURE

1. Foulkes WD, Smith IE, Reis-Filho JS. Triple-negative breast cancer. *N Engl J Med* 2010;363(20):1938-48. doi: 10.1056/NEJMra1001389

2. Schmid P, Cortes J, Dent R, et al. Overall Survival with Pembrolizumab in Early-Stage Triple-Negative Breast Cancer. *N Engl J Med* 2024 doi: 10.1056/NEJMoa2409932 [published Online First: 20240915]

3. Cortes J, Rugo HS, Cescon DW, et al. Pembrolizumab plus Chemotherapy in Advanced Triple-Negative Breast Cancer. *N Engl J Med* 2022;387(3):217-26. doi: 10.1056/NEJMoa2202809

4. Cappell KM, Kochenderfer JN. Long-term outcomes following CAR T cell therapy: what we know so far. *Nat Rev Clin Oncol* 2023;20(6):359-71. doi: 10.1038/s41571-023-00754-1 [published Online First: 20230413]

5. Albelda SM. CAR T cell therapy for patients with solid tumours: key lessons to learn and unlearn. *Nat Rev Clin Oncol* 2024;21(1):47-66. doi: 10.1038/s41571-023-00832-4 [published Online First: 20231030]

6. Newick K, O'Brien S, Moon E, Albelda SM. CAR T Cell Therapy for Solid Tumors. *Annu Rev Med* 2017;68:139-52. doi: 10.1146/annurev-med-062315-120245 [published Online First: 20161117]

7. Geng P, Chi Y, Yuan Y, et al. Novel chimeric antigen receptor T cell-based immunotherapy: a perspective for triple-negative breast cancer. *Front Cell Dev Biol* 2023;11:1158539. doi: 10.3389/fcell.2023.1158539 [published Online First: 20230629]

8. Hammerl D, Martens JWM, Timmermans M, et al. Spatial immunophenotypes predict response to anti-PD1 treatment and capture distinct paths of T cell evasion in triple negative breast cancer. *Nat Commun* 2021;12(1):5668. doi: 10.1038/s41467-021-25962-0 [published Online First: 2021/09/29]

9. Li F, Li C, Cai X, et al. The association between CD8+ tumor-infiltrating lymphocytes and the clinical outcome of cancer immunotherapy: A systematic review and meta-analysis. *EClinicalMedicine* 2021;41:101134. doi: 10.1016/j.eclinm.2021.101134 [published Online First: 20210916]

10. Gruosso T, Gigoux M, Manem VSK, et al. Spatially distinct tumor immune microenvironments stratify triple-negative breast cancers. *J Clin Invest* 2019;129(4):1785-800. doi: 10.1172/JCI96313 [published Online First: 2019/02/13]

11. Galon J, Bruni D. Approaches to treat immune hot, altered and cold tumours with combination immunotherapies. *Nat Rev Drug Discov* 2019;18(3):197-218. doi: 10.1038/s41573-018-0007-y

12. de Jong VMT, Wang Y, Ter Hoeve ND, et al. Prognostic Value of Stromal Tumor-Infiltrating Lymphocytes in Young, Node-Negative, Triple-Negative Breast Cancer Patients Who Did Not Receive (neo)Adjuvant Systemic Therapy. *J Clin Oncol* 2022;40(21):2361-74. doi: 10.1200/JCO.21.01536 [published Online First: 20220330]

13. Park JH, Jonas SF, Bataillon G, et al. Prognostic value of tumor-infiltrating lymphocytes in patients with early-stage triple-negative breast cancers (TNBC) who did not receive adjuvant chemotherapy. *Ann Oncol* 2019;30(12):1941-49. doi: 10.1093/annonc/mdz395

14. Nederlof I, Isaeva OI, de Graaf M, et al. Neoadjuvant nivolumab or nivolumab plus ipilimumab in early-stage triple-negative breast cancer: a phase 2 adaptive trial. *Nat Med* 2024 doi: 10.1038/s41591-024-03249-3 [published Online First: 20240916]

15. Keren L, Bosse M, Marquez D, et al. A Structured Tumor-Immune Microenvironment in Triple Negative Breast Cancer Revealed by Multiplexed Ion Beam Imaging. *Cell* 2018;174(6):1373-87 e19. doi: 10.1016/j.cell.2018.08.039 [published Online First: 2018/09/08]

16. Combes AJ, Samad B, Tsui J, et al. Discovering dominant tumor immune archetypes in a pan-cancer census. *Cell* 2022;185(1):184-203 e19. doi: 10.1016/j.cell.2021.12.004 [published Online First: 20211227]

17. Franklin RA, Liao W, Sarkar A, et al. The cellular and molecular origin of tumor-associated macrophages. *Science* 2014;344(6186):921-5. doi: 10.1126/science.1252510 [published Online First: 20140508]

18. Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev Immunol* 2008;8(12):958-69. doi: 10.1038/nri2448

19. Toney NJ, Opdenaker LM, Cicek K, et al. Tumor-B-cell interactions promote isotype switching to an immunosuppressive IgG4 antibody response through upregulation of IL-10 in triple negative breast cancers. *J Transl Med* 2022;20(1):112. doi: 10.1186/s12967-022-03319-5 [published Online First: 20220307]

20. Chen Z, Wu J, Wang L, et al. Tumor-associated macrophages of the M1/M2 phenotype are involved in the regulation of malignant biological behavior of breast cancer cells through the EMT pathway. *Med Oncol* 2022;39(5):83. doi: 10.1007/s12032-022-01670-7 [published Online First: 20220516]

21. Ding MJ, Su KE, Cui GZ, et al. Association between transforming growth factor-beta1 expression and the clinical features of triple negative breast cancer. *Oncol Lett* 2016;11(6):4040-44. doi: 10.3892/ol.2016.4497 [published Online First: 20160426]

22. Qian BZ, Pollard JW. Macrophage diversity enhances tumor progression and metastasis. *Cell* 2010;141(1):39-51. doi: 10.1016/j.cell.2010.03.014

23. Engstrom A, Erlandsson A, Delbro D, Wijkander J. Conditioned media from macrophages of M1, but not M2 phenotype, inhibit the proliferation of the colon cancer cell lines HT-29 and CACO-2. *Int J Oncol* 2014;44(2):385-92. doi: 10.3892/ijo.2013.2203 [published Online First: 2013/12/04]

24. Pahl JH, Kwappenberg KM, Varypataki EM, et al. Macrophages inhibit human osteosarcoma cell growth after activation with the bacterial cell wall derivative liposomal muramyl tripeptide in combination with interferon-γ. *Journal of Experimental & Clinical Cancer Research* 2014;33

25. Georgouli M, Herraiz C, Crosas-Molist E, et al. Regional Activation of Myosin II in Cancer Cells Drives Tumor Progression via a Secretory Cross-Talk with the Immune Microenvironment. *Cell* 2019;176(4):757-74 e23. doi: 10.1016/j.cell.2018.12.038 [published Online First: 20190131]

26. Yuan A, Hsiao YJ, Chen HY, et al. Opposite Effects of M1 and M2 Macrophage Subtypes on Lung Cancer Progression. *Sci Rep* 2015;5:14273. doi: 10.1038/srep14273 [published Online First: 20150924]

27. Zhang Y, Chen H, Mo H, et al. Single-cell analyses reveal key immune cell subsets associated with response to PD-L1 blockade in triple-negative breast cancer. *Cancer Cell* 2021;39(12):1578-93 e8. doi: 10.1016/j.ccell.2021.09.010 [published Online First: 20211014]

28. Schlenker R, Schwalie PC, Dettling S, et al. Myeloid-T cell interplay and cell state transitions associated with checkpoint inhibitor response in melanoma. *Med* 2024 doi: 10.1016/j.medj.2024.03.015 [published Online First: 20240404]

29. van Elsas MJ, Middelburg J, Labrie C, et al. Immunotherapy-activated T cells recruit and skewlate-stage activated M1-like macrophagesthat are critical for therapeutic efficacy. *Cancer Cell* 2024 doi: 10.1016/j.ccell.2024.04.011 [published Online First: 20240509]

30. Rannikko JH, Hollmen M. Clinical landscape of macrophage-reprogramming cancer immunotherapies. *Br J Cancer* 2024 doi: 10.1038/s41416-024-02715-6 [published Online First: 20240603]

31. Chen X, Zaro JL, Shen WC. Fusion protein linkers: property, design and functionality. *Adv Drug Deliv Rev* 2013;65(10):1357-69. doi: 10.1016/j.addr.2012.09.039 [published Online First: 20120929]

32. van Rosmalen M, Krom M, Merkx M. Tuning the Flexibility of Glycine-Serine Linkers To Allow Rational Design of Multidomain Proteins. *Biochemistry* 2017;56(50):6565-74. doi: 10.1021/acs.biochem.7b00902 [published Online First: 20171207]

33. Hallgren J, Tsirigos K, Pedersen MD, et al. DeepTMHMM predicts alpha and beta transmembrane proteins using deep neural networks: bioRxiv, 2022.

34. Pertel T, Reinhard C, Luban J. Vpx rescues HIV-1 transduction of dendritic cells from the antiviral state established by type 1 interferon. *Retrovirology* 2011;8(49) doi: https://doi.org/10.1186/1742-4690-8-49

35. Lehmann BD, Colaprico A, Silva TC, et al. Multi-omics analysis identifies therapeutic vulnerabilities in triple-negative breast cancer subtypes. *Nat Commun* 2021;12(1):6276. doi: 10.1038/s41467-021-26502-6 [published Online First: 20211101]

36. Mounir M, Lucchetta M, Silva TC, et al. New functionalities in the TCGAbiolinks package for the study and integration of cancer data from GDC and GTEx. *PLoS Comput Biol* 2019;15(3):e1006701. doi: 10.1371/journal.pcbi.1006701 [published Online First: 20190305]

37. Hatakeyama M, Opitz L, Russo G, et al. SUSHI: an exquisite recipe for fully documented, reproducible and reusable NGS data analysis. *BMC Bioinformatics* 2016;17(1):228. doi: 10.1186/s12859-016-1104-8 [published Online First: 20160602]

38. Andrews S. FastQC: A Quality Control Tool for High Throughput Sequence Data 2010 [Available from: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

39. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* 2018;34(17):i884-i90. doi: 10.1093/bioinformatics/bty560

40. Dobin A, Davis CA, Schlesinger F, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 2013;29(1):15-21. doi: 10.1093/bioinformatics/bts635 [published Online First: 20121025]

41. Frankish A, Diekhans M, Jungreis I, et al. Gencode 2021. *Nucleic Acids Res* 2021;49(D1):D916-D23. doi: 10.1093/nar/gkaa1087

42. Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read mapping by seed-and-vote. *Nucleic Acids Res* 2013;41(10):e108. doi: 10.1093/nar/gkt214 [published Online First: 20130404]

43. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014;15(12):550. doi: 10.1186/s13059-014-0550-8

44. Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* 2012;16(5):284-7. doi: 10.1089/omi.2011.0118 [published Online First: 20120328]

45. Liberzon A, Birger C, Thorvaldsdottir H, et al. The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* 2015;1(6):417-25. doi: 10.1016/j.cels.2015.12.004

46. House IG, Savas P, Lai J, et al. Macrophage-Derived CXCL9 and CXCL10 Are Required for Antitumor Immune Responses Following Immune Checkpoint Blockade. *Clin Cancer Res* 2020;26(2):487-504. doi: 10.1158/1078-0432.CCR-19-1868 [published Online First: 20191021]

47. Qu Y, Wen J, Thomas G, et al. Baseline Frequency of Inflammatory Cxcl9-Expressing Tumor-Associated Macrophages Predicts Response to Avelumab Treatment. *Cell Rep* 2020;32(1):107873. doi: 10.1016/j.celrep.2020.107873

48. Litchfield K, Reading JL, Puttick C, et al. Meta-analysis of tumor- and T cell-intrinsic mechanisms of sensitization to checkpoint inhibition. *Cell* 2021;184(3):596-614 e14. doi: 10.1016/j.cell.2021.01.002 [published Online First: 20210127]

49. Bill R, Wirapati P, Messemaker M, et al. CXCL9:SPP1 macrophage polarity identifies a network of cellular programs that control human cancers. *Science* 2023;381(6657):515-24. doi: 10.1126/science.ade2292 [published Online First: 20230803]

50. Sloas C, Beghi S, Huangfu Y, et al. Abstract 5249: Macrophages expressing synthetic cytokine receptors reverse IL10-mediated immunosuppression within solid tumors and promote adaptive immunity. *Cancer Research* 2024;84(6_Supplement):5249-49. doi: 10.1158/1538-7445.Am2024-5249

51. Sloas C, Beghi S, Huangfu Y, et al. Macrophages expressing synthetic cytokine receptors reverse IL-10 mdiated Immunosuppression within solid tumors and promote adaptive immunity https://74dfcf.p3cdn1.secureserver.net/wp-content/uploads/2024/04/AACR_2024_EM-C_FINAL.pdf2024 [accessed 01.10.2024.

52. Muller E, Christopoulos PF, Halder S, et al. Toll-Like Receptor Ligands and Interferon-gamma Synergize for Induction of Antitumor M1 Macrophages. *Front Immunol* 2017;8:1383. doi: 10.3389/fimmu.2017.01383 [published Online First: 20171026]

53. Lin Y, Xu J, Lan H. Tumor-associated macrophages in tumor metastasis: biological roles and clinical therapeutic applications. *J Hematol Oncol* 2019;12(1):76. doi: 10.1186/s13045-019-0760-3 [published Online First: 20190712]

54. Bernabei P, Coccia EM, Rigamonti L, et al. Interferon-γ receptor 2 expression as the deciding factor in human T, B, and myeloid cell proliferation or death. *Journal of Leukocyte Biology* 2001;70(6):950-60. doi: 10.1189/jlb.70.6.950

55. Walter MR. The molecular basis of IL-10 function: from receptor structure to the onset of signaling. *Curr Top Microbiol Immunol* 2014;380:191-212. doi: 10.1007/978-3-662-43492-5_9

56. Wrana JL, Attisano L, Carcamo J, et al. TGF beta signals through a heteromeric protein kinase receptor complex. *Cell* 1992;71(6):1003-14. doi: 10.1016/0092-8674(92)90395-s

FIGURE LEGENDS

Figure 1 *IL10* and *TGFB1* are expressed in TNBC patients. Shown is the percentile expression of *IL10* and *TGFB1*

expression across all transcripts in TNBC samples from TCGA database. Each dot represents one patient. n=195 TNBC

patients.

Figure 2 IL-10 / TGFβ stimulation induces STAT1 signaling in chimeric cytokine receptors (ChCR) expressing

THP-1 and primary macrophages A) Scheme of lentiviral constructs that were used. B) Scheme of ChCR. Binding of

IL-10 or TGFβ to the respective ChCR should induce pSTAT1 signaling. Created in BioRender. Traxel, S. (2024)

BioRender.com/y85b161 C/D) THP-1 were either left untransduced or transduced with mock, IL-10 ChCR or TGFβ ChCR

construct. C) Expression of IL-10Rα, IL-10Rβ and TGFβR2 was assessed by flow cytometry. Shown are representative

histograms. Cells are gated on tNGFR+ cells. D) THP-1 were either left untreated, stimulated with 1, 10 or 100ng/ml IL-

10 or TGFβ, or 5ng/ml IFNγ for 30min. pSTAT1 signaling was assessed by Phosphoflow. Shown are median fluorescence

intensities (MFI's, mean ± SD) in single cells for untransduced or in tNGFR+ cells for mock, IL-10 ChCR or TGFβ ChCR.

N=2-3 E/F) Primary human monocytes were either transduced with mock, IL-10 ChCR or TGFβ ChCR construct. E)

Transduction efficiency (tNGFR+) was assessed by flow cytometry upon differentiation to macrophages. Shown are %

tNGFR+ cells (mean ± SD). N=14-33 F) Expression of IL-10Rα, IL-10Rβ and TGFβR2 was assessed by flow cytometry.

Shown are MFIs gated on tNGFR+ cells (mean ± SD) from 4-5 healthy donors from 3-4 independent experiments G)

Transduced macrophages were either left untreated or stimulated as indicated for 30min. pSTAT1 signaling was assessed

by Phosphoflow. Shown are pSTAT1 MFIs in tNGFR+ cells (mean ± SD) from 4 donors in 4 independent experiments

(IL-10 ChCR) or 5 donors in 3 independent experiments (TGFβ ChCR). p-values were calculated by ratio paired t-tests. *

p < 0.05, ** p<0.01, *** p<0.001, **** p<0.00001

Figure 3 ChCR expressing macrophages have an inflammatory phenotype upon IL-10 or TGFβ stimulation A/B)

Primary human monocytes were either left untransduced or transduced with mock, IL-10 ChCR or TGFβ ChCR construct.

After differentiation to macrophages, cells were either left unstimulated or stimulated with 100ng/ml IL-10, 10 or 100ng/ml

TGFβ, or 10ng/ml IFNγ. Expression of surface marker was assessed by flow cytometry. A) Shown are median fluorescence

intensities (MFIs, mean ± SD) after subtracting isotype MFI or % CD38+ or % CD163+ gated on CD11b+ cells from 3-4

healthy donors in 4 independent experiments. B) Shown are MFI (mean ± SD) after subtracting isotype MFI or % CD38+

gated on CD11b+ (untransduced) or NGFR+ (transduced) cells from 3 healthy donors in 2 independent experiments. For

MFIs, p-values were calculated using ratio paired t-tests. For % positive cells, p-values were calculated using paired t-tests.

* p < 0.05, ** p<0.01, *** p<0.001, **** p<0.00001

Figure 4 ChCR expressing macrophages have a IFNγ-like transcriptome A-D) Monocytes were either transduced with

mock, IL-10 ChCR or TGFβ ChCR lentivirus. After differentiation to macrophages, cells were either left unstimulated or

stimulated with 100ng/ml IL10, 100ng/ml TGFβ or 10ng/ml IFNγ for 2 days. mRNA was isolated and subjected to bulk

RNA sequencing. A) Gene expression principal component analysis, n=3 donors B) Differential gene expression analysis

of mock transduced versus IL-10 ChCR transduced macrophages upon IL-10 stimulation and mock transduced versus

TGFβ ChCR transduced macrophages upon TGFβ stimulation. Genes highlighted in orange are part of hallmark IFNγ

pathway. Threshold for Log2 fold changes was set to <-1 or > 1. P-values were calculated using the Wald test and adjusted

for multiple comparison using the Benjamini-Hochberg method. Threshold of false discovery rate (FDR) was set to 0.05.

n=3 donors. C) Gene set enrichment analysis (GSEA) of hallmark gene sets. Shown is the normalized enrichment score

(NES) of the top 5 enriched hallmark gene sets. D) GSEA of Gene ontology (GO) / biological processes (BP) gene set.

Shown is the NES of top 10 enriched BP.

Figure 5 ChCR expressing macrophages upregulate genes associated with good response to anti-PDL-1 treatment

A-D) Monocytes were either transduced with mock, IL-10 ChCR or TGFβ ChCR lentivirus. After differentiation to

macrophages, cells were either left unstimulated or stimulated with 100ng/ml IL10, 100ng/ml TGFβ or 10ng/ml IFNγ for

2 days. A) mRNA was isolated and subjected to bulk RNA sequencing. *CXCL9* and *CXCL10* expression analyzed by bulk

RNAseq. Shown are normalized log2 counts (mean ± SD). N=3 donors. P-values were calculated using the Wald test and

adjusted for multiple comparison using the Benjamini-Hochberg method. * False discovery rate (FDR) < 0.05, **

FDR<0.01, *** FDR<0.001, **** FDR<0.00001 B) Supernatants were collected and analyzed by CXCL10 ELISA. Shown

are CXCL10 concentration (mean ± SD). N=4 donors analyzed in 2-3 independent experiments. p-values were calculated

using paired t-tests. * p < 0.05, ** p<0.01, *** p<0.001, **** p<0.00001 C-D) mRNA was isolated and subjected to bulk

RNA sequencing. C) Heatmap of genes enriched in response associated with response to anti-PD-L1 treatment in TNBC

patients.[27] Expression in genetically engineered macrophages is shown. D) Gene set enrichment analysis of genes expressed

in macrophages and associated with response (R) and no response (NR) to anti-PD-L1 treatment in TNBC patients.[27]

Figure 6 ChCR expressing macrophages have anti-tumoral activity A-F) Monocytes were either left untransduced or

transduced with mock, IL-10 or TGFβ ChCR and were differentiated to macrophages. Spheroids were formed by mixing

1000 MDA-MB-231 TNBC cells and 1500 MRC-5 fibroblasts. 1 day later, genetically engineered macrophages were added

at a 1:10 ratio (MDA-MB-231: macrophages). Spheroids were either left unstimulated, stimulated with IL-10 (100ng/ml),

TGFβ (10ng/ml), or IFNγ (10ng/ml). A/B) Quantification of tumor cells in spheroids by measuring mean EGFP intensity

(mean ± SD) over time. Shown are results from 3 donors from 2 independent experiments. C/D) Quantification of tumor

cells in spheroids by measuring mean EGFP intensity (mean ± SD) 6d after adding macrophages. Shown are results from

3 donors from 2 independent experiments. E/F) Quantification of tumor cells in spheroid by measuring luciferase activity

6d after adding macrophages. Shown are relative light units (RLU) (mean ± SD) after subtraction of background RLU from

3 donors analyzed in 2 independent experiments. p-values were calculated using paired t-tests comparing the ChCRs with

mock condition. * p < 0.05, ** p<0.01, *** p<0.001, **** p<0.00001

FIGURES

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

ONLINE SUPPLEMENTAL FIGURES

Online supplemental figure S1 IL10 ChCR induces pSTAT1 signaling in HEK-Blue IFNγ reporter cell line. A) Scheme of IL10 ChCR constructs with single chain constructs (IL10RA- IFNGR1and IL10RB-IFNGR2) and the construct containing both chains (IL10RA-IFNGR1 P2A IL10RB-IFNGR2). B) IL10Ra and IL10Rb expression in HEK-Blue transduced with IL10Ra ChCR, IL10Rb ChCR and IL10Rab ChCR. Expression of IL10Ra and IL10Rb was analyzed by flow cytometry. C) HEK-Blue cells were either left untransduced or transduced with IL10Ra ChCR, IL10Rb ChCR or IL10Rab ChCR. Cells were stimulated as indicated for 24h. IFNγ signaling activity was measured using Quanti-Blue assay. Shown is OD 620nm from three independent experiments. P-values were calculated using 2-way ANOVA and multiple comparison with Tukey correction for multiple comparison. * p < 0.05, ** p<0.01, *** p<0.001, **** p<0.00001

Online supplemental figure S2: Conditioned medium of ChCR expressing macrophages affects viability of MDA-MB-231 and BT-549. Monocytes were transduced, differentiated to macrophages and stimulated for 2 days as indicated. Generated conditioned medium was used to treat MDA-MB-231 or BT-549 cell line for 3 days. Viability was measured by WST-1 assay. Shown are results from using conditioned medium from 3-5 donors. p-values were calculated using paired t-test by comparing ChCR vs. mock for each stimulation. * $p < 0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.00001$

SUPPLEMENTARY TABLE

Supplementary Table 1: Uniprot aminoacid (AA) sequences used for the design of the ChCR chains

	Signal peptide	Extracellular domain	Linker	Transmembrane domain	Cytosolic domain
IL10RA ChCR	IFNGR1 AA1-17	IL10RA AA22-235	Glycine serine	IFNGR1 AA246-266	IFNGR1 AA267-489
IL10RB ChCR	IFNGR2 AA1-21	IL10RB AA20-220	Glycine serine	IFNGR2 AA248-268	IFNGR2 AA269-337
TGFBR1 ChCR	IFNGR1 AA1-17	TGFBR1 AA34-126	Glycine serine	IFNGR1 AA246-266	IFNGR1 AA267-489
TGFBR2 ChCR	IFNGR2 AA1-21	TGFBR2 AA23-166	Glycine serine	IFNGR2 AA248-268	IFNGR2 AA269-337