Macrophages expressing chimeric cytokine receptors have an inflammatory phenotype and anti-tumoral activity upon IL-10 or TGFβ stimulation

Sabrina Traxel[1], Florian Schmidt[1], Corina Beerli[1], Dinh-Van Vuong[1], Roberto F. Speck[1*], Simon Bredl[1*]

[1] Department of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, University of Zurich, Raemistrasse 100, 8091 Zurich, Switzerland

*co-senior authors

ORCID IDS:

ST 0000-0002-9193-5586

FS 0009-0006-4040-3316

CB 0000-0002-0062-2833

RFS 0000-0002-8453-1137

SB 0000-0001-6822-755X

CORRESPONDING ADDRESSES:

Dr. rer. nat. Simon Bredl

Department of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, University of Zurich, Raemistrasse 100, 8091 Zurich, Switzerland

e-mail: simon.bredl@gmail.com

tel: +41 43 253 32 78

Dr. sc. nat. Sabrina Traxel,

Department of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, University of Zurich, Raemistrasse 100, 8091 Zurich, Switzerland

e-mail: sabrinatraxel@gmail.com

tel: +41 43 253 44 15

ABSTRACT

Background

Triple-negative breast cancer (TNBC) is a highly aggressive subtype of breast cancer that lacks hormone receptors and

HER2 amplification, making it unresponsive to standard hormone or HER2-targeted therapies. Although immune

checkpoint inhibitors (ICIs) have shown promise in tumors with high lymphocyte infiltration, their efficacy remains

limited in tumors with minimal lymphocyte infiltration. To overcome this challenge, we developed a novel approach to

reprogram tumor-associated macrophages (TAMs) from an immunosuppressive to an inflammatory phenotype within the

tumor microenvironment (TME), aiming to enhance therapeutic outcomes in TNBC.

Methods

We designed a chimeric cytokine receptor (ChCR) that triggers STAT1 signaling upon stimulation with IL-10 or TGFβ,

cytokines prevalent in the TME that typically drive immunosuppression. Human primary macrophages were transduced

with a lentiviral vector expressing the ChCR. Changes in their phenotype, secretome, and transcriptome were analyzed

following stimulation. The anti-tumoral activity of these reprogrammed macrophages was assessed using co-culture

assays with 3D TNBC spheroids.

Results

ChCR-expressing macrophages showed robust STAT1 activation in response to IL-10 or TGFβ stimulation, resulting in

an inflammatory phenotype similar to IFNγ activation, as confirmed by phenotypic markers, and transcriptomic profiling.

These ChCR-stimulated macrophages demonstrated significant anti-tumoral effects in 3D TNBC spheroids. Moreover,

ChCR stimulation led to the upregulation of *CXCL9* and *CXCL10*, chemokines essential for lymphocyte recruitment, and

genes associated with good response to ICIs.

Conclusion

We successfully engineered ChCRs that reprogram TAMs within an IL-10- and TGFβ-rich environment, inducing an

inflammatory phenotype and anti-tumoral activity. The expression of *CXCL9* and *CXCL10* further supports lymphocyte

recruitment, potentially facilitating greater lymphocyte infiltration and disrupting the immunosuppressive TME. This

approach offers a promising strategy to improve immunotherapy outcomes in TNBC patients, particularly those with low

immune cell infiltration, thereby addressing a critical unmet clinical need.

What is already known on this topic

Low T cell infiltration in triple negative breast cancer is associated with bad prognosis and non-responsiveness to

immune checkpoint inhibitor in combination with chemotherapy. The presence of macrophages with an inflammatory

phenotype is associated with T cell infiltration and better prognosis. The tumor microenvironment (TME), however, is

rich in immunosuppressive cytokines like IL-10 and TGFβ, rendering macrophages immunosuppressive. Thus,

reprogramming macrophages in the TME could benefit patients with low T cell infiltration.

What this study adds

We have designed a chimeric cytokine receptors (ChCR) that bind IL-10 or TGFβ and induces an IFNγ-like

inflammatory signaling and thus utilizes the local presence of immunosuppressive cytokines to reprogram macrophages.

ChCR expressing macrophages show an inflammatory phenotype in the presence of immunosuppressive cytokines IL-10

or TGFβ and have an anti-tumoral activity *in vitro*.

How this study might affect research, practice or policy –

Our study is a starting point to explore ChCR-expressing macrophages as an adoptive cell therapy. Such therapy has the

potential to attract T cells into the tumor and thereby boost the adaptive anti-tumoral immune response and response to

immune checkpoint inhibitors. Moreover, our data suggest that such genetically-engineered macrophages have direct

anti-tumoral activity.

LIST OF ABBREVIATIONS

CAR   Chimeric antigen receptor

ChCR   Chimeric cytokine receptor

EF1α   Elongation factor 1 alpha core promoter

FDR   False discovery rate

GO:BP   Gene Ontology biological process

hPGK   Human phosphoglycerate kinase promoter

ICI   Immune checkpoint inhibitor

MFI   Median fluorescence intensity

MOI   Multiplicity of infection

NES   Normalized enrichment score

NR   No response

P/S   Penicillin-Streptomycin

PCA   Principle component analysis

R   Response

RH10   RPMI with P/S and 10% human AB serum

RLU   Relative light units

RNAseq   RNA sequencing

SEAP   Secreted embryonic alkaline phosphatase

SFFV   Spleen focus-forming virus promoter

TAM   Tumor associated macrophage

TCGA   The Cancer Genome Atlas

TME   Tumor microenvironment

TNBC   Triple negative breast cancer

tNGFR   Truncated nerve growth factor receptor

99 INTRODUCTION

Triple negative breast cancer (TNBC) is an aggressive subtype of breast cancer that lacks both hormone receptor

expression and HER2 amplification and is therefore not amenable to hormone or HER2 targeted therapies (reviewed[1]).

Current first line treatment for both high-risk early-stage TNBC and advanced stage, PD-L1 positive (combined positive

score ≥10) TNBC is the immune checkpoint inhibitor (ICI) pembrolizumab in combination with chemotherapy. Early

stage TNBC has a good prognosis with 86.6% overall survival at 60 months when treated neoadjuvant with

pembrolizumab and chemotherapy.[2] In patients with advanced stage TNBC the pembrolizumab and chemotherapy

combination minimally improves the median overall survival to 23 months, compared to 16.1 months in the placebo and

chemotherapy group.[3] Thus, there is clearly an unmet need for patients with advanced TNBC.

Adoptive cell therapy using chimeric antigen receptor (CAR) T cells has achieved remarkable success in hematological

malignancies[4] and there was great hope that CAR T cell therapy would also change the therapeutic landscape for solid

tumors. To date, CAR T cells had only limited success in solid tumors.[5] Several phase I clinical trials of CAR-based

immunotherapies for TNBC are currently underway.[6] While we eagerly await the first results of adoptive CAR T cell

therapies in TNBC, alternative innovative treatments must be sought.

A critical determinant of disease progression of solid tumors including TNBCs is their immunophenotype, which is based

on the number and localization of CD8+ T cells in the tumor microenvironment (TME).[7-9] Three main

immunophenotypes have been defined: hot (high number of intratumoral T cells), excluded (T cells restricted to the

tumor margin) or cold (lack of T cells) (reviewed[10]). Patients with a hot TME, including patients suffering from TNBC,

have a better prognosis and show better response to ICI treatment than patients with an excluded or cold TME. [7 9 11 12] In

TNBC, 42-54% of patients have a tumor with excluded or cold immunophenotype.[7 9 13] Therefore, shifting a cold into a

hot TME is of great therapeutic interest.

Monocytes are actively recruited into the TME and there, differentiate to tumor associated macrophages (TAMs), which,

together with other myeloid cells, they constitute up to 30% of the tumor mass.[14 15] Macrophages can adopt a wide range

of phenotypes depending on the stimuli received and on the microenvironment.[16] Consequently, their activities include

immunosuppression, tissue repair, phagocytosis, inflammation, and cytotoxicity. *In vitro*, we can model extreme

macrophage phenotypes: Inflammatory macrophages are induced by pro-inflammatory signals like IFNγ and/or LPS,

whereas immunosuppressive macrophages are induced by immunosuppressive cytokines like IL-4, IL-13, IL-10, or

TGFβ. Notably, these extreme phenotypes cannot describe the complexity of phenotypes in macrophages found *in vivo*,

where features of both extremes can be found within the same cell (reviewed[17]). The presence of IL-10 and TGFβ in the

TME of solid tumors, such as in TNBC[18-20], however, pushes TAMs towards an immunosuppressive phenotype. These

pro-tumoral TAMs inhibit the immune system, support tumor growth, angiogenesis, and tumor cell migration

(reviewed[21]). In contrast, inflammatory macrophages have anti-tumoral activity by attracting T cells, by activating

adaptive immune response, and by their direct tumoricidal effects.[22][23] (reviewed[24]) In fact, tumors with high number of

inflammatory macrophages are more likely to have a hot TME, a better prognosis and better response to ICIs.[7][22][25][26]

Thus, reprogramming TAMs towards an inflammatory phenotype is a hot topic in the field of cancer immunotherapy

(reviewed[17]).

We believe that macrophages should be reprogrammed to switch to inflammatory and anti-tumoral macrophages locally

in the TME. Systemically activated macrophages may be retained in various tissues, unable to reach their destination.

Furthermore, local activation would be less likely to have off-target effects. Therefore, we developed chimeric cytokine

receptor (ChCR) expressing macrophages that utilize immunosuppressive cytokines, prevalent in the TME, to switch to

inflammatory macrophages. Specifically, we engineered macrophages to express a ChCR that binds either IL-10 or

TGFβ and induces an IFNγ signaling.

METHODS

Ethics statement

Monocytes used were obtained from buffy coats, anonymously provided by the Blood Donation Service Zurich, Swiss

Red Cross, Schlieren. Written consent for the use of buffy coats for research purposes was obtained from blood donors

by the Blood Donation Centre.

Cell lines

HEK293T (ATCC) were maintained in DMEM (high glucose, Sigma-Aldrich, D6429) with 10% heat-inactivated FBS

(Gibco, 10270-106) (hereafter called D10). HEK-Blue™ IFN-γ cells (Invivogen, hkb-infg)) were maintained in D10 with

100μg/ml Zeocin (Invivogen, ant-zn-1) and 30μg/ml Blasticidin (Invivogen, ant-bl-1). MDA-MB-231 (ATCC, HTB-26)

were maintained in RPMI-1640 (Sigma-Aldrich, R8758) with 10% heat-inactivated FBS (hereafter called R10). BT-549

(ATCC, HTB-122) were maintained in R10 with 10ug/ml insulin (Sigma- Aldrich, I9278). THP-1 (ATCC) were

maintained in R10 with 2-mercaptoethanol (Gibco, 31350010). MRC-5 (ATCC, CCL-171) were maintained in EMEM

(ATCC, 30-2003) with 10% heat-inactivated FBS. All cells were maintained at 5% CO_2 and 37°C in a humidified

incubator.

Monocyte derived macrophages

Peripheral blood mononuclear cells were isolated from buffy coats of healthy donors (Blood Donation Service Zurich,

Swiss Red Cross, Schlieren) using density gradient centrifugation on lymphoprep (Axon Lab, 12053230). Monocytes

were isolated using CD14 microbeads (Miltenyi Biotec, 130-050-201) and plated at 150'000 cells/cm^2 in F+ high

attachment plates (Sarstedt) in RPMI-1640 with Penicillin-Streptomycin (P/S, Gibco, 15140-122). Two hours after

plating, medium was replaced with RPMI-1640 with P/S and 10% human AB serum (Sigma Aldrich, H6914), hereafter

called RH10. Monocytes were differentiated for 6-11 days with medium change every 2-3 days. In some experiments,

macrophages were stimulated for additional 48h with IFNγ (Biolegend, 570204), IL-10 (Biolegend, 571004) or TGFβ1

(Peprotech, 100-21) at indicated concentrations.

Plasmid construction

To generate ChCR plasmids, amino acid sequences of the human IL10RA, IL10RB, TGFβR1, TGFβR2, IFNGR1 and

IFNGR2 receptor were obtained from UniProt (Supplementary Table 1). The ChCR were constructed as follows: The

extracellular domains of IL10RA, IL10RB, TGFβR1, or TGFβR2 were fused to the transmembrane and extracellular

domain of the IFNGR1 or IFNGR2, including a glycine-serine linker for flexibility.[27] As an N-terminal signal peptide,

we used the one from the corresponding intracellular IFNGR. All variants were analyzed with DeepTMHMM to verify *in*

silico signal peptide cleavage, transmembrane domains and surface expression.[28] Sequences were ordered codon-

optimized at GenScript Biotech Corporation. For expression of single chains, ChCR sequences were sub-cloned into a

pCWX Dest lentiviral vector (Gift from Patrick Salmon) under the control of an elongation factor 1 alpha core promoter

(EF1α). To simultaneously express both chains, ChCR sequences were subcloned into a pCWX Dest lentiviral vector

separated by a P2A sequence and under the control of a spleen focus‐forming virus (SFFV) promoter. All lentiviral

vectors also encode for a truncated nerve growth factor receptor (tNGFR) driven by a human phosphoglycerate kinase

(hPGK) promoter as a reporter for transduction. Mock lentiviral vector only encodes for tNGFR.

To generate the Luciferase-EGFP plasmid, luciferase and EGFP sequences, separated by a T2A sequence, were ordered

as gene blocks and sub-cloned into a pCWX Dest lentiviral vector under the control of EF1α promoter.

Lentivirus production and titration

Lentiviral vectors were produced in HEK293T. Briefly, 13 million cells were plated in T150 flasks. The day after, cells

were co-transfected using PEI (Polysciences, 23966) with pMD2.G, p8.9NDSB-DPAVDLL[29] (gift from Jeremy Luban),

pcDNA3.1 Vpx239[29] (gift from Jeremy Luban) and pCWX Dest transfer vector at a molar ratio of 251:107.5:756.1:45.

4h after transfection, medium was changed and supplemented with 5mM sodium butyrate (Sigma Aldrich, B5887). Virus

was harvested 52h after transfection and clarified by centrifugation. Lentiviral supernatants were concentrated by

ultrafiltration using 100kDa cut-off Amicon filters (Merck Milipore, UFC910024). Titration of lentiviruses was done by

transducing HEK293T. Three days after transduction, transduced cells were stained for tNGFR and analyzed by flow

cytometry on a LSR II Fortessa (BD Biosciences).

Stable cell line generation

THP-1 were transduced with lentiviruses at a multiplicity of infection (MOI) 1 by spinoculation (800g, 45min, 32°C) in

presence of protamine sulfate (100ug/ml, Sigma-Aldrich, P4020). Transduced cells (tNGFR+) were sorted using

MACSelect LNGFR microbeads (Miltenyi Biotec, 130-091-330).

MDA-MB-231 cell line was transduced with the lentiviral construct to express Luciferase and EGFP. Cells were sorted

for high EGFP expression on a BD FACSAria III (BD Biosciences).

Monocyte transduction

To transduce monocytes, lentivirus was added 2h after plating of monocytes in RPMI with P/S and 5% human AB serum at a MOI of 10 or 15 in presence of protamine sulfate (100ug/ml, Sigma-Aldrich, P4020). After overnight incubation, medium was changed to RH10. Transduction efficiency was assessed between day 6 and day 12 of differentiation and was defined as %tNGFR+ cells. In general, macrophages with transduction rates < 65% were excluded from experiments. In early experiments assessing IL-10 ChCR macrophage polarization, transduction rates ≥ 55% were accepted.

IFNγ signaling reporter cells

HEK-Blue^{TM} IFN-γ were transduced with lentiviruses by spinoculation in the presence of 8ug/ml polybrene (SantaCruz, sc-134220). To measure secreted embryonic alkaline phosphatase (SEAP), transduced cells were detached mechanically and seeded in 96-well plate. Cells were incubated with cytokines as indicated for 24h. Thereafter, 30ul of the cell supernatant was incubated with 170ul Quanti-Blue solution (Invivogen, rep-qbs) and incubated for 1h at 37°C. OD was measured at 620nm using Spectramax iD3 plate reader (Molecular Devices).

Flow cytometry analysis

To assess transduction rate and polarization marker expression, macrophages were harvested for flow cytometry by incubation with 5mM EDTA in PBS on ice for 45min, followed by gentle pipetting. Detached macrophages were first treated with True-Stain FcX (Biolegend, 422301) and thereafter stained with antibodies (Table 1). Stained samples were acquired on LSR II Fortessa (BD Life Sciences). Untransduced or mock transduced cells were used as gating control for tNGFR and IL-10Rα, IL-10Rβ, and TGFβR2, respectively. Data was analyzed using FlowJo v10.10 software (BD Life Sciences). Corresponding isotypes were used to normalize median fluorescence intensities (MFI) by subtraction or as a gating control.

Table 1: Flow cytometry antibodies

Epitope	Fluorochrome	Company	Category Number
anti-mouse	BV421	BioLegend	405317
anti-rat	BV421	BioLegend	405414
CD11b m/h	FITC	BioLegend	101206
CD163	BV711	BioLegend	333630
CD206	PE	BioLegend	321105
CD38	BV605	BioLegend	303532
CD86	BV785	BioLegend	305441
HLA-DR	BV510	BioLegend	307646

IgG1 Isotype	BV711	BioLegend	400167
IgG1 Isotype	PE	BD Pharmingen	555749
IgG1 Isotype	BV605	BioLegend	400161
IgG2b Isotype	BV785	BioLegend	400355
IgG2a Isotype	BV510	BioLegend	400267
IL-10Rα	unconjugated	BioLegend	308802
IL-10Rα	PE	Miltenyi Biotec	130-121-341
IL-10Rβ	unconjugated	R&D Systems	MAB874
IL-10Rβ	APC	R&D Systems	FAB874A
NGFR (CD271)	Alexa Fluor 647	BD Biosciences	560326
NGFR (CD271)	BV510	BD Biosciences	563451
NGFR (CD271)	APC	BioLegend	345108
pSTAT1 (Tyr701)	PE	BioLegend	666403
TGFβR2	PE	BioLegend	399703

STAT1 phosphoflow

THP-1 or human primary macrophages were used for STAT1 phosphoflow experiments. THP-1 were plated in round

bottom 96-well plates in R10. Macrophages were detached as described for flow cytometry analysis and plated in round

bottom ultra low attachment 96-well plates (Corning, 7007) in RH10. Cells were rested for 1h at 37°C and thereafter

stimulated for 30min as indicated. Cells were fixed by adding one volume of 4% PFA (Santa Cruz, sc-281692) and

15min incubation at 37°C. Next, cell surface markers were stained before cells were permeabilized with True-Phos Perm

Buffer (Biolegend, 425401). Cells were stained for intracellular pSTAT1 (Tyr701) and analyzed on LSR II Fortessa (BD

Biosciences).

IL10 and *TGFB1* expression in TNBC

Gene expression data from 195 TNBC patients were obtained from The Cancer Genome Atlas (TCGA). TNBC patients

in the TCGA dataset were previously identified by Lehmann et al.[30] Raw counts were retrieved, preprocessed and

normalized using the R package TCGABiolinks (2.33.0) without filtration.[31] Percentile expression of *IL10* and *TGFB1*

across all genes was calculated by dividing the rank expression with the total analyzed genes.

RNA sequencing (RNAseq)

RNAseq and preprocessing

For bulk RNAseq, monocytes were transduced and differentiated into macrophages. Macrophages were stimulated for

48h as indicated. Total RNA was extracted using Quiashredder (Qiagen, 79654)) and RNeasy Mini Kit (Qiagen, 74104)

with on column DNA digestion (Qiagen, 79254). RNAseq was performed at the Functional Genomics Center Zurich

(FGCZ). The NovaSeq X Plus (Illumina, Inc, California, USA) was used for cluster generation and sequencing according

to standard protocol. Sequencing was paired end 150 bp. Between 28 and 86 million reads were generated per sample.

RNAseq analysis was performed using the SUSHI framework,[32] which encompassed the following steps: Read quality

was inspected using FastQC,[33] and sequencing adaptors removed using fastp,[34] alignment of the RNA-Seq reads using

the STAR aligner,[35] and with the GENCODE human genome build GRCh38.p13 (Release 42) as the reference[36], the

counting of gene-level expression values using the featureCounts function of the R package Rsubread.[37] Unless stated

otherwise, all R functions were executed on R version 4.3.2 (R Core Team, 2020) and Bioconductor version 3.17.

RNAseq analysis

Differential expression analysis was performed using the generalized linear model as implemented by the DESeq2

Bioconductor R package.[38] Principle component analysis (PCA) was performed using the prcomp function from stats R

package. For gene set enrichment analysis (GSEA), differentially expressed genes were ranked by log2 fold change.

Gene Ontology (GO) GSEA tests were performed using the gseGO function respectively of the clusterProfiler

Bioconductor R package.[39] Hallmark and custom pathways GSEA was performed using the GSEA function of the

Bioconductor R package clusterProfiler (4.13.0)[39] and the hallmark gene sets (MSigDb 2023.2)[40] on R version (4.4.0).

Analysis of previously published scRNAseq data

We compared the transcriptome of ChCR macrophages with previously described scRNAseq data of macrophages from

TNBC patients with or without response to anti-PD-L1 atezolizumab.[25] Using the R package VennDiagram (1.7.3), we

determined the overlap of significantly upregulated genes in ChCR macrophages (log2FC>0.5 and false discovery rate

(FDR)<0.05) and genes upregulated in baseline tumor macrophages of anti-PD-L1 responders previously published.

Moreover, we performed GSEA with costum pathways generated from genes upregulated in baseline tumor macrophages

from anti-PD-L1 responders and non-responders, respectively. GSEA was performed using the GSEA function of the

Bioconductor clusterProfiler (4.13.0)[39] on R version (4.4.0).[25]

CXCL10 ELISA

To analyze CXCL10 secretion, differentiated macrophages were stimulated for 48h as indicated. Supernatants were

collected, cell debris removed by centrifugation, and stored at -80°C. CXCL10 in supernatant was measured using the

ELISA MAX™ Deluxe Set Human CXCL10 (IP-10) kit (Biolegend, 439904). OD450nm and OD570 as background

were measured using Multiskan SkyHigh plate reader (Thermo Scientific™).

3D spheroid assay

Heterotypic spheroids were generated by mixing 1000 MDA-MB-231, expressing EGFP and luciferase, and 1500 MRC-

5 cells in Akura™ 96 Spheroid Microplates (InSphero, CS-PB15) in R10. Plates were centrifuged at 100g for 5 min to

remove air bubbles and incubated at 37°C, 5% CO_2 overnight. To support the aggregation and maturation of the

spheroids, the plates were placed on a tilting stand at a 30° angle. The following day, macrophages were detached as

described for flow cytometry analysis, added at a 10:1 ratio (macrophages:MDA-MB-231 at seeding), and stimulated as

indicated. Medium and stimuli were replaced every 2-3 days. Fluorescent (EGFP) images were captured using the

Cytation 3 (BioTek) with a 4x objective. Images were processed and analyzed with the Gen5 (BioTek) software. The

mean fluorescence intensity of the full image was determined and the background of a medium control well subtracted.

Luciferase assay

As an endpoint quantification of tumor cells in the spheroids, luciferase activity of the remaining MDA-MB-231 was

determined at day 6. Spheroids were washed twice with PBS to remove luciferase in the supernatant and left with 50 ul

residual PBS per well. 50 ul of BrightGloTM substrate (Promega, E2610) was added per well to lyse remaining cells.

The lysate was transferred into a 96-well white plate (Pierce™, 15042) and incubated for minimum 2min. Luciferase

activity was determined using GloMax® Explorer luminometer (Promega, GM3500) and background signal from a

medium control well was subtracted.

WST1 assay

To assess the ability of genetically-engineered macrophages to affect TNBC proliferation, macrophages were stimulated

for 2d with IFNγ, IL-10 or TGFβ at indicated concentrations. Supernatants were harvested, clarified by centrifugation

and stored at -80°C. For WST1 assay, 5'000 BT-549 or 10'000 MDA-MB-231 cells were plated in 96-well plates. The

next day, medium was replaced with supernatants from stimulated macrophages or RH10 as untreated control. 72h after

addition of supernatants, viability was measured by incubating cells for 1h with cell proliferation reagent WST-1 (Roche,

5015944001). OD440nm and OD650nm as background were measured using Multiskan SkyHigh plate reader (Thermo

Scientific™).

Statistical analysis

The statistical analysis of TCGA and previously published single cell RNAseq data was performed in R (4.4.0). The

statistical analysis of bulk RNAseq data was performed using the Sushi framework of the FGCZ.[32] The remaining

statistical analysis was performed using Graphpad Prism 10.2.0 using statistical tests specified in the figure legends.

Data availability

RNA-Seq data are deposited on the Gene Expression Omnibus database under accession number GSE280810.

RESULTS

IL10 and *TGFB1* are expressed in TNBC

IL-10 and TGFβ1 have previously been shown to be expressed in TNBC patients.[18-20] To confirm this data, we analyzed

publicly available RNAseq data from 195 TNBC patients for *IL10* and *TGFB1* expression. We found that *IL10* and

TGFB1 were expressed among the top 73% (median) and top 90% (median) genes, respectively (figure 1A). Thus,

TCGA gene expression analysis confirms previously published results of IL-10 and TGFβ expression in TNBC patients.

IL-10 and TGFβ ChCR induce STAT1 signaling

To convert IL-10 or TGFβ signals into IFNγ signals, we designed two ChCR, which consist of the extracellular domains

of IL-10 receptor or TGFβ receptor, respectively, fused to the intracellular domains of the IFNγ receptor (figure1B and

C). To control for transduction efficacy, we also incorporated the tNGFR in the lentiviral vector.

In a first instance, we verified IFNγ signaling through the IL-10 ChCR in HEK-Blue IFNγ reporter cell line. In the HEK-

Blue IFNγ reporter cell line, IFNγ-induced STAT1 signaling leads to expression of SEAP, which can be measured in the

supernatant using a colorimetric assay. We stably expressed either the full IL-10 ChCR or the individual components, i.e.

IL-10RA fused to IFNGR1 and IL-10RB fused to IFNGR2 in the HEK-Blue IFNγ reporter cells (online supplemental

figure S1A). All cell lines were transduced at >90% efficiency (data not shown) and ChCR expression was confirmed by

flow cytometry (online supplemental figure S1B). We found that IL-10 induced IFNγ signaling in a dose-dependent

manner when the full IL-10 ChCR was expressed (online supplemental figure S1C). Surprisingly, IL-10RA-IFNGR1

alone triggered moderate IFNγ signaling, but both chains of the ChCR were necessary to achieve maximum STAT1

signaling. We also verified that the expression of the full TGFβ ChCR in HEK-Blue IFNγ reporter cell line induced

STAT1 signaling (data not shown). Next, we proceeded to the characterization of both ChCRs in the monocyte cell line

THP-1, which more closely approximates the properties of primary human macrophages. We found a prominent

upregulation of IL-10Rα and TGFβR2 and a moderate one of IL10Rβ using flow cytometry (figure1D). We tested

several TGFβR1 antibodies, all suffered from non-specificity and therefore we do not have expression data for TGFβR1.

Since TGFβR1 is on the first position of the polycistronic vector, and TGFβR2 expression is detectable, we assume that

TGFβR1 is expressed as well. Importantly, IL-10 and TGFβ induced a dose-dependent activation of STAT1 in IL-10 or

TGFβ ChCR-expressing THP-1 cells, respectively (figure 1E). As expected, untransduced and mock transduced THP-1

cells only showed pSTAT1 signaling upon IFNγ stimulation. To investigate the ChCRs in primary macrophages, we first

optimized the protocol for transducing monocytes. We achieved a transduction efficiency of 70.2 ± 14.5% (mean±SD)

across all vectors (figure 1F). Similar to ChCR expression in THP-1, we saw a very prominent upregulation of IL-10Rα

and TGFβR2 and a lesser one of IL-10Rβ (figure 1G). IL-10 induced STAT1 activation to a similar level like IFNγ,

whereas TGFβ induced only a little, but consistent STAT1 activation in the macrophages expressing the ChCRs

(figure1H). Thus, both IL-10 and TGFβ ChCR variants trigger the IFNγ signaling pathway.

ChCR expression induces an inflammatory phenotype upon IL-10 / TGFβ stimulation

We next assessed whether triggering the ChCRs induces an inflammatory phenotype in primary human macrophages. We

focused on the expression of polarization markers typically upregulated on macrophages either upon IFNγ (i.e., HLA-

DR, CD38, CD86) or IL-10 stimulation (i.e., CD163). Importantly, both, IL-10 and TGFβ induced the expression of

HLA-DR, CD38 and CD86 in the macrophages expressing the corresponding ChCRs (figure 2A and B). Moreover, IL-

10 blocked the upregulation of CD163 in IL-10 ChCR transduced macrophages, while IL-10 massively increased its

expression in the controls (figure 2A). Notably, CD163 is upregulated in response to IL-10 but not to TGFβ and therefore

we did not measure CD163 expression in TGFβ ChCR-expressing macrophages. Overall, we found that, ChCR-

expressing macrophages indeed show a phenotype similar to IFNγ stimulated macrophages upon stimulation with IL-10

or TGFβ.

ChCR-expressing macrophages have an IFNγ-like transcriptome

Next, we performed RNAseq to extend the characterization of ChCR-expressing macrophages. PCA showed that ChCR-

expressing macrophages stimulated with their corresponding ligand clustered together with IFNγ stimulated-

macrophages, indicating an IFNγ-like transcriptome profile induced by stimulation of the ChCRs (figure 4A). As

expected, differential gene expression analysis of ChCR-expressing macrophages showed the upregulation of many IFNγ

stimulated genes upon IL-10 or TGFβ stimulation when compared to mock transduced macrophages stimulated equally

(figure 3B). Similarly, gene set enrichment analysis (GSEA) of hallmark gene sets showed that IFNγ stimulated genes,

among other *CXCL9, CXCL10, and CIITA,* were significantly enriched in ChCR-expressing macrophages upon

stimulation (figure 3C). Moreover, GSEA of gene ontology biological processes (GO:BP) showed enrichment of genes

associated with antigen presentation (MHC-II HLA genes) and several immune related pathways (figure 3D). Thus, we

could confirm that ChCR-expressing macrophages have an IFNγ-like transcriptome.

ChCR-expressing macrophage upregulate genes associated with anti-PD-L1 response in TNBC patients

CXCL9 and CXCL10 have been associated with better response to immunotherapy and a hot TME in TNBC and other

tumors.[7 25 41-43] Interestingly, both *CXCL9* and *CXCL10* were upregulated in ChCR-expressing macrophages in RNAseq,

which we confirmed for CXCL10 by ELISA (figure 4A and B). Zhang et al have performed single cell RNAseq of

biopsies from TNBC patients treated with chemotherapy and anti-PD-L1 and have identified a TAM transcriptome

profile associated with anti-PD-L1 response.[25] Thus, we assessed the expression of other genes associated with good

response to anti-PD-L1 treatment in our ChCR-expressing macrophages upon stimulation. Indeed, 46 (IL-10 ChCR) and

55 (TGFβ ChCR) of 195 response associated genes were also significantly upregulated in ChCR-expressing macrophages

upon stimulation with the respective ligand in comparison to mock transduced macrophages (Supplementary table 2).

Moreover, the heatmap analyzing response associated genes identified two cluster that are separated depending to the

presence of IFNγ or ChCR-induced IFNγ-like signaling (figure 4C). Note that 14 (IL-10 ChCR) and 22 (TGFβ ChCR)

response associated genes were downregulated in samples with ChCR-induced IFNγ-like signaling. Responder genes that

were upregulated in ChCR-expressing macrophages upon stimulation belong for example to hallmark pathways of IFNγ

response and allograft rejection (i.e. *CD74, CXCL9, STAT1, SERPING1, B2M…*) and TNFα signaling via NF-κB (i.e.

TNF, SAT1…). In addition, we have performed GSEA of custom responder (R, upregulated in anti-PD-L1 responders)

and non-responder (NR, downregulated in anti-PD-L1 responders) gene sets. We found that the responder gene set was

indeed enriched in ChCR-expressing macrophages upon stimulation with the respective ligand (figure 4D). In contrast,

the non-responder gene set was not enriched in ChCR-expressing macrophages. Thus, the transcriptome of our

genetically-engineered macrophages could be beneficial for the response to anti-PD-L1 therapy.

ChCR-expressing macrophages have anti-tumoral activity

Inflammatory macrophages have long been known to have anti-tumoral effects.[23 44] Therefore, we explored whether

ChCR engineered macrophages have an anti-tumoral effect in a heterotypic TNBC spheroid model. This spheroid model

consisted of the TNBC cell line MDA-MB-231 and MRC-5 fibroblasts. The presence of fibroblasts was essential as

MDA-MB-231 cells alone did not form spheroids. One day after formation of MDA-MB-231 and MRC-5 spheroids, we

added the genetically-engineered macrophages. To monitor the viability of MDA-MB-231 cells, we engineered the cell

line to express both EGFP and luciferase. We used EGFP to measure MDA-MB-231 number by microscopy over time

and luciferase activity to determine the end-point cell number. Indeed, IL-10- or TGFβ ChCR-expressing macrophages

limited the expansion of MDA-MB-231 spheroid when added together with the corresponding cytokines, as assessed by

mean EGFP intensity (figure 5A-B). We observed similar results when measuring the remaining luciferase activity on

day 6, even though there was only a trend for TGFβ ChCR-expressing macrophages (figure 5C). Next, we were

wondering whether the anti-tumoral effect of ChCR-expressing macrophages was mediated by soluble factors or cell-

contact. We found that supernatant from stimulated ChCR-expressing macrophages alone led to reduced MDA-MB-231

and BT-549 cell number after three days incubation (online supplemental figure S2), indicating secreted factors, are, at

least in part, responsible for the anti-tumoral effect observed in the spheroid model.

DISCUSSION

The cold TME is highly suppressive and makes tumors non-responsive to ICI. The presence of inflammatory

macrophages in the TME is a favorable prognostic marker and increases also the responsiveness to ICIs.[7 25 42] and

therefore, TAMs or macrophages recruited to the TME represent a therapeutic target. Here, we developed ChCR-

expressing macrophages that are designed to switch to inflammatory macrophages in the presence of IL-10 and TGFβ in

the TME. We show that ChCR-expressing macrophages indeed have a phenotype similar to IFNγ-induced inflammatory

macrophages. Importantly, ChCR-expressing macrophages also have a direct anti-tumoral activity.

In some ways, the TME is a standalone area for the best tumor growth. Immunosuppressive cytokines are abundantly

present, among other IL-10 and TGFβ, in the TME. In TCGA RNAseq data, we confirmed that the transcripts of these

cytokines are present in TNBC. Therefore, we constructed ChCRs by combining the extracellular domains of the IL-10

and TGFβ receptors with the intracellular signaling domains of the IFNγ receptor to activate macrophages in the TME,

leveraging the local enrichment of IL-10 and TGFβ. Please, note that IFNγ is the crucial cytokine for inducing the

inflammatory activation of macrophages. To provide flexibility to the ChCRs, we added a glycine-serine linker between

the extracellular domains and the transmembrane domains of the ChCRs.[27] To ensure targeting the ChCR to the cell

surface, we used the signal peptides from the corresponding IFNγR subunit. The characterization in terms of surface

expression and signaling clearly showed that we have created functional ChCRs.

Our results suggest that ChCR-expressing macrophages become inflammatory in response to IL-10 or TGFβ. These

ChCR-expressing macrophages exhibit several attributes that would allow them to disrupt an immunosuppressive TME.

Firstly, RNAseq showed that ChCR-expressing macrophages upregulated the expression of the chemokines *CXCL9* and

CXCL10 upon IL-10 or TGFβ stimulation, which we confirmed for CXCL10 by ELISA. Both in TNBC and beyond,

there are several studies that underline the importance of CXCL9 and CXCL10 expression in the TME for the efficacy of

a checkpoint inhibitor treatment.[7 22 25 41-43] It is noteworthy that Sloas et al. recently reported that macrophages engineered

with an IL-10/IFNλ switch receptor, similar to our IL-10 ChCR, led to an increased T cell infiltration into the tumor *in*

vivo in a syngeneic tumor mouse model.[45 46] Secondly, HLA-DR and CD86 were also upregulated as determined by flow

cytometry. These surface proteins are crucial for T cell activation and antigen presentation. RNAseq analysis also

showed enrichment of pathways associated with antigen presentation. Therefore, we believe that ChCR-expressing

macrophages could lead to attraction, infiltration and activation of T cells and thereby initiating a positive feedback loop

that results in T cells secreting IFNγ, which in turn leads to TAMs adopting an inflammatory phenotype.[26] Ultimately, we

show that ChCR-expressing macrophages have a direct anti-tumoral effect when adding to a TNBC spheroid composed

of MDA-MB-231 cells and MRC-5 fibroblasts. The anti-tumoral effect is, at least partially, mediated by secreted factors,

as conditioned medium obtained from ChCR-expressing macrophages reduced the viability of the TNBC cells MDA-

MB-231 and BT-549. It is well known that macrophages exhibit anti-tumoral activity upon IFNγ and LPS stimulation.[23]

[47] Here we show that IFNγ-like signaling alone via the ChCR renders macrophages anti-tumorally active.

We noticed that spheroids co-cultured with IL-10 or TGFβ ChCR-expressing macrophages showed a reduced spheroid

dissemination compared to the controls (data not shown). It is highly speculative but this reduced dissemination may be

due to the anti-tumoral activity of the ChCR-expressing macrophages, while one of the pro-tumoral roles of TAMs is to

promote metastasis (reviewed[21]).

We observed differences in the performance of the two ChCR variants evaluated: the TGFβ ChCR responded to a lower

dose of TGFβ than the IL-10 ChCR to IL-10 as shown by the STAT1 activation in THP-1 and CXCL10 secretion in

primary macrophages. These differences could be due to distinct affinities to the corresponding cytokine or distinct

activation thresholds of the ChCRs. The difference in activation threshold might be dependent on the combination of

extracellular domain with the intracellular domain of the IFNGR2, which is the limiting subunit in inducing signaling.[48]

For instance, in the case of the IL-10 ChCR, we fused the low affinity IL-10Rb to IFNGR2 (reviewed[49]), whereas in the

case of TGFβ ChCR, we fused the high affinity TGFβR2 to the IFNGR2.[50] Importantly, in the complex setting of

spheroids, macrophages expressing IL-10 ChCR or TGFβ ChCR performed similar, when saturating concentrations were

used (figure 5). Ultimately, *in vivo* experiments will be needed to assess and compare activity and toxicity of both ChCR

variants.

As mentioned above, patients with advanced, PD-L1 positive TNBC benefit from adding pembrolizumab to

chemotherapy.[3] TAMs isolated from patients who responded to anti-PD-L1 treatment show a very unique transcriptomic

pattern.[25] The RNAseq data we obtained from ChCR-expressing macrophages in response to the corresponding

cytokines, showed upregulation of key genes associated with better response to PD-L1 ICI. Therefore, we have good

reasons to assume that PD-L1 ICI treatment in concert with adoptively transferred ChCR-expressing macrophages could

result in an additive or synergistic anti-tumoral activity.

The ChCR-expressing macrophages present several attributes that may render them anti-tumorally active. Our data,

however, have limitations. Our experiments, including the spheroid assay, do not fully recapitulate the complexity of a

tumor with its rich cell content, low pH, high levels of adenosine and hypoxia. Thus, we will further investigate the use

ChCR-expressing macrophages as an adoptive cell therapy for TNBC *in vivo* in a humanized mouse model.

DECLARATIONS

Ethics approval and consent to participate

Written consent for the use of buffy coats for research purposes was obtained from blood donors by the Blood Donation

Centre.

Consent for publication

Not applicable.

Competing interests

SB and RFS hold a patent associated to the ChCR (WO2024133517). SB received an initial startup coaching from the

Innosuisse and a convertible loan from Venture kick to develop the project about gene engineered macrophages to a

potential startup. The convertible loan was not used for the scientific work, presented in this manuscript. No potential

competing interests were declared by the other authors.

Funding

This project was funded by a grant from Gilead Sciences Switzerland Sàrl, UZH Postdoc grant, Innosuisse (103.487 IP-

LS), Claudia von Schilling Stiftung, Lotte und Adolf Hotz-Sprenger Stiftung, OPO Stiftung, Gebauer Stiftung, Novartis

Foundation for Medical Biological Research (#19B139), USZ Foundation, UZH Entrepreneur Fellowship, Arnold U. und

Susanne Huggenberger-Bischoff Stiftung, Peter Bockhoff Stiftung, and Mach-Gaensslen Stiftung.

Authors' contributions

Conceptualization: ST, RFS, SB

Chimeric cytokine receptor conceptualization and design: SB

Experiments: ST, FS, CB, VDV

Study supervision: RFS, SB

Manuscript writing: ST, RFS, SB

Funding acquisition: ST, CB, RFS, SB

Acknowledgments

We thank the anonymous volunteers for donating the buffy coats. Flow cytometry was performed with equipment of the
Flow Cytometry Facility, University of Zurich. We thank for the assistance in cell sorting experiments by the Flow
Cytometry Facility, University of Zurich. We thank the Functional Genomics Center Zurich (FGCZ) of University of
Zurich and ETH Zurich, and in particular Timothy Sykes, Hubert Rehrauer, and Peter Leary, for the support on RNA
sequencing analysis. We thank Jeremy Luban for providing us with pcDNA3.1 Vpx and p8.9NDSB-DPAVDLL
plasmids for virus production and Patrick Salmon for providing us with pCWX Dest backbone. We thank Patrick Turko
for his advice on analysis of TCGA data. We thank Sarah Brünigk for her advice with the RNAseq data analysis. The
results shown here are in part based upon data generated by the TCGA Research Network: https://www.cancer.gov/tcga.
ChatGPT was used to improve wording, grammar and spelling of this manuscript.
472

LITERATURE

1. Foulkes WD, Smith IE, Reis-Filho JS. Triple-negative breast cancer. *N Engl J Med* 2010;363(20):1938-48. doi: 10.1056/NEJMra1001389

2. Schmid P, Cortes J, Dent R, et al. Overall Survival with Pembrolizumab in Early-Stage Triple-Negative Breast Cancer. *N Engl J Med* 2024 doi: 10.1056/NEJMoa2409932 [published Online First: 20240915]

3. Cortes J, Rugo HS, Cescon DW, et al. Pembrolizumab plus Chemotherapy in Advanced Triple-Negative Breast Cancer. *N Engl J Med* 2022;387(3):217-26. doi: 10.1056/NEJMoa2202809

4. Cappell KM, Kochenderfer JN. Long-term outcomes following CAR T cell therapy: what we know so far. *Nat Rev Clin Oncol* 2023;20(6):359-71. doi: 10.1038/s41571-023-00754-1 [published Online First: 20230413]

5. Daei Sorkhabi A, Mohamed Khosroshahi L, Sarkesh A, et al. The current landscape of CAR T-cell therapy for solid tumors: Mechanisms, research progress, challenges, and counterstrategies. *Front Immunol* 2023;14:1113882. doi: 10.3389/fimmu.2023.1113882 [published Online First: 20230320]

6. Geng P, Chi Y, Yuan Y, et al. Novel chimeric antigen receptor T cell-based immunotherapy: a perspective for triple-negative breast cancer. *Front Cell Dev Biol* 2023;11:1158539. doi: 10.3389/fcell.2023.1158539 [published Online First: 20230629]

7. Hammerl D, Martens JWM, Timmermans M, et al. Spatial immunophenotypes predict response to anti-PD1 treatment and capture distinct paths of T cell evasion in triple negative breast cancer. *Nat Commun* 2021;12(1):5668. doi: 10.1038/s41467-021-25962-0 [published Online First: 2021/09/29]

8. Li F, Li C, Cai X, et al. The association between CD8+ tumor-infiltrating lymphocytes and the clinical outcome of cancer immunotherapy: A systematic review and meta-analysis. *EClinicalMedicine* 2021;41:101134. doi: 10.1016/j.eclinm.2021.101134 [published Online First: 20210916]

9. Gruosso T, Gigoux M, Manem VSK, et al. Spatially distinct tumor immune microenvironments stratify triple-negative breast cancers. *J Clin Invest* 2019;129(4):1785-800. doi: 10.1172/JCI96313 [published Online First: 2019/02/13]

10. Galon J, Bruni D. Approaches to treat immune hot, altered and cold tumours with combination immunotherapies. *Nat Rev Drug Discov* 2019;18(3):197-218. doi: 10.1038/s41573-018-0007-y

11. Park JH, Jonas SF, Bataillon G, et al. Prognostic value of tumor-infiltrating lymphocytes in patients with early-stage

triple-negative breast cancers (TNBC) who did not receive adjuvant chemotherapy. *Ann Oncol*

2019;30(12):1941-49. doi: 10.1093/annonc/mdz395

12. Nederlof I, Isaeva OI, de Graaf M, et al. Neoadjuvant nivolumab or nivolumab plus ipilimumab in early-stage triple-

negative breast cancer: a phase 2 adaptive trial. *Nat Med* 2024 doi: 10.1038/s41591-024-03249-3 [published

Online First: 20240916]

13. Keren L, Bosse M, Marquez D, et al. A Structured Tumor-Immune Microenvironment in Triple Negative Breast

Cancer Revealed by Multiplexed Ion Beam Imaging. *Cell* 2018;174(6):1373-87 e19. doi:

10.1016/j.cell.2018.08.039 [published Online First: 2018/09/08]

14. Franklin RA, Liao W, Sarkar A, et al. The cellular and molecular origin of tumor-associated macrophages. *Science*

2014;344(6186):921-5. doi: 10.1126/science.1252510 [published Online First: 20140508]

15. Combes AJ, Samad B, Tsui J, et al. Discovering dominant tumor immune archetypes in a pan-cancer census. *Cell*

2022;185(1):184-203 e19. doi: 10.1016/j.cell.2021.12.004 [published Online First: 20211227]

16. Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev Immunol* 2008;8(12):958-

69. doi: 10.1038/nri2448

17. Pittet MJ, Michielin O, Migliorini D. Clinical relevance of tumour-associated macrophages. *Nat Rev Clin Oncol*

2022;19(6):402-21. doi: 10.1038/s41571-022-00620-6 [published Online First: 20220330]

18. Toney NJ, Opdenaker LM, Cicek K, et al. Tumor-B-cell interactions promote isotype switching to an

immunosuppressive IgG4 antibody response through upregulation of IL-10 in triple negative breast cancers. *J

Transl Med* 2022;20(1):112. doi: 10.1186/s12967-022-03319-5 [published Online First: 20220307]

19. Chen Z, Wu J, Wang L, et al. Tumor-associated macrophages of the M1/M2 phenotype are involved in the regulation

of malignant biological behavior of breast cancer cells through the EMT pathway. *Med Oncol* 2022;39(5):83.

doi: 10.1007/s12032-022-01670-7 [published Online First: 20220516]

20. Ding MJ, Su KE, Cui GZ, et al. Association between transforming growth factor-beta1 expression and the clinical

features of triple negative breast cancer. *Oncol Lett* 2016;11(6):4040-44. doi: 10.3892/ol.2016.4497 [published

Online First: 20160426]

21. Qian BZ, Pollard JW. Macrophage diversity enhances tumor progression and metastasis. *Cell* 2010;141(1):39-51.
doi: 10.1016/j.cell.2010.03.014

22. Bill R, Wirapati P, Messemaker M, et al. CXCL9:SPP1 macrophage polarity identifies a network of cellular
programs that control human cancers. *Science* 2023;381(6657):515-24. doi: 10.1126/science.ade2292 [published
Online First: 20230803]

23. Pace JL, Russell SW, Schreiber RD, et al. Macrophage activation: priming activity from a T-cell hybridoma is
attributable to interferon-gamma. *Proc Natl Acad Sci U S A* 1983;80(12):3782-6. doi: 10.1073/pnas.80.12.3782

24. Mantovani A, Allavena P, Marchesi F, Garlanda C. Macrophages as tools and targets in cancer therapy. *Nature
Reviews Drug Discovery* 2022;21(11):799-820. doi: 10.1038/s41573-022-00520-5

25. Zhang Y, Chen H, Mo H, et al. Single-cell analyses reveal key immune cell subsets associated with response to PD-
L1 blockade in triple-negative breast cancer. *Cancer Cell* 2021;39(12):1578-93 e8. doi:
10.1016/j.ccell.2021.09.010 [published Online First: 20211014]

26. van Elsas MJ, Middelburg J, Labrie C, et al. Immunotherapy-activated T cells recruit and skewlate-stage activated
M1-like macrophagesthat are critical for therapeutic efficacy. *Cancer Cell* 2024 doi:
10.1016/j.ccell.2024.04.011 [published Online First: 20240509]

27. Chen X, Zaro JL, Shen WC. Fusion protein linkers: property, design and functionality. *Adv Drug Deliv Rev*
2013;65(10):1357-69. doi: 10.1016/j.addr.2012.09.039 [published Online First: 20120929]

28. Hallgren J, Tsirigos K, Pedersen MD, et al. DeepTMHMM predicts alpha and beta transmembrane proteins using
deep neural networks: bioRxiv, 2022.

29. Pertel T, Reinhard C, Luban J. Vpx rescues HIV-1 transduction of dendritic cells from the antiviral state established
by type 1 interferon. *Retrovirology* 2011;8(49) doi: https://doi.org/10.1186/1742-4690-8-49

30. Lehmann BD, Colaprico A, Silva TC, et al. Multi-omics analysis identifies therapeutic vulnerabilities in triple-
negative breast cancer. *Nat Commun* 2021;12(1):6276. doi: 10.1038/s41467-021-26502-6 [published
Online First: 20211101]

31. Mounir M, Lucchetta M, Silva TC, et al. New functionalities in the TCGAbiolinks package for the study and
integration of cancer data from GDC and GTEx. *PLoS Comput Biol* 2019;15(3):e1006701. doi:
10.1371/journal.pcbi.1006701 [published Online First: 20190305]

32. Hatakeyama M, Opitz L, Russo G, et al. SUSHI: an exquisite recipe for fully documented, reproducible and reusable

NGS data analysis. *BMC Bioinformatics* 2016;17(1):228. doi: 10.1186/s12859-016-1104-8 [published Online

First: 20160602]

33. Andrews S. FastQC: A Quality Control Tool for High Throughput Sequence Data 2010 [Available from:

http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

34. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* 2018;34(17):i884-

i90. doi: 10.1093/bioinformatics/bty560

35. Dobin A, Davis CA, Schlesinger F, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 2013;29(1):15-

21. doi: 10.1093/bioinformatics/bts635 [published Online First: 20121025]

36. Frankish A, Diekhans M, Jungreis I, et al. Gencode 2021. *Nucleic Acids Res* 2021;49(D1):D916-D23. doi:

10.1093/nar/gkaa1087

37. Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read mapping by seed-and-vote. *Nucleic*

Acids Res 2013;41(10):e108. doi: 10.1093/nar/gkt214 [published Online First: 20130404]

38. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.

Genome Biol 2014;15(12):550. doi: 10.1186/s13059-014-0550-8

39. Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological themes among gene

clusters. *OMICS* 2012;16(5):284-7. doi: 10.1089/omi.2011.0118 [published Online First: 20120328]

40. Liberzon A, Birger C, Thorvaldsdottir H, et al. The Molecular Signatures Database (MSigDB) hallmark gene set

collection. *Cell Syst* 2015;1(6):417-25. doi: 10.1016/j.cels.2015.12.004

41. House IG, Savas P, Lai J, et al. Macrophage-Derived CXCL9 and CXCL10 Are Required for Antitumor Immune

Responses Following Immune Checkpoint Blockade. *Clin Cancer Res* 2020;26(2):487-504. doi: 10.1158/1078-

0432.CCR-19-1868 [published Online First: 20191021]

42. Schlenker R, Schwalie PC, Dettling S, et al. Myeloid-T cell interplay and cell state transitions associated with

checkpoint inhibitor response in melanoma. *Med* 2024 doi: 10.1016/j.medj.2024.03.015 [published Online First:

20240404]

43. Litchfield K, Reading JL, Puttick C, et al. Meta-analysis of tumor- and T cell-intrinsic mechanisms of sensitization to

checkpoint inhibition. *Cell* 2021;184(3):596-614 e14. doi: 10.1016/j.cell.2021.01.002 [published Online First:

20210127]

44. Cameron DJ, Churchill WH. Cytotoxicity of human macrophages for tumor cells: enhancement by bacterial

lipopolysaccharides (LPS). *The Journal of Immunology* 1980;124(2):708-12. doi: 10.4049/jimmunol.124.2.708

45. Sloas C, Beghi S, Huangfu Y, et al. Abstract 5249: Macrophages expressing synthetic cytokine receptors reverse

IL10-mediated immunosuppression within solid tumors and promote adaptive immunity. *Cancer Research*

2024;84(6_Supplement):5249-49. doi: 10.1158/1538-7445.Am2024-5249

46. Sloas C, Beghi S, Huangfu Y, et al. Macrophages expressing synthetic cytokine receptors reverse IL10-mediated

Immunosuppression   within   solid   tumors   and   promote   adaptive   immunity

https://74dfcf.p3cdn1.secureserver.net/wp-content/uploads/2024/04/AACR_2024_EM-C_FINAL.pdf2024

[accessed 01.10.2024.

47. Muller E, Christopoulos PF, Halder S, et al. Toll-Like Receptor Ligands and Interferon-gamma Synergize for

Induction of Antitumor M1 Macrophages. *Front Immunol* 2017;8:1383. doi: 10.3389/fimmu.2017.01383

[published Online First: 20171026]

48. Bernabei P, Coccia EM, Rigamonti L, et al. Interferon-γ receptor 2 expression as the deciding factor in human T, B,

and myeloid cell proliferation or death. *Journal of Leukocyte Biology* 2001;70(6):950-60. doi:

10.1189/jlb.70.6.950

49. Walter MR. The molecular basis of IL-10 function: from receptor structure to the onset of signaling. *Curr Top*

Microbiol Immunol 2014;380:191-212. doi: 10.1007/978-3-662-43492-5_9

50. Wrana JL, Attisano L, Carcamo J, et al. TGF beta signals through a heteromeric protein kinase receptor complex.

Cell 1992;71(6):1003-14. doi: 10.1016/0092-8674(92)90395-s

FIGURE LEGENDS

Figure 1 IL-10 / TGFβ stimulation induces STAT1 signaling in chimeric cytokine receptors (ChCR) expressing

THP-1 and primary macrophages A) Expression of *IL10* and *TGFB1* in TNBC. Shown is the percentile expression of

IL10 and *TGFB1* expression across all transcripts in TNBC samples from TCGA database. Each dot represents one

patient. n=195 TNBC patients. B) Scheme of lentiviral constructs that were used. C) Scheme of ChCR. Binding of IL-10

or TGFβ to the respective ChCR induces pSTAT1 signaling. C/D) THP-1 were either left untransduced or transduced

with mock, IL-10 ChCR or TGFβ ChCR construct. C) Expression of IL-10Rα, IL-10Rβ and TGFβR2 was assessed by

flow cytometry. Shown are representative histograms. Cells are gated on tNGFR+ cells. D) THP-1 were either left

untreated, stimulated with 1, 10 or 100ng/ml IL-10 or TGFβ, or 5ng/ml IFNγ for 30min. pSTAT1 signaling was assessed

by Phosphoflow. Shown are median fluorescence intensities (MFI's, mean ± SD) in single cells for untransduced or in

tNGFR+ cells for mock, IL-10 ChCR or TGFβ ChCR. N=2-3 E/F) Primary human monocytes were either transduced

with mock, IL-10 ChCR or TGFβ ChCR construct. E) Transduction efficiency (tNGFR+) was assessed by flow

cytometry upon differentiation to macrophages. Shown are % tNGFR+ cells (mean ± SD). N=14-33 F). Expression of IL-

10Rα, IL-10Rβ and TGFβR2 was assessed by flow cytometry. Shown are MFIs gated on tNGFR+ cells (mean ± SD)

from 4-5 healthy donors from 3-4 independent experiments G) Transduced macrophages were either left untreated or

stimulated as indicated for 30min. pSTAT1 signaling was assessed by Phosphoflow. Shown are pSTAT1 MFIs in

tNGFR+ cells (mean ± SD) from 4 donors in 4 independent experiments (IL-10 ChCR) or 5 donors in 3 independent

experiments (TGFβ ChCR). p-values were calculated by ratio paired t-tests. * p ≤ 0.05, ** p≤0.01, *** p≤0.001, ****

p≤0.00001

Figure 2 ChCR-expressing macrophages have an inflammatory phenotype upon IL-10 or TGFβ stimulation A/B)

Primary human monocytes were either left untransduced or transduced with mock, IL-10 ChCR or TGFβ ChCR

construct. After differentiation to macrophages, cells were either left unstimulated or stimulated with 100ng/ml IL-10, 10

or 100ng/ml TGFβ, or 10ng/ml IFNγ. Expression of surface marker was assessed by flow cytometry. A) Shown are

median fluorescence intensities (MFIs, mean ± SD) after subtracting isotype MFI or % CD38+ or % CD163+ gated on

CD11b+ cells from 3-4 healthy donors in 4 independent experiments. B) Shown are MFI (mean ± SD) after subtracting

isotype MFI or % CD38+ gated on CD11b+ (untransduced) or NGFR+ (transduced) cells from 3 healthy donors in 2

independent experiments. For MFIs, p-values were calculated using ratio paired t-tests. For % positive cells, p-values

were calculated using paired t-tests. * p ≤ 0.05, ** p≤0.01, *** p≤0.001, **** p≤0.00001

Figure 3 ChCR-expressing macrophages have an IFNγ-like transcriptome A-D) Monocytes were either transduced

with mock, IL-10 ChCR or TGFβ ChCR lentivirus. After differentiation to macrophages, cells were either left

unstimulated or stimulated with 100ng/ml IL-10, 100ng/ml TGFβ or 10ng/ml IFNγ for 2 days. mRNA was isolated and

subjected to bulk RNA sequencing. A) Gene expression principal component analysis (PCA), n=3 donors B) Differential

gene expression analysis of mock transduced versus IL-10 ChCR transduced macrophages upon IL-10 stimulation and

mock transduced versus TGFβ ChCR transduced macrophages upon TGFβ stimulation. P-values were calculated using

the Wald test and adjusted for multiple comparison using the Benjamini-Hochberg method. Significantly down- or

upregulated genes were defined as log2 fold change (log2 FC) $\leq$-1 or $\geq$1 and FDR < 0.05. Genes highlighted in red are

part of the hallmark IFNγ pathway. n=3 donors. C) Gene set enrichment analysis (GSEA) of hallmark gene sets. Shown

is the normalized enrichment score (NES) of the top 5 enriched hallmark gene sets. D) GSEA of Gene ontology

biological processes (GO:BP) gene set. Shown is the NES of top 10 enriched BP.

Figure 4 ChCR-expressing macrophages upregulate genes associated with good response to anti-PDL-1 treatment

A-D) Monocytes were either transduced with mock, IL-10 ChCR or TGFβ ChCR lentivirus. After differentiation to

macrophages, cells were either left unstimulated or stimulated with 100ng/ml IL-10, 100ng/ml TGFβ or 10ng/ml IFNγ

for 2 days. A) mRNA was isolated and subjected to bulk RNA sequencing. *CXCL9* and *CXCL10* expression analyzed by

bulk RNAseq. Shown are normalized log2 counts (mean ± SD). N=3 donors. P-values were calculated using the Wald

test and adjusted for multiple comparison using the Benjamini-Hochberg method. * False discovery rate (FDR) $\leq$ 0.05,

** FDR$\leq$0.01, *** FDR$\leq$0.001, **** FDR$\leq$0.00001 B) Supernatants were collected and analyzed by CXCL10 ELISA.

Shown are CXCL10 concentration (mean ± SD). N=4 donors analyzed in 2-3 independent experiments. p-values were

calculated using paired t-tests, comparing ChCRs with mock condition. * p $\leq$ 0.05, ** p$\leq$0.01, *** p$\leq$0.001, ****

p$\leq$0.00001 C-D) mRNA was isolated and subjected to bulk RNA sequencing. C) Heatmap of genes enriched in response

associated with response to anti-PD-L1 treatment in TNBC patients.[25] Expression in genetically-engineered macrophages

is shown. D) Gene set enrichment analysis of genes expressed in macrophages and associated with response (R) and no

response (NR) to anti-PD-L1 treatment in TNBC patients.[25] P-values were adjusted for multiple comparison using the

Benjamini Hochberg method.

Figure 5 ChCR-expressing macrophages have anti-tumoral activity A-C) Monocytes were either left untransduced or

transduced with mock, IL-10 or TGFβ ChCR and were differentiated to macrophages. Spheroids were formed by mixing

1000 MDA-MB-231 TNBC cells (expressing EGFP and luciferase) and 1500 MRC-5 fibroblasts. 1 day later, genetically-

engineered macrophages were added at a 1:10 ratio (MDA-MB-231: macrophages). Spheroids were either left

unstimulated, stimulated with IL-10 (100ng/ml), TGFβ (10ng/ml), or IFNγ (10ng/ml). A) Quantification of tumor cells in

spheroids by measuring mean EGFP intensity (mean ± SD) over time. Shown are results from 3 donors from 2

independent experiments. B) Quantification of tumor cells in spheroids by measuring mean EGFP intensity (mean ± SD)

6d after adding macrophages. Shown are results from 3 donors from 2 independent experiments. C) Quantification of

tumor cells in spheroid by measuring luciferase activity 6d after adding macrophages. Shown are relative light units

(RLU) (mean ± SD) after subtraction of background RLU from 3 donors analyzed in 2 independent experiments. p-

values were calculated using paired t-tests comparing the ChCRs with mock condition. * $p \leq 0.05$, ** $p \leq 0.01$, ***

$p \leq 0.001$, **** $p \leq 0.00001$

Figure 1

Figure 2

Figure 3

A

Construct
- mock
- IL-10 ChCR
- TGFβ ChCR

Stimulation
- unstim.
- IL-10
- TGFβ
- IFNγ

PC2: 12% variance

PC1: 58% variance

B

Mock vs. IL-10 ChCR
upon IL-10 stimulation

- IFNγ hallmark genes
- significantly up- or downregulated
- not significant

−log10 P-value

log2 FC

ITGB7, SERPING1, HLA−DQA1, LGALS3BP, VAMP5, STAT1, CD38, GBP4, CXCL9, CIITA, IDO1, CXCL10

Mock vs. TGFβ ChCR
upon TGFβ stimulation

- IFNγ hallmark genes
- significantly up- or downregulated
- not significant

−log10 P-value

log2 FC

ITGB7, SECTM1, GBP4, SERPING1, SOCS3, VAMP5, HLA−DQA1, CFB, TAP1, RSAD2, CXCL9, C1S, IDO1, ISG20, CD38, CXCL10

C

IL-10 ChCR Hallmark

- INTERFERON_GAMMA_RESPONSE
- INTERFERON_ALPHA_RESPONSE
- IL6_JAK_STAT3_SIGNALING
- INFLAMMATORY_RESPONSE
- ALLOGRAFT_REJECTION

NES
0.0 0.5 1.0 1.5 2.0 2.5 3.0

TGFβ ChCR Hallmark

- INTERFERON_GAMMA_RESPONSE
- INTERFERON_ALPHA_RESPONSE
- INFLAMMATORY_RESPONSE
- TNFA_SIGNALING_VIA_NFKB
- IL6_JAK_STAT3_SIGNALING

NES
0.0 0.5 1.0 1.5 2.0 2.5 3.0

D

IL10 ChCR GO:BP

- defense response to virus
- antigen processing and presentation of exogenous peptide antigen via MHC class II
- antigen processing and presentation of peptide or polysaccharide antigen via MHC class II
- negative regulation of viral genome replication
- antigen processing and presentation
- immune system process
- cellular response to type II interferon
- peptide antigen assembly with MHC class II protein complex
- response to type II interferon
- defense response

NES
0.0 0.5 1.0 1.5 2.0 2.5

TGFβ ChCR GO:BP

- cellular response to type II interferon
- defense response to virus
- immune system process
- adaptive immune response
- response to virus
- neutrophil chemotaxis
- response to bacterium
- cellular response to interleukin−1
- antigen processing and presentation
- immune response

NES
0.0 0.5 1.0 1.5 2.0 2.5

Figure 4

Figure 5