Non-equilibrium strategies for ligand specificity in signaling networks

Andrew Goetz[1]*, Jeremy Barrios[2], Ralitsa Madsen[3], Purushottam Dixit[1,4]*

*For correspondence:
andrew.goetz@yale.edu (AG);
purushottam.dixit@yale.edu (PD)

[1]Department of Biomedical Engineering, Yale University, New Haven, CT; [2]Department of Physics, Yale University, New Haven, CT; [3]MRC Protein Phosphorylation and Ubiquitylation Unit, University of Dundee, Dundee, DD1 5EH; [4]Systems Biology Institute, Yale University, West Haven, CT

Abstract Signaling networks often encounter multiple ligands and must respond selectively to generate appropriate, context-specific outcomes. At thermal equilibrium, ligand specificity is limited by the relative affinities of ligands for their receptors. Here, we present a non-equilibrium model showing how signaling networks can overcome thermodynamic constraints to preferentially signal from specific ligands while suppressing others. In our model, ligand-bound receptors undergo sequential phosphorylation, with progression restarted by ligand unbinding or receptor degradation. High-affinity complexes are *kinetically* sorted toward degradation-prone states, while low-affinity complexes are sorted towards inactivated states, both limiting signaling. As a result, network activity is maximized for ligands with intermediate affinities. This mechanism explains paradoxical experimental observations in receptor tyrosine kinase (RTK) signaling, including non-monotonic relationships between ligand affinity, kinase activity, and signaling output. Given the ubiquity of multi-site phosphorylation and ligand-induced degradation across signaling pathways, we propose that *kinetic sorting* provides a general non-equilibrium strategy for ligand discrimination in cellular networks.

Introduction

Cells routinely encounter a wide variety of extracellular ligands and decode their identity with remarkable precision to generate context-specific responses. This selective processing of environmental cues is essential for regulating diverse biological processes, including development, immune surveillance, and tissue homeostasis (*Cantley et al., 2014*). Failures in ligand discrimination underlie many diseases, including diabetes and cancer (*Madsen et al., 2025*; *Madsen and Vanhaesebroeck, 2020*).

A key determinant of ligand specificity in biochemical networks is the thermodynamic stability of molecular complexes, such as ligand–receptor or substrate–enzyme pairs. At thermal equilibrium, the abundance of complexes is determined by their equilibrium binding constants. This imposes a fundamental limit on specificity: high-affinity ligands are inevitably favored over lower-affinity competitors, with complex abundances scaling in proportion to their association constants.

However, many signaling networks display paradoxical behaviors that cannot be explained by equilibrium affinity alone (*Clark et al., 1999*; *Coombs et al., 2002*; *Freed et al., 2017*; *Madsen et al., 2025*; *Myers et al., 2023*). For example, receptor tyrosine kinases (RTKs) can produce stronger downstream signaling outputs in response to intermediate-affinity ligands than to high-affinity ligands (*Freed et al., 2017*; *Madsen et al., 2025*; *Myers et al., 2023*). Additionally, partially inhibiting kinase activity can paradoxically increase receptor phosphorylation levels (*Kiyatkin et al., 2020*; *Kleiman et al., 2011*). These observations raise a fundamental question: how do cells overcome

thermodynamic constraints to achieve nuanced, ligand-specific responses?

A classic solution to bypass equilibrium limits is kinetic proofreading (KPR), a mechanism first proposed by Hopfield (*Hopfield, 1974*) and Ninio (*Ninio, 1975*). KPR enhances specificity of high affinity ligands by introducing energy-consuming, irreversible steps—such as phosphorylation—that amplify differences between competing ligands. KPR has been invoked in diverse systems, including DNA replication (*Hopfield, 1980*), mRNA surveillance (*Hilleren and Parker, 1999*), protein folding (*Gulukota and Wolynes, 1994*), and immune receptor signaling (*McKeithan, 1995*; *Huang et al., 2019*; *Lever et al., 2014*). Yet, most KPR models (with a few exceptions (*Lever et al., 2014*; *Murugan et al., 2014*), see below) assume that correct recognition correlates with the highest affinity - a premise that fails in systems where low affinity ligands dominate signaling outputs compared to high affinity ligands.

In this study, we investigate how signaling networks achieve ligand specificity through non-equilibrium mechanisms that go beyond classical KPR. We develop a simple, biologically grounded model combining two ubiquitous features of cellular signaling: sequential multi-site phosphorylation and ligand-induced receptor degradation. These two motifs are found in many major receptor systems, including RTKs (*Furdui et al., 2006*; *Sorkin and Goh, 2009*), G protein-coupled receptors (GPCRs)(*Koenig and Edwardson, 1997*; *Tobin, 2008*), T cell receptors(*McKeithan, 1995*; *Charpentier and King, 2021*), and interleukin receptors (*Kollewe et al., 2004*; *Cendrowski et al., 2016*).

Our model shows that high-affinity ligand-receptor complexes are preferentially sorted toward degradation-prone states, while low-affinity complexes repeatedly dissociate the ligand, resulting in maximal signaling output only from intermediate-affinity ligands. Notably, this ligand specificity can be tuned by tuning easily controllable cellular parameters, e.g. enzyme abundances. This non-equilibrium *kinetic sorting* mechanism explains the paradoxical non-monotonic dependence of signaling activity on ligand affinity and phosphorylation rate observed in RTKs. More broadly, we propose that kinetic sorting provides a general strategy for achieving ligand discrimination across diverse natural and synthetic signaling networks.

Results

Figure 1. Reaction scheme of kinetic proofreading models. Chemical species and rate constants are shown in the figure. R denote ligand-free receptors, B denote ligand-bound inactive receptors, and $P_n, n \in [1, N]$ are phosphorylated receptors. The ultimate phosphorylated species P_N (marked red) is assumed to be signaling competent. (a) shows the traditional model first proposed by McKeithan (*McKeithan, 1995*). (b, c) show the sustained signaling model and the limited signaling model (*Lever et al., 2014*) which introduce additional receptor states, P_N^0 and I respectively, directly following receptor activation.

Figure 2. Ligand discrimination in kinetic proofreading models. (a) Activity P_N plotted as a function of non-dimensional ligand dissociation rate δ for the traditional KPR scheme (Fig. 1a). (b) Activity P_N plotted as a function of non-dimensional ligand dissociation rate δ for the limited signaling model (Fig. 1b). (c) The dependence of the activity on the dimensionless phosphorylation rate ω for the limited signaling model. All figures plotted for a sequence of $N = 1, 5,$ and 10 phosphorylation sites.

Classic kinetic proofreading always favors high-affinity ligands

Kinetic proofreading (KPR) is the *standard model* for non-equilibrium ligand discrimination. To set the stage, we first revisited the classic KPR model originally proposed by McKeithan to explain how T cell receptors avoid activation downstream of weak ligands (*McKeithan, 1995*) (Fig. 1a; see Supplementary Materials for equations).

In this model, ligand-bound receptors undergo a series of phosphorylation steps, with the final state P_N representing the active, signaling-competent form. Importantly, ligand unbinding at any phosphorylation stage returns the receptor to the unbound state R. We parameterized the model using dimensionless quantities: the ligand dissociation rate $\delta = k_d\tau$, phosphorylation rate $\omega = k_p\tau$, and ligand concentration $u = L/K_D$, where $K_D = k_d/k_{on}$. Assuming saturating ligand ($u \to \infty$), the steady-state abundance of the active state is:

$$P_N = \frac{\delta^N}{(\omega + \delta)^N}. \tag{1}$$

As expected, increasing the phosphorylation cascade length N amplifies the preference for low-dissociation (high-affinity) ligands (Fig. 2a), reflecting the classical KPR outcome.

Modified KPR schemes cannot explain paradoxical RTK behavior

Next, we examined two previously proposed extensions of KPR: the sustained signaling model and the limited signaling model (*Lever et al., 2014*) (Fig. 1b,c). Both models introduce an additional state to Mckeithan's KPR scheme. The sustained signaling model adds an active but ligand-free state P_N^0, while the limited signaling model introduces an inactivated state I downstream of P_N.

Of these, only the limited signaling model exhibits non-monotonic dependence on ligand dissociation rates at saturating ligand concentrations (*Lever et al., 2014*) (Fig. 2b), consistent with some paradoxical features observed in RTKs (*Freed et al., 2017*; *Madsen et al., 2025*; *Myers et al., 2023*). However, it fails to reproduce a second key observation: phosphorylation in this model increases monotonically with kinase activity, whereas RTK experiments show that partial kinase inhibition can paradoxically increase phosphorylation (*Kiyatkin et al., 2020*; *Kleiman et al., 2011*) (Fig. 2c). Thus, these models are insufficient to explain RTK signaling dynamics.

A kinetic sorting model integrates ligand-induced degradation

To address these gaps, we present a model incorporating two widespread signaling motifs: sequential multi-site phosphorylation and ligand-induced receptor degradation (Fig. **??**). In our model, receptors are delivered to the surface at a constant rate, internalized at a basal rate k_{int}, and degraded more rapidly when highly phosphorylated ($k_{int}^* > k_{int}$). Ligand-bound receptors undergo reversible phosphorylation. A key feature of our model is that *all* phosphorylated species are signaling competent. Indeed, in many signaling pathways all phosphorylation sites on the receptor (*Schulze et al.,*

Figure 3. Kinetic sorting of receptor species. Abundances of networks species B (ligand bound inactive receptor) and $P_n, n \in [1, 5]$ for a signaling receptor with $N = 5$ phosphorylation sites. Abundances are shown for ligands of three different affinities. The inset shows the activity of the first phosphorylation site A_1. Species abundances below 10^{-3} are not shown.

---

[102] *2005*; *Tobin, 2008*; *Kollewe et al., 2004*; *Lemmon and Schlessinger, 2010*; *Latorraca et al., 2020*)
[103] have downstream effects. Therefore, we define the net activity A_n of phosphorylation site n as all
[104] receptor states where the site n is phosphorylated: $A_n = \sum_{m \geq n} P_m$.

[105] **Parameter ranges.** To ensure that the phenomena captured by our model are relevant to real
[106] signaling networks, we selected ranges for the dimensionless parameters based on direct experi-
[107] mental measurements and model fits. Importantly, many of these kinetic processes have compa-
[108] rable rates across diverse receptor systems (*Koenig and Edwardson, 1997*; *Subtil et al., 1994*; *Liu*
[109] *et al., 2000*). Specifically, basal receptor internalization occurs at rates of $k_{\mathrm{int}} \approx 10^{-4}$–$10^{-3}, \mathrm{s}^{-1}$(*Wiley,*
[110] *2003*), while ligand-induced internalization is faster, at $k_{\mathrm{int}}^* \approx 10^{-3}$–$10^{-2}, \mathrm{s}^{-1}$(*Wiley, 2003*; *Lyashenko*
[111] *et al., 2020*). Ligand dissociation rates typically fall in the range $k_{\mathrm{d}} \approx 10^{-2}$–$10^{-1}, \mathrm{s}^{-1}$(*Chen et al.,*
[112] *2009*; *Lyashenko et al., 2020*), and receptor phosphorylation and dephosphorylation occur at $\sim$
[113] 10^{-1}–$10^{0}, \mathrm{s}^{-1}$(*Kleiman et al., 2011*; *Chen et al., 2009*; *Lyashenko et al., 2020*). For EGFR, equilibrium
[114] dissociation constants range from $\sim 0.1, \mathrm{nM}$ for the high-affinity ligand Betacellulin to $\sim 25, \mathrm{nM}$ for
[115] the low-affinity ligand AREG (*Hu et al., 2022*; *Macdonald-Obermann and Pike, 2014*). Based on these
[116] values, we set the following ranges for our dimensionless parameters: $\beta \in [1, 100]$, $\rho \in [0.01, 100]$,
[117] $\omega \in [1, 1000]$, and $\delta \in [1, 1000]$. Finally, the number of phosphorylation sites with known func-
[118] tional roles typically ranges from 5 to 25 (*Schulze et al., 2005*). These broad ranges comfortably
[119] encompass experimentally measured estimates. Unless otherwise specified, our default parame-
[120] ter values are $\delta = 20$, $\omega = 200$, $\rho = 0.01$, $\beta = 50$, and $N = 10$.

[121] Before examining how phosphorylation levels depend on model parameters, we illustrate the
[122] mechanism of kinetic sorting of receptor states, which tunes ligand specificity beyond pure thermo-
[123] dynamic preference, using a simple example. To that end, we consider a signaling network with
[124] $N = 5$ phosphorylation sites interacting with three ligands of distinct affinities—high, medium,
[125] and low. We assume the dissociation rates for these ligands are $\delta_H = 20$, $\delta_M = 200$, and $\delta_L = 1000$,
[126] respectively. In order to compare our model with aforementioned paradoxical experimental ob-
[127] servations which have been performed at saturating ligand concentration, we take the limit $u \to \infty$.

[128] Fig. 3 shows that low affinity ligands ($\delta_L = 1000$) predominantly sort receptors towards the in-
[129] active state B and early phosphorylation states $P_n, n \sim 1$ as frequent ligand unbinding prevents
[130] progression to later phosphorylation states. This behavior resembles the traditional KPR mecha-
[131] nism described by McKeithan (*McKeithan, 1995*). In contrast, receptors bound to high affinity lig-
[132] ands are sorted toward later phosphorylation states, which mark them for enhanced degradation.

Figure 4. Kinetic sorting model predicts ligand specificity. (a) The activity A_n of the n^{th} phosphorylation site as a function of dimensionless dissociation rate δ. The activity is normalized to the maximum activity. The maximum A_n as a function of n is shown in the inset. (b) Activity of the first phosphorylation site A_1 plotted as a function of the dissociation rate δ for different values of the phosphorylation rate ω. (c, d) Activity of the first phosphorylation site A_1 plotted as a function of phosphorylation rate ω (dephosphorylation rate ρ in panel d) for different values of the dissociation rate δ.

Here, similar to traditional KPR, the fraction of receptors reaching the final phosphorylation state is highest. Yet, the overall receptor pool is reduced due to ligand-induced degradation, lowering net phosphorylation activity. Strikingly, receptors bound to intermediate affinity ligands ($\delta_M = 200$) are sorted towards intermediate phosphorylation states, resulting in maximal phosphorylation output. Below, we kinetic parameters govern the ability of the network to overcome thermodynamic preference and acquire ligand specificity.

Early phosphorylation sites show ligand-specificity. Figure 4a illustrates how total phosphorylation activity at each site, $A_n, n \in [1, N]$ varies with ligand dissociation rate δ. Notably, early phosphorylation sites ($n \sim 1$) exhibit maximal activity at intermediate values of δ while both high- and low-affinity ligands suppress net receptor phosphorylation. Our model predicts that this ligand-specificity diminishes for later sites, where outputs increasingly resemble traditional KPR which favors high-affinity ligands.

To examine how model parameters shape ligand specificity, we focused on the activity at the first phosphorylation site, A_1, which exhibits the strongest discriminatory behavior (Fig.4a). As shown in Fig.4b, achieving ligand specificity at high dissociation rates δ requires sufficiently high phosphorylation rates ω. Notably, our model captures a puzzling observation from EGFR signaling: the high-affinity ligand EGF produces lower/comparable steady-state phosphorylation compared to lower-affinity ligands such as Epigen and Epiregulin (*Freed et al., 2017*; *Myers et al., 2023*; *Madsen et al., 2025*). Experimental estimates place the basal EGFR internalization rate at $k_{\text{int}} \approx 1.3 \times 10^{-3}, \text{s}^{-1}$ (*Chen et al., 2009*), the EGF dissociation rate at $k_{\text{d}} \approx 3 \times 10^{-2}, \text{s}^{-1}$ (*Chen et al., 2009*), and the phosphorylation rate at $k_{\text{p}} \approx 10^{-1} - 10^{0}, \text{s}^{-1}$, yielding $\delta_{\text{EGF}} \approx 10\text{–}20$ and $\omega_{\text{EGFR}} \approx 100\text{–}1000$.

Low-affinity ligands such as Epigen (EPGN) and Epiregulin (EREG) have equilibrium dissociation constants about 10-fold higher than EGF (*Hu et al., 2022*), corresponding to $\delta_{\text{EPGN}} \approx \delta_{\text{EREG}} \approx 100\text{–}200$. The effective degradation rate of fully activated receptors is estimated to be 10–50 times higher than that of inactive receptors (*Lyashenko et al., 2020*), implying $\beta = 50$. Under these conditions, our model predicts a switch in phosphorylation levels: as δ increases from δ_{EGF} to δ_{EPGN}, receptor phosphorylation increases—reversing the expectation based purely on thermodynamic affinity. This effect arises because EGF-bound receptors are efficiently sorted towards degradation-prone states compared to those bound to lower-affinity ligands.

Our model also explains another paradox in EGFR signaling. Experimental studies have shown that EGF-stimulated receptors exhibit higher steady-state phosphorylation when kinase activity is partially inhibited (*Kiyatkin et al., 2020*; *Kleiman et al., 2011*). As shown in Fig.4c, at low δ values (e.g., $\delta = 16$), decreasing the phosphorylation rate ω from levels typical of EGFR ($\omega_{\text{EGFR}} \approx 100\text{–}1000$) paradoxically increases overall receptor phosphorylation. A similar effect is observed when receptor dephosphorylation is enhanced (Fig.4d). Importantly, our model makes a testable prediction: the reversal of thermodynamic preference observed between EGF and EPGN/EREG will disappear when kinase activity is mildly suppressed (see, e.g., the curves for $\omega = 256$ and $\omega = 16$ over $\delta \in [10, 100]$), such as by treatment with low doses of the kinase inhibitor gefitinib (*Herbst et al., 2004*).

Multi-site phosphorylation and ligand-induced degradation are both essential for ligand-specificity. To assess the importance of sequential multi-site phosphorylation on ligand specificity, we analyzed A_1^N, the phosphorylation of the first site for signaling networks with N phosphorylation sites. Fig. 5a shows that multi-site phosphorylation is essential to endow signaling networks with ligand specificity and ligand-induced receptor degradation alone is not sufficient. This is because the non-monotonic preference for intermediate affinity ligands arises only when the receptors can be sorted among multiple phosphorylation sites: earlier ones for low affinity ligands and later ones for high affinity ligands.

To assess how receptor degradation shapes ligand specificity for a multi-site phosphorylation network, we examined how altering receptor turnover influences model behavior. As shown in Fig. 5b, the optimal dissociation rate δ_{opt} —which maximizes receptor phosphorylation levels—increases with ligand-induced degradation rate β. When kinase activity ω, low affinity ligands (high δ) over high affinity ligands. Crucially, this optimal δ_{opt} emerges only when receptor degradation is strong ($\beta \gg 1$). These predictions can be tested by blocking receptor degradation, e.g., via mutation of ubiquitination sites (*Gerritsen et al., 2023*).

To quantify ligand specificity, we computed receptor phosphorylation in response to ligands differing by at least one $k_{\text{B}}T$ in binding free energy from the optimal ligand. Figure 5c shows that as β increases, phosphorylation downstream of suboptimal ligands (red line in inset) declines relative to the optimal ligand. This enhanced specificity is further amplified by increasing kinase activity ω.

These results show that both multi-site phosphorylation and ligand induced degradation are essential for ligand specificity.

Discussion

Cells face the formidable task of decoding multiple extracellular signals to generate appropriate, context-specific responses. This challenge is especially acute for receptors like RTKs, GPCRs, and interleukin receptors, which bind multiple cognate ligands and yet elicit distinct downstream outcomes. While equilibrium affinity provides a baseline for ligand specificity, it cannot fully explain the rich and often counterintuitive behaviors observed in many signaling systems.

Here, we show that non-equilibrium mechanisms—specifically, *kinetic sorting* through multi-site phosphorylation and ligand-induced degradation—can explain how signaling networks achieve ligand specificity beyond equilibrium limits. In the model, high-affinity ligand–receptor complexes are sorted toward degradation-prone states, low-affinity complexes are sorted towards inactivated

Figure 5. Multiple phosphorylation sites and receptor degradation dictate ligand specificity. (a) Activity A_1 of the first phosphorylation site as a function of the dissociation rate δ for signaling networks with different number of phosphorylation sites. (b) The optimal dissociation rate δ_{opt} that leads to maximum phosphorylation activity as a function of dimensionless degradation rate β for different values of ω. δ_{opt} is shown only if $\delta_{\text{opt}} \in [1, 1000]$. (c) The relative activity of a ligand with dissociation rate that differs by $k_{\text{B}}T$ compared to δ_{opt} plotted as a function of β for different values of ω (see inset). Of the two ligands that differ in stability by $k_{\text{B}}T$, the ligand exhibiting maximum activity is considered.

states, and intermediate-affinity ligands strike the optimal balance between progression and degradation to maximize signaling. This framework explains paradoxical features observed in RTK systems, including the non-monotonic dependence of phosphorylation on ligand affinity and kinase activity.

Importantly, our model predicts that early phosphorylation sites show the strongest ligand discrimination, consistent with recent experimental observations. It also makes the testable prediction that impairing receptor degradation should reduce specificity by eliminating the kinetic sorting effect.

More broadly, our findings suggest that ligand-specific sorting of multidimensional receptor states—across phosphorylation, degradation, localization, and engagement fates—may be a general strategy for encoding ligand identity. While our model focused on the activity–degradation axis, real-world receptors operate in even richer state spaces, offering exciting directions for future work.

Our findings complement prior studies on mechanisms of ligand specificity that operate at thermal equilibrium, such as those described in the Bone Morphogenetic Protein (BMP) pathway (*Antebi et al., 2017*; *Su et al., 2022*; *Parres-Gold et al., 2025*). BMP signaling relies on promiscuous ligand–receptor interactions, with specificity emerging from differences in receptor abundance, binding affinity, and complex activity. By contrast, our work shows that non-equilibrium mechanisms—such as phosphorylation cycles and ligand-induced receptor degradation—can achieve ligand discrimination even for a single receptor type. Given that ligand–receptor promiscuity, multisite phosphorylation, and receptor turnover are common features across signaling systems (e.g., in the EGFR/ErbB family (*Linggi and Carpenter, 2006*)), it is likely that biological networks integrate both equilibrium and non-equilibrium strategies to achieve robust and tunable ligand specificity.

In recent years, there has been growing interest in engineering synthetic physical and chemical circuits capable of carrying out complex computational tasks, including input discrimination, classification, prediction, and the generation of multiple stable cell states (*Shakiba et al., 2021*; *Ma et al., 2022*; *Benzinger et al., 2022*; *Zhu et al., 2022*; *Floyd et al., 2024*; *Parres-Gold et al., 2025*; *Aoki et al., 2019*). Some of these synthetic strategies rely on equilibrium thermodynamics (*Parres-Gold et al., 2025*), while others exploit non-equilibrium steady states (*Floyd et al., 2024*). We propose that non-equilibrium kinetic sorting, which harnesses receptor synthesis and degradation, could provide synthetic biologists with a powerful framework for achieving precise control over molecular abundances and dynamic system behavior.

Finally, we address a major concern in non-equilibrium signaling circuits: the energetic cost of operation. Previous theoretical work has shown that free energy dissipation places fundamental constraints on the performance of signaling networks (*Bryant and Machta, 2023*; *Lan et al.,*

2012; *Mehta and Schwab, 2012*; *Qian and Reluga, 2005*; *Cao et al., 2015*; *Azeloglu and Iyengar, 2015*; *Floyd et al., 2024*; *Mahdavi et al., 2024*). These studies typically focus on futile cycles of reversible modifications such as phosphorylation or methylation. In contrast, ligand-induced receptor degradation—a central feature of many signaling networks—is a far more energy-intensive process. For example, MCF10A cells maintain approximately 10^5 EGFR molecules on the surface (each 1,210 amino acids in length)(*Shi et al., 2016*), with a synthesis rate of about 15 receptors per second(*Lyashenko et al., 2020*), corresponding to an energetic cost of roughly $\sim 8 \times 10^4$ ATP/sec (assuming 4.5 ATP per peptide bond (*Milo et al., 2010*)). By comparison, EGFR dephosphorylation occurs over ~ 15 seconds (*Kleiman et al., 2011*), and only $5-10\%$ of receptors are phosphorylated at steady state (*Shi et al., 2016*; *Feng et al., 2023*), resulting in a much lower energetic cost of $\sim 6 \times 10^2$ ATP/sec for dephosphorylation. Thus, the energetic burden of receptor turnover can exceed that of reversible modification cycles by up to two orders of magnitude. These estimates suggest that, at least in eukaryotic systems, the energetic demands of non-equilibrium modification cycles are unlikely to pose a fundamental limitation on the functionality of signaling networks.

References

Antebi YE, Linton JM, Klumpe H, Bintu B, Gong M, Su C, McCardell R, Elowitz MB. Combinatorial signal perception in the BMP pathway. Cell. 2017; 170(6):1184–1196.

Aoki SK, Lillacci G, Gupta A, Baumschlager A, Schweingruber D, Khammash M. A universal biomolecular integral feedback controller for robust perfect adaptation. Nature. 2019; 570(7762):533–537.

Azeloglu EU, Iyengar R. Signaling networks: information flow, computation, and decision making. Cold Spring Harbor perspectives in biology. 2015; 7(4):a005934.

Benzinger D, Ovinnikov S, Khammash M. Synthetic gene networks recapitulate dynamic signal decoding and differential gene expression. Cell Systems. 2022; 13(5):353–364.

Bryant SJ, Machta BB. Physical constraints in intracellular signaling: The cost of sending a bit. Physical review letters. 2023; 131(6):068401.

Cantley LC, Hunter T, Sever R, Thorner JW. Signal transduction: principles, pathways, and processes. Cold Spring Harbor Laboratory Press Cold Spring Harbor, NY, USA:; 2014.

Cao Y, Wang H, Ouyang Q, Tu Y. The free-energy cost of accurate biochemical oscillations. Nature physics. 2015; 11(9):772–778.

Cendrowski J, Mamińska A, Miaczynska M. Endocytic regulation of cytokine receptor signaling. Cytokine & growth factor reviews. 2016; 32:63–73.

Charpentier JC, King PD. Mechanisms and functions of endocytosis in T cells. Cell Communication and Signaling. 2021; 19(1):92.

Chen WW, Niepel M, Sorger PK. Classic and contemporary approaches to modeling biochemical reactions. Genes & development. 2010; 24(17):1861–1875.

Chen WW, Schoeberl B, Jasper PJ, Niepel M, Nielsen UB, Lauffenburger DA, Sorger PK. Input–output behavior of ErbB signaling pathways as revealed by a mass action model trained against dynamic data. Molecular systems biology. 2009; 5(1):239.

Clark RB, Knoll BJ, Barber R. Partial agonists and G protein-coupled receptor desensitization. Trends in pharmacological sciences. 1999; 20(7):279–286.

Coombs D, Kalergis AM, Nathenson SG, Wofsy C, Goldstein B. Activated TCRs remain marked for internalization after dissociation from pMHC. Nature immunology. 2002; 3(10):926–931.

Feng S, Sanford JA, Weber T, Hutchinson-Bunch CM, Dakup PP, Paurus VL, Attah K, Sauro HM, Qian WJ, Wiley HS. A Phosphoproteomics Data Resource for Systems-level Modeling of Kinase Signaling Networks. bioRxiv. 2023; .

Floyd C, Dinner AR, Murugan A, Vaikuntanathan S. Limits on the computational expressivity of non-equilibrium biophysical processes. arXiv preprint arXiv:240905827. 2024; .

Freed DM, Bessman NJ, Kiyatkin A, Salazar-Cavazos E, Byrne PO, Moore JO, Valley CC, Ferguson KM, Leahy DJ,
Lidke DS, et al. EGFR ligands differentially stabilize receptor dimers to specify signaling kinetics. Cell. 2017;
171(3):683–695.

Furdui CM, Lew ED, Schlessinger J, Anderson KS. Autophosphorylation of FGFR1 kinase is mediated by a se-
quential and precisely ordered reaction. Molecular cell. 2006; 21(5):711–717.

Gerritsen JS, Faraguna JS, Bonavia R, Furnari FB, White FM. Predictive data-driven modeling of C-terminal
tyrosine function in the EGFR signaling network. Life Science Alliance. 2023; 6(8).

Gulukota K, Wolynes PG. Statistical mechanics of kinetic proofreading in protein folding in vivo. Proceedings
of the National Academy of Sciences. 1994; 91(20):9292–9296.

Herbst RS, Fukuoka M, Baselga J. Gefitinib—a novel targeted approach to treating cancer. Nature Reviews
Cancer. 2004; 4(12):956–965.

Hilleren P, Parker R. mRNA surveillance in eukaryotes: kinetic proofreading of proper translation termination
as assessed by mRNP domain organization? Rna. 1999; 5(6):711–719.

Hopfield JJ. Kinetic proofreading: a new mechanism for reducing errors in biosynthetic processes requiring
high specificity. Proceedings of the National Academy of Sciences. 1974; 71(10):4135–4139.

Hopfield JJ. The energy relay: a proofreading scheme based on dynamic cooperativity and lacking all character-
istic symptoms of kinetic proofreading in DNA replication and protein synthesis. Proceedings of the National
Academy of Sciences. 1980; 77(9):5248–5252.

Hu C, Leche CA, Kiyatkin A, Yu Z, Stayrook SE, Ferguson KM, Lemmon MA. Glioblastoma mutations alter EGFR
dimer structure to prevent ligand bias. Nature. 2022; 602(7897):518–522.

Huang WY, Alvarez S, Kondo Y, Lee YK, Chung JK, Lam HYM, Biswas KH, Kuriyan J, Groves JT. A molecu-
lar assembly phase transition and kinetic proofreading modulate Ras activation by SOS. Science. 2019;
363(6431):1098–1103.

Huang Y, Bharill S, Karandur D, Peterson SM, Marita M, Shi X, Kaliszewski MJ, Smith AW, Isacoff EY, Kuriyan J.
Molecular basis for multimerization in the activation of the epidermal growth factor receptor. elife. 2016;
5:e14107.

Irannejad R, Von Zastrow M. GPCR signaling along the endocytic pathway. Current opinion in cell biology.
2014; 27:109–116.

Kiyatkin A, van Alderwerelt van Rosenburgh IK, Klein DE, Lemmon MA. Kinetics of receptor tyrosine kinase
activation define ERK signaling dynamics. Science Signaling. 2020; 13(645):eaaz5267.

Kleiman LB, Maiwald T, Conzelmann H, Lauffenburger DA, Sorger PK. Rapid phospho-turnover by receptor
tyrosine kinases impacts downstream signaling and drug binding. Molecular cell. 2011; 43(5):723–737.

Koenig JA, Edwardson JM. Endocytosis and recycling of G protein-coupled receptors. Trends in pharmacological
sciences. 1997; 18(8):276–287.

Kollewe C, Mackensen AC, Neumann D, Knop J, Cao P, Li S, Wesche H, Martin MU. Sequential autophos-
phorylation steps in the interleukin-1 receptor-associated kinase-1 regulate its availability as an adapter in
interleukin-1 signaling. Journal of Biological Chemistry. 2004; 279(7):5227–5236.

Lan G, Sartori P, Neumann S, Sourjik V, Tu Y. The energy–speed–accuracy trade-off in sensory adaptation.
Nature physics. 2012; 8(5):422–428.

Latorraca NR, Masureel M, Hollingsworth SA, Heydenreich FM, Suomivuori CM, Brinton C, Townshend RJ, Bou-
vier M, Kobilka BK, Dror RO. How GPCR phosphorylation patterns orchestrate arrestin-mediated signaling.
Cell. 2020; 183(7):1813–1825.

Lemmon MA, Schlessinger J. Cell signaling by receptor tyrosine kinases. Cell. 2010; 141(7):1117–1134.

Lever M, Maini PK, Van Der Merwe PA, Dushek O. Phenotypic models of T cell activation. Nature Reviews
Immunology. 2014; 14(9):619–629.

Limbird LE, De Meyts P, Lefkowitz RJ. β-Adrenergic receptors: evidence for negative cooperativity. Biochemical
and biophysical research communications. 1975; 64(4):1160–1168.

Linggi B, Carpenter G. ErbB receptors: new insights on mechanisms and biology. Trends in cell biology. 2006;
16(12):649–656.

Liu H, Rhodes M, Wiest DL, Vignali DA. On the dynamics of TCR: CD3 complex cell surface expression and
downmodulation. Immunity. 2000; 13(5):665–675.

Lyashenko E, Niepel M, Dixit PD, Lim SK, Sorger PK, Vitkup D. Receptor-based mechanism of relative sensing
and cell memory in mammalian signaling networks. Elife. 2020; 9:e50342.

Ma Y, Budde MW, Mayalu MN, Zhu J, Lu AC, Murray RM, Elowitz MB. Synthetic mammalian signaling circuits for
robust cell population control. Cell. 2022; 185(6):967–979.

Macdonald JL, Pike LJ. Heterogeneity in EGF-binding affinities arises from negative cooperativity in an aggre-
gating system. Proceedings of the National Academy of Sciences. 2008; 105(1):112–117.

Macdonald-Obermann JL, Pike LJ. Different epidermal growth factor (EGF) receptor ligands show distinct
kinetics and biased or partial agonism for homodimer and heterodimer formation. Journal of Biological
Chemistry. 2014; 289(38):26178–26188.

Madsen RR, Le Marois A, Mruk ON, Voliotis M, Yin S, Sufi J, Qin X, Zhao SJ, Gorczynska J, Morelli D, et al. Oncogenic
PIK3CA corrupts growth factor signaling specificity. Molecular Systems Biology. 2025; 21(2):126–157.

Madsen RR, Vanhaesebroeck B. Cracking the context-specific PI3K signaling code. Science Signaling. 2020;
13(613):eaay2940.

Mahdavi SD, Salmon GL, Daghlian P, Garcia HG, Phillips R. Flexibility and sensitivity in gene regulation out of
equilibrium. Proceedings of the National Academy of Sciences. 2024; 121(46):e2411395121.

McKeithan TW. Kinetic proofreading in T-cell receptor signal transduction. Proceedings of the national
academy of sciences. 1995; 92(11):5042–5046.

Mehta P, Schwab DJ. Energetic costs of cellular computation. Proceedings of the National Academy of Sciences.
2012; 109(44):17978–17982.

Milo R, Jorgensen P, Moran U, Weber G, Springer M. BioNumbers—the database of key numbers in molecular
and cell biology. Nucleic acids research. 2010; 38(suppl_1):D750–D753.

Mudumbi KC, Burns EA, Schodt DJ, Petrova ZO, Kiyatkin A, Kim LW, Mangiacapre EM, Ortiz-Caraveo I, Ortiz HR,
Hu C, et al. Distinct interactions stabilize EGFR dimers and higher-order oligomers in cell membranes. Cell
reports. 2024; 43(1).

Murugan A, Huse DA, Leibler S. Discriminatory proofreading regimes in nonequilibrium systems. Physical
Review X. 2014; 4(2):021016.

Myers PJ, Lee SH, Lazzara MJ. An integrated mechanistic and data-driven computational model predicts cell
responses to high-and low-affinity EGFR ligands. bioRxiv. 2023; p. 2023–06.

Ninio J. Kinetic amplification of enzyme discrimination. Biochimie. 1975; 57(5):587–595.

Parres-Gold J, Levine M, Emert B, Stuart A, Elowitz MB. Contextual computation by competitive protein dimer-
ization networks. Cell. 2025; .

Qian H, Reluga TC. Nonequilibrium thermodynamics and nonlinear kinetics in a cellular signaling switch. Phys-
ical review letters. 2005; 94(2):028101.

Qiao L, Khalilimeybodi A, Linden-Santangeli NJ, Rangamani P. The evolution of systems biology and systems
medicine: From mechanistic models to uncertainty quantification. Annual Review of Biomedical Engineering.
2025; 27.

Schulze WX, Deng L, Mann M. Phosphotyrosine interactome of the ErbB-receptor kinase family. Molecular
systems biology. 2005; 1(1):2005–0008.

Shakiba N, Jones RD, Weiss R, Del Vecchio D. Context-aware synthetic biology by controller design: Engineering
the mammalian cell. Cell systems. 2021; 12(6):561–592.

Shi T, Niepel M, McDermott JE, Gao Y, Nicora CD, Chrisler WB, Markillie LM, Petyuk VA, Smith RD, Rodland KD,
et al. Conservation of protein abundance patterns reveals the regulatory architecture of the EGFR-MAPK
pathway. Science Signaling. 2016; 9(436):rs6–rs6.

Sorkin A, Goh LK. Endocytosis and intracellular trafficking of ErbBs. Experimental cell research. 2009;
315(4):683–696.

Su CJ, Murugan A, Linton JM, Yeluri A, Bois J, Klumpe H, Langley MA, Antebi YE, Elowitz MB. Ligand-receptor
promiscuity enables cellular addressing. Cell systems. 2022; 13(5):408–425.

Subtil A, Hémar A, Dautry-Varsat A. Rapid endocytosis of interleukin 2 receptors when clathrin-coated pit
endocytosis is inhibited. Journal of cell science. 1994; 107(12):3461–3468.

Tobin A. G-protein-coupled receptor phosphorylation: Where, when and by whom. British journal of pharma-
cology. 2008; 153(S1):S167–S176.

Wiley HS. Trafficking of the ErbB receptors and its influence on signaling. The EGF Receptor Family. 2003; p.
81–91.

Zhu R, del Rio-Salgado JM, Garcia-Ojalvo J, Elowitz MB. Synthetic multistability in mammalian cells. Science.
2022; 375(6578):eabg9765.

Supplementary materials

Equations for proofreading models

The equations describing species abundances in the traditional KPR model similar to that of McK-
eithan (*McKeithan, 1995*) are as follows:

$$\frac{dR}{dt} = -k_{\mathrm{on}}LR + k_{\mathrm{d}}B + k_{\mathrm{d}}\sum_{i=1}^{N}P_i \tag{2}$$

$$\frac{dB}{dt} = +k_{\mathrm{on}}LR - k_{\mathrm{d}}B - k_{\mathrm{p}}B \tag{3}$$

$$\frac{dP_1}{dt} = k_{\mathrm{p}}B - k_{\mathrm{p}}P_1 - k_{\mathrm{d}}P_1 \tag{4}$$

$$\frac{dP_i}{dt} = k_{\mathrm{p}}P_{i-1} - k_{\mathrm{p}}P_i - k_{\mathrm{d}}P_i \ \forall \ i \in [2, N-1] \tag{5}$$

$$\frac{dP_N}{dt} = k_{\mathrm{p}}P_{N-1} - k_{\mathrm{d}}P_N \tag{6}$$

395 For the limited signaling model, the dynamics of B, and $P_i, i \in [1, N-1]$ are identical to the
396 traditional KPR model. The dynamics of R and P_N are modified as follows:

$$\frac{dR}{dt} = -k_{\mathrm{on}}LR + k_{\mathrm{d}}B + k_{\mathrm{d}}\sum_{i=1}^{N}P_i + k_{\mathrm{d}}I \tag{7}$$

$$\frac{dP_N}{dt} = k_{\mathrm{p}}P_{N-1} - k_{\mathrm{d}}P_N - k_{\mathrm{in}}P_N \tag{8}$$

$$\frac{dI}{dt} = k_{\mathrm{in}}P_N - k_{\mathrm{d}}P_I \tag{9}$$

Equations for the model with receptor degradation

Signaling receptors participate in a variety of complex regulatory processes, including non-linear
ligand binding dynamics (*Limbird et al., 1975*; *Macdonald and Pike, 2008*), receptor oligomeriza-
tion (*Mudumbi et al., 2024*; *Huang et al., 2016*), context-specific interactions with adapter pro-
teins (*Madsen and Vanhaesebroeck, 2020*; *Feng et al., 2023*), and trafficking between cellular com-
partments leading to degradation (*Sorkin and Goh, 2009*; *Wiley, 2003*; *Irannejad and Von Zastrow,*
2014).

While computational models that incorporate these mechanistic details are powerful tools for
hypothesis generation (*Chen et al., 2010*; *Qiao et al., 2025*), they often require large-scale datasets
for accurate parameterization (*Feng et al., 2023*). As an alternative, simplified models that inten-
tionally omit certain mechanistic details can still yield deep qualitative insights, even if they cannot
quantitatively reproduce experimental data.

In this study, we present such a simplified model aimed at explaining two paradoxical features of receptor tyrosine kinase (RTK) signaling: (1) the non-monotonic relationship between ligand-receptor affinity and steady-state receptor phosphorylation (*Freed et al., 2017*; *Madsen et al., 2025*; *Myers et al., 2023*), and (2) the counterintuitive increase in receptor phosphorylation following mild kinase inhibition (*Kleiman et al., 2011*; *Kiyatkin et al., 2020*).

To keep the model simple and tractable, we neglect receptor recycling and oligomerization. Previously, we showed that the combined effects of endocytosis, recycling, and degradation can be captured by a single effective dimensionless parameter, β in this study, which reflects the degradation bias of fully phosphorylated receptors compared to partially phosphorylated receptors(*Lyashenko et al., 2020*). Similarly, receptor dimerization and negative cooperativity can be abstracted into a Hill coefficient $\eta < 1$ (*Lyashenko et al., 2020*). For the phenomena explored here, including oligomerization would modify the shape of the response curves but not their qualitative behavior.

Under these assumptions, the governing equations for the model are given by:

$$
\frac{dR}{dt} = k_{\text{delivery}} - k_{\text{on}} L R + k_{\text{d}} B + k_{\text{d}} \sum_{i=1}^{N} P_i
$$
$$
- \quad k_{\text{int}} R \tag{10}
$$

$$
\frac{dB}{dt} = k_{\text{on}} L R - k_{\text{d}} B - k_{\text{p}} B - k_{\text{int}} B \tag{11}
$$

$$
\frac{dP_1}{dt} = k_{\text{p}} B - k_{\text{p}} P_1 - k_{\text{d}} P_1 - k_{\text{int}} P_1 \tag{12}
$$

$$
\frac{dP_i}{dt} = k_{\text{p}} P_{i-1} - k_{\text{p}} P_i - k_{\text{d}} P_i
$$
$$
- \quad k_{\text{int}} P_i \ \forall \ i \in [2, N-1] \tag{13}
$$

$$
\frac{dP_N}{dt} = k_{\text{p}} P_{N-1} - k_{\text{d}} P_N - k_{\text{int}}^* P_N \tag{14}
$$

All equations are solved at steady state and in the limit $u \to \infty$. All codes required to generate the figures in the manuscript can be found at https://github.com/BarriosJer0/KineticSorting.