A *Drosophila* screen of schizophrenia-related genes highlights the requirement of

neural and glial matrix metalloproteinases for neuronal remodeling

Shir Keret[1,2], Hagar Meltzer[1,2*], Neta Marmor-Kollet[1,2,3] Oren Schuldiner[1,2*]

[1] Department of Molecular Cell Biology, Weizmann Institute of Science, Rehovot, Israel

[2] Department of Molecular Neuroscience, Weizmann Institute of Science, Rehovot,

Israel

[3] Current affiliation: Department of Biology, Brandeis University, MA, USA

* Correspondence to: hagar.meltzer@weizmann.ac.il; oren.schuldiner@weizmann.ac.il

Abstract

Schizophrenia (SCZ) is a multifactorial neuropsychiatric disorder of complex and mostly unknown etiology, affected by genetic, developmental and environmental factors. Neuroanatomical abnormalities, such as loss of grey matter, are apparent prior to the onset of symptoms, suggesting neurodevelopmental origin. Indeed, it has been hypothesized, and recently experimentally supported, that SCZ is associated with dysregulation of developmental synaptic pruning. Here, we explore the molecular link between SCZ-associated genes and developmental neuronal remodeling, focusing on the *Drosophila* mushroom body (MB), which undergoes stereotypic remodeling during metamorphosis. We conducted a loss-of-function screen in which we knocked down *Drosophila* homologs of human genes that genomic studies have associated with SCZ. Out of our 'positive hits', we focused on matrix metalloproteinases (MMPs), mostly known for their role in remodeling of the extracellular matrix. We found that both Mmp1 and Mmp2, which are closely-related to mammalian MMPs, are required in neurons and in glia for the pruning of MB axons. Our combinatorial loss-of-function experiments suggest that Mmp2 secretion from glia is the most important promoter of axon pruning. Our results shed new light on potential molecular players underlying neurodevelopmental defects in SCZ and highlight the advantage of genetically tractable model organisms in the study of human neurodevelopmental disorders.

Introduction

Following its initial establishment, the developing nervous system of both vertebrates and invertebrates undergoes remodeling to shape its mature connectivity. In humans, developmental neuronal remodeling occurs postnatally and includes degenerative events such as retraction of synapses and large-scale axon elimination. While these processes occur most prominently during the first two years of life, they continue until after adolescence (Luo & O'Leary, 2005; Schuldiner & Yaron, 2015). Defects in neuronal remodeling have long been hypothesized to be one of the underlying causes of neuropsychiatric disorders, including schizophrenia (SCZ; Feinberg, 1982; Penzes et al., 2011).

SCZ is a severe, multifactorial mental health condition affecting ~1% of the population. It is characterized by delusions, hallucinations, social withdrawal and cognitive deficits, typically onsetting during the late teens to early adulthood (McCutcheon et al., 2020). It is known from MRI studies that structural abnormalities in the brain, such as reduction in grey matter and ventricular enlargement, may precede the onset of psychotic symptoms, implying neurodevelopmental roots (Brent et al., 2013; Suzuki et al., 2002; Zikidi et al., 2020). It has been suggested, for many years, that the characteristic neuroanatomical alterations in individuals with SCZ are due to dysregulation of neuronal remodeling, specifically over-pruning of synapses (Feinberg, 1982; Hoffman & Dobscha, 1989; Keshavan et al., 1994). The first experimental evidence to support this hypothesis came several years ago, and associated SCZ with excessive expression of complement components leading to increased synapse elimination by microglia (Sekar et al., 2016). This work as well as others highlighted glia, including microglia and astrocytes, as key mediators of synapse pruning in normal development as well as in SCZ and other neuropathologies (Reviewed in Laricchiuta et al., 2024; Notter, 2021; Scott-Hewitt et al., 2023). Still, the molecular basis of SCZ risk and its link to abnormalities in neuronal remodeling remains mostly unknown.

Drosophila melanogaster is an ideal model to study neuronal remodeling, due to its unparalleled genetic toolkit combined with its massive stereotypic circuit remodeling during metamorphosis (Truman, 1990; Yaniv & Schuldiner, 2016). Our focus is on the *Drosophila* mushroom body (MB), a brain structure that functions as a center of olfactory

learning and memory (Heisenberg, 1998). The MB is comprised of three types of intrinsic
neurons, known as Kenyon cells (KCs), which are sequentially born from identical
neuroblasts (Fig. 1A). Out of the three KC types (γ, α'/β' and α/β), only the first-born γ-
KCs undergo stereotypic remodeling. When they initially grow, in the embryonic and early
larval stages, γ-KCs project axons that bifurcate to form vertical and medial lobes. At the
early pupal stage (~6 hours after puparium formation; h APF) both axonal lobes begin to
undergo pruning up to their branchpoint (which is over at ~18h APF). Subsequently, γ-
KCs initiate axon regrowth to create the adult-specific medially projecting γ-lobe (Lee et
al., 1999; Yaniv & Schuldiner, 2016; Fig. 1A). Importantly, glia were shown, by our lab and
others, to actively participate in the remodeling of γ-KCs (Awasaki et al., 2011; Awasaki
& Ito, 2004; Boulanger et al., 2021; Hakim et al., 2014; Marmor-Kollet et al., 2023; Watts
et al., 2004). Three glial subtypes are present around the MB: cortex glia, ensheathing
glia and astrocyte-like glia (hereafter referred to as astrocytes; Freeman, 2015). Cortex
glia were shown to have an instructive role in pruning via secretion of the TGF-β ligand
myoglianin, which subsequently activates a pruning program in the γ-KCs (Awasaki et al.,
2011). Astrocytes, however, are the main cleaners of axonal debris following pruning
(Hakim et al., 2014; Tasdemir-Yilmaz & Freeman, 2014). Moreover, we recently
demonstrated that astrocytes actively infiltrate the axon bundle prior to pruning to facilitate
axon defasciculation and pruning (Marmor-Kollet et al., 2023).

Importantly, *Drosophila* is routinely used to study human neurodegenerative disorders
(Lessing & Bonini, 2009), and the Disrupted-in-Scizophrenia-1 (DISC1) gene has been
shown to affect MB structure and its regulation of sleep (Furukubo-Tokunaga, 2009;
Sawamura et al., 2008). Here, we harness the fly genetic toolkit and the well-
characterized remodeling of the MB to delve into the molecular underpinnings of the long-
standing hypothesis, and recent evidence supporting it, that defects in neuronal
remodeling contribute to SCZ etiology. We systematically examine how MB remodeling
is affected by perturbations in *Drosophila* genes whose human homologs have been
associated with SCZ. Our findings highlight novel players in axon pruning, and provide
potential candidates for future research on the neurodevelopmental molecular origin of
SCZ.

Results

Drosophila homologs of human SCZ-related SNPs are dynamically expressed during the development of the mushroom body γ-KCs

We searched genome-wide association studies (GWAS) for candidate genes that have SNPs in SCZ patients (Wu et al., 2017; http://www.szdb.org/index.html). SNP-containing genes were converted to their homolog *Drosophila* genes using HumanMine (Lyne et al., 2022; Fig. 1B). To narrow our candidate list to genes that are potentially required for pruning, we used our previously generated transcriptional atlas of developing γ-KCs (Alyagor et al., 2018) to identify genes that are specifically upregulated before the onset of (0-6h APF), or during (6-18h APF), γ-axon pruning (Fig. 1B, Fig. 1C clusters III-VI highlighted). This process resulted in 32 candidate genes, out of which 29 had available RNAi lines for screening (Fig.1B).

Drosophila homologs of human SCZ-associated genes are required for γ-KC axon pruning

If gene upregulation prior to γ axon pruning is due to requirement in the pruning process, then knockdown (KD) is expected to inhibit pruning and result in persistence of larval γ axons in the adult brain. Therefore, we screened candidate genes using a loss-of-function (LOF) strategy by tissue-specific expression of RNAi-species via the Gal4-UAS system (Brand & Perrimon, 1993). For KD in γ-KCs, we used the pan-KC driver OK107-Gal4, which is strongly expressed in all KC types (α/β, α'/β' and γ), combined with a second, independent binary system (Riabinina & Potter, 2016) to visualize γ-KCs (using the more specific R71G10-QF2 driver, which is expressed in all γ-KCs and stochastically in α/β-KCs, to drive mtdT-HA expression). Screening the 29 RNAi lines using OK107-Gal4 resulted in 11 genes displaying varying degrees of pruning defects upon KD, and one gene whose KD led to abnormal MB morphology (Fig. 2). For 4 genes, KD resulted in lethality (Fig. 2K).

Drosophila Mmp2 is required in both KCs and glia for γ axon pruning

Many of the 'positive hits' in the screen are promising directions for further exploration. However, before deciding which genes to focus on in this study, we also reviewed the expression of the SCZ-related genes in glia - due to their known roles in neuronal

remodeling and synaptic mis-regulation in SCZ (Scott-Hewitt et al., 2023; Sekar et al.,

2016). Thus, we turned to our previously generated transcriptional profiling of astrocytes

in the *Drosophila* brain (Marmor-Kollet et al., 2023), and analyzed the larval vs. adult

expression of all of the SCZ homolog *Drosophila* genes (Fig. 3A). Genes that are more

highly expressed in late larval compared to adult astrocytes are good candidates to be

involved in γ axon pruning, which is initiated at early pupal stages. Interestingly, two genes

that came up in our KC screen as required for axon pruning, are also significantly

upregulated in larval astrocytes: Syntaxin13 and Matrix Metalloproteinase 2 (Mmp2; Fig.

3A and Fig. 2B,D). We decided to delve deeper into Mmp2, as a previous, CRISPR-based

screen done by our lab also found it to be required in γ-KCs for axon pruning (Meltzer et

al., 2019). To test whether Mmp2 is required in glia for γ axon pruning, we used the pan-

glial driver Repo-Gal4 to KD Mmp2 (in combination with R71G10-QF2-driven QUAS-

mtdT-HA to visualize the γ-KCs). Indeed, this resulted in a γ axon pruning defect,

indicating that Mmp2 originating from glia promotes MB γ axon pruning (Fig. 3C-D). We

further knocked down Mmp2 using the astrocyte-specific driver Alrm-Gal4, which also

resulted in a pruning defect, although milder (Fig. S1 A,C), validating its requirement in

astrocytes. The milder defect could be due to weaker RNAi expression in astrocytes by

the Alrm-Gal4 driver compared to Repo-Gal4, or because of Mmp2 secretion by additional

glial subtypes (e.g., cortex or ensheathing glia). Notably, the initial growth of γ-KCs, as

evident in the late larval stage, is unaffected by Mmp2 KD in either KCs or glia (Fig. S1

D,F,G,I).

Mmp1 is required in glia, and to a lesser extent in KCs, for γ axon pruning

MMPs are extracellular proteases that can cleave extracellular matrix (ECM) components

(Page-McCaw et al., 2007). In mouse, there are 23 MMPs, while the zebrafish genome

encodes 25 MMPs[26]. The *Drosophila* genome, in contrast, encodes only two MMPs –

Mmp1 and Mmp2 – which together have 13 predicted isoforms, most of which are likely

secreted, while each Mmp also has a glycosylphosphatidylinositol (GPI)-anchored

isoform (LaFever et al., 2017). The significantly reduced complexity makes *Drosophila*

MMPs an ideal model system to uncover novel insights into their roles in neuronal

remodeling. Since both *Drosophila* MMPs are functionally similar, we tested the potential

requirement of Mmp1 in KCs as well as glia. Indeed, KD of Mmp1 in glia results in

significant γ axon pruning defect (Fig. 4A-C), while its KD in KCs results in a mild γ axon

pruning defect (Fig. 4D-F). As with Mmp2, astrocyte-specific Mmp1 knockdown with Alrm-

Gal4 also results in a pruning defect, but milder than with Repo-Gal4 (Fig. S1 A-B). Initial

growth of the γ-KCs is unaffected by Mmp1 KD in either KCs nor glia (Fig. S1 D-E, G-H).

While axon pruning of γ-KCs relies on both glial and neuronal MMPs, the main

requirement is glial Mmp2

Since most Mmp1 and Mmp2 isoforms are secreted, we speculated that both are secreted

in parallel from KCs and glia to promote axon pruning. Therefore, we preformed different

KD combinations of Mmp1 and Mmp2 in KCs and/or glia (Fig. 5). Our results draw two

main conclusions: first, while both Mmp1 and Mmp2 are required for pruning, the

contribution of Mmp2 is more significant as, in all conditions, the double KD of Mmp1 plus

Mmp2 did not further exacerbate the pruning defect of Mmp2 KD alone (Fig. 5 A-L).

Second, while Mmp1/2 expression from both neurons (KCs) and glia is required for

pruning of KC-axons, the main source is likely glia, as KD of a single or double Mmp in

both glia and neurons was not significantly more severe than the respective KD in glia

only (Fig. 5, quantified in M-O).

Taken together, our results suggest that while both MMPs affect pruning via secretion

from both neurons and glia, the strongest requirement for MB γ axon pruning comes from

glial-secreted Mmp2.

Discussion

Four decades ago, Feinberg hypothesized that SCZ is caused by aberrant synaptic

pruning during adolescence: "too many, too few or the wrong synapses are eliminated"

(Feinberg, 1982). In recent years, accumulating findings are providing experimental

support of this hypothesis (Keshavan et al., 2020; Sekar et al., 2016; Sellgren et al.,

2019). In this work, we aimed to further explore the association of SCZ with defective

neuronal remodeling, by utilizing the stereotypic remodeling of the *Drosophila* MB.

In a genetic LOF screen for *Drosophila* homologs of human genes that contain SNPs

associated with SCZ, over 40% of the genes we screened (12 out of 29) were found to

have a role in γ-KC development, most notably (11 out of 12) in axon pruning. This high

rate of 'positive hits' strengthens the link between SCZ and defects in developmental

neuronal remodeling. Notably, we focused on genes with upregulated expression prior to
pruning, but future efforts to overexpress genes that are downregulated prior to pruning
might also reveal interesting insights.

While many of the positive hits are fascinating candidates for further research, in this
study we decided to focus on the role of MMPs in neuronal remodeling, for several
reasons. First, in our screen, Mmp2 was found to be required for γ axon pruning (Fig. 2B),
but is also highly expressed in larval astrocytes compared to adult ones. This is of
particular interest due to the speculated role of glia in abnormal synaptic pruning in SCZ
patients. Second, numerous reports have highlighted the link between MMPs and the
pathophysiology of SCZ. Specifically, elevated levels of MMPs, most prominently Mmp9,
were shown to correlate with SCZ risk and cognitive impairment (Dickerson et al., 2023;
Kudo et al., 2020; Seitz-Holland et al., 2022), emphasizing their potential as biomarkers
and therapeutic targets. While it has been suggested that this correlation is via MMP-
dependent changes in dendritic spine morphology (Lepeta et al., 2017), the precise
mechanism remains poorly understood. Third, the *Drosophila* MMPs appear to be quite
evolutionary conserved (LaFever et al., 2017; Llano et al., 2002), highlighting the potential
relevance to more translational contexts. As there are only 2 fly MMPs compared to 23 in
mammals, *Drosophila* provides a simplified and powerful model system for mechanistic
exploration.

Here, we found that the two *Drosophila* MMPs, expressed from both KCs and glia, are
required for γ axon pruning in the MB. Mmp2, specifically from glia, seems to be the most
significant requirement in this process.

MMPs are known for their roles in tissue remodeling by cleaving substrates in the
extracellular matrix (ECM; Page-McCaw et al., 2007), and are well studied in the cancer
field (Overall & López-Otín, 2002), but their mechanistic function in the developing
nervous system is less characterized (e.g.,Broadie et al., 2011; Dear et al., 2016; Gore et
al., 2021). MMPs are known to be secreted by neurons and glia, and are thought to
facilitate synaptic remodeling of dendritic spines (Huntley, 2012). In the adult brains of
rodent models, pharmacologically- (Szklarczyk et al., 2002) or injury- (Pijet et al., 2019)
induced dendritic remodeling is suggested to occur via Mmp9. In the developmental
remodeling of *Drosophila* dendritic arborization (da) neurons, both Mmp1 and Mmp2 were

shown to be required for the elimination of severed dendritic branches in a non-cell

autonomous manner (Kuo et al., 2005). Their source, as well as the mechanism by which

they promote dendrite removal, remain unknown (the authors speculate they are secreted

from phagocytic cells – at the time they suggested blood cells, but given more recent

studies from the same lab, epidermal cells are more likely; Han et al., 2014). Interestingly,

MMPs were also shown to mediate dendrite reshaping of da neurons in the mature fly via

local degeneration of the basement membrane, further reinforcing their significance in

refining the mature nervous system (Yasunaga et al., 2010).

The mechanism by which MMPs promote remodeling of MB γ-KCs, and specifically their

substrates in this context, remain to be uncovered. Since the main known role of MMPs

is cleaving ECM substrates, they may act as 'path cleaners' for the migration of cells and

molecules (Page-McCaw et al., 2007). This function can be relevant to axon pruning in

several, non-mutually exclusive ways. One option is that MMPs allow the migration of

Myoglianin (Myo) in the ECM. Myo is part of the pruning initiation cascade and was shown

to be secreted by cortex glia and astrocytes (Awasaki et al., 2011), and bind the Baboon

receptor on γ-KCs to eventually promote expression of EcR-B1, a key regulator of MB

remodeling (Yu et al., 2013; Zheng et al., 2003). Moreover, it was shown that the human

Mmp9 can proteolytically activate latent TGF-β, which is the human homolog of the

Drosophila Myo (Yu & Stamenkovic, 2000). However, we found that the expression of

EcR-B1 in γ-KC cell bodies in unaffected by Mmp1/Mmp2 knockdown, in neither glia nor

KCs (Fig. S2). Another option is that MMPs facilitate astrocyte infiltration. Astrocytes are

known to infiltrate the γ-KC axon bundle at the onset of pruning, likely assisting in its

defasciculation (Marmor-Kollet et al., 2023), and later engulf the axonal debris (Hakim et

al., 2014; Tasdemir-Yilmaz & Freeman, 2014). Thus, the pruning defects caused by MMP

KD in γ-KCs, as well as non-cell autonomously by KD in glia (including the astrocytes

themselves), may be due to failure of the astrocytes to infiltrate and defasciculate the

axonal bundle. MMPs were also shown to degrade adhesion molecules (Page-McCaw et

al., 2007), and our lab previously showed that adhesion molecules – specifically Fasciclin

II (FasII) – must be downregulated for pruning to occur properly (Bornstein et al., 2015).

It is thus possible that MMPs cleave FasII, or other adhesion molecules, to promote axon

pruning. Since most *Drosophila* MMP isoforms are secreted (LaFever et al., 2017),

together with the finding that they are required from both KC and glia origin, it is possible

that MMPs are also secreted from additional cells within or around the MB circuit. All the

above-mentioned options are promising directions for future research, in hopes of

uncovering the mechanisms by which MMPs promote neuronal remodeling.

Notably, the main hypothesis in the literature is that SCZ is associated with increased

synapse pruning, while in our study knockdown of SCZ-associated genes actually

resulted in inhibition of pruning. However, SNPs can result in loss- but also gain-of-

function, which could account for this apparent conflict. In fact, it was previously shown

that the SCZ-related SNP in Mmp9 is in the 3' UTR, and affects Mmp9 mRNA folding

which eventually results in higher Mmp9 expression compared to WT (Lepeta et al.,

2017).

Understating the neurodevelopmental function of MMPs may provide insights into the

changes that the ECM in the brain undergoes throughout development, and specifically

during neuronal remodeling. The ECM consists of different components that allow efficient

cell-cell communication as well as cell-ECM communication (Miller et al., 2020). It is

therefore likely that MMPs, secreted from both glia and neurons, have a profound effect

on the ECM during development, and thus have a pivotal role in the coordination of

developmental processes and the etiology of neuropsychiatric disorders.

Materials and methods

Gene selection for the screen

Human genes that contain SNPs that are associated with SCZ were collected from two

studies: PFC2 & CLOZUK (Wu et al., 2017; Supplemental excel 'screen data' sheet

'Human SCZ SNPs'). These genes were to their *Drosophila melanogaster* homologs

converted using HumanMine (Supplemental excel 'screen data' sheet 'Fly homologs to

human genes'). In cases of multiple fly homologs for a single human gene, we decided to

include all.

Drosophila melanogaster rearing and strains

All fly strains were reared under standard laboratory conditions at 25°C on molasses-

containing food. Males and females were chosen at random. Unless specifically stated

otherwise, the relevant developmental stage is adult, which refers to 3-5 days post eclosion. All RNAi lines for the screen were from the TRiP collection (Zirin et al., 2020) and obtained for the Bloomington *Drosophila* Stock Center (BDSC) – the full list is detailed in table 1 below. The following constructs were also obtained from BDSC: Luciferase RNAi (used as control for RNAi experiments; #35788), Mmp1 RNAi (#31489), Repo-Gal4 (#7415), OK107-Gal4 (#854) and QUAS-mtdTomato-3xHA (#30004). R71G10-QF2 was previously generated by our lab (Bornstein et al., 2021).

Gene	TRiP #	BDSC #
tartan	HM05011	28525
Fish-lips	HM05054	28568
retinal degeneration A	JF03371	29435
Ppr-Y	HMS00819	33882
FER tyrosine kinase	HMS00249	33375
Matrix metalloproteinase 2	HMJ23143	61309
Phospholipase C at 21C	HMS00436	32438
Proteasome alpha3 subunit	HMS05889	77145
Regulatory particle non-ATPase 7	HMS00096	34787
Syntaxin 13	HMS01723	38525
anterior pharynx defective 1	HMS01693	38249
BNIP3	HMJ02058	42494
liquid facets	HMJ22079	58130
eukaryotic translation release factor 1	HMS05812	67900
DIP-kappa	HM04050	31740
MYPT-75D	HMS01166	34688
Protein phosphatase 19C	HMS01841	38372
GTPase activating protein and centrosome-associated	HMS01132	34976
Ran-binding protein M	HMC05142	61172
Na[+]-driven anion exchanger 1	HMC05184	62177
minidiscs	HMC05214	62207
Serine-arginine protein kinase at 79D	HMC04154	55881
Ypel	HM05209	29530
Akt1	HMS00007	33615
absent, small, or homeotic discs 2	HMC05816	64942
Fmr1	HMS00248	34944
Arginine methyltransferase 1	GL01072	36891
Ge-1	HMS00340	32349
crooked legs	HMS02202	41669

Table 1. RNAi lines used in the screen

Immunohistochemistry and imaging

Drosophila brains were dissected in cold ringer solution and placed on ice. Followed by fixation in 4% paraformaldehyde (PFA) solution for 20 min in room temperature (RT). Fixed brains were then washed with PB supplemented with 0.3% Triton X-(PBT) – 3 times,

20 min each. Next the brains were blocked for non-specific staining using 5% heat
inactivated normal goat serum (NGS). Antibody staining was performed as follows:
primary antibodies (4°C, overnight), PBT washes (X3 in RT, 20 min each), secondary
antibodies (RT, 2h), PBT washes (X3 in RT, 20 min each), mounting in SlowFade
(Invitrogen). Primary antibodies included mouse anti-FasII 1:25 (1D4; DSHB), mouse
anti-EcRB1 1:25 (AD4.4; DSHB) and rat anti-HA 1:250 (Sigma Aldrich, 11867423001).
Secondary antibodies included Alexa fluor 647 goat anti-mouse 1:300 (A-21236;
Invitrogen) and Alexa fluor 568 goat anti-rat 1:300 (A-21247; Invitrogen). All brains were
imaged on a Zeiss LSM980 confocal microscope. The images were processed using
ImageJ 2.14.0 (NIH).

Quantification and statistical analysis
Blind ranking of the pruning phenotypes for Fig. 3-5 was performed by an independent
scorer who reviewed the Z-projection. Ranking scores were as follows: 1 = WT-like
phenotype, 2 = very mild pruning defect (PD), 3 = mild PD, 4 = moderate PD, 5 = severe
PD.
A non-parametric statistical analysis was preformed: Kruskal Wallis H test followed by a
post hoc Mann Whitney U test with FDR correction.
*** = p value<0.001, ** = p value<0.01, * = p value<0.05, N.S = not significant. Specific p
values are indicated in the figure legends.
309

Acknowledgements

We thank Andrea Page-McCaw (Vanderbilt University) for kindly sharing reagents, and
the Bloomington Drosophila Stock Center for reagents. Monoclonal antibodies were
obtained from the Developmental Studies Hybridoma Bank developed under the auspices
of the NICHD and maintained by the University of Iowa. We thank R. Rothkopf for
assistance with statistics. This work was supported by by European Research
Council (ERC) advanced grant #101054886 "NeuRemodelBehavior," and by the ISF-
NSFC joint research program (grant no. 2573/18). O.S is an incumbent of the Prof. Erwin
Netter Professorial Chair of Cell Biology.

Author contributions

320 S.K. conceptualized the project, designed, performed and analyzed experiments,
321 performed bioinformatic and statistical analyses, and wrote the manuscript. H.M.
322 designed and performed experiments, interpreted results, and wrote the manuscript. N.M-
323 K. designed experiments and assisted with bioinformatic analyses. O.S. led the project,
324 designed experiments, interpreted results, wrote the manuscript, and procured funding.

Declaration of interests

326 The authors declare that they have no conflict of interest.

A

Larva | Pupa (18h) | Pupa (24h) | Adult

Kenyon Cells

Calyx

γ

γ pruning

Astrocytes

γ regrowth

α' α

γ

β'

β

B

339
Human SNPs associated with schizophrenia (GWAS)

257
Drosophila homolog genes

195
Expressed in γ-KCs

32
Upregulated in γ-KCs during pruning

29
KD screen

C

Genes Dynamically expressed in γ-KCs

hours after pupa formation

L2 L3 0h ⎯ every 3h ⎯ 30h Adult

II

III

IV

V

VI

VII

Min   Max

Figure 1. *Drosophila* homologs of SCZ-related genes are dynamically expressed in developing γ-KCs.

(A) Schematic representation of *Drosophila* MB development. During early pupal stages, the γ-KCs (green) prune their dendrites completely and their vertical and medial axonal branches up to their branch-point. Subsequently, they reform dendrites in the calyx and regrow axons to form the adult-specific medial lobe.

(B) A schematic representation describing the logic for selection of candidates for the loss-of-function screen.

(C) A heatmap representing the expression of dynamically expressed genes along γ-KC development (based on Alyagor et al., 2018). Each row represents a gene, with red and blue indicating high and low relative expression, respectively. The genes in clusters III-VI were selected as candidates for screening.

KC-Gal4> RNAi

Control
R71G10>mtdT
FasII
9/12 20μm

Mmp2
7/10

Plc21C
7/10

Syx13
9/10

Pp4-19C
13/13

RanBPM
6/6

Ypel
8/11

Ge-1
9/9

crol
8/12

BNIP3
8/8

Abnormal MB phenotype
mild pruning defect
WT-like phenotype 13
pruning defect 9
Lethal 4
1
2

Fig 2. *Drosophila* homologs of SCZ-associated genes are required for γ-axon pruning.

(A-J) Confocal z-projections of brains in which the indicated UAS-RNAi transgenes were expressed in all KCs using the strong driver OK107-Gal4, while R71G10-QF2-drives the expression of membranal tandem tomato (mtdT; green) in γ-KCs and stochastically in α/β-KCs. The RNAi transgenes target the following genes: Matrix Metalloproteinase 2 (Mmp2; B), Phospholipase C at 21C (Plc21C; B), Syntaxin 13 (Syx13; D), Protein phosphatase 19C (Pp4-19C; E), Ran-binding protein M (RanBPM; F), Yippee-like (Ypel; G), Ge-1 (Ge-1; H), crooked legs (crol; I), BCL2 Interacting Protein 3 (BNIP3; J). Control is RNAi targeting Luciferase (A). FasII antibody (magenta) strongly labels α/β-KCs and weakly labels γ-KCs.

Arrows indicate unpruned axons. Scale bar corresponds to 20µm. The number of MBs (each from an individual brain) showing the presented phenotype out of the total n for each genotype is indicated.

(K) Pie chart representing the overall screen results.

A Expression in larval vs. adult astrocytes

-log10(p. adj) vs log2 Fold Change (L3/Adult)

Syx13
Mmp2

B Control
C Mmp2-RNAi

Glia-Gal4 > RNAi

R71G10>mtdT
FasII

20μm

D

Control
Mmp2-RNAi

0%   50%   100%

***

WT-like
Very mild
Mild
Moderate
Severe

Fig. 3. Mmp2 is upregulated in larval astrocytes and required in glia for γ-axon

pruning.

(A) Volcano plot representing the expression of candidates in astrocytes (based on

Marmor-Kollet et al., 2023). The x-axis represents log2 fold change of 3^{rd} instar larvae

(L3) vs. adult astrocytes. The y-axis represents the significance of the change (-log10 of

p. adjust). The annotated genes are upregulated in larval compared to adult astrocytes,

as well as required in KCs for γ-axon pruning (See Fig. 2 and Supplemental excel 'screen

data' sheet 'Astrocytes - genes exp. data').

(B-C) Confocal z-projections of control (B) and Mmp2-RNAi (C) driven by the pan-glial

Repo-Gal4 while R71G10-QF2 drives expression of mtdT (green) to label the γ-KCs plus

some of the α/β-KCs.

FasII antibody (magenta) strongly labels α/β-KCs and weakly labels γ-KCs.

Arrows indicate unpruned axons. Scale bar corresponds to 20μm.

(D) Quantification of the pruning defect phenotypes, ranging from 1 (WT-like phenotype)

to 5 (severe pruning defect). *** p=0.0002

Fig. 4. **Mmp1 is required in glia, and to a lesser extent in KCs, for γ-axon pruning.**

(A-B, D-E) Confocal z-projections of adult MBs in which γ-KCs are labeled by R71G10-

QF2-driven QUAS-mtdT (green), and control or Mmp1-RNAi is expressed in all glia using

Repo-Gal4 (A, B), or in all KCs using OK107-Gal4 (D, E).

FasII antibody (magenta) strongly labels α/β-KCs and weakly labels γ-KCs.

Arrows indicate unpruned axons. Scale bar corresponds to 20μm.

(C, F) Quantification of the pruning defect phenotypes, ranging from 1 (WT-like

phenotype) to 5 (severe pruning defect). ** $p=0.002$, n.s = not significant.

Fig. 5. While axon pruning of γ-KCs relies on both glial and neuronal MMPs, the

main requirement is glial Mmp2.

(A-C, E-G, I-K) Confocal z-projections of adult MBs in which γ-KCs are labeled by

R71G10-QF2-driven QUAS-mtdT (green), and UAS-RNAi species targeting MMP1 (A, E,

I), MMP2 (B, F J), or both MMPs (C, G, K) driven by Repo-Gal4 (A-C), OK107-Gal4 (E-

G), or both Gal4s simultaneously (I-K). FasII antibody (magenta) strongly labels α/β-KCs

and weakly labels γ-KCs. Arrows indicate unpruned axons.

(D, H, L-O) Quantification of the pruning defect phenotypes, ranging from 1 (WT-like

phenotype) to 5 (severe pruning defect). Genotypes are indicated with the letter of the

corresponding image within this figure. *** = $p < 0.001$, * = $p < 0.05$, n.s = not significant.

A vs. B $p = 0.019$, B vs. C $p = 0.814$, E vs. F $p = 0.041$, F vs. G $p = 0.659$, I vs. J $p = 0.019$, J

vs. K $p = 0.905$, A vs. E $p = 0.0006$, A vs. I $p = 0.088$, B vs. F $p = 0.0007$, B vs. J $p = 0.809$, C

vs. G $p = 0.0002$, C vs. K $p = 0.937$

References

Alyagor, I., Berkun, V., Keren-Shaul, H., Marmor-Kollet, N., David, E., Mayseless, O., Issman-Zecharya, N., Amit, I., & Schuldiner, O. (2018). Combining Developmental and Perturbation-Seq Uncovers Transcriptional Modules Orchestrating Neuronal Remodeling. *Dev Cell*, *47*(1), 38-52 e36. https://doi.org/10.1016/j.devcel.2018.09.013

Awasaki, T., Huang, Y., O'Connor, M. B., & Lee, T. (2011). Glia instruct developmental neuronal remodeling through TGF-beta signaling. *Nat Neurosci*, *14*(7), 821-823. https://doi.org/10.1038/nn.2833

Awasaki, T., & Ito, K. (2004). Engulfing action of glial cells is required for programmed axon pruning during Drosophila metamorphosis. *Curr Biol*, *14*(8), 668-677. https://doi.org/10.1016/j.cub.2004.04.001

Bornstein, B., Meltzer, H., Adler, R., Alyagor, I., Berkun, V., Cummings, G., Reh, F., Keren-Shaul, H., David, E., Riemensperger, T., & Schuldiner, O. (2021). Transneuronal Dpr12/DIP-delta interactions facilitate compartmentalized dopaminergic innervation of Drosophila mushroom body axons. *EMBO J*, *40*(12), e105763. https://doi.org/10.15252/embj.2020105763

Bornstein, B., Zahavi, E. E., Gelley, S., Zoosman, M., Yaniv, S. P., Fuchs, O., Porat, Z., Perlson, E., & Schuldiner, O. (2015). Developmental Axon Pruning Requires Destabilization of Cell Adhesion by JNK Signaling. *Neuron*, *88*(5), 926-940. https://doi.org/10.1016/j.neuron.2015.10.023

Boulanger, A., Thinat, C., Züchner, S., Fradkin, L. G., Lortat-Jacob, H., & Dura, J.-M. (2021). Axonal chemokine-like Orion induces astrocyte infiltration and engulfment during mushroom body neuronal remodeling. *Nature Communications*, *12*(1), 1849. https://doi.org/10.1038/s41467-021-22054-x

Brand, A. H., & Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development*, *118*(2), 401-415. https://doi.org/10.1242/dev.118.2.401

Brent, B. K., Thermenos, H. W., Keshavan, M. S., & Seidman, L. J. (2013). Gray matter alterations in schizophrenia high-risk youth and early-onset schizophrenia: a review of structural MRI findings. *Child Adolesc Psychiatr Clin N Am*, *22*(4), 689-714. https://doi.org/10.1016/j.chc.2013.06.003

Broadie, K., Baumgartner, S., & Prokop, A. (2011). Extracellular matrix and its receptors in Drosophila neural development. *Dev Neurobiol*, *71*(11), 1102-1130. https://doi.org/10.1002/dneu.20935

Dear, M. L., Dani, N., Parkinson, W., Zhou, S., & Broadie, K. (2016). Two classes of matrix metalloproteinases reciprocally regulate synaptogenesis. *Development*, *143*(1), 75-87. https://doi.org/10.1242/dev.124461

Dickerson, F., Vaidya, D., Liu, Y., & Yolken, R. (2023). Levels of Matrix Metalloproteinase 9 Are Elevated in Persons With Schizophrenia or Bipolar Disorder: The Role of Modifiable Factors. *Biol Psychiatry Glob Open Sci*, *3*(4), 766-772. https://doi.org/10.1016/j.bpsgos.2023.02.007

Feinberg, I. (1982). Schizophrenia: caused by a fault in programmed synaptic elimination during adolescence? *J Psychiatr Res*, *17*(4), 319-334. https://doi.org/10.1016/0022-3956(82)90038-3

Freeman, M. R. (2015). Drosophila Central Nervous System Glia. *Cold Spring Harb Perspect Biol*, *7*(11). https://doi.org/10.1101/cshperspect.a020552

Furukubo-Tokunaga, K. (2009). Chapter 12 - Modeling schizophrenia in flies. In S. Akira (Ed.), *Progress in Brain Research* (Vol. 179, pp. 107-115). Elsevier. https://doi.org/https://doi.org/10.1016/S0079-6123(09)17912-8

Gore, S. V., James, E. J., Huang, L. C., Park, J. J., Berghella, A., Thompson, A. C., Cline, H. T., & Aizenman, C. D. (2021). Role of matrix metalloproteinase-9 in neurodevelopmental deficits and experience-dependent plasticity in Xenopus laevis. *Elife*, *10*. https://doi.org/10.7554/eLife.62147

Hakim, Y., Yaniv, S. P., & Schuldiner, O. (2014). Astrocytes play a key role in Drosophila mushroom body axon pruning. *PLoS One*, *9*(1), e86178. https://doi.org/10.1371/journal.pone.0086178

Han, C., Song, Y., Xiao, H., Wang, D., Franc, N. C., Jan, L. Y., & Jan, Y. N. (2014). Epidermal cells are the primary phagocytes in the fragmentation and clearance of degenerating dendrites in Drosophila. *Neuron*, *81*(3), 544-560. https://doi.org/10.1016/j.neuron.2013.11.021

Heisenberg, M. (1998). What do the mushroom bodies do for the insect brain? an introduction. *Learn Mem*, *5*(1-2), 1-10.

Hoffman, R. E., & Dobscha, S. K. (1989). Cortical pruning and the development of schizophrenia: a computer model. *Schizophr Bull*, *15*(3), 477-490. https://doi.org/10.1093/schbul/15.3.477

Huntley, G. W. (2012). Synaptic circuit remodelling by matrix metalloproteinases in health and disease. *Nat Rev Neurosci*, *13*(11), 743-757. https://doi.org/10.1038/nrn3320

Keshavan, M., Lizano, P., & Prasad, K. (2020). The synaptic pruning hypothesis of schizophrenia: promises and challenges. *World Psychiatry*, *19*(1), 110-111. https://doi.org/https://doi.org/10.1002/wps.20725

Keshavan, M. S., Anderson, S., & Pettergrew, J. W. (1994). Is Schizophrenia due to excessive synaptic pruning in the prefrontal cortex? The Feinberg hypothesis revisited. *Journal of Psychiatric Research*, *28*(3), 239-265. https://doi.org/https://doi.org/10.1016/0022-3956(94)90009-4

Kudo, N., Yamamori, H., Ishima, T., Nemoto, K., Yasuda, Y., Fujimoto, M., Azechi, H., Niitsu, T., Numata, S., Ikeda, M., Iyo, M., Ohmori, T., Fukunaga, M., Watanabe, Y., Hashimoto, K., & Hashimoto, R. (2020). Plasma levels of matrix metalloproteinase-9 (MMP-9) are associated with cognitive performance in patients with schizophrenia. *Neuropsychopharmacol Rep*, *40*(2), 150-156. https://doi.org/10.1002/npr2.12098

Kuo, C. T., Jan, L. Y., & Jan, Y. N. (2005). Dendrite-specific remodeling of <i>Drosophila</i> sensory neurons requires matrix metalloproteases, ubiquitin-proteasome, and ecdysone signaling. *Proceedings of the National Academy of Sciences*, *102*(42), 15230-15235. https://doi.org/doi:10.1073/pnas.0507393102

LaFever, K. S., Wang, X., Page-McCaw, P., Bhave, G., & Page-McCaw, A. (2017). Both Drosophila matrix metalloproteinases have released and membrane-tethered forms but have different substrates. *Scientific Reports*, *7*(1), 44560. https://doi.org/10.1038/srep44560

Laricchiuta, D., Papi, M., Decandia, D., Panuccio, A., Cutuli, D., Peciccia, M., Mazzeschi, C., & Petrosini, L. (2024). The role of glial cells in mental illness: a systematic review on astroglia and microglia as potential players in schizophrenia and its cognitive and emotional aspects. *Front Cell Neurosci*, *18*, 1358450. https://doi.org/10.3389/fncel.2024.1358450

Lee, T., Lee, A., & Luo, L. (1999). Development of the Drosophila mushroom bodies: sequential generation of three distinct types of neurons from a neuroblast. *Development*, *126*(18), 4065-4076. https://www.ncbi.nlm.nih.gov/pubmed/10457015

Lepeta, K., Purzycka, K. J., Pachulska-Wieczorek, K., Mitjans, M., Begemann, M., Vafadari, B., Bijata, K., Adamiak, R. W., Ehrenreich, H., Dziembowska, M., & Kaczmarek, L. (2017). A normal genetic variation modulates synaptic MMP-9 protein levels and the severity of schizophrenia symptoms. *EMBO Mol Med*, *9*(8), 1100-1116. https://doi.org/10.15252/emmm.201707723

Lessing, D., & Bonini, N. M. (2009). Maintaining the brain: insight into human neurodegeneration from Drosophila melanogaster mutants. *Nat Rev Genet*, *10*(6), 359-370. https://doi.org/10.1038/nrg2563

Llano, E., Adam, G., Pendás, A. M., Quesada, V. c., Sánchez, L. M., Santamaría, I., Noselli, S., & López-Otín, C. (2002). Structural and Enzymatic Characterization of Drosophila Dm2-MMP, a Membrane-bound Matrix Metalloproteinase with Tissue-specific Expression*. *Journal of Biological Chemistry*, *277*(26), 23321-23329. https://doi.org/https://doi.org/10.1074/jbc.M200121200

Luo, L., & O'Leary, D. D. (2005). Axon retraction and degeneration in development and disease. *Annu Rev Neurosci*, *28*, 127-156. https://doi.org/10.1146/annurev.neuro.28.061604.135632

Lyne, R., Bazaga, A., Butano, D., Contrino, S., Heimbach, J., Hu, F., Kalderimis, A., Lyne, M., Reierskog, K., Stepan, R., Sullivan, J., Wise, A., Yehudi, Y., & Micklem, G. (2022). HumanMine: advanced data searching, analysis and cross-species comparison. *Database*, *2022*, baac054. https://doi.org/10.1093/database/baac054

Marmor-Kollet, N., Berkun, V., Cummings, G., Keren-Shaul, H., David, E., Addadi, Y., & Schuldiner, O. (2023). Actin-dependent astrocytic infiltration is a key step for axon defasciculation during remodeling. *Cell Reports*, *42*(2), 112117. https://doi.org/https://doi.org/10.1016/j.celrep.2023.112117

McCutcheon, R. A., Reis Marques, T., & Howes, O. D. (2020). Schizophrenia-An Overview. *JAMA Psychiatry*, *77*(2), 201-210. https://doi.org/10.1001/jamapsychiatry.2019.3360

Meltzer, H., Marom, E., Alyagor, I., Mayseless, O., Berkun, V., Segal-Gilboa, N., Unger, T., Luginbuhl, D., & Schuldiner, O. (2019). Tissue-specific (ts)CRISPR as an efficient strategy for in vivo screening in Drosophila. *Nat Commun*, *10*(1), 2113. https://doi.org/10.1038/s41467-019-10140-0

Miller, A. E., Hu, P., & Barker, T. H. (2020). Feeling Things Out: Bidirectional Signaling of the Cell-ECM Interface, Implications in the Mechanobiology of Cell Spreading, Migration, Proliferation, and Differentiation. *Adv Healthc Mater*, *9*(8), e1901445. https://doi.org/10.1002/adhm.201901445

Notter, T. (2021). Astrocytes in schizophrenia. *Brain Neurosci Adv*, *5*, 23982128211009148. https://doi.org/10.1177/23982128211009148

Overall, C. M., & López-Otín, C. (2002). Strategies for MMP inhibition in cancer: innovations for the post-trial era. *Nat Rev Cancer*, *2*(9), 657-672. https://doi.org/10.1038/nrc884

Page-McCaw, A., Ewald, A. J., & Werb, Z. (2007). Matrix metalloproteinases and the regulation of tissue remodelling. *Nat Rev Mol Cell Biol*, *8*(3), 221-233. https://doi.org/10.1038/nrm2125

Penzes, P., Cahill, M. E., Jones, K. A., VanLeeuwen, J.-E., & Woolfrey, K. M. (2011). Dendritic spine pathology in neuropsychiatric disorders. *Nature Neuroscience*, *14*(3), 285-293. https://doi.org/10.1038/nn.2741

Pijet, B., Stefaniuk, M., & Kaczmarek, L. (2019). MMP-9 Contributes to Dendritic Spine Remodeling Following Traumatic Brain Injury. *Neural Plast*, *2019*, 3259295. https://doi.org/10.1155/2019/3259295

Riabinina, O., & Potter, C. J. (2016). The Q-System: A Versatile Expression System for Drosophila. *Methods Mol Biol*, *1478*, 53-78. https://doi.org/10.1007/978-1-4939-6371-3_3

Sawamura, N., Ando, T., Maruyama, Y., Fujimuro, M., Mochizuki, H., Honjo, K., Shimoda, M., Toda, H., Sawamura-Yamamoto, T., Makuch, L. A., Hayashi, A., Ishizuka, K., Cascella, N. G., Kamiya, A., Ishida, N., Tomoda, T., Hai, T., Furukubo-Tokunaga, K., & Sawa, A. (2008). Nuclear DISC1 regulates CRE-mediated gene transcription and sleep homeostasis in the fruit fly. *Mol Psychiatry*, *13*(12), 1138-1148, 1069. https://doi.org/10.1038/mp.2008.101

Schuldiner, O., & Yaron, A. (2015). Mechanisms of developmental neurite pruning. *Cell Mol Life Sci*, *72*(1), 101-119. https://doi.org/10.1007/s00018-014-1729-6

Scott-Hewitt, N., Huang, Y., & Stevens, B. (2023). Convergent mechanisms of microglia-
mediated synaptic dysfunction contribute to diverse neuropathological conditions. *Annals*
of the New York Academy of Sciences, *1525*(1), 5-27.
https://doi.org/https://doi.org/10.1111/nyas.15010
Seitz-Holland, J., Seethaler, M., Makris, N., Rushmore, J., Cho, K. K., Rizzoni, E., Vangel, M.,
Sahin, O. S., Heller, C., Pasternak, O., Szczepankiewicz, F., Westin, C. F., Lošák, J.,
Ustohal, L., Tomandl, J., Vojtíšek, L., Kudlička, P., Jáni, M., Woo, T. W., . . . Kubicki, M.
(2022). The association of matrix metalloproteinase 9 (MMP9) with hippocampal volume
in schizophrenia: a preliminary MRI study. *Neuropsychopharmacology*, *47*(2), 524-530.
https://doi.org/10.1038/s41386-021-00997-5
Sekar, A., Bialas, A. R., de Rivera, H., Davis, A., Hammond, T. R., Kamitaki, N., Tooley, K.,
Presumey, J., Baum, M., Van Doren, V., Genovese, G., Rose, S. A., Handsaker, R. E.,
Schizophrenia Working Group of the Psychiatric Genomics, C., Daly, M. J., Carroll, M. C.,
Stevens, B., & McCarroll, S. A. (2016). Schizophrenia risk from complex variation of
complement component 4. *Nature*, *530*(7589), 177-183.
https://doi.org/10.1038/nature16549
Sellgren, C. M., Gracias, J., Watmuff, B., Biag, J. D., Thanos, J. M., Whittredge, P. B., Fu, T.,
Worringer, K., Brown, H. E., Wang, J., Kaykas, A., Karmacharya, R., Goold, C. P.,
Sheridan, S. D., & Perlis, R. H. (2019). Increased synapse elimination by microglia in
schizophrenia patient-derived models of synaptic pruning. *Nat Neurosci*, *22*(3), 374-385.
https://doi.org/10.1038/s41593-018-0334-7
Suzuki, M., Nohara, S., Hagino, H., Kurokawa, K., Yotsutsuji, T., Kawasaki, Y., Takahashi, T.,
Matsui, M., Watanabe, N., Seto, H., & Kurachi, M. (2002). Regional changes in brain gray
and white matter in patients with schizophrenia demonstrated with voxel-based analysis
of MRI. *Schizophrenia Research*, *55*(1), 41-54.
https://doi.org/https://doi.org/10.1016/S0920-9964(01)00224-9
Szklarczyk, A., Lapinska, J., Rylski, M., McKay, R. D., & Kaczmarek, L. (2002). Matrix
metalloproteinase-9 undergoes expression and activation during dendritic remodeling in
adult hippocampus. *J Neurosci*, *22*(3), 920-930. https://doi.org/10.1523/jneurosci.22-03-
00920.2002
Tasdemir-Yilmaz, O. E., & Freeman, M. R. (2014). Astrocytes engage unique molecular
programs to engulf pruned neuronal debris from distinct subsets of neurons. *Genes Dev*,
28(1), 20-33. https://doi.org/10.1101/gad.229518.113
Truman, J. W. (1990). Metamorphosis of the central nervous system of Drosophila. *J*
Neurobiol, *21*(7), 1072-1084. https://doi.org/10.1002/neu.480210711
Watts, R. J., Schuldiner, O., Perrino, J., Larsen, C., & Luo, L. (2004). Glia engulf degenerating
axons during developmental axon pruning. *Curr Biol*, *14*(8), 678-684.
https://doi.org/10.1016/j.cub.2004.03.035
Wu, Y., Yao, Y. G., & Luo, X. J. (2017). SZDB: A Database for Schizophrenia Genetic Research.
Schizophr Bull, *43*(2), 459-471. https://doi.org/10.1093/schbul/sbw102
Yaniv, S. P., & Schuldiner, O. (2016). A fly's view of neuronal remodeling. *Wiley Interdiscip Rev*
Dev Biol, *5*(5), 618-635. https://doi.org/10.1002/wdev.241
Yasunaga, K., Kanamori, T., Morikawa, R., Suzuki, E., & Emoto, K. (2010). Dendrite reshaping
of adult Drosophila sensory neurons requires matrix metalloproteinase-mediated
modification of the basement membranes. *Dev Cell*, *18*(4), 621-632.
https://doi.org/10.1016/j.devcel.2010.02.010
Yu, Q., & Stamenkovic, I. (2000). Cell surface-localized matrix metalloproteinase-9
proteolytically activates TGF-beta and promotes tumor invasion and angiogenesis. *Genes*
Dev, *14*(2), 163-176.
Yu, X. M., Gutman, I., Mosca, T. J., Iram, T., Ozkan, E., Garcia, K. C., Luo, L., & Schuldiner,
O. (2013). Plum, an immunoglobulin superfamily protein, regulates axon pruning by

facilitating TGF-beta signaling. *Neuron*, *78*(3), 456-468.
https://doi.org/10.1016/j.neuron.2013.03.004
Zheng, X., Wang, J., Haerry, T. E., Wu, A. Y., Martin, J., O'Connor, M. B., Lee, C. H., & Lee, T.
(2003). TGF-beta signaling activates steroid hormone receptor expression during
neuronal remodeling in the Drosophila brain. *Cell*, *112*(3), 303-315.
https://doi.org/10.1016/s0092-8674(03)00072-2
Zikidi, K., Gajwani, R., Gross, J., Gumley, A. I., Lawrie, S. M., Schwannauer, M., Schultze-
Lutter, F., Fracasso, A., & Uhlhaas, P. J. (2020). Grey-matter abnormalities in clinical high-
risk participants for psychosis. *Schizophr Res*, *226*, 120-128.
https://doi.org/10.1016/j.schres.2019.08.034
Zirin, J., Hu, Y., Liu, L., Yang-Zhou, D., Colbeth, R., Yan, D., Ewen-Campen, B., Tao, R., Vogt,
E., VanNest, S., Cavers, C., Villalta, C., Comjean, A., Sun, J., Wang, X., Jia, Y., Zhu, R.,
Peng, P., Yu, J., . . . Perrimon, N. (2020). Large-Scale Transgenic Drosophila Resource
Collections for Loss- and Gain-of-Function Studies. *Genetics*, *214*(4), 755-767.
https://doi.org/10.1534/genetics.119.302964
613