Decreased hippocampal neurite density in middle-aged/aging adults following prenatal exposure to higher levels of maternal inflammation

Raana A. Mohyee*[1], Blake L. Elliott*[1], Madeline R. Pike[1], Emma Smith[1], Ann M. Kring[4], Ingrid R. Olson[1], Elizabeth C. Breen [2], Barbara A. Cohn[3], Piera M. Cirillo[3], Nickilou Y. Krigbaum[3], Thomas M. Olino[1], Mark D'Esposito[4], Ashby B. Cogan[4], Bhakti P. Patwardan[4], Lauren M. Ellman[1]

[1]Department of Psychology & Neuroscience, Temple University, [2]Cousins Center for Psychoneuroimmunology, University of California, Los Angeles, [3]Child Health and Development Studies, Public Health Institute, [4]Department of Psychology, University of California, Berkeley

*Denotes shared first authorship

Corresponding author: Lauren M. Ellman, Ph.D.

Email: ellman@temple.edu

Author Contributions: Raana Mohyee and Dr. Blake Elliott are co-first authors of this manuscript, including conducting all the imaging analyses, conceiving of the study aims, writing initial drafts of the manuscript, and contributing to data analyses. Madeline Pike contributed to writing, editing and overseeing analyses as a PNMI consultant. Emma Smith contributed to writing, editing and providing feedback as a PNMI consultant. Dr. Lauren Ellman wrote and was Principal Investigator of the grant that the analyses were based on, she contributed to the conception of the analyses, she provided feedback at each stage of manuscript preparation, and she provided edits for the manuscript. Dr. Ingrid Olson provided feedback at each stage of manuscript preparation, helped with the design and methods of the study, and she provided edits for the manuscript, specifically as they related to the neuroimaging data analytic plan and methods. Dr. Ann Kring was integral in the conception, design, and methods of the study, oversaw all data collection as the site PI, and provided feedback and edits at the final stage of manuscript preparation. Dr. Elizabeth Breen was the primary immunologist on the study and contributed to the conception of the design of the study and to decisions about methods for serological analyses, oversaw all cytokine analyses, and provided feedback and edits at the final stage of manuscript preparation. Barbara Cohn, Piera Cirillo, and Nickilou Krigbaum were integral in the design and conception of the original study, oversee the CHDS cohort and were integral in participant recruitment and selection, provided epidemiological consultant throughout the study and during manuscript development, and provided feedback and edits at the final stage of manuscript preparation. Dr. Thomas Olino contributed to edits and statistical consultation. Mark T. D'Esposito[4] contributed to the methods and design for neuroimaging collection and provided edits at the final stage of manuscript preparation. Ashby B. Cogan[4] and Bhakti P. Patwardan[4] completed the data collection.

Competing Interest Statement: The authors do not have any competing interests to disclose.

Classification: Neuroscience; Immunology and Inflammation

Keywords: prenatal maternal inflammation, hippocampal neurite density, middle-aged adults

This PDF file includes:

> Main Text
> Figures 1 to 2
> Tables 1 to 5

Abstract

In animal models, exposure to heightened maternal inflammation in utero is associated with altered offspring hippocampal development, including reduced dendritic arborization and density. However, the effects of prenatal maternal inflammation (PNMI) on offspring hippocampal microstructure in humans remains unclear. Here, we examined the relationship between exposure to PNMI and neurite density in the hippocampus and its subfields among offspring during late middle age. Participants included 72 mother-offspring dyads from the Child Health and Development Studies (CHDS) cohort. Data for four inflammatory biomarkers (IL-6, IL-8, IL-1 receptor antagonist [IL-1RA], and soluble TNF receptor-II [sTNF-RII]) were available from first and second trimester maternal sera. Neurite density in the offspring hippocampus and its subfields was estimated using microstructural modeling of offsprings' diffusion-weighted Magnetic Resonance Imaging data (mean age of offspring at imaging = 59 years; 51% male). We estimated the relationship between each biomarker and region-of-interest's neurite density. Higher first trimester maternal IL-1RA and IL-6 levels were associated with lower offspring hippocampal neurite density. These relationships were specific to the CA3, CA4, dentate gyrus, and subiculum subfields. In addition, higher second trimester IL-6 was associated with lower subiculum neurite density. Our findings reveal that exposure to heightened prenatal levels of maternal inflammation is linked to altered offspring hippocampal microstructure in late middle age, which could have implications for memory decreases during this period and may be relevant for understanding risk of aging-related cognitive changes.

Significance Statement

The contribution of prenatal maternal inflammation (PNMI) to offspring brain microstructure in later life is well established in animal models, but poorly understood in humans. Our study discovered long-lasting impacts of elevated PNMI during early-mid gestation on the structural integrity of the hippocampus in offspring during late middle age. Our findings underscore the potential role of

developmental programming in aging-related neurological and cognitive decline, as the observed degradation in hippocampal microstructure is present over half a century following exposure.

Main Text

Introduction

During fetal development, a cascade of biological events results in the formation of distinct cortical and subcortical brain structures, which have been shown to be influenced by the intrauterine environment (1–4). Evidence from human epidemiological studies has linked prenatal maternal inflammation (PNMI) to offspring neuropsychiatric and neurodevelopmental disorders (5). For example, studies have associated PNMI with increased prevalence of autism spectrum disorder (6–9), psychotic disorders (10–12), depression (13), and childhood internalization and externalization symptoms (14,15).

Importantly, alterations in the structure and function of the hippocampus, a subcortical region integral to learning and memory, emotion, and regulation of physiological stress responses (16–22), have been implicated in the etiology and course of many of these psychiatric conditions, as well as types of dementia including Alzheimer's disease (23–25). The hippocampus, which completes much of its cytoarchitectural maturation and subfield differentiation during the second and third trimesters of pregnancy, is particularly sensitive to proinflammatory exposures during the fetal period (26–28). The hippocampus contains some of the highest densities of microglia and several proinflammatory marker receptors in the brain, especially the interleukin-1β (IL-1β) receptor (26,29). As such, the hippocampus serves as a prime avenue for exploring the longevity of the developmental outcomes associated with prenatal exposure to heightened maternal inflammation (30,31).

The preclinical literature has identified several mechanisms by which the microstructure of the developing hippocampus is altered by exposure to heightened maternal immune activation, including reduced neurogenesis (32), reduced hippocampal expression of several key neuromodulatory neurotransmitters (33), and altered glial development (34). One clinical study that supports these findings observed that prenatal exposure to increased maternal interleukin-8 (IL-8) in the second and third trimesters was related to decreases in left entorhinal cortex volume, which directs input to and output from the hippocampus, in offspring diagnosed with schizophrenia spectrum disorders compared with healthy controls (35).

In addition, both clinical and preclinical studies demonstrate the

importance of timing of PNMI exposure in relation to offspring neurodevelopment, though findings differ somewhat across species. For example, epidemiological studies have linked both first and the second-trimester PNMI to increased risk of offspring depression and schizophrenia (15,36), and at least one rodent study has linked late gestational (corresponding to the second trimester in humans) exposure to PNMI to behavioral deficits mirroring negative symptoms of schizophrenia, as well as altered dopaminergic, GABAergic, and glutamatergic neurotransmission in the hippocampus (33). Another rodent study found that key cellular processes affecting neuronal architecture, including neurogenesis, reelin immunoreactivity, and apoptosis, were more drastically altered in offspring following PNMI during middle gestation (corresponding to the first trimester in humans) than in late gestation, with various behavioral deficits observed following exposure at both timepoints (36). In the present study we will investigate timing effects by examining PNMI during both the first and second trimesters.

       Along with timing of PNMI exposure, preclinical studies suggest that offspring age at the time of neurological measurement provides another source of variation in existing preclinical findings on PNMI and hippocampal development, although this has yet to be explored in human research. For instance, PNMI has been shown to impair hippocampal neurogenesis in juvenile rodents; yet, changes in neurogenesis observed in adulthood, as a result of PNMI, appear to be spatially restricted and sex-specific (31,37). Furthermore, PNMI-related changes in offspring expression of several immunogenic chemosignals, both peripheral and intracranial, have been observed in proximal stages of development, but surprisingly *not* in adulthood (38).

       To our knowledge, very few human studies have investigated the effects of PNMI on hippocampus microarchitecture past the neonatal period, giving rise to questions regarding the translation of preclinical findings across the human lifespan(36). Moreover, existing human studies have either depended on indirect measures of PNMI (39), utilized postmortem methods (36), or used nonspecific metrics of structural change, like total volume or diffusion tensor-derived metrics (35,40). Furthermore, no human studies (to our knowledge) have investigated the effects of PNMI on the hippocampus in late middle age in a population-based sample unselected for psychiatric outcomes (36). Determining the association between PNMI and offspring hippocampal microstructure in later stages of human development is crucial given the relevance for aging-related memory decline and neurodegenerative diseases in this age group (41).

       We also sought to investigate PNMI's effects on cytoarchitecture in the hippocampal subfields, each of which has unique histological and functional characteristics (42). Animal studies have shown that PNMI decreases neuron

density and the number of pyramidal cells in CA1, CA2, and CA3 (43,44) reduces CA1 axonal size and myelin thickness (34), and impairs cell proliferation and neurogenesis in the dentate gyrus, leading to memory deficits (45–47). Furthermore, PNMI results in reduced CA3 fiber density (48), aberrant CA3-to-CA1 synaptic transmission (49), and altered excitatory postsynaptic potentials in CA1 pyramidal neurons, resulting in abnormal object information processing (50). These findings highlight the need to investigate whether PNMI differentially affects specific hippocampal subfields to better understand its impact on neuronal functioning and behavior.

Consequently, in a sample of late middle age human adults who have been followed since the fetal period as part of a longitudinal cohort, we investigated the association between PNMI on adult offspring's hippocampal microstructure using noninvasive, *in vivo* brain imaging and novel microstructural modeling techniques. The Neurite Orientation Dispersion and Density Imaging (NODDI) model uses diffusion magnetic resonance imaging (MRI) to measure density and spatial distribution of the orientation of neurites (e.g., dendrites, axons, and their branches) (51). The NODDI model was developed to overcome challenges in measuring the brain microstructure, lending itself to the study of the hippocampus particularly well. We hypothesized that heightened PNMI would be associated with reduced neurite density in the hippocampus of adult offspring. Additionally, we hypothesized that these effects would be specific to the CA1, CA3, and Dentate Gyrus.

Results

Descriptives and Bivariate Correlations

Demographic information is presented in Table 1, and bivariate correlations for first and second trimester variables are reported in Tables 2 and 3. Maternal education and intracranial volume were not significantly correlated with neurite density, nor with any prenatal biomarkers at T1 or T2. Offspring sex (female) was negatively correlated with CA1 and CA3 neurite density. Offspring age at time of imaging was negatively correlated with maternal serum IL-8 at T1.

Prenatal Maternal Inflammation and Hippocampal Neurite Density

We next conducted analyses to determine whether individual differences in hippocampal microstructure (neurite density) were related to maternal inflammatory biomarkers using hierarchical multiple regression. Parameter estimates can be found in Table 4. At T1, higher maternal IL-1RA and IL-6 were

associated with lower offspring neurite density in the hippocampus. At T2, we found no significant associations between any biomarker and the whole hippocampus.

We next investigated whether the relationship between PNMI and hippocampal neurite density was specific to particular hippocampal subfields (Figure 2B) using hierarchical multiple regression. Parameter estimates can be found in Table 5. When individual hippocampal subfields were analyzed, at T1 both IL-1RA (Figure 2C) and IL-6 (Figure 2E) were associated with lower offspring neurite density in the CA3 and dentate gyrus subfields. Additionally, higher IL-1RA was associated with lower subiculum subfield neurite density and higher IL-6 was associated with lower CA4 subfield neurite density. There were no significant correlations between any biomarker and the CA1 subfield. At T2, we found that IL-6 was associated with lower offspring neurite density in the subiculum. No other associations were observed between any biomarker and any hippocampal subfield at T2 (Figure 2D, F).

Discussion

To our knowledge, this is the first human study to find an association between exposure to heightened prenatal inflammation and offspring neurite density in the hippocampus and its subfields during late middle age. We found that higher levels of maternal inflammatory biomarkers, specifically IL-1RA and IL-6 during the first trimester, were associated with reduced neurite density in the hippocampus and certain hippocampal subfields, including CA3, CA4, dentate gyrus, and subiculum. Interestingly, no significant relationship was observed in the CA1 subfield, and only IL-6 from the second trimester was associated with the subiculum subfield. These results indicate that PNMI is associated with reduced neurite density in the hippocampus and that these effects depend on the timing of exposure, are distributed differentially across subfields of the hippocampus, and are evident in late middle age. These findings are consistent with previous studies that demonstrate microstructural alterations in the hippocampus of nonhuman animals following PNMI (31,37,52).

Our results implicate heightened prenatal exposure to IL-1β (via measurement of IL-1RA), in altered development of the offspring hippocampus. As noted earlier, research has shown that IL-1RA is a strong, valid, and reliable estimate of IL-1β activity that is present at much higher concentrations in sera than the cytokine itself in pregnancy, which is necessary for improving detectable values of cytokine activity in the prenatal period (14,15,53). Extensive evidence from animal models of maternal prenatal immune activation have demonstrated reduced hippocampal neurogenesis in exposed offspring (31). An *in vitro* study of

human hippocampal cells identified upregulation of kynurenine production and gene expression for several related neurotoxic enzymes following experimental exposure to heightened IL-1β, demonstrating at least one mechanism of action through which the observed effects on hippocampal neurite density may have been borne out by heightened exposure to prenatal, maternal IL-1β (54). Furthermore, IL-1β is an integral molecular mediator of the cellular processes in the hippocampus that support long-term memory consolidation, the expression of which in the healthy hippocampus is tightly regulated (26,55). Importantly, blocking IL-1β signaling in models of Alzheimer's Disease reduces tau pathology, microglial activation, and secretion of cytokines by astrocytes (55).

Additionally, in vivo human studies have shown IL-1β linked to cognitive abilities. In a prospective birth cohort in the Southeastern United States, heightened maternal IL-1β during pregnancy was negatively associated with several cognitive abilities including overall intelligence and nonverbal and spatial abilities [measured with the Differential Abilities Scale [DAS-II]; (56)], in offspring in early childhood (57). The hippocampus may play a crucial role in these abilities. Traditionally the hippocampus has been shown to be critical in the formation of episodic memories, spatial memories, and the ability to navigate through space (16,58). However, the hippocampus has now been shown to be crucial for many broader aspects of cognition, such as perception, working memory, semantic search, concept formation, future planning, and social behavior (59–67). While IQ and broader cognitive abilities such as nonverbal reasoning engage multiple brain regions, including the hippocampus, our results suggest that the observed associations may be driven, at least partially, by the hippocampal contribution to these specific domains. Together, these findings suggest that neurite density is sensitive to cytokine-specific effects on human hippocampal cytoarchitecture and that these microstructural changes observed in the hippocampus in response to heightened IL-1β expression could drive functional impairments related to learning and memory at both early and later stages of development.

The significant relationship between neurite density in offspring hippocampi and maternal IL-6 from both trimesters (though evidence in the second trimester was only associated with the subiculum subfield) is consistent with the extensive preclinical findings implicating IL-6 in hippocampal development and function. Similar to IL-1β, IL-6 has been associated with the maintenance of long-term potentiation, a cellular process key to the consolidation of long-term memory(26), and appears to be a key molecular mediator of symptoms of neuropsychiatric disorders in which hippocampal abnormalities have been regularly observed (68,69). Elevated maternal prenatal IL-6 in rodents has been shown to upregulate glutamate receptor expression, associated with a

significant increase in excitatory postsynaptic currents, in the hippocampus of offspring, causing an imbalance of excitatory and inhibitory activity in the region (70). This imbalance likely has downstream effects on the hippocampus. Exposure to heightened maternal IL-6 in utero has been linked to several subfield-specific microstructural changes in offspring hippocampi, including decreased cell proliferation and neuronal survival in the granule cell layer of the dentate gyrus, neuronal loss in the hilus region of the dentate gyrus, and decreased neuronal density in the CA1 subfield of female offspring and in CA2 and CA3 subfields of the male offspring (31). As such, our findings with IL-6 confirm not only the importance of the cytokine in the developing hippocampus, but also highlight the potential regional specificity of subfields of the hippocampus most heavily involved in learning and memory function, namely the dentate gyrus and CA3 subfields.

Previous research in humans has shown that both prenatal IL-1ra and IL-6 are related to offspring depression (15). The hippocampus could be a critical region linking PNMI and offspring depression. Major depressive disorder (MDD) is clinically recognized as a stress sensitive disorder, and the hippocampus is highly sensitive to stress (24,71–73). Additionally, the greatest cognitive impairments observed in MDD are memory impairments associated with hippocampal function (74). The current finding of increased prenatal IL-1RA and IL-6 associated with decreased offspring hippocampal neurite density (particularly in CA3 and dentate gyrus regions) could be critical for understanding the neurobiological mechanisms contributing to depression and memory disorders.

The dentate gyrus and CA3 subfields, both of which have been shown to be particularly susceptible to microstructural changes following PNMI in the present study, are critical for memory function. The dentate gyrus, which receives dense input from the entorhinal cortex via the perforant pathway, and connects with the CA3 subfield via the mossy fiber pathway. The CA3 subfield has reciprocal connections back to the dentate gyrus, which are critical for memory function (75). The dentate gyrus is thought to play a central role in pattern separation, a process that distinguishes and decorrelates incoming input patterns from the entorhinal cortex, ensuring the encoding of distinct memories. The CA3 region is believed to contribute to pattern completion, aiding in the retrieval of memories by associating and reinstating stored information (76–82). APOE ε4, the best-known genetic risk factor for Alzheimer's Disease, has been shown to have a selective effect on the dentate gyrus and CA3 subfields (83,84). Importantly, our sample ranged from 57 to 63 years old, which is when the first signs of cognitive decline start to emerge (85). Thus, this study's findings indicating a strong association between heightened PNMI and reduced neurite

density in the DG and CA3 subfields offer valuable insights into the potential long-term consequences of maternal prenatal inflammation on cognitive health in aging and neurodegenerative disease risk, emphasizing the need for further translational research into the underlying mechanisms driving these specific associations.

In this sample, we found no significant associations with hippocampal or subfield neurite density with measures of sTNF-RII from either trimester. sTNF-RII is a valid and reliable estimate of TNF-α activity in pregnancy. TNF-α plays an important role in modulating hippocampal-dependent cognitive functions, including sleep and memory function (55). Furthermore, TNF-α demonstrably alters hippocampal microstructure through activity-dependent regulation of hippocampal synaptic expression of AMPA and GABA receptors in the hippocampus (26,86), and heightened expression of TNF-α has been shown to reduce cell proliferation and neuronal differentiation in the hippocampus (30). Even so, the robust relationship between TNF-α and hippocampal function has largely been characterized by the cross-sectional study of endogenously produced TNF-α in the brain, and current research findings suggest that this cytokine may not be essential to producing the behavioral deficits typically observed in maternal immune activation models (86). As such, more work is needed to definitively characterize the effects of maternal TNF-α expression on offspring hippocampal development.

Despite mixed evidence supporting different mechanisms of action for the translation of inflammation from mother to offspring in utero, findings from preclinical studies increasingly converge on the possibility of altered development of offspring microglia as a notable underlying mechanism driving the neuroimmunological changes observed in offspring as a consequent of PNMI (26,55,68). Some evidence also supports the idea that peripheral cytokines expressed by the mother can cross or upregulate cytokine production by the placenta, which then enter the fetal bloodstream (30,87–89); however, further human studies would be necessary to parse out specific mechanisms of action.

The prominence of first trimester exposures in the significant relationships observed between PNMI and hippocampal microstructure in this study and others points to the potential mediating role of offspring microglia. The first trimester is the period during which microglial populate the burgeoning central nervous system in the embryo, and evidence suggests that exposure of developing microglia to dysregulated cytokine expression results in the accelerated maturation and morphological activation of these cells, such that when exposed to a secondary immune insult, these cells mount stronger and more rapid cellular responses (26,90). It is possible that PNMI alters the susceptibility of microglia-mediated developmental processes in the

hippocampus, like decreased neurogenesis postnatally, resulting in lasting changes in hippocampal microstructure (26,55,91). Indeed, Graciarena et al. (92) found that the impaired neurogenesis in the dentate gyrus of adult mice prenatally exposed to maternal inflammation was associated with increased activation of microglia in particular. Thus, the long-term impact of PNMI on hippocampal microstructure observed in this study could be partially related to a cascade of alterations to neurodevelopment, with offspring microglia activation contributing to neurodevelopmental sequelae.

In addition to the potential mediating role of offspring microglia activation, a second alternative explanation for our results may be found in the placenta. Specifically, susceptibility of the placenta to maternal inflammation during pregnancy likely alters prenatal development through a number of physiological pathways independent of the offspring's immune system, including disruption of the flow of oxygen or nutrients to the fetus (93). Furthermore, maternal inflammation during pregnancy can succeed a number of lifestyle and environmental factors that alter offspring hippocampal development through several non-neurodevelopmental pathways (29). Thus, an alternative interpretation of this data could entertain PNMI as a proxy measure for other physiological and psychological insults experienced by the mother driving the observed changes in offspring hippocampal development through further alternative pathways.

Strengths, Limitations and Future Directions

This study's strengths include the use of a unique cohort with prenatal serum data *and* modern neuroimaging data collected during late middle age. The availability of trimester-specific, serologically defined biomarkers of PNMI enables an investigation of its systemic immunological effects. Despite the age of our samples, we successfully detected biomarkers in all cases and focused on analytes present at higher concentrations in pregnant serum. Additionally, our study implemented the innovative Neurite Orientation Dispersion and Density Imaging (NODDI) model with diffusion imaging. As demonstrated by Zhang et al. (2012), NODDI provides greater specificity than fractional anisotropy (FA) in diffusion imaging (51). Voxels with distinct NODDI parameters can produce identical FA values, making them indistinguishable using FA alone. Our use of this innovative technique offers a more detailed understanding of hippocampal structure in late middle age, providing insights into the intricate neural mechanisms affected by PNMI.

The current study has several limitations to consider. First, although this is one of the largest neuroimaging studies of a birth cohort of this kind, the sample

size still only allowed for modest power to detect significant associations and, therefore, examining interactions were not possible (e.g., sex differences in associations) (94).Second, it was beyond the scope of this study to investigate potential sources of maternal inflammation, such as infection (95), stress (96), obesity (97), nutrition (98), and/or toxic exposures (99) which could all exert additional independent and/or interactive teratogenic effects on brain development, limiting the conclusions about environmental factors leading to increased PNMI. Similarly, given the complex immunological changes experienced during each stage of pregnancy (100), the single measurements of maternal cytokine expression from each trimester offer limited insight into maternal immune function in this sample. Additionally, although we focused on one specific subcortical region, there could be widespread effects of PNMI throughout the cortex and subcortically. Although this is an interesting avenue for future research, controlling for multiple comparisons with this approach is a necessity, making this approach difficult in practice with the current sample size. Finally, degradation of the samples is always possible; however multiple steps were taken to ensure that degradation of samples did not influence cytokine analyses (see 15 for more details) and there is no reason to believe that degradation of samples would occur in a directional capacity to influence results.

Although the current study discovered later life impacts of elevated maternal prenatal inflammation on hippocampal cytoarchitecture, future PNMI research holds promise in several directions. Examining how PNMI interplays with hippocampal-dependent processes (e.g. episodic memory) can illuminate cognitive implications (101). Additionally, more fine-grained longitudinal studies across the lifespan can track developmental trajectories of the hippocampus and immune system of offspring together. Furthermore, given the differential effects of PNMI observed across male and female offspring in animal studies, an examination of sex differences in hippocampal microstructure in humans is warranted as well (31,33,102).

Conclusion

Our study represents a significant advancement in our understanding of the enduring effects of PNMI on offspring hippocampal microstructure in later life in humans. The large cohort, combined with the longitudinal nature of our study, with retention rates allowing for analysis decades after pregnancy, have illuminated effects of maternal prenatal inflammation on hippocampal microstructure in a period of late middle age (late 50's-early 60's) that represents a key time for early identification and intervention for cognitive decline. By revealing a specific link between heightened maternal prenatal inflammation and

alterations in offspring neurite density within key hippocampal subfields, our research highlights the potential implications of programming on aging-related neurological and cognitive decline. These insights underscore the importance of further investigation into the mechanisms underlying these associations and their broader implications for public health.

Materials and Methods

Participants

Participants were recruited from the Healthy Brains Project, a follow-up study of the Child Health and Development Studies (CHDS). CHDS is a prospective multi-generational cohort originally recruited for the purpose of studying health outcomes across the life span following prenatal exposure to environmental factors that affect human development (N = 19,044 live births) (103). This cohort enrolled pregnant women who were living in the greater Alameda County, California, United States area, between 1959 and 1966, and who received prenatal care from medical facilities within the Kaiser Foundation Health Plan.

Adult offspring who participated in the CHDS follow-up during adolescence (see 14 for details of this study), and for whom archived maternal serum samples from the first (T1) and/or second (T2) trimester had been previously analyzed for markers of inflammation (n=737), were invited to participate in the Healthy Brains Project adult follow up study. Recruitment followed a systematic, tiered sampling method and prioritized individuals who resided within close enough proximity to the neuroimaging facility (Berkeley, California) to participate in the neuroimaging portion of the study. Current analyses drew from the 114 individuals enrolled in the Healthy Brains Project cohort at the time of present analyses (circa January 2023). Of these 114 cohort members, 85 had completed the MRI portion of the study. Three participants from whom high-resolution MRI scans of the hippocampus, T1-weighted structural MRI scans, or diffusion-weighted MRI scans were not collected were excluded from analysis. An additional five participants whose MRI scans were deemed too low quality after visual inspection were excluded. From the remaining 77 participants, 5 were missing T1 sera data, and 8 were missing T2 sera data, leaving a final sample of 72 participants with T1 sera data and full MRI and 69 participants with T2 sera data and full MRI. 72 participants with MRI and T1 sera data, and 69 participants with MRI and T2 sera data were included in the final analyses (Figure 1).

The research protocol was approved by the Institutional Review Boards

(IRBs) at Temple University, University of California, Berkeley, and the Public Health Institute of California and all adult offspring enrolled in the Healthy Brains Project provided full informed written consent. However, the original CHDS predated IRBs; therefore mothers and offspring in the original CHDS voluntarily participated in in-person interviews and gave oral permission to researchers for medical record access for themselves and their children.

Demographics Variables

Detailed sociodemographic information was collected for all mothers during the second trimester of pregnancy (mean gestation = 15.3 weeks, SD 6.8 weeks). Maternal education was used as a proxy for postnatal advantages because it 1) was shown to be correlated with other measures of socioeconomic status (SES; e.g., annual household income) in similar published studies (14,104–106), 2) has been previously used in this way by multiple similar studies (107), 3) had the most complete data of the available SES measures in this dataset, and 4) accounts for variations in hippocampal functioning related to maternal education. Maternal education was categorized as "Did not complete High School," "Completed High School," or "Completed more than High School " and coded as a factor variable for analyses. Offspring race was categorized as "American Indian/Alaska Native", "Asian", "Black", and "White." Offspring sex assigned at birth (herein referred to as "sex") was categorized as "Male" or "Female." See Table 1 for more detailed demographic information.

Prenatal Maternal Inflammation (PNMI)

Prenatal Maternal Inflammation (PNMI) was quantified from archived prenatal first [T1; mean (SD) gestation = 11.21 (2.50) weeks] and second [T2; - mean (SD) gestation = 23.17 (2.37) weeks] trimester maternal serum as previously described in detail (see 15). PNMI biomarker assays were conducted using high-sensitivity (IL-6, IL-8) and regular sensitivity (sTNF-RII, IL-1RA) ELISAs (R&D Systems, Inc., Minneapolis, MN), and raw concentrations were reported in picograms per milliliter (pg/mL).

MRI Data acquisition

MRI scans were acquired on a 3 Tesla Siemens MAGNETOM Trio scanner using a 32-channel head coil at the University of California, Berkeley, Henry H. Wheeler Jr. Brain Imaging Center. A T1-weighted structural image was acquired using multi-echo, magnetization-prepared, 180-degree radio-frequency

pulses and rapid gradient-echo (MEMPRAGE) sampling (TR=2,530.0 ms, TE=1.64-7.22 ms, FOV=256.0 mm, flip angle=7°, slice thickness=1.0 mm, slices per slab=176.0, voxel size=1.0 × 1.0 × 1.0 mm, matrix size=176 × 256 × 256 voxels, scan time [min]=6:03). Additionally, a high-resolution, T2-weighted scan with partial brain coverage focused on the medial temporal lobe and an oblique coronal slice orientation (positioned orthogonally to the longitudinal axis of the hippocampus) was acquired from each participant for use in parcellation of the subfields of the hippocampus (TR= 8,020.0 ms, TE= 51.0 ms, FOV=175 mm, flip angle=122°, slice thickness=1.5 mm, slices=40, voxel size=0.4 × 0.4 × 1.5 mm, matrix size=448 × 448 × 40 voxels, scan time [min]=3:54). Diffusion-weighted images were collected along 111 directions using a multi-shell sequence (b = 1000 s/mm2 and b = 2000 s/mm2) and sensitivity encoding image reconstruction (SENSE) (TR=3,400.0 ms, TE=94.8 ms, FOV=210 mm, flip angle=78°, slice thickness=2.0 mm, slices=69, voxel size=2.0 × 2.0 × 2.0 mm, matrix size=104 × 104 × 69 voxels), scan time [min]=6:31). Additional b0 (no diffusion weighting) images were collected in the reverse phase encoding direction for estimation of susceptibility-induced distortions in image preprocessing (described in subsequent sections).

MRI Processing

Hippocampal Subfield Segmentation. Region-of-interest (ROI) masks were generated for the following subfields of the hippocampus using T1-weighted and high-resolution, T2-weighted structural scans with FreeSurfer Version 7.2.0: Cornu Ammonis (CA) subregions 1, 3, and 4; the dentate gyrus; and the subiculum (51). Due to the difficulty in reliably segmenting CA2 and CA3 separately, the Free Surfer algorithm opts to subsume CA2 in the segmentation for CA3, an approach we adhered to in this study. For the purposes of these analyses, the Dentate Gyrus/Granule Cell Layer and Molecular Layer segmentations were combined, as cytoarchitectural delineation of the subfields of the hippocampus suggest that the Molecular and Granule Cell Layers are structures within the Dentate Gyrus (108). The three subregions of the subiculum were similarly analyzed together as a single structure.

Diffusion-weighted Imaging (DWI). Diffusion-weighted scans were denoised using dwidenoise from MRtrix3 Version 3.0.2 (109–112). Susceptibility-induced distortions were then calculated using topup from FSL Version 6.0.5.1(52,113,114). Eddy current distortion correction and susceptibility-induced distortion correction was performed with slice-to-volume motion correction and outlier replacement using eddy from FSL Version 6.0.5.1 (115–117). The distortion-corrected volumes were then visually inspected for remaining image

quality issues before NODDI metrics were estimated.

NODDI Metrics. The NODDI model was fitted to participants' preprocessed diffusion images using the AMICO pipeline Version 1.4.2 to produce estimates of neurite density for the whole brain (118). The NODDI model hybridizes the tensor-based and ball-and-stick modeling approaches to consider three contributions to the overall diffusion signal: intra-neurite fluid, modeled as 'sticks', or highly-restricted diffusion; fluid in cell bodies and the immediate extracellular space, modeled as anisotropic, or partially restricted diffusion; and free water, modeled as isotropic diffusion (51). NODDI calculates the relative contributions of each component of the tissue to produce a whole-brain map of intra-neurite fluid, termed the Intracellular Volume Fraction (ICVF). The ICVF has been validated against gold-standard histological measures of neurite density. These have shown positive correlation with myelin expression (119) and dendritic complexity and length (120), and highly negative correlation with measures of cell body density (119). ICVF is referred to as the neurite density index (NDI) hereafter in this manuscript. The average NDI was computed bilaterally for each hippocampal subfield for use in the statistical analyses.

Data Analysis Plan

All analyses were conducted in R version 4.2.1 using a combination of base R and tidyverse functions (121,122). The dependent variables were examined for normality by examining skewness and kurtosis values, and visually, through inspection of graphic representations of these data. All independent variables (PNMI biomarkers) were natural log-transformed due to skewed values (as is typical with these biomarkers) (14,15,49,106).

Bivariate analyses using Spearman's correlations were conducted between the biomarkers of PNMI and hippocampal data (113). Covariates (maternal education, offspring intracranial volume, offspring age at time of imaging, offspring sex, and offspring handedness) were identified as either variables significantly associated with both the independent and dependent variables or variables implicated in the focal processes (i.e. hippocampal neurite density or PNMI).

Following previous research (15), separate hierarchical linear regressions were conducted to estimate the association between each T1 and T2 inflammatory biomarker and overall hippocampal neurite density. For each model, covariates (maternal education and offspring age, sex, handedness, and intracranial volume) were entered in the first step of the model. The second step added the biomarker of interest.

To investigate the effect of PNMI on hippocampal subfield

microarchitecture, we repeated our hierarchical linear regressions to estimate the association between each T1 and T2 inflammatory biomarker and neurite density of each hippocampal subregion. For each subregion, four hierarchical multiple regressions were conducted to predict neurite density, one for each biomarker (IL-1RA, IL-6, IL-8, and sTNF-RII, respectively).

All model results were examined for evidence of and met regression assumptions, including linearity, homoscedasticity, independence between observations, and normality of residuals.

Acknowledgments

The authors thank the Child Health and Development Studies and Healthy Brains Research Teams and the participating families for their ongoing dedication to our studies. This work was supported by grants NIMH R01 MH118545 and NIMH R01 MH096478 (PI, Ellman) from the National Institute of Mental Health, and grants N01HD13334 and N01HD63258 from The Eunice Kennedy Shriver National Institute of Child Health and Human Development (PI, Cohn).

References

1. De Asis-Cruz J, Andescavage N, Limperopoulos C. Adverse Prenatal Exposures and Fetal Brain Development: Insights From Advanced Fetal Magnetic Resonance Imaging. Biol Psychiatry Cogn Neurosci Neuroimaging. 2022;7:480–90.

2. Doi M, Usui N, Shimada S. Prenatal Environment and Neurodevelopmental Disorders. Front Endocrinol. 2022;13:860110.

3. Fitzgerald E, Hor K, Drake AJ. Maternal influences on fetal brain development: The role of nutrition, infection and stress, and the potential for intergenerational consequences. Early Hum Dev. 2020 Nov;150:105190.

4. Stiles J, Jernigan TL. The Basics of Brain Development. Neuropsychol Rev. 2010;20:327–48.

5. Han VX, et al. Maternal acute and chronic inflammation in pregnancy is associated with common neurodevelopmental disorders: a systematic review. Transl Psychiatry. 2021;11:1–21.

6. Brown AS, Sourander A, Hinkka-Yli-Salomäki S, McKeague IW, Sundvall J, Surcel HM. Elevated Maternal C-Reactive Protein and Autism in a National Birth Cohort. Mol Psychiatry. 2014 Feb;19(2):259–64.

7. Zawadzka A, Cieślik M, Adamczyk A. The Role of Maternal Immune Activation in the Pathogenesis of Autism: A Review of the Evidence, Proposed Mechanisms and Implications for Treatment. Int J Mol Sci. 2021 Oct 26;22(21):11516.

8. Sato A, Kotajima-Murakami H, Tanaka M, Katoh Y, Ikeda K. Influence of Prenatal Drug Exposure, Maternal Inflammation, and Parental Aging on the Development of Autism Spectrum Disorder. Front Psychiatry. 2022;13:821455.

9. Beversdorf DQ, Stevens HE, Jones KL. Prenatal Stress, Maternal Immune Dysregulation, and Their Association With Autism Spectrum Disorders. Curr Psychiatry Rep. 2018 Aug 9;20(9):76.

10. Brown AS, Begg MD, Gravenstein S, Schaefer CA, Wyatt RJ, Bresnahan M, et al. Serologic Evidence of Prenatal Influenza in the Etiology of Schizophrenia. Arch Gen Psychiatry. 2004 Aug 1;61(8):774–80.

11. Goldstein JM, Cherkerzian S, Seidman LJ, Donatelli J a. L, Remington AG, Tsuang MT, et al. Prenatal maternal immune disruption and sex-dependent risk for psychoses. Psychol Med. 2014 Nov;44(15):3249–61.

12. Allswede DM, Yolken RH, Buka SL, Cannon TD. Cytokine concentrations throughout pregnancy and risk for psychosis in adult offspring: a longitudinal case-control study. Lancet Psychiatry. 2020 Mar 1;7(3):254–61.

13. Kappelmann N, Perry BI, Khandaker GM. Prenatal and Childhood Immuno-Metabolic Risk Factors for Adult Depression and Psychosis. Harv Rev Psychiatry. 2022 Feb;30(1):8.

14. Mac Giollabhui N, Breen EC, Murphy SK, Maxwell SD, Cohn BA, Krigbaum NY, et al. Maternal inflammation during pregnancy and offspring psychiatric symptoms in childhood: Timing and sex matter. J Psychiatr Res. 2019 Apr;111:96–103.

15. Lipner E, Mac Giollabhui N, Breen EC, Cohn BA, Krigbaum NY, Cirillo PM, et al. Sex-Specific Pathways From Prenatal Maternal Inflammation to Adolescent Depressive Symptoms | Adolescent Medicine. JAMA Psychiatry [Internet]. 2024 [cited 2024 Feb 26]; Available from: https://jamanetwork.com/journals/jamapsychiatry/article-abstract/2814653

16. Burgess N, Maguire EA, O'Keefe J. The human hippocampus and spatial and episodic memory. Neuron. 2002 Aug 15;35(4):625–41.

17. Elliott BL, D'Ardenne K, Murty VP, Brewer GA, McClure SM. Midbrain–Hippocampus Structural Connectivity Selectively Predicts Motivated Memory Encoding. J Neurosci. 2022 Dec 14;42(50):9426–34.

18. Jacobson L, Sapolsky R. The role of the hippocampus in feedback regulation of the hypothalamic-pituitary-adrenocortical axis. Endocr Rev. 1991 May;12(2):118–34.

19. Murty VP, Adcock RA. Enriched encoding: reward motivation organizes cortical networks for hippocampal detection of unexpected events. Cereb Cortex N Y N 1991. 2014 Aug;24(8):2160–8.

20. Phelps EA. Human emotion and memory: interactions of the amygdala and hippocampal complex. Curr Opin Neurobiol. 2004 Apr;14(2):198–202.

21. Scoville WB, Milner B. LOSS OF RECENT MEMORY AFTER BILATERAL HIPPOCAMPAL LESIONS. J Neurol Neurosurg Psychiatry. 1957 Feb;20(1):11–21.

22. Squire LR, Zola-Morgan S. The medial temporal lobe memory system. Science. 1991 Sep 20;253(5026):1380–6.

23. Lieberman JA, Girgis RR, Brucato G, Moore H, Provenzano F, Kegeles L, et al. Hippocampal dysfunction in the pathophysiology of schizophrenia: a

selective review and hypothesis for early detection and intervention. Mol Psychiatry. 2018 Aug;23(8):1764–72.

24. MacQueen G, Frodl T. The hippocampus in major depression: evidence for the convergence of the bench and bedside in psychiatric research? Mol Psychiatry. 2011 Mar;16(3):252–64.

25. Banker SM, Gu X, Schiller D, Foss-Feig JH. Hippocampal contributions to social and cognitive deficits in autism spectrum disorder. Trends Neurosci. 2021 Oct 1;44(10):793–807.

26. Bilbo SD, Schwarz JM. The immune system and developmental programming of brain and behavior. Front Neuroendocrinol. 2012 Aug;33(3):267–86.

27. Meyer U, Yee BK, Feldon J. The Neurodevelopmental Impact of Prenatal Infections at Different Times of Pregnancy: The Earlier the Worse? The Neuroscientist. 2007 Jun;13(3):241–56.

28. Arnold SE, Trojanowski JQ. Human fetal hippocampal development: II. The neuronal cytoskeleton. J Comp Neurol. 1996;367(2):293–307.

29. Goldstein JM, Cohen JE, Mareckova K, Holsen L, Whitfield-Gabrieli S, Gilman SE, et al. Impact of prenatal maternal cytokine exposure on sex differences in brain circuitry regulating stress in offspring 45 years later. Proc Natl Acad Sci. 2021 Apr 13;118(15):e2014464118.

30. Musaelyan K, Egeland M, Fernandes C, Pariante CM, Zunszain PA, Thuret S. Modulation of Adult Hippocampal Neurogenesis by Early-Life Environmental Challenges Triggering Immune Activation. Neural Plast. 2014;2014:1–10.

31. Green HF, Nolan YM. Inflammation and the developing brain: Consequences for hippocampal neurogenesis and behavior. Neurosci Biobehav Rev. 2014 Mar 1;40:20–34.

32. Couch ACM, Berger T, Hanger B, Matuleviciute R, Srivastava DP, Thuret S, et al. Maternal immune activation primes deficiencies in adult hippocampal neurogenesis. Brain Behav Immun. 2021 Oct;97:410–22.

33. Bitanihirwe BK, Peleg-Raibstein D, Mouttet F, Feldon J, Meyer U. Late Prenatal Immune Activation in Mice Leads to Behavioral and Neurochemical Abnormalities Relevant to the Negative Symptoms of Schizophrenia. Neuropsychopharmacology. 2010 Nov;35(12):2462–78.

34. Makinodan M, Tatsumi K, Manabe T, Yamauchi T, Makinodan E, Matsuyoshi H, et al. Maternal immune activation in mice delays myelination

and axonal development in the hippocampus of the offspring. J Neurosci Res. 2008 Aug 1;86(10):2190–200.

35. Ellman LM, Deicken RF, Vinogradov S, Kremen WS, Poole JH, Kern DM, et al. Structural brain alterations in schizophrenia following fetal exposure to the inflammatory cytokine interleukin-8. Schizophr Res. 2010 Aug;121(1–3):46–54.

36. Meyer U, Feldon J, Schedlowski M, Yee BK. Immunological stress at the maternal–foetal interface: A link between neurodevelopment and adult psychopathology. Brain Behav Immun. 2006 Jul 1;20(4):378–88.

37. Mouihate A, Kalakh S. Maternal Interleukin-6 Hampers Hippocampal Neurogenesis in Adult Rat Offspring in a Sex-Dependent Manner. Dev Neurosci. 2021;43(2):106–15.

38. Garay PA, Hsiao EY, Patterson PH, McAllister AK. Maternal immune activation causes age- and region-specific changes in brain cytokines in offspring throughout development. Brain Behav Immun. 2013 Jul;31:54–68.

39. Graham AM, Doyle O, Tilden EL, Sullivan EL, Gustafsson HC, Marr M, et al. Effects of Maternal Psychological Stress During Pregnancy on Offspring Brain Development: Considering the Role of Inflammation and Potential for Preventive Intervention. Biol Psychiatry Cogn Neurosci Neuroimaging. 2022 May 1;7(5):461–70.

40. Demers CH, Aran Ö, Glynn LM, Davis EP. Prenatal Programming of Neurodevelopment: Structural and Functional Changes. In: Wazana A, Székely E, Oberlander TF, editors. Prenatal Stress and Child Development [Internet]. Cham: Springer International Publishing; 2021 [cited 2023 Mar 14]. p. 193–242. Available from: https://doi.org/10.1007/978-3-030-60159-1_9

41. Mufson EJ, Mahady L, Waters D, Counts SE, Perez SE, DeKosky S, et al. Hippocampal Plasticity During the Progression of Alzheimer's disease. Neuroscience. 2015 Nov 19;309:51–67.

42. Andersen P, Morris R, Amaral D, Bliss T, O'Keefe J. The hippocampus book. Oxford university press; 2006.

43. Hao LY, Hao XQ, Li SH, Li XH. Prenatal exposure to lipopolysaccharide results in cognitive deficits in age-increasing offspring rats. Neuroscience. 2010;166(3):763–70.

44. Samuelsson AM, Jennische E, Hansson HA, Holmäng A. Prenatal exposure to interleukin-6 results in inflammatory neurodegeneration in hippocampus

with NMDA/GABAA dysregulation and impaired spatial learning. Am J Physiol-Regul Integr Comp Physiol. 2006 May;290(5):R1345–56.

45. Belnoue L, Grosjean N, Ladevèze E, Abrous DN, Koehl M. Prenatal stress inhibits hippocampal neurogenesis but spares olfactory bulb neurogenesis. PloS One. 2013;8(8):e72972.

46. Girard S, Sébire H, Brochu ME, Briota S, Sarret P, Sébire G. Postnatal administration of IL-1Ra exerts neuroprotective effects following perinatal inflammation and/or hypoxic-ischemic injuries. Brain Behav Immun. 2012 Nov 1;26(8):1331–9.

47. Graciarena M, Depino AM, Pitossi FJ. Prenatal inflammation impairs adult neurogenesis and memory related behavior through persistent hippocampal TGFβ1 downregulation. Brain Behav Immun. 2010;24(8):1301–9.

48. Wu D, Lei J, Xie H, Dong J, Burd I. Diffusion MRI revealed altered inter-hippocampal projections in the mouse brain after intrauterine inflammation. Brain Imaging Behav. 2020;14:383–95.

49. Kelley MH, Wu WW, Lei J, McLane M, Xie H, Hart KD, et al. Functional changes in hippocampal synaptic signaling in offspring survivors of a mouse model of intrauterine inflammation. J Neuroinflammation. 2017;14:1–8.

50. Ito HT, Smith SE, Hsiao E, Patterson PH. Maternal immune activation alters nonspatial information processing in the hippocampus of the adult offspring. Brain Behav Immun. 2010;24(6):930–41.

51. Zhang H, Schneider T, Wheeler-Kingshott CA, Alexander DC. NODDI: Practical in vivo neurite orientation dispersion and density imaging of the human brain. NeuroImage. 2012 Jul;61(4):1000–16.

52. Mouihate A, Kalakh S. Breastfeeding promotes oligodendrocyte precursor cells division and myelination in the demyelinated *corpus callosum*. Brain Res. 2023 Dec 15;1821:148584.

53. Irwin MR, Olmos L, Wang M, Valladares EM, Motivala SJ, Fong T, et al. Cocaine Dependence and Acute Cocaine Induce Decreases of Monocyte Proinflammatory Cytokine Expression across the Diurnal Period: Autonomic Mechanisms. J Pharmacol Exp Ther. 2007;320(2):507–15.

54. Zunszain PA, Anacker C, Cattaneo A, Choudhury S, Musaelyan K, Myint AM, et al. Interleukin-1β: A New Regulator of the Kynurenine Pathway Affecting Human Hippocampal Neurogenesis. Neuropsychopharmacology. 2012 Mar;37(4):939–49.

55. Salvador AF, de Lima KA, Kipnis J. Neuromodulation by the immune system: a focus on cytokines. Nat Rev Immunol. 2021 Aug;21(8):526–41.

56. Beran TN. Elliott, C. D. (2007). Differential Ability Scales (2nd ed.). San Antonio, TX: Harcourt Assessment. Can J Sch Psychol. 2007 Jun 1;22(1):128–32.

57. Dozmorov MG, Bilbo SD, Kollins SH, Zucker N, Do EK, Schechter JC, et al. Associations between maternal cytokine levels during gestation and measures of child cognitive abilities and executive functioning. Brain Behav Immun. 2018;70:390–7.

58. Eichenbaum H, Cohen NJ. Can we reconcile the declarative memory and spatial navigation views on hippocampal function? Neuron. 2014;83(4):764–70.

59. Yonelinas AP. The hippocampus supports high-resolution binding in the service of perception, working memory and long-term memory. Behav Brain Res. 2013;254:34–44.

60. Olson IR, Page K, Moore KS, Chatterjee A, Verfaellie M. Working memory for conjunctions relies on the medial temporal lobe. J Neurosci. 2006;26(17):4596–601.

61. Duff MC, Covington NV, Hilverman C, Cohen NJ. Semantic memory and the hippocampus: Revisiting, reaffirming, and extending the reach of their critical relationship. Front Hum Neurosci. 2020;13:471.

62. Solomon SH, Schapiro AC. Semantic search as pattern completion across a concept. Trends Cogn Sci. 2020;24(2):95–8.

63. Morton NW, Preston AR. Concept formation as a computational cognitive process. Curr Opin Behav Sci. 2021;38:83–9.

64. Moscovitch M, Cabeza R, Winocur G, Nadel L. Episodic memory and beyond: the hippocampus and neocortex in transformation. Annu Rev Psychol. 2016;67(1):105–34.

65. Buckner RL. The role of the hippocampus in prediction and imagination. Annu Rev Psychol. 2010;61(1):27–48.

66. Biderman N, Bakkour A, Shohamy D. What are memories for? The hippocampus bridges past experience with future decisions. Trends Cogn Sci. 2020;24(7):542–56.

67. Rubin RD, Watson PD, Duff MC, Cohen NJ. The role of the hippocampus in flexible cognition and social behavior. Front Hum Neurosci. 2014;8:742.

68. Knuesel I, Chicha L, Britschgi M, Schobel SA, Bodmer M, Hellings JA, et al. Maternal immune activation and abnormal brain development across CNS disorders. Nat Rev Neurol. 2014 Nov;10(11):643–60.

69. Smith SEP, Li J, Garbett K, Mirnics K, Patterson PH. Maternal Immune Activation Alters Fetal Brain Development through Interleukin-6. J Neurosci. 2007 Oct 3;27(40):10695–702.

70. Mirabella F, Desiato G, Mancinelli S, Fossati G, Rasile M, Morini R, et al. Prenatal interleukin 6 elevation increases glutamatergic synapse density and disrupts hippocampal connectivity in offspring. Immunity. 2021 Nov 9;54(11):2611-2631.e8.

71. Kessler RC, Davis CG, Kendler KS. Childhood adversity and adult psychiatric disorder in the US National Comorbidity Survey. Psychol Med. 1997 Sep;27(5):1101–19.

72. Thomas RM, Hotsenpiller G, Peterson DA. Acute psychosocial stress reduces cell survival in adult hippocampal neurogenesis without altering proliferation. J Neurosci Off J Soc Neurosci. 2007 Mar 14;27(11):2734–43.

73. Malberg JE, Duman RS. Cell proliferation in adult hippocampus is decreased by inescapable stress: reversal by fluoxetine treatment. Neuropsychopharmacol Off Publ Am Coll Neuropsychopharmacol. 2003 Sep;28(9):1562–71.

74. Konstantine, Zakzanis K, Leach L, Kaplan E. On the Nature and Pattern of Neurocognitive Function in Major Depressive Disorder. Cogn Behav Neurol. 1998 Jul;11(3):111.

75. Lisman JE. Relating Hippocampal Circuitry to Function: Recall of Memory Sequences by Reciprocal Dentate–CA3 Interactions. Neuron. 1999 Feb 1;22(2):233–42.

76. Bakker A, Kirwan CB, Miller M, Stark CEL. Pattern separation in the human hippocampal CA3 and dentate gyrus. Science. 2008 Mar 21;319(5870):1640–2.

77. Cerasti E, Treves A. The spatial representations acquired in CA3 by self-organizing recurrent connections. Front Cell Neurosci [Internet]. 2013 [cited 2023 Sep 22];7. Available from: https://www.frontiersin.org/articles/10.3389/fncel.2013.00112

78. Kesner R. A process analysis of the CA3 subregion of the hippocampus. Front Cell Neurosci [Internet]. 2013 [cited 2023 Sep 22];7. Available from: https://www.frontiersin.org/articles/10.3389/fncel.2013.00078

79. Rolls E. The mechanisms for pattern completion and pattern separation in the hippocampus. Front Syst Neurosci [Internet]. 2013 [cited 2023 Sep 22];7. Available from: https://www.frontiersin.org/articles/10.3389/fnsys.2013.00074

80. Rolls ET. Pattern separation, completion, and categorisation in the hippocampus and neocortex. Neurobiol Learn Mem. 2016 Mar;129:4–28.

81. Treves A, Rolls ET. Computational analysis of the role of the hippocampus in memory. Hippocampus. 1994 Jun;4(3):374–91.

82. Yassa MA, Stark CEL. Pattern separation in the hippocampus. Trends Neurosci. 2011 Oct;34(10):515–25.

83. Mueller SG, Weiner MW. Selective effect of age, Apo e4, and Alzheimer's disease on hippocampal subfields. Hippocampus. 2009 Jun;19(6):558–64.

84. Mueller S, Schuff N, Raptentsetsang S, Elman J, Weiner M. Selective Effect of Apo e4 on CA3 and Dentate in Normal Aging and Alzheimer's Disease using high resolution MRI at 4Tesla. NeuroImage. 2008 Aug 1;42(1):42–8.

85. Salthouse TA. When does age-related cognitive decline begin? Neurobiol Aging. 2009 Apr 1;30(4):507–14.

86. Konefal SC, Stellwagen D. Tumour necrosis factor-mediated homeostatic synaptic plasticity in behavioural models: testing a role in maternal immune activation. Philos Trans R Soc B Biol Sci. 2017 Mar 5;372(1715):20160160.

87. Antonson AM. Prenatal programming of the brain: The maternal-fetal interface takes center stage. Brain Behav Immun. 2023 Mar 1;109:219–20.

88. Girard S, Tremblay L, Lepage M, Sébire G. IL-1 Receptor Antagonist Protects against Placental and Neurodevelopmental Defects Induced by Maternal Inflammation. J Immunol. 2010 Apr 1;184(7):3997–4005.

89. Hsiao EY, Patterson PH. Placental regulation of maternal-fetal interactions and brain development. Dev Neurobiol. 2012;72(10):1317–26.

90. Town T, Nikolic V, Tan J. The microglial "activation" continuum: from innate to adaptive responses. J Neuroinflammation. 2005 Oct 31;2(1):24.

91. Humphreys KL, King LS, Sacchet MD, Camacho MC, Colich NL, Ordaz SJ, et al. Evidence for a sensitive period in the effects of early life stress on hippocampal volume. Dev Sci. 2019 May;22(3):e12775.

92. Graciarena M, Roca V, Mathieu P, Depino AM, Pitossi FJ. Differential vulnerability of adult neurogenesis by adult and prenatal inflammation: Role of TGF-β1. Brain Behav Immun. 2013 Nov 1;34:17–28.

93. Tong M, Abrahams VM. Immunology of the Placenta. Obstet Gynecol Clin. 2020 Mar 1;47(1):49–63.

94. Schönbrodt FD, Perugini M. At what sample size do correlations stabilize? J Res Personal. 2013 Oct 1;47(5):609–12.

95. Van Dyke TE, van Winkelhoff AJ. Infection and inflammatory mechanisms. J Clin Periodontol. 2013;40(s14):S1–7.

96. Rohleder N. Stimulation of Systemic Low-Grade Inflammation by Psychosocial Stress. Psychosom Med. 2014 Apr;76(3):181.

97. Karczewski J, Śledzińska E, Baturo A, Jończyk I, Maleszko A, Samborski P, et al. Obesity and inflammation. Eur Cytokine Netw. 2018 Sep 1;29(3):83–94.

98. Stumpf F, Keller B, Gressies C, Schuetz P. Inflammation and Nutrition: Friend or Foe? Nutrients. 2023 Jan;15(5):1159.

99. Mozzoni P, Iodice S, Persico N, Ferrari L, Pinelli S, Corradi M, et al. Maternal air pollution exposure during the first trimester of pregnancy and markers of inflammation and endothelial dysfunction. Environ Res. 2022 Sep 1;212:113216.

100. Meyyazhagan A, Kuchi Bhotla H, Pappuswamy M, Tsibizova V, Al Qasem M, Di Renzo GC. Cytokine see-saw across pregnancy, its related complexities and consequences. Int J Gynecol Obstet. 2023;160(2):516–25.

101. Brown AS, Vinogradov S, Kremen WS, Poole JH, Deicken RF, Penner JD, et al. Prenatal Exposure to Maternal Infection and Executive Dysfunction in Adult Schizophrenia. Am J Psychiatry. 2009 Jun;166(6):683–90.

102. Braun AE, Carpentier PA, Babineau BA, Narayan AR, Kielhold ML, Moon HM, et al. "Females Are Not Just 'Protected' Males": Sex-Specific Vulnerabilities in Placenta and Brain after Prenatal Immune Disruption. eNeuro. 2019 Nov 6;6(6):ENEURO.0358-19.2019.

103. van den Berg BJ, Christianson RE, Oechsli FW. The California Child Health and Development Studies of the School of Public Health, University of California at Berkeley*. Paediatr Perinat Epidemiol. 1988;2(3):265–82.

104. Fineberg AM, Ellman LM, Schaefer CA, Maxwell SD, Shen L, Chaudhury NH, et al. Fetal exposure to maternal stress and risk for schizophrenia

spectrum disorders among offspring: Differential influences of fetal sex. Psychiatry Res. 2016 Feb 28;236:91–7.

105. Lipner E, Murphy SK, Breen EC, Cohn BA, Krigbaum NY, Cirillo PM, et al. Infection and higher cortisol during pregnancy and risk for depressive symptoms in adolescent offspring. Psychoneuroendocrinology. 2022 Jul;141:105755.

106. Pike MR, Lipner E, O'Brien KJ, Breen EC, Cohn BA, Cirillo PM, et al. Prenatal maternal Inflammation, childhood cognition and adolescent depressive symptoms. Brain Behav Immun. 2024 Jul 1;119:908–18.

107. Schlotz W, Phillips DIW. Fetal origins of mental health: Evidence and mechanisms. Brain Behav Immun. 2009 Oct;23(7):905–16.

108. Amaral DG, Scharfman HE, Lavenex P. The dentate gyrus: fundamental neuroanatomical organization (dentate gyrus for dummies). Prog Brain Res. 2007;163:3–22.

109. Cordero-Grande L, Christiaens D, Hutter J, Price AN, Hajnal JV. Complex diffusion-weighted image estimation via matrix recovery under general noise models. NeuroImage. 2019 Oct 15;200:391–404.

110. Jelescu IO, Veraart J, Adisetiyo V, Milla SS, Novikov DS, Fieremans E. One diffusion acquisition and different white matter models: how does microstructure change in human early development based on WMTI and NODDI? NeuroImage. 2015 Feb 15;107:242–56.

111. Tournier JD, Smith R, Raffelt D, Tabbara R, Dhollander T, Pietsch M, et al. MRtrix3: A fast, flexible and open software framework for medical image processing and visualisation. NeuroImage. 2019 Nov 15;202:116137.

112. Veraart J, Fieremans E, Novikov DS. Diffusion MRI noise mapping using random matrix theory. Magn Reson Med. 2016;76(5):1582–93.

113. Spearman C. The proof and measurement of association between two things. Am J Psychol. 1904;15(1):72–101.

114. Mouihate A, Kalakh S, AlMutairi R, Alashqar A. Prenatal Inflammation Dampens Neurogenesis and Enhances Serotonin Transporter Expression in the Hippocampus of Adult Female Rats. Med Princ Pract. 2019;28(4):352–60.

115. Andersson JLR, Graham MS, Drobnjak I, Zhang H, Filippini N, Bastiani M. Towards a comprehensive framework for movement and distortion correction of diffusion MR images: Within volume movement. Neuroimage. 2017 May 15;152:450–66.

116. Andersson JLR, Sotiropoulos SN. An integrated approach to correction for off-resonance effects and subject movement in diffusion MR imaging. Neuroimage. 2016 Jan 15;125:1063–78.

117. Andersson JLR, Graham MS, Zsoldos E, Sotiropoulos SN. Incorporating outlier detection and replacement into a non-parametric framework for movement and distortion correction of diffusion MR images. NeuroImage. 2016 Nov 1;141:556–72.

118. Daducci A, Canales-Rodríguez EJ, Zhang H, Dyrby TB, Alexander DC, Thiran JP. Accelerated Microstructure Imaging via Convex Optimization (AMICO) from diffusion MRI data. NeuroImage. 2015 Jan 15;105:32–44.

119. Wang S, Zhao Y, Zhang L, Wang X, Wang X, Cheng B, et al. Stress and the brain: Perceived stress mediates the impact of the superior frontal gyrus spontaneous activity on depressive symptoms in late adolescence. Hum Brain Mapp. 2019 Aug 9;40(17):4982–93.

120. Crombe A, Planche V, Raffard G, Bourel J, Dubourdieu N, Panatier A, et al. Deciphering the microstructure of hippocampal subfields with *in vivo* DTI and NODDI: Applications to experimental multiple sclerosis. NeuroImage. 2018 May 15;172:357–68.

121. Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al. Welcome to the Tidyverse. J Open Source Softw. 2019 Nov 21;4(43):1686.

122. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2022. Available from: https://www.R-project.org/

Figures and Tables

Figure 1. Participants Retained in Subsamples of the Child Health and Development Studies (CHDS) and Healthy Brain Project (HBP). T1 = first trimester of pregnancy; T2 = second trimester of pregnancy.

Figure 2. (A) Color maps for B-F; (B) map of subfields of the hippocampus; (C) significance levels of relationship between prenatal exposure to first-trimester maternal IL-1β and subfield neurite density in the first trimester; (D) significance levels of relationship between prenatal exposure to second-trimester maternal IL-1β and subfield neurite density; (E) significance levels of relationship between prenatal exposure to first-trimester maternal IL-6 and subfield neurite density in the first trimester; (F) significance levels of relationship between prenatal exposure to second-trimester maternal IL-6 and subfield neurite density in the second trimester; CA = Cornu Ammonis, DG = Dentate Gyrus, Molecular Layer, and Granule Cell Layer.

Table 1. *Participant Demographics.*

Age (years) M (SD) [range]	59.6 (1.1) [57.3-62.3]
Sex % (n) Male	48.6 (36)
Ethnicity % (n) Hispanic	8.1 (6)
Race % (n)	
American Indian/Alaska Native	2.7 (2)
Asian	6.8 (5)
Black	12.2 (9)
White	70.3 (52)
More than one race	6.8 (5)
Unknown	1.4 (1)
Maternal Education % (n)	
< high school	6.8 (5)
= high school	33.8 (25)
> high school	59.5 (44)

Note. N = 74; demographics are from all participants who were included in at least one hierarchical regression analysis.

Table 2. *Spearman's correlations for variables of interest: first trimester.*

Spearman's ρ	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1. Maternal Education	1.00	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2. Age	0.850	1.00	-	-	-	-	-	-	-	-	-	-	-	-	-
3. Sex	-0.091	0.017	1.00	-	-	-	-	-	-	-	-	-	-	-	-
4. Handedness	-0.201	-0.118	-0.056	1.00	-	-	-	-	-	-	-	-	-	-	-
5. Intracranial Volume	0.200	-0.011	**-0.059****	-0.132	1.00	-	-	-	-	-	-	-	-	-	-
6. IL-1RA	-0.036	0.014	-0.035	-0.150	0.225	1.00	-	-	-	-	-	-	-	-	-
7. IL-6	-0.059	-0.058	0.081	-0.091	0.015	**0.428****	1.00	-	-	-	-	-	-	-	-
8. IL-8	0.080	**-0.363****	0.026	-0.026	0.017	**0.283***	**0.606****	1.00	-	-	-	-	-	-	-

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
9. sTNF-RII	-0.012	0.063	-0.109	-0.094	0.190	**0.464****	**0.311****	0.156	1.00	-	-	-	-	-	-
10. Whole HPC	0.056	-0.031	-0.096	0.053	-0.110	-0.172	-0.204	0.010	-0.153	1.00	-	-	-	-	-
11. CA1	0.141	0.053	**-0.252***	0.021	0.171	-0.025	-0.207	-0.098	-0.113	**0.737****	1.00	-	-	-	-
12. CA3	-0.028	-0.147	**-0.232***	0.125	-0.089	**-0.267***	-0.229	-0.085	**-0.265***	**0.755****	**0.672****	1.00	-	-	-
13. CA4	0.034	-0.063	0.068	0.109	-0.204	-0.178	-0.172	-0.071	**-0.267***	**0.778****	**0.535****	**0.690****	1.00	-	-
14. DG	0.059	-0.124	-0.030	0.096	-0.147	**-0.243***	-0.206	-0.004	-0.224	**0.936****	**0.650****	**0.740****	**0.869****	1.00	-
15. Subiculum	0.027	-0.132	0.025	-0.046	-0.138	-0.118	0.016	0.203	-0.057	**0.696****	**0.334****	**0.296***	**0.347****	**0.599****	1.00

Note. Covariates included were maternal education at birth, offspring age, sex, handedness, and intracranial volume; IL = Interleukin, RA = receptor antagonist, sTNF-RII = soluble tumor necrosis factor receptor-II, HPC = hippocampus, CA = Cornu Ammonis, DG = Dentate Gyrus (including the molecular and granule cell layers); * $p < 0.05$, ** $p < 0.01$

Table 3. *Spearman's correlations for variables of interest: second trimester.*

Spearman's ρ	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1. Maternal Education	1.00	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2. Age	0.042	1.00	-	-	-	-	-	-	-	-	-	-	-	-	-
3. Sex	-0.104	-0.063	1.00	-	-	-	-	-	-	-	-	-	-	-	-
4. Handedness	-0.181	-0.199	-0.098	1.00	-	-	-	-	-	-	-	-	-	-	-
5. Intracranial Volume	0.185	0.007	**-0.622****	-0.052	1.00	-	-	-	-	-	-	-	-	-	-
6. IL-1RA	-0.040	0.168	0.020	0.072	0.077	1.00	-	-	-	-	-	-	-	-	-
7. IL-6	0.005	-0.014	0.074	0.022	0.048	**0.414****	1.00	-	-	-	-	-	-	-	-

8. IL-8	0.040	-0.115	0.058	-0.031	0.047	**0.323****	**0.630****	1.00	-	-	-	-	-	-	-
9. sTNF-RII	-0.149	-0.194	0.006	0.173	0.054	0.158	**0.337****	0.143	1.00	-	-	-	-	-	-
10. Whole HPC	0.077	0.000	-0.067	0.062	-0.139	-0.101	-0.041	-0.031	0.013	1.00	-	-	-	-	-
11. CA1	0.127	0.039	**-0.245***	0.013	0.145	0.018	-0.120	-0.164	-0.028	**0.733****	1.00	-	-	-	-
12. CA3	-0.049	-0.109	-0.195	0.158	-0.120	-0.179	-0.132	-0.106	0.005	**0.739****	**0.685****	1.00	-	-	-
13. CA4	0.025	-0.041	0.093	0.152	**-0.256***	-0.153	-0.190	-0.114	-0.204	**0.784****	**0.542****	**0.682****	1.00	-	-
14. DG	0.071	-0.088	0.001	0.142	-0.190	-0.139	-0.099	-0.046	-0.017	**0.938****	**0.653****	**0.725****	**0.867****	1.00	-
15. Subiculum	0.081	-0.133	0.026	-0.075	-0.143	-0.148	0.047	0.177	0.102	**0.703****	**0.317****	**0.283***	**0.359****	**0.613****	1.00

Note. Covariates included were maternal education at birth, offspring age, sex, handedness, and intracranial volume; IL = Interleukin, RA = receptor antagonist, sTNF-RII = soluble tumor necrosis factor receptor-II, HPC = hippocampus, CA = Cornu Ammonis, DG = Dentate Gyrus (including the molecular and granule cell layers); * $p < 0.05$, ** $p < 0.01$

Table 4. *Results of hierarchical linear regression analyses for prenatal maternal inflammation and offspring hippocampal microstructure during late middle age.*

	Independent Variable	β	SE	t	p	ΔR²
	Step 1: Covariates					0.040
	Sex	-0.181	0.004	-1.438	0.155	
	Age	-0.084	0.002	-0.827	0.411	
	Intracranial Volume	-0.163	1.210e-08	-1.590	0.117	
	Maternal Education, = High School	0.522	0.008	2.160	**0.034***	
	Maternal Education, > High School	0.458	0.007	1.976	0.052	
	Step 2: Additive Effect					
First Trimester	**IL-1ra**	-0.269	0.004	-2.333	**0.023***	**0.061***
	IL-6	-0.281	0.002	-2.444	**0.017***	**0.067***
	IL-8	-0.087	0.001	-0.672	0.504	0.029
	sTNF-RII	-0.091	0.006	-0.762	0.449	-0.006
Second Trimester	IL-1ra	-0.122	0.004	-0.990	0.326	-0.001
	IL-6	0.088	0.002	0.691	0.492	-0.008
	IL-8	0.012	0.001	0.094	0.925	-0.015
	sTNF-RII	0.107	0.009	0.852	0.398	-0.004

Note. Each of the additive effects were assessed using separate models; IL = Interleukin, RA = receptor antagonist, sTNF-RII = soluble tumor necrosis factor receptor-II, HPC = hippocampus, CA = Cornu Ammonis, DG = Dentate Gyrus (including the molecular and granule cell layers), * $p < 0.05$.

Table 5. *Results of hierarchical linear regression analyses for prenatal maternal inflammation and offspring hippocampal microstructure during late middle age for each hippocampal subfield.*

	Independent Variable	CA1 β (SE)	CA1 t	CA1 ΔR²	CA3 β (SE)	CA3 t	CA3 ΔR²	CA4 β (SE)	CA4 t	CA4 ΔR²	Dentate Gyrus β (SE)	DG t	DG ΔR²	Subiculum β (SE)	Subic t	Subic ΔR²
Step 1: Covariates				0.05			0.10*			0.06			0.02			-0.02
	Sex	-0.217 (0.003)	-1.624		-0.340 (0.010)	**-2.754***		-0.027 (0.005)	-0.308		-0.107 (0.004)	-0.933		0.040 (0.005)	0.131	
	Age	0.060 (0.001)	0.473		-0.152 (0.004)	-1.499		-0.092 (0.002)	-0.872		-0.130 (0.001)	-1.210		-0.147 (0.002)	-1.318	
	Intracranial volume	0.009 (1.08e-8)	-0.096		0.225 (2.68e-8)	-2.266		-0.231 (1.52e-8)	-1.933		-0.166 (1.59e-8)	-1.622		-0.035 (1.598-8)	-0.651	
	Maternal Education, = High School	0.476 (.006)	**2.033***		0.441 (0.016)	1.845		0.532 (0.009)	**2.266***		0.445 (0.008)	1.813		0.280 (0.009)	1.087	
	Maternal Education, > High School	0.462 (0.006)	**1.998***		0.333 (0.016)	1.498		0.484 (0.009)	**2.110***		0.398 (0.008)	1.705		0.291 (0.009)	1.225	
Step 2: Additive Effects																
First Trimester	T1 IL-1RA	-0.096 (.003)	-0.81	-0.004	-0.364 (0.008)	**-3.42****	**0.12****	-0.170 (0.004)	0.15	0.02	-0.280 (0.004)	**-2.42***	**0.07***	-0.255 (0.005)	**-2.13***	**0.05***
First Trimester	T1 IL-6	-0.220 (0.002)	-1.9	0.04	-0.332 (0.005)	**-3.07****	**0.10****	-0.276 (0.002)	**-2.44***	**0.07***	-0.319 (0.002)	**-2.79****	**0.09****	-0.071 (0.003)	-0.58	-0.01
First Trimester	T1 IL-8	-0.142 (0.001)	-1.12	0.004	-0.230 (0.002)	-1.89	0.03	-0.173 (0.001)	-1.38	0.01	-0.149 (0.001)	-1.15	0.005	0.112 (0.001)	0.85	-0.004
First Trimester	T1 sTNF-RII	-0.090 (0.006)	-0.76	-0.01	-0.202 (0.016)	0.08	0.03	-0.128 (0.008)	-1.09	0.003	-0.160 (0.007)	-1.33	0.01	0.027 (0.009)	0.22	-0.01
Second Trimester	T2 IL-1RA	-0.007 (0.004)	-0.06	-0.02	-0.209 (0.010)	-1.79	0.03	-0.099 (0.005)	-0.83	-0.004	-0.113 (0.004)	-0.91	-0.003	-0.145 (0.005)	-1.15	0.005
Second Trimester	T2 IL-6	-0.064 (0.001)	-0.5	-0.01	-0.003 (0.004)	-0.03	-0.01	-0.051 (0.002)	-0.41	-0.01	0.016 (0.002)	0.12	-0.02	0.273 (0.002)	**2.14***	**0.06***
Second Trimester	T2 IL-8	-0.117 (0.001)	-0.94	-0.002	-0.017 (0.003)	-0.15	-0.01	-0.093 (0.001)	-0.77	-0.01	-0.030 (0.001)	-0.22	-0.01	0.193 (0.001)	1.52	0.02
Second Trimester	T2 sTNF-RII	0.043 (0.008)	0.34	-0.01	0.121 (0.020)	1	-3.2e-05	-0.035 (0.011)	-0.29	-0.01	0.101 (0.009)	0.79	-0.01	0.081 (0.012)	0.62	-0.01

Note. Each of the additive effects were assessed using separate models; IL = Interleukin, RA = receptor antagonist, sTNF-RII = soluble tumor necrosis factor receptor-II, HPC = hippocampus, CA = Cornu Ammonis, DG = Dentate Gyrus (including the molecular and granule cell layers), * $p < 0.05$, **$p < 0.01$ FDR corrected for multiple comparisons.