On pathogenicity with varying microbial virulence and host resistance

Anjali Gupta[1], Robert L. Unckless[2*]

[1]Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS, USA

[2]Department of Molecular Biosciences, University of Kansas, Lawrence, KS, USA

[*]Corresponding author

Email: unckless@ku.edu

Running title: Comparative analysis of survival models and pathogen potential

Keywords: Innate immunity, *Drosophila*, host susceptibility, cox proportional hazards model,

pathogenic potential, risk score, fraction symptomatic.

Acknowledgements: We thank Brian Lazzaro for providing the *Providencia* strains, and

Unckless lab members for helpful comments and advice. We also thank two anonymous

reviewers who provided feedback that ultimately greatly improved the manuscript. The work

was supported by NSF 2330095 and NIH R01-AI139154 to RLU and funds from KU

Entomology Endowment to AG.

Data   accessibility: Data   and   code   is   archived   on   Zenodo/GitHub

(https://doi.org/10.5281/zenodo.15116399   /   https://github.com/anjaligupta1210/Pathogenicity-

across-variation-in-microbial-virulence-and-host-resistance.git).

Abstract:

The pathogenicity of a microbe is difficult to define in a comparative context: across microbes

with varying virulence or across host genotypes with varying susceptibilities. Several different

statistical approaches have been employed to investigate pathogenicity and susceptibility.

Measures like proportional mortality or morbidity at a given time are attractive due to their

simplicity but represent a single snapshot. Survival curve approaches, such as the Cox

proportional hazards model and risk scores provide a more nuanced picture of the course of

infection. More recently, Casadevall introduced the concept of pathogenic potential, a composite

measure encompassing both host susceptibility and pathogen virulence, which focuses on the

pathogenicity of a single pathogenic microbe, and later expanded to include additional nuances.

Using *Drosophila melanogaster*, we conducted infection experiments with diverse species of

Providencia bacteria that naturally vary in virulence. We also used several infectious doses to

tune infections. We employed different host genotypes that vary in susceptibility to *Providencia*

infection. Our analysis incorporates factors such as host genotype, pathogen type, inoculum load,

symptomatic fraction, and mortality to compare host- and pathogen-based measures of

pathogenicity. We discuss the advantages and limitations of each method, providing insights into

their applicability in deciphering the intricacies of host-pathogen interactions and guiding the

choice of analytical approaches in infectious disease research.

Introduction:

Virulence is the relative capacity of a microbe to cause damage in a host (Casadevall & Pirofski, 1999). It is intricately linked to host susceptibility, and its measurement is context-dependent, varying across host systems or for interspecific microbial comparisons (Casadevall & Pirofski, 2001). But virulence lacks an absolute measure, emphasizing the need for a relative assessment of a microbe's impact on a host (Casadevall & Pirofski, 1999). Hosts may display symptoms when infected with a specific microbial threshold, exhibiting observable disruptions in normal physiological functions, including mortality in severe cases. This relationship is further complicated by the fact that virulence is not a static characteristic but can evolve in response to changes in host resistance, environmental conditions, and microbial interactions (Ewald, 1994; Read, 1994). Host resistance refers to the ability to limit pathogen burden, while tolerance describes the capacity to minimize damage caused by infection (Råberg et al., 2009). Understanding these distinctions is essential to interpreting the genetic and evolutionary underpinnings of infection outcomes. Pathogen virulence is similarly modulated by evolutionary trade-offs between transmission and host survival. For instance, highly virulent strains may dominate in environments where transmission opportunities are frequent and host mortality does not significantly impede pathogen spread (Frank, 1996). Conversely, in environments where hosts are scarce or transmission is limited, less virulent strains that prolong host survival may be favored (Ebert & Bull, 2003). These trade-offs illustrate the intricate evolutionary dynamics governing host-pathogen interactions and the maintenance of genetic variation within both host and pathogen populations. Additionally, the expression of virulence factors can be modulated by the host immune response, with some pathogens evolving mechanisms to evade or suppress host

defenses, thereby enhancing their pathogenic potential (Casadevall & Pirofski, 2003; Mims et al.,

2001). Understanding these dynamics is crucial for developing effective disease management

strategies and predicting the emergence of more virulent or drug-resistant strains (Anderson &

May, 1982; Antia et al., 2003).

Moreover, the interplay between microbial virulence and host resistance underscores the

importance of considering both pathogen and host genetics in the study of infectious diseases.

Genetic variations in hosts can influence susceptibility to infections, leading to diverse clinical

outcomes even within the same species (Hill, 2001; Schmid-Hempel & Ebert, 2003). Similarly,

microbial genetic diversity can result in different virulence profiles, impacting the severity and

spread of infections (Mackinnon & Read, 2004; Woolhouse et al., 2002). Therefore, a

comprehensive understanding of virulence requires an integrated approach that examines both

microbial and host factors, as well as their interactions within specific ecological and

evolutionary contexts (Casadevall & Pirofski, 2001; Galvani, 2003).

Infections often progress most critically during early time points, when host responses are most

active, and pathogen replication is rapid. The early infection period is crucial in determining the

trajectory of disease outcomes, as it captures the dynamic interplay between host defenses and

pathogen strategies (Duneau et al., 2017). Metrics that focus solely on survival or endpoint

measures may miss these critical early events, underscoring the need for more nuanced

approaches to study host-pathogen dynamics. In many studies, the quantification of host

susceptibility and pathogen virulence relies on host mortality rates, often using the Cox

proportional hazards or related models (Cox, 1972; Klein & Moeschberger, 2003). This model

offers a robust framework for survival analysis, combining a non-parametric baseline hazard

with a parametric model for covariates proportional to the baseline hazard (Therneau et al.,

2000). Risk scores, crucial in survival analysis, are calculated based on the Cox model (Hosmer

Jr et al., 2008). While risk scores have been employed to assess risk factors, they do not

encompass a quantitative measure of pathogen virulence and solely rely on host mortality

(Bradburn et al., 2003).

Simple statistics such as the proportion of hosts alive at different time points and the time to

death (or survival time) are commonly used to summarize survival data. The proportion alive

provides a straightforward measure of survival at specific times but lacks the nuance to account

for varying follow-up times and censoring (Aalen et al., 2008; Cutler & Ederer, 1958). Time to

death offers more detailed information but is limited by the need for complete survival data, may

not accurately reflect the impact of covariates (Altman, 1990) and is inappropriate for pathogens

where hosts can make a full recovery. The Kaplan-Meier estimator is another widely used

method in survival analysis that provides an estimate of the survival function from lifetime data

(Kaplan & Meier, 1958). This non-parametric statistic handles censored data effectively and

allows for the estimation of survival probabilities at various time points. However, it does not

account for the effects of multiple covariates simultaneously, limiting its utility in multifactorial

analyses (Collett, 2023; Pocock et al., 2002). To address these limitations, the Cox proportional

hazards model offers a more comprehensive approach by incorporating multiple covariates into

the survival analysis, thus providing insights into the factors that influence host mortality rates

(Cox, 1972). However, while this model is powerful for identifying risk factors, it does not

directly quantify pathogen virulence, necessitating additional methods to capture the full

complexity of host-pathogen interactions (Hosmer Jr et al., 2008).

Casadevall, 2017 introduced the concept of the pathogenic potential of a (single) microbe,

highlighting the variable nature of host damage during host-microbe interactions. Clinical

symptoms manifest when host damage disrupts homeostasis, leading to death when damage

surpasses the point of repair and return to homeostasis becomes impossible. Pathogenic potential

is conceptualized as a composite of host susceptibility and pathogen virulence, expressed as

$PP_T = \frac{F_S}{I} 10^M \frac{1}{T}$, where PP represents pathogenic potential, F_S denotes the fraction symptomatic

(host susceptibility factor), I is the infective dose (or pathogen burden), M is the mortality

fraction and T represents time to disease manifestation (Casadevall, 2017). In this model, the

fraction symptomatic host (F_S) is a measure of host susceptibility, referring to the inherent or

acquired traits of the host that affect its vulnerability to infection and disease progression. These

traits can include genetic factors, immune status, and existing health conditions that influence

how severely the host is affected by the pathogen (Casadevall, 2017). The infective dose (I) is a

measure of pathogen virulence, pertaining to the ability of the pathogen to cause damage to the

host, which is determined by its mechanisms of infection, replication rate, and production of

virulence factors that harm the host (Casadevall, 2017). The mortality fraction (M) is a measure

of the lethality of the infection, indicating the proportion of infected hosts that succumb to the

disease (Casadevall, 2017). The time to disease (T) captures the duration from infection to the

onset of clinical symptoms, adding a temporal dimension to the pathogenic potential and

reflecting the speed of disease progression (Casadevall, 2017).

To fully understand the dynamics of host-pathogen interactions, is it crucial to distinguish between host-based and pathogen-based measures of pathogenicity? Or, do these metrics ultimately capture similar aspects of disease dynamics? Host-based measures focus on the response of the host organism to the pathogen, including metrics such as survival rates, risk scores, and the proportion of symptomatic individuals (Schmid-Hempel, 2009). These measures provide insight into how the host's health and survival are affected by infection, often using tools like Kaplan-Meier survival curves and Cox proportional hazards models to analyze mortality data (Cox, 1972; Kaplan & Meier, 1958). But do they truly reflect the virulence of the pathogen itself, or are they more indicative of the host's capacity to withstand infection (Kleinbaum & Klein, 1996)? In contrast, pathogen-based measures aim to quantify the inherent virulence of the pathogen, independent of the host's response. This includes assessing the pathogen's ability to cause disease at various doses and its potential to induce host mortality (Mideo et al., 2008; Read, 1994). Are these distinctions meaningful, particularly in cases of high-dose infections where it becomes difficult to attribute mortality to a specific pathogen due to overwhelming pathogen loads (Frank & Schmid☐Hempel, 2008)? By exploring these measures, can we better understand the relationship between host resilience and pathogen virulence, or are these categories less distinct than they initially appear?

Here, we set out to investigate the informativeness of quantitative measures of pathogenicity, including simple metrics of survival, cox survival analysis, and pathogen potential and its related measures. We conducted infections of *Drosophila melanogaster* using *Providencia* species as a pathogen due to their varied levels of virulence and interaction with the host immune system.

The *Providencia* genus offers a compelling system for investigating these dynamics. These

bacteria exhibit significant genetic diversity in virulence, with some species causing highly lethal

infections, while others result in moderate or minimal host damage (Galac & Lazzaro, 2011). *P.*

sneebia and *P. alcalifaciens* are highly pathogenic to *D. melanogaster*, with even low doses

leading to lethal infections (Galac & Lazzaro, 2011). In contrast, *P. burhodogranariea* Strain D

exhibits lower virulence, making it a suitable model for studying infections that result in

moderate survival rates (Galac & Lazzaro, 2011). This variation across species provides a

powerful framework to examine the genetic and evolutionary mechanisms underlying differences

in pathogen virulence and host susceptibility. *Providencia rettgeri* causes differential survival

outcomes in *D. melanogaster* depending on the host's *dpt* genotype, making it an excellent

candidate for examining host genotype impact influences on infection dynamics (Hanson et al.,

2019, 2023; Mullinax et al., 2023; Unckless et al., 2015, 2016). Additionally, the genetic

diversity within the host population for the *dpt* genotypes, further enriches this model, allowing

us to dissect the contributions of specific alleles to infection outcomes (Hanson et al., 2023;

Mullinax et al., 2023). The wildtype serine allele (dpt^{S69}) is associated with higher resistance to

P. rettgeri, while the segregating arginine allele (dpt^{S69R}) confers increased susceptibility

(Hanson et al., 2023; Mullinax et al., 2023; Unckless et al., 2016). The null allele (Δdpt) results

in nearly all flies succumbing to infection within two days, highlighting the crucial role of the

dpt gene in mediating immune responses (Hanson et al., 2023; Mullinax et al., 2023; Unckless et

al., 2016). This variation allows for a robust analysis of how host genetics influence

susceptibility and resistance to bacterial infections. Our approach allowed us to vary infection

outcomes along three axes: the intrinsic virulence of the pathogen, the dose of the pathogen, and

the susceptibility of the host. We then compared pathogenicity metrics across these three axes.

We measured survival against bacterial infections for four different species of *Providencia*

injected in variable doses (different ODs) in flies of three different *dpt* genotypes (homozygous

serine *dpt* (wildtype, dpt^{S69}), homozygous arginine *dpt* (dpt^{S69R}), and *dpt* null (Δdpt)) in a

controlled genetic background. The *Providencia* strains we used represent a range from avirulent

(*P. burhodogranariea* strain D) to very virulent (*P. alcalifaciens* and *P. sneebia*). The *dpt*

genotypes are an allelic series at least in relation to *P. rettgeri* (Mullinax et al., 2023; Unckless et

al., 2016). Finally, we considered advantages and limitations of our different pathogenicity

measures, offering recommendations on when to use each method of analysis.

Figure 1: A. Experimental design (made using BioRender). 10 flies/vial were infected with variable pathogen strains (*P. burhodograneria* Strain D, *P. rettgeri, P. sneebia, P. alcalifaciens*) with different loads of infective inoculum (OD) and each vial was recorded for upto 7 days post-infection for mortality and fraction symptomatic (using a climbing assay – how many flies can climb up the vial in 5 seconds). Risk scores are further computed using mortality and pathogen potential is calculated using inoculum load, mortality, and fraction symptomatic. **B.** Plot of the proportion of flies alive on day 4 post infection for various host genotypes (homozygous serine *dpt* (wildtype, *dpt^{S69}*), homozygous arginine *dpt* (*dpt^{S69R}*), and *dpt* null (*Δdpt*)), and variable pathogen strains (*P. burhodograneria* Strain D, *P. rettgeri, P. sneebia, P. alcalifaciens*) with different loads of infective inoculum (OD). The error bars represent the 95% confidence interval around the mean of the proportion of flies alive on day 4 post infection.

Materials & methods:

<u>Fly stocks and maintenance</u>: We used CRISPR/Cas9 edited flies with modified *diptericin A* gene

(Mullinax et al., 2023). Briefly, after editing, flies were isogenized for the X and 3^{rd}

chromosomes using a marked strain that allowed us to replace with a specific line's

chromosomes. We did not attempt to do this for the 2^{nd} chromosome. Thus, the second

chromosome may contain off-target edits or be an altogether different chromosome if the

injected line was segregating for multiple 2^{nd} chromosomes. The latter possibility is unlikely

because we Sanger sequenced several immune genes on the 2^{nd} chromosome for our edited lines

and all sequences were identical across all the lines. As with any CRISPR/Cas9 editing

experiment, we cannot rule out off-target edits, but if they exist, it only slightly changes our

interpretation: instead of measuring variation in *dpt* genotype, we are measuring variation in a

second chromosome composite genotype. We used three lines of *D. melanogaster,* with the same

genetic background, but polymorphic for *diptericin A* gene – homozygous serine *dpt* (wildtype,

dpt^{S69}), homozygous arginine *dpt* (*dpt^{S69R}*), and *dpt* null (*Δdpt,* refers to lines with a 3 base pair

deletion). Flies were maintained in a 23°C incubator with a 12h light:12h dark schedule on a

cornmeal-molasses-yeast diet (64.3g/L cornmeal, 79.7mL/L molasses, 35.9g/L yeast, 8g/L agar,

15.4mL of food acid mix (50mL Phosphoric Acid + 418mL Propionic Acid + 532mL deionized

water) and 1g/L Tegosept.

<u>Bacterial strains</u>: The following bacteria were used for systemic infections for this study:

Providencia rettgeri (isolated from wild-caught *D. melanogaster* by B. Lazzaro in State College,

PA, USA), *Providencia burhodograneria* Strain D (isolated from the hemolymph of wild

Drosophila; P. Juneja and B. Lazzaro), *Providencia sneebia* (isolated from the hemolymph of

wild *Drosophila*; P. Juneja and B. Lazzaro), and *Providencia alcalifaciens* (isolated from wild-

caught *D. melanogaster* by P. Juneja and S. Short in Ithaca, NY, USA) (Galac & Lazzaro, 2011;

Juneja & Lazzaro, 2009; Lazzaro, 2002). Bacteria were grown from glycerol stocks on LB plates

at 37°C overnight. Individual colonies of bacteria were picked and grown in 2mL LB broth

shaking overnight at 37°C for preparing infection cultures. The optical density (A_{600nm}) of the

suspension was adjusted according to the desired infection dose.

Infection assays: We used bacterial suspensions for *P. rettgeri, P. burhodograneria* Strain D, *P.*

sneebia, and *P. alcalifaciens* at an Optical Density (A_{600nm}) of approximately 0.01, 0.1, 0.5, 1,

and 8 for our infections (we use "approximate" concentrations because it is difficult to achieve an

exact OD_{600} in bacterial culture preparation, however, the exact concentrations for each treatment

are recorded and available in our data file in the column "OD"). We infected a total of 50 flies

(10 flies per vial) per genotype per bacteria per optical density. We also infected a total of 50

flies (10 flies per vial) per genotypes with LB media as a control. To induce systemic infection,

5-9 day old flies were pricked in the thorax with a needle dipped in a bacterial suspension. We

also used LB media instead of bacterial suspension to poke flies as a control. Flies were

incubated post infection at 23°C at a density of 10flies/vial with a 12h light:12h dark schedule

and survival was tracked daily for 7 days post infection. The experiments were randomized and

conducted across 8 different days.

We used a climbing assay to estimate the fraction of symptomatic (sick) flies following

infection, with mortality included as the ultimate symptom. Each day, for seven days post-

infection, we assessed the proportion of flies in each vial capable of climbing the vial walls

within 5 seconds after being tapped to the bottom. Flies that failed to climb were recorded as

symptomatic, and those that had died were also included in the calculation as symptomatic, given

that mortality is indicative of severe disease progression. Figure 1A represents a schematic of our

experimental design.

Bacterial load: To measure actual dosing in newly infected flies, single wildtype, dpt^{S69} flies

were pricked using *P. rettgeri* and *P. sneebia* cultures at an optical density (A_{600nm}) of

approximately 0.01, 0.1, 0.5, 1, and 8, (exact concentrations for each treatment are recorded and

available in our data file in the column "OD") followed by surface sterilization by washing in

70% ethanol and molecular grade water, and then, homogenized in 500μL LB and plated on LB

plates immediately after the infection using a WASP touch spiral plater from Don Whitely

Scientific. The LB agar plates were incubated overnight at 37°C. Gut commensal bacteria grow

more slowly than *Providencia* under these conditions, so by limiting incubation to overnight we

exclude any commensal bacteria from our assay. The number of colony forming units (CFU) on

each plate was recorded using a Neutec Flash and Go colony counter. We used 5 replicates

(using single flies) per bacteria per A_{600nm} for quantifying the systemic bacterial load in infected

flies. We chose to measure bacterial load for only two species in this study (*P. rettgeri* and *P.*

sneebia) because, based on previous studies (Beal et al., 2020), it is established that bacterial

load does not vary significantly across species within the same genus at the same optical

densities (ODs). As a result, we did not perform bacterial load assays for the other two

pathogens, *P. burhodograneria* Strain D and *P. alcalifaciens*, as we did not expect different

outcomes at equivalent ODs.

<u>Data and statistical analysis</u>: We fitted our data to a mixed effects cox regression model to

analyze the effect of the type and dose of pathogen and the host genotype. We performed a

principal component analysis and a correlation matrix to look at the variation between the

different measures of virulence and pathogenicity. Data analysis performed done in R version

4.3.1 (R Core Team, 2013). Data visualization was performed using R packages – ggplot2

(Wickham, 2011), survminer (Kassambara et al., 2017), wesanderson (Ram & Wickham, 2018),

ggpubr (Kassambara & Kassambara, 2020), corrplot (Wei et al., 2017), RColorBrewer (Neuwirth

& Neuwirth, 2014) and ggrepel (Slowikowski et al., 2018). Data was fitted to a cox-proportional

hazards model to analyze the effect of host genotype, pathogen type and dose on survival [Table

S2]. Statistical analysis was performed using R packages – lme4 (Bates, 2010) and coxme

(Therneau & Therneau, 2015). We have included date as a random factor in all statistical

analyses to account for variability across experimental days.

Results:

Host genotype, dose and pathogen strain interact to influence survival against infection

We measured survival against bacterial infections for four different species of *Providencia*

injected in variable doses (different ODs) in flies of three different *dpt* genotypes (homozygous

serine *dpt* (wildtype, dpt^{S69}), homozygous arginine *dpt* (dpt^{S69R}), and *dpt* null (Δdpt)) in a

controlled genetic background. These *Providencia* strains represent a range from avirulent (*P.*

burhodogranereia) to highly virulent (*P. alcalifaciens* and *P. sneebia*) bacteria (Galac &

Lazzaro, 2011). The host *dpt* genotypes are an allelic series at least in relation to *P. rettgeri*

where the wildtype serine allele is most resistant, the segregating arginine allele is more

susceptible, and the null allele results in almost all flies dying within two days of infection

(Mullinax et al., 2023). Our results from our infection experiments align with previously

published findings - (i) flies homozygous for the serine allele survive systemic infection with *P.*

rettgeri much better than those homozygous for the arginine allele (Unckless et al., 2016), (ii) *P.*

alcalifaciens and *P. sneebia* are highly pathogenic to *D. melanogaster*, where small doses of

infection with these bacteria can result in lethal infections (Galac & Lazzaro, 2011), (iii) most

flies survive infections against *P. burhodograneria* Strain D at moderate doses (Galac &

Lazzaro, 2011) [Figure 1B, Figure 2A].

Figure 2: Host-based measures of pathogenicity: A. Survival curves show proportion alive 0-7 days post infection in various host genotypes (homozygous serine *dpt* (wildtype, *dpt^S69^*), homozygous arginine *dpt* (*dpt^S69R^*), and *dpt* null (*Δdpt*)), and variable pathogen strains (*P. burhodograneria* Strain D, *P. rettgeri, P. sneebia, P. alcalifaciens*) with different loads of infective inoculum (OD) ranging from OD_{600} = 0-8. **B.** Fraction symptomatic four days post-infection in the same assays as A calculated following climbing assays. **C.** Risk scores for the same assays as A calculated based on the Cox models.

We surveyed survival against infection using Kaplan-Meier survival curves and proportion alive

on day 4 post-infection to assess the effect of host genotype, dose, and pathogen strain [Figure

1B, Figure 2A]. We rationalized that 4 days post infection was a reasonable snapshot because

most flies destined to die had died by that point so the interaction between host immunity and

pathogen was mostly over. We calculated the fraction of flies that died between day 4 and day 7

post-infection to justify this timepoint and only 2% of flies die between day 4 and day 7 post-

infection. Very small doses of *P. alcalifaciens* were sufficient to cause high mortality in flies

regardless of their genotype. We saw a range of rates of mortality for survival against *P. sneebia*

across genotypes and across different doses of infection. While for survival against *P.*

burhodograneria Strain D, even high doses of injection did not prove to be lethal for most of the

flies. We also quantified the bacterial load immediately after infection for the range of doses

(optical densities) used in our assays. Dose was not linear with the optical density, instead the

optical density was linearly related to log transformation of the dose (Figure S1, Table S1). We

looked at the effect of host genotype, pathogen type and dose on survival using a cox-

proportional hazards model [Table S2]. First, both the dose (OD) and pathogen species

significantly affect survival, with *P. alcalifaciens* showing the highest pathogenicity (Table S2:

coef = 1.273, p < 0.001). Second, host genotype markedly influences survival outcomes: *Δdpt*

flies showed a higher mortality against all pathogens except *P. sneebia*, while the genotypic

difference between dpt^{S69} and dpt^{S69R} did not significantly influence survival against the different

pathogen strains except for *P. rettgeri* where flies with the serine allele (dpt^{S69}) have significantly

better survival rates compared to those with the arginine allele (dpt^{S69R}) or null allele (Δdpt)

(Table S2: GenotypeDptS: coef = -1.116, p = 0.001). Third, interactions between OD, treatment,

and genotype underscore the complexity of these relationships, highlighting specific

combinations that significantly alter survival rates (Table S2: OD1 * Treatment*P. rettgeri* *

GenotypeDptS (coef = -1.566, p = 0.005)).

Figure 3: Pathogen-based measures of pathogenicity: A. Pathogen potential four days post-infection calculated based on the inoculum load, host mortality and fraction symptomatic for various host genotypes (homozygous serine *dpt* (wildtype, dpt^{S69}), homozygous arginine *dpt* (dpt^{S69R}), and *dpt* null (*Δdpt*)), and variable pathogen strains (*P. burhodograneria* Strain D, *P. rettgeri, P. sneebia, P. alcalifaciens*) with different loads of infective inoculum (OD) ranging from OD_{600} = 0-8. **B.** Pathogen potential on median survival day (T) calculated based on the inoculum load, host mortality, fraction symptomatic and day post-infection for the same assays as A. **C.** Fraction symptomatic on median survival day (T) for the same assays as A calculated following climbing assays.

Limitations of measures of pathogenicity

Host-based measures: We examined Kaplan-Meier survival curves, risk scores, and fraction

symptomatic to assess host-based measures of pathogenicity [Figure 2]. Both risk scores and

fraction symptomatic follow an approximately inverse relationship to host survival upon

infection [Figure 2]. Fraction symptomatic is least sensitive to host genotype out of the three

measures of host susceptibility [Figure 2]. Risk scores show variation with both the dose and

virulence of pathogens but fail to resolve the interaction between the dose of a pathogen and the

host genotype [Figure 2]. These host-based measures of pathogenicity are easy to interpret, but

they tell us little about pathogen virulence of individual microbes. It becomes difficult to

determine whether the differences in survival seen in the host is due to a highly virulent pathogen

or infection with a high dose of a pathogen. Most pathogens become virulent enough to cause

visible symptoms/mortality when administered to a single host at high doses.

Pathogen-based measures: We looked at the pathogen potential and related metrics to assess

pathogen-based measures of pathogenicity [Figure 3]. We quantified the pathogen potential on

day 4 post-infection, and on the day of median survival for each vial. Pathogen potential was

proposed as a tool for adding quantitative rigor to comparisons in microbial pathogenesis

(Casadevall, 2017, 2022; Casadevall et al., 2011; Casadevall & Pirofski, 1999, 2001, 2003;

Smith & Casadevall, 2022). Pathogen potential is helpful in analyzing the dose-dependent effect

of pathogen on host mortality, but there are limitations to the measure. Pathogen potential failed

to capture the variation between host genotypes [Figure 4A]. The limitation in assessing

pathogen potential arises when evaluating high doses or highly virulent microbes at moderate

doses, wherein the likelihood of a specific pathogenic microbe being deemed the primary cause

of mortality approaches zero, even though all individuals die. This counterintuitive phenomenon

stems from the low probability that any specific pathogen cell within the high dose is the sole

contributor to mortality. While this concept aligns with the understanding that the pathogen

responsible for mortality may not be easily pinpointed, it presents a challenge in practical utility

in our context, as the assessment becomes less informative with more virulent pathogens.

Figure 4: A. P-values from ANOVA to indicate the significance of each factor corresponding to each measure of pathogenicity (see Table S3 for more information), B. Heatmap showing R^2 (R=correlation coefficient) values to assess the differences between different measures of host susceptibility (survival, risk scores, and fraction symptomatic) and pathogen virulence (pathogen potential) (see Table S4 for more information). [Host-based measures are labelled purple while pathogen-based measures are labelled orange]

In this context, it is important to consider the transparency of the CRISPR methodology

employed in these experiments. Specifically, the application of CRISPR-Cas9 in generating the

dpt mutant strains (Δ*dpt*, *dpt^{S69}*, and *dpt^{S69R}*) could introduce concerns about off-target effects,

particularly since chromosome 2, which houses many immune-related genes, was not

backcrossed. We acknowledge the possibility of off-target effects, particularly considering the

sensitivity of the *dpt^{S69R}* mutant to Gram-positive bacteria, a pathogen type that *dpt* does not

typically impact (Carboni et al., 2022; Hanson et al., 2019). To address these concerns, we

conducted Sanger sequencing to evaluate potential off-target effects. While we cannot entirely

exclude such effects, our data suggest they are minimal. We opted against extensive

backcrossing due to potential confounding genetic variability and focused instead on robust

sequencing to reduce possible confounders. Importantly, even if off-target effects are present,

each of our genotypes represents a distinct chromosome 2 background. Thus, we are not

attributing effects to a single mutation but rather assessing the cumulative impact of the entire

second chromosome.

Comparative analysis: We took a comparative approach to determine how different measures of

host susceptibility (survival, risk scores, and fraction symptomatic) and pathogen virulence

(pathogen potential) were correlated with each other to assess how different they are in their

analytical information. From the R^2 matrix, it was evident that all the different host-based

measures of virulence and susceptibility were highly correlated to each other, but not to the

pathogen-based measures (pathogen potential) [Figure 4B]. To further examine the relationship

among metrics, we performed a principal component analysis (PCA). This revealed three main

groupings: risk score (alone), fraction alive and fraction symptomatic metrics, and pathogen-

potential related metrics [Figure S2]. Pathogen potential metrics clustered away from the host-

based measures as they incorporate quantitative information about the infecting pathogen as well

as host response. Interestingly, risk scores also clustered away from all the different

pathogenicity measures as risk scores were able to capture the information for the variation

between host genotype, pathogen type and dose of the pathogen [Figure 4A, Figure S2].

Discussion:

Our study investigated the complex dynamics of host-pathogen interactions, focusing on the

quantification of host susceptibility and pathogenicity. Traditionally, host mortality rates,

analyzed through the Cox proportional hazards models, have been widely used to assess

virulence. An alternative, pathogenic potential concept, introduced by Casadevall, 2017 offers a

more nuanced perspective by considering the composite nature of host damage. In this study, we

explore the advantages and limitations of traditional survival analysis and risk scores compared

to the quantitative measures like pathogen potential. Our results suggest that, at least for our

study system, the best measure depends on overall host mortality in relation to dose.

Our infection experiments revealed intricate interactions between host genotype, pathogen strain,

and infection dose [Table S2]. First, in most cases, mortality increased with increasing dose and

this was reflected in most of our measures of infection outcomes. However, because pathogen

potential (and related metric) explicitly control for the effect of dose, those measures are less

sensitive to dose variation. Second, host genotype influences the outcomes of infection for most,

but not all infections and infection outcome measures. The observed higher mortality in Δ*dpt*

flies against all pathogens, except *P. sneebia*, underscores the role of the diptericin A gene in

systemic infection resilience (Hanson et al., 2023; Mullinax et al., 2023; Unckless et al., 2016).

Finally, pathogen species consistently displayed variation in pathogenicity (*sensu latu*) across

infection outcome measures. *P. alcalifaciens* demonstrated high pathogenicity even at low doses,

emphasizing the importance of considering both pathogen virulence and infection dose in

assessments.

Traditional survival analysis and risk scores, rooted in Cox models, proved informative for

overall trends and risk assessment. However, they fall short in providing a comprehensive

understanding of pathogen virulence, particularly for avirulent or less virulent pathogens. The

limitations of these host-based measures highlight the necessity for additional quantitative

insights. The pathogen-based measure, pathogen potential, offers a quantitative lens but

encounters limitations at high pathogen doses, where pinpointing the primary cause of mortality

becomes less informative. It becomes evident that each measure contributes unique insights,

emphasizing the importance of a nuanced and multifaceted approach in analyzing host-pathogen

interactions. A combination of pathogen potential and risk scores seemed to be particularly

informative in separating the variation between host susceptibility and pathogen virulence.

Additionally, concepts such as Systemic Pathogen Burden Load (SPBL) and Bacterial Load

Undergoing Dynamics (BLUD) could further enrich the biological context of the manifestation

of symptoms during infection (Duneau et al., 2017). These measures, which focus on pathogen

load and its dynamics over time, offer insights into infection progression and immune response

that complement the survival data and risk scores we used. Although our current study does not

include these measures, this could provide a more detailed understanding of immune responses

in the early phases of infection, which aligns with recent work on time-specific infection

dynamics.

We propose several recommendations for the optimal use of each measure, considering their

strengths and limitations –

(i) <u>Kaplan-Meier Curves (to visualize survival functions):</u> Kaplan-Meier curves are a

robust measure and should be used to assess the overall survival trends over time. It is

particularly useful for understanding the impact of host genotype, dose, and pathogen strain on

the course of infection. **Limitations:** While survival analysis provides valuable information

about the overall outcome, it may not distinguish between the effects of pathogen virulence and

the dose administered.

(ii) <u>Risk Scores from Cox Proportional Hazards Models:</u> Risk scores derived from Cox

proportional hazards models are informative for assessing variation in both dose and virulence of

pathogens. They are especially useful when trying to understand how different combinations of

host genotype, dose, and pathogen strain contribute to the risk of mortality. **Limitations:** Risk

scores may not provide information about the virulence of avirulent or less virulent pathogens,

and their utility diminishes for high doses of less virulent pathogens.

(iii) <u>Fraction Symptomatic:</u> Fraction symptomatic is useful for insights into the

proportion of individuals exhibiting symptoms, giving a qualitative assessment of the infection.

Limitations: Fraction symptomatic is less sensitive to pathogen virulence and may not provide

detailed information about the severity of the infection.

(iv) <u>Pathogen Potential:</u> Pathogen potential is valuable for understanding the dose-

dependent effects of pathogens on host mortality. **Limitations:** Pathogen potential may be less

informative when mortality is high since it loses resolution, where the likelihood of pinpointing a

specific pathogen as the primary cause of mortality decreases.

Survival analysis is robust for overall trends, risk scores offer detailed information on variation,

fraction symptomatic provides qualitative insights, and pathogen potential contributes

quantitative rigor. A comparative analysis aids in choosing the most relevant measure based on

experimental goals.

References:

Aalen, O., Borgan, O., & Gjessing, H. (2008). *Survival and event history analysis: a process point of view*. Springer Science & Business Media.

Altman, D. G. (1990). *Practical statistics for medical research*. Chapman and Hall/CRC.

Anderson, R. M., & May, R. M. (1982). Coevolution of hosts and parasites. *Parasitology*, *85*(2), 411–426.

Antia, R., Regoes, R. R., Koella, J. C., & Bergstrom, C. T. (2003). The role of evolution in the emergence of infectious diseases. *Nature*, *426*(6967), 658–661.

Bates, D. M. (2010). *lme4: Mixed-effects modeling with R*. Springer New York.

Beal, J., Farny, N. G., Haddock-Angelli, T., Selvarajah, V., Baldwin, G. S., Buckley-Taylor, R., Gershater, M., Kiga, D., Marken, J., & Sanchania, V. (2020). Robust estimation of bacterial cell count from optical density. *Communications Biology*, *3*(1), 512.

Bradburn, M. J., Clark, T. G., Love, S. B., & Altman, D. G. (2003). Survival analysis part II: multivariate data analysis–an introduction to concepts and methods. *British Journal of Cancer*, *89*(3), 431–436.

Carboni, A. L., Hanson, M. A., Lindsay, S. A., Wasserman, S. A., & Lemaitre, B. (2022). Cecropins contribute to Drosophila host defense against a subset of fungal and Gram-negative bacterial infection. *Genetics*, *220*(1), iyab188.

Casadevall, A. (2017). The Pathogenic Potential of a Microbe. *MSphere*, *2*(1). https://doi.org/10.1128/msphere.00015-17

Casadevall, A. (2022). *Expanding the Pathogenic Potential Concept To Incorporate Fulminancy,*

Time, and Virulence Factors. https://journals.asm.org/journal/msphere

Casadevall, A., Fang, F. C., & Pirofski, L. (2011). Microbial virulence as an emergent property:

consequences and opportunities. *PLoS Pathogens*, *7*(7), e1002136.

Casadevall, A., & Pirofski, L. (1999). Host-pathogen interactions: redefining the basic concepts of

virulence and pathogenicity. *Infection and Immunity*, *67*(8), 3703–3713.

Casadevall, A., & Pirofski, L. (2001). Host□pathogen interactions: the attributes of virulence. *The*

Journal of Infectious Diseases, *184*(3), 337–344.

Casadevall, A., & Pirofski, L. (2003). The damage-response framework of microbial pathogenesis.

Nature Reviews Microbiology, *1*(1), 17–24.

Collett, D. (2023). *Modelling survival data in medical research*. Chapman and Hall/CRC.

Cox, D. R. (1972). Regression models and life□tables. *Journal of the Royal Statistical Society: Series*

B (Methodological), *34*(2), 187–202.

Cutler, S. J., & Ederer, F. (1958). Maximum utilization of the life table method in analyzing survival.

Journal of Chronic Diseases, *8*(6), 699–712.

Duneau, D., Ferdy, J.-B., Revah, J., Kondolf, H., Ortiz, G. A., Lazzaro, B. P., & Buchon, N. (2017).

Stochastic variation in the initial phase of bacterial infection predicts the probability of survival

in D. melanogaster. *Elife*, *6*, e28298.

Ebert, D., & Bull, J. J. (2003). Challenging the trade-off model for the evolution of virulence: is

virulence management feasible? *Trends in Microbiology*, *11*(1), 15–20.

Ewald, P. W. (1994). *Evolution of infectious disease*. Oxford University Press, USA.

Frank, S. A. (1996). Models of parasite virulence. *The Quarterly Review of Biology*, *71*(1), 37–78.

Frank, S. A., & Schmid☐Hempel, P. (2008). Mechanisms of pathogenesis and the evolution of

parasite virulence. *Journal of Evolutionary Biology*, *21*(2), 396–404.

Galac, M. R., & Lazzaro, B. P. (2011). Comparative pathology of bacteria in the genus Providencia to

a natural host, Drosophila melanogaster. *Microbes and Infection*, *13*(7), 673–683.

Galvani, A. P. (2003). Epidemiology meets evolutionary ecology. *Trends in Ecology & Evolution*,

18(3), 132–139.

Hanson, M. A., Grollmus, L., & Lemaitre, B. (2023). Ecology-relevant bacteria drive the evolution of

host antimicrobial peptides in Drosophila. *Science*, *381*(6655), eadg5725.

Hanson, M. A., Lemaitre, B., & Unckless, R. L. (2019). Dynamic evolution of antimicrobial peptides

underscores trade-offs between immunity and ecological fitness. *Frontiers in Immunology*, *10*,

2620.

Hill, A. V. S. (2001). The genomics and genetics of human infectious disease susceptibility. *Annual*

Review of Genomics and Human Genetics, *2*(1), 373–400.

Hosmer Jr, D. W., Lemeshow, S., & May, S. (2008). *Applied survival analysis: regression modeling*

of time-to-event data (Vol. 618). John Wiley & Sons.

Juneja, P., & Lazzaro, B. P. (2009). Providencia sneebia sp. nov. and Providencia burhodogranariea

sp. nov., isolated from wild Drosophila melanogaster. *International Journal of Systematic and*

Evolutionary Microbiology, *59*(5), 1108–1111.

Kaplan, E. L., & Meier, P. (1958). Nonparametric estimation from incomplete observations. *Journal*

of the American Statistical Association, *53*(282), 457–481.

Kassambara, A., & Kassambara, M. A. (2020). Package 'ggpubr.' *R Package Version 0.1*, *6*(0).

Kassambara, A., Kosinski, M., Biecek, P., & Fabian, Sjdsc. (2017). Package 'survminer.' *Drawing*

Survival Curves Using 'Ggplot2'(R Package Version 03 1), *3*.

Klein, J. P., & Moeschberger, M. L. (2003). *Survival analysis: techniques for censored and truncated*

data (Vol. 1230). Springer.

Kleinbaum, D. G., & Klein, M. (1996). *Survival analysis a self-learning text*. Springer.

Lazzaro, B. P. (2002). *A population and quantitative genetic analysis of the Drosophila melanogaster*

antibacterial immune response. The Pennsylvania State University.

Mackinnon, M. J., & Read, A. F. (2004). Immunity promotes virulence evolution in a malaria model.

PLoS Biology, *2*(9), e230.

Mideo, N., Alizon, S., & Day, T. (2008). Linking within-and between-host dynamics in the

evolutionary epidemiology of infectious diseases. *Trends in Ecology & Evolution*, *23*(9), 511–

517.

Mims, C. A., Nash, A. A., & Stephen, J. (2001). *Mims' pathogenesis of infectious disease*. Gulf

Professional Publishing.

Mullinax, S. R., Darby, A. M., Gupta, A., Chan, P., Smith, B., & Unckless, R. (2023). A suite of

selective pressures supports the maintenance of alleles of a Drosophila immune peptide. *BioRxiv*,

2023–2028.

Neuwirth, E., & Neuwirth, M. E. (2014). Package 'RColorBrewer.' *ColorBrewer Palettes*.

Pocock, S. J., Clayton, T. C., & Altman, D. G. (2002). Survival plots of time-to-event outcomes in

clinical trials: good practice and pitfalls. *The Lancet*, *359*(9318), 1686–1689.

R Core Team, R. (2013). *R: A language and environment for statistical computing*.

Råberg, L., Graham, A. L., & Read, A. F. (2009). Decomposing health: tolerance and resistance to

parasites in animals. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

364(1513), 37–49.

Ram, K., & Wickham, H. (2018). wesanderson: a Wes Anderson palette generator. *R Package*

Version 0.3, *6*, 2018.

Read, A. F. (1994). The evolution of virulence. *Trends in Microbiology*, *2*(3), 73–76.

Schmid-Hempel, P. (2009). Immune defence, parasite evasion strategies and their relevance for

'macroscopic phenomena' such as virulence. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, *364*(1513), 85–98.

Schmid-Hempel, P., & Ebert, D. (2003). On the evolutionary ecology of specific immune defence.

Trends in Ecology & Evolution, *18*(1), 27–32.

Slowikowski, K., Schep, A., Hughes, S., Lukauskas, S., Irisson, J.-O., Kamvar, Z. N., Ryan, T.,

Christophe, D., Hiroaki, Y., & Gramme, P. (2018). Package ggrepel. *Automatically Position*

Non-Overlapping Text Labels with 'ggplot2.

Smith, D. F. Q., & Casadevall, A. (2022). On the relationship between Pathogenic Potential and

Infective Inoculum. *PLoS Pathogens*, *18*(6). https://doi.org/10.1371/journal.ppat.1010484

Therneau, T. M., Grambsch, P. M., Therneau, T. M., & Grambsch, P. M. (2000). *The cox model*.

Springer.

Therneau, T. M., & Therneau, M. T. M. (2015). Package 'coxme.' *R Package Version*, *2*(5).

Unckless, R. L., Howick, V. M., & Lazzaro, B. P. (2016). Convergent Balancing Selection on an

Antimicrobial Peptide in Drosophila. *Current Biology*, *26*(2), 257–262.

https://doi.org/10.1016/j.cub.2015.11.063

Unckless, R. L., Rottschaefer, S. M., & Lazzaro, B. P. (2015). The Complex Contributions of

Genetics and Nutrition to Immunity in Drosophila melanogaster. *PLoS Genetics*, *11*(3).

https://doi.org/10.1371/journal.pgen.1005030

Wei, T., Simko, V., Levy, M., Xie, Y., Jin, Y., & Zemla, J. (2017). Package 'corrplot.' *Statistician*,

56(316), e24.

Wickham, H. (2011). ggplot2. *Wiley Interdisciplinary Reviews: Computational Statistics*, *3*(2), 180–

185.

Woolhouse, M. E. J., Webster, J. P., Domingo, E., Charlesworth, B., & Levin, B. R. (2002).

Biological and biomedical implications of the co-evolution of pathogens and their hosts. *Nature*

Genetics, *32*(4), 569–577.