Coordinate post-transcriptional regulation by microRNAs and RNA binding proteins is critical for early embryonic cell fate decisions

Carolyn Sangokoya [1,2] * and Robert Blelloch [1,2,3] *

[1]The Eli and Edythe Broad Center of Regeneration Medicine and Stem Cell Research, Center for Reproductive Sciences, University of California San Francisco, San Francisco, CA, USA
[2]Department of Pathology, University of California San Francisco, San Francisco, CA, USA
[3]Department of Urology, University of California San Francisco, San Francisco, CA, USA
*Correspondence should be addressed to: Carolyn.Sangokoya@ucsf.edu, Robert.Blelloch@ucsf.edu

Abstract/Summary

Post-transcriptional control by RNA binding proteins (RBPs) and microRNAs play central roles in mRNA stability and translation (1). However, little is known about how RBPs and microRNAs coordinate in developmental time to regulate cell fate. Here, we show that coordinate RBP and microRNA control of a single transcript, Profilin 2 (Pfn2), is essential for differentiation of embryonic stem cells (ESCs) into the primary germ layer lineages. The Pfn2 3'untranslated region has a binding site for Iron Regulatory Proteins and a nearby binding site for ESC enriched microRNAs (2,3). Deletion of this microRNA site leads to increased PFN2 and reduced FGF signaling during pluripotency transition prior to germ layer formation (4). In contrast, deletion of the iron response element leads to decreased PFN2 and a Wnt signaling defect, reduced nuclear beta-catenin, and a subsequent block in mesendodermal lineages during early germ layer formation. The choreographed microRNA-IRE axis of control on the Pfn2 transcript is essential for two key signal transduction steps during ESC differentiation.

Keywords

Post-transcriptional, microRNA, beta-catenin, cell fate, RNA binding proteins, Wnt signaling, embryonic stem cells

Introduction

Mouse embryonic stem cells (ESCs) have the remarkable ability to self-renew indefinitely while retaining the potential to make all downstream lineages of adult body including the germline. To do so, they are typically grown in the presence of LIF and two inhibitors (2i) toward MEK (PD0325901) and GSK3 (CHIR99021) (5). MEK inhibition blocks FGF signaling while GSK3 inhibition promotes canonical Wnt signaling. GSK3 represses canonical signaling by phosphorylating the secondary messenger beta-catenin which directs it to the proteosome for degradation (6–8). ESCs grown under these conditions are called naïve ESCs and are transcriptionally similar to the *in vivo* pluripotent epiblast cells of the peri-implantation embryonic day 4.5 (E4.5) blastocyst (9). Upon removal of LIF and 2i, the naïve ESCs initiate differentiation, first transitioning to a more epithelial-like state resembling the pluripotent epithelial cells of the post-implantation E5.5 egg cylinder stage embryo, also called the formative pluripotent state (10). The cells can be retained in formative-like state by inhibiting Wnt signaling, providing low level Activin signaling, and allowing for autonomous FGF signaling (11). Following release of Wnt signaling, these cells than initiate differentiation into somatic germ layers, recapitulating gastrulation in vivo. This differentiation can occur spontaneously *in vitro* in the form of embryoid bodies or in more directed fashion in presence of specific factors (12–14). These well-defined methods of recapitulating early embryo development in a dish allows for an unprecedented ability to understand fundamental molecular and cellular mechanisms driving cell fate.

Post-transcriptional regulation of mRNA stability and protein translation is thought to play a central role in cell fate decisions, although only a limited number of examples exist. One such example is the role of the embryonic stem cell (ESC)-enriched (ESCC) family of microRNAs that promote pluripotency including the dedifferentiation of somatic cells to induced pluripotent stem cells (2,15,16). MicroRNAs are short non-coding RNAs that function by binding the 3'UTR of target mRNAs, which in turn leads to both destabilization of the RNA and the inhibition of their translation (17–21). A functional target of the ESCC family in promoting pluripotency is the actin/dynamin binding protein Profilin-2 (PFN2) (2). Deeper analysis of PFN2's role showed that deletion of the binding site for the ESCC miRNAs in the 3'UTR of PFN2 transcript leads to upregulation of PFN2 mRNA and protein levels, which in turn inhibits endocytosis-directed FGF signaling and thus the transition from naïve to formative pluripotency (4). There is also an iron response element (IRE) in the 3'UTR of PFN2 (3). This element is less from than 100bp from the ESCC miRNA binding element and is conserved through humans (Fig. 1a). IREs are target sites for the RNA-binding proteins (RBPs) Iron Response Proteins (IRP) 1 and 2. In contrast to microRNAs, binding of IRPs to IRE sites in the 3'UTR of a transcript typically results in its stabilization (22).

The presence of both a microRNA and an RBP site in close proximity on the 3' UTR on the PFN2 target transcript raises the question of how they might be interacting to regulate ESC self-renewal and differentiation. Therefore, we deleted the IRE element in the PFN2 3'UTR and compared the consequences to loss of the ESCC site. Unlike loss of the ESCC site, disruption of the IRE site had no impact on PFN2 mRNA and protein levels in naïve ESCs. However, upon differentiation to the formative state and further into the three germ layers, disruption of the IRE element results in a reduction of PFN2 mRNA and protein levels. This reduction results in an inhibition of differentiation toward the three somatic germ layers, especially those of the primitive streak, mesoderm and endoderm. Additionally, this defect is associated with a decrease in Wnt signaling that is cell autonomous and is downstream of GSK3β directed degradation of beta-catenin. There is a decrease in beta-catenin in the nucleus, where it would normally be required to induce the gene expression program of mesendoderm differentiation. Therefore, these data show how the 3'UTR of a single gene can help direct the switch between pluripotency and the trilineage differentiation associated with gastrulation.

Results

Divergent post-transcriptional regulation of Profilin-2

To evaluate the role for the *Pfn2* 3'UTR iron response element (IRE) in ESC self-renewal and differentiation, we used CRISPR-Cas9 mutagenesis to delete this IRE site, generating PFN2-3UTR ΔIRE mutant ESCs. These cells were subcloned to produce a stable mutant cell line (Extended Data Fig. 1a,b)). The PFN2-3UTR ΔIRE mutant ESCs were compared to matching wild-type control ESCs and previously produced PFN2-3UTR ΔESCC mutant ESCs (4). When cultured in naïve pluripotency media conditions (LIF+2i), the PFN2-3UTR ΔIRE mutant ESCs showed no significant difference in PFN2 mRNA and protein levels relative to corresponding wild-type cells (Fig. 1b-d). In contrast the PFN2-3UTR ΔESCC mutant ESCs showed elevated PFN2 mRNA and protein levels as previously described (4). To evaluate changes during ESC differentiation, we performed a time course of embryoid body (EB) differentiation (Fig 1b). Upon EB differentiation, wild-type ESCs show an approximately two-fold increase in Profilin-2 mRNA levels and three to four-fold increase in protein levels starting at day 2 (EBD2, representing the naïve to formative pluripotency transition) through day 6 (EBD6, representing germ layer specification) (Fig. 1c,d). This normal increase in Pfn2 with differentiation is lost in PFN2-3UTR ΔIRE mutant cells (Fig. 1c,d). In contrast, PFN2-3UTR ΔESCC cells show similar levels of PFN2 mRNA and protein as wild-type cells at EBD2 and EBD6. These findings uncover divergent roles for the ESCC miRNA binding and IRE sites in the 3'UTR in PFN2. The ESCC miRNA site suppresses PFN2 levels in naïve ESCs, while the IRE site is essential for the stability of the mRNA and upregulation of PFN2 protein upon ESC differentiation.

Loss of the PFN2-3UTR IRE site inhibits ESC differentiation

To characterize the impact of the PFN2-3UTR IRE site on the formation of the three germ layers, we performed embryoid body differentiation of wild-type and PFN2-3UTR ΔIRE mutants at EBD6 followed by RNA-seq. Over 4000 genes were dysregulated in the mutants (1815 up, 2261 down, FDR <0.05) (Fig. 2a). Compared to wild-type cells, PFN2-3UTR ΔIRE mutants showed reduced expression of markers for all 3 germ layers including mesoderm (*Mixl1, Col4a1, Col4a2*), endoderm (*Pdgfra, Sox17, Hnf4a, Hnf1b*) and ectoderm (*Sox11, Pax6*), (Fig. 2a). In contrast, pluripotency markers were abnormally elevated. Gene ontology analysis showed enrichment for categories such as cell migration involved in gastrulation, endoderm formation, and mesoderm formation (Fig. 2b). A closer look at well-described markers of naïve and formative pluripotency and of the three germ layers confirmed elevated expression of naive ESC markers and reduced expression of formative and germ layer markers (Fig. 2c). Specifically, naïve-specific markers such as *Esrrb, Dppa4, Utf1,* and *Prdm14* were up, while formative markers such as *Fgf5* and *Dnmt3b* were down. Mesoderm (e.g. *Mixl1, Col4a1, Col4a2, Eomes*, and *T*), endoderm (e.g. *Sox17, Gata4, Hnf4a/b, Foxa2*), and ectoderm markers (e.g. *Pax6, Ncam1, Sox1* and *Krt18*) were consistently down (Fig. 2d,e). qRT-PCR analysis for a subset of these markers confirmed these findings. In contrast, while the ESCC site mutants showed reduced expression of the ectoderm marker Pax6, they showed normal levels of mesoderm (Mixl1) and endoderm (Gata4, Ttr) markers (Extended Data Fig. 2). Similarly, immunoblot analysis showed that proteins associated with mesoderm (ACTA2) and endoderm (FOXA2, GATA4) were reduced in the IRE site mutant cells at EBD6 (Fig 2f). A similar phenotype was seen with *Pfn2* knockout cells. In contrast, the ESCC site mutants showed abnormal upregulation of these proteins. Together, these results show a striking defect in ESC differentiation with the loss of the IRE site in the *Pfn2* 3'UTR. In contrast, loss of the ESCC site appears to result in a defect in ectoderm differentiation consistent with the prior report of an FGF-related defect in the naïve to formative pluripotency differentiation (4). Therefore, these two 3'UTR elements have divergent impact on both PFN2 levels and cell fate during ESC differentiation.

Loss of the PFN2-3UTR IRE site results in a cell autonomous defect in Wnt signaling

We noticed the PFN2-3UTR ΔIRE mutant EBs were morphologically abnormal compared to wild-type between EBD6 and EBD6.5, showing a loose dissociation at the outer surface of the EBs followed by failure to differentiate or cavitate (Fig. 3a-d). This dysmorphic defect appeared similar to a previously reported morphological defect associated with EB differentiation of ESCs deficient in *Ctnnb1* (beta-catenin) (23). Beta-catenin plays important roles in both cell adhesion through its interaction with cadherins at cell-cell junctions and in cell signaling as a secondary messenger in canonical Wnt signaling (23,24). Importantly, Wnt signaling is essential for mesendoderm differentiation during ESC differentiation (25). Given the combined PFN2-3UTR ΔIRE defects in EB morphogenesis, differentiation into mesoderm and endoderm lineages, we hypothesized a role for increased PFN2 in regulating beta-catenin and enabling differentiation of ESCs into the three germ layers.

Beta-catenin levels are regulated by GSK3β-directed phosphorylation, a post-translational modification that directs beta-catenin to the proteosome for degradation. In the absence of Wnt signaling, beta-catenin is associated with the destruction complex that includes sequential phosphorylation of Thr41, Ser37, and Ser33 by GSK3β (26). The presence of Wnt-signaling blocks this beta-catenin phosphorylation and degradation (27). Wnt/β-catenin signaling and subsequent β-catenin/TCF transcriptional activation are specifically mediated through the molecular form of β-catenin that remains unphosphorylated at residues 37 and 41 (28,29) often referred to as the signaling form of β-catenin, or Active β-Catenin (ABC). Therefore, to test our hypothesis, we performed analysis of intracellular unphosphorylated (active) beta catenin levels at single cell resolution by flow cytometry at EBD2 and at EBD6 (Fig. 3e), At EBD2, the intracellular levels of active beta-catenin in wild-type and PFN2-3UTR ΔIRE mutant EBs were not significantly different, while at EBD6, active beta-catenin levels were significantly lower in the PFN2-3UTR ΔIRE mutant cells. (Fig. 3f-h). In contrast, active beta-catenin levels were not impacted in PFN2-3UTR ΔESCC mutants at either EBD2 or EBD6 (Extended Data Fig. 3a-c).

Next, we asked if the decrease in active beta-catenin was cell autonomous (e.g. defects in signaling downstream of the Wnt receptor or non-autonomous (e.g. defect in Wnt secretion). To do so, we produced chimeric EBs mixing mutant with wild-type cells expressing distinct color labels (Extended Data Fig. 3d). Chimeric EBs composed of both wild-type and PFN2-3UTR ΔIRE showed less of a morphological phenotype than EBs solely made up of PFN2-3UTR ΔIRE cells, although their surfaces still appeared less sharp than WT+WT or WT+PFN2-3UTR ΔESCC chimeric EBs (Extended Data Fig. 3e-g). Also, there was skewing of PFN2-3UTR ΔIRE cells toward the center of the EBs (Extended Data Fig. 3h-j). Next, we dissociated the EBs into single cells and performed flow cytometric analysis for the color labels, intracellular beta-catenin, and intracellular EOMES as a marker of mesendoderm differentiation (Fig. 3i). This analysis showed that the PFN2-3UTR ΔIRE, but not wild-type or PFN2-3UTR ΔESCC cells isolated from the chimeras showed reduced active and total beta-catenin (Fig. 3j,k, Extended Data Fig. 3k,l). Furthermore, unlike the wild-type or PFN2-3UTR ΔESCC cells, the PFN2-3UTR ΔIRE cells derived from the chimeras failed to upregulate EOMES, (Fig. 3l,m). Together, these data uncover a cell-autonomous defect in active beta-catenin signaling downstream of the Wnt receptor in the PFN2-3UTR ΔIRE background.

PFN2-3UTR IRE site functions downstream of GSK3 to stabilize beta-catenin

To look further into mechanism by which PFN2 impacts beta-catenin and cell differentiation, we transitioned to a directed mesendodermal differentiation protocol (Fig. 4a). EBs were produced as above, but at day 3, CHIR and Activin were added to the media to specifically induce mesendoderm differentiation (25,30–33). T Brachyury (*Tbxt*) upregulation was used as the readout of successful

differentiation into early mesendodermal progenitors (25,32). T Brachyury is also a direct target of Wnt induced beta-catenin signaling (34). Wild-type cells showed an over 30x increase in *Tbxt* (Fig. 4b). In contrast, PFN2-3UTR ΔIRE cells showed greatly reduced induction (~2x). Intracellular flow cytometry of active beta-catenin remained reduced in PFN2-3UTR ΔIRE relative to wild-type cells even in the presence of the GSK3 inhibitor (Fig. 4c,d). To further confirm the cell autonomous nature of the defect, we repeated the chimera experiments, now under directed mesendodermal culture conditions. Similar to undirected EB differentiation, combining PFN2-3UTR ΔIRE and wild-type cells failed to rescue either active beta-catenin or total beta-catenin levels in directed differentiation conditions (Fig. 4e-g). By contrast, there were no significant differences between wild-type and PFN2-3UTR ΔESCC mutants within chimeras at EBD4M. Together, these results show that PFN2 is acting downstream of GSK3β directed beta-catenin degradation to regulate Wnt-signaling and thus mesendodermal differentiation.

PFN2-3UTR IRE regulates nuclear beta-catenin levels
Wnt/beta-catenin signaling requires the transport of beta-catenin into the nucleus where it acts as a co-transcription factor in association with TCF/LEF proteins (Fig 5a). Given PFN2's known association with the cytoskeletal network (35), we hypothesized that destabilization of beta-catenin could be secondary to a failure of beta-catenin transport to or retention in the nucleus. To test this hypothesis, we once again performed directed mesendoderm differentiation. EBs were then fractioned into nuclear, cytoplasmic, and membrane fractions. Immunoblot analysis showed a reduction in nuclear active beta-catenin and T Brachyury (Fig. 5b-d). To further characterize, we performed a time-course immediately following CHIR-Activin induction at EBD3, collecting samples at 2, 6, and 18 hours post-induction (Fig. 5e-g). Nuclear active beta-catenin peaked at 18H in wild-type cells and was ~3x fold reduced in PFN2-3UTR ΔIRE cells (Fig. 5h). In contrast, cytoplasmic levels of either active or total beta-catenin were not significantly different between the wild-type and mutant cells (Fig. 5i-j). Nuclear T Brachyury levels similarly peaked at 16 hours and were dramatically reduced in PFN2-3UTR ΔIRE cells (Fig. 5k). ELISA of PFN2 itself showed reduced levels in both the cytoplasmic and nuclear compartments during the induction time course and at the EBD4M endpoint (Fig. 5l-m). Together, these results show a defect in either the transport or retention of nuclear beta-catenin associated with the reduced levels of PFN2 in PFN2-3UTR ΔIRE cells.

Discussion
Together with the recent dissection of the ESCC site in the 3'UTR of PFN2 (4), the findings reported here uncover a coordinate role of a miRNA and an RNA binding protein on the 3'UTR of a single gene in regulating the switch from pluripotency to somatic differentiation (Fig. 6). In pluripotent cells, the ESCC family of miRNAs destabilize PFN2, allowing endocytosis which is essential for normal FGF/ERK signaling (4). This signaling is important for the normal transition from naïve to formative pluripotency (4,36–39). Following the transition, an IRE which is bound by iron response proteins is required for an increase in PFN2 levels, in turn promoting Wnt/beta-catenin signaling. Defects in the Wnt/beta-catenin due to deletion of the IRE result in a block in activation of the mesendoderm transcriptional program that normally occurs with gastrulation. The impact of PFN2 on beta-catenin signaling is downstream of GSK driven degradation of the protein and is associated with alterations in the nuclear/cytoplasmic localization of the protein suggesting a role in the transport and/or retention of beta-catenin in the nucleus. Thus, our findings show how the combinatorial post-transcriptional control of PFN2 plays a critical role in early mammalian cell fate decisions.

Previous examples of microRNA and RBP coordination on a 3'UTR largely involve RBPs enhancing miRNA targeting by improving target site accessibility (40–42) or suppressing microRNA targeting by competitive engagement of overlapping binding sites (42–47). In the setting of development, for example, the RNA-binding protein Dead end (Dnd1) prohibits the function of several miR-430 and miR-372-family miRNAs by blocking their accessibility to transcript target sites, and this regulation is essential for proper development of primordial germ cells (43). In the case of PFN2, the RBP and microRNA binding sites are not overlapping, but instead are ~100 bases apart and are acting at distinct developmental times. Therefore, it seems unlikely their opposing impacts on PFN2 levels are due to competitive binding. Instead, these sites reveal a developmentally relevant switch where the actions of the ESCC microRNAs on PFN2 during pluripotency 'switches' to predominant role for IRP binding during somatic differentiation into the three germ layers. We propose this model could provide a paradigm for other miRNA-RBP pairs playing critical roles in cell fate decisions throughout development.

Our data further shows that PFN2 regulates the transport or retention of beta-catenin in the nucleus. Reduced PFN2 levels during ESC differentiation by deletion of the IRE in its 3'UTR results in reduced nuclear beta-catenin levels without significantly impacting cytoplasmic levels. Nuclear beta-catenin is critical for the differentiation of embryonic stem cells down the meso-endoderm lineages (25). Therefore PFN2's role in regulating nuclear beta-catenin levels can explain the requirement for elevated PFN2 levels during ESC differentiation into mesendoderm. These findings raise the important question of how PFN2 regulates beta-catenin nuclear import and/or retention. A small number of beta-catenin interacting proteins have been shown to play a role in its nuclear import, but otherwise very little is known (48–50). Our attempts at co-immunoprecipitation of PFN2 and beta-catenin failed to uncover a direct interaction (data not shown). Therefore, it remains unclear if the role of PFN2 in beta-catenin cellular localization is direct or indirect. However both PFN2 and beta-catenin interact with actin (51–53) and, therefore, it could be that a direct role for PFN2 in actin control is indirectly impacting beta-catenin nuclear import and/or retention. Indeed, nuclear transport of beta catenin and actin have been proposed to be tightly intertwined (54). Beta-catenin is a member of the Armadillo family, whose members share the ability to move into the nucleus independent of the classic nuclear localization signal (NLS) which uses an Importin-dependent mechanism for transport (55). Therefore, it is possible that PFN2 will be found to play important roles in the transport of other Armadillo families as well. In either case, future studies on PFN2's role in beta-catenin nuclear transport/retention are likely identify further novel insights into how the Wnt pathway is controlled in time and space during development.

In sum, our findings reveal that post-transcriptional regulation of the cytoskeletal protein Profilin-2 within a conserved short region of its 3'UTR that includes both an Ago2/ESCC miRNA site and an IRP/IRE site is essential for its tight molecular control during early ESC differentiation. This tight control of Profilin-2 expression tunes the FGF/ERK signaling pathway, which is integral to the naïve to formative pluripotency transition, as well as promotes the WNT/Beta-catenin signaling pathway, which is integral to lineage commitment to mesendoderm. This model provides a paradigm for how the coordination of miRNA and RBP binding control on a single transcript over development time can regulate cell fate decisions.

Acknowledgments

Flow Cytometry was performed at the UCSF Parnassus Flow Cytometry Core, supported in part by the DRC Center Grant NIH P30 DK063720. Sequencing was performed at the UCSF Center for Advanced Technology, supported by UCSF PBBR, RRP IMIA, and NIH 1S10OD028511-01 grants. This research was supported by funding from NIH/NICHD (K08HD105017) and Burroughs Wellcome Fund (to C.S.), and R01 GM125089 and GM122439 (to R.B.).

Author Contributions:

C.S. and R.B. developed the concept and designed experiments. C.S. conducted all experiments and all initial data analysis. C.S. and R.B. interpreted the data, wrote and edited the manuscript.

Declaration of Interests:

The authors declare no competing interests.

Figure legends

Figure 1
IRP/IRE site regulates PFN2 levels during ESC differentiation

(a) Schematic map of PFN2-3'UTR showing locations and evolutionary conservation of the neighboring IRP/IRE and Ago/miR-290 binding sites. **(b)** Schematic of (ESC) differentiation as embryoid bodies (EB) with descriptors of developmental stages at days 2, 5, and 7. **(c)** Expression analysis of Profilin-2 by qPCR in wild-type, PFN2-3UTR ΔIRE, and PFN2-3UTR ΔESCC mutants at naïve pluripotency (ESC): wild-type N=29; PFN2-3UTR ΔIRE N=38; PFN2-3UTR ΔESCC N=23; formative pluripotency (EBD2): wild-type N=8; PFN2-3UTR ΔIRE N=19; PFN2-3UTR ΔESCC N=7;, and early trilineage differentiation (EBD6): wild-type N=7; PFN2-3UTR ΔIRE N=19; PFN2-3UTR ΔESCC N=7 relative to wild-type ESC. Biological replicates for each cell line are shown. Error bars represent SEM. ****$P<0.0001$, unpaired two-tailed t test. **(d)** Absolute Profilin-2 protein levels (ng/mL) by ELISA in wild-type, PFN2-3UTR ΔIRE, and PFN2-3UTR ΔESCC mutants at naïve pluripotency (ESC): wild-type N=3; PFN2-3UTR ΔIRE N=3; PFN2-3UTR ΔESCC N=3;, formative pluripotency (EBD2): wild-type N=3; PFN2-3UTR ΔIRE N=3; PFN2-3UTR ΔESCC N=3;, and early trilineage differentiation (EBD6): wild-type N=5; PFN2-3UTR ΔIRE N=5; PFN2-3UTR ΔESCC N=3; relative to wild-type ESC. Biological replicates for each cell line are shown. Error bars represent SEM. *$P< 0.05$, **$P< 0.01$, ****$P<0.0001$, unpaired two-tailed t test.

Figure 2
Deletion of IRE in PFN2-3UTR impairs early mesendodermal lineage specification

(a) Volcano plot of transcriptomic data PFN2-3UTR ΔIRE versus wild-type at EBD6 showing downregulated (left side) and upregulated (right side) differentially-expressed genes (FDR<0.05). Integration of n=3 biological replicates for each genotype. **(b)** Gene ontology enrichment analysis of PFN2-3UTR ΔIRE versus wild-type at EBD6. **(c)** Heatmap display showing mean-scaled normalized gene expression, measured by RNA-seq, of representative pluripotency and trilineage differentiation markers in wild-type and PFN2-3UTR ΔIRE mutants at EBD6; (bottom right) color coding scheme for known developmental and pluripotency markers **(d)** Dot plot of normalized counts of select representative transcripts of mesoderm, endoderm and **(e)** ectoderm, pluripotency-related markers from bulk transcriptomic studies of wild-type and PFN2-3UTR ΔIRE at EBD6. N=3 biological replicates for each genotype. Error bars represent SEM. *$P< 0.05$, **$P< 0.01$, ***$P<0.001$, ****$P<0.0001$, unpaired two-tailed t test. **(f)** Representative immunoblot (left) and immunoblot summary of protein levels (right) of mesoderm (ACTA2) and endoderm (FOXA2) markers in wild-type versus PFN2-3UTR ΔIRE, PFN2-3UTR ΔESCC, and PFN2-KO mutants at EBD6. Scanned images of unprocessed blots are shown in Extended Data Fig. 2b.

Figure 3
Deletion of IRE inPFN2-3UTR leads to morphologic and molecular defects in embryoid bodies during differentiation

(a) Schematic of experimental setup of stem cells (ESC) to embryoid body (EB) formation and differentiation (EBD2-6) with associated developmental trajectory. Representative images of **(b)** wild-type and **(c)** PFN2-3UTR ΔIRE mutant EBD6 embryoid bodies in culture, with **(d)** focus inset highlighting morphologic defect **(e)** Schematic of EB experimental culture and analysis process **(f)** Flow cytometry analysis of dissociated cells from wild-type versus PFN2-3UTR ΔIRE mutant embryoid

bodies stained for active (non-phosphorylated) beta-catenin at **(f)** EBD2 and **(g)** EBD6, with **(h)** summary plot of n=3 independent experiments (single cells of representative experiment shown), ***P<0.001, ****P<0.0001, unpaired two-tailed t test. **(i)** Schematic depicting chimera formation and EB time course, intracellular protein analysis. Representative flow cytometry analysis of dissociated **(j)** wild-type versus PFN2-3UTR ΔIRE mutant chimeric embryoid bodies and wild-type versus PFN2-3UTR ΔESCC mutant chimeric embryoid bodies stained for intracellular active beta-catenin at EBD6, with associated **(k)** summary plot of n=3 independent experiments (single cells of representative experiment shown), ****P<0.0001, unpaired two-tailed t test. Representative flow cytometry analysis of dissociated **(l)** wild-type versus PFN2-3UTR ΔIRE mutant chimeric embryoid bodies and wild-type versus PFN2-3UTR ΔESCC mutant embryoid bodies stained for intracellular EOMES protein at EBD6, with associated **(m)** summary plot of n=3 independent experiments (single cells of representative experiment shown), ****P<0.0001, unpaired two-tailed t test.

Figure 4
Deletion of IRE in PFN2-3UTR IRE impairs beta catenin levels and Brachyury expression during primitive streak-like (Wnt/Nodal) induction

(a) Schematic experimental setup for directed mesoderm differentiation. **(b)** Relative expression of Brachyury (T) in PFN2-3UTR ΔIRE mutants versus wild-type by quantitative RT-PCR at ESC, EBD2, and EBD4M. N= 3 biological replicates for each genotype. ESC vs EBD4M, **** = P<0.0001, unpaired two-tailed t test. **(c)** Representative flow cytometry analysis of one experiment of dissociated wild-type versus PFN2-3UTR ΔIRE mutant embryoid bodies stained for active (non-phosphorylated) beta-catenin at EBD4M, with associated **(d)** summary plot of n=3 independent experiments, ***P<0.001, unpaired two-tailed t test. Representative flow cytometry analysis of dissociated **(e)** summary plot of wild-type versus PFN2-3UTR ΔIRE mutant chimeric embryoid bodies (left) and wild-type versus PFN2-3UTR ΔESCC mutant chimeric embryoid bodies (right) stained for active (non-phosphorylated) beta-catenin at EBD4M, **(f)** wild-type versus PFN2-3UTR ΔIRE mutant chimeric embryoid bodies (left) and wild-type versus PFN2-3UTR ΔESCC mutant chimeric embryoid bodies (right) stained for total beta-catenin at EBD4M, (single cells of representative experiment shown), with **(g)** summary of multiple experiments,   ***P<0.001, unpaired two-tailed t test.

Figure 5
Deletion of IRE in PFN2-3UTR is associated with decreased Profilin2 levels, decreased nuclear retention of active beta-catenin, and failure to induce T Brachyury

(a) Schematic of beta-catenin signaling pathway showing including cytoplasmic-nuclear transit of beta-catenin following activation leading to TCF/LEF target gene transcription inducing mesendodermal differentiation. **(b)** Representative immunoblot analysis of active beta-catenin (ABC) and Brachyury (T) protein levels in cytoplasmic, nuclear, and membrane subcellular compartments in wild-type and **(c)** PFN2-3UTR ΔIRE mutants at EBD4M. Scanned images of unprocessed blots are shown in Extended Data Fig. 5b. **(d)** Summary of nuclear Brachyury (T) protein levels from multiple immunoblots. Error bars represent SEM. **P< 0.01, unpaired two-tailed t test. **(e)** Schematic experimental setup for time course collection following addition of CHIR+Activin. **(f)** Immunoblot analysis of active beta-catenin (ABC), total beta-catenin (TBC), and Brachyury (T) protein levels in cytoplasmic and nuclear subcellular compartments in wild-type and **(g)** PFN2-3UTR ΔIRE mutants over 18 hour differentiation timecourse. Scanned images of unprocessed blots are shown in Extended Data Fig. 5. Protein levels of **(h)** nuclear active beta-catenin, **(i)** cytoplasmic active beta-catenin, **(j)** cytoplasmic total beta-catenin, and **(k)** nuclear Brachyury (T), and in wildtype (white) and PFN2-3UTR

ΔIRE mutants (red) over 18 hour timecourse. **(l)** Subcellular cytoplasmic and nuclear PFN2 protein by ELISA in wild-type (white) and PFN2-3UTR ΔIRE mutants (red) at 6H and 18H timepoints. **(m)** Nuclear PFN2 protein by ELISA in EBD4 lysate from wild-type (white) and PFN2-3UTR ΔIRE mutants (red). (h-m) N=3. Error bars represent SEM. *P< 0.05, ****P<0.0001, unpaired two-tailed t test.

Figure 6
Impact of post-transcriptional control of the Profilin-2 transcript on stem cell biology during pluripotency, pluripotency transition, and early trilineage differentiation

Schematic summary of the impact of Profilin-2 transcript Ago2/miR-290 and IRP/IRE binding sites on stem cell biology during pluripotency, pluripotency transition, and early trilineage differentiation.

Methods

Cell Culture and ESC Monolayer Differentiation
Mouse ESCs were grown and maintained at 37C and 5%CO2 on 0.2% gelatin-coated plates, cultured in ESC medium (custom DMEM) supplemented with 15% FBS (Corning Mediatech), LIF (1000U/mL, custom) and 2i (1uM PD0325901 (Axon MedChem) and 3uM CHIR99021 (Axon MedChem). Differential states of pluripotency from naïve to EpiLC transition were generated by removal of LIF and 2i. In brief, ESCs were plated in ESC medium (as described above). To initiate differentiation, LIF and 2i were removed. Cells were collected at indicated time points following LIF/2i removal.

Embryoid Body Generation and Differentiation
For embryoid body (EB) generation, ESCs were cultured in rotary (100 rpm) suspension culture on a low attachment plate allowing for self-aggregation and regular reproducible spontaneous differentiation over a course of 7 to 10 days. EB differentiation was evaluated by quantitative reverse transcriptase PCR, immunoblot analysis, and bulk RNA-sequencing. Directed differentiation to mesendoderm was performed by induction with 10 ng mL^{-1} Activin (GF300, Millipore) and 3 uM CHIR99021 (Axon MedChem). Chimeric embryoid bodies were formed by combining equal numbers of cells prior to rotary suspension culture.

Quantitative RT -PCR
RNA was extracted from cells using TRIzol (Invitrogen) and quantified on a Nanodrop spectrophotometer (ThermoFisher Scientific). Reverse transcription and quantitative PCR were performed according to manufacturer's instructions (Maxima, K1641, ThermoScientific; PowerSYBR, Applied Biosystems). Primer sequences are listed in [Supplementary Table 1].

ELISA immunoassay
Sandwich ELISA immunoassay for in vitro quantitative measurement of PFN2. A microplate pre-coated with an antibody specific for Mouse PFN2 was used for sandwich ELISA immunoassay in vitro quantitative measurement of PFN2 (RK06647, ABclonal Science) according to manufacturer's instructions.

Generation of Site-based Mutant Cell Lines by CRISPR-CAS9 Gene Editing
Mutant ESC cell lines were generated by cloning gRNAs into a U6 vector (pGuide, Origene) and co-transfection with a Cas9-expressing vector (pCas-Guide-EF1a-GFP, Origene) using Fugene6 (Roche) as described in (4). Guide sequences are listed in [Supplementary Table 2].

RNA-seq library preparation and sequencing
RNA (total or poly-adenylated) was extracted using TRIzol (Invitrogen) or Oligo(dT)25-coated Dynabeads (Invitrogen), respectively. RNA quality was examined by Qubit 2.0 Fluorometer (Life Technologies). cDNA library generation was done using the KAPA RNA Hyperprep kit (KK8541, Roche) following the manufacturer's instructions. cDNA library quality was assessed using Qubit 2.0 fluorometer to determine concentrations. The 4200 TapeStation system (Agilent) was used to determine insert size following the manufacturer's instructions. cDNA libraries were then sequenced using the HiSeq 4000 (Illumina) and NovaSeq X Systems (Illumina).

Bulk RNA-Seq data pre-processing and analysis
Raw fastq files were filtered and preprocessed by quality trimming using FASTX-Toolkit (http://hannonlab.cshl.edu/fastx_toolkit). Adapters were removed and reads collapsed using unique molecular identifiers using FASTX-Toolkit and Cutadapt (56). RNA-seq read mapping and alignment with alignment to mm10 reference genome assembly was performed using STAR (57). Read count tables were generated using the RSEM program (58). Differential gene expression was performed using R packages edgeR (59) and DESeq2 (60). FDR was calculated using the Benjamini–Hochberg method. DE genes were defined with FDR less than 0.05 in comparison to the control group.

Gene Ontology and GO Enrichment Analysis
Gene Ontology reference and GO Enrichment Analysis were accessed and used to perform ontology and fold enrichment analysis using the GO Ontology database DOI: 10.5281/zenodo.10536401 (61,62)

Immunoblot
Total cell extracts were obtained by lysing cell pellets with ice-cold RIPA lysis buffer containing EDTA-free protease inhibitor mixture (Roche) and PhosSTOP phosphatase inhibitor (Roche), with DNA shearing by 21-gauge needle. Lysates were incubated at 4C for 15 min and centrifuged at 20,000 x g at 4 C on a table-top centrifuge. The protein concentration of the supernatant was measured by BCA assay, with 30 micrograms of protein resolved by sodium dodecyl sulfate-polyacrylamide gel electrophoresis. After electrophoresis, proteins were transferred onto an Immobilon-FL PVDF membrane (Millipore) and processed for immunodetection. Primary antibodies were incubated overnight at 4C. Secondary antibodies were incubated for 1h at RT. Images were digitally acquired using a Li-COR Odyssey imager (Li-COR Instruments) and its proprietary software. Antibodies are listed in Supplementary Table 3. Bands were quantified using Image J (63).

Subcellular Fractionations
Subcellular fractionations were performed according to manufacturer instructions (78840, Thermo Scientific), and followed by ELISA and/or immunoblot.

Intracellular Staining Assay
Cells were first fixed with 4% paraformaldehyde for 15 min at room temperature, then permeabilized on ice with 0.1% Triton-X for 5 min and stained for the indicated primary antibody for 30 min at 4C at dilutions as listed in Supplementary Table 3. This was followed by incubation for 30 minutes at 4C with species-specific secondary antibodies at dilutions as listed in Supplementary Table 3. Finally the cells were washed with cold PBS and prepared for analysis by flow cytometry. Antibodies are listed in Supplementary Table 3.

Flow Cytometry

After dissociation into single-cell suspension using Trypsin or TrypLE (ThermoFisher Scientific), cells were resuspended in HBSS/1% BSA and filtered for size by cell strainer (35 micron mesh). Cells were processed by flow cytometry on an LSR II flow cytometer (Becton Dickinson), with events collected by FACS Diva software (BD Biosciences) using a hierarchical gating strategy selecting for a homogenous cell population (forward-scatter area vs. side-scatter area) of single cells (forward-scatter width vs. forward-scatter area). Dot plots and mean fluorescence intensities for gated events were generated using FlowJo analysis software.

Statistical Analysis

No data were excluded from the analyses. Statistical tests were performed in GraphPad Prism (v10). Data collection and analysis were not performed blind to the conditions of the experiments, but data analyses have been performed with identical parameters and software. Data representation and statistical analyses were also performed using R software. The number of biological replicates and independent experiments, both equal to or greater than 3, is indicated in figure legends. The statistical tests used are indicated in the figure legends.

Supplemental information

Document S1. Figures S1-S5. Tables S1-S3.

Data availability

Sequencing data that support the findings of this study have been deposited in the Gene Expression Omnibus (GEO) and are publicly available as of the date of publication. Primers, oligonucleotide sequences and antibodies are presented in Supplementary Tables 1-3. Source data are provided with this paper. All other data supporting the findings of this study are available from the corresponding author upon reasonable request.

Code availability

This paper does not report original code.

References

1.   Ye J, Blelloch R. Regulation of pluripotency by RNA binding proteins. Cell Stem Cell. 2014 Sep 4;15(3):271–80.

2.   Judson RL, Greve TS, Parchem RJ, Blelloch R. MicroRNA-based discovery of barriers to dedifferentiation of fibroblasts to pluripotent stem cells. Nat Struct Mol Biol. 2013 Oct;20(10):1227–35.

3.   Luscieti S, Galy B, Gutierrez L, Reinke M, Couso J, Shvartsman M, et al. The actin-binding protein profilin 2 is a novel regulator of iron homeostasis. Blood. 2017 Oct 26;130(17):1934–45.

4.   Sangokoya C, Blelloch R. MicroRNA-dependent inhibition of PFN2 orchestrates ERK activation and pluripotent state transitions by regulating endocytosis. Proc Natl Acad Sci USA. 2020 Aug 25;117(34):20625–35.

5.   Ying Q-L, Wray J, Nichols J, Batlle-Morera L, Doble B, Woodgett J, et al. The ground state of embryonic stem cell self-renewal. Nature. 2008 May 22;453(7194):519–23.

6.   Aberle H, Bauer A, Stappert J, Kispert A, Kemler R. beta-catenin is a target for the ubiquitin-proteasome pathway. EMBO J. 1997 Jul 1;16(13):3797–804.

7.   Kitagawa M, Hatakeyama S, Shirane M, Matsumoto M, Ishida N, Hattori K, et al. An F-box protein, FWD1, mediates ubiquitin-dependent proteolysis of beta-catenin. EMBO J. 1999 May 4;18(9):2401–10.

8.   Hart M, Concordet JP, Lassot I, Albert I, del los Santos R, Durand H, et al. The F-box protein beta-TrCP associates with phosphorylated beta-catenin and regulates its activity in the cell. Curr Biol. 1999 Feb 25;9(4):207–10.

9.   Boroviak T, Loos R, Bertone P, Smith A, Nichols J. The ability of inner-cell-mass cells to self-renew as embryonic stem cells is acquired following epiblast specification. Nat Cell Biol. 2014 Jun;16(6):516–28.

10.   Smith A. Formative pluripotency: the executive phase in a developmental continuum. Development. 2017 Feb 1;144(3):365–73.

11.   Kinoshita M, Barber M, Mansfield W, Cui Y, Spindlow D, Stirparo GG, et al. Capture of Mouse and Human Stem Cells with Features of Formative Pluripotency. Cell Stem Cell. 2021 Mar 4;28(3):453-471.e8.

12.   Carpenedo RL, Sargent CY, McDevitt TC. Rotary suspension culture enhances the efficiency, yield, and homogeneity of embryoid body differentiation. Stem Cells. 2007 Sep;25(9):2224–34.

13.   Dang SM, Kyba M, Perlingeiro R, Daley GQ, Zandstra PW. Efficiency of embryoid body formation and hematopoietic development from embryonic stem cells in different culture systems. Biotechnol Bioeng. 2002 May 20;78(4):442–53.

14.   Arias AM, Marikawa Y, Moris N. Gastruloids: Pluripotent stem cell models of mammalian gastrulation and embryo engineering. Dev Biol. 2022 Aug;488:35–46.

15.   Wang Y, Baskerville S, Shenoy A, Babiarz JE, Baehner L, Blelloch R. Embryonic stem cell-specific microRNAs regulate the G1-S transition and promote rapid proliferation. Nat Genet. 2008 Dec 1;40(12):1478–83.

16.   Judson RL, Babiarz JE, Venere M, Blelloch R. Embryonic stem cell-specific microRNAs promote induced pluripotency. Nat Biotechnol. 2009 May;27(5):459–61.

17.   Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. Cell. 2004 Jan 23;116(2):281–97.

18.   Hutvágner G, Zamore PD. A microRNA in a multiple-turnover RNAi enzyme complex. Science. 2002 Sep 20;297(5589):2056–60.

19.   Liu J, Carmell MA, Rivas FV, Marsden CG, Thomson JM, Song J-J, et al. Argonaute2 is the catalytic engine of mammalian RNAi. Science. 2004 Sep 3;305(5689):1437–41.

20.   Meister G, Landthaler M, Patkaniowska A, Dorsett Y, Teng G, Tuschl T. Human Argonaute2 mediates RNA cleavage targeted by miRNAs and siRNAs. Mol Cell. 2004 Jul 23;15(2):185–97.

21.   Pillai RS, Artus CG, Filipowicz W. Tethering of human Ago proteins to mRNA mimics the miRNA-mediated repression of protein synthesis. RNA. 2004 Oct;10(10):1518–25.

22.   Galy B, Conrad M, Muckenthaler M. Mechanisms controlling cellular and systemic iron homeostasis. Nat Rev Mol Cell Biol. 2024 Feb;25(2):133–55.

23.   Lyashenko N, Winter M, Migliorini D, Biechele T, Moon RT, Hartmann C. Differential requirement for the dual functions of β-catenin in embryonic stem cell self-renewal and germ layer formation. Nat Cell Biol. 2011 Jun 19;13(7):753–61.

24.   Valenta T, Hausmann G, Basler K. The many faces and functions of β-catenin. EMBO J. 2012 Jun 13;31(12):2714–36.

25.   Gadue P, Huber TL, Paddison PJ, Keller GM. Wnt and TGF-beta signaling are required for the induction of an in vitro model of primitive streak formation using embryonic stem cells. Proc Natl Acad Sci USA. 2006 Nov 7;103(45):16806–11.

26.   Liu C, Li Y, Semenov M, Han C, Baeg GH, Tan Y, et al. Control of beta-catenin phosphorylation/degradation by a dual-kinase mechanism. Cell. 2002 Mar 22;108(6):837–47.

27.   Orford K, Crockett C, Jensen JP, Weissman AM, Byers SW. Serine phosphorylation-regulated ubiquitination and degradation of beta-catenin. J Biol Chem. 1997 Oct 3;272(40):24735–8.

28.   van Noort M, Meeldijk J, van der Zee R, Destree O, Clevers H. Wnt signaling controls the phosphorylation status of beta-catenin. J Biol Chem. 2002 May 17;277(20):17901–5.

29.   van Noort M, Weerkamp F, Clevers HC, Staal FJT. Wnt signaling and phosphorylation status of -catenin: importance of the correct antibody tools. Blood. 2007 Oct 1;110(7):2778–9.
30.   Arnold SJ, Hofmann UK, Bikoff EK, Robertson EJ. Pivotal roles for eomesodermin during axis formation, epithelium-to-mesenchyme transition and endoderm specification in the mouse. Development. 2008 Feb;135(3):501–11.

31.   Russ AP, Wattler S, Colledge WH, Aparicio SA, Carlton MB, Pearce JJ, et al. Eomesodermin is required for mouse trophoblast development and mesoderm formation. Nature. 2000 Mar 2;404(6773):95–9.

32.   Turner DA, Rué P, Mackenzie JP, Davies E, Martinez Arias A. Brachyury cooperates with Wnt/β-catenin signalling to elicit primitive-streak-like behaviour in differentiating mouse embryonic stem cells. BMC Biol. 2014 Aug 13;12(1):63.

33.	Tosic J, Kim G-J, Pavlovic M, Schröder CM, Mersiowsky S-L, Barg M, et al. Eomes and Brachyury control pluripotency exit and germ-layer segregation by changing the chromatin state. Nat Cell Biol. 2019 Dec 2;21(12):1518–31.

34.	Yamaguchi TP, Takada S, Yoshikawa Y, Wu N, McMahon AP. T (Brachyury) is a direct target of Wnt3a during paraxial mesoderm specification. Genes Dev. 1999 Dec 15;13(24):3185–90.

35.	Ree R, Kind L, Kaziales A, Varland S, Dai M, Richter K, et al. PFN2 and NAA80 cooperate to efficiently acetylate the N-terminus of actin. J Biol Chem. 2020 Dec 4;295(49):16713–31.

36.	Burdon T, Stracey C, Chambers I, Nichols J, Smith A. Suppression of SHP-2 and ERK signalling promotes self-renewal of mouse embryonic stem cells. Dev Biol. 1999 Jun 1;210(1):30–43.

37.	Stavridis MP, Lunn JS, Collins BJ, Storey KG. A discrete period of FGF-induced Erk1/2 signalling is required for vertebrate neural specification. Development. 2007 Aug;134(16):2889–94.

38.	Yang S-H, Kalkan T, Morrisroe C, Smith A, Sharrocks AD. A genome-wide RNAi screen reveals MAP kinase phosphatases as key ERK pathway regulators during embryonic stem cell differentiation. PLoS Genet. 2012 Dec 13;8(12):e1003112.

39.	Kunath T, Saba-El-Leil MK, Almousailleakh M, Wray J, Meloche S, Smith A. FGF stimulation of the Erk1/2 signalling cascade triggers transition of pluripotent embryonic stem cells from self-renewal to lineage commitment. Development. 2007 Aug;134(16):2895–902.

40.	Kim S, Kim S, Chang HR, Kim D, Park J, Son N, et al. The regulatory impact of RNA-binding proteins on microRNA targeting. Nat Commun. 2021 Aug 20;12(1):5057.

41.	Kedde M, van Kouwenhove M, Zwart W, Oude Vrielink JAF, Elkon R, Agami R. A Pumilio-induced RNA structure switch in p27-3' UTR controls miR-221 and miR-222 accessibility. Nat Cell Biol. 2010 Oct;12(10):1014–20.

42.	Sternburg EL, Estep JA, Nguyen DK, Li Y, Karginov FV. Antagonistic and cooperative AGO2-PUM interactions in regulating mRNAs. Sci Rep. 2018 Oct 17;8(1):15316.

43.	Kedde M, Strasser MJ, Boldajipour B, Oude Vrielink JAF, Slanchev K, le Sage C, et al. RNA-binding protein Dnd1 inhibits microRNA access to target mRNA. Cell. 2007 Dec 28;131(7):1273–86.
44.	Eiring AM, Harb JG, Neviani P, Garton C, Oaks JJ, Spizzo R, et al. miR-328 functions as an RNA decoy to modulate hnRNP E2 regulation of mRNA translation in leukemic blasts. Cell. 2010 Mar 5;140(5):652–65.

45.	Bhattacharyya SN, Habermacher R, Martine U, Closs EI, Filipowicz W. Relief of microRNA-mediated translational repression in human cells subjected to stress. Cell. 2006 Jun 16;125(6):1111–24.

46.	Chang S-H, Lu Y-C, Li X, Hsieh W-Y, Xiong Y, Ghosh M, et al. Antagonistic function of the RNA-binding protein HuR and miR-200b in post-transcriptional regulation of vascular endothelial growth factor-A expression and angiogenesis. J Biol Chem. 2013 Feb 15;288(7):4908–21.

47.   Jafarifar F, Yao P, Eswarappa SM, Fox PL. Repression of VEGFA by CA-rich element-binding microRNAs is modulated by hnRNP L. EMBO J. 2011 Apr 6;30(7):1324–34.

48.   Goto T, Sato A, Adachi S, Iemura S, Natsume T, Shibuya H. IQGAP1 protein regulates nuclear localization of β-catenin via importin-β5 protein in Wnt signaling. J Biol Chem. 2013 Dec 20;288(51):36351–60.

49.   Komiya Y, Mandrekar N, Sato A, Dawid IB, Habas R. Custos controls β-catenin to regulate head development during vertebrate embryogenesis. Proc Natl Acad Sci USA. 2014 Sep 9;111(36):13099–104.

50.   Griffin JN, Del Viso F, Duncan AR, Robson A, Hwang W, Kulkarni S, et al. RAPGEF5 Regulates Nuclear Translocation of β-Catenin. Dev Cell. 2018 Jan 22;44(2):248-260.e4.

51.   Witke W, Podtelejnikov AV, Di Nardo A, Sutherland JD, Gurniak CB, Dotti C, et al. In mouse brain profilin I and profilin II associate with regulators of the endocytic pathway and actin assembly. EMBO J. 1998 Feb 16;17(4):967–76.

52.   Pilo Boyl P, Di Nardo A, Mulle C, Sassoè-Pognetto M, Panzanelli P, Mele A, et al. Profilin2 contributes to synaptic vesicle exocytosis, neuronal excitability, and novelty-seeking behavior. EMBO J. 2007 Jun 20;26(12):2991–3002.

53.   Yamada S, Pokutta S, Drees F, Weis WI, Nelson WJ. Deconstructing the cadherin-catenin-actin complex. Cell. 2005 Dec 2;123(5):889–901.

54.   Sen B, Xie Z, Howard S, Styner M, van Wijnen AJ, Uzer G, et al. Mechanically Induced Nuclear Shuttling of β-Catenin Requires Co-transfer of Actin. Stem Cells. 2022 Apr 29;40(4):423–34.

55.   Fagotto F. Looking beyond the Wnt pathway for the deep nature of β-catenin. EMBO Rep. 2013 May;14(5):422–33.

56.   Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal, 17*(1), pp. 10-12. doi:https://doi.org/10.14806/ej.17.1.200

57.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013 Jan 1;29(1):15–21.

58.   Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinformatics. 2011 Aug 4;12:323.

59.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics. 2010 Jan 1;26(1):139–40.

60.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

61.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene Ontology: tool for the unification of biology. Nat Genet. 2000 May;25(1):25–9.

62.   Thomas PD, Ebert D, Muruganujan A, Mushayahama T, Albou L-P, Mi H. PANTHER: Making genome-scale phylogenetics accessible to all. Protein Sci. 2022 Jan;31(1):8–22.

63.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. Nat Methods. 2012 Jul;9(7):671–5.

Fig 1

Fig 2

Fig 3

a

ESC → EBD2 → EBD5 → EBD6

-Lif/2i EB formation

[formative pluripotency] [early trilineage differentiation] [mid-trilineage differentiation]

b **c** **d**

EBD6

WT

PFN2 3UTR MT-IRE

[INSET] PFN2 3UTR MT-IRE

e

EBD2 → EBD5 → EBD6

dissociate intracellular protein analysis (flow)

f

EBD2 intracellular Activated Beta-Catenin

WT   PFN2-ΔIRE

g

EBD6 intracellular Activated Beta-Catenin

WT   PFN2-ΔIRE

h

intracellular Activated Beta-Catenin

EBD2   EBD6

[WT] [ΔIRE] [WT] [ΔIRE]

Fluorescence Intensity relative to WT

*** **** ****

WT PFN2ΔIRE WT PFN2ΔIRE

i

ESC
WT MT

EB CHIMERA

EBD2 → EBD5 → EBD6

dissociate intracellular protein analysis (flow)

j

EBD6 CHIMERA intracellular Activated Beta-Catenin

WT PFN2-ΔIRE WT PFN2-ΔESCC

k

EBD6 CHIMERA intracellular Activated Beta-Catenin

WT+PFN2ΔIRE WT+PFN2ΔESCC
[WT] [ΔIRE] [WT] [ΔESCC]

Fluorescence Intensity relative to WT

****

[WT]+ΔIRE WT+[ΔIRE] [WT]+ΔESCC WT+[ΔESCC]

l

EBD6 CHIMERA intracellular EOMES

WT PFN2-ΔIRE WT PFN2-ΔESCC

m

EBD6 CHIMERA intracellular EOMES

WT+PFN2ΔIRE WT+PFN2ΔESCC
[WT] [ΔIRE] [WT] [ΔESCC]

Fluorescence Intensity relative to WT

****

[WT]+ΔIRE WT+[ΔIRE] [WT]+ΔESCC WT+[ΔESCC]

Fig 4

Fig 5

Fig 6