The assembly of microbial communities on red sandstone surfaces was shaped by dispersal limitation and heterogeneous selection

Bowen Wang [a, b], Chengshuai Zhu [a], Xin Wang [c], Tianyu Yang [d], Bingjian Zhang [a, c] and Yulan Hu [a*]

[a] Department of Archaeology, Cultural Heritage and Museology, Zhejiang University, Hangzhou, 310028, China.
[b] Department of Biology, Johns Hopkins University, Baltimore, MD, 21218, USA.
[c] Department of Chemistry, Zhejiang University, Hangzhou, 310028, China.
[d] Leshan Giant Buddha Grottoes Research Institute, Leshan Giant Buddha Scenic Area Management Committee, Leshan, 614000, China.

[*]Corresponding: hu_yulan@zju.edu.cn

Abstract

Understanding the role of microbiota on stone surface is essential for developing effective grottoes conservation strategies. However, ecological feature of microbial communities on stone surface have been rarely investigated systematically. In this study, we explored diversity, assembly, and functional profiles of microbial communities on the red sandstone surface of the Leshan Giant Buddha from microbial ecology perspective. The results show that Proteobacteria, Actinobacteria, Cyanobacteria, and Ascomycota are the dominant phyla. Fundamental metabolic pathways are maintained during the formation of visually distinguishable microbial communities, but gene profiles vary across microbial communities of different colors. Ecological modelling suggests that selective pressure from the harsh stone surface environment fostered the interplay of dispersal limitation and heterogeneous selection during community assembly. The assembly of visually distinct microbial communities is linked to a narrower ecological niche, higher proportion of habitat specialists, elevated complexity, and increased resilience of the prokaryotic network to disturbances. Microbial-mediated ammonium assimilation and nitrogen mineralization might be the two prominent processes that contribute to stone biodeterioration. This study deepens our understanding of the assembly mechanisms and functional potentials of microbial communities on stone cultural heritage surfaces, provides microbial ecological insights for the conservation of these cultural treasures.

Importance: Minimal systematically research on the ecological interpretation of stone biodeterioration. This study reports dispersal limitation and heterogeneous selection shape the microbial community assembly responsible for the biodeterioration of red sandstone. Furthermore, fundamental metabolic processes of microbial communities, such as ammonium assimilation and nitrogen mineralization, are identified as contributors to stone biodeterioration. This study improves our understanding of microbial community assembly and their functional roles, providing a microbial ecological basis for developing effective strategies for the conservation of stone cultural heritage.

Keywords: Community assembly; Microbial community; Microbial diversity; Microbial ecology; Stone; Cultural Heritage

1. Introduction

Grottoes are a common form of stone cultural heritage. The stone surface of grottoes are integral components of ecosystems, providing critical habitats for microbial communities, serving as crucial habitats for microorganisms organized into diverse microbial communities[1-3]. Under specific conditions, these microbial communities can reinforce the stone through metabolic processes like biomineralization[1,4-7]. However, in most cases, the microbial communities on stone surface not only diminish the aesthetic value of the artifacts by obscuring and deteriorating the carvings but may also, in some cases, compromise the structural integrity of the grottoes through severe erosion process[3,8-11]. Therefore, cleaning and controlling the formation of microbial communities on stone surface are essential tasks in the conservation of stone cultural heritage. Understanding the characteristics of microbial communities on stone surface is crucial for developing effective strategies to mitigate microbial deterioration. Currently, research in this field primarily focuses on the taxonomic diversity of microorganisms and their biogeochemical cycling and metabolic capabilities. These studies have revealed the key microbial groups involved in the biodeterioration of stone cultural heritage and their associated biogeochemical processes and metabolic responses[10,12-17]. However, ecological understanding of how microbial communities assemble on the stone surface remains significantly limited.

The ecological mechanisms that govern microbial community assembly are one of the main scientific questions in ecological research[18-20]. It is generally believed that the assembly of local microbial communities is controlled by the combined effects of deterministic and stochastic processes[21-23]. Deterministic processes are used to describe the traditional niche-based theory, which hypothesize that species traits, interspecific interactions, and environmental factors govern community structure[24,25]. In contrast, stochastic processes are based on the neutral theory, which illustrate that community assembly is driven by stochastic events such as birth, death, colonization, extinction, and speciation[26,27]. The niche on the stone surface exposes microorganisms to environmental stresses (abiotic and biotic constraints that limit species productivity and ecosystem development[28]) from nutrient and water scarcity, as well as extreme fluctuations in temperature, humidity, and solar UV radiation[1]. Generally, environmental stress increases deterministic assembly while reducing stochastic assembly, because species with higher tolerance or adaptability to these stressors tend to survive and thrive[28-30]. Furthermore, under high-stress conditions, microbial dispersal limitations are enhanced because elevated stress impedes colonization and successful establishment[28,31,32]. A recent study on microbial communities on limestone surface revealed that prokaryotes and fungi in this environment are primarily dominated by the mutual effect of dispersal limitation (stochastic process) and homogeneous selection (deterministic process)[3]. However, it is currently unknown whether such community assembly mechanisms are specific to certain stone type, geographic location, or

influenced by other factors.

The Leshan Giant Buddha, a world-renowned UNESCO World Heritage site, is a typical
example of red sandstone grotto cultural heritage. Previous studies found abundant
nitrogen-metabolizing microorganisms and ammonia monooxygenase genes, linked to
biodeterioration, on the surface of the Leshan Giant Buddha, which suggests that the
deterioration of the stone is closely related to microbial growth[33-35]. The cleaning of stone
surface is a crucial task for conservation of the grotto. The current issue is that despite regular
cleaning of the stone surface by heritage conservators, visually distinguishable microbial
communities persistently reappear after each cleaning. However, our understanding of how
microorganisms assemble microbial communities on the stone surface of the Leshan Giant
Buddha, as well as on other red sandstone surfaces, remains limited. There is an essential need
to investigate the mechanisms by which microbes survive and how they form microbial
communities within such ecosystems under various environmental stresses from the perspective
of microbial ecology. Therefore, this study uses the Leshan Giant Buddha, a UNESCO World
Heritage site, as a case study to provide a comprehensively microbial ecological evaluation to the
microbial communities on the surface of this red sandstone cultural heritage. The goal of this
study is to examine whether the ecological characteristics of microbial communities on red
sandstone surfaces are associated with adaptation to harsh environmental conditions.
Specifically, we investigate how these microorganisms assemble into stable communities under
repeated wet-dry cycles, nutrient limitations, and UV radiation, and what functional capacities
they exhibit. Our focus extends beyond the interactions within the microbiota to include the
interactions among microorganisms, environmental factors, and the stone substrate.
107

2. Results

2.1. Taxonomic composition varies by the color of microbial communities
Samples of three different types of visually distinguishable microbial communities, along
with blank samples, were collected from the stone surface of the Leshan Giant Buddha. In
general, the total 27 samples were grouped and named according to their respective colors (Bla,
black; Gre, green; Whi, white; and CK, blank control) (Fig. 1). We then performed amplicon
sequencing of conserved phylogenetic marker genes (16S and ITS rRNA) to identify bacteria and
fungi within the microbial communities. After quality filtering and rarefaction, a total of 8,791
and 2,404 non-zero ASVs were obtained from the 16S and ITS sequences, respectively. Each
group of data overlap with others, but they are not entirely identical or inclusive of each other
(Fig. S1). In addition, metagenome sequencing was applied for each group and obtained
206,491,166 clean reads in total (Table S1).
120
121

Figure 1. Representative examples and sampling of microbial communities on the red sandstone
surface of the Leshan Giant Buddha. (a). Location of the Leshan Giant Buddha. (b). Sampling
regions of the samples. Red boxes here indicate the specific sampling regions. (c). Representative
examples of CK samples. (d). Representative examples of Black samples. (e). Representative
examples of Green samples. (f). Representative examples of White samples. Notes: "n" in picture
c-f represents the number of samples used in this study within each group.

To get taxonomic insights into the microbiota, we annotated the sequencing data, then
visualized top 10 dominant phyla, top 20 dominant genus of bacteria (Fig. 2 a; Fig. S2 a; Table S2,
S3) and fungi (Fig. 2 b; Fig. S2 b; Table S4, S5) among all samples. We also constructed overall
taxonomic community composition trees of top 5% dominant bacteria, and top 30% dominant
fungi (Fig. 2 c, d). The results show that among all samples, Proteobacteria, Actinobacteria, and
Cyanobacteria are the dominant top three phyla in bacteria, comprising at least 70% of the total
bacteria in each sample. Ascomycota is the most dominant phylum in fungi, comprising at least
50% of the total fungi in each sample.
To estimate the separation pattern between each group of microbial communities,
non-metric multidimensional scaling (NMDS) based on Bray-Curtis dissimilarity at the ASV level
was performed and then conducted Anosim (analysis of similarities) and PERMANOVA (*i.e.*,
adonis) tests (Fig. 2 e, f). The results show that both bacteria and fungi among the four groups of
samples reveal significant and major differences ($P < 0.001$). In summary, the structure of
microbial communities varies according to the coloration of the samples, which indicates distinct
microbial compositions among samples of different colors.

Figure 2. Taxonomic composition and distinct taxonomic structures of the microbial communities.
(a). Relative abundances of dominant bacteria among all samples in phylum level. (b). Relative
abundances of dominant fungi among all samples in genus level. (c). The overall taxonomic
community composition tree of top 5% dominant bacteria. (d). The overall taxonomic community
composition tree of top 30% dominant fungi. (e). Non-metric multidimensional scaling (NMDS)
ordination based on Bray-Curtis dissimilarities at the ASV level for bacteria. (f). Non-metric
multidimensional scaling ordination based on Bray-Curtis dissimilarities at the ASV level for fungi.
Notes: The triple asterisks *** represents the P < 0.001.

2.2. Dispersal limitation and hetero selection dominants the assembly of red sandstone surface microbial communities

To characterize the assembly mechanism of the red sandstone surface microbial communities, we applied null model analysis[36,37] and dissimilarity-overlap curve (DOC) model analysis (Fig. 3)[38].

Null model analysis showed that both deterministic and stochastic processes controlled the assembly of microbial communities, but the relative importance of the two processes in community assembly vary among sample types (Fig. 3 a, b). In general, dispersal limitation dominated the assembly of both bacteria communities (60.11%) and fungi communities (55.56%) on stone surface. Bacteria revealed overall elevated deterministic process than eukaryotes. Hetero selection and dispersal limitation can explain about 91% of the assembly mechanism of the bacteria communities. While hetero selection and dispersal limitation were also main process for the assembly process of fungi communities; drift played a more pivotal role in the assembly process of fungi communities (20.23%) than in the bacteria ones (5.41%). Bacteria samples revealed significantly higher value of betaNTI than fungi ones (Fig. 3 c). Significant elevation of betaNTI was detected among White groups of bacteria samples comparing with CK (Fig. 3 d). By contrast, significant elevation of betaNTI was detected among Black as well as Green groups of fungi samples comparing with CK (Fig. 3 e).

In addition, the DOC model analysis demonstrated a significantly negative correlation between overlap and dissimilarity in both bacteria and fungi communities (P < 0.001), which indicated a general dynamic and interspecific interaction within these microbial communities (Fig. 3 f, g). These complex dynamic interactions among microorganisms in the microbiota might lead to competition within the shared niche for survival.

Figure 3. Assembly dynamics of the red sandstone surface microbial communities. (a). The relative importance of assembly processes in bacteria communities. (b). The relative importance of assembly processes in fungi communities. (c). The differences of betaNTI between bacteria and fungi communities. (d). Changes of betaNTI among three groups of bacteria samples comparing with CK. (e). Changes of betaNTI among three groups of fungi samples comparing with CK. (f). Dissimilarity-overlap curve (DOC) analysis of the universality of bacteria dynamics among samples. (g). DOC analysis of the universality of fungi dynamics among samples.

2.3. Specialization prevails over flexibility in microbial community formation on red sandstone surface

To delve deeper into the dynamic interactions among microorganisms within the microbiota, we further investigated the niche characterization of microbial communities on red sandstone surface. According to Levins niche breadth theory, the distribution of microbiota composition can be used as an indicator of the environmental conditions[39]. Here, we calculated niche breadth (*Bcom*) and niche overlap (*Ocom*) as indicators to describe the habitat niche characterization of both bacteria and fungi making up the stone surface microbial communities. First, the *Bcom* was calculated with Levins index (Fig. 4 a-c). The results showed that the visually distinguishable microbial communities revealed decreased overall *Bcom* comparing with CK group. Such results were also further validated with Shannon index (Fig. S3 a-c). Then, the *Ocom* was calculated with Schoener index. The *Ocom* for both bacteria and fungi displayed a similar trend that CK samples revealed the highest *Ocom*, followed by Black, Green, and White, in descending order. All

pairwise comparisons were statistically significant, indicating distinct niche overlaps among the
different samples. When comparing *Ocom* across all samples, bacteria revealed a significantly
higher *Ocom* than fungi (Fig. 4 d-f). Such results were also further validated with Pianka index
(Fig. S3 d-f). In summary, both bacteria and fungi communities exhibit significant variations in
Bcom and *Ocom* across different sample types (CK, Black, Green, White). In general, CK samples
displayed widest niche breadth and highest overlap level compared with the rest groups of
samples. These results suggested that the formation of visually distinguishable microbial
communities on stone surface requires species to be more specialized, rather than relying on
diverse and flexible species with metabolic flexibility and redundancy.

To further explore the inference, we also calculated the proportion of generalists and
specialists among the microbial communities (Fig. 4 g-k). Generalists are species that are better
suited to survive in diverse and challenging environments and often play pivotal roles within
their communities. Specialists, on the other hand, have a more limited range of habitats and are
specifically adapted to environmental conditions[40,41]. The results showed that at the ASVs level,
the proportion of specialists in each type of sample was higher than that of generalists in both
bacteria and fungi (Fig. 4 g). To determine whether there were differences between all bacteria
and fungi, we also compared the proportion of specialists and generalists within these groups.
The results showed that in each group of both bacteria and fungi, the proportion of specialists
was higher than that of generalists. But, bacteria revealed overall higher proportion of
generalists than that of specialists (Fig. 4 g). At the phylum level, up to 69.61 % of the bacteria
were classified as the middle; the proportion of specialists (29.95 %) was higher than that of
generalists (0.44 %) (Fig. 4 i, Table S6). Myxococcota revealed the largest fraction of specialists in
bacteria (52.4%) among the top 10 dominant phyla. (Fig. 4 h, Table S6). For the fungi, 59.5% were
classified as middle, 40.5% as specialists, and none of the ASV is classified as generalists. (Fig. 4 k,
Table S7). This suggested that specialists prevail over generalists in the dominant phyla within
the assembly of microbial communities. Notably, the non-dominant phyla (others) of both
bacteria and fungi exhibited the highest proportion of specialists (Fig. 4 h, j; Table S6, S7). This
suggested that the non-dominant phyla within the community might have even lower metabolic
flexibility compared to the dominant phyla.
234

Figure 4. Niche characterization of the red sandstone surface microbial communities based on Levins niche breadth index and Schoener niche overlap index. (a). Average habitat niche breadth (*Bcom*) differences among three groups of bacteria samples comparing with CK in bacteria. (b). *Bcom* differences among three groups of fungi samples comparing with CK in fungi. (c). *Bcom* differences between bacteria and fungi. (d). Average habitat niche breadth (*Ocom*) differences among three groups of bacteria samples comparing with CK in bacteria. (e). Average *Ocom* differences among three groups of fungi samples comparing with CK in fungi. (f). Average *Ocom* differences between bacteria and fungi. (g). Proportional distribution of specialists, middle, and generalists in each sample group, and between bacteria and fungi. (h). Proportional distribution

of specialists, middle, and generalists of bacteria at phylum level based on Levins niche breadth
index. (i). Overall proportion of the habitat niche breadth of bacteria at phylum level. (j).
Proportional distribution of specialists, middle, and generalists of fungi at phylum level. (k).
Overall proportion of the habitat niche breadth of fungi at phylum level. Notes: The *Bocm* was
calculated with Levins index. The *Ocom* was calculated with Schoener index. All ASVs were
defined as specialists, middle or generalists according to Levins niche breadth index. The
asterisks in the boxplots stand for the significance of the Wilcoxon signed-rank test. The "ns"
stands for no significance, single asterisk stands for $P < 0.05$, double asterisks stand for $P < 0.01$,
Triple asterisks stand for $P < 0.001$, Four asterisks stand for $P < 0.0001$.

2.4. Elevated network complexity and resilience to disturbances during the formation of
visually distinguishable microbial communities on red sandstone surface
Given the non-random assembly patterns observed in red sandstone surface microbial
communities, we proceeded to construct microbial co-occurrence networks of both bacteria and
fungi in different types of microbial communities to further explore the interaction patterns
within these communities (Fig. 5).
Topological features analysis of the networks revealed that bacteria network showed higher
links (Fig. 5 a). This indicated that the elevation of the complexity of the bacteria network was
necessary for the formation of visually distinguishable microbial communities. To evaluate the
resilience of the bacteria network to disturbances, we also analyzed the robustness of the
network with different approaches including targeted and randomly removal of key species,
removal of nodes, as well as the calculation of negative correlation ratio. The results showed that
the bacteria network robustness decreased as both targeted and randomly removal of the key
species, with Black, Green, and White groups showing greater vulnerability compared to CK (Fig.
5 b, c). The bacteria networks of the three visually distinguishable microbial communities also
revealed greater vulnerability to the removal of nodes than CK (Fig. 5 d). In addition, the CK
network also revealed higher negative correlation ratio than the Black and White networks of
visually distinguishable microbial communities (Fig. 5 e). This implied the formation of the two
visually distinguishable microbial communities could mitigate the mutual exclusion of microbial
populations in resource competition.
By contrast, the topological features of fungi networks were different from those of bacteria
networks. The number of links in the fungi Green group decreased to half compared to the CK
group, while the number of links in the White group increased by approximately half (Fig. 5 f).
The network robustness analysis results indicated that only the White group demonstrated
higher stability across all test categories (Fig. 5 g-j). This suggests that the role of fungi in
community formation might be more complex and variable depending on the morphology of
communities.

Figure 5. Topological features of the co-occurrence network within microbial communities on the stone surface. (a). Co-occurrence network of bacteria communities. (b). Robustness of bacteria networks under targeted removal of key species. (c). Robustness of bacteria networks under random removal of key species. (d). Vulnerability of bacteria networks under removal of key node. (e). Negative correlation ratio of the bacteria co-occurrence networks within each group. (f). Co-occurrence network of fungi communities. (g). Robustness of fungi networks under targeted removal of key species. (h). Robustness of fungi networks under random removal of key species. (i). Vulnerability of fungi networks under removal of key node. (j). Negative correlation ratio of the fungi co-occurrence networks within each group.

2.5. Fundamental metabolic pathways are maintained but gene profiles vary

To elucidate the metabolic processes that enable the survival of microbial communities on the stone surface, we performed metagenomic sequencing for each type of microbial communities. Ten Gene Ontology (GO) terms that could potentially contribute to the assembly of microbial communities on red sandstone surface were identified in all four types of microbial communities[3]. These terms were associated with functions related to organism interactions, microbial survival, and trophic processes. Each of the ten GO terms was detectable in every group (Fig. 6 a). Next, we used PICRUSt2 to predict the functional potential of the bacterial communities. The results showed significant differences in both enzyme functions (ECs) and metabolic pathway genes (KOs) among the groups (PERMANOVA, P < 0.05) (Fig. 6 b, Fig. S4, Table S8, S9), suggesting that the microbial communities in each group adapted to their specific environmental conditions through distinct metabolic details.

Nitrogen cycle processes have been broadly reported as closely related with the deterioration of stone[3,9,42-44]. Hence, we performed detailed insight into nitrogen cycle processes. The results showed that relative gene abundance of each nitrogen cycle pathway was consistent across all four types of microbial communities (Fig. S5). Ammonium assimilation and nitrogen mineralization were the two dominant processes among the nitrogen cycle. The two processes possessed accounted for up to 38.52% and 31.39% of the overall abundance of reads within the nitrogen cycle pathways, respectively, which imply the intensive nitrogen-exchanging interactions between microbial communities and the stone surface substrate. Nevertheless, the specific high-abundance genes involved in major nitrogen cycling processes differed across groups, indicating that each group harbors distinct metabolic features within individual nitrogen cycling pathways (Fig. 6 c).

Figure 6. Functional profiles of the stone surface microbiota. (a). GO functional annotation of the
stone surface microbiota. (b). Significant differences of enzyme functions (ECs) among four
groups. (c). Nitrogen cycling potential of the stone surface microbiota. Arrow thickness
corresponds to the proportion of functional genes for corresponding metabolic processes.

3. Discussion

Uncovering the diversity, assembly mechanisms, and functional traits of microbial
communities is a central focus in microbial ecology[18,45,46]. Recently, the role of microbial
communities on stone surface has garnered increasing attention, particularly in the field of stone
cultural heritage conservation[47]. In this study, we present a pioneering case study on the
ecological roles of microbiota on red sandstone surface, using the Leshan Giant Buddha as an
example. The results indicate that the taxonomic composition of microbial communities varies
according to their coloration. During the formation of visually distinguishable microbial
communities, core metabolic pathways of microbial communities are maintained but gene
profiles vary across different types of microbial communities, particularly in nitrogen
cycle-related biogeochemical processes. Dispersal limitation and heterogeneous selection
dominate the assembly of stone surface microbial communities. Habitat specialists play a more
significant role than generalists during the formation of microbial communities on the red
sandstone surface. As visible microbial communities form on stone surface, network complexity

increases, and resistance to disturbances is enhanced. By comparing our findings with previous
studies on microbial communities on stone surface, we inferred the following commonalities and
unique characteristics in the ecological features of stone surface microbial communities.
345
3.1. Harsh living conditions on the stone surface might induced conserved core ecological
functions of stone surface microbes
In this study, the microbial diversity analysis revealed that Proteobacteria, Actinobacteria,
Cyanobacteria, Chloroflexi, and Acidobacteriota dominate the top phyla among the prokaryotes
in all samples, while Ascomycota stands as the most dominant phylum in eukaryotes, despite the
distinct microbial community morphology across different sample groups. Such phylum-level
diversity patterns are largely consistent with previous reports on taxonomic diversity of microbial
communities on the surface of other stone types. However, significant differences were observed
at the genus level[2-4,34,48-50]. These variations might be attributed to site-specific factors, including
the local climate and the types of stone cultural heritage[51]. Comparison with previous studies
suggests that core ecological functions of stone surface microbes are conserved across the
surfaces of various types of stone. The capability to withstand pressures such as fluctuations
between wet and dry conditions, endure UV stress on stone surface, and facilitate ion exchange
with the stone appears essential for the persistence of microbial communities on the outdoor
stone surface[3,9].
Nitrogen cycle is a crucial way of ion exchange between microbial communities and the
stone substrate. In previous study, the two dominated types of nitrogen cycling genes in
limestone surface microbial communities are nitrogen assimilation and mineralization, which is
coherent with what we found in red sandstone surface microbial communities in this study[3].
Nitrifying bacteria genera *Nitrospira* and *Nitrosomonas* were significantly enriched in limestone
surface microbial communities[3]. In another case study involving regular sandstone,
Nitrosopumilus and *Nitrospira* were also identified from deteriorated sediments[9]. Furthermore,
Methylobacterium, another dominant genus identified in our study, comprises species recognized
as plant growth-promoting rhizobacteria, with genomes encoding enzymes involved in nitrogen
metabolism and ammonia assimilation[52]. Notably, in previous study, the two dominated types of
nitrogen cycling genes in limestone surface microbial communities are nitrogen assimilation and
mineralization, which is coherent with what we found in red sandstone surface microbial
communities in this study[3]. Despite this, nitrogen metabolic genes are diverse across different
groups, which suggests that members of each type of group microbial community could adapted
to the rock surface niche through distinct metabolic details[53].
Conserved core functions among microbial communities within similar niches is not unique
to microbial communities on stone surface. For example, several gut microbiota studies have
reported functional, but not compositional, convergence of gut microbiomes[54-57]. Some soil
microbiota studies also reported the stable functional structure despite high taxonomic
variability[58-60]. In addition, microbial functional genes of ocean microbiota that execute certain
functions, such as B_{12} biosynthesis, are widely distributed, while the microbial taxa that carry
these genes vary significantly[61]. These results corroborate the ecological hypothesis that
environment truly selects for function, rather than taxonomy[62]. Different taxonomic groups could
reveal convergent functions as a result of functional redundancy among microbial systems[63]. On
stone surface, conserved core functions among various types of stones might be attributed to the

selective pressure imposed by the harsh living conditions on the stone surface, favoring the
survival of microbial communities with specific functions[58,64-66].
388
3.2. Assembly of stone surface microbial communities requires a general pattern of dynamic
interactions
In this study, we witnessed the extensive microbe-microbe and microbe-environment
interactions within the microbial communities on the stone surface. The mutual effect of hetero
selection and dispersal limitation controls the assembly process of both prokaryotic and
eukaryotic communities on red sandstone surface. Such results differ from previous studies on
the assembly process of microbial communities on limestone surface, where the assembly
process is driven by hetero selection and dispersal limitation[3]. This suggests that the assembly
mechanisms of microbial communities on stone surface might vary based on stone types,
geographical locations, climatic characteristics, or other environmental factors. In addition, the
results of the DOC model analysis suggest a general pattern of dynamic interspecific interactions
within the communities on the red sandstone surface and high selective pressure on the red
sandstone surface[38,67-69]. Such complexity was further testified in this study by the topological
feature analysis of the microbial networks. The active biogeochemical metabolism such as
ammonium assimilation and nitrogen mineralization processes by the microbial communities also
suggests their active interactions with the red sandstone substrate[70]. The elevated complexity of
microbial communities might arise from widely functional complementarity and the resulting
extensive interactions among microbes[71].
Here, we propose a hypothesis that a possible model capable of precisely interpreting the
assembly processes of stone surface microbial communities is the classic MacArthur
consumer-resource model, where species-species interactions are mediated through resource
dynamics[72-74]. Because the nutrient-poor nature of stone surface, the colonization by pioneer
species is primarily driven by deterministic processes linked to nutrient availability, which aligns
with commonly observed among natural communities from various locations[75]. Some pioneer
species utilize inorganic energy from the stone to establish themselves. These species then leak
metabolic by-products into the environment, increasing the diversity of resources and enhancing
the biocompatibility of the stone surface. Other species can subsequently utilize these leaked
metabolites, in turn releasing new by-products into the environment, further enhancing the
biocompatibility of the stone surface, and facilitating the formation of microbial communities.
The observed decrease in ecological niche breadth among the visually distinguishable microbial
communities in this study also suggests that the utilization of specific resources is more efficient
within these communities. We also observed that specialization prevails over flexibility during
the formation of microbial communities on red sandstone surface. This is consistent with a
previous study on the assembly of microbial communities in a wide range of well-defined
resource environments, where habitat specialists also drive the increase of diversity in such
microbial communities[19].
425
3.3. Summary
In summary, this study focuses on microbial communities on the surface of red sandstones
of the Leshan Giant Buddha, which reveals their ecological and functional characteristics. Under
the interplay of dispersal limitation and hetero selection, the microbiota on the red sandstone

were assembles into distinct types of visually distinguishable microbial communities.,
significantly differentiate the taxonomic composition of these communities. However, under the
harsh habitant on red sandstone surface, such as UV radiation, fluctuations between dry and wet
conditions, and limited nutrient availability, the core functions of the microbe among different
types of microbial communities are convergent. These functions are particularly enriched in
biogeochemical processes related to nitrogen cycling and responses to environmental stress. The
formation of visually distinguishable microbial communities is associated with a narrower niche
width, elevated complexity, and increased resilience of the prokaryotic network to disturbances.
Furthermore, a higher proportion of habitat specialists are found in microbes forming these
visually distinguishable microbial communities (Fig. 7). Given the nutrient limitations of the red
sandstone surface, the classic MacArthur consumer-resource model might provide a potential
framework to further explain the assembly mechanisms of these microbial communities. Future
research into the ecological traits and functional potential of these microbial communities will
provide critical insights for managing microbial communities on red sandstone surface. Moreover,
microbial deterioration affects not only stone cultural heritage but also a wide range of different
cultural heritage materials[76-80]. The microbial ecology-based approach used in this study will help
advance research on microbial deterioration mechanisms in other cultural heritage materials,
providing a model for future work.

Figure 7. Schematic diagram illustrates the characteristics of microbial communities on the red
sandstone surface of the Leshan Giant Buddha

4. Materials and Methods

4.1. Sampling processes

The Leshan Giant Buddha is a world-famous, typical red sandstone-made, UNESCO World
Heritage site. It was built between 713 and 803 during the Tang dynasty, located in the Leshan
city, Sichuan province, China. Samples used in this study were collected from the stone surface of
the Leshan Giant Buddha between 2019 and 2022. Among these, 6 samples have been used in
our previous study[34]. The other 21 samples were newly collected in 2022. Surface samples were
scraped using a pre-cleaned steel scalpel and then stored in low temperature until DNA

extraction.

4.2. DNA extraction and NGS sequencing

The DNA extraction was applied using a genomic DNA extraction kit from TIANGEN
(DP302-02) with 0.5 g of sample processed for extraction. The amplicon sequencing was
performed as followed. The 16S V3-V4 region rDNA and ITS2 region were amplified by PCR (initial
denaturation at 98 ℃ for 30 seconds; 32cycles of denaturation at 98 ℃ for 10 seconds, annealing
at 54℃ for 30 seconds, and extension at 72 ℃ for 45 seconds; and then final extension at 72 ℃
for 10 minutes.) with following primers: 0341-F (CCT ACG GGN GGC WGC AG) and 806-R (GAC
TAC HVG GGT ATC TAA TCC) for 16S[81], ITS1FI2 (GTG ART CAT CGA ATC TTT G) and ITS2 (TCC TCC
GCT TAT TGA TAT GC) for ITS2. The PCR products were purified by AMPure XT beads (Beckman
Coulter Genomics, Danvers, MA, USA) and quantified by Qubit (Invitrogen, USA). The amplicon
pools were prepared for sequencing and the size and quantity of the amplicon library were
assessed on Agilent 2100 Bioanalyzer (Agilent, USA) and with the Library Quantification Kit for
Illumina (Kapa Biosciences, Woburn, MA, USA), respectively, and then sequenced on Illumina
NovaSeq 6000 PE250 platform in LC-Bio Technology Co., Ltd, Hang Zhou, Zhejiang Province,
China. The raw sequence data were then deposited in the Genome Sequence Archive[82] in
National Genomics Data Center[83], China National Center for Bioinformation / Beijing Institute of
Genomics, Chinese Academy of Sciences (GSA: CRA013382) that are publicly accessible at
https://ngdc.cncb.ac.cn/gsa.
The DNA libraries were sequenced on the illumina NovaseqTM 6000 platform by LC Bio
Technology CO., Ltd (Hangzhou, China). We are grateful to LC Bio Technology CO., Ltd for assisting
in sequencing. DNA concentration was measured using Qubit. Based on the measured
concentration, equal-mass DNA samples were pooled from all samples within each group by
mixing 100 ng of DNA from each sample. The metagenome sequencing was performed as
followed: DNA library was constructed by TruSeq Nano DNA LT Library Preparation Kit
(FC-121-4001). DNA was fragmented by dsDNA Fragmentase (NEB, M0348S) by incubate at 37℃
for 30min. Library construction begins with fragmented cDNA. Blunt-end DNA fragments are
generated using a combination of fill-in reactions and exonuclease activity, and size selection is
performed with provided sample purification beads. An A-base is then added to the blunt ends of
each strand, preparing them for ligation to the indexed adapters. Each adapter contains a T-base
overhang for ligating the adapter to the A-tailed fragmented DNA. These adapters contain the full
complement of sequencing primer hybridization sites for single, paired-end, and indexed reads.
Single- or dual- index adapters are ligated to the fragments and the ligated products are amplified
with PCR. The raw sequence data were then deposited in the Genome Sequence Archive[82] in
National Genomics Data Center[83], China National Center for Bioinformation / Beijing Institute of
Genomics, Chinese Academy of Sciences (GSA: CRA013393) that are publicly accessible at
https://ngdc.cncb.ac.cn/gsa.

4.3. Taxonomic and functional annotations

For the amplicon sequencing data, the DADA2 R package[84] in QIIME 2[85] was used to conduct
a complete pipeline to generate raw paired-end fastq sequencing files into merged, denoised,
chimera-free sequences and output ASV. The ASV tables for both 16S and ITS were rarified to a
minimum sequencing depth (13,622 for 16S and 16,232 for ITS) to mitigate the impact of

sequencing depth on subsequent analysis. Representative ASV sequences were classified into different organisms using a naive Bayesian model with RDP classifier[86], which is based on the SILVA database[87] (version 138) for 16S or UNITE database[88] (version 9.0) for ITS.

Metagenome raw sequencing reads were processed to obtain valid reads for further analysis. First, sequencing adapters were removed from sequencing reads using cutadapt (version 1.9). Secondly, low quality reads were trimmed by fqtrim (version 0.94) using a sliding-window algorithm. Window size is 6 bp, and base quality must not be lower than 20. Once quality-filtered reads were obtained, they were de novo assembled to construct the metagenome for each sample by MEGAHIT with the following parameters: *--k-list 21,29,39,59,79,99,119,141 -m 0.8 -t 8 --min-contig-len 200 --continue*. All coding regions (CDS) of metagenomic contigs were predicted by MetaGeneMark (version 3.26). CDS sequences of all samples were clustered by CD-HIT (version 4.6.1)[89] to obtain unigenes with the following parameters: *-c 0.95 -aS 0.9 -d 0 -g 1 -M 20000 -T 20*. The CDS sequences from all samples are clustered together to group highly similar or identical sequences, and a single representative sequence from each cluster is chosen as a unigene. Unigene abundance for a certain sample were estimated by TPM based on the number of aligned reads by bowtie2 (version 2.2.0)[90]. The TPM value for each Unigene was calculated by dividing its RPK (Reads Per Kilobase) by the sum of RPK values across all Unigenes in the sample, then multiplying the result by 1,000,000 to normalize expression levels to a per-million scale.

The figures in this study was widely plotted in R project ggplot2 package[91]. Top 10 phyla were visualized in R with ggplot2 and ggalluvial as our previous publications[69,76]. Bray-Curtis distance matrix was calculated in R with Vegan package[92] (version 2.5.7). NMDS was then generated with Vegan and plotted with ggplot2 package. Adonis and anosim test were used to calculate the significance of β diversity differences. The tree shape visualization of the taxonomic diversity was performed by Metacoder package in R[93]. The functional annotations (GO, KEGG, CARD, MGEs, VFDBs) of unigenes were obtained by aligning them against the NCBI NR database (version 20191121) by DIAMOND (version 0.9.14)[94] with the following parameters: *-k 25 -e 0.00001 --query-cover 0 --min-score 0 --id 0 -p 20'; Command: srun -p all -c 20 singularity exec --bind /public1/Programs,/public1/Softwares:/Softwares,/public*. Nitrogen cycling genes were annotated with the NCycDB database[95]. Heatmaps were visualized with pheatmap package in R. The functional deduction of bacteria enzyme functions (ECs) and metabolic pathway genes (KOs) was performed by PICRUSt2[96].

4.4. Microbial communities assembly analyses

A dissimilarity-overlap curve (DOC) model was used to determine whether microbiota on the red sandstone surface of the Leshan Giant Buddha have unique ecological dynamics in R with DOC package (https://github.com/Russel88/DOC) according to previous reports[3,68,69]. The DOC was obtained by plotting the dissimilarity (Jensen–Shannon divergence) against the overlap fraction of taxa for each pairwise community[38] with 1000 bootstrap replicates to evaluate the significance and initiation of negative slopes.

Null model analysis was performed as described by Stegen *et al.* [36,37] and visualized as previous research[69]. The deviation of phylogenetic and taxonomic diversity was estimated with 999 randomizations. Beta mean nearest taxon distance (βMNTD) and beta nearest taxon index (βNTI) were calculated with picante package in R[97]. Phylogenetic analysis were performed by iqtree2 with 1000 bootstrap[98]. A significant deviation indicated the dominance of deterministic

processes, including homogeneous or heterogeneous selection processes when βNTI < −2 or
βNTI > 2, respectively. The Bray-Curtis based Raup-Crick (RCbray), the deviation between
Bray-Curtis and the null distribution, was then calculated to partition pairwise comparisons that
assigned the stochastic processes when |βNTI| < 2. The relative influence of homogenizing
dispersal and dispersal limitation was quantified as the fraction of pairwise comparisons with
RCbray < −0.95 and RCbray > 0.95, respectively. In addition, dispersal was seen as a stochastic
process in this research, not restricted to a deterministic or stochastic process as previously
described[3,18]. Low-occurrence ASVs with a prevalence less than 50% and low abundance ASVs
with total read counts lower than the number of samples were excluded to ensure the accuracy
in taxon-specific assembly characterization and reduce type I errors as presented in previous
research[3,69].

4.5. Niche characteristics analysis

Levins niche breadth index, Shannon niche breadth index, Schoener niche overlap index, and Pianka niche overlap index were calculated with spaa package in R[99]. Generalists were defined by ASVs with *Bcom* > 5, whereas specialists were defined by those with *Bcom* < 1.5[3,100]. Significance of the differences was tested by Wilcoxon signed-rank test.

4.6. Network construction and topological feature analysis

Microbial co-occurrence networks were constructed according to previous publication[101] with modifications by ggClusterNet R package[102]. Topological features were also further visualized with ggClusterNet in R. Spearman's correlation was applied to assess the relationships between microbial taxa. Correlation coefficients with a threshold of r >= 0.6 and P values <= 0.05 were retained for network construction to ensure significant and robust associations. The network was further clustered using the *cluster_fast_greedy* algorithm, which identifies densely connected sub-networks within the microbial community.

4.7. Acknowledgements of English edition

During the preparation of this work the authors used ChatGPT in order to check the English grammar. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

Availability of data and materials

The raw sequence data reported in this paper have been deposited in the Genome Sequence Archive in National Genomics Data Center, China National Center for Bioinformation / Beijing Institute of Genomics, Chinese Academy of Sciences (GSA: CRA013382 & GSA: CRA013393) that are publicly accessible at https://ngdc.cncb.ac.cn/gsa.

All scripts used in this study are publicly available at the GitHub repository (https://github.com/Bowenw6/Microbiome_Buddha_Grotto_2023).

Author Contributions Statement

Bowen Wang: Conceptualization, Data curation, Formal analysis, Investigation,
Methodology, Resources, Visualization, Writing - Original Draft, Writing - Review & Editing.
Chengshuai Zhu: Conceptualization, Data curation, Investigation, Resources. **Xin Wang**:
Conceptualization, Investigation, Resources. **Tianyu Yang**: Conceptualization, Investigation,
Resources. **Bingjian Zhang**: Conceptualization, Funding acquisition, Supervision. **Yulan Hu**:
Conceptualization, Funding acquisition, Project administration, Supervision, Writing - Review &
Editing.
598

Competing interests

600   No, we declare that the authors have no competing interests, or other interests that might
601   be perceived to influence the results and/or discussion reported in this paper.
602
603

Funding

605   This work is supported by the National Key Research and Development Program
606   (2019YFC1520503)
607
608

Supplementary Files

Figure S1. The overall taxonomic community composition among all the samples.
Figure S2. Taxonomic community composition of the microbial communities in genus level.
Figure S3. Niche characterization of the stone surface microbial communities based on Shannon
niche breadth index and Pianka niche overlap index.
Figure S4. Significant differences of metabolic pathway genes (KOs) among four groups.
Figure S5. Gene abundance within each nitrogen cycle pathway of the stone surface microbiota.
Table S1. Summaries for the details from the metagenome sequencing
Table S2. Relative abundances of dominant bacteria among 27 samples at the phylum level.
Table S3. Relative abundances of dominant bacteria among 27 samples at the genus level.
Table S4. Relative abundances of dominant fungi among 27 samples at the phylum level.
Table S5. Relative abundances of dominant fungi among 27 samples at the genus level.
Table S6. Habitat niche breadth of bacteria at the phylum level.
Table S7. Habitat niche breadth of fungi at the phylum level.
Table S8. Enzyme functions (ECs) among all samples.
Table S9. Metabolic pathway genes (KOs) among all samples.
625
626

627   **References:**
628

1   Liu, X. B., Koestler, R. J., Warscheid, T., Katayama, Y. & Gu, J. D. Microbial deterioration and
sustainable conservation of stone monuments and buildings. *Nat Sustain* **3**, 991-1004 (2020).
https://doi.org/10.1038/s41893-020-00602-5

2   Brewer, T. E. & Fierer, N. Tales from the tomb: the microbial ecology of exposed rock surfaces.
Environ Microbiol **20**, 958-970 (2018). https://doi.org/10.1111/1462-2920.14024
3   He, J. *et al.* From surviving to thriving, the assembly processes of microbial communities in
stone biodeterioration: A case study of the West Lake UNESCO World Heritage area in China.
Sci Total Environ **805**, 150395 (2022). https://doi.org/10.1016/j.scitotenv.2021.150395
4   Qiang, L., Zhang, B. B., Yang, B. X. & Ged, C. Q. Deterioration-Associated Microbiome of Stone
Monuments: Structure, Variation, and Assembly. *Appl Environ Microbiol* **84**,
e02680-02617-02617 (2018).
5   Li, T., Hu, Y. & Zhang, B. Biomineralization Induced by Colletotrichum acutatum: A Potential
Strategy for Cultural Relic Bioprotection. *Front Microbiol* **9**, 1884 (2018).
6   Jroundi, F., Gonzalez-Muñoz, M. T. & Rodriguez-Navarro, C. in *Microorganisms in the*
Deterioration and Preservation of Cultural Heritage   (ed Edith Joseph)   281-299 (Springer
International Publishing, 2021).
7   Liu, X. *et al.* Biofilms on stone monuments: biodeterioration or bioprotection? *Trends*
Microbiol **30**, 816-819 (2022). https://doi.org/https://doi.org/10.1016/j.tim.2022.05.012
8   Li, J. *et al.* The active microbes and biochemical processes contributing to deterioration of
Angkor sandstone monuments under the tropical climate in Cambodia – A review. *J Cult Herit*
47, 218-226 (2021). https://doi.org/https://doi.org/10.1016/j.culher.2020.10.010
9   Ding, X. *et al.* Microbiome and nitrate removal processes by microorganisms on the ancient
Preah Vihear temple of Cambodia revealed by metagenomics and N-15 isotope analyses.
Appl Microbiol Biotechnol **104**, 9823-9837 (2020).
https://doi.org/10.1007/s00253-020-10886-4
10   Wu, F. *et al.* Metagenomic and metaproteomic insights into the microbiome and the key
geobiochemical potentials on the sandstone of rock-hewn Beishiku Temple in Northwest
China. *Sci Total Environ* **893**, 164616 (2023).
https://doi.org/https://doi.org/10.1016/j.scitotenv.2023.164616
11   Fidanza, M. R. & Caneva, G. Natural biocides for the conservation of stone cultural heritage: A
review. *J Cult Herit* **38**, 271-286 (2019).
https://doi.org/https://doi.org/10.1016/j.culher.2019.01.005
12   Li, Q., Wu, C., He, J. & Zhang, B. Unraveling the microbiotas and key genetic contexts
identified on stone heritage using illumina and nanopore sequencing platforms. *Int Biodeter*
Biodegr **185**, 105688 (2023). https://doi.org/https://doi.org/10.1016/j.ibiod.2023.105688
13   Ding, X. *et al.* Microbiome characteristics and the key biochemical reactions identified on
stone world cultural heritage under different climate conditions. *Journal of Environmental*
Management **302**, 114041 (2022).
https://doi.org/https://doi.org/10.1016/j.jenvman.2021.114041
14   Ennis, N. J., Dharumaduri, D., Bryce, J. G. & Tisa, L. S. Metagenome Across a Geochemical
Gradient of Indian Stone Ruins Found at Historic Sites in Tamil Nadu, India. *Microb Ecol* **81**,
385-395 (2021). https://doi.org/10.1007/s00248-020-01598-3
15   Louati, M. *et al.* Elucidating the ecological networks in stone-dwelling microbiomes. *Environ*
Microbiol **22**, 1467-1480 (2020). https://doi.org/https://doi.org/10.1111/1462-2920.14700
16   Li, T., Cai, Y. & Ma, Q. Microbial Diversity on the Surface of Historical Monuments in Lingyan
Temple, Jinan, China. *Microb Ecol* (2022). https://doi.org/10.1007/s00248-021-01955-w
17   Cattò, C. *et al.* Biofilm colonization of stone materials from an Australian outdoor sculpture:

Importance of geometry and exposure. *Journal of Environmental Management* **339**, 117948
(2023). https://doi.org/https://doi.org/10.1016/j.jenvman.2023.117948

18   Zhou, J. & Ning, D. Stochastic Community Assembly: Does It Matter in Microbial Ecology?
MMBR **81**, e00002-00017 (2017). https://doi.org/10.1128/MMBR.00002-17

19   Dal Bello, M., Lee, H., Goyal, A. & Gore, J. Resource–diversity relationships in bacterial
communities reflect the network structure of microbial metabolism. *Nature Ecology &*
Evolution **5**, 1424-1434 (2021). https://doi.org/10.1038/s41559-021-01535-8

20   Goldford, J. E. *et al.* Emergent simplicity in microbial community assembly. *Science* **361**,
469-474 (2018). https://doi.org/10.1126/science.aat1168

21   Ning, D. *et al.* A quantitative framework reveals ecological drivers of grassland microbial
community assembly in response to warming. *Nat Commun* **11**, 4717 (2020).
https://doi.org/10.1038/s41467-020-18560-z

22   Chase, J. M. Stochastic Community Assembly Causes Higher Biodiversity in More Productive
Environments. *Science* **328**, 1388-1391 (2010). https://doi.org/10.1126/science.1187820

23   Ofițeru, I. D. *et al.* Combined niche and neutral effects in a microbial wastewater treatment
community. *PNAS* **107**, 15345-15350 (2010). https://doi.org/10.1073/pnas.1000604107

24   Chesson, P. Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology,*
Evolution,   and   Systematics   **31**,   343-366   (2000).
https://doi.org/https://doi.org/10.1146/annurev.ecolsys.31.1.343

25   Fargione, J., Brown, C. S. & Tilman, D. Community assembly and invasion: An experimental
test   of   neutral   versus   niche   processes.   *PNAS*   **100**,   8916-8920   (2003).
https://doi.org/10.1073/pnas.1033107100

26   Chave, J. Neutral theory and community ecology. *Ecol Lett* **7**, 241-253 (2004).
https://doi.org/https://doi.org/10.1111/j.1461-0248.2003.00566.x

27   Vega, N. M. & Gore, J. Stochastic assembly produces heterogeneous communities in the
Caenorhabditis   elegans   intestine.   *PLOS*   *Biology*   **15**,   e2000633   (2017).
https://doi.org/10.1371/journal.pbio.2000633

28   Ning, D. *et al.* Environmental stress mediates groundwater microbial community assembly.
Nature Microbiology **9**, 490-501 (2024). https://doi.org/10.1038/s41564-023-01573-x

29   He, Q. *et al.* Temperature and microbial interactions drive the deterministic assembly
processes   in   sediments   of   hot   springs.   *Sci   Total   Environ*   **772**,   145465   (2021).
https://doi.org/https://doi.org/10.1016/j.scitotenv.2021.145465

30   Tolkkinen, M. *et al.* Multi-stressor impacts on fungal diversity and ecosystem functions in
streams: natural   vs.   anthropogenic   stress.   *Ecology*   **96**,   672-683   (2015).
https://doi.org/https://doi.org/10.1890/14-0743.1

31   Huot, O. B. & Tamborindeguy, C. Drought stress affects olanum lycopersicum susceptibility to
actericera cockerelli colonization. *Entomologia Experimentalis et Applicata* **165**, 70-82 (2017).
https://doi.org/https://doi.org/10.1111/eea.12627

32   Luo, C., Lü, F., Shao, L. & He, P. Application of eco-compatible biochar in anaerobic digestion
to relieve acid stress and promote the selective colonization of functional microbes. *Water*
Res **68**, 710-718 (2015). https://doi.org/https://doi.org/10.1016/j.watres.2014.10.052

33   Bai, F. *et al.* Microbial biofilms on a giant monolithic statue of Buddha: The symbiosis of
microorganisms and mosses and implications for bioweathering. *Int Biodeter Biodegr* **156**
(2021). https://doi.org/ARTN 105106 10.1016/j.ibiod.2020.105106

34   Zhu, C. *et al.* Analysis of the Microbiomes on Two Cultural Heritage Sites. *Geomicrobiol J*, 1-10
(2022). https://doi.org/10.1080/01490451.2022.2137604
35   Zhu, C. *et al.* Unveiling the dual role of biocolonization: a case study on the deterioration and
preservation of sandstone monuments in Leshan Giant Buddha, China. *World Journal of*
Microbiology and Biotechnology **41**, 25 (2025). https://doi.org/10.1007/s11274-024-04237-y
36   Stegen, J. C., Lin, X., Konopka, A. E. & Fredrickson, J. K. Stochastic and deterministic assembly
processes in subsurface microbial communities. *ISME J* **6**, 1653-1664 (2012).
https://doi.org/10.1038/ismej.2012.22
37   Stegen, J. C. *et al.* Quantifying community assembly processes and identifying features that
impose them. *ISME J* **7**, 2069-2079 (2013). https://doi.org/10.1038/ismej.2013.93
38   Bashan, A. *et al.* Universality of human microbial dynamics. *Nature* **534**, 259-262 (2016).
https://doi.org/10.1038/nature18301
39   Levins, R. *Evolution in Changing Environments Some Theoretical Explorations. (MPB-2)*.
(Princeton University Press, 1968).
40   Sexton, J. P., Montiel, J., Shay, J. E., Stephens, M. R. & Slatyer, R. A. Evolution of Ecological
Niche Breadth. *Annual Review of Ecology, Evolution, and Systematics* **48**, 183-206 (2017).
https://doi.org/10.1146/annurev-ecolsys-110316-023003
41   von Meijenfeldt, F. A. B., Hogeweg, P. & Dutilh, B. E. A social niche breadth score reveals niche
range strategies of generalists and specialists. *Nature Ecology & Evolution* **7**, 768-781 (2023).
https://doi.org/10.1038/s41559-023-02027-7
42   Meng, H., Zhang, X., Katayama, Y., Ge, Q. & Gu, J.-D. Microbial diversity and composition of
the Preah Vihear temple in Cambodia by high-throughput sequencing based on genomic DNA
and RNA. *Int Biodeter Biodegr* **149** (2020).
43   Meng, S. *et al.* Community structures and biodeterioration processes of epilithic biofilms
imply the significance of micro-environments. *Sci Total Environ* **876**, 162665 (2023).
https://doi.org/https://doi.org/10.1016/j.scitotenv.2023.162665
44   Chen, X. *et al.* The Organisms on Rock Cultural Heritages: Growth and Weathering.
Geoheritage **13** (2021). https://doi.org/ARTN 5610.1007/s12371-021-00588-2
45   Escalas, A. *et al.* Microbial functional diversity: From concepts to applications. *Ecology and*
Evolution **9**, 12000-12016 (2019). https://doi.org/https://doi.org/10.1002/ece3.5670
46   Prosser, J. I. *et al.* The role of ecological theory in microbial ecology. *Nat Rev Microbiol* **5**,
384-392 (2007). https://doi.org/10.1038/nrmicro1643
47   Sterflinger, K. *et al.* Future directions and challenges in biodeterioration research on historic
materials and cultural properties. *Int Biodeter Biodegr* **129**, 10-12 (2018).
https://doi.org/https://doi.org/10.1016/j.ibiod.2017.12.007
48   Wu, Y. *et al.* Monitoring the Deterioration of Masonry Relics at a UNESCO World Heritage Site.
KSCE Journal of Civil Engineering **25**, 3097-3106 (2021).
https://doi.org/10.1007/s12205-021-1716-z
49   Zhang, X. W., Ge, Q. Y., Zhu, Z. B., Deng, Y. M. & Gu, J. D. Microbiological community of the
Royal Palace in Angkor Thom and Beng Mealea of Cambodia by Illumina sequencing based on
16S rRNA gene. *Int Biodeter Biodegr* **134**, 127-135 (2018).
https://doi.org/10.1016/j.ibiod.2018.06.018
50   Schröer, L., De Kock, T., Cnudde, V. & Boon, N. Differential colonization of microbial
communities inhabiting Lede stone in the urban and rural environment. *Sci Total Environ* **733**,

139339 (2020). https://doi.org/https://doi.org/10.1016/j.scitotenv.2020.139339

51 Qian, Y., Liu, X., Hu, P., Gao, L. & Gu, J.-D. Identifying the major metabolic potentials of
microbial-driven carbon, nitrogen and sulfur cycling on stone cultural heritage worldwide. *Sci*
Total   Environ   954,   176757   (2024).
https://doi.org/https://doi.org/10.1016/j.scitotenv.2024.176757

52 Grossi, C. E. M. *et al.* Methylobacterium sp. 2A Is a Plant Growth-Promoting Rhizobacteria
That Has the Potential to Improve Potato Crop Yield Under Adverse Conditions. *Frontiers in*
Plant Science **11** (2020). https://doi.org/10.3389/fpls.2020.00071

53 Ramond, P., Galand, P. E. & Logares, R. Microbial functional diversity and redundancy: moving
forward.   *FEMS   Microbiology   Reviews   49*,   fuae031   (2025).
https://doi.org/10.1093/femsre/fuae031

54 Tian, L. *et al.* Deciphering functional redundancy in the human microbiome. *Nat Commun* **11**,
6217 (2020). https://doi.org/10.1038/s41467-020-19940-1

55 Li, H. *et al.* Food provisioning results in functional, but not compositional, convergence of the
gut microbiomes of two wild Rhinopithecus species: Evidence of functional redundancy in the
gut   microbiome.   *Sci   Total   Environ   858*,   159957   (2023).
https://doi.org/https://doi.org/10.1016/j.scitotenv.2022.159957

56 Moya, A. & Ferrer, M. Functional Redundancy-Induced Stability of Gut Microbiota Subjected
to   Disturbance.   *Trends   Microbiol   24*,   402-413   (2016).
https://doi.org/10.1016/j.tim.2016.02.002

57 Tian, L. & Zhang, J. Dynamics of Rabies Epidemics in Vampire Bats. *Complexity* **2020**, 7032451
(2020). https://doi.org/https://doi.org/10.1155/2020/7032451

58 Talbot, J. M. *et al.* Endemism and functional convergence across the North American soil
mycobiome. *PNAS* **111**, 6341-6346 (2014). https://doi.org/10.1073/pnas.1402584111

59 Zeng, Q. *et al.* Stable functional structure despite high taxonomic variability across fungal
communities in soils of old-growth montane forests. *Microbiome* **11**, 217 (2023).
https://doi.org/10.1186/s40168-023-01650-7

60 Chen, H. *et al.* Functional Redundancy in Soil Microbial Community Based on Metagenomics
Across the Globe. *Front Microbiol* **13** (2022).

61 Zhou, J. *et al.* The diversity and ecological significance of microbial traits potentially involved
in   B12   biosynthesis   in   the   global   ocean.   *mLife*   **2**,   416-427   (2023).
https://doi.org/https://doi.org/10.1002/mlf2.12095

62 Burke, C., Steinberg, P., Rusch, D., Kjelleberg, S. & Thomas, T. Bacterial community assembly
based   on   functional   genes   rather   than   species.   *PNAS*   **108**,   14288-14293   (2011).
https://doi.org/10.1073/pnas.1101591108

63 Louca, S. *et al.* Function and functional redundancy in microbial systems. *Nature Ecology &*
Evolution **2**, 936-943 (2018). https://doi.org/10.1038/s41559-018-0519-1

64 Chikina, M., Robinson, J. D. & Clark, N. L. Hundreds of Genes Experienced Convergent Shifts in
Selective Pressure in Marine Mammals. *Molecular Biology and Evolution* **33**, 2182-2192
(2016). https://doi.org/10.1093/molbev/msw112

65 Winemiller, K. O., Fitzgerald, D. B., Bower, L. M. & Pianka, E. R. Functional traits, convergent
evolution,   and   periodic   tables   of   niches.   *Ecol   Lett   18*,   737-751   (2015).
https://doi.org/https://doi.org/10.1111/ele.12462

66 Nie, Y. *et al.* Species Divergence vs. Functional Convergence Characterizes Crude Oil Microbial

Community Assembly. *Front Microbiol* **7** (2016).

67   Li, P. *et al.* Global diversity and biogeography of potential phytopathogenic fungi in a changing
world. *Nat Commun* **14**, 6482 (2023). https://doi.org/10.1038/s41467-023-42142-4

68   Gao, C. *et al.* Fungal community assembly in drought-stressed sorghum shows stochasticity,
selection, and universal ecological dynamics. *Nat Commun* **11**, 34 (2020).
https://doi.org/10.1038/s41467-019-13913-9

69   Wang, B., Zhu, C., Hu, Y., Zhang, B. & Wang, J. Dynamics of microbial community composition
during degradation of silks in burial environment. *Sci Total Environ* **883**, 163694 (2023).
https://doi.org/https://doi.org/10.1016/j.scitotenv.2023.163694

70   Meng, H., Luo, L., Chan, H. W., Katayama, Y. & Gu, J.-D. Higher diversity and abundance of
ammonia-oxidizing archaea than bacteria detected at the Bayon Temple of Angkor Thom in
Cambodia. *Int Biodeter Biodegr* **115**, 234-243 (2016).
https://doi.org/https://doi.org/10.1016/j.ibiod.2016.08.021

71   Puente-Sánchez, F., Pascual-García, A., Bastolla, U., Pedrós-Alió, C. & Tamames, J.
Cross-biome microbial networks reveal functional redundancy and suggest genome reduction
through functional complementarity. *Communications Biology* **7**, 1046 (2024).
https://doi.org/10.1038/s42003-024-06616-5

72   MacArthur, R. Species packing and competitive equilibrium for many species. *Theoretical*
Population Biology **1**, 1-11 (1970).
https://doi.org/https://doi.org/10.1016/0040-5809(70)90039-0

73   Advani, M., Bunin, G. & Mehta, P. Statistical physics of community ecology: a cavity solution
to MacArthur's consumer resource model. *Journal of Statistical Mechanics: Theory and*
Experiment **2018**, 033406 (2018). https://doi.org/10.1088/1742-5468/aab04e

74   Marsland, R., III *et al.* Available energy fluxes drive a transition in the diversity, stability, and
functional structure of microbial communities. *PLOS Computational Biology* **15**, e1006793
(2019). https://doi.org/10.1371/journal.pcbi.1006793

75   Sriswasdi, S., Yang, C.-c. & Iwasaki, W. Generalist species drive microbial dispersion and
evolution. *Nat Commun* **8**, 1162 (2017). https://doi.org/10.1038/s41467-017-01265-1

76   Wang, B., Qi, M., Ma, Y., Zhang, B. & Hu, Y. Microbiome Diversity and Cellulose
Decomposition Processes by Microorganisms on the Ancient Wooden Seawall of Qiantang
River of Hangzhou, China. *Microb Ecol* **86**, 2109-2119 (2023).
https://doi.org/10.1007/s00248-023-02221-x

77   Wang, B., Zhu, C., Wang, B., Zhang, B. & Hu, Y. Analysis of the biocorrosion community from
ancient wooden constructions at Tianluoshan (7000–6300 cal BP), Zhejiang Province, China.
Heritage Science **12**, 189 (2024). https://doi.org/10.1186/s40494-024-01304-3

78   Zhu, C. *et al.* Application and evaluation of a new blend of biocides for biological control on
cultural heritages. *Int Biodeter Biodegr* **178**, 105569 (2023).
https://doi.org/https://doi.org/10.1016/j.ibiod.2023.105569

79   Li, J. *et al.* Changes in soil microbial communities at Jinsha earthen site are associated with
earthen site deterioration. *BMC Microbiology* **20**, 147 (2020).
https://doi.org/10.1186/s12866-020-01836-1

80   Wang, X., Wang, B., Hu, Y., Zhang, Z. & Zhang, B. Activity-based protein profiling technology
reveals malate dehydrogenase as the target protein of cinnamaldehyde against Aspergillus
niger. *Int J Food Microbiol* **417**, 110685 (2024).

https://doi.org/https://doi.org/10.1016/j.ijfoodmicro.2024.110685

81   Logue, J. B. *et al.* Experimental insights into the importance of aquatic bacterial community
composition to the degradation of dissolved organic matter. *ISME J* **10**, 533-545 (2016).
https://doi.org/10.1038/ismej.2015.131

82   Chen, T. *et al.* The Genome Sequence Archive Family: Toward Explosive Data Growth and
Diverse Data Types. *Genomics, Proteomics & Bioinformatics* **19**, 578-583 (2021).
https://doi.org/https://doi.org/10.1016/j.gpb.2021.08.001

83   Members, C.-N. & Partners. Database Resources of the National Genomics Data Center, China
National Center for Bioinformation in 2022. *Nucleic Acids Research* **50**, D27-D38 (2022).
https://doi.org/10.1093/nar/gkab951

84   Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon data.
Nat Methods **13**, 581-583 (2016). https://doi.org/10.1038/nmeth.3869

85   Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data science
using QIIME 2. *Nat Biotechnol* **37**, 852-857 (2019).
https://doi.org/10.1038/s41587-019-0209-9

86   Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Nave Bayesian Classifier for Rapid
Assignment of rRNA Sequences into the New Bacterial Taxonomy. *Appl Environ Microbiol* **73**,
5261 (2007).

87   Pruesse, E. *et al.* SILVA: a comprehensive online resource for quality checked and aligned
ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res* **35**, 7188-7196 (2007).
https://doi.org/10.1093/nar/gkm864

88   Nilsson, R. H. *et al.* The UNITE database for molecular identification of fungi: handling dark
taxa and parallel taxonomic classifications. *Nucleic Acids Research* **47**, D259-D264 (2019).
https://doi.org/10.1093/nar/gky1022

89   Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-generation
sequencing data. *Bioinformatics* **28**, 3150-3152 (2012).
https://doi.org/10.1093/bioinformatics/bts565

90   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**,
357-359 (2012). https://doi.org/10.1038/nmeth.1923

91   Wickham, H. ggplot2. *WIREs Computational Statistics* **3**, 180-185 (2011).
https://doi.org/https://doi.org/10.1002/wics.147

92   Dixon, P. VEGAN, a package of R functions for community ecology. *Journal of Vegetation*
Science **14**, 927-930 (2003).
https://doi.org/https://doi.org/10.1111/j.1654-1103.2003.tb02228.x

93   Foster, Z. S. L., Sharpton, T. J. & Grünwald, N. J. Metacoder: An R package for visualization and
manipulation of community taxonomic diversity data. *PLOS Computational Biology* **13**,
e1005404 (2017). https://doi.org/10.1371/journal.pcbi.1005404

94   Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND. *Nat*
Methods **12**, 59-60 (2015). https://doi.org/10.1038/nmeth.3176

95   Tu, Q., Lin, L., Cheng, L., Deng, Y. & He, Z. NCycDB: a curated integrative database for fast and
accurate metagenomic profiling of nitrogen cycling genes. *Bioinformatics* **35**, 1040-1048
(2019). https://doi.org/10.1093/bioinformatics/bty741

96   Douglas, G. M. *et al.* PICRUSt2 for prediction of metagenome functions. *Nat Biotechnol* **38**,
685-688 (2020). https://doi.org/10.1038/s41587-020-0548-6

97   Kembel, S. W. *et al.* Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* **26**,
1463-1464 (2010). https://doi.org/10.1093/bioinformatics/btq166

98   Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A Fast and Effective
Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and*
Evolution **32**, 268-274 (2015). https://doi.org/10.1093/molbev/msu300

99   Zhang, J. M., KP. spaa: An R package for computing species association and niche overlap. .
Research Progress of Biodiversity Conservation in China (in Chinese) **X**, 165-174 (2014).

100 Pandit, S. N., Kolasa, J. & Cottenie, K. Contrasts between habitat generalists and specialists:
an empirical extension to the basic metacommunity framework. *Ecology* **90**, 2253-2262
(2009). https://doi.org/https://doi.org/10.1890/08-0851.1

101 Hao, X. *et al.* Effects on community composition and function Pinus massoniana infected by
Bursaphelenchus   xylophilus.   *BMC*   *Microbiology*   **22**,   157   (2022).
https://doi.org/10.1186/s12866-022-02569-z

102 Wen, T. *et al.* ggClusterNet: An R package for microbiome network analysis and
modularity-based   multiple   network   layouts.   *iMeta*   **1**,   e32   (2022).
https://doi.org/https://doi.org/10.1002/imt2.32

912