Red sandstone surface microbiota: functional convergence despite taxonomic

divergence under dispersal limitation and hetero selection

Bowen Wang[a, b], Chengshuai Zhu [a], Xin Wang[c], Tianyu Yang[d], Bingjian Zhang [a, c] and Yulan Hu[a*]

[a] Department of Archaeology, Cultural Heritage and Museology, Zhejiang University, Hangzhou,
310028, China.
[b] Department of Biology, Johns Hopkins University, Baltimore, MD, 21218, USA.
[c] Department of Chemistry, Zhejiang University, Hangzhou, 310028, China.
[d] Leshan Giant Buddha Grottoes Research Institute, Leshan Giant Buddha Scenic Area
Management Committee, Leshan, 614000, China.

[*]Corresponding: hu_yulan@zju.edu.cn

Abstract

Understanding the role of microbiota on stone surfaces is essential for developing effective
grottoes conservation strategies. However, ecological feature of microbial communities on stone
surfaces remains underexplored. In this study, we explored diversity, ecological feature, and
functional profiles of microbial communities on the red sandstone surface of the Leshan Giant
Buddha from microbial ecology perspective. The results show that Proteobacteria, Actinobacteria,
and Cyanobacteria are the dominant prokaryotic phyla, and Ascomycota is the most dominant
eukaryotic phylum. Interestingly, despite taxonomic divergence, the functional profiles of different
types of the microbial communities remain convergent across all samples. As indicated by the
ecological modelling, the interplay of dispersal limitation and hetero selection might have
contributed to such species divergence, while selective pressure from the harsh environment on
the stone surface promoted functional convergence. The assembly of visually distinct microbial
communities is linked to a narrower ecological niche, higher proportion of habitat specialists,
elevated complexity, and increased resilience of the prokaryotic network to disturbances.
Microbial-mediated ammonium assimilation and nitrogen mineralization might be the two
prominent processes that contribute to stone biodeterioration. This study deepens our
understanding of microbial community assembly mechanism on stone cultural heritage surfaces
and functional potentials, which provides microbial ecological insights for the conservation of
these cultural treasures.

Keywords: Microbiota; Biodeterioration; Stone; Cultural heritage; Functional convergence;
Community assembly; Microbial ecology

1. Introduction

Grottoes are a common form of stone cultural heritage. The stone surfaces of grottoes are integral components of ecosystems, providing critical habitats for microbial communities, serving as crucial habitats for microorganisms organized into diverse microbial communities[1-3]. These microbial communities not only diminish the aesthetic value of the artifacts by obscuring and deteriorating the carvings but may also, in some cases, compromise the structural integrity of the grottoes through severe erosion process[3-7]. Conversely, some studies have shown that under specific conditions, microbial communities on stone surfaces can contribute to the reinforcement of the stone through metabolic processes such as biomineralization[1,8-11]. Understanding the role of microorganisms on stone surfaces is essential for formulating better conservation strategies for grotto heritage. Current research in this field primarily focuses on the taxonomic diversity of microorganisms and their biogeochemical cycling and metabolic capabilities. These studies have revealed the key microbial groups involved in the biodeterioration of stone cultural heritage and their associated biogeochemical processes and metabolic responses[6,12-17]. However, ecological understanding of microbial communities on stone surfaces remains significantly limited.

The ecological mechanisms that govern microbial community assembly are one of the main scientific questions in ecological research[18-20]. It is generally believed that the assembly of local microbial communities is controlled by the combined effects of deterministic and stochastic processes[21-23]. Deterministic processes are used to describe the traditional niche-based theory, which hypothesize that species traits, interspecific interactions, and environmental factors govern community structure[24,25]. In contrast, stochastic processes are based on the neutral theory, which illustrate that community assembly is driven by stochastic events such as birth, death, colonization, extinction, and speciation[26,27]. A recent study on microbial communities on limestone surface revealed that prokaryotes and fungi in this environment are primarily dominated by the mutual effect of dispersal limitation (stochastic process) and homogeneous selection (deterministic process) [3]. However, it is currently unknown whether such community assembly mechanisms are specific to certain stone type, geographic location, or influenced by other factors.

The Leshan Giant Buddha, a world-renowned UNESCO World Heritage site, is a typical example of red sandstone grotto cultural heritage. The cleaning of stone surface is a crucial task for conservation of the grotto. However, despite regular cleaning of the stone surfaces by heritage conservators, microbial communities persistently reappear after each cleaning. To gain deeper insights into these continuously reappearing microbial communities on the stone surface of the Leshan Giant Buddha, several studies have explored their basic diversity and tested the abundance of certain nitrogen metabolic-related genes[28,29]. However, beyond this, our current knowledge regarding the microbial communities on the stone surface of the Leshan Giant Buddha, or even on other red sandstone surfaces, remains limited. The ecological mechanism of how microbes survive and form microbial communities within such ecosystem is still unknown. Therefore, this study uses the Leshan Giant Buddha, a UNESCO World Heritage site, as a case study to comprehensively evaluate the characteristics of microbial communities on the surface of this red sandstone cultural heritage from a microbial ecological perspective. Our goal is to gain ecological insights into the microbial communities on stone surfaces, providing a pioneering case study for understanding the

85　ecological roles of the stone surface microbiota. We aim to investigate how microorganisms on
86　stone surfaces cope with frequent wet-dry cycles, nutrient limitations, and UV radiation stress,
87　how they assemble microbial communities in such environments, and what functional potential
88　and characteristics they exhibit. Our focus extends beyond the interactions within the microbiota
89　to include the interactions among microorganisms, environmental factors, and the stone substrate.

90　2. Results

91　2.1. Taxonomic composition varies by the color of microbial communities

Samples of three different types of visually distinguishable microbial communities, along with
blank samples, were collected from the stone surface of the Leshan Giant Buddha. In general, the
total 27 samples were grouped and named according to their respective colors (Bla, black; Gre,
green; Whi, white; and CK, blank control) (Fig. 1). We then performed amplicon sequencing of
conserved phylogenetic marker genes (16S and ITS rRNA) to identify prokaryotes and eukaryotes
within the microbial communities. After quality filtering and rarefaction, a total of 8928 and 2419
ASVs were obtained from the 16S and ITS sequences, respectively. Each group of data overlap with
others, but they are not entirely identical or inclusive of each other (Fig. S1). In addition,
metagenome sequencing was applied for each group and obtained 206, 491, 166 clean reads in
total (Table S1).
102

Figure 1. Representative examples and sampling of microbial communities on the red sandstone
surface of the Leshan Giant Buddha. (a). Location of the Leshan Giant Buddha. (b). Sampling
regions of the samples. Red boxes here indicate the specific sampling regions. (c). Representative
examples of CK samples. (d). Representative examples of Black samples. (e). Representative
examples of Green samples. (f). Representative examples of White samples. Notes: "n" in picture
c-f represents the number of samples used in this study within each group.

To get taxonomic insights into the microbiota, we annotated the sequencing data, then

visualized top 10 dominant phyla, top 20 dominant genus of prokaryotes (Fig. 2 a; Fig. S2 a; Table

S2, S3) and eukaryotes (Fig. 2 b; Fig. S2 b; Table S4, S5) among all samples. We also constructed

overall taxonomic community composition trees of top 5% dominant prokaryotes, and top 30%

dominant eukaryotes (Fig. 2 c, d). The results show that among all samples, Proteobacteria,

Actinobacteria, and Cyanobacteria are the dominant top three phyla in prokaryotes, comprising at

least 70% of the total prokaryotic community in each sample. Ascomycota is the most dominant

phylum in eukaryotes, comprising at least 50% of the total eukaryotic community in each sample.

To estimate the separation pattern between each group of microbial communities, non-metric

multidimensional scaling (NMDS) based on Bray-Curtis dissimilarity at the ASV level was performed

and then conducted Anosim (Analysis of similarities) and PERMANOVA (i.e., adonis) tests (Fig. 2 e,

f). The results show that both prokaryotes and eukaryotes among the four groups of samples reveal

significant and major differences ($P < 0.001$). In summary, the structure of microbial communities

varies according to the coloration of the samples, which indicates distinct microbial compositions

among samples of different colors.

Figure 2. Taxonomic composition and distinct taxonomic structures of the microbial communities.
(a). Relative abundances of dominant prokaryotes among all samples in phylum level. (b). Relative
abundances of dominant prokaryotes among all samples in genus level. (c). The overall taxonomic
community composition tree of top 5% dominant prokaryotes. (d). The overall taxonomic
community composition tree of top 30% dominant eukaryotes. (e). Non-metric multidimensional
scaling (NMDS) ordination based on Bray-Curtis dissimilarities at the ASV level for prokaryotic
communities. (f). Non-metric multidimensional scaling ordination based on Bray-Curtis
dissimilarities at the ASV level for eukaryotic communities. Notes: The triple asterisks ***
represents the $P < 0.001$.

2.2. Dispersal limitation and hetero selection dominants the assembly of red sandstone surface microbial communities

139
140

To characterize the assembly mechanism of the red sandstone surface microbial communities,
we applied null model analysis[30,31] and dissimilarity-overlap curve (DOC) model analysis (Fig. 3)[32].

Null model analysis shows that both deterministic and stochastic processes controlled the
assembly of microbial communities, but the relative importance of the two processes in
community assembly vary among sample types (Fig. 3 a, b). In general, dispersal limitation
dominants the assembly of both prokaryotic communities (56.41%) and eukaryotic communities
(55.56%) on stone surface. Prokaryotes reveal overall elevated deterministic process than
eukaryotes. Hetero selection and dispersal limitation can explain about 91% of the assembly
mechanism of the prokaryotic communities. While hetero selection and dispersal limitation are
also main process for the assembly process of eukaryotic communities; drift plays a more pivotal
role in the assembly process of eukaryotic communities (20.23%) than in the prokaryotic ones
(5.41%). Prokaryotic samples reveal significantly higher value of betaNTI than eukaryotic ones (Fig.
3 c). No significant change of betaNTI is detected among any groups of prokaryotic samples
comparing with CK (Fig. 3 d). On the contrary, significant elevation of betaNTI is detected among
Black as well as Green groups of eukaryotic samples comparing with CK (Fig. 3 e).

In addition, the DOC model analysis demonstrates a significantly negative correlation
between overlap and dissimilarity in both prokaryotic and eukaryotic communities (P < 0.001),
which indicates a general dynamic and interspecific interaction within these microbial
communities (Fig. 3 f, g). These complex dynamic interactions among microorganisms in the
microbiota might lead to competition within the shared niche for survival.

Figure 3. Assembly dynamics of the red sandstone surface microbial communities. (a). The relative importance of assembly processes in prokaryotic communities. (b). The relative importance of assembly processes in eukaryotic communities. (c). The differences of betaNTI between prokaryotic and eukaryotic communities. (d). Changes of betaNTI among three groups of prokaryotic samples comparing with CK. (e). Changes of betaNTI among three groups of eukaryotic samples comparing with CK. (f). Dissimilarity-overlap curve (DOC) analysis of the universality of prokaryotic dynamics among samples. (g). DOC analysis of the universality of eukaryotic dynamics among samples.

2.3. Specialization prevails over flexibility in microbial community formation on red sandstone surfaces

To delve deeper into the dynamic interactions among microorganisms within the microbiota, we further investigated the niche characterization of microbial communities on red sandstone surfaces. According to Levins' niche breadth theory, the distribution of microbiota composition can be used as an indicator of the environmental conditions[33]. Here, we calculated niche breadth (*Bcom*) and niche overlap (*Ocom*) as indicators to describe the habitat niche characterization of both prokaryotes and eukaryotes making up the stone surface microbial communities. First, the *Bcom* was calculated with Levins' index (Fig. 4 a-c). The results show that the visually distinguishable microbial communities reveal decreased overall *Bcom* comparing with CK group. Such results were also further validated with Shannon index (Fig. S3 a-c). Then, the *Ocom* was calculated with Schoener index. The *Ocom* for both prokaryotes and eukaryotes displayed a similar

trend that CK samples had the highest *Ocom*, followed by Black, Green, and White, in descending
order. All pairwise comparisons were statistically significant, indicating distinct niche overlaps
among the different samples. When comparing *Ocom* across all samples, prokaryotes had a
significantly higher *Ocom* than eukaryotes (Fig. 4 d-f). Such results were also further validated with
Pianka index (Fig. S3 d-f). In summary, both prokaryotic and eukaryotic communities exhibit
significant variations in *Bcom* and *Ocom* across different sample types (CK, Black, Green, White).
In general, CK samples displayed widest niche breadth and highest overlap level compared with
the rest groups of samples. These results suggest that the formation of visually distinguishable
microbial communities on rock surfaces requires species to be more specialized, rather than
relying on diverse and flexible species with metabolic flexibility and redundancy.

To further explore the inference, we also calculated the proportion of generalists and
specialists among the microbial communities (Fig. 4 g-k). Generalists are species that are better
suited to survive in diverse and challenging environments and often play pivotal roles within their
communities. Specialists, on the other hand, have a more limited range of habitats and are
specifically adapted to environmental conditions[34,35]. The results show that at the ASVs level, the
proportion of specialists in each type of sample is higher than that of generalists in both
prokaryotes and eukaryotes (Fig. 4 g). To determine whether there are differences between all
prokaryotes and eukaryotes, we also compared the proportion of specialists and generalists within
these groups. The results show that in prokaryotes, the proportion of specialists is lower than that
of generalists. Conversely, in eukaryotes, the proportion of specialists is higher than that of
generalists (Fig. 4 g). At the phylum level, up to 70.22 % of the prokaryotes are classified as the
middle; the proportion of specialists (29.26 %) is higher than that of generalists (0.52 %) (Fig. 4 i,
Table S6). Myxococcota have the largest fraction of specialists in prokaryotes (50%) among the top
10 dominant phyla. (Fig. 4 h, Table S6). For the eukaryotes, 59.5% are classified as middle, 40.5%
as specialists, and none of the ASV is classified as generalists. (Fig. 4 k, Table S7). This suggests that
specialists prevail over generalists in the dominant phyla within the assembly of microbial
communities. Notably, the non-dominant phyla (others) of both prokaryotes and eukaryotes
exhibit the highest proportion of specialists (Fig. 4 h, j; Table S6, S7). This suggests that the non-
dominant phyla within the community might have even lower metabolic flexibility compared to
the dominant phyla.

Figure 4. Niche characterization of the red sandstone surface microbial communities. (a). Average habitat niche breadth (*Bcom*) differences among three groups of prokaryotic samples comparing with CK in prokaryotes. (b). *Bcom* differences among three groups of prokaryotic samples comparing with CK in eukaryotes. (c). *Bcom* differences between prokaryotes and eukaryotes. (d). Average habitat niche breadth (*Ocom*) differences among three groups of prokaryotic samples comparing with CK in prokaryotes. (e). Average *Ocom* differences among three groups of prokaryotic samples comparing with CK in eukaryotes. (f). Average *Ocom* differences between prokaryotes and eukaryotes. (g). Proportional distribution of specialists, middle, and generalists in each sample group, and between prokaryotes and eukaryotes. (h). Proportional distribution of

specialists, middle, and generalists of prokaryotes at phylum level based on Levins' niche breadth
index. (i). Overall proportion of the habitat niche breadth of prokaryotes at phylum level. (j).
Proportional distribution of specialists, middle, and generalists of eukaryotes at phylum level. (k).
Overall proportion of the habitat niche breadth of eukaryotes at phylum level. Notes: The *Bocm*
was calculated with Levins' index. The *Ocom* was calculated with Schoener index. All ASVs were
defined as specialists, middle or generalists according to Levins' niche breadth index. The asterisks
in the boxplots stand for the significance of the Wilcoxon signed-rank test. The "ns" stands for no
significance, single asterisk stands for P < 0.05, double asterisks stand for P < 0.01, Triple asterisks
stand for P < 0.001, Four asterisks stand for P < 0.0001.
235
236

2.4. Elevated network complexity and resilience to disturbances during the formation of visually
distinguishable microbial communities on red sandstone surface
Given the non-random assembly patterns observed in red sandstone surface microbial
communities, we proceeded to construct microbial co-occurrence networks of both prokaryotes
and eukaryotes in different types of microbial communities to further explore the interaction
patterns within these communities (Fig. 5).
Topological features analysis of the networks reveals that prokaryotic network showed higher
total nodes and links (Fig. 5 a). This indicates that the elevation of the complexity of the prokaryotic
network is necessary for the formation of visually distinguishable microbial communities. To
evaluate the resilience of the prokaryotic network to disturbances, we also analyzed the
robustness of the network with different approaches including targeted and randomly removal of
key species, removal of nodes, as well as the calculation of negative correlation ratio. The results
show that the prokaryotic network robustness decreases as both targeted and randomly removal
of the key species, with Black, Green, and White groups showing greater vulnerability compared
to CK (Fig. 5 b, c). The prokaryotic networks of the three visually distinguishable microbial
communities also have greater vulnerability to the removal of nodes than CK (Fig. 5 d). In addition,
the CK network also revealed higher negative correlation ratio than the other three networks of
visually distinguishable microbial communities, especially higher than both the Black and the
White samples (Fig. 5 e). This implies the formation of visually distinguishable microbial
communities could mitigate the mutual exclusion of microbial populations in resource competition.
In contrast, the topological features of eukaryotic networks differ from those of prokaryotic
networks. The number of links in the Green group decreased to half compared to the CK group,
while the number of links in the White group increased by approximately half (Fig. 5 f). The
network robustness analysis results indicate that only the White group demonstrated higher
stability across all test categories (Fig. 5 g-j). This suggests that the role of prokaryotes in
community formation might be more complex and variable depending on the morphology of
communities.
264

Figure 5. Topological features of the co-occurrence network within microbial communities on the stone surface. (a). Co-occurrence network of prokaryotic communities. (b). Robustness of prokaryotic networks under targeted removal of key species. (c). Robustness of prokaryotic networks under random removal of key species. (d). Vulnerability of prokaryotic networks under removal of key node. (e). Negative correlation ratio of the prokaryotic co-occurrence networks within each group. (f). Co-occurrence network of eukaryotic communities. (g). Robustness of eukaryotic networks under targeted removal of key species. (h). Robustness of eukaryotic networks under random removal of key species. (i). Vulnerability of eukaryotic networks under removal of key node. (j). Negative correlation ratio of the eukaryotic co-occurrence networks within each group.

2.5. Functions of the microbial communities are convergent despite taxonomic variations

To elucidate the metabolic processes that enable the survival of microbial communities on the stone surface, we performed metagenomic sequencing to each type of microbial communities. Ten Gene Ontology (GO) terms that could potentially contribute to the assembly of microbial communities on red sandstone surfaces were identified in all four types of microbial communities[3]. These terms are associated with functions related to organism interactions, microbial survival, and trophic processes. (Fig. 6 a). We then performed more detailed insight into nitrogen cycle processes because they have been broadly reported as closely related with the deterioration of stone[3,5,36-38]. The results showed that despite the variation in taxonomic structure, relative gene abundance within each nitrogen cycle pathway remained consistent across all four types of microbial communities (Fig. 6 b). Notably, the specific high-abundance genes regulating the four major nitrogen cycling processes (ammonium assimilation, nitrogen mineralization, ammonia assimilatory nitrate reduction, and dissimilatory nitrate reduction) might vary among different groups. However, each group consistently possesses high-abundance genes capable of regulating these respective nitrogen cycling processes. Afterwards, we analyzed the overall abundance distribution of the genes involving in the nitrogen cycle process (Fig. 6 c). The results showed that ammonium assimilation and nitrogen mineralization were the two dominant processes among the nitrogen cycle. The two processes possessed accounted for up to 38.52% and 31.39% of the overall abundance of reads within the nitrogen cycle pathways, respectively, which imply the intensive nitrogen-exchanging interactions between microbial communities and stone surface substrate. In addition, we performed Kyoto Encyclopedia of Genes and Genomes (KEGG) functional enrichment annotation of all the genes to gain further insights (Fig. S4 a). The result showed that stone surface microbial communities revealed activating metabolic processes in environmental information processing, cellular processes, and basic metabolism, *etc*. Notably, about 2.4 % of unigenes were annotated as human diseases-related functions, which suggests that the potential human health risks posed by these microorganisms on stone surfaces might worth a thorough evaluation.

Figure 6. Functional profiles of the stone surface microbiota. (a). GO functional annotation of the stone surface microbiota. (b). Gene abundance within each nitrogen cycle pathway of the stone surface microbiota. (c). Nitrogen cycling potential of the stone surface microbiota. Arrow thickness indicates the proportion of functional genes for corresponding metabolic processes.

To get further insights into the potential human health risks from the stone surface microbiota, we further analyzed the human diseases-related unigenes (Fig. S4 b, c; Table S8). The results showed that the 7763 unigenes belongs to 166 types of antimicrobial genes and bacterial infectious disease-related genes by the KEGG annotation (Fig S4 b). The most abundant genes include the *mrcA*, encoding penicillin-binding protein; *lepA*, encoding GTP-binding protein; and *namH*, encoding UDP-MurNAc hydroxylase, *etc*. In addition, we annotated the virulence factors within metagenome sequencing data using the Virulence Factor Database (VFDB). The results showed that there are in total 1,324,923 unigenes can be annotated as the virulence factors. These unigenes can be classified into 14 different categories including biofilm formation, motility, immune modulation, and notably, antimicrobial activities (Fig. S4 c). Similar to the genes in the nitrogen cycle pathway, each category of virulence factors is evenly distributed across the four different types of stone surface microbial communities.

Environmental antimicrobial resistance (AMR) genes poses a rising global threat, stemming from human activities and the widespread use of antimicrobials, leading to contamination and

proliferation of resistant bacteria and genes in nature[39]. On this basis, we then further annotated
the AMR genes within the stone surface microbiota using the Comprehensive Antibiotic Resistance
Database (CARD). The results showed that in our sequencing data, 26,862 unigenes were
annotated as 166 AMR genes, classified into 9 different resistance mechanisms (Fig. S5, Table S9).
These genes, especially the dominant genes with antibiotic efflux, antibiotic target alteration, and
antibiotic target protection mechanism are evenly distributed across the four different types of
red sandstone surface microbial communities. The presence of mobile genetic elements (MGEs) is
closely linked to the horizontal gene transfer of AMR genes among microbial communities[40-42].
Hence, we also performed a metagenome-wide search against the MGEs within the stone surface
microbiota using the Mobile Genetic Element Database. In total, 483 unigenes were annotated as
the MGEs. These unigenes can be further annotated as 57 genes, classified into 14 different types
such as transposase, integrase, insertion sequences and plasmids, *etc*. (Fig. S6, Table S10). The
most abundant gene, *tnpA*, encodes the only known transposase that facilitates both transposition
and target immunity[43,44]. This gene, similar to all the above-mentioned genes within the stone
surface microbiota, are also evenly distributed across all the four different types of microbial
communities.
342

3. Discussion

Uncovering the diversity, assembly mechanisms, and functional traits of microbial
communities is a central focus in microbial ecology[18,45,46]. Recently, the role of microbial
communities on stone surfaces has garnered increasing attention, particularly in the field of stone
cultural heritage conservation[47]. In this study, we present a pioneering case study on the ecological
roles of microbiota on red sandstone surfaces, using the Leshan Giant Buddha as an example. The
results indicate that despite the taxonomic composition of microbial communities varies according
to their coloration, their functions are convergent, particularly in terms of nitrogen cycle-related
biogeochemical processes. Dispersal limitation and heterogeneous selection dominate the
assembly of stone surface microbial communities. Habitat specialists play a more significant role
than generalists during the formation of microbial communities on rock surface. As visible
microbial communities form on stone surfaces, network complexity increases, and resistance to
disturbances is enhanced. By comparing our findings with previous studies on microbial
communities on stone surfaces, we inferred both commonalities and unique characteristics in the
ecological features of stone surface microbial communities.
358

3.1. Harsh habitant on stone surface might have facilitated the functional convergence of
microbial communities
In this study, the microbial diversity analysis shows that Proteobacteria, Actinobacteria,
Cyanobacteria, Chloroflexi, and Acidobacteriota dominate the top phyla among the prokaryotes in
all samples, while Ascomycota stands as the most dominant phylum in eukaryotes, despite the
distinct microbial community morphology across different sample groups. Such phylum-level
diversity patterns are largely consistent with previous reports on taxonomic diversity of microbial
communities on the surface of other stone types, but the diversity differ significantly at the genus
level[2,3,8,29,48-50]. Interestingly, in this study, despite the distinct taxonomic structures observed
among different sample types based on NMDS analysis, there is a high degree of functional

consistency across all sample groups. By comparing the results of studies on microbial communities
on the surfaces of several other types of stone, such characteristic of functional convergence
despite taxonomic divergence is widely observed across these surfaces. These microbial
communities all possess the capability to withstand pressures such as fluctuations between wet
and dry conditions, endure UV stress on stone surfaces, and facilitate ion exchange with the stone
substrate[3,5].

Previous study in limestone surface microbial communities reveals that nitrifying bacteria
genera *Nitrospira* and *Nitrosomonas* were significantly enriched in such habitant[3]. In another case
study involving regular sandstone, *Nitrosopumilus* and *Nitrospira* were also identified from
deteriorated sediments[5]. However, in our study, several dominant genera with ability to participate
in nitrogen cycling were also identified from the red sandstone microbial communities, yet their
taxonomy differs from that of the microorganisms in previously studies in other types of stones.
For example, in this study, a dominant prokaryotic genus, *Sphingomonas*, possesses the ability to
remove ammonia nitrogen and nitrite[51]. Additionally, species from the genus *Sphingomonas* are
involved in nitrification processes under low dissolved oxygen conditions[52]. Furthermore, another
dominant genus in our study, *Methylobacterium*, includes species known as plant growth-
promoting rhizobacteria, whose genomes contain enzymes involved in nitrogen metabolism and
ammonia assimilation[53]. Notably, the two dominated types of nitrogen cycling genes in limestone
surface microbial communities are nitrogen assimilation and mineralization, which is coherent with
what we found in red sandstone surface microbial communities in this study[3]. In summary, this
characteristic - functional convergence despite taxonomic divergence - might be widely applicable
among microbial communities inhabiting various types of stone surfaces.

Functional convergence despite taxonomic divergence is not unique to microbial communities
on stone surfaces. Several gut microbiota studies have reported functional, but not compositional,
convergence of gut microbiomes[54-57]. Some soil microbiota studies also reported the stable
functional structure despite high taxonomic variability[58-60]. In addition, microbial functional genes
of ocean microbiota that execute certain functions, such as B_{12} biosynthesis, are widely distributed,
while the microbial taxa that carry these genes vary significantly[61]. These results corroborate the
ecological hypothesis that environment truly selects for function, rather than taxonomy[62]. Different
taxonomic groups could reveal convergent functions as a result of functional redundancy among
microbial systems[63]. On stone surface, such functional convergence phenomenon might be
attributed to the selective pressure imposed by the harsh living conditions on the stone surface,
favoring the survival of microbial communities with specific functions[58,64-66]. This study suggests
that the interplay of deterministic and stochastic processes might have led to significant variations
in the taxonomic structure of the stone surface microbiota, which is consistent with previous cases
regarding crude oil microbial community assembly[66].

3.2. Assembly of stone surface microbial communities requires a general pattern of dynamic
interactions
In this study, we witnessed the extensive microbe-microbe and microbe-environment
interactions within the microbial communities on the stone surface. The mutual effect of hetero
selection and dispersal limitation controls the assembly process of both prokaryotic and eukaryotic
communities on red sandstone surfaces. Such results differ from previous studies on the assembly
process of microbial communities on limestone surfaces, where the assembly process is driven by

hetero selection and dispersal limitation[3]. This suggests that the assembly mechanisms of
microbial communities on stone surfaces might vary based on stone types, geographical locations,
climatic characteristics, or other environmental factors. In addition, the results of the DOC model
analysis suggest a general pattern of dynamic interspecific interactions within the communities on
the red sandstone surface and high selective pressure on the red sandstone surface[32,67-69]. Such
complexity was further testified in this study by the topological feature analysis of the microbial
networks. The active biogeochemical metabolism such as ammonium assimilation and nitrogen
mineralization processes by the microbial communities also suggests their active interactions with
the red sandstone substrate[70].

Here, we propose a hypothesis that a possible model capable of precisely interpreting the
assembly processes of stone surface microbial communities is the classic MacArthur consumer-
resource model, where species-species interactions are mediated through resource dynamics[71-73].
Because the nutrient-poor nature of stone surfaces, the colonization by pioneer species is
primarily driven by deterministic processes linked to nutrient availability, which aligns with
commonly observed among natural communities from various locations[74]. Some pioneer species
utilize inorganic energy from the stone to establish themselves. These species then leak metabolic
by-products into the environment, increasing the diversity of resources and enhancing the
biocompatibility of the stone surface. Other species can subsequently utilize these leaked
metabolites, in turn releasing new by-products into the environment, further enhancing the
biocompatibility of the stone surface, and facilitating the formation of microbial communities. The
observed decrease in ecological niche breadth among the visually distinguishable microbial
communities in this study also suggests that the utilization of specific resources is more efficient
within these communities. We also observed that specialization prevails over flexibility during the
formation of microbial communities on red sandstone surfaces. This is consistent with a previous
study on the assembly of microbial communities in a wide range of well-defined resource
environments, where habitat specialists also drive the increase of diversity in such microbial
communities[19].

3.3. Potential resources and bio-risks in stone surface microbial communities
The elimination of deterioration-causing microorganisms is one of the fundamental goals in
the conservation of stone cultural heritage. This study demonstrates that the microbial community
assembly on the surface of red sandstone is driven by dispersal limitation and hetero selection
processes. The harsh conditions on the stone surface create a challenging environment, exerting
strong selective pressure. Additionally, ecological modelling from this study reveals widely
negative correlation between species, all of which suggest that intense competition is occurring in
these microbial communities. One possible source of the negative correlation between species in
microbial communities is that certain species can gain a population advantage in competitions
during community formation by secreting antagonistic substances. Recently, there have been
numerous reports of metagenomic studies that explore new antagonistic substances as
antimicrobial resources from various habitats, including novel antibiotics, and antimicrobial
peptides, among others[75-77]. If machine learning and other methods can be used for in-depth
mining of metagenomic sequencing data from microbiota on red sandstone, it is possible to
develop novel antimicrobial agents for the removal of stone deterioration-causing microorganisms
in the future.

The "One Health" concept emphasizes the interconnectedness between human health and
environmental health[78]. Previous study provided a description of AMR genes in the microbial
communities on the surface of two stone cultural heritage sites[12,79], which indicates that stone
surfaces might be an overlooked ecosystem capable of harboring AMR genes as well as other
potentially risky factors for human health. In this study, an in-depth analysis of the potential
human health risks associated with microbial communities on stone surfaces was conducted. As
expected, virulence factors, AMR genes, and MGEs were observed from red sandstone surface.
We suggest that AMR genes could aid microorganisms in gaining an advantage in intense
competition within the microbial communities, both within and between species competition,
thereby promoting the enrichment of AMR genes within the stone surface microbial communities.
Human activities and the extensive use of antimicrobials might have facilitate this process, notably
through mechanisms like the water cycle[79-82]. Furthermore, the presence of MGEs on stone
surfaces indicates the potential for horizontal gene transfer of AMR genes among microbial
communities, potentially facilitating the dissemination and persistence of resistant
microorganisms on stone surface as in previous cases across various environments[83-85]. This
underscores the potential for stone surface microbial communities to be reservoirs for antibiotic-
resistant bacteria. Currently, natural environmental health assessments predominantly focus on
habitats including soil, water, and air, *etc*[86-88]. Although stone surfaces are also vital components
of ecosystems, research on whether these stone surface microbes pose health risks remains
limited. Our study suggests that more detailed assessments of the environmental health of stone
surfaces are necessary in future research.
478
3.4. Summary
In summary, this study focuses on microbial communities on the surface of red sandstones of
the Leshan Giant Buddha, which reveals their ecological and functional characteristics. Under the
interplay of dispersal limitation and hetero selection, the microbiota on the red sandstone were
assembles into distinct types of visually distinguishable microbial communities., significantly
differentiate the taxonomic composition of these communities. However, under the harsh habitant
on red sandstone surface, such as UV radiation, fluctuations between dry and wet conditions, and
limited nutrient availability, the functions of the microbe among different types of microbial
communities are convergent. These functions are particularly enriched in biogeochemical
processes related to nitrogen cycling and responses to environmental stress. The formation of
visually distinguishable microbial communities is associated with a narrower niche width, elevated
complexity, and increased resilience of the prokaryotic network to disturbances. Furthermore, a
higher proportion of habitat specialists are found in microbes forming these visually
distinguishable microbial communities. Given the nutrient limitations of the red sandstone surface,
the classic MacArthur consumer-resource model might provide a potential framework to further
explain the assembly mechanisms of these microbial communities. A significant accumulation of
AMRs and MGEs within these communities might help enhances their competitive advantage
during community formation, enabling them to resist antagonistic substances secreted by
competing organisms. The sources of these AMRs and MGEs might include water cycles and human
activities (Fig. 7). Future research into the ecological traits and functional potential of these
microbial communities will provide critical insights for managing microbial communities on red
sandstone surfaces. Moreover, microbial deterioration affects not only stone cultural heritage but

also a wide range of different cultural heritage materials[89-92]. The microbial ecology-based
approach used in this study will help advance research on microbial deterioration mechanisms in
other cultural heritage materials, providing a model for future work.

Figure 7. Schematic diagram illustrates the characteristics of microbial communities on the red
sandstone surface of the Leshan Giant Buddha

4. Materials and Methods

509 See detailed materials and methods in **Supplementary information 1**. The authors used
510 ChatGPT to check the English grammar during the writing of this manuscript. Authors are fully
511 responsible for the content of this manuscript, even those parts produced by ChatGPT, and are
512 thus liable for any breach of publication ethics.

Supplementary Files

Supplementary information 1. Supplementary Materials and Methods

Figure S1. The overall taxonomic community composition among all the samples.

Figure S2. Taxonomic community composition of the microbial communities in genus level.

Figure S3. Niche characterization of the stone surface microbial communities based on Shannon

niche breadth index and Pianka niche overlap index.

Figure S4. Human health risk potentials within the stone surface microbiota.

Figure S5. Antimicrobial resistance (AMR) genes within the stone surface microbiota.

Figure S6. The mobile genetic elements (MGEs) within the stone surface microbiota.

Table S1. Summaries for the details from the metagenome sequencing

Table S2. Relative abundances of dominant prokaryotes among 27 samples at the phylum level.

Table S3. Relative abundances of dominant prokaryotes among 27 samples at the genus level.

Table S4. Relative abundances of dominant eukaryotes among 27 samples at the phylum level.

Table S5. Relative abundances of dominant eukaryotes among 27 samples at the genus level.

Table S6. Habitat niche breadth of prokaryotes at the phylum level.

Table S7. Habitat niche breadth of eukaryotes at the phylum level.

Table S8. Human disease-related genes within the stone surface microbiota based on KEGG annotation.

Table S9. AMR genes within the stone surface microbiota.

Table S10. The mobile genetic elements within the stone surface microbiota.

References:

Liu, X. B., Koestler, R. J., Warscheid, T., Katayama, Y. & Gu, J. D. Microbial deterioration and sustainable conservation of stone monuments and buildings. *Nat Sustain* **3**, 991-1004 (2020). https://doi.org/10.1038/s41893-020-00602-5

Brewer, T. E. & Fierer, N. Tales from the tomb: the microbial ecology of exposed rock surfaces. *Environ Microbiol* **20**, 958-970 (2018). https://doi.org/10.1111/1462-2920.14024

He, J. *et al.* From surviving to thriving, the assembly processes of microbial communities in stone biodeterioration: A case study of the West Lake UNESCO World Heritage area in China. *Sci Total Environ* **805**, 150395 (2022). https://doi.org/10.1016/j.scitotenv.2021.150395

Li, J. *et al.* The active microbes and biochemical processes contributing to deterioration of Angkor sandstone monuments under the tropical climate in Cambodia – A review. *J Cult Herit* **47**, 218-226 (2021). https://doi.org/https://doi.org/10.1016/j.culher.2020.10.010

Ding, X. *et al.* Microbiome and nitrate removal processes by microorganisms on the ancient Preah Vihear temple of Cambodia revealed by metagenomics and N-15 isotope analyses. *Appl Microbiol Biotechnol* **104**, 9823-9837 (2020). https://doi.org/10.1007/s00253-020-10886-4

Wu, F. *et al.* Metagenomic and metaproteomic insights into the microbiome and the key geobiochemical potentials on the sandstone of rock-hewn Beishiku Temple in Northwest China. *Sci Total Environ* **893**, 164616 (2023). https://doi.org/https://doi.org/10.1016/j.scitotenv.2023.164616

Fidanza, M. R. & Caneva, G. Natural biocides for the conservation of stone cultural heritage: A review. *J Cult Herit* **38**, 271-286 (2019). https://doi.org/https://doi.org/10.1016/j.culher.2019.01.005

Qiang, L., Zhang, B. B., Yang, B. X. & Ged, C. Q. Deterioration-Associated Microbiome of Stone Monuments: Structure, Variation, and Assembly. *Appl Environ Microbiol* **84**, e02680-02617-02617 (2018).

Li, T., Hu, Y. & Zhang, B. Biomineralization Induced by Colletotrichum acutatum: A Potential Strategy for Cultural Relic Bioprotection. *Front Microbiol* **9**, 1884 (2018).

Jroundi, F., Gonzalez-Muñoz, M. T. & Rodriguez-Navarro, C. in *Microorganisms in the Deterioration and Preservation of Cultural Heritage* (ed Edith Joseph) 281-299 (Springer

International Publishing, 2021).

11 Liu, X. *et al.* Biofilms on stone monuments: biodeterioration or bioprotection? *Trends Microbiol*
30, 816-819 (2022). https://doi.org/https://doi.org/10.1016/j.tim.2022.05.012

12 Li, Q., Wu, C., He, J. & Zhang, B. Unraveling the microbiotas and key genetic contexts identified
on stone heritage using illumina and nanopore sequencing platforms. *Int Biodeter Biodegr* **185**,
105688 (2023). https://doi.org/https://doi.org/10.1016/j.ibiod.2023.105688

13 Ding, X. *et al.* Microbiome characteristics and the key biochemical reactions identified on stone
world cultural heritage under different climate conditions. *Journal of Environmental*
Management   **302**,   114041   (2022).
https://doi.org/https://doi.org/10.1016/j.jenvman.2021.114041

14 Ennis, N. J., Dharumaduri, D., Bryce, J. G. & Tisa, L. S. Metagenome Across a Geochemical
Gradient of Indian Stone Ruins Found at Historic Sites in Tamil Nadu, India. *Microb Ecol* **81**, 385-
395 (2021). https://doi.org/10.1007/s00248-020-01598-3

15 Louati, M. *et al.* Elucidating the ecological networks in stone-dwelling microbiomes. *Environ*
Microbiol **22**, 1467-1480 (2020). https://doi.org/https://doi.org/10.1111/1462-2920.14700

16 Li, T., Cai, Y. & Ma, Q. Microbial Diversity on the Surface of Historical Monuments in Lingyan
Temple, Jinan, China. *Microb Ecol* (2022). https://doi.org/10.1007/s00248-021-01955-w

17 Cattò, C. *et al.* Biofilm colonization of stone materials from an Australian outdoor sculpture:
Importance of geometry and exposure. *Journal of Environmental Management* **339**, 117948
(2023). https://doi.org/https://doi.org/10.1016/j.jenvman.2023.117948

18 Zhou, J. & Ning, D. Stochastic Community Assembly: Does It Matter in Microbial Ecology?
MMBR **81**, e00002-00017 (2017). https://doi.org/10.1128/MMBR.00002-17

19 Dal Bello, M., Lee, H., Goyal, A. & Gore, J. Resource–diversity relationships in bacterial
communities reflect the network structure of microbial metabolism. *Nature Ecology &*
Evolution **5**, 1424-1434 (2021). https://doi.org/10.1038/s41559-021-01535-8

20 Goldford, J. E. *et al.* Emergent simplicity in microbial community assembly. *Science* **361**, 469-
474 (2018). https://doi.org/10.1126/science.aat1168

21 Ning, D. *et al.* A quantitative framework reveals ecological drivers of grassland microbial
community assembly in response to warming. *Nat Commun* **11**, 4717 (2020).
https://doi.org/10.1038/s41467-020-18560-z

22 Chase, J. M. Stochastic Community Assembly Causes Higher Biodiversity in More Productive
Environments. *Science* **328**, 1388-1391 (2010). https://doi.org/10.1126/science.1187820

23 Ofiţeru, I. D. *et al.* Combined niche and neutral effects in a microbial wastewater treatment
community. *PNAS* **107**, 15345-15350 (2010). https://doi.org/10.1073/pnas.1000604107

24 Chesson, P. Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology,*
Evolution,   and   Systematics **31**,   343-366   (2000).
https://doi.org/https://doi.org/10.1146/annurev.ecolsys.31.1.343

25 Fargione, J., Brown, C. S. & Tilman, D. Community assembly and invasion: An experimental test
of   neutral   versus   niche   processes. *PNAS* **100**,   8916-8920   (2003).
https://doi.org/10.1073/pnas.1033107100

26 Chave, J. Neutral theory and community ecology. *Ecol Lett* **7**, 241-253 (2004).
https://doi.org/https://doi.org/10.1111/j.1461-0248.2003.00566.x

27 Vega, N. M. & Gore, J. Stochastic assembly produces heterogeneous communities in the
Caenorhabditis   elegans   intestine. *PLOS   Biology* **15**,   e2000633   (2017).

https://doi.org/10.1371/journal.pbio.2000633

28 Bai, F. *et al.* Microbial biofilms on a giant monolithic statue of Buddha: The symbiosis of
microorganisms and mosses and implications for bioweathering. *Int Biodeter Biodegr* **156**
(2021). https://doi.org/ARTN 105106 10.1016/j.ibiod.2020.105106

29 Zhu, C. *et al.* Analysis of the Microbiomes on Two Cultural Heritage Sites. *Geomicrobiol J*, 1-10
(2022). https://doi.org/10.1080/01490451.2022.2137604

30 Stegen, J. C., Lin, X., Konopka, A. E. & Fredrickson, J. K. Stochastic and deterministic assembly
processes in subsurface microbial communities. *ISME J* **6**, 1653-1664 (2012).
https://doi.org/10.1038/ismej.2012.22

31 Stegen, J. C. *et al.* Quantifying community assembly processes and identifying features that
impose them. *ISME J* **7**, 2069-2079 (2013). https://doi.org/10.1038/ismej.2013.93

32 Bashan, A. *et al.* Universality of human microbial dynamics. *Nature* **534**, 259-262 (2016).
https://doi.org/10.1038/nature18301

33 Levins, R. *Evolution in Changing Environments Some Theoretical Explorations. (MPB-2).*
(Princeton University Press, 1968).

34 Sexton, J. P., Montiel, J., Shay, J. E., Stephens, M. R. & Slatyer, R. A. Evolution of Ecological Niche
Breadth. *Annual Review of Ecology, Evolution, and Systematics* **48**, 183-206 (2017).
https://doi.org/10.1146/annurev-ecolsys-110316-023003

35 von Meijenfeldt, F. A. B., Hogeweg, P. & Dutilh, B. E. A social niche breadth score reveals niche
range strategies of generalists and specialists. *Nature Ecology & Evolution* **7**, 768-781 (2023).
https://doi.org/10.1038/s41559-023-02027-7

36 Meng, H., Zhang, X., Katayama, Y., Ge, Q. & Gu, J.-D. Microbial diversity and composition of the
Preah Vihear temple in Cambodia by high-throughput sequencing based on genomic DNA and
RNA. *Int Biodeter Biodegr* **149** (2020).

37 Meng, S. *et al.* Community structures and biodeterioration processes of epilithic biofilms imply
the significance of micro-environments. *Sci Total Environ* **876**, 162665 (2023).
https://doi.org/https://doi.org/10.1016/j.scitotenv.2023.162665

38 Chen, X. *et al.* The Organisms on Rock Cultural Heritages: Growth and Weathering. *Geoheritage*
13 (2021). https://doi.org/ARTN 5610.1007/s12371-021-00588-2

39 Larsson, D. G. J. & Flach, C.-F. Antibiotic resistance in the environment. *Nat Rev Microbiol* **20**,
257-269 (2022). https://doi.org/10.1038/s41579-021-00649-x

40 Guo, J., Li, J., Chen, H., Bond, P. L. & Yuan, Z. Metagenomic analysis reveals wastewater
treatment plants as hotspots of antibiotic resistance genes and mobile genetic elements.
Water Res **123**, 468-478 (2017). https://doi.org/https://doi.org/10.1016/j.watres.2017.07.002

41 Baker, M. *et al.* Machine learning and metagenomics reveal shared antimicrobial resistance
profiles across multiple chicken farms and abattoirs in China. *Nature Food* **4**, 707-720 (2023).
https://doi.org/10.1038/s43016-023-00814-w

42 Arnold, B. J., Huang, I. T. & Hanage, W. P. Horizontal gene transfer and adaptive evolution in
bacteria. *Nat Rev Microbiol* **20**, 206-218 (2022). https://doi.org/10.1038/s41579-021-00650-4

43 Nicolas, E. *et al.* The Tn3-family of Replicative Transposons. *Microbiology Spectrum* **3**,
10.1128/microbiolspec.mdna1123-0060-2014   (2015).
https://doi.org/10.1128/microbiolspec.mdna3-0060-2014

44 Shkumatov, A. V. *et al.* Structural insight into Tn3 family transposition mechanism. *Nat Commun*
13, 6155 (2022). https://doi.org/10.1038/s41467-022-33871-z

45   Escalas, A. *et al.* Microbial functional diversity: From concepts to applications. *Ecology and*
Evolution **9**, 12000-12016 (2019). https://doi.org/https://doi.org/10.1002/ece3.5670

46   Prosser, J. I. *et al.* The role of ecological theory in microbial ecology. *Nat Rev Microbiol* **5**, 384-
392 (2007). https://doi.org/10.1038/nrmicro1643

47   Sterflinger, K. *et al.* Future directions and challenges in biodeterioration research on historic
materials and cultural properties. *Int Biodeter Biodegr* **129**, 10-12 (2018).
https://doi.org/https://doi.org/10.1016/j.ibiod.2017.12.007

48   Wu, Y. *et al.* Monitoring the Deterioration of Masonry Relics at a UNESCO World Heritage Site.
KSCE Journal of Civil Engineering **25**, 3097-3106 (2021). https://doi.org/10.1007/s12205-021-
1716-z

49   Zhang, X. W., Ge, Q. Y., Zhu, Z. B., Deng, Y. M. & Gu, J. D. Microbiological community of the
Royal Palace in Angkor Thom and Beng Mealea of Cambodia by Illumina sequencing based on
16S rRNA gene. *Int Biodeter Biodegr* **134**, 127-135 (2018).
https://doi.org/10.1016/j.ibiod.2018.06.018

50   Schröer, L., De Kock, T., Cnudde, V. & Boon, N. Differential colonization of microbial
communities inhabiting Lede stone in the urban and rural environment. *Sci Total Environ* **733**,
139339 (2020). https://doi.org/https://doi.org/10.1016/j.scitotenv.2020.139339

51   Yun, L. *et al.* Ammonia nitrogen and nitrite removal by a heterotrophic Sphingomonas sp. strain
LPN080 and its potential application in aquaculture. *Aquaculture* **500**, 477-484 (2019).
https://doi.org/https://doi.org/10.1016/j.aquaculture.2018.10.054

52   Fitzgerald, C. M., Camejo, P., Oshlag, J. Z. & Noguera, D. R. Ammonia-oxidizing microbial
communities in reactors with efficient nitrification at low-dissolved oxygen. *Water Res* **70**, 38-
51 (2015). https://doi.org/https://doi.org/10.1016/j.watres.2014.11.041

53   Grossi, C. E. M. *et al.* Methylobacterium sp. 2A Is a Plant Growth-Promoting Rhizobacteria That
Has the Potential to Improve Potato Crop Yield Under Adverse Conditions. *Frontiers in Plant*
Science **11** (2020). https://doi.org/10.3389/fpls.2020.00071

54   Tian, L. *et al.* Deciphering functional redundancy in the human microbiome. *Nat Commun* **11**,
6217 (2020). https://doi.org/10.1038/s41467-020-19940-1

55   Li, H. *et al.* Food provisioning results in functional, but not compositional, convergence of the
gut microbiomes of two wild Rhinopithecus species: Evidence of functional redundancy in the
gut microbiome. *Sci Total Environ* **858**, 159957 (2023).
https://doi.org/https://doi.org/10.1016/j.scitotenv.2022.159957

56   Moya, A. & Ferrer, M. Functional Redundancy-Induced Stability of Gut Microbiota Subjected to
Disturbance. *Trends Microbiol* **24**, 402-413 (2016). https://doi.org/10.1016/j.tim.2016.02.002

57   Tian, L. & Zhang, J. Dynamics of Rabies Epidemics in Vampire Bats. *Complexity* **2020**, 7032451
(2020). https://doi.org/https://doi.org/10.1155/2020/7032451

58   Talbot, J. M. *et al.* Endemism and functional convergence across the North American soil
mycobiome. *PNAS* **111**, 6341-6346 (2014). https://doi.org/10.1073/pnas.1402584111

59   Zeng, Q. *et al.* Stable functional structure despite high taxonomic variability across fungal
communities in soils of old-growth montane forests. *Microbiome* **11**, 217 (2023).
https://doi.org/10.1186/s40168-023-01650-7

60   Chen, H. *et al.* Functional Redundancy in Soil Microbial Community Based on Metagenomics
Across the Globe. *Front Microbiol* **13** (2022).

61   Zhou, J. *et al.* The diversity and ecological significance of microbial traits potentially involved in

B12 biosynthesis in the global ocean. *mLife* **2**, 416-427 (2023).
https://doi.org/https://doi.org/10.1002/mlf2.12095

62   Burke, C., Steinberg, P., Rusch, D., Kjelleberg, S. & Thomas, T. Bacterial community assembly
based on functional genes rather than species. *PNAS* **108**, 14288-14293 (2011).
https://doi.org/10.1073/pnas.1101591108

63   Louca, S. *et al.* Function and functional redundancy in microbial systems. *Nature Ecology &*
Evolution **2**, 936-943 (2018). https://doi.org/10.1038/s41559-018-0519-1

64   Chikina, M., Robinson, J. D. & Clark, N. L. Hundreds of Genes Experienced Convergent Shifts in
Selective Pressure in Marine Mammals. *Molecular Biology and Evolution* **33**, 2182-2192 (2016).
https://doi.org/10.1093/molbev/msw112

65   Winemiller, K. O., Fitzgerald, D. B., Bower, L. M. & Pianka, E. R. Functional traits, convergent
evolution, and periodic tables of niches. *Ecol Lett* **18**, 737-751 (2015).
https://doi.org/https://doi.org/10.1111/ele.12462

66   Nie, Y. *et al.* Species Divergence vs. Functional Convergence Characterizes Crude Oil Microbial
Community Assembly. *Front Microbiol* **7** (2016).

67   Li, P. *et al.* Global diversity and biogeography of potential phytopathogenic fungi in a changing
world. *Nat Commun* **14**, 6482 (2023). https://doi.org/10.1038/s41467-023-42142-4

68   Gao, C. *et al.* Fungal community assembly in drought-stressed sorghum shows stochasticity,
selection, and universal ecological dynamics. *Nat Commun* **11**, 34 (2020).
https://doi.org/10.1038/s41467-019-13913-9

69   Wang, B., Zhu, C., Hu, Y., Zhang, B. & Wang, J. Dynamics of microbial community composition
during degradation of silks in burial environment. *Sci Total Environ* **883**, 163694 (2023).
https://doi.org/https://doi.org/10.1016/j.scitotenv.2023.163694

70   Meng, H., Luo, L., Chan, H. W., Katayama, Y. & Gu, J.-D. Higher diversity and abundance of
ammonia-oxidizing archaea than bacteria detected at the Bayon Temple of Angkor Thom in
Cambodia. *Int Biodeter Biodegr* **115**, 234-243 (2016).
https://doi.org/https://doi.org/10.1016/j.ibiod.2016.08.021

71   MacArthur, R. Species packing and competitive equilibrium for many species. *Theoretical*
Population Biology **1**, 1-11 (1970). https://doi.org/https://doi.org/10.1016/0040-
5809(70)90039-0

72   Advani, M., Bunin, G. & Mehta, P. Statistical physics of community ecology: a cavity solution to
MacArthur's consumer resource model. *Journal of Statistical Mechanics: Theory and*
Experiment **2018**, 033406 (2018). https://doi.org/10.1088/1742-5468/aab04e

73   Marsland, R., III *et al.* Available energy fluxes drive a transition in the diversity, stability, and
functional structure of microbial communities. *PLOS Computational Biology* **15**, e1006793
(2019). https://doi.org/10.1371/journal.pcbi.1006793

74   Sriswasdi, S., Yang, C.-c. & Iwasaki, W. Generalist species drive microbial dispersion and
evolution. *Nat Commun* **8**, 1162 (2017). https://doi.org/10.1038/s41467-017-01265-1

75   Wang, Z. *et al.* A naturally inspired antibiotic to target multidrug-resistant pathogens. *Nature*
601, 606-611 (2022). https://doi.org/10.1038/s41586-021-04264-x

76   Dulya, O. *et al.* A trait-based ecological perspective on the soil microbial antibiotic-related
genetic machinery. *Environment International* **190**, 108917 (2024).
https://doi.org/https://doi.org/10.1016/j.envint.2024.108917

77   Santos-Júnior, C. D. *et al.* Discovery of antimicrobial peptides in the global microbiome with

machine learning. *Cell* **187**, 3761-3778.e3716 (2024). https://doi.org/10.1016/j.cell.2024.05.013

78 Promoting the science of One Health. *Nat Commun* **14**, 4735 (2023). https://doi.org/10.1038/s41467-023-40293-y

79 He, J., Zhang, N., Shen, X., Muhammad, A. & Shao, Y. Deciphering environmental resistome and mobilome risks on the stone monument: A reservoir of antimicrobial resistance genes. *Sci Total Environ* **838**, 156443 (2022). https://doi.org/https://doi.org/10.1016/j.scitotenv.2022.156443

80 Guo, Z.-F. *et al.* Data-driven discoveries on widespread contamination of freshwater reservoirs by dominant antibiotic resistance genes. *Water Res* **229**, 119466 (2023). https://doi.org/https://doi.org/10.1016/j.watres.2022.119466

81 Balaban, N. Q. *et al.* Definitions and guidelines for research on antibiotic persistence. *Nat Rev Microbiol* **17**, 441-448 (2019). https://doi.org/10.1038/s41579-019-0196-3

82 Lin, Z.-J. *et al.* Behavior of antibiotic resistance genes in a wastewater treatment plant with different upgrading processes. *Sci Total Environ* **771**, 144814 (2021). https://doi.org/https://doi.org/10.1016/j.scitotenv.2020.144814

83 Smalla, K., Cook, K., Djordjevic, S. P., Klümper, U. & Gillings, M. Environmental dimensions of antibiotic resistance: assessment of basic science gaps. *FEMS Microbiol Ecol* **94**, fiy195 (2018). https://doi.org/10.1093/femsec/fiy195

84 Ellabaan, M. M. H., Munck, C., Porse, A., Imamovic, L. & Sommer, M. O. A. Forecasting the dissemination of antibiotic resistance genes across bacterial genomes. *Nat Commun* **12**, 2435 (2021). https://doi.org/10.1038/s41467-021-22757-1

85 Ezugworie, F. N., Igbokwe, V. C. & Onwosi, C. O. Proliferation of antibiotic-resistant microorganisms and associated genes during composting: An overview of the potential impacts on public health, management and future. *Sci Total Environ* **784**, 147191 (2021). https://doi.org/https://doi.org/10.1016/j.scitotenv.2021.147191

86 Banerjee, S. & van der Heijden, M. G. A. Soil microbiomes and one health. *Nat Rev Microbiol* **21**, 6-20 (2023). https://doi.org/10.1038/s41579-022-00779-w

87 Hernando-Amado, S., Coque, T. M., Baquero, F. & Martínez, J. L. Defining and combating antibiotic resistance from One Health and Global Health perspectives. *Nature Microbiology* **4**, 1432-1442 (2019). https://doi.org/10.1038/s41564-019-0503-9

88 Li, S. *et al.* Linking biodiversity and ecological function through extensive microeukaryotic movement across different habitats in six urban parks. *iMeta* **2**, e103 (2023). https://doi.org/https://doi.org/10.1002/imt2.103

89 Wang, B., Qi, M., Ma, Y., Zhang, B. & Hu, Y. Microbiome Diversity and Cellulose Decomposition Processes by Microorganisms on the Ancient Wooden Seawall of Qiantang River of Hangzhou, China. *Microb Ecol* **86**, 2109-2119 (2023). https://doi.org/10.1007/s00248-023-02221-x

90 Wang, B., Zhu, C., Wang, B., Zhang, B. & Hu, Y. Analysis of the biocorrosion community from ancient wooden constructions at Tianluoshan (7000–6300 cal BP), Zhejiang Province, China. *Heritage Science* **12**, 189 (2024). https://doi.org/10.1186/s40494-024-01304-3

91 Zhu, C. *et al.* Application and evaluation of a new blend of biocides for biological control on cultural heritages. *Int Biodeter Biodegr* **178**, 105569 (2023). https://doi.org/https://doi.org/10.1016/j.ibiod.2023.105569

92 Li, J. *et al.* Changes in soil microbial communities at Jinsha earthen site are associated with earthen site deterioration. *BMC Microbiology* **20**, 147 (2020). https://doi.org/10.1186/s12866-

785   [020-01836-1](020-01836-1)

786