Stimulus history, not expectation, drives sensory prediction errors in mammalian cortex

Jacob A. Westerberg[1,2,,*], Yihan S. Xiong[1,*], Hamed Nejat[1], Eli Sennesh[1], Séverine Durand[3], Hannah Cabasco[3], Hannah Belski[3], Ryan Gillis[3], Henry Loeffler[3], Ahad Bawany[3], Carter R. Peene[3], Warren Han[3], Katrina Nguyen[3], Vivian Ha[3], Tye Johnson[3], Conor Grasso[3], Ben Hardcastle[3], Ahrial Young[3], Jackie Swapp[3], Ben Ouellete[3], Shiella Caldejon[3], Ali Williford[3], Peter A. Groblewski[3], Shawn R. Olsen[3], Carly Kiselycznyk[3], Jerome A. Lecoq[3], Alexander Maier[1], and André M. Bastos[1,†]

[1]*Department of Psychology, Vanderbilt Brain Institute, Vanderbilt Vision Research Center, Vanderbilt University, Nashville, Tennessee, United States*
[2]*Department of Vision and Cognition, Netherlands Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, Amsterdam, The Netherlands*
[3]*Allen Institute for Brain Science, Seattle, Washington, United States.*

[†]Correspondence: andre.bastos@vanderbilt.edu
[*]These authors contributed equally

ORCiD:
J.A.W. - 0000-0001-5331-8707
Y.S.X. - 0009-0007-1298-3672
H.N.   - 0009-0003-5963-5801
E.S.   - 0000-0001-7014-8471
S.D.   - 0000-0001-7788-0200
H.C.   - 0009-0001-7899-2463
H.B.   - 0009-0001-5422-5964
H.L.   - 0009-0007-3935-6863
A.B.   - 0009-0002-5495-5411
C.R.P. - 0009-0000-6660-2264
W.H.   - 0009-0006-0104-0380
K.N.   - 0009-0009-8547-9703
V.H.   - 0009-0005-0093-3825
T.J.   - 0009-0006-7239-7571
C.G.   - 0009-0003-6384-4075
B.H.   - 0000-0002-4819-5631
J.S.   - 0009-0008-4965-6242
R.G.   - 0009-0000-1879-6519
B.O.   - 0000-0003-4913-554X
A.W.   - 0009-0000-7282-7515
P.A.G. - 0000-0002-8415-1118
S.R.O. - 0000-0002-9568-7057
C.K.   - 0009-0006-8410-1192
J.A.L. - 0000-0002-0131-0938
A.M.   - 0000-0002-7250-502X
A.M.B. - 0000-0003-1804-4418

Summary

Predictive coding (PC) is a popular framework to explain cortical responses. PC states that the brain computes internal models of expected events and responds robustly to unexpected stimuli with prediction errors that feed forward. This can be studied using oddball tasks. A repetitive sequence interrupted by a novel stimulus is a "local" oddball. "Global" oddballs defy expectations while repeating the local context. This dissociates expectation-driven neuronal prediction error responses vs. merely stimulus-history-driven release from adaptation. We recorded neuron spiking activity across the visual hierarchy in mice and monkeys viewing these oddballs. Local oddball responses largely followed PC hypotheses. They were robust, emerged early in layers 2/3, and fed forward. Global oddball responses challenged PC. They were weak, absent in most visual areas, more robust in prefrontal cortex, and emerged in non-granular layers. Contrary to the most popular PC models that emphasize prediction error computations occurring in early sensory areas and feeding forward, this implies that genuine predictive coding computations take place in more cognitive, higher-order areas.

Keywords

Cognition, Expectation, Macaque, Mouse, Oddball, Optogenetics, Perception, Predictive Coding, Visual Cortex

Highlights

- Core principles of Predictive Coding hold for local, but not global oddballs
- Global oddballs are sparsely encoded in visual cortex and do not engage feedforward mechanisms
- Parvalbumin and somatostatin interneurons do not support sensory global oddball detection
- Global oddballs are earliest and most robustly encoded in prefrontal brain areas

Significance

Global oddballs defy our expectations without defying their local context. Whether they are encoded in sensory brain areas was uncertain. Through multiarea neurophysiological recordings in mice and monkeys, Westerberg et al. show that these oddballs are sparsely encoded in sensory cortex, but their mechanism does not emerge from classical Predictive Coding.

Introduction

Expectations shape our perception and it is often immediately recognizable when our predictions are violated (Itti and Baldi, 2009). Predictive coding (PC) has emerged as one of the most popular theoretical frameworks to explain how predictions are instantiated in the brain (Friston, 2010; Rao and Ballard, 1999). One likely reason for the appeal of PC is its wide explanatory power despite its simplicity: less than a handful of theoretical constructs (prior probabilities/predictions, incoming data, prediction errors) can account for a large host of neurophysiological and psychological phenomena, ranging from the mismatch negativity (Garrido et al., 2007), sensory processing (Schwiedrzik and Freiwald, 2017), and adaptation (Hosoya et al., 2005) to attentional modulation (Brown and Friston, 2013; Kok et al., 2011; Smout et al., 2019; Spratling, 2008). It has even been suggested the PC can explain conscious experience (Seth et al., 2012).

Predictive coding (PC) models make a series of hypotheses about the nature and underlying mechanisms of neuronal processing (Bastos et al., 2012; Friston, 2005; Friston et al., 2017; Keller and Mrsic-Flogel, 2018; Rao and Ballard, 1999; Schultz et al., 1997; Spratling, 2010; Srinivasan et al., 1982). PC (schematized in Figure 1A) proposes that the brain compares incoming sensory information against a prediction signal. This prediction is based on an internal model about the statistics of the environment that is generated in higher-order cortex; these statistics are typically taken to include the mean expected stimulus (typically represented mathematically as μ) and the precision of the expected stimulus (typically represented mathematically as Π). Predictions are transmitted to sensory areas, and if an incoming sensory signal matches the prediction, the two signals cancel. Thus, expected sensory data are "explained away" and leave the brain unexcited. In other words, PC is subtractive and multiplicative: when sensory signals do not match the prediction, subtraction results in a larger value, called the prediction error, which is weighted by the precision. This error signal is computed in superficial layers 2 and 3 of sensory cortex, which propagate the signal feed forward up the cortical hierarchy (Bastos et al., 2012). Being a second-order statistic, the precision Π is always a positive, nonzero quantity, ensuring that prediction errors are always fed up the cortical hierarchy, even if attenuated. As they ascend, prediction errors instigate updates to the internal model to improve predictions. Associated prediction update signals flow back down the hierarchy.

PC has also been proposed to occur over a series of computational steps, each of which compares predictions to prediction errors from a previous step (Rao and Ballard, 1999). This "hierarchical predictive coding model" has since been further adjusted, expanded, and modified to fit both the structural and functional architecture of the neocortex (Bastos et al., 2012; Friston, 2010) (Figure 1A). These models efficiently predict and explain the cortical responses to first order pattern violations ("local oddballs"), such as a change from repeated stimulation (e.g., a stimulus sequence such as x-x-x-Y) (Chao et al., 2018; Gallimore et al., 2023; Xiong et al., 2023). Such local oddball responses depend on a simplified predictive model that the current stimulus will continue to repeat. However, local oddball responses persist during states of deep anesthesia when feedback connections are functionally weak (Nourski et al., 2018; Uhrig et al., 2016; Xiong et al., 2023). As such, they by themselves do not provide strong evidence for PC, which envisions rich and multi-faceted predictions feeding back and modulating sensory processing. It is an open question whether sensory cortex is engaged by more complex prediction errors (Gabhart et al., 2024). More complex prediction error computations and higher-order predictions can be studied using "global oddballs". These second-order pattern violations occur when a repeated presentation of first order pattern violations is interrupted (e.g., x-x-x-Y / x-x-x-Y / x-x-x-Y / x-x-x-<u>X</u>). Global oddballs are themselves a repeated stimulus, so responses cannot be explained by recent stimulus history, i.e., release from adaptation.

Two prior studies failed to find global oddball encoding in the spiking activity in sensory cortex (areas V1 and V4 of macaques (Solomon et al., 2021) and auditory area Tpt (Xiong et al., 2023)). In addition, two other studies reported neuronal signatures of global oddballs in higher-order prefrontal cortex (Bellet et al., 2024; Xiong et al., 2023), which were dependent on the conscious state (Xiong et al., 2023). However, these studies did not include an extensive period of habituation to the stimulus, which may be required for the brain to learn an internal

model that can drive predictive coding in sensory cortex. These prior studies also did not sample widely across cortex to determine at what stage of cortical processing hierarchical predictive coding emerges. Here, we tested core tenets of the hierarchical predictive coding model using Multi-Area, high-Density, Laminar-resolved Neurophysiological (MaDeLaNe) recordings of spiking activity in mice and monkeys. We sampled across multiple stages of the visual cortical processing pathway, ranging from primary visual cortex to high-level visual and prefrontal areas. Using optogenetic "photo tagging" in mice, we tested whether specific inhibitory cell types (Parvalbumin- and Somatostatin-positive interneurons) were involved in oddball processing.

These techniques allowed us to test three core hypotheses postulated by hierarchical predictive coding theories (Figure 1A, Bastos et al., 2012; Friston, 2010; Rao and Ballard, 1999). These hypotheses pertain to the proposed canonical microcircuit for prediction computations. We tested these hypotheses for both local and global oddballs. The first hypothesis (H1) tests the proposal that prediction error computations are a feedforward cortical mechanism. If so, then prediction errors should (a) be encoded early in time in the sensory response (within the first 50-150 ms of processing), (b) emerge early in the sensory cortical processing hierarchy and then subsequently in higher areas, (c) evoke activity in pyramidal neurons superficial layers (L2/3) of cortex, as feedforward projections anatomically project from these neurons, and (d) evoke a feedforward pattern of directed functional connectivity between neurons, with earlier areas of the hierarchy exerting greater influence on later areas than vice-versa. The second hypothesis (H2) is that if prediction error is a core cortical computation, neuronal oddball processing should be robust and widespread (observed widely in different cortical areas and species). The third hypothesis (H3) is that if predictions are subtractive, they should involve the activation of inhibitory interneuron cell populations, and a release from inhibition during prediction errors.

Consistent with feedforward prediction error computations, we found that local oddballs were first detected in primary visual cortex (V1) followed by higher areas, were initiated by L2/3 neurons, and recruited a substantial neuronal population in each area. In contrast, global oddball sensory prediction error responses were absent in most visual areas. They evoked a later, weaker, and much less widespread neuronal response. Superficial L2/3 did not preferentially encode global oddballs. We tested the hypothesis that prediction errors would feed forward by analyzing neuronal spiking time courses with Granger causality (Dhamala et al., 2008). We did not find a feedforward pattern of activation evoked by global oddballs, but did for local oddballs. Neither global nor local oddballs involved a release from predictive suppression mediated by PV+/SST+ inhibitory interneurons. These results indicate that while local oddballs follow many of the principles proposed by predictive coding, global oddballs do not. These results challenge the core hypothesis that cortical neurons are engaged by complex predictive coding computations in a widespread fashion and with a cortical microcircuit dedicated to computing prediction error.

Results

To evaluate the ubiquity of Predictive Coding (PC) as a mechanism for explaining stimulus-history-driven vs. expectation-driven sensory responses, we habituated animals to a local oddball sequence for several days before recording neuronal responses to those local oddballs. We then introduced 2nd order global oddballs on the recording day (Figure 1B-E). This extensive habituation procedure was designed in order to ensure that animals had sufficient exposure to a sequence (e.g., x-x-x-Y) before the start of the experiments. On the recording day for mouse experiments, 6 Neuropixels probes were inserted into visual cortical areas (Figure 1F) where measurable visual responses were identified with expected response latencies by area (Figure 1G). Measurable responses were present across each presentation in the sequence (Figure 1H).

Two cohorts of mice participated, the key difference being the stimulus sequence they were habituated to. One cohort was habituated to a sequence of 3 rightward oriented gratings followed by a leftward oriented grating (x-x-x-Y), the other cohort to 3 leftward then a rightward (y-y-y-X) to counterbalance the stimulus set across the population of animals. Following a 500 ms baseline, each stimulus was presented for 500 ms with 500 ms interstimulus interval, for a total trial duration of 4500 ms (Figure 1E). Mice were habituated to these

sequences (n=8 mice for x-x-x-Y and n=8 mice for y-y-y-X sequences) over the course of 5 sessions for a total
of 2000 trials. For monkey experiments, a total of two macaques were habituated to these sequences with the
additional requirement of central fixation of the eyes on the screen. Monkeys were habituated over the course of
3 sessions for a total of 2850 complete trials prior to recordings. Then, in the experimental session,
mice/monkeys were exposed to their habituated sequence before a block where the habituated sequence was
intermixed with the never-before-experienced global oddball sequence with a lesser frequency (20%). After that
experimental (global-local oddball, or "GLO") block, mice/monkeys experienced a block of randomly orientation
sequences ("Random" block) before the "Control" block where animals viewed repeated x-x-x-x sequences
followed by a y-y-y-y sequences, rendering the same stimuli used in global oddball trials predictable. These
controls were included for evaluating oddball detection when used in conjunction with the responses in the GLO
experimental block.

The primary comparison of interest is composed of the response to the oddball stimulus from the GLO
block (i.e., presentation 4 (P4) of either the local or global oddball sequences) and the response to the identical
stimulus in the same sequence position of the Control block. To account for any non-specific differences between
blocks, we normalize the responses by subtracting the activity to the stimulus prior (presentation 3, P3). For
global oddballs (x_1-x_2-x_3-X), this results in X - x_3 in the GLO block as compared to X - x_3 in the Control block.
Global oddball status is defined by the expectation of the 4^{th} stimulus in the block where X in the experimental
block is unexpected and X in the control block is expected. For local oddballs (x_1-x_2-x_3-Y), this results in Y - x_3
in the GLO block as compared to Y - y_3 in the Control block where local oddball status is defined by the stimulus
deviation (oddball), or lack thereof (control), of the 4^{th} stimulus in the block relative to its prior stimuli. Significant
oddball detection is then determined through statistical comparison of the difference of differences for each
oddball type (P4-P3 GLO block vs. P4-P3 control block) using a nonparametric, cluster-based permutation test
to control for multiple comparisons (see Methods). We use this comparison at the population and single-unit
level to evaluate the role of areas, layers, and cell types for oddball detection.

Local Oddballs

Based on recent work (Hamm and Yuste, 2016; Pak et al., 2021; Westerberg et al., 2021, 2023; Yan et al.,
2018), we expected local oddballs to initially evoke enhanced spiking responses at the earliest stages of visual
processing (H1, Figure 1A). We specifically hypothesized that local oddball detection occurs first in earlier visual
cortical areas (i.e., V1) before being fed forward to higher areas with a measurable delay.

We found that the local oddball evoked an increase in neuronal spiking throughout the visual hierarchy
(Fig. 3). Note that this increased response cannot be explained by the fact that the oddball is a different stimulus
(compared to the rest of the stimulus sequence) since we averaged across all neurons (thus eliminating any
tuning-based preferences for one or the other stimulus), and about half of the animals saw the orthogonal
orientation as the oddball stimulus. Also, across all recorded single-units in the dataset, we found there was a
highly similar number of units strongly preferring each of the two orientations used in the task (LO-orientation:
21.2% of all single-units, GO-orientation: 22.3% of all single-units). This indicates there was not a strong
response bias for one oddball feature over the other within the population of measured units. To summarize,
despite completely randomizing neuronal biases, the local oddball stimulus consistently evoked a larger
response when we examined spiking activity across layers, areas, and animals.

The grand average local oddball response (Figure 2C) also featured another remarkable difference to all
other responses we compared it to - a profound oscillatory component with a cycle roughly resembling 5 Hz.
While the magnitude of this oscillation diminished over the course of the 1 s stimulus presentation, it remained
present until stimulus offset, and could even be observed for another cycle or so after stimulus offset (Figure
1H).

Local oddball detection was present in the majority of single-units measured across brain areas (Figure
2A), supporting our hypothesis (H2, Figure 1A). At the population level, local oddball detection was measured
earlier in lower-order visual areas before emerging in higher-order visual areas (Figure 2B). This was

concomitant with the median latency of visual onset responses of these areas, indicating feedforward
propagation (see also directed functional connectivity analysis, Figure 4). When assessing cell types identified
as critical for the neural instantiation of PC (Bastos et al., 2012) (H1, H3, Figure 1A, layer 2/3 pyramidal cells,
inhibitory interneurons), we found that these populations show enhanced activation to the local oddball as
compared to the control condition (Figure 2C, D). In the case of the layer 2/3 pyramidal cells of area V1 at the
population level, we found greater local oddball detection within those units as compared to other V1 units,
consistent with a feedforward propagation of the local oddball prediction error. Altogether, we largely found
support for the core predictions of the neural instantiation of PC in the local oddball responses.

Hierarchical Profile of Global Oddballs do not reveal feedforward prediction error

Global oddball responses were markedly different. First, detection of global oddballs was restricted to only a few
brain areas at the population level (areas LM, AM, and PM, Figure 3A). We also observed that higher-order brain
areas detected the global oddball (AM: 336 ms post-stimulus, PM: 384 ms post-stimulus) earlier than the
hierarchically earlier area (LM: 394 ms post-stimulus). At the individual unit level, only about 6% of all units
showed significant global oddball detection across areas (Figure 3B) and there was little consistency when the
detection emerged across units (examples in Figure 3C). Previous reports have suggested that sensory
prediction errors signaling might be isolated to neuronal populations which are selective for the features that
define the error producing stimulus (Furutachi et al., 2024). As such, we isolated the population of GO-orientation
selective neurons within our recorded populations and determined whether they specifically harbored a global
oddball detection signal. Of the 22.3% of single-units that were strongly selective for the GO-orientation (relative
to the LO-orientation), only 6.1% of those single units were also significant GO detectors. This is comparable to
the rate of GO detectors we observed in the general population (6.4% of single-units). In other words, GO
detection was not isolated to feature selective populations nor were all units highly selective for the GO stimulus
found to be GO detectors.
Next, we evaluated laminar differences in global oddball detection. For this, we limited analysis to higher-
(AM, PM) vs. lower-order (LM) cortex where we observed significant population global oddball detection. We
divided the units into extragranular vs. granular compartments which should primarily reflect bottom-up vs. top-
down driven factors contributing to global oddball detection. We observed significant population global oddball
detection in only the extragranular compartments, and earlier in higher-order cortex (337 ms post-stimulus) than
lower-order cortex (401 ms post-stimulus) (Figure 3D).
To address the direction of propagation of oddball responses throughout the hierarchy, we next
calculated Granger causality (GC, see Methods) to quantify directed functional connectivity amongst neurons.
For each cortical area and mouse, we combined all available single neurons to create a population spiking time
series (a minimum of N=12 neurons per area were used to create this population spiking time series). We then
calculated GC between each pair of cortical areas, using the anatomical hierarchical ordering: V1, RL, LM, AL,
PM, AM (in order from earliest to latest area in the mouse visual hierarchy, Harris et al., 2019). This generated
a total of 30 inter-areal GC values, one for each area pair and direction of influence. For each pair of areas, we
computed the GC hierarchical asymmetry by subtracting feedforward GC minus feedback GC. For example, the
feedforward GC from V1 to RL was subtracted from the feedback GC from RL to V1. A positive difference
indicates that feedforward GC > feedback GC. A positive difference for both local and global oddballs would be
consistent with the hypothesized (H1, Figure 1A) feedforward propagation of prediction error. We compared
oddball GC responses during stimulus processing in both the early (100-300 ms post-oddball onset) and late
(300-500 ms post-oddball onset) time windows between local and global oddballs and compared them to
baseline (100 ms pre-oddball onset to 100 ms post-oddball onset). GC became significantly hierarchically
asymmetric in the feedforward direction for local, but not global oddballs during both early and late time periods
(Figure 4B-D, Wilcoxon ranksum test of GC hierarchical asymmetry values across mice vs. pre-oddball baseline,
P<0.01). In contrast, GC hierarchical asymmetry did not change during global oddballs compared to baseline

(Figure 4B-D, Wilcoxon ranksum test of GC hierarchical asymmetry values across mice vs. pre-oddball baseline,
all comparisons, P>0.01).

To determine whether top-down prediction signatures throughout all sensory cortical areas might be
hidden in synaptic dynamics rather than spiking activity, we examined the Current Source Density (CSD) profiles
during global oddball detection. CSD is frequently used as a proxy for synaptic activity occurring along the cortical
column (Cox et al., 2019; Godlove et al., 2014; Kajikawa et al., 2017; Maier et al., 2010, 2011; Mehta et al.,
2000a, 2000b; Ninomiya et al., 2015; Schroeder et al., 1998; Self et al., 2013; van Kerkoerle et al., 2014a, 2017;
Westerberg et al., 2019, 2021, 2022, 2023) which reliably reflects columnar processing dynamics, including
those that may be missed by spiking activity (Herrera et al., 2022; Tovar et al., 2023, 2020). We computed CSD
from the LFP responses to the GLO and control sequences (Figure 5A) and then performed the same global
oddball contrast that was done with the spiking data. We hypothesized (H1, Figure 1A) that a difference in
current sinks in the upper or deep layers of cortex might manifest when the prediction is different (i.e., when the
global oddball stimulus is predicted in the control vs. when it is not in the experimental block). In the global
oddball condition, we did not observe reliable changes in synaptic activation at the earliest stage of visual cortical
processing (i.e. in V1, Figure 5B), neither before nor after visual stimulation. Specifically, we did not observe
significant, reliable synaptic changes in the extragranular compartments which would be indicative of top-down
influences deploying a prediction to earlier areas. This observation can also be made in downstream visual
cortical areas (Figure 5C). This indicates the predictions are not being deployed widely through synaptic
modulations nor in the spiking activity as shown earlier.

In summary, we did not find support for a global oddball detection mechanism across all visual cortical
regions, nor that detection propagated to the earliest stages of sensory cortical processing (e.g., in V1), findings
that did not support our hypothesis (H2, Figure 1A). Instead, we found sparse global oddball limited to higher
cortical regions and in the laminar compartments more often associated with cortical feedback than feedforward
activity. Local oddballs but not global oddballs evoked feedforward connectivity amongst visual cortical neurons.

Global Oddballs: Lacking Expected Cell-type Involvement

While the observability of PC elements in global sensory prediction errors appears far more limited in scope than
in the case of local oddballs, it may be that the specific cell types critical to the neural instantiation of PC still play
a critical role in global oddball detection in sensory cortex. Therefore, we next reevaluated global oddball
detection with more careful consideration of the layer 2/3 pyramidal cells, which should reflect the feedforward
element of the prediction error, and the inhibitory interneuron cells, which should show differential activation
reflecting an instantiation of the subtractive prediction.

To identify layer 2/3 pyramidal units we eliminated single-units that were found to be inhibitory
interneurons during the optotagging session for each animal, we eliminated units found outside layer 2 or 3, and
we ensured included units had spike waveform duration of longer than 0.4 ms (Figure 6A), which primarily reflects
broad-spiking pyramidal cells (Foehring et al., 1991; González-Burgos et al., 2004; Henze et al., 2000;
McCormick et al., 1985; Mitchell et al., 2007; Nowak et al., 2003; Povysheva et al., 2006; Tasker et al., 1996).
At the population level, we found that putative layer 2/3 pyramidal cells did not signal global oddball detection
(Figure 6B). In fact, we saw that an even smaller proportion of putative layer 2/3 pyramidal units (5% of single
units) significantly detected than global oddball as compared to the unrestricted set of all units (6% of all single
units) (Figure 6C, failing to support H1, Figure 1A). However, there were sparse units which did signal the global
oddball in all measured areas except area AL (an example single unit response is shown in Figure 6D).

In PC, a subtractive element exists to explain away expected input. This has been posited to be
neuronally instantiated at the level of inhibitory interneuron populations (Bastos et al., 2012). Somatostatin-
(SST) and Parvalbumin- (PV) expressing interneuron populations were identified with optotagging (Figure 7A,
see Methods). PV+/SST+ units were identified across all brain areas of interest (Figure 7B) and on average
showed discernable responses to the stimulus sequences (Figure 7C). However, we did not observe a
substantial population of either interneuron type acting as global oddball detectors, although sparse individual

examples could be observed (Figure 7D). The population averages showed no significant global oddball detection, indicating a minor role, if any role at all, for these populations in detecting global oddballs in sensory cortex (failing to support H3, Figure 1A).

Global Oddballs: Higher-order Detection Outside Sensory Cortex in Macaque Monkeys

To determine whether these results are ubiquitous across species, we recorded an additional data set from a cohort of monkeys (n=2). On the recording day for the monkey experiments, 1-4 high-density deep array probes (Diagnostic Biochips, see Methods) and/or V probes (Plexon) were inserted into visual cortex and frontal cortex. The monkeys experienced identical patterns of oddball sequences (Figure 8A). Electrode arrays were implanted in frontal cortical regions (FEF, PFC) as well as six visual cortical areas (V1, V2, V3, V4, MT, MST). Visual cortex recordings can be used to test whether predictive coding is a widespread, ubiquitous phenomenon across species (H2, Figure 1A).

The first and most apparent observation is that the pattern of local vs. global oddball detection is largely consistent between species. Local oddballs were detected across all measured brain areas in the monkey (Figure 8B, left). Similar to the mouse, global oddballs followed a roughly inverse hierarchical pattern. They emerged first in FEF (32 ms), next in V3/PFC (91/128 ms), then in MT (160 ms) (using the same statistical tests as in the mouse). Global oddball representation was not detected at the earliest stages of visual processing (V1 and V2). Once again this demonstrates the relatively sparse representation of global oddballs in sensory cortex (this fails to support H2, Figure 1A). What is also worth noting is that there was significant global oddball detection in both of the prefrontal areas and the associated modulation is stronger than that observed in sensory cortex. In sum, findings were largely consistent across species and moreover, we observed that global oddball detection is robustly signaled in prefrontal cortex even though its propagation through the visual cortex is relatively limited, weak, and late in the temporal evolution of the response.

Discussion

Local vs. Global Oddballs

In this study, we tested the neuronal implementation of predictive coding computations across visual and higher-order cortex in mice and monkeys. We implemented an extensive habituation period that exposed subjects to a predictable sequence (e.g., x-x-x-Y) for multiple sessions. This enabled subjects to create an internal model. We aimed to test whether deviations from this model (prediction errors) were signaled when subjects were shown global oddballs (e.g., x-x-x-x) on 20% of trials on recording days. We performed an extensive survey of spiking responses throughout the visual cortex of mice and monkeys using MaDeLaNe (Multi-Area, high-Density, Laminar Neurophysiology) recordings. We tested three core hypotheses of current PC models (Figure 1A, Bastos et al., 2012; Friston, 2005; Rao and Ballard, 1999) by examining coding of local and global oddballs. These models hypothesized prediction error computations to invoke feedforward mechanisms, be robust and widespread, and engage release from predictive suppression by modulating inhibitory interneuron activity.

Local oddballs largely followed this pattern. They emerged early in time, in superficial layers, and fed forward. The feedforward nature of local oddballs was evident both in temporal order of spiking response and as measured by directed functional connectivity analysis. Local oddballs engaged a robust and wide-spread response, recruiting around half the neurons in all visual cortical areas in both species. However, local oddballs did not result from a reduction in levels of PV+/SST+ inhibitory neuron activity - these inhibitory neurons also *increased* their firing rates during local oddballs. This implies that the predictive subtraction computation implied by predictive coding is not implemented via these inhibitory interneurons. Instead it seems more likely that local oddballs in this temporal context rely mostly on membrane-specific adaptation (Sanchez-Vives et al., 2000a, 2000b) and do not require feedback, as they survive (or are even enhanced) during anesthetized states (Nourski

et al., 2018; Uhrig et al., 2016; Xiong et al., 2024). Therefore, local oddball responses by themselves do not
provide strong evidence for ubiquitous PC, which envisions that prediction errors are the result of a subtraction
between top-down prediction and bottom-up sensory drive under varying sensory and cognitive conditions.
In contrast to local oddballs, neuronal responses to global oddballs did not support any of the PC
hypotheses (Figure 1A) that we tested. In sensory areas, global oddballs emerged late in the sensory response.
Global oddballs emerged first in higher-order areas AM/PM in mice, and first in FEF in monkeys, and next in
lower-order areas LM in mice and V3/MT in monkeys. This temporal progression resembled feedback processing
more than it did feedforward. Global oddballs were not encoded in the population of putative pyramidal neurons
in L2/3. Instead, they emerged weakly and sparsely in non-granular layers. This laminar pattern of neuronal
response is more indicative of cortical feedback, not feedforward as hypothesized. Global oddballs should have
elicited prediction error responses. But they were only detected in a small number of neurons in each area (3-
9%). Inhibitory interneurons did not selectively encode the global oddballs.

Predictive Coding: a ubiquitous and feedforward or sparse and feedback process?

Predictive coding (PC) models have hypothesized that a core cortical computation is prediction error (Mumford,
1992; Rao and Ballard, 1999; Srinivasan et al., 1982). We set up our task so that subjects were highly habituated
to a particular sequence (e.g., x-x-x-Y) such that its violation (e.g., x-x-x-X) should drive robust neuronal
representations of prediction error. In PC the local oddballs (e.g., x-x-x-Y) should not drive much of a prediction
error signal because they were fully predictable. To reiterate: under PC, neuronal encoding of the local oddball
should be weak (as it is predictable, so should be subtracted away) and encoding of the global oddball should
be robust (as it is a violation to a well-established pattern). We found the opposite. Spiking responses of cortical
neurons robustly encoded unsurprising local oddballs but weakly encoded surprising global oddballs. Neither
oddball type engaged a release of inhibition from PV/SST+ interneurons. Global oddballs, when they elicited
responses, first engaged higher cortical areas typically associated with cognitive (e.g., PFC), not sensory
functions (Miller and Cohen, 2001). This argues against the ubiquity of predictive coding signals in the cortex,
especially in sensory cortex.
Another core tenant of predictive coding models is that prediction error computations were hypothesized
to emerge early in sensory processing and feed forward. This hypothesis describes local oddball processing
relatively well. However, global oddball processing was better described by a feedback pattern of processing.
Global oddballs emerged late in the sensory response and in higher-order areas first, then lower-order areas (in
order of increasing hierarchical level for mice: 394 ms in LM, 384 ms in PM, 336 ms in AM, in order of increasing
hierarchical level for macaques: 91 ms in V3, 160 ms in MT, 32 ms in FEF, 128 ms in PFC). Putative excitatory
pyramidal neurons in L2/3 (which project feedforward connections) and their associated current sinks did not
encode global oddballs. During local oddballs, Granger causality analysis showed a robust increase in
feedforward directed functional connectivity, with V1 increasing its functional influence on downstream areas.
This enhanced feedforward connectivity was not observed during global oddballs. Therefore, by four
independent metrics, including hierarchical area, temporal order, Granger causality and laminar compartment of
spiking/current flow, we failed to detect a feedforward component during global oddball processing. This violates
a core tenet of PC models that hypothesized a substantial engagement of L2/3 pyramidal neurons in sensory
cortex that transmit prediction errors in the feedforward direction.
Instead, our findings pointed to the engagement of cortical feedback as the predominant pathway by
which global oddballs are signaled. This was seen in the temporal latencies of global oddball responses as well
as the laminar pattern of activation: global oddballs were found to emerge at the population level in the non-
granular layers. These non-granular layers are associated with feedback processing, not feedforward
processing.

Caveats

While the findings already demonstrate that the hypothesized ubiquitous elements of Predictive Coding are actually more nuanced, there are other crucial considerations that must be taken into account. For one, we have chosen to study pure global oddballs, which are deviations from expectations that are induced by the lack of a stimulus change. It is possible that other forms of expectation that do depend on local stimulus change (e.g., such as a local oddball which is also a global oddball) are encoded in sensory cortex (e.g., Furutachi et al., 2024; Jamali et al., 2024). However, more complex forms of expectation processing which were envisioned by PC models, such as the global oddball, do not appear to robustly engage sensory cortical representations.

Another important consideration is the degree of attention that is paid to the oddballs. Previous work suggested that global oddballs cannot be observed under certain states of unconsciousness (Nourski et al., 2018; Xiong et al., 2023). Given the close relationship between conscious awareness and attention (Maier and Tsuchiya, 2021), it seems plausible that attention may be a prerequisite for global oddball processing (Bekinschtein et al., 2009). Indeed, hierarchical predictive coding models have postulated that prediction error signals are modulated by a gain term that takes on a functional interpretation in terms of precision (Π in Figure 1A, see also (Feldman and Friston, 2010)). That being said, global oddballs were observed in prefrontal areas in monkeys and in higher-visual areas in mice, and thus despite being available to the animals, did not have a profound effect on sensory responses. Models that consider gain in terms of precision require that precision must be positive and nonzero (Bolstad and Curran, 2016), entailing that prediction error signals might be attenuated, but are never entirely quashed, by gain modulation. Future studies should explicitly consider the role of gain modulation on prediction error responses by manipulating the attentional state.

Updates to Predictive Coding models

Our results challenge core hypotheses of current PC models (Figure 1A). Global oddballs emerged in hierarchically late areas in both species. They did not feedforward throughout the visual cortex and did not preferentially engage feedforward projecting layers 2/3. Local oddballs, given their extensive habituation, should have been an unsurprising stimulus evoking minimal prediction error. Yet, they engaged activity in over half of recorded cortical neurons. Global oddballs were designed to be a surprising stimulus to evoke strong prediction errors. But they did not engage a robust response in visual cortex, neither in mice nor in monkeys. Instead, global oddballs were initially signaled late in hierarchical processing, area AM in mice, prefrontal area FEF in monkeys. So although global oddballs are available at high levels of the hierarchy and could in principle have informed and guided sensory processing, they did not drive a prediction error response in sensory cortex as hypothesized by PC.

While our results rule out most PC models from the existing modeling literature, the broader family of predictive processing theories have resources for new models that could capture the effects we observe. The most prominent such resource involves the brain representing a hierarchy of time-scales (Kiebel et al., 2008) ranging from the very short in primary sensory cortex to the more long-term in higher-order areas such as PFC. Such a theory would imply that we only observe prediction-error encoding in regions of the brain representing the stimulus or task at a sufficiently long timescale to capture the prediction error induced by the oddball stimulus. Forward models based on Slow Feature Analysis have previously captured observed properties of a number of sensory and higher-order cell types (Berkes and Wiskott, 2005; Dähne et al., 2014; Franzius et al., 2007); PC could serve as a learning mechanism for such models.

Rather than being a computation that originates at the level of sensory processing, these data thus lead us to propose that prediction errors are computed in higher-order areas such as PFC. Higher areas such as the PFC display the property of mixed selectivity (Rigotti et al., 2013). Neurons in these areas have response properties that are not based on sensory bottom-up features and are more flexible with regard to response preferences. These higher-order neurons can re-map "on the fly" based on experience-dependent learning. This property is ideal for representing predictions, which map out the statistical structure of a changing and context-

dependent environment. Therefore, the neuronal instantiation of prediction itself requires a flexible, dynamic mechanism to adapt to changing environments. Neurons in sensory cortex, with receptive field properties defined at evolutionary time scales and further refined during development, seem ill-suited for such a computation.

We propose instead that ensembles of PFC neurons (and neurons in other higher-order areas) generate predictions based on experience-dependent learning. This ensemble learning may be facilitated by beta-rhythmic oscillations that have properties seemingly ideal for formation of predictions. Namely, they can form neuronal ensembles in a dynamic and cognition-dependent manner (Antzoulatos and Miller, 2016; Buschman et al., 2012; Miller et al., 2018) and are associated with feedback communication in cortex (Bastos and Schoffelen, 2015; Michalareas et al., 2016; van Kerkoerle et al., 2014b; Vezoli et al., 2021). This pattern of predictive activity (as well as violations to prediction) can then be communicated selectively and sparsely via top-down feedback to sensory and other areas that "need to know" it. This is consistent with global oddballs evoking a feedback pattern of activity. It is also consistent with our observation of sparse and less wide-spread patterns of global oddball responses and the absence of predictive information in primary and secondary sensory areas. In contrast, unsurprising local oddballs were robustly encoded in cortex and drove strong feedforward processing. To summarize, based on the global oddball paradigm and an extensive spiking survey of cortical activity in mice and monkeys, our data indicate that sensory feedforward processing is not in the business of primarily reporting prediction errors as traditional PC models hypothesized. Sensory processing is for the most part sensory in nature, i.e., representing stimulus features without perturbation by predictive information. Predictive codes are not a ubiquitous property of cortical processing that drive cortical responses. They are largely a property of higher-order neurons and are conveyed to sensory areas in a selective and sparse manner and with a modulatory nature.

Methods

Animals

Mouse Experiments at the Allen Institute

All experiments were performed in 16 C57/B6 mice (*mus musculus*) of both sexes (n_{male}=6, n_{female}=10), aged 95-128 days. Optogenetics experiments were conducted with Sst-IRES-Cre/wt; Ai32(RCL-ChR2(H134R)_EYFP)/wt mice (n=9) and Pvalb-IRES-Cre/wt; Ai32(RCL-ChR2(H134R)_EYFP)/wt) mice (n=7). All experiments on animals were conducted with approval of the Allen Institute's Institutional Animal Care and Use Committee.

Macaque Monkey Experiments at Vanderbilt University

All procedures were approved by the Vanderbilt University Institutional Animal Care and Use Committee, in compliance with the regulations set by the Association for the Assessment and Accreditation of Laboratory Animal Care, and followed the guidelines of the United States National Institutes of Health. Twomacaque monkeys were used in this study: one male adult bonnet macaque (*Macaca radiata*) aged 19 years (Ca, 7.4kg), and one male adult rhesus macaque (*Macaca mulatta*) aged 11 years (Jo, 12.0 kg).

Surgery

Mouse Experiments at the Allen Institute

Mice were anesthetized with isoflurane (2%) and mounted on a stereotaxic apparatus. Buprenorphine (0.05 mg/kg, analgesic) and dexamethasone (2 mg/kg, anti-inflammatory) were injected subcutaneously. Body temperature was maintained at 36.5°C. After hair removal and disinfection with 70% ethanol and 5% iodine, scalp was removed and facia was retracted. A circular craniotomy of 3–5 mm was made on the left hemisphere

using a biopsy punch (Robbins Instruments) and/or a dental drill (Foredom) with a 0.5 mm drill bit (FST Item No.
19007–05). Finally, a titanium head plate was fixed to the skull with dental cement (Metabond C&B).

For optogenetics experiments, we injected the GCaMP6m-p2A-ChRmine-Kv2.1 virus during the surgery
(500 nL per 2 depths and 2 sites). Injection speed was 100–250 nL/min, and after injection in each site, pipette
was held in place for 5 minutes. Injection sites were centered at 2.7 mm lateral, 0.2 mm anterior relative to
lambda, and injection depths were 120 μm and 250 μm.

The 5 mm craniotomy was sealed with a glass coverslip, the bottom of which was coated with silicone to
reduce adhesion to the brain surface. Glass coverslip held in place with Vetbond and Kwik-Cast (World Precision
Instruments) until the day of the recording (several weeks later). These glass coverslips, or 'windows', were
made beforehand by gluing 2 layers of round cover glass (e.g., 1 x 3 mm + 1 x 5 mm, Warner Instruments, #1
thickness) with Norland Optical Adhesive #71 (UV-cured).

Intrinsic signal imaging was performed to identify visual area boundaries (Juavinett et al., 2017). Based
on the identified visual areas, an insertion window was designed for each mouse. Mice were habituated to head-
fixation and visual stimulation over a period of two weeks.

Macaque monkey experiments at Vanderbilt University

Monkeys were implanted with a head post and recording chambers during an anesthetized procedure. Monkeys
were anesthetized with isoflurane. Body temperature will be maintained near 38°C with a circulating water or
warm air recirculating heating blanket. Peripheral oxygen levels as well as heart rate, temperature, ECG and
expired CO_2 levels were monitored continuously during surgery. The animals' head was shaved and scrubbed
(with Betadine and ethanol) after mounting the animal in a stereotaxic device. Local anesthetic (Lidocaine) was
injected subcutaneously before opening and also placed into ear canals before fitting the animal into the
stereotaxic device.

A skin flap was made over the skull. The skull was cleaned and dried, and orthopedic screws were placed
in the skull. The head restraint device, held in position with a stereotaxic holder, was fixed in place with orthopedic
screws and methyl methacrylate cement. Custom-made recording chambers (~20 mm in diameter, Crist
Instruments) were placed on the skull to allow access to all targeted brain areas in one hemisphere and secured
in place with orthopedic screws and dental cement. Craniotomies (10-20 mm diameter) were subsequently
performed using piezosurgery drills (Piezosurgery, Columbus, Ohio) to permit access to the underlying cortex.
Monkey Ca had two recording chambers, one placed over the prefrontal cortex (giving access to areas FEF and
LPFC) and one placed over the temporal cortex (giving access to areas MT, MST, V4, and TEO). Monkey Ca
had one recording chamber positioned over the occipital cortex (giving access to areas V1, V2, V3). After
surgery, analgesics were given for pain management (Buprenorphine and NSAIDs for 3 days), and animals
showed no signs of pain or distress.

Electrophysiology Experiments

Mouse Experiments at the Allen Institute

The Neuropixels data was acquired at the Allen Institute as part of the OpenScope project that allows the
community to apply for observation on the Allen Brain Observatory platform
(https://alleninstitute.org/division/mindscope/openscope/). The Neuropixels pipeline at the Allen Institute is
described in detail in previous reports (Durand et al., 2023; Siegle et al., 2021).

On the day of recording, the cranial coverslip was removed and replaced with the insertion window
containing holes aligned to six cortical visual areas. Mice were allowed to recover for 1–2 hours after insertion
window placement, before being head-fixed in the recording rig.

Six Neuropixels probes were targeted to each of the six visual cortical areas (V1, LM, RL, AL, PM, AM).
Probes were doused with CM-Dil (1 mM in ethanol; Thermo Fisher, V22888) for post hoc ex vivo probe

localization. Each probe was mounted on a 3-axis micromanipulator (New Scale Technologies). The tip of each probe was aligned to its associated opening in the insertion window using a coordinate transformation obtained via a previous calibration procedure. The XY locations of the six visual area targets were translated into XYZ coordinates for each 3-axis manipulator using a custom Python script. The operator then moved each probe into place with a joystick, with the probes fully retracted along the insertion axis, approximately 2.5 mm above the brain surface. The probes were then manually lowered one by one to the brain surface until spikes were visible on the electrodes closest to the tip. After the probes penetrated the brain to a depth of around 100 µm, they were inserted automatically at a rate of 200 µm/min (total of 3.5 mm or less in the brain). After the probes reached their targets, they were allowed to settle for 5–10 min.

Neuropixels data was acquired at 30 kHz (spike band, 500 Hz high-pass filter) and 2.5 kHz (LFP band, 1,000 Hz low-pass filter) using the Open Ephys GUI (Siegle et al., 2017). Videos of the eye and body were acquired at 30 Hz. The angular velocity of the running wheel was recorded at the time of each stimulus frame, at approximately 60 Hz.

The spike-band data was median-subtracted, first within-channel to center the signal around zero, then across channels to remove common-mode noise. The median-subtracted data file was sent through the Kilosort2 MATLAB package (https://github.com/mouseland/Kilosort2) (Pachitariu et al., 2023), which applies a 150 Hz high-pass filter, followed by whitening in blocks of 32 channels. Kilosort2 models this filtered, whitened data as a sum of spike 'templates'. The shape and locations of each template was iteratively refined until the data can be accurately reconstructed from a set of N templates at M spike times. Finally, Kilosort2 output was curated to remove putative double-counted spikes and units with artefactual waveforms. All units not classified as noise were packaged into Neurodata Without Borders (NWB) files (Rübel et al., 2022) and analyzed in this paper.

Current source density (CSD) (Freeman and Nicholson, 1975; Mitzdorf, 1985; Nicholson and Freeman, 1975; Nicholson and Llinas, 1971) was estimated from the LFP band data by epoching the LFPs, median-filtering with a filter-width of 3 channels, and then downsampling to every 4^{th} channel to retrieve a single vertical/laminar column of responses. CSD was then estimated as the 2^{nd} spatial derivative across multiple contact points according to the equation of (Mitzdorf and Singer, 1978):

$$CSD(t,c) = -\sigma\,(V(t, c - 2) - 2V(t,c) + V(t, c + 2))/(2s^2)$$

where $V(t, c)$ is the extracellular voltage at time t measured at an electrode contact at position c, s is the inter-contact distance of the electrode array, and σ is the tissue conductivity (here taken as 0.4 S/m). CSD contrasts were computed by grand-averaging the raw LFPs over each condition, then calculating the CSD as above and subtracting across conditions. We corrected for time in task as a confounder via double contrasts, first subtracting the 3^{rd} (fully adapted) presentation response from that of the 4^{th}, and then pooling trials, within task conditions. These subtractions were treated as normal random variates for a two-tailed, nonparametric spatiotemporal cluster-test that controls the family-wise error rate to $p < 0.05$ to determine significant cross-condition contrasts (Maris and Oostenveld, 2007). Both LFP and CSD are smoothed with a width 4 Gaussian filter in both the spatial (laminar) and temporal dimensions for plotting.

Macaque Monkey Experiments at Vanderbilt University

We acquired data using between 1-4 linear array multi-channel recording probes. In most experiments, we recorded with 128-channel "deep array" probes where the inter-contact spacing was either 25 or 40 µm (Diagnostic Biochips, Glen Burnie, MD) and the coverage was 3.175 mm and 5.08 mm, respectively. In some experiments, we used 32-channel V probes with an inter-contact spacing of 100 µm (Plexon, Dallas, TX) with a coverage of 3.1 mm. To insert electrodes into the appropriate depth and area we used a mechanical Microdrive (Narishige, Tokyo, Japan) that was mounted onto the chambers. A guide tube was lowered through a recording grid (Crist Instrument Co., Inc.) to penetrate the dura mater and granulation tissue. We then acutely introduced the recording probe into the respective area, and waited ~1 hour for the signal to settle prior to starting the

recordings. Recordings were acquired using a RHD System (Intan, Los Angeles, CA) sampling at 30 kHz. Recordings were referenced to the guide tube which remained in contact with tissue. Grid positions were pre-determined based on a pre-recording MRI scan with the recording chambers and grid in place, with water to mark the trajectories of each grid hole position. We advanced the electrode until there was visually responsive neuronal activity on most channels and until the local field potentials showed a distinct spectral signature of the cortical sheet, characterized by large amplitude alpha/beta (10-30 Hz) oscillations in deep channels and gamma (40-150 Hz) oscillations in superficial channels (Maier et al., 2010; Mendoza-Halliday et al., 2024; Tovar et al., 2023, 2020; Westerberg et al., 2019). Offline, using MATLAB the data was filtered between 500-5000 Hz, full-wave rectified, low-pass filtered below 250 Hz, and downsampled to 1000 Hz. We refer to this signal as the MUAe signal and it reflects the activity of nearby surrounding neurons (Logothetis et al., 2001; Self et al., 2013; Supèr and Roelfsema, 2005; Tovar et al., 2020; Trautmann et al., 2019; Westerberg et al., 2023, 2020).

Visual Stimulation Parameters

Mouse Experiments at the Allen Institute

All Neuropixels recordings were according to the standardized Neuropixels visual coding pipeline. Visual stimuli were generated using custom scripts based on PsychoPy62 and were displayed using an ASUS PA248Q LCD monitor, with 1920 X 1200 pixels (55.7 cm wide, 60 Hz refresh rate). Stimuli were presented monocularly, and the monitor was positioned 15 cm from the right eye of the mouse and spanned 120 dva × 95 dva. Monitor placement was standardized across rigs such that mouse right eye gaze would typically be directed at the center of the monitor.

Macaque Monkey Experiments at Vanderbilt University

We used MonkeyLogic (developed and maintained at the National Institute for Mental Health) to control the behavioral task (Hwang et al., 2019). Visual stimuli were displayed using PROPixx Pro projectors (VPixx Technologies, Quebec, Canada) with a resolution of 1920 x 1080 at a 120 Hz refresh rate. The projector screen was positioned 57 cm in front of the monkey's eyes.

The monkey was trained to fixate its eyes around a central fixation dot (radius of fixation window: 1.5 dva). Eye position was monitored using an Eyelink 1000 (SR Research, Ottawa, Canada). A task began with an isoluminant gray screen. Once the monkey maintained fixation for 500ms, a sequence of drifting gratings appeared (12 dva radius, 2 Hz drift rate, 1 cycle/degree, 0.8 Michelson contrast, angle of drifting grating was 45 or 135 degrees from the vertical midline) for 500 ms, and was replaced by a blank screen with the fixation dot remaining for 500 ms. This sequence was repeated 4 times in a trial, for a total trial length of 4500 ms (500 ms pre-sequence fixation + 4x1000 ms stimulus presentations). The monkey was rewarded for maintaining fixation throughout the trial with a small juice reward.

Habituation to predicted sequence

Mouse Experiments at the Allen Institute

Mice were habituated over the course of 5 behavior-only sessions in which only the predicted sequence (either xxxY or yyyX) was presented. This resulted in 2000 habituation trials prior to recordings.

Macaque Monkey Experiments at Vanderbilt University

Monkeys were habituated to a predicted sequence over the course of 3 behavior-only sessions in which only the predicted sequence (either xxxY or yyyX) was presented. This resulted in 2850 complete habituation trials prior

to recording sessions. Subsequently after a recording session, monkeys were again habituated to one of the
predicted sequences prior to the next recording session.

Data Analysis

Mouse Experiments at the Allen Institute

We recorded 29977 total putative single-units, with an average of 359 (+/-42) units per animal and an average
of 60 (+/-3) units per probe. We further restricted the units for analysis, consistent with previous reports using
the same methods for analyses (Siegle et al., 2021), to those which exhibit a visual response that exceeded 5
standard deviations of the baseline firing rate, minimum of 0.1 sp/s, inter-spike interval (ISI) violations below
0.33, a presence ratio above 0.95, an amplitude cutoff below 0.1, and were found to be in areas V1, LM, RL, AL,
PM, AM, or LP. This resulted in a sample of 5389 units from 15 of 16 mice. No units from 1 mouse passed criteria
due to a destabilizing event during the recording. Additionally, data from another mouse were omitted from
analysis due to issues with the visual stimulus timing. As a result, our final sample consisted of 5062 single-units
from 14 mice.

Macaque Monkey Experiments at Vanderbilt University

We recorded a total of 8,672 sites with MUAe signals across 11 cortical areas: 640 in PFC, 416 in FEF, 1472 in
PMD, 512 in MST, 512 in MT, 608 in TEO, 1056 in V4, 1536 in V3, 896 in V2, and 1024 in V1. These MUAe
signals were extracted from broadband signal with a butterworth filter of stopbands at 450Hz and 5500Hz, and
passbands at 500Hz and 5000Hz. For analysis, we restricted these sites that had visually evoked activity, which
was determined by whether the time period from 50-250 ms post-stimulus was significantly greater than pre-
stimulus baseline (using a threshold of p<0.05, rank sum test). This resulting population that was used for
analysis consisted of 508 sites in PFC, 177 in FEF, 161 in MST, 252 in MT, 230 in V4, 569 in V3, 328 in V2, and
270 in V1.

Granger causality analysis

In mice, we began by defining all available single units in each area. We took all cortical neurons for this analysis,
and per mouse and probe, included an area in the analysis if a minimum of 12 single neurons were
simultaneously acquired. This resulted in a total of six neuronal spiking time series, one per area. Granger
causality was calculated separately per mouse and separately for local and global oddball trials, using all
available trials for the calculation. We calculated Granger causality using a non-parametric implementation
(Dhamala et al., 2008). For this, we computed fourier coefficients using the Fieldtrip toolbox (Oostenveld et al.,
2011) using a sliding window approach utilizing 200ms time windows for analysis. We computed the Fourier
coefficients using multitapers (3 multitapers per trial, time window size = 200ms, smoothing = 10 Hz, function
'ft_freqanalysis', using method 'mtmfft', frequencies 0-100 Hz were calculated). One time window was on the
baseline, from 100ms pre-oddball onset to 100ms post-oddball onset. Two time windows reflecting stimulus
processing were also used, one centered from 100ms to 300ms post-oddball onset, another from 300ms to
500ms post-oddball onset. Bivariate Granger causality was computed pair-wise between all available neuronal
spiking time series (using ft_connectivity analysis with method = 'granger' and granger.sfmethod = 'bivariate').

Statistics and Visualization

All data was normalized by its average level of activity to produce an arbitrary value that was less variable across
units with differing firing rates. Unless otherwise noted, all statistics were performed using nonparametric, cluster-
based permutation testing (Maris and Oostenveld, 2007) of the spiking activity at either the single-(multi)unit
level or at the population level with alpha thresholds set to 0.05 for both stages of the test. We additionally
required that a statistically significant difference as indicated by this test persist for at least 25 ms. Data was

smoothed with a 50 ms moving mean window. Confidence clouds, where displayed, are computed for a 95% confidence interval across units' condition-wise mean activity included in the population.

Feature selectivity was measured in the same way as a previous report (Furutachi et al., 2024). First, we determined the mean response to each of the stimuli during visual stimulation (0-500 ms following stimulus appearance) and during a separate presentation block where stimulus order was randomized. For each single-unit, we subtracted the mean response to GO-orientation stimulus from the LO-orientation stimulus and divided by the sum of those two means. Single-units with this index above 0.8 or below -0.8 were defined as highly selective for the GO-orientation or the LO-orientation, respectively.

Data and Code Availability

All mouse data used in this work are openly available through the DANDI Archive (dandiset id: 000253, doi: https://doi.org/10.48324/dandi.000253/0.240503.0152). Software resources for the analysis of this dataset are provided here: https://alleninstitute.github.io/openscope_databook/projects/glo.html. The code to generate the global-local oddball sequences is provided here: https://github.com/AllenInstitute/openscope-glo-stim.

Acknowledgements

This work was funded by the National Institutes of Health [grant number: R00MH116100] (A.M.B.), through Vanderbilt University Startup Funds (A.M.B.), and via funding from the Vanderbilt Brain Institute Faculty Fellow award (A.M.B.). J.A.W. was funded by a fellowship from the International Human Frontier Science Program Organization [grant number: LT0001/2023-L]. The Neuropixels dataset was obtained at the Allen Brain Observatory as part of the OpenScope program, which is operated by the Allen Institute, Neural Dynamics. OpenScope was supported by the National Institutes of Health [grant number: U24NS113646]. We also thank the Allen Institute founder, Paul G. Allen, for his vision, encouragement, and support.

Declaration of Interests

The authors declare no competing interests.

Author contributions

Conceptualization: J.A.W., A.M., A.M.B.; Data curation: J.A.W., A.B., B.H., J.A.L.; Formal analysis: J.A.W., Y.S.X., E.S., A.M.B; Funding acquisition: J.A.W., J.A.L., A.M.B.; Investigation: J.A.W., Y.X., H.N., S.D., H.C., H.B., H.L. A.M.B.; Methodology: J.A.W., W.H., K.N., V.H., T.J., C.G., B.H., A.Y., J.S., R.G., B.O., S.C., A.W., P.A.G., J.A.L., A.M., A.M.B.; Project administration: J.A.W., C.K., J.A.L., A.M., A.M.B.; Software: J.A.W., A.B., C.R.P.; Supervision: S.R.O., J.A.L., A.M., A.M.B.; Validation: J.A.W., A.B., C.R.P., J.A.L.; Visualization: J.A.W., Y.S.X.; Writing - original draft: J.A.W., A.M., A.M.B.; Writing - review and editing: all authors.

References

Antzoulatos, E.G., Miller, E.K., 2016. Synchronous beta rhythms of frontoparietal networks support only behaviorally relevant representations. eLife 5, e17822. https://doi.org/10.7554/eLife.17822

Bastos, A.M., Schoffelen, J.-M., 2015. A Tutorial Review of Functional Connectivity Analysis Methods and Their Interpretational Pitfalls. Front. Syst. Neurosci. 9, 175. https://doi.org/10.3389/fnsys.2015.00175

Bastos, A.M., Usrey, W.M., Adams, R.A., Mangun, G.R., Fries, P., Friston, K.J., 2012. Canonical Microcircuits for Predictive Coding. Neuron 76, 695–711. https://doi.org/10.1016/j.neuron.2012.10.038

Bekinschtein, T.A., Dehaene, S., Rohaut, B., Tadel, F., Cohen, L., Naccache, L., 2009. Neural signature of the conscious processing of auditory regularities. Proc. Natl. Acad. Sci. 106, 1672–1677. https://doi.org/10.1073/pnas.0809667106

Bellet, M.E., Gay, M., Bellet, J., Jarraya, B., Dehaene, S., Kerkoerle, T. van, Panagiotaropoulos, T.I., 2024. Spontaneously emerging internal models of visual sequences combine abstract and event-specific information in the prefrontal cortex. Cell Rep. 43. https://doi.org/10.1016/j.celrep.2024.113952

Berkes, P., Wiskott, L., 2005. Slow feature analysis yields a rich repertoire of complex cell properties. J. Vis. 5, 9. https://doi.org/10.1167/5.6.9

Bolstad, W.M., Curran, J.M., 2016. Introduction to Bayesian Statistics, 3rd ed. Wiley. https://doi.org/10.1002/9781118593165

Brown, H.R., Friston, K.J., 2013. The functional anatomy of attention: a DCM study. Front. Hum. Neurosci. 7, 784. https://doi.org/10.3389/fnhum.2013.00784

Buschman, T.J., Denovellis, E.L., Diogo, C., Bullock, D., Miller, E.K., 2012. Synchronous oscillatory neural ensembles for rules in the prefrontal cortex. Neuron 76, 838–846. https://doi.org/10.1016/j.neuron.2012.09.029

Chao, Z.C., Takaura, K., Wang, L., Fujii, N., Dehaene, S., 2018. Large-Scale Cortical Networks for Hierarchical Prediction and Prediction Error in the Primate Brain. Neuron 100, 1252-1266.e3. https://doi.org/10.1016/j.neuron.2018.10.004

Cox, M.A., Dougherty, K., Westerberg, J.A., Schall, M.S., Maier, A., 2019. Temporal dynamics of binocular integration in primary visual cortex. J. Vis. 19, 13. https://doi.org/10.1167/19.12.13

Dähne, S., Wilbert, N., Wiskott, L., 2014. Slow Feature Analysis on Retinal Waves Leads to V1 Complex Cells. PLOS Comput. Biol. 10, e1003564. https://doi.org/10.1371/journal.pcbi.1003564

Dhamala, M., Rangarajan, G., Ding, M., 2008. Estimating Granger Causality from Fourier and Wavelet Transforms of Time Series Data. Phys. Rev. Lett. 100, 018701. https://doi.org/10.1103/PhysRevLett.100.018701

Durand, S., Heller, G.R., Ramirez, T.K., Luviano, J.A., Williford, A., Sullivan, D.T., Cahoon, A.J., Farrell, C., Groblewski, P.A., Bennett, C., Siegle, J.H., Olsen, S.R., 2023. Acute head-fixed recordings in awake mice with multiple Neuropixels probes. Nat. Protoc. 18, 424–457. https://doi.org/10.1038/s41596-022-00768-6

Feldman, H., Friston, K., 2010. Attention, Uncertainty, and Free-Energy. Front. Hum. Neurosci. 4. https://doi.org/10.3389/fnhum.2010.00215

Foehring, R.C., Lorenzon, N.M., Herron, P., Wilson, C.J., 1991. Correlation of physiologically and morphologically identified neuronal types in human association cortex in vitro. J. Neurophysiol. 66, 1825–1837. https://doi.org/10.1152/jn.1991.66.6.1825

Franzius, M., Sprekeler, H., Wiskott, L., 2007. Slowness and Sparseness Lead to Place, Head-Direction, and Spatial-View Cells. PLOS Comput. Biol. 3, e166. https://doi.org/10.1371/journal.pcbi.0030166

Freeman, J.A., Nicholson, C., 1975. Experimental optimization of current source-density technique for anuran cerebellum. J. Neurophysiol. 38, 369–382. https://doi.org/10.1152/jn.1975.38.2.369

Friston, K., 2010. The free-energy principle: a unified brain theory? Nat. Rev. Neurosci. 11, 127–138. https://doi.org/10.1038/nrn2787

Friston, K., 2005. A theory of cortical responses. Philos. Trans. R. Soc. B Biol. Sci. 360, 815–836. https://doi.org/10.1098/rstb.2005.1622

Friston, K.J., Parr, T., de Vries, B., 2017. The graphical brain: belief propagation and active inference. Netw. Neurosci. 1–78. https://doi.org/10.1162/NETN_a_00018

Furutachi, S., Franklin, A.D., Aldea, A.M., Mrsic-Flogel, T.D., Hofer, S.B., 2024. Cooperative thalamocortical circuit mechanism for sensory prediction errors. Nature 633, 398–406. https://doi.org/10.1038/s41586-024-07851-w

Gallimore, C.G., Ricci, D.A., Hamm, J.P., 2023. Spatiotemporal dynamics across visual cortical laminae
support a predictive coding framework for interpreting mismatch responses. Cereb. Cortex 33, 9417–
9428. https://doi.org/10.1093/cercor/bhad215
Garrido, M.I., Kilner, J.M., Kiebel, S.J., Friston, K.J., 2007. Evoked brain responses are generated by feedback
loops. Proc. Natl. Acad. Sci. U. S. A. 104, 20961–20966. https://doi.org/10.1073/pnas.0706274105
Godlove, D.C., Maier, A., Woodman, G.F., Schall, J.D., 2014. Microcircuitry of Agranular Frontal Cortex:
Testing the Generality of the Canonical Cortical Microcircuit. J. Neurosci. 34, 5355–5369.
https://doi.org/10.1523/JNEUROSCI.5127-13.2014
González-Burgos, G., Krimer, L.S., Urban, N.N., Barrionuevo, G., Lewis, D.A., 2004. Synaptic Efficacy during
Repetitive Activation of Excitatory Inputs in Primate Dorsolateral Prefrontal Cortex. Cereb. Cortex 14,
530–542. https://doi.org/10.1093/cercor/bhh015
Hamm, J.P., Yuste, R., 2016. Somatostatin Interneurons Control a Key Component of Mismatch Negativity in
Mouse Visual Cortex. Cell Rep. 16, 597–604. https://doi.org/10.1016/j.celrep.2016.06.037
Harris, J.A., Mihalas, S., Hirokawa, K.E., Whitesell, J.D., Choi, H., Bernard, A., Bohn, P., Caldejon, S., Casal,
L., Cho, A., Feiner, A., Feng, D., Gaudreault, N., Gerfen, C.R., Graddis, N., Groblewski, P.A., Henry,
A.M., Ho, A., Howard, R., Knox, J.E., Kuan, L., Kuang, X., Lecoq, J., Lesnar, P., Li, Y., Luviano, J.,
McConoughey, S., Mortrud, M.T., Naeemi, M., Ng, L., Oh, S.W., Ouellette, B., Shen, E., Sorensen,
S.A., Wakeman, W., Wang, Q., Wang, Y., Williford, A., Phillips, J.W., Jones, A.R., Koch, C., Zeng, H.,
2019. Hierarchical organization of cortical and thalamic connectivity. Nature 575, 195–202.
https://doi.org/10.1038/s41586-019-1716-z
Henze, D.A., Borhegyi, Z., Csicsvari, J., Mamiya, A., Harris, K.D., Buzsáki, G., 2000. Intracellular Features
Predicted by Extracellular Recordings in the Hippocampus In Vivo. J. Neurophysiol. 84, 390–400.
https://doi.org/10.1152/jn.2000.84.1.390
Herrera, B., Westerberg, J.A., Schall, M.S., Maier, A., Woodman, G.F., Schall, J.D., Riera, J.J., 2022.
Resolving the mesoscopic missing link: Biophysical modeling of EEG from cortical columns in primates.
NeuroImage 263, 119593. https://doi.org/10.1016/j.neuroimage.2022.119593
Hosoya, T., Baccus, S.A., Meister, M., 2005. Dynamic predictive coding by the retina. Nature 436, 71–77.
https://doi.org/10.1038/nature03689
Hwang, J., Mitz, A.R., Murray, E.A., 2019. NIMH MonkeyLogic: Behavioral control and data acquisition in
MATLAB. J. Neurosci. Methods 323, 13–21. https://doi.org/10.1016/j.jneumeth.2019.05.002
Itti, L., Baldi, P., 2009. Bayesian surprise attracts human attention. Vision Res. 49, 1295–1306.
https://doi.org/10.1016/j.visres.2008.09.007
Jamali, S., Bagur, S., Brémont, E., Kerkoerle, T.V., Dehaene, S., Bathellier, B., 2024. Parallel mechanisms
signal a hierarchy of sequence structure violations in the auditory cortex.
https://doi.org/10.1101/2024.08.21.609026
Juavinett, A.L., Nauhaus, I., Garrett, M.E., Zhuang, J., Callaway, E.M., 2017. Automated identification of
mouse visual areas with intrinsic signal imaging. Nat. Protoc. 12, 32–43.
https://doi.org/10.1038/nprot.2016.158
Kajikawa, Y., Smiley, J.F., Schroeder, C.E., 2017. Primary Generators of Visually Evoked Field Potentials
Recorded in the Macaque Auditory Cortex. J. Neurosci. 37, 10139–10153.
https://doi.org/10.1523/JNEUROSCI.3800-16.2017
Keller, G.B., Mrsic-Flogel, T.D., 2018. Predictive Processing: A Canonical Cortical Computation. Neuron 100,
424–435. https://doi.org/10.1016/j.neuron.2018.10.003
Kiebel, S.J., Daunizeau, J., Friston, K.J., 2008. A Hierarchy of Time-Scales and the Brain. PLOS Comput. Biol.
4, e1000209. https://doi.org/10.1371/journal.pcbi.1000209
Kok, P., Rahnev, D., Jehee, J.F.M., Lau, H.C., de Lange, F.P., 2011. Attention Reverses the Effect of
Prediction in Silencing Sensory Signals. Cereb. Cortex 22, 2197–2206.
Logothetis, N.K., Pauls, J., Augath, M., Trinath, T., Oeltermann, A., 2001. Neurophysiological investigation of
the basis of the fMRI signal. Nature 412, 150–157. https://doi.org/10.1038/35084005
Maier, A., Adams, G., Aura, C., Leopold, D., 2010. Distinct Superficial and Deep Laminar Domains of Activity in
the Visual Cortex during Rest and Stimulation. Front. Syst. Neurosci. 4.
Maier, A., Aura, C.J., Leopold, D.A., 2011. Infragranular Sources of Sustained Local Field Potential Responses
in Macaque Primary Visual Cortex. J. Neurosci. 31, 1971–1980.
https://doi.org/10.1523/JNEUROSCI.5300-09.2011
Maier, A., Tsuchiya, N., 2021. Growing evidence for separate neural mechanisms for attention and
consciousness. Atten. Percept. Psychophys. 83, 558–576. https://doi.org/10.3758/s13414-020-02146-4

Maris, E., Oostenveld, R., 2007. Nonparametric statistical testing of EEG- and MEG-data. J. Neurosci. Methods 164, 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024

McCormick, D.A., Connors, B.W., Lighthall, J.W., Prince, D.A., 1985. Comparative electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. J. Neurophysiol. 54, 782–806. https://doi.org/10.1152/jn.1985.54.4.782

Mehta, A.D., Ulbert, I., Schroeder, C.E., 2000a. Intermodal selective attention in monkeys. I: distribution and timing of effects across visual areas. Cereb. Cortex 10, 343–358. https://doi.org/10.1093/cercor/10.4.343

Mehta, A.D., Ulbert, I., Schroeder, C.E., 2000b. Intermodal Selective Attention in Monkeys. II: Physiological Mechanisms of Modulation. Cereb. Cortex 10, 359–370. https://doi.org/10.1093/cercor/10.4.359

Mendoza-Halliday, D., Major, A.J., Lee, N., Lichtenfeld, M.J., Carlson, B., Mitchell, B., Meng, P.D., Xiong, Y.S., Westerberg, J.A., Jia, X., Johnston, K.D., Selvanayagam, J., Everling, S., Maier, A., Desimone, R., Miller, E.K., Bastos, A.M., 2024. A ubiquitous spectrolaminar motif of local field potential power across the primate cortex. Nat. Neurosci. 27, 547–560. https://doi.org/10.1038/s41593-023-01554-7

Michalareas, G., Vezoli, J., van Pelt, S., Schoffelen, J.-M., Kennedy, H., Fries, P., 2016. Alpha-Beta and Gamma Rhythms Subserve Feedback and Feedforward Influences among Human Visual Cortical Areas. Neuron 89, 384–397. https://doi.org/10.1016/j.neuron.2015.12.018

Miller, E.K., Cohen, J.D., 2001. An integrative theory of prefrontal cortex function. Annu. Rev. Neurosci. 24, 167–202. https://doi.org/10.1146/annurev.neuro.24.1.167

Miller, E.K., Lundqvist, M., Bastos, A.M., 2018. Working Memory 2.0. Neuron 100, 463–475. https://doi.org/10.1016/j.neuron.2018.09.023

Mitchell, J.F., Sundberg, K.A., Reynolds, J.H., 2007. Differential Attention-Dependent Response Modulation across Cell Classes in Macaque Visual Area V4. Neuron 55, 131–141. https://doi.org/10.1016/j.neuron.2007.06.018

Mitzdorf, U., 1985. Current source-density method and application in cat cerebral cortex: investigation of evoked potentials and EEG phenomena. Physiol. Rev. 65, 37–100. https://doi.org/10.1152/physrev.1985.65.1.37

Mitzdorf, U., Singer, W., 1978. Prominent excitatory pathways in the cat visual cortex (A 17 and A 18): A current source density analysis of electrically evoked potentials. Exp. Brain Res. 33, 371–394. https://doi.org/10.1007/BF00235560

Mumford, D., 1992. On the computational architecture of the neocortex. II. The role of cortico-cortical loops. Biol. Cybern. 66, 241–251. https://doi.org/10.1007/BF00198477

Nicholson, C., Freeman, J.A., 1975. Theory of current source-density analysis and determination of conductivity tensor for anuran cerebellum. J. Neurophysiol. 38, 356–368. https://doi.org/10.1152/jn.1975.38.2.356

Nicholson, C., Llinas, R., 1971. Field Potentials in Alligator Cerebellum and Theory of Their Relationship to Purkinje Cell Dendritic Spikes. J. Neurophysiol. 34, 509-+. https://doi.org/10.1152/jn.1971.34.4.509

Ninomiya, T., Dougherty, K., Godlove, D.C., Schall, J.D., Maier, A., 2015. Microcircuitry of agranular frontal cortex: contrasting laminar connectivity between occipital and frontal areas. J. Neurophysiol. 113, 3242–3255. https://doi.org/10.1152/jn.00624.2014

Nourski, K.V., Steinschneider, M., Rhone, A.E., Kawasaki, H., Howard, M.A., Banks, M.I., 2018. Auditory Predictive Coding across Awareness States under Anesthesia: An Intracranial Electrophysiology Study. J. Neurosci. 38, 8441–8452. https://doi.org/10.1523/JNEUROSCI.0967-18.2018

Nowak, L.G., Azouz, R., Sanchez-Vives, M.V., Gray, C.M., McCormick, D.A., 2003. Electrophysiological Classes of Cat Primary Visual Cortical Neurons In Vivo as Revealed by Quantitative Analyses. J. Neurophysiol. 89, 1541–1566. https://doi.org/10.1152/jn.00580.2002

Oostenveld, R., Fries, P., Maris, E., Schoffelen, J.-M., 2011. FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput. Intell. Neurosci. 2011, 156869. https://doi.org/10.1155/2011/156869

Pachitariu, M., Sridhar, S., Stringer, C., 2023. Solving the spike sorting problem with Kilosort. https://doi.org/10.1101/2023.01.07.523036

Pak, A., Kissinger, S.T., Chubykin, A.A., 2021. Impaired Adaptation and Laminar Processing of the Oddball Paradigm in the Primary Visual Cortex of Fmr1 KO Mouse. Front. Cell. Neurosci. 15. https://doi.org/10.3389/fncel.2021.668230

Povysheva, N.V., Gonzalez-Burgos, G., Zaitsev, A.V., Kröner, S., Barrionuevo, G., Lewis, D.A., Krimer, L.S., 2006. Properties of Excitatory Synaptic Responses in Fast-spiking Interneurons and Pyramidal Cells

from Monkey and Rat Prefrontal Cortex. Cereb. Cortex 16, 541–552.
https://doi.org/10.1093/cercor/bhj002
Rao, R. P. N., Ballard, D.H., 1999. Predictive coding in the visual cortex: a functional interpretation of some
extra-classical receptive-field effects. Nat. Neurosci. 2, 79–87. https://doi.org/10.1038/4580
Rao, R. P.N, Ballard, D.H., 1999. Predictive coding in the visual cortex: a functional interpretation of some
extra-classical receptive-field effects. Nat. Neurosci. 2, 79–87.
Rigotti, M., Barak, O., Warden, M.R., Wang, X.-J., Daw, N.D., Miller, E.K., Fusi, S., 2013. The importance of
mixed selectivity in complex cognitive tasks. Nature 497, 585–590. https://doi.org/10.1038/nature12160
Rübel, O., Tritt, A., Ly, R., Dichter, B.K., Ghosh, S., Niu, L., Baker, P., Soltesz, I., Ng, L., Svoboda, K., Frank,
L., Bouchard, K.E., 2022. The Neurodata Without Borders ecosystem for neurophysiological data
science. eLife 11, e78362. https://doi.org/10.7554/eLife.78362
Sanchez-Vives, M.V., Nowak, L.G., McCormick, D.A., 2000a. Membrane mechanisms underlying contrast
adaptation in cat area 17 *in vivo*. J. Neurosci. 20, 4267–4285.
Sanchez-Vives, M.V., Nowak, L.G., McCormick, D.A., 2000b. Cellular mechanisms of long-lasting adaptation in
visual cortical neurons *in vitro*. J. Neurosci. 20, 4286–4299.
Schroeder, C.E., Mehta, A.D., Givre, S.J., 1998. A spatiotemporal profile of visual system activation revealed
by current source density analysis in the awake macaque. Cereb. Cortex 8, 575–592.
https://doi.org/10.1093/cercor/8.7.575
Schultz, W., Dayan, P., Montague, P.R., 1997. A neural substrate of prediction and reward. Science 275,
1593–1599. https://doi.org/10.1126/science.275.5306.1593
Self, M.W., van Kerkoerle, T., Supèr, H., Roelfsema, P.R., 2013. Distinct Roles of the Cortical Layers of Area
V1 in Figure-Ground Segregation. Curr. Biol. 23, 2121–2129. https://doi.org/10.1016/j.cub.2013.09.013
Seth, A.K., Suzuki, K., Critchley, H.D., 2012. An interoceptive predictive coding model of conscious presence.
Front. Psychol. 3, 395. https://doi.org/10.3389/fpsyg.2011.00395
Siegle, J.H., Jia, X., Durand, S., Gale, S., Bennett, C., Graddis, N., Heller, G., Ramirez, T.K., Choi, H., Luviano,
J.A., Groblewski, P.A., Ahmed, R., Arkhipov, A., Bernard, A., Billeh, Y.N., Brown, D., Buice, M.A., Cain,
N., Caldejon, S., Casal, L., Cho, A., Chvilicek, M., Cox, T.C., Dai, K., Denman, D.J., de Vries, S.E.J.,
Dietzman, R., Esposito, L., Farrell, C., Feng, D., Galbraith, J., Garrett, M., Gelfand, E.C., Hancock, N.,
Harris, J.A., Howard, R., Hu, B., Hytnen, R., Iyer, R., Jessett, E., Johnson, K., Kato, I., Kiggins, J.,
Lambert, S., Lecoq, J., Ledochowitsch, P., Lee, J.H., Leon, A., Li, Y., Liang, E., Long, F., Mace, K.,
Melchior, J., Millman, D., Mollenkopf, T., Nayan, C., Ng, L., Ngo, K., Nguyen, T., Nicovich, P.R., North,
K., Ocker, G.K., Ollerenshaw, D., Oliver, M., Pachitariu, M., Perkins, J., Reding, M., Reid, D.,
Robertson, M., Ronellenfitch, K., Seid, S., Slaughterbeck, C., Stoecklin, M., Sullivan, D., Sutton, B.,
Swapp, J., Thompson, C., Turner, K., Wakeman, W., Whitesell, J.D., Williams, D., Williford, A., Young,
R., Zeng, H., Naylor, S., Phillips, J.W., Reid, R.C., Mihalas, S., Olsen, S.R., Koch, C., 2021. Survey of
spiking in the mouse visual system reveals functional hierarchy. Nature 592, 86–92.
https://doi.org/10.1038/s41586-020-03171-x
Siegle, J.H., López, A.C., Patel, Y.A., Abramov, K., Ohayon, S., Voigts, J., 2017. Open Ephys: an open-source,
plugin-based platform for multichannel electrophysiology. J. Neural Eng. 14, 045003.
https://doi.org/10.1088/1741-2552/aa5eea
Smout, C.A., Tang, M.F., Garrido, M.I., Mattingley, J.B., 2019. Attention promotes the neural encoding of
prediction errors. PLOS Biol. 17, e2006812. https://doi.org/10.1371/journal.pbio.2006812
Solomon, S.S., Tang, H., Sussman, E., Kohn, A., 2021. Limited Evidence for Sensory Prediction Error
Responses in Visual Cortex of Macaques and Humans. Cereb. CORTEX 31, 3136–3152.
https://doi.org/10.1093/cercor/bhab014
Spratling, M.W., 2010. Predictive coding as a model of response properties in cortical area V1. J. Neurosci.
Off. J. Soc. Neurosci. 30, 3531–3543. https://doi.org/10.1523/JNEUROSCI.4911-09.2010
Spratling, M.W., 2008. Reconciling predictive coding and biased competition models of cortical function. Front.
Comput. Neurosci. 2, 4. https://doi.org/10.3389/neuro.10.004.2008
Srinivasan, M.V., Laughlin, S.B., Dubs, A., 1982. Predictive coding: a fresh view of inhibition in the retina. Proc.
R. Soc. Lond. B Biol. Sci. 216, 427–459. https://doi.org/10.1098/rspb.1982.0085
Supèr, H., Roelfsema, P.R., 2005. Chronic multiunit recordings in behaving animals: advantages and
limitations. Prog. Brain Res., Development, Dynamics and Pathology of Neuronal Networks: from
Molecules to Functional Circuits 147, 263–282. https://doi.org/10.1016/S0079-6123(04)47020-4
Tasker, J.G., Hoffman, N.W., Kim, Y.I., Fisher, R.S., Peacock, W.J., Dudek, F.E., 1996. Electrical properties of
neocortical neurons in slices from children with intractable epilepsy. J. Neurophysiol. 75, 931–939.

901   https://doi.org/10.1152/jn.1996.75.2.931

902 Tovar, D.A., Westerberg, J.A., Cox, M.A., Dougherty, K., Carlson, T.A., Wallace, M.T., Maier, A., 2020.
903   Stimulus Feature-Specific Information Flow Along the Columnar Cortical Microcircuit Revealed by
904   Multivariate Laminar Spiking Analysis. Front. Syst. Neurosci. 14.

905 Tovar, D.A., Westerberg, J.A., Cox, M.A., Dougherty, K., Wallace, M.T., Bastos, A.M., Maier, A., 2023. Near-
906   field potentials index local neural computations more accurately than population spiking.
907   https://doi.org/10.1101/2023.05.11.540026

908 Trautmann, E.M., Stavisky, S.D., Lahiri, S., Ames, K.C., Kaufman, M.T., O'Shea, D.J., Vyas, S., Sun, X., Ryu,
909   S.I., Ganguli, S., Shenoy, K.V., 2019. Accurate Estimation of Neural Population Dynamics without
910   Spike Sorting. Neuron 103, 292-308.e4. https://doi.org/10.1016/j.neuron.2019.05.003

911 Uhrig, L., Janssen, D., Dehaene, S., Jarraya, B., 2016. Cerebral responses to local and global auditory novelty
912   under general anesthesia. NeuroImage 141, 326–340.
913   https://doi.org/10.1016/j.neuroimage.2016.08.004

914 van Kerkoerle, T., Self, M.W., Dagnino, B., Gariel-Mathis, M.-A., Poort, J., Togt, C. van der, Roelfsema, P.R.,
915   2014a. Alpha and gamma oscillations characterize feedback and feedforward processing in monkey
916   visual cortex. Proc. Natl. Acad. Sci. 111, 14332–14341. https://doi.org/10.1073/pnas.1402773111

917 van Kerkoerle, T., Self, M.W., Dagnino, B., Gariel-Mathis, M.-A., Poort, J., van der Togt, C., Roelfsema, P.R.,
918   2014b. Alpha and gamma oscillations characterize feedback and feedforward processing in monkey
919   visual cortex. Proc. Natl. Acad. Sci. U. S. A. 111, 14332–14341.
920   https://doi.org/10.1073/pnas.1402773111

921 van Kerkoerle, T., Self, M.W., Roelfsema, P.R., 2017. Layer-specificity in the effects of attention and working
922   memory on activity in primary visual cortex. Nat. Commun. 8, 13804.
923   https://doi.org/10.1038/ncomms13804

924 Vezoli, J., Vinck, M., Bosman, C.A., Bastos, A.M., Lewis, C.M., Kennedy, H., Fries, P., 2021. Brain rhythms
925   define distinct interaction networks with differential dependence on anatomy. Neuron S0896-
926   6273(21)00725-X. https://doi.org/10.1016/j.neuron.2021.09.052

927 Westerberg, J.A., Cox, M.A., Dougherty, K., Maier, A., 2019. V1 microcircuit dynamics: altered signal
928   propagation suggests intracortical origins for adaptation in response to visual repetition. J.
929   Neurophysiol. 121, 1938–1952. https://doi.org/10.1152/jn.00113.2019

930 Westerberg, J.A., Maier, A., Schall, J.D., 2020. Priming of Attentional Selection in Macaque Visual Cortex:
931   Feature-Based Facilitation and Location-Based Inhibition of Return. eNeuro 7.
932   https://doi.org/10.1523/ENEURO.0466-19.2020

933 Westerberg, J.A., Schall, J.D., Woodman, G.F., Maier, A., 2023. Feedforward attentional selection in sensory
934   cortex. Nat. Commun. 14, 5993. https://doi.org/10.1038/s41467-023-41745-1

935 Westerberg, J.A., Schall, M.S., Maier, A., Woodman, G.F., Schall, J.D., 2022. Laminar microcircuitry of visual
936   cortex producing attention-associated electric fields. eLife 11, e72139.
937   https://doi.org/10.7554/eLife.72139

938 Westerberg, J.A., Sigworth, E.A., Schall, J.D., Maier, A., 2021. Pop-out search instigates beta-gated feature
939   selectivity enhancement across V4 layers. Proc. Natl. Acad. Sci. 118.
940   https://doi.org/10.1073/pnas.2103702118

941 Xiong, Y. (Sophy), Donoghue, J.A., Lundqvist, M., Mahnke, M., Major, A.J., Brown, E.N., Miller, E.K., Bastos,
942   A.M., 2023. Propofol-mediated loss of consciousness disrupts predictive routing and local field phase
943   modulation of neural activity. https://doi.org/10.1101/2023.09.02.555990

944 Yan, Y., Zhaoping, L., Li, W., 2018. Bottom-up saliency and top-down learning in the primary visual cortex of
945   monkeys. Proc. Natl. Acad. Sci. U. S. A. 115, 10499–10504. https://doi.org/10.1073/pnas.1803854115
946

Figure 1. Predictive coding and the global-local oddball experimental paradigm

A. A stimulus enters the system through the excitatory neurons (gray triangles) in the middle input layers of cortex. That ascends the cortical column to the superficial layers (L2/3), excitatory then inhibitory cells (red circles), before reaching the deep layers (L5/6). The superficial excitatory neurons also innervate the next hierarchical stage of cortical processing (left to right) through the middle input layers. The intracolumnar processing motif then repeats. Crucially, feedback information (with respect to the expectations regarding the stimulus), is sent from deep layers to the superficial layers of the preceding area. The activation from area to area, layer to layer, and cell to cell can be computed from the equation at the top with the terms again indicated in the graphic as to where they are pertinent.

B. Visual stimuli were presented in 4-stimulus sequences consisting of 2-oriented, drifting bar grating, full-screen stimuli (symbolized as 'x'/'X' or 'y'/'Y' where capitalization indicates oddball status).

C. 16 transgenic mice were used in the experiment, genotypes were selected to allow for optotagging of either somatostatin- (magenta) or parvalbumin-positive (cyan) interneurons. Mice were split into 2 training groups, 1 group exposed to xxxY sequences during the entirety of training, 1 group to yyyX.

D. Neuropixels recording sessions were divided into 3 blocks. In the experiment block, mice experienced their trained 'Local' oddball sequence (xxxY or yyyX) for a 50-trial habituation period. Then, on 20% of the next ~500 sequences, they observed a never previously seen 'Global' oddball sequence (xxxX or yyyY). A block of random sequences of 4 stimuli followed the experiment block. A control block followed where alternating sequences of xxxx, yyyy were experienced.

E. Stimuli were presented for 500 ms with a 500 ms inter-stimulus interval and 1 s inter-trial intervals.

F. Neuropixels were introduced into 6 visual cortical regions in all mice (V1, LM, RL, AL, PM, AM).

G. Visual response latencies averaged across all single-units for each visual area followed the expected cascade given visual hierarchical position.

H. Population average response to the 4-stimulus sequences with the global oddball indicated in red and the local oddball in green.

Figure 2. Local oddball responses support experimental predictions of Predictive Coding.

A. Percentage of all single-units in an area where a significant detection of the local oddball was found using nonparametric cluster-based permutation testing (see Methods) at the single-unit level.

B. Early period of spiking activation averaged across units for hierarchical area categories (early, light; middle, medium; late, dark green) averaged across single units for a given area. Data area GLO(P4-P3) - CONTROL(P4-P3) demonstrating population enhanced neuronal activation to the local oddball. Detection is later and stronger in higher-order areas.

C. Average response of putative layer 2/3 V1 units to the local oddball (solid green) as compared to the non-layer 2/3 V1 units (dashed green) in the GLO block (P4-P3) and their equivalent responses in the control block (gray solid, gray dashed, respectively). Putative layer 2/3 V1 units have, on average, greater response to the local oddball than other V1 units.

D. Average response of optotagged (see Methods) inhibitory interneurons to the local oddball (green) in the GLO block (P4-P3) and their equivalent response in the control block (gray). Inhibitory interneurons are more active in the oddball condition.

Figure 3. Sparse global oddball detection across visual cortical hierarchy.

A. Global oddball detection across visual cortical areas. Lower-order areas to higher-order areas bottom to top. Clouds are 95% confidence intervals across units in an area. On left, red clouds are P4-P3 in GLO block, gray clouds are P4-P3 in control block. On right, 95% confidence interval clouds for the responses across units per area for the component presentations for the left. Top, GLO experiment with red P4 and gray P3. Bottom, control experiment with red P4 and gray P3. Yellow highlights in LM, PM, AM reflect periods of significant population global oddball detection with the latency indicated.

B. Percentage of all single-units in an area where a significant detection of the global oddball was found using nonparametric cluster-based permutation testing (see Methods) at the single-unit level.

C. Single-unit examples of units that were found to have significant global oddball detection in each of the 3 areas that showed significant global oddball detection at the population level. Bottom left of each subpanel shows the P4-P3 difference in the GLO block (red) and control block (gray) with the period of significant detection highlighted in yellow. Right of each subpanel shows the component responses in the GLO block (top) and control block (bottom) for P4 (red) and P3 (gray).

D. Global oddball detection for the areas found to have significant detection at the population level (LM, PM, AM) divided into lower-order (left, LM) and higher-order (right, PM, AM) categories with units further divided by layer. Layer 4 units defined to be in "granular" layer category (purple) and layer 2, 3, 5, and 6 units defined to be in extragranular category (orange). 95% confidence interval clouds plotted identical to panel (A) for P4-P3 differences in each block (red, GLO; gray, control). Yellow highlight indicates period of significant global oddball detection. Significant global oddball detection only present in extragranular layer populations.

003 Figure 4. Granger causality
004

A. Granger causality asymmetry (Feedforward minus feedback GC) between all possible areas in the baseline (pre-oddball time window).Green bars denote local oddballs, red bars denote global oddballs. Mean across mice, +/ SEM across mice.

B. Granger causality asymmetry (Feedforward minus feedback GC) between all possible areas in the initial sensory processing period (100ms to 300ms post-oddball onset).Green bars denote local oddballs, red bars denote global oddballs. Mean across mice, +/ SEM across mice. Asterisks indicate connections that are significantly different from baseline at P<0.01 (black asterisks) and at P<0.001 (blue asterisks).

C. Same as B, but for the late sensory processing period (300ms to 500ms post-oddball onset).

D. Hierarchical summary of significant changes in GC asymmetry from pre-oddball baseline for local oddballs. Significant interactions at both P<0.01 and P<0.001 are shown, and both early (B) and late (C) time periods are depicted. Global oddballs are not shown because there were no significant observed chances in GC asymmetry.

Figure 5. Synaptic activity is not indicative of top-down predictions being fed back to early sensory cortex.

A. Laminar CSD is calculated from the LFP response (top) to sequence presentations averaged across trials and across mice (n=9) for area V1 after aligning cortical depth between animals. CSD is calculated for both the global oddball experimental condition (middle, red highlighted region) and control condition (bottom, gray highlighted region). Epoch between white dotted lines indicates presentation 3, and between the black dotted lines indicates the global oddball presentation (middle) or the presentation 4 in the control (bottom).

B. Evaluating the global oddball contrast for evidence of differences in synaptic activity with global oddball detection in V1. Difference of differences identical to previous global oddball contrasts for the CSD data. Top image indicates all values following the contrast and normalization between -2 and +2 standard deviations. Bottom image shows the same data with a mask applied to only display significant values (Welch's t-test, p<0.01). Color map intensity indicated by scale bar in bottom right of figure.

C. Difference of differences for global oddball contrast for the remaining visual cortical areas. Formatting identical to B.

Figure 6. Restricting to putative layer 2/3 neurons does not reveal hidden feedforward global oddball signals.

A. Histogram of waveform durations of single-units. Threshold for narrow vs. broad spiking neurons set to 0.4 (orange dashed line). Units included as putative layer 2/3 pyramidal cells highlighted in orange, but further reduced by other criteria to obtain final population (see Results).

B. Lack of global oddball detection in layer 2/3 pyramidal cell subpopulation. Clouds are 95% confidence intervals across units. Top right, red cloud is P4-P3 in GLO block, gray cloud is P4-P3 in control block. Bottom, 95% confidence interval clouds for the responses across units for the component presentations for the top right. Left, GLO experiment with red P4 and gray P3. Right, control experiment with red P4 and gray P3.

C. Percentage of specifically putative layer 2/3 pyramidal single-units in an area where a significant detection of the global oddball was found using nonparametric cluster-based permutation testing (see Methods) at the single-unit level.

D. Single-unit example of a putative layer 2/3 unit that was found to have significant global oddball detection. Bottom shows P3 and P4 of the sequence response in the GLO block (red) and control block (gray) with the period of significant detection highlighted in yellow. Above P4 shows the P4-P3 difference for the GLO block (red) vs. the control block (gray) with the yellow highlight reflecting the period of significant detection.

Figure 7. Somatostatin- and parvalbumin-positive interneurons do not support Predictive Coding of global oddballs.

A. Response of optotagged SST- (left, magenta) and PV- (right, cyan) units as compared to non-optotagged units (black) to 40 Hz laser stimulation in the high power condition (level 1.35) during the optotagging block.
B. Relative proportion of each inhibitory interneuron type across areas (left) and layers (right).
C. Population average response to the oddball sequences for both interneuron types. P1 to P3 in black, local oddball in green, global oddball in red.
D. Percentage of either SST- (left) or PV- (right) tagged interneurons where a significant detection of the global oddball was found using nonparametric cluster-based permutation testing (see Methods) at the single-unit level. Also, example single-unit responses during the sequence presentation (specifically presentations 3 (P3) and 4 (P4)) in the experimental (red) or control (gray) blocks (below) as well as the differences (P4-P3) for those blocks (above). Period of significant difference highlighted in yellow.
E. Lack of global oddball detection in SST and PV cell subpopulations. Clouds are 95% confidence intervals across units. Top, red cloud is P4-P3 in GLO block, gray cloud is P4-P3 in control block. Middle and bottom, 95% confidence interval clouds for the responses across units for the component presentations for the top. Middle, GLO experiment with red P4 and gray P3. Bottom, control experiment with red P4 and gray P3.

Figure 8. Distinct profiles of local vs. global oddball detection across cortical hierarchy in monkeys.

A. Macaque monkeys (n=2) viewed sequences of drifting grating stimuli while fixating centrally. Sequences were temporally organized identically to the mouse experiment.

B. Recordings were performed across 8 brain areas at different stages of the visual cortical processing hierarchy (right). Multi-unit spiking responses were recorded at different sites across multiple recording sessions (n=32). Responses are shown as 95% confidence intervals after performing the same correction as with the mouse data: Green, local oddball - stimulus response prior; Red, global oddball - stimulus previous; Gray, local/global stimulus control - stimulus previous. Yellow highlights indicate periods of significant enhanced oddball activation at the population level.