A multi-omics census reveals obesity-associated microRNA *miR-let-7* as novel instigator of

adipose mitochondrial dysfunction and of intergenerational metabolic decline.

Chien Huang[1,2#], Joo Hyun Park[1#], Ali Altintas[3], Natasa Stanic[1], Hande Topel Batarlar[1], Rocio

Valdebenito Malmros[1], Jesper Havelund[1], Nils Joakim Færgeman[1], Bjørk Ditlev Marcher Larsen[1],

Karolina Szczepanowska[4,5], Jan-Wilm Lackmann[4], Aleksandra Trifunovic[4], Eva Kildall Hejbøl[6], Sönke

Detlefsen[6,7], Ida Engberg Jepsen[8], Stefanie Hansborg Kolstrup[9], Ricardo Laguna-Barraza[10], Javier-

Martin Gonzalez[10], Konstantin Khodosevich[11], Romain Barrès[3] and Jan-Wilhelm Kornfeld[1,12*]

[1]Department of Biochemistry and Molecular Biology (BMB), University of Southern Denmark (SDU),

Odense, Denmark. [2]Laboratory of Animal Physiology, Department of Animal Science and Technology,

National Taiwan University, Taipei, Taiwan. [3]Novo Nordisk Foundation Center for Basic Metabolic

Research (CBMR), Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen,

Denmark. [4]Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases

(CECAD) & Center for Molecular Medicine (CMMC), University of Cologne, Cologne, Germany.

[5]IMol Polish Academy of Sciences, Warsaw, Poland. [6]Department of Pathology, Odense University

Hospital, Odense, Denmark. [7]Department of Clinical Research, Faculty of Health Science, University

of Southern Denmark, Odense, Denmark. [8]The Fertility Clinic, Department of Obstetrics and

Gynecology, Zealand University Hospital, Koege, Denmark. [9]Biomedical Laboratory, Institute of

Molecular Medicine, University of Southern Denmark, Odense, Denmark. [10]Core Facility for

Transgenic Mice, Department of Experimental Medicine, Faculty of Health and Medical Sciences,

University of Copenhagen, Copenhagen, Denmark. [11]Biotech Research & Innovation Centre (BRIC),

University of Copenhagen, Copenhagen, Denmark. [12]Novo Nordisk Foundation Center for Adipocyte

Signaling, University of Southern Denmark, Odense, Denmark.

#These authors contributed equally to this work,

*Corresponding author

Abstract

We here describe that obesity and weight loss in male mice causes reversible abnormalities in glucose and

lipid metabolism, serum metabolomes and reduced expression of genes controlling mitochondrial energy

dissipation in white adipose tissue. When mating obese male mice with lean females, we observed concordant

reductions in adipose mitochondrial gene expression and translation in offspring (F_1) that resemble those

observed in the paternal (F_0) generation. When mapping miRNA differential expression across somatic

organs and the germline (i.e., liver, adipose tissue, sperm) and generations, we found that obesity and weight

loss reversible affected miRNA levels, and that *miR-let7* isoforms were induced in adipose tissues of obese

F_0 and F_1 adipose tissue and sperm of obese F_0 mice. Overexpressing *miR-let-7* in adipocytes silenced

DICER1, a cellular rheostat required for adipose tissue adaptation in obesity as evidenced by deficiencies in

mitochondrial function following DICER1 loss in adipocytes. Delivery of *miR-let-7* into lean oocytes elicited

glucose intolerance and impediments in adipose mitochondrial gene expression in mice sired from miRNA-

injected embryos, thus phenocopying important aspects of obesity heredity. When performing single-cell

RNA-Seq of miRNA-injected embryos, *miR-let7* impaired mitochondrial gene expression, suggesting altered

energy metabolism following alterations in zygotic miRNAs. When studying miRNA alterations in human

semen, lifestyle-induced weight loss downregulated *MIR-LET-7* in human subjects, suggesting similar roles

for human *MIR-LET-7* in gametic epigenomes and embryogenesis.

Introduction:

Intergenerational epigenetic inheritance of metabolic dysfunction

Genome-wide association studies have demonstrated that genetic (i.e., DNA) variants only explain 20% of obesity heritability[1]. Given the pandemic rise in obesity within evolutionarily short timescales, non-genetic or ‚*epigenetic*' processes, i.e., processes spurred by obesogenic lifestyles, increase disease susceptibility and could contribute to obesity development. This type of epigenetic inheritance is particuarly intriguing for paternal programming where fathers contribute only sperm as biological material to fertilized embryos (‚intergenerational inheritance') or no genetic/biological material at all (‚transgenerational inheritance'). Epidemiological support for the existence of hereditary, paternally-acquired traits comes from historical records demonstrating that ample ancestral food supplies affects cardiometabolic traits for up to two generations[2,3], showing that individuals carry increased risks of cardiometabolic dysfunction if their parents or grandparents were obese[4]. In line with these correlative observations in human, mechanistic studies in isogenic mice demonstrated that paternal obesity impairs metabolism in offsprings (i.e., F_1 generation and beyond) and, for instance, low-protein[5] or high-fat diet feeding[6-8], depression-like behavior[9], fear-conditioning[10] and psychological traumata[11] all impart epigenetic behavioral and metabolic effects in offsprings. Recently, also beneficial types of epigenetic programming were shown, and rodent offspring conceived following cold temperature exposure display increased amounts of metabolically favorable brown fat[12], suggesting a high plasticity of the molecular and cellular mechanisms underlying intergenerational epigenetic responses.

The molecular basis for epigenetic inheritance of paternal obesity in male gametes

The molecular and cellular basis for epigenetic heredity remains enigmatic: Although changes in epigenetic marks such as DNA methylation (DNAme), histone post-translational (PTM) modifications and small Noncoding RNAs (sncRNAs) all control epigenetic processes in male gametes[13], and despite lifestyle/diet-induced changes in DNAme and histone PTMs being reported, evidence for causatively implicating these marks in eliciting paternal response in progeny is rare for higher organism like rodents and humans. Other instigators of intergenerational response are found amongst the different types of

gametic sncRNAs like microRNAs (miRNAs): miRNAs were initially described in nematodes and fruit

flies[14-17] and sncRNAs-evoked gametic gene silencing is well-understood [18]. Due to their mobile nature

and microvesicular exchange between cell types[19], high abundance in gametes and dynamic changes in

expression following exposure to diet[20,21], sperm-derived sncRNAs have the potential for transmitting

lifestyle choices to zygotes and progeny. Consistent with this, metabolic trait-associated miRNAs affect

intergenerational phenotypes and phenocopy parts of F_0 lifestyles if microinjected [22,23] or removed from

zygotes[24]. Beyond controlling gametic epigenomes, miRNAs are essential for germ cell development

processes like spermatogenesis[25] and embryogenesis[26], and miRNA can be exchanges between zygotes

and other cell types [11,23,27-29]. Taken together, it is plausible to hypothesize that obesity-evoked changes

in sperm miRNAs alter zygotic miRNA pools and embryonic development, instigating epigenetic

phenotypic traits and somatic gene expression changes in offsprings. As diverse paternal lifestyle

exposures ranging from inflammation[14], nutrition[27,29] to stress [11,28] alter gametic miRNA repertoires and

F_1 phenotypes in non-overlapping ways, the intriguing possibility exists that environmental experiences

are encoded in individual gametic sncRNA signature and that these sperm sncRNAs have the potential

to transmit complex lifestyle exposures given the high bandwidth of RNA-mediated effects[30].

Regulation of adipose tissue (mitochondrial) function by miRNAs

Adipose tissue is a multifunctional endocrine organ that coordinates tissue-level (autocrine, paracrine)

and system-wide (endocrine) responses by release of adipokines, lipids and inflammatory factors and,

shown recently, microvesicles (exosomes)[31-33]. Adipose tissue can safely store ingested nutrients as

lipids, although long-term calorie excess causes insulin resistance[34], aberrant gene expression in

adipocytes[35], vascular remodeling and immune cell infiltration[31]. Mitochondria in adipose tissue exert

important functions for tissue/energy homeostasis, particularly during nutrient deprivation/excess[33]:

Whereas brown adipose tissue (BAT) contains large numbers of mitochondria to generate heat by

dissipation of the electron transport chain (ETC) proton gradients[36], white adipose tissue (WAT)

protects from calorie overload by removing nutrients from circulation and storing as energetic

macromolecules. Mitochondrial function in WAT is impaired in obesity [37] and reduced mitochondrial

function in fat itself causes metabolic imbalance[38-41]. Mitochondrial DNA content, oxidative

phosphorylation (OxPhos), abundance and composition of ETC complexes are functionally impaired in

obesity and morphological aberrations are found in obesity[37,41] and aging[42]. Conversely, treatment with

AMP-activated protein kinase (AMPK) agonist Metformin improves mitochondrial function in adipose

tissue, as do exercise or calorie restriction, and these interventions correlate with metabolic

improvement[43]. MiRNAs are important regulators of mitochondrial function and *miR-143, miR-181a,*

miR-378 instigate, whereas *miR-532* negatively affects mitochondrial function by post-transcriptionally

silencing mitochondrial translocation and RNA homeostasis[44-46]. We and others showed that removing

miRNAs from adipocytes by ablating DICER1, a central component of the conserved miRNA

processing machinery, impairs adipogenesis, thus triggering lipodystrophy and metabolic dysfunction

in mice[47-49].

Collectively, we here report that obesity in the male mice is linked to reversible alterations in adipose,

but not liver, mitochondrial gene regulation that are inherited to adipose tissue of F_1 progeny. We

mechanistically implicate obesity-associated *miR-Let-7*-driven suppression of DICER1 and

concomitant declines in mitochondrial function in adipocytes, and established sperm-derived *miR-Let-*

7 as intergenerational driver of mitochondrial dysfunction and glucose impairments in F_1 offspring.

Results

Paternal obesity and weight loss elicit reversible changes in mitochondrial metabolites

The epigenetic and phenotypical consequences of altering paternal metabolism by experimental high-fat diet feeding have been documented, yet the plasticity of hereditary dietary effects remains elusive. To investigate the potential reversibility of metabolic dysfunction in the paternal (F_0) and offspring (F_1) generation, we used diet-induced obese (DIO) 6-week-old male C57BL/6J mice as rodent model. We fed animals with i. low-fat diet (F_0 LF) ii. micronutrient-matched high-fat diet (F_0 HF) for 18 weeks or iii. fed HF for 9 weeks, followed by 9 weeks of LF-induced weight loss feeding (F_0 HF-LF), and F_1 progeny sired from each group received LF to specifically interrogate the hereditary sequelæ of F_0 metabolism (**Fig.1a**). We found that HF-LF-evoked weight loss corrected HF-induced body weight gains (**Fig.1b**), glucose intolerance (**Fig.1c,d**), insulin resistance (**Fig.1e**), and reduced serum glucose (**Fig.1f**), insulin (**Fig.1g**), and leptin levels (**Fig.1h**). We examined organ/body weight ratios to evaluate tissue-level effects of weight gain and regression, and found that weight loss corrected the HF-induced increase in epididymal WAT (eWAT) and inguinal WAT (iWAT) mass, yet did not affect liver and BAT to body weight ratios (**Fig.1i**). To gain system-level insights into the effects of obesity and weight loss, we performed i. serum metabolomics ii. serum lipidomics, iii. transcriptomics (mRNAseq and small RNAseq, sRNAseq) iv. proteomics and v. histological profiling in liver and eWAT of F_0 groups (**Fig.1j**). By metabolomics, we detected 600 unique features (**Table S1**) that segregated animals into three groups (**Fig.1k**). When focussing on metabolites that were (re)-reversibly affected by weight loss (**Fig.1l**), we found that mitochondria-associated metabolites such as the tricarbon acid (TCA) cycle intermediates citrate, malate, fumarate, alpha-ketoglutarate and citrate, were reduced by obesity and restored by weight loss (**Fig.1m**, **Fig.S1a,b**). Using lipidomics, we detected 117 lipid species (**Table S2**) that correlated with body weight trajectories and were able to segragated F_0 groups (**Fig.S2a,b**). Amongst these, we observed that saturated, short-chain ceramides, i.e., inhibitors of mitochondrial FA oxidation in adipose tissue[50], were reversibly induced, whereas more indolent, long-chain, unsaturated ceramides were reversibly downregulated by HF feeding (**Fig.S2c**). *Thus, dietary obesity and weight loss cause reversible alterations in energy metabolism in mice that correlate with serum metabolites that are instigators of tissue-level mitochondrial dysfunction.*

Obesity triggers reversible mitochondrial mRNA-miRNA gene alteration in adipose tissue.

Beyond instigating mitochondrial dysfunction at the systemic metabolite level, obesity and weight loss reversibly increased eWAT adipocyte size (**Fig.2a,b**) and prompted Metabolic Dysfunction Associated Liver Disease (MASLD) hallmarks such as micro- and macrovasicular steatosis, fibrosis and lobular inflammation (**Fig.S3a-e**). Given the pronounced lipid accumulation in adipose tissue and liver, we next asked if the observed reduction in serum mitochondrial metabolites in obese mice coincided with tissue-level mitochondrial dysfunction in these organs, and performed global transcriptomic profiling in liver and eWAT of F_0 groups. Analysis of liver mRNAseq and group-level plotting using multidimensional scaling (MDS) showed that the F_0 LF group segregated from others (**Fig.S3f**). When comparing F_0 HF vs F_0 LF, we found 3,495 up-regulated and 1,639 down-regulated genes (**Fig.S3g**), mirroring the effects observed when comparing F_0 HF-LF vs F_0 LF that yielded 3,095 up-regulated and 955 down-regulated genes (**Fig.S3h**). Conversely, comparing F_0 HF vs F_0 HF-LF, yielded only 808 up-regulated and 960 down-regulated genes (**Fig.S3i**), highlighting that the expression changes triggered by HF feeding in liver were to large parts irreversible despite the profound weight loss (**Table S3**). Gene Set Enrichment (GSEA)[51] analyses of differentially expressed genes (DEG) revealed upregulated processes such as lipid metabolism and organelle formation, whereas repressed processes include lipoprotein trafficking, endoplasmatic reticulum and ribosome function (**Fig.S3j**), yet we observed no transcriptional evidence of altered mitochondrial processes. Next, we turned our attention to adipose tissue and performed RNAseq followed by DEG analysis (**Table S4)**. MDS analysis showed transcriptional segregation of diet groups (**Fig.2c, Fig.S4a**) and, comparing F_0 HF vs F_0 LF, we found 2,337 up-regulated and 3,867 down-regulated genes (**Fig.2d**). These effects that were to a large degree reversible, as evidenced by analysing the comparison between F_0 HF vs F_0 HF-LF that yielded 1,986 up-regulated and 2,858 down-regulated genes (**Fig.2e**), whereas the comparision F_0 LF and F_0 HF-LF mice only yielded 262 up-regulated and 459 down-regulated genes (**Fig.2f**). When performing GSEA on DEG we found gene ontology (GO) processes related to cell proliferation and immune activation; known transcriptional processes reflecting adipocyte hyperplasia and inflammation associated with obesity. Importantly, we also detected GO terms related to lipid catabolism, mitochondrial ATP synthesis and respiration and

ETC downregulated in obese adipose tissue (**Fig.2g**). When specifically interrogating the expression of

mRNA encoding ETC Complexes I-V, we confirmed the pervasive repression of nuclear-expressed

ETC mRNAs (**Fig.S4b**).

The down-regulation of mitochondrial programs in obese eWAT at least prompted the question whether

mitochondria-associated ,mito-miRs[1]' like *miR-let-7*[2], *miR-378*[3], *miR-181*[4], *miR-143*[5] that are known to

support mitochondrial formation and bioenergetics, contribute to these effects. When thus conducted

small RNA-Seq (sRNAseq, **Table S5**) and found profound changes between F_0 adipose miRNomes

(**Fig.2h**). As expected, mito-miRs promoting mitochondrial biogenesis and function, for instance *miR-*

143-3p, *miR-181a-2-3p*, *miR-378a-3p* and *miR-378* where (reversibly) repressed, whilst mitochondria-

inactivating *miR-532* was reversibly induced in F_0 HF eWAT (**Fig.2i**), suggesting that miRNA

deregulation contributed to the observed mitochondrial impediments observed in obese fat. Integration

of our miRNA-mRNA datasets with miRNA-mRNA interactomes from fat-specific AGO-HITS-CLIP[6]

confirmed that HF-associated mito-miRs might be an important driver of the mitochondria dysfunction

observed in obese eWAT (**Fig.2j**). *Thus, obesity caused a pervasive, reversible repression of*

mitochondrial ETC genes, FA oxidation and OxPhos, and miRNA-mRNA network analysis implicates

obesity-repressed mito-miRs in this process.

Paternal obesity causes concordant mitochondrial changes in offspring adipose tissue.

Unresolved questions in the field of epigenetic phenotypic inheritance revolve around the 'bandwidth'

and 'plasticity' of heritable intergenerational responses: Whilst the former tries to answer if paternal

hereditary processes like fear conditioning[7], temperature homeostasis[8], and adipose activation can elicit

concordant molecular, metabolic and behavioral alterations in offsprings [9], 'plasticity' refers to the

reversibility of paternally acquired and inherited traits. To address both points, we next mated lean,

obese and weight-regressed male mice with lean virgin female mice. Following mating, female mice

gave risen to three offspring generations: F_1 of F_0 LF (n=44), F_1 of F_0 HF (n=52), and F_1 of F_0 HF-LF

(n=38) and average litter sizes (7.33, 5.7, 6.33, respectively) and rate of successful matings (100%,

75%, 100%, respectively) were not different between groups whilst gender ratios differed (i.e., 58.7%,

57.7%, 45.5% being female mice, respectively).

To interrogate physiological and molecular effects of ancestral diet feeding without confounding HF

challenges, F_1 offsprings groups were fed only LF; and body weight in F_1 groups did not differ (**Fig.3a**).

In contrast, progeny sired by obese father exhibited glucose intolerance (**Fig.3b,c**) and impaired insulin

sensitivity (**Fig.3d**) despite similar random-fed glycemia and serum insulin levels (**Fig.3e,f**).

Intriguingly, offsprings of fathers undergoing pre-conception weight loss (i.e., the F_1 [F_0 HF-LF] group)

did not exhibit metabolic aberrations, highlighting that intergenerational responses are plastic and thus

reflect dynamic changes in paternal lifestyle and metabolic improvement. Concurrently with aberrations

in glucose metabolism, F_1 [F_0 HF-LF] mice had higher serum leptin adipokine levels (**Fig.3g**), trended

towards increased fat mass (**Fig.3h**), and exhibited higher relative eWAT tissue weight (**Fig.3i**), and

adipocyte hypertrophy (**Fig.3j,k**), ressembling the adipose effects seen in F_0 HF mice. Prompted by the

reversible eWAT-hypertrophy in F_1 mice, we next performed RNAseq in eWAT of F_1 groups: Although

calculation DEG of transcriptional changes using similar statistical stringencies as in F_0 yielded only

few DEG (**Fig.S5a**), GSEA demonstrated reversible induction in germ cells-associated processes,

whereas mitochondrial processes were reversibly repressed in eWAT of F_1 [F_0 HF] mice (**Fig.3l**) and

similar patterns of ETC complex I-V (CI-IV) gene repression were observed as in F_0, albeit with less

robust statistical significance and smaller effect sizes (**Fig.S5b**), yet qualitatively mimicking gene

changes observed in the F_0 HF-LF fathers themselves. To interrogate if these transcriptional changes

caused biologically meaningful changes in mitochondrial proteins, we conducted bulk proteomics by

mass-spectrometry in eWAT of F_0 and F_1 groups (**Fig.3m**) and found an enrichment of mitochondrial

GO processes in downregulated cluster II (**Fig.S7a**), validating repression of important ETC proteins

NDUFA6, COX6B1 (**Fig.3n**) and NDUFA8, NDUFS6, and mitochondrial FA carrier proteins like

Carnithin Palmitoyltransferase 1 (CPT1, **Fig.S5c**) in F_0 HF and F_1 [F_0 HF-LF] eWAT. Of note, mRNA-

seq experiments in F_1 livers did not yield similar transcriptional (i.e. mitochondria-associated, reversible

by weight loss) and morphological (HF-induced cellular hypertrophy) sequelæ of paternal obesity

(**Fig.S6a-j**), emphasizing the specificity of paternal programming and metabolic and mitochondrial

decline for epididymal adipose tissue. *Thus, we here demonstrate decreased glucose homeostasis and*

adipose hypertrophy in mice sired from obese fathers which was corrected by paternal pre-conception

weight loss, and that correlated with hereditary patterns of mitochondrial dysfunction in adipose tissue.

Paternal obesity reversibly affects the germeline miRNome and induces miR-let-7 in F_0 and F_1.

One particularly abundant molecular entity in male germ cells (spermatozoa) are various sncRNAs such

as miRNAs, transfer RNA (tRNA) fragments[24] and, recently, mitochondrial mt-RNA-derived cleavage

products[21]. As small RNAs are subject to dietary cues in humans[20], it is conceivable that diet can alters

spermatic miRNAs which, following fertilisation, are transferred into oocytes and affect zygotic and

embryonic development, thereby contributing to the altered phenotypic traits and gene expression we

observed in offspring from obese fathers.

To test the effects of obesity and weight gain on male gametic miRNAs, we next isolated motile sperm

by swim-up assays, and mapped F_0 sperm as well as F_0 and F_1 eWAT miRNA alterations by sRNA-Seq

(**Fig.4a,Fig.S7b**). Amongst miRNAs induced in F_1 eWAT, we did not observe mito-miR deregulation

as for F_0 eWAT (**Fig.S7c**) argueing against their causatively involvement in heritable mitochondrial

dysfunction. In contrast, by focussing on miRNAs that were concurrently regulated in F_0 and F_1, we

detected *miR-let-7* family members induced in F_0 sperm, and F_0 and F_1 eWAT (**Fig.4b,c**;**Fig.S7d**). *miR-*

let-7 has been described in intergenerational obesity studies by us [7] and others[52], and is known to impair

embryo implantation and glucose homeostasis when overexpressed [53,54]. Amongst *miR-let-7* isoforms,

miR-let-7d-5p and *miR-let-7e-5p* were consistently induced across generations (F_0, F_1) and in spermand

eWAT (**Fig.4c**), so chose to functionally dissect their role in sperm and eWAT.

miR-let-7i has also been described in the context of adipocyte differentiation[55], yet its roles in

mitochondrial homeostasis of mature adipocytes are unreported . To mimic the obesity-associated

increased in *miR-let-7d/e-5p in vitro*, we delivered mature miRNA mimetics by reverse transfection of

stromal vascular fraction (SVF)-derived epididymal white adipocytes (1°eWA, **Fig.4d,e**). When we

interrogated the effects of ectopic *miR-let-7* delivery by mRNAseq, we found that both *miR-Let-7*

isoforms triggered similar transcriptional effects 1°eWA as expected owing to their similar nucleotide sequence (**Fig.S8a**). Compared to Negative Control (NC) oligonucleotides, *miR-let-7d-5p* caused more pronounced changes (977 down-regulated, 1,301 up-regulated DEG) than *miR-let-7e-5p* (759 down-regulated, 635 up-regulated DEG) and both downregulated GO processes related to lipid metabolism, Insulin-AKT Serine/Threonine Kinase (AKT) signaling and carbohydrate metabolism (**Fig.4f,g, Fig.S8b**). To test if these gene expression changes also affect mitochondrial function, we predicted *miR-let-7* target gene networks using AGO-HITS-CLIP datasets [56]and found that *miR-let-7* induction in eWAT was linked to repression of several MitoCarta 3.0-annotated mitochondrial genes (**Fig.4h,Table S8**). Amongst a wider range of predicted targets, we validated *miR-let-7* mRNA targets linked to mitochondrial FA uptake (*Slc25a24*), glycolysis (*Hk2*, *Pdk2*) and miRNA processing (*Dicer1*) using dual luciferase reporter assays (DLRA)-based miRNA-mRNA interaction studies in HEK293 cells (**Fig.4i**). We found that *Dicer1* 3'UTRs harboured two independent *miR-let-7* seeds (**Fig.4h**) that, upon mutation, became refractory to *miR-let-7* repression (**Fig.4j**). As 1°eWA overexpressing *miR-let-7d/e-5p* reduced DICER1 protein, but only caused mild reductions in mitochondrial/TCA-associated protein itself (**Fig.4k,Fig.S8c,d**), we hypothesized that the chronic *miR-let-*7 induction seen in obesity might repress mitochondrial function via sustained DICER1 loss; and that *Dicer1* silencing is a better proxy for mimicking the prolonged *miR-let-7d/e-5p* induction in obesity. To test if *Dicer1* knockdown can phenocopy the obesity-induced impairment in adipocyte mitochondrial function, we next designed two *Dicer1*-targeting locked nucleic acid (LNA) oligonucleotide inhibitors that repressed *Dicer1* (**Fig.S8e**) and protein (**Fig.S8f**) and interrogated mitochondrial function using Seahorse bioenergetics in *Dicer1*-deficient 1°eWA. We found that *Dicer1* silencing repressed oxygen consumption rates (OCR, **Fig.4k**) as proxy for OxPhos activity, and reduced extracellular acidification rates (ECAR) which reflects lactate production as proxy for glycolytic rates in 1°eWA (**Fig.4l**). *Thus, paternal obesity changes miRNA expression in germline and somatic tissues and obesity-associated miR-let-7 impairs mitochondrial oxidative respiration by silencing of DICER1 in mature adipocytes.*

Zygotic delivery of miR-Let-7 impairs glucose tolerance and adipose mitochondrial gene expression in sired offsprings.

Prompted by the profound, yet reversible effects of obesity and weight loss on sperm miRNAs, we next

aimed to test the functional relevance of sperm-evoked *miR-let-7* changes for offspring phenotypes, as

environmentally induced and artificial alterations in sperm and zygote miRNA pools were previously

shown to trigger metabolic abnormalities in progeny[23,57]. We focussed on *miR-let7d/e-5p* given their

concordant increase in F_0 and F_1 eWAT as well as in F_0 sperm, and because *miR-let-7* is essential for

embryonic fate decisions[58] which partially dependent of post-transcriptional silencing of *Dicer1*[59]. To

test if zygotic delivery of *miR-let-7* in itself triggers phenotypical changes in F_1 that resemble those seen

as consequence of paternal F_0 obesity, we injected 50 femtomol of Negative Control (NC)

oligonucleotides, miRNA mimetics encoding *mmu-miR-let7-d/-e-5p* and *mmu-miR-699-5p* and *miR-*

699a-5p, a miRNA that was expressed at similar levels as *miR-let7-d* in sperm, but not regulated by HF

feeding, into zygotes conceived from lean males and females using public protocols[27] (**Fig.5a**). We

injected 126/138/110/106 zygotes with NC, *miR-let-7d-5p*, *miR-let-7e-5p* and *miR-699a-5p*,

respectively, and observed similar implantation rates, litter sizes and successful pregnancies (**Table S9**).

When analysing at least 19 pups per group from at least four pregnancies per group, we observed no

differences in BW(**Fig.5b**), relative eWAT (**Fig.5c**) and liver weights (**Fig.5d**) between F_m groups.

When interrogating random-fed glucose levels, we observed significant increases in progeny sired from

miR-let-7d-5p, *miR-let-7e-5p* and *miR-699a-5p*-injected embryos (**Fig.5e**). Importantly, we observed

impaired glucose tolerance only in mice sired from *miR-let-7d/e-5p*-injected embryos (**Fig.5f-h**) and

mRNA-seq profiling in eWAT, despite mild DEG changes, revealed down-regulation of mitochondrial

GO processes in these mice. *Thus, zygotic microinjection of miR-let-7 isoforms that accumulate in obese*

sperm, causes glucose intolerance and impairment of mitochondrial gene regulation in mice derived

from these embryos.

Obesity-associated miR-let-7 rewires mitochondrial metabolism during embryogenesis.

Intergenerational epigenetic programming in isogenic rodent background has been reported before, yet

it is elusive if these effects are translatable to species with different reproductive apparatuses, genetic

backgrounds, and timescales of obesity development such as humans. Although epidemiological

evidence[3,60] suggests heritable effects in humans ('Dutch Hunger Winter'[61]), direct epigenetic effects

are difficult to delineate. It is becoming appreciated though, that human sperm is sensitive to nutritional

changes, and that nutritional and metabolic cues are sufficient to shape human sncRNA expression[20].

To test the effect of voluntary weight loss in semen in a cohort of (n=15) morbidly obese (BMI, 39.49

± 1.80) men undergoing voluntary weight loss[62], we quantified *HSA-MIR-LET7D/E* using TaqMan

assays. We found that changes in *HSA-MIR-LET7D/E* correlated with weight loss (ΔBMI), suggesting

similar correlations between metabolic status and MIR-LET-7D/E in human and mice (**Fig.6a,b**).

As miRNA pertubation in sperm i. gives rise to incompletely developed embryos, suggesting roles for

sperm-derived miRNAs in embryonic development [19,63,64] and ii. zygotes can differentiate potential into

virtually any cell type, early zygotic / embryogenesis represents an important window of developmental

plasticity that might affect offspring phenotypes if perturbed. It is thus tenable that alterations in sperm

miRNAs change zygotic miRNA pools, which will have far-reaching consequences causing changes to

epigenetic and metabolic pathways in embryos[64]. Based on the observd mild but specific mitochondrial

alterations in F_m mice, we hypothesized that *miR-let7d/e* might inflict gene alterations during cell fate

decisions during embryo development, we repeated the miRNA injections, allowed miRNA-injected

embryos to divide until the 8-cell stage, dissociated embryos into single blastomeres, and performed

using custom made single-cell RNA-Seq using Smart-seq2 protocols[65] (**Fig.6c**). We found that, after

three cell divisions, the miRNA-injected blastomers were still transcriptionally distinct, reflecting the

profound transcriptional and developmental effects of perturbing miRNA in zygotes (**Fig.6d**). When

performing GSEA analysis of miRNA-induced DEG genesets, we found that *miR-let-7d* repressed GO

terms linked to mitochondrial processes such as ATP synthesis, TCA cycle and ETC/CI biogenesis.

Thus*, zygotic delivery of obesity-associated miR-let-7d-5p epigenetically rewires embryogenesis and*

impairs mitochondrial gene regulation in blastomeres.

Discussion:

High transcriptional plasticity of mitochondrial dysfunction in adipose tissue:

In our study we found that hepatic expression changes inflicted by HFD feeding where to a large extend irreversible, whereas adipose tissue miRNA/mRNA transcriptomes and translatomes, and particularly mitochondrial OxPhos and TCA genesets, reverted to a state *ante*. Similar studies have investigated the effects of reversible (i.e., diet- and exercise-evoked weight gain and loss) on hepatic and adipose transcriptomes: Gonzalez-Francesa et al. observed[40] that sustained diet-induced weight loss correlated with appreciable adipose transcriptional plasticity and mitochondrial gene expression, where short-term, exercise induced did not. Thus, durations of weight loss, and differences in control diet exposure might reflect the importance of nutritional and exposure factors during weight loss interventions on tissue responses in obesity. Despite not performing profiling of chromatic accessibility, transcriptional factor (TF) binding site enrichment analysis revealed that transcriptional activities of mitochondrial biogenesis factors like Nuclear Respiratory Factor 1 (*Nrf1*) and expression of transcriptional (co)-regulators such as Peroxisome proliferator-activated receptor gamma coactivator 1-alpha (*Ppargc1a*) and *Nrf1*[66] were repressed in obese fat (not shown). As Brandao et al. reported [67] that ablation of DICER1 in adipocytes repressed *Ppargc1a*/*Nrf1*, we can conceive a scenario where reduced DICER1 (for instance due to increased *miR-let-7d/e-5p*) drives and/or synergizes with reduced *Ppargc1a*/*Nrf1* levels to impair mitochondrial biogenesis in hypercaloric adipocytes.

Reprogramming of embryonal development by gametic miRNAs:

In *C. elegans* and *D. melanogaster*, germline gene silencing by small Noncoding RNAs (sncRNA) is well-understood and similar processes are suggested for mammals [18]. miRNAs are not only critical for sperm cell formation [25] and embryogenesis [26], but can also be transferred from sperm to zygotes and it is plausible that oocyte-sperm fusion alters zygotic miRNA pools, affect embryonic development, and instructs phenotypic traits and somatic gene expression in offspring as observed in our intergenerational paradigm. In contrast to DNA-intrinsic epigenetic modifications like DNA methylation (DNAme) and histone post-translational modifications, miRNAs diffuse between and across cellular compartments, are highly abundant in most cells, and, crucially, can be exchanged between cells through extravesicular

transport [30]. As mature sperm is transcriptionally inert, and no mechanism exists how sperm senses

endocrine information and translates this into differential miRNA levels, sperm relies on 'decoding

somatic cell types' that possess these sentinel features and are capable of instructing sperm by transfer

of miRNAs and other sncRNAs like mitochondrial tRNA fragments that are exchanged between sperm

and oocyte at fertilisation [21]. Furthermore, epididymal cells, somatic epithelial cells of the extragonadal

male reproductive tracts, secrete *epididysomes*, microvesicles that fuse with nascent sperm [68], and as

we observed several miRNAs condordantly regulated in F_0 adipose tissue and sperm, horizontal miRNA

transfer from adipocytes to sperm, potentially via epididysomes is a plausible scenario that still lacks

experimental proof in our paradigm. Finally, altering miRNA levels like *miR-199a-5p* in zygotes by

microinjection is sufficient to reprogram embryonal metabolism [69], highlighting that miRNA alterations

can rewire embryonal development and metabolism which are still compatible with fetal development.

Methods:

Mice

All animal experiments were approved by the Ministry of Environment and Agriculture Denmark (Miljø- og Fødevarestyrelsen), the license no. 2018-15-0201-01544. All mice were housed under a 12-h light/ 12-h dark cycle in a temperature and humidity-controlled facility and had *ad libitum* access to diets and drinking water. 4-week-old C57BL/6J male mice (Janvier Labs) were fed with a chow diet (NIH-31, Zeigler Brothers Inc., 8% calories from fat) for 2 weeks of acclimatization and then were fed with either a low-fat diet (LF, D12450J, 10% kcal from fat, Research Diet) or a high-fat diet (HF, D12492, 60% kcal from fat, Research Diet) for 9 weeks. After 9 weeks of feeding, half of HF-fed mice were randomly assigned to the diet regression group which was given LFD.

Mouse husbandry

All male virgin mice were maintained on the assigned diet until 21 weeks of age and mated with 10-week-old C57BL/6J female virgin mice (Janvier Labs). 8-week-old C57BL/6J female mice were fed with a chow diet for 2 weeks of acclimatization and then were fed with LF during breeding and lactation. Mating was set up for 5 days in a female's cage and vaginal plugs were checked every morning. All mice, including the F_0 HF group, were maintained on LF during breeding to remove confounding effects from maternal diets. Males were sent from breeding cages to the original cage to be fed with the assigned diet for additional 2 weeks and then sacrificed. The offspring from all groups of founders consumed only LF. Representative male offspring from six litters per group were included in all analyses.

Intraperitoneal glucose and insulin tolerance tests:

Mice were fasted for 6 h for glucose-tolerance tests. Fasting glucose levels were measured and then mice were intraperitoneally injected with 2 g/kg body weight of 20 % glucose solution (B. Braun Medical A/S). For intraperitoneal insulin-tolerance tests, random-fed glucose levels were measured, and mice were intraperitoneally injected with 0.75 IU/kg body weight of 100IU stock insulin (Novo

Nordisk). Blood glucose levels were measured at 15, 30, 60, 90, and 120 min after injecting glucose or

insulin using a CONTOUR®NEXT glucometer (Bayer).

Dual-energy x-ray absorbance:

Fat and lean mass were measured by Dual-energy x-ray absorbance (DXA) scan (PIXImus 2, GE Lunar

Corp.). Mice were anesthetized by isoflurane inhalation and placed in a prone position with all four

limbs stretched out. All body composition was measured on the total body excluding the head and tail

by version 2.10.041 of the proprietary software.

Tissue collection:

Mice were sacrificed by carbon dioxide euthanasia. Livers, eWAT, iWAT and BAT were weighed,

snap-frozen in liquid nitrogen, and stored in -80 °C. Pieces from liver and eWAT were fixed in 4%

paraformaldehyde (PFA) for histology analysis. For serum collection, blood collected by the cardiac

puncture was centrifuged at 2,000 g for 15 min. The supernatant was collected and stored in -80 °C.

Sperm collection:

Mature sperm cells were collected from the cauda epididymis. Briefly, sperm cells are allowed to swim

out from the cauda epididymis and incubate in PBS at 37 °C for 15 min. Supernatant containing sperms

was passed through a 40-μm cell strainer and then incubated in a somatic cell lysis buffer (0.5 % Triton-

X and 0.1 % SDS) for 40 min on ice. Sperms were centrifuged at 600 g for 5 min and the pellets were

washed with PBS and pelted again at 600 g for 5 min. The supernatant was discarded, and the pellets

were snap-frozen in liquid nitrogen and stored at -80 °C.

Zygotic miRNA microinjection and embryo implantation:

Ovulation of prepubescent (4-week-old) C57BL/6J females was induced by intraperitoneal injection of

PMSG, 5 IU/female, followed by IP injection of hCG, 5IU/female, 47h later. After the second injection,

females were set in breeding with C57BL/6J stud males. Next morning, they were monitored for

formation of a vaginal plug (indicating mating has occurred during the night). Females were euthanized

and oviducts were dissected to harvest the cumuli containing zygotes and unfertilized oocytes. Cumuli

were disaggregated by incubation in a 10mg/ml solution of Hyaluronidase (Sigma-Aldrich ref. H4272)

in M2 medium (Sigma-Aldrich ref. M7167) at RT for 10 minutes. Zygotes were sorted out, assessed by

the presence of the second polar body and cultured in KSOM medium (Merck Millipore ref. MR-106-

D). Oocytes were discarded.

Each miRNAs was diluted at 10 ng/ul in microinjection buffer (5 mM Tris, 0.1 mM EDTA, pH 7.4) as

stock solution and kept at -80 °C. The day of microinjection, 1,5 ng/ul solutions were prepared from

the stock, and injected in the pronuclei of the zygotes following standard microinjection procedures.

Injected zygotes were cultured in KSOM until the microinjection session was completed and then

cryopreserved by the "Slow Freezing" Method (Manipulating the Mouse Embryo, a Laboratory Manual,

fourth edition, Chapter 16, Richard Behringer et al. Cold Spring Harbor Laboratory Press, 2014) and

transported to SDU at -150 °C for embryo transfer and recovery.

The frozen microinjected embryos were thawed at SDU and cultured in KSOM (Merck, MR-121-D)

for 40 minutes (Manipulating the Mouse Embryo, a Laboratory Manual, fourth edition, Chapter 16,

Richard Behringer et al. Cold Spring Harbor Laboratory Press, 2014). Viable embryos with intact zona

pellucida were selected for further transfer. These embryos were then washed in 10 drops of M2 and

surgically transferred to the oviducts of 0.5 dpc pseudopregnant females. Pseudopregnancy was induced

in the females by mating them with a vasectomized male. Prior to implantation, the pseudopregnant

females were anesthetized with ketamine (100 mg/kg) and xylazine (10 mg/kg) through intraperitoneal

injection, and the surgical embryo transfer procedure was performed following the protocol described

by Behringer et al. (Manipulating the Mouse Embryo, a Laboratory Manual, fourth edition, Chapter 6,

Richard Behringer et al. Cold Spring Harbor Laboratory Press, 2014). Postoperative analgesic treatment

with carprofen (5 mg/kg) was administered subcutaneously for 48 hours.

8-cell-morula blastomere disaggregation:

miRNAs were injected in the embryo at zygote stage as described above. After injection, embryos were

incubated in KSOM (KSOM-Embryomax advanced embryo medium, Merck Millipore #MD-101-D)

for 48-56 hours at 37 °C, in 5% CO_2. The culture medium was refreshed daily. In the afternoon of the

third day of culture, when the embryos had reached the 8-cell morula stage, they were incubated at RT

in Tyrod's solution (Sigma-Aldrich T1788) for up to 30 seconds until the zona pellucida was completely

dissolved. Afterwards the naked morulas were washed through three drops of M2 medium (Sigma-

Aldrich M7167), and finally incubated in KSOM without Ca^{2+} and Mg^{2+} at RT for 20 minutes. Then,

embryos were moved to a new drop of fresh M2 medium and there, pipetted up and down repeatedly to

facilitate the blastomere disaggretation. When single blastomeres were obtained, they were individually

transferred to single wells of a 96-well single-cell-sequencing plate from VWR (Cat. No. 82006-636)

with 2 ul of RNAse inhibitor mix (containing 0.12 ul Triton X-100 10 %, 0.1 ul RNAse inhibitor (Takara

(Cat. No. 2313B, 40 U/ul), 1.78 ul MilliQ water (RNAse-free)). Plates were them frozen at -80 °C until

the sequencing was performed.

Lipodomics and Metabolomics:

Serum Extraction:

100 µL of serum was mixed with 1000 µL of a 2:1 chloroform/methanol solution and 200 µL of water.

Four blank samples, each containing 100 µL of water, were processed in parallel. The mixtures were

shaken at 1000 rpm for 30 minutes at 4°C, followed by centrifugation at 16,000g for 10 minutes at 4

°C to separate the phases. The aqueous phase was re-extracted with 350 µL of an 86:14:1

chloroform/methanol/water solution. The organic phase from this re-extraction was combined with the

corresponding organic phase from the initial extraction. Aqueous fraction was dried using SpeedVac,

while the organic phase was dried under a stream of nitrogen. For subsequent analysis, the dried aqueous

fractions were reconstituted in 30 µL of 1 % formic acid for metabolomics. The organic phase was

reconstituted in 30 µL of lipidomics solvent A (5:1:4 isopropanol/methanol/water with 5 mM

ammonium acetate and 0.1 % acetic acid). A pooled quality control sample was prepared for each

analytical setup by combining 5 µL from each sample, excluding blanks.

 Lipidomic analysis:

For lipidomics analysis, 2.5 µL of each sample was injected into an Agilent 1290 Infinity UPLC system

(Agilent Technologies, Santa Clara, CA, USA) equipped with a Zorbax Eclipse Plus C18 guard column

(2.1 × 5 mm, 1.8 µm) and an analytical column (2.1 × 150 mm, 1.8 µm), maintained at 50 °C. Analytes

were eluted at a flow rate of 400 µL/min using eluent A (water with 5 mM ammonium acetate and 0.1

% acetic acid) and eluent B (99:1 isopropanol/water with 5 mM ammonium acetate and 0.1 % acetic

acid). The gradient was as follows: 0 % B from 0 to 1 min, 0-25 % B from 1 to 1.5 min, 25-95 % B

from 1.5 to 12 min, 95 % B from 12 to 14 min, and 95-0 % B from 14 to 15 min, followed by a 3-

minute equilibration.

For metabolomics analysis, 1.5 µL of each sample was injected into the same UPLC-MS system, but

with a ZORBAX Eclipse Plus C18 guard column (2.1 × 50 mm, 1.8 µm) and an analytical column (2.1

× 150 mm, 1.8 µm) maintained at 40°C. Eluent A was 0.1 % formic acid, and eluent B was 0.1 % formic

acid in acetonitrile. The gradient was as follows: 3 % B from 0 to 1.5 min, 3-40 % B from 1.5 to 4.5

min, 40-95 % B from 4.5 to 7.5 min, 9 5% B from 7.5 to 10.1 min, and 95-3 % B from 10.1 to 10.5 min,

followed by a 3.5-minute equilibration.

Both analyses were performed on a 6530B quadrupole time-of-flight mass spectrometer (Agilent

Technologies, Santa Clara, CA, USA) for mass spectrometric detection in positive and negative ion

modes. MS1 mode settings included a scanning range of 50-1700 m/z for lipidomics and 10-1050 m/z

for metabolomics, with 2 scans/sec. Internal calibration was performed using Hexakis(1H,1H,3H-

tetrafluoropropoxy) phosphazene delivered through a second needle in the ion source via an isocratic

pump running at 20 µL/min. Iterative data-dependent MS2 mode was used for fragmentation in both

analyses with the following settings: collision energy at 40 V (lipidomics) and 20-40 V (metabolomics),

precursor threshold at 5000 counts, and active exclusion after 2 spectra, with re-inclusion after 0.5 min.

Data Processing and Normalization:

All data was converted. mzML format using ProteoWizard[70]. Lipidomics data was annotated in MS-

Dial (v. 4.9) against the incorporated lipid-blast[71] with an MS1/MS2 mass tolerance of 0.005/0.01 Da

and a minimum identification score of 70 %. The annotated compounds were exported to PCDL

manager B.08.00 (Agilent Technologies) to create a database for area extraction in Profinder 10.0

(Agilent Technologies). Metabolomics data was processed in MzMine (v2.59) utilizing modules such

as ADAP chromatogram builder and deconvolution. Compounds were annotated at Metabolomics

Standards Initiative (MSI) levels 3 using the libraries of National Institute of Standards and Technology

2017 (NIST17) and MassBank of North America (MoNa). Features with signals less than 5 times those

in blanks or missing in more than 20 % of QC samples were removed and signals were QC corrected

using statTarget[72].

Serum insulin and leptin ELISA:

Serum insulin and leptin levels were measured by mouse insulin ELISA kit (Mercodia, Uppsala,

Sweden) and mouse leptin ELISA kit (Crystal Chem, IL, USA) according to manufactures' protocol.

Histology and staining:

Liver and adipose tissues fixed with 4 % of paraformaldehyde were used for histology analyses.

Formalin-fixed, paraffin-embedded tissues were cut into 3 μm sections. Sections were stained by

hematoxylin and eosin (H&E) and Sirius Red by standard procedures at the Department of Pathology

at Odense University Hospital, Odense, Denmark. Immunohistochemistry for CD45 was performed on

Ventana BenchMark Ultra (Roche) with ready-to-use CD45 (LCA) (2B11 & PD7/26) Mouse

Monoclonal Antibody. A liver pathologist (SD) assigned steatosis grades as S0 (< 5% hepatocytes with

large fat vacuoles), S1 (5-33%), S2 (>33-66%), and S3 (>66% hepatocytes with large fat vacuoles)[73].

Lobular inflammation (0-3) and hepatocellular ballooning (0-2) were assessed according to the NAS-

CRN Activity Score. In addition, portal inflammation was assessed, using a 4-tiered score. Liver fibrosis

stage was assessed according to NAS-CRN for fibrosis; F0, no fibrosis; F1, perisinusoidal or periportal

fibrosis; F2, perisinusoidal and portal/periportal fibrosis; F3, bridging fibrosis; F4, cirrhosis.

Measurement of adipocyte volumes:

The volume weighted mean volume of adipocytes was determined from the H&E sections with

newCAST stereology software (Visiopharm), autodisector; measurements on systematic, randomly

selected fields were performed on by at least 75 adipocyte profiles in each tissue.

Bulk proteomics from epididymal white adipose tissue:

Deep frozen fat pads (200 mg of epididymal adipose tissue) were powdered with mortar-pestle and

subjected to homogenization with ceramic beads (Percellys, 5500 rpm, 2 x 10 seconds, 30 seconds

break) in the buffer containing 50 mM HEPES pH 7.4, 1 % Triton X-100, 100 mM NaF, 10 mM sodium

orthovanadate, 10 mM EDTA, 0.2 % SDS, 100 mM NaCl. Tissue homogenates were further processed

to remove the nucleic acid traces (Diagenode SA; 10 min, cycle 30/30 seconds, at 4 °C). Floating fat

and tissue debris were removed from protein extracts by centrifugation at 20000 x g. 150 mg of proteins

were precipitated with 4x volume of ice-cold acetone and washed with ice-cold acetone twice. Air-dry

pellets were resuspended in 8M Urea/50 mM triethylammonium bicarbonate (TAEB) buffer

supplemented with a protease inhibitor cocktail (Sigma). 50 mg of the samples were incubated with 5

mM dithiothreitol at 25°C for 60 min, followed by treatment with 40 mM chloroacetamide for 30 min

(room temperature, light-protected). Lys-C protease (enzyme: substrate ratio of 1:75) was added for 4

hours at 25 °C. The TAEB-diluted sample (2M Urea final) was supplemented with trypsin (1:75 ratio)

followed by overnight digestion at 25 °C. Acidified peptides (1 % formic acid final) were purified on

the SDR-RP (C18) multi-stop-and-go-tip (Stage Tip)[74].

LC MS/MS Label-Free Quantitative Proteomic Analysis from epididymal white adipose tissue:

All samples were analyzed by the Proteomics Facility on a Q Exactive Plus Orbitrap mass spectrometer

coupled to an EASY nLC 1000 (both Thermo Scientific). Peptides were loaded with solvent A (0.1 %

formic acid in water) onto an in-house packed analytical column (50 cm - 75 µm I.D., filled with 2.7

µm Poroshell EC120 C18, Agilent) and separated with 150 min gradients. Peptides were

chromatographically separated at a constant flow rate of 250 nL/min using the following gradient: 4-6

% solvent B (0.1 % formic acid in 80 % acetonitrile) within 5.0 min, 6-23 % solvent B within 120.0

min, 23-54 % solvent B within 7.0 min, 54-85 % solvent B within 6.0 min, followed by washing and

column equilibration. The mass spectrometer was operated in data-dependent acquisition mode. The

MS1 survey scan was acquired from 300-1750 m/z at a resolution of 70,000. The top 10 most abundant

peptides were isolated within a 2.1 Th window and subjected to HCD fragmentation at a normalized

collision energy of 27 %. The AGC target was set to 5E5 charges, allowing a maximum injection time

of 60 ms. Product ions were detected in the Orbitrap at a resolution of 17,500. Precursors were

dynamically excluded for 20.0 s. All mass spectrometric raw data were processed with MaxQuant[75]

(version 1.5.3.8, using default parameters and searched against the canonical murine Uniprot reference

proteome (UP589, downloaded on 28/08/2020). Label-free quantification option was enabled as was

the match-between runs option between replicates. Follow-up analysis was done in Perseus[76] 1.6.15.

Hits from the decoy database, the contaminant list, and those only identified by modified peptides were

removed. Afterwards, results were filtered for data completeness in replicate groups and LFQ values

imputed using sigma downshift with standard settings. Finally, FDR-controlled T-tests between sample

groups were performed with s0 = 0.2. Proteins were annotated on GOBP, GOCC, KEGG and Pfam

terms. Afterwards, all one-way ANOVA (s= 0.2) significant candidates were Zscored and hierarchically

clustered using the Euclidean distance model. Resulting clusters were merged onto the whole data set

and a Fisher Exact Test calculated on cluster level with the whole dataset as background.

RNA isolation from mouse tissue and primary adipocytes:

Total RNAs from tissues, sperm, and primary adipocytes were isolated using the TRI reagent (Sigma)

followed by clean-up with RPE buffer (Qiagen, Germany). The quality of RNA was validated by the

Agilent RNA 6000 Nano-Kit in Agilent 2100 Bioanalyzer according to the manufacturer's protocol.

(Agilent Technologies, Waldbronn, Germany).

RNA sequencing:

Small RNA sequencing:

(1) For sRNA-seq from F_0 sperm, library preparation and sRNA-seq was performed at Cologne Center

for Genomics, Germany. 50 ng of total RNA was used for library preparation followed by size selection

(Small RNA-Seq Library Prep Kit, Lexogen). The library was sequenced in 1x50-bp single-end reads

on NovaSeq6000 serie no. A00316. (2) Small RNA sequencing from F_0 and F_1 eWAT was carried out

after quality control, 125 ng of total RNA was used for library preparation (Nextflex library kit)

followed by sequencing by 1x50-bp single-end reads on Illumina Novaseq 6000.

mRNA sequencing:

mRNA sequencing was performed in-house. After quality control, 1 µg of total RNA was used for

library preparation (NEBNext RNA prep kit) followed by 2x50bp paired-end sequencing on Illumina

Novaseq 6000.

RNAseq data analysis:

Sequencing reads were aligned to the mouse reference genome from Gencode (vM25)[77] using STAR

aligner (v 2.7.9a)[78] and gene features were counted by using featureCountsv (2.0.3)[79]. The genes with

low expression were filtered out using *filterByExpr* function with default settings (apart from "Embryo

Inject" dataset, where parameter min.count was set to 40 as more rigorous filtering was required) from

edgeR (v4.0.16)[80] R package. Differential expression analyses were performed using edgeR's

voomLmFit [81]unction with sample quality weights (sample.weights = TRUE) and linear model "0 +

group", whereas group represents the combination of all experimental groups of interest. The

experimental conditions and linear models used for each dataset are summarized below. Genes with

adjusted p-values below 0.05 were considered as differentially expressed. Gene enrichment analyses

were performed by clusterProfiler (v4.10.1)[82] R package using both over-representation analysis (ORA)

and gene-set enrichment analysis (GSEA) using differentially expressed genes and log-fold-change

(logFC) ranked genes ($n_{permutation}$ = 1,000,000), respectively. Gene sets were retrieved from Gene

Ontology (GO.db v3.18.0) and Reactome (reactome.db, v1.86.2) databases.

Smart-seq2 mRNA sequencing of murine blastomeres:

Custom-made Smart-seq2 protocol was implemented as before[65]. Briefly, after thawing the plate, 1 µL

of   dNTP   mix   (10 mM),   0.1 µL   of   oligo-dT   (100   µM,

/5Biosg/AAGCAGTGGTATCAACGCAGAGTAC(T)30VN-3′) and 0.9 ul nuclease-free water was

added to each well and the lysis protocol was run when the samples were incubated at 72°C for 3 min

in thermocycler. Samples were then put on ice for at least 1 min. Next, 5.60 µL of reverse-transcription

mix containing 0.25 µL of RNase I, 2 µL of 5× first-strand buffer, 2 µL of betaine (5 M), 0.06 µL of

MgCl$_2$ (1 M), 0.5 µL of DTT (100 mM), 0.5 µL of SuperScript III (200 U/µL), and 0,29 ul of nuclease-

free water was added to each well and reverse transcription was carried out as follows: 42 °C-90:00, 42

°C-hold, 42 °C—12:20, 10 cycles (50 °C-2:00, 42 °C-2:00), 39 °C-12:00, 70 °C-15:00, 4 °C hold. After

the first 90 min of RT, 0.4 µL of 12,5 µM TSO (0.83 µM final concentration) was added to each reaction

at room temperature. The plate was resealed, centrifuged for 1 min at 700 g, and placed back on the

thermocycler to resume cycling from 42 °C-12:20. After reverse transcription, 15 µL of cDNA

enrichment PCR reaction mix containing 12.5 µL of 2× KAPA HiFi HotStart ReadyMix, 0.25 µL of

ISPCR primer (10 µM, /5Biosg/AAGCAGTGGTATCAACGCAGAGT-3′), and 2.25 µL of nuclease-

free water was added to each well and cDNA enrichment PCR was performed as follows for 21 cycles:

98 °C-3:00, 7 cycles (98 °C-0:20, 60 °C-4:00, and 72 °C-6:00), 7 cycles (98 °C-0:20, 64 °C-0:30, and

72 °C-6:00), and 7 cycles (98 °C-0:20, 67 °C-0:30, and 72 °C-7:00), 72 °C-10:00, 4 °C-hold. The cDNA

purification was performed using AMPure XP beads at a ratio of 1:1 (Beckmann). For elution, Qiagen

elution buffer (EB) was used. cDNA concentrations were measured using the Qubit 3.0 fluorometer

according to the manufacturer's protocol, and all cDNA samples were analyzed on Agilent Tape station

using the High Sensitivity DNA assay.

Dual-indexed Illumina Nextera XT sequencing libraries were prepared using 25 % of the recommended

reaction volumes of Nextera XT components. All Nextera XT libraries were prepared using 125 pg of

input cDNA for tagmentation, and were subsequently enriched for 12 PCR cycles, and purified using

AMPure XP beads (Beckmann) at a sample:bead ratio of 0.6:1. Library concentrations were measured

using Qubit 3.0 fluorometer, and the fragments were analyzed on the Agilent Tape station using the

High Sensitivity DNA assay. All libraries were individually normalized and diluted to a concentration

of pooled library to 5.4 ng/µL with average size ~515bp (16 nM). The libraries were sequenced on

Illumina NextSeq500 in a single-end 75-bp format (FC-404-2005, Illumina). Single-nuclei libraries

were sequenced at an average depth of 3 mio reads.

Data availability – NGS:

(1) Illumina datasets (sRNA-seq, mRNA-seq) from F_0 / F_1 liver and eWAT are available as Gene Expression Omnibus series under accession number [GEOxxx].

(2) Illumina datasets (sRNA-seq) from F_0 sperm are available as Gene Expression Omnibus series under accession number [GEOxxx].

(3) Illumina datasets (mRNA-seq) from offsprings of miRNA-injected zygotes (F_m) eWAT are available as Gene Expression Omnibus series under accession number [GEOxxx].

(4) Illumina datasets (mRNA) from 1°eWA overexpressing Negative Control, *miR-let-7d-5p, miR-let-7e-5p* mimics are available as Gene Expression Omnibus series under accession number [GEOxxx].

(5) Smart-Seq2 single-cell mRNA-seqdata from miRNA-injected 8-cell stage embryos are available as Gene Expression Omnibus series under accession number [GEOxxx]

Data availability – Proteomics:

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE [1] partner repository with the dataset identifier PXD042800. The data are available to the reviewer discretion: **Username: reviewer_pxd042800@ebi.ac.uk, Password:** QHj9fv33

Data availability – Metabolomics:

[]

Male participants, sperm samples and analysis

The intervention study was approved by the Danish National Committee on Health Research Ethics (SJ-769), the Danish Data Protection Agency (REG-048- 2019) and performed at the Fertility Clinic, Zealand University Hospital, Denmark[62]. The semen samples were collected from obese patients (≥ 30 kg/m2) via masturbation following two days of abstinence. The sperm analyses were performed in the IVF laboratory in an authorized tissue establishment. The other semen samples were diluted with

MHM-C (Fujifilm) and frozen immediately for further analysis. The semen samples were collected via

masturbation following two days of abstinence, and the samples were analyzed a maximum of two to

three hours after production. After liquefaction, ejaculate volume and sperm concentration (total, motile

and progressive) were assessed using a microscope (Leica DM750). Sperm concentration was

determined with a CellVision counting chamber (CellVision Technology, Heerhugowaard, the

Netherlands). Total sperm count was calculated as the product between ejaculate volume and sperm

concentration. The sperm analyses were performed in the IVF laboratory in an authorized tissue

establishment.

Study design and approvals

This prospective intervention study was conducted between May 2020 and April 2022 at the Fertility

Clinic, Zealand University Hospital, Denmark. The study was approved by the Danish National

Committee on Health Research Ethics (SJ-769), the Danish Data Protection Agency (REG-048- 2019)

and registered on the Clinical Trial website (www. ClinicalTrials.gov identifier NCT04721938). All

participants provided written informed consent.

Primary adipocyte culture and transfection

Epididymal white adipose tissues (eWAT) from 6- to 8-week-old male C57BL/6J mice were removed,

minced, and digested with pre-warmed collagenase media (500 U/mL collagenase type 2 (LS004183,

Worthington) and 3.75U/mL DNase I (10104159001, Sigma). Isolated cells were seeded in 6-well

plates and grown in DMEM/Ham's F12 medium (21331, Vendor: Gibco), supplemented with 0.1 %

Biotin/D-Pantothenate (33 mM/17 mM), 1 % penicillin-streptomycin, and 20 % FBS. Until reaching

80 % confluency, primary cells were induced by induction medium with 1 µM rosiglitazone, 850 nM

insulin, 1 µM dexamethasone, 250 µM 3-isobutyl-1-methyl-xanthin (IBMX) and 10 % FBS.

Subsequently, the medium was changed every other day by a differentiation medium containing 1 µM

rosiglitazone, 850 nM insulin and 10 % FBS. 100 nM of Negative Control and *mmu-miR-let-7-d/e-5p*

mimics or LNA oligos tageting *Dicer1* were reversely transfected using Lipofectamine RNAiMAX

(Thermo Fisher Scientific) at day 4 after induction. Completely differentiated cells were harvested at 7

days after induction for further analysis.

Plasmid design and dual luciferase assay

For the generation of reporter constructs, the specific region of murine *Ago2*, *Dicer-1*, *Hif1an*, *Hk2*,

Cpt2, *Crat*, *Pcx*, *Idh2*, *Sdha*, *Sdhaf3*, *Pdk2*, *Slc25a22* and *Slc25a24* containing either the wildtype (wt)

or mutated (mut) *miR-let-7* binding sites was cloned behind the translation stop codon of the firefly-

luciferase gene in pmirGLO Dual-Luciferase vector using custom DNA oligos provided by IDT. Renilla

luciferase coding in the same plasmid was used as a control reporter for normalization purposes. To

report the miRNA activity in mammalian cells, HEK 293 cells were reversely transfected in 96-well

plates with 60 ng DNA using the TransIT transfection reagent (mirusbio) for 3 days and forwardly

transfected with *miR-let-7d/e-5p* mimics for 2 days. HEK-293 cells were cultured in DMEM medium

(21013, Gibco) supplemented with 5.6 mM L-glutamine, 1 % penicillin-streptomycin, and 10 % FBS.

After 72-h vector transfection and 48-h miRNA mimics transfection, dual luciferase reporter assays

(Promega) were performed. The relative activities of firefly luciferase indicating the interaction

between miRNA and specific region of targeted mRNA were normalized by the activities of Renilla

luciferase.

TaqMan assay and general qPCR

To quantify the levels of mature miRNAs, TaqMan MicroRNA Reverse Transcription Kit (4366596,

Thermo Fisher), TaqMan MicroRNA Assays and TaqMan Universal Master Mix II (4440042, Thermo

Fisher) were applied according to manufacturer's protocols. Level of mature *hsa-miR-16* was used as

internal control in human samples. The relative expression of mature miRNA was calculated using a

comparative method ($2^{-\partial\partial CT}$) according to the ABI Relative Quantification Method.

Determination of oxygen consumption rates and measurement of glycolytic activity:

Primary adipocytes from epididymal SVF (1°eWA) were seeded into Agilent Seahorse XFe96

Bioanalyzer microplates. Induction and differentiation were performed as described above. After four

days of differentiation, 16,000 per well mature adipocytes were re-seeded and transfected with LNA

oligos targeting *Dicer1* in DMEM/Ham's F-12 medium plus 10 % Fetal Bovine Serum, 1 % P/S, 0.1 %

Biotin and 0.1 % Pantothenic acid (Growth medium) at 37° C and 5 % CO_2. Seahorse measurement

was conducted at day 7 from differentiation. For each seahorse plate, the corresponding calibration plate

was prepared 24 hours prior to experiments using 200 µl XF Seahorse Calibrant Agilent per well.

Corresponding media were prepared before the experiment and consisted of Assay Medium

supplemented with 25 mM Glucose, 1 mM Glutamine, 2 mM Sodium Pyruvate, set to pH = 7.4 and

filtered sterile. The calibration plate possessed a cartridge having 4 pockets per well. Shortly before the

measurement pocket A was filled with 20 µl 20 µM Oligomycin, pocket B with 22 µl 20 µM FCCP and

pocket C with 25 µl 5 µM Antimycin A and Rotenone. One hour before measurements, plates were

washed with 1x PBS and changed to 180 µl Assay Medium in a non-CO_2 incubator at 37° C. Calibration

was started using calibration plates, measuring O_2 and pH LED Value/emission/Initial reference Delta

for each well. All measurements started with measuring basal values, followed by injection of

Oligomycin, FCCP, and Rotenone plus Antimycin A (MitoStressKit). Measurement parameters were:

Mix 3 min, wait 0 min, measure 3 min with each reagent's effect assessed within three consecutive

measurement cycles with a total duration of 18 min per reagent injection.

Protein extraction and western blot analysis

After being washed once by phosphate-buffered saline (PBS), the treated cells at harvest time were

lysed by RIPA buffer supplemented with cOmplete and PhosSTOP and quantified by Pierce BCA

Protein Assay Kit (23227, Thermo Fisher). Western blot analyses were carried out according to standard

protocols with primary antibodies listed in **Table S1**. Relative protein quantification was normalized

by HSC70.

Statistical analysis

Data were shown as mean ± standard error. One-way ANOVA and two-way ANOVA followed by

Tukey's multiple comparisons test were used for statistical analysis in *in vivo* experiments. Significant

differences between groups were noted in figure legends respectively.

Acknowledgements:

We thank the CECAD Proteomics Core Facility for support with proteomics analyses, the Laboratory

Manager, Stine Ravn, at The Zeland Fertility Clinic for handling and analyzing the sperm samples. We

wish to thank Bente Møller and Irina Korshuniova from BRIC Single Cell Genomics Core Facility for

excellent help for Smart-seq2 experiments. Smart-seq2 work was partly supported by CellX (The

Danish Single Cell Examination Platform) funded by the Danish Research Agency through the Danish

national research infrastructure program (5229-0009B). JWK, JHP and BDML received funding from

the European Research Council Starting Grant TransGenRNA (#675014), JWK, CH and NS from

Sygeforsikring Denmark, University of Southern Denmark and the Danish Diabetes Academy (DDA),

which is funded by the Novo Nordisk Foundation (NNF17SA0031406). JWK received support from

the NNF Challenge (33444) and Bioscience and Basic Biomedicine Programs of the NNF (28416).

  Figure legends

Figure 1: Obesity and weight loss elicit reversible changes in glucose metabolism and

mitochondrial metabolites. (**a**) Illustration of intergenerational *in vivo* cohorts depicting F_0 groups of

obese and weight-regression C57BL/6N male mice, F_1 offsprings and the overall analytical workflow.

(**b**) Body weights of F_0 LF (n=6), F_0 HF (n=13; n=6 for 15-24 weeks of age,), F_0 HF-LF (n=7). (**c,d**)

Blood glucose levels (**c**) and area under the curve (AUC) of glucose levels in (**d**) during intraperitoneal

glucose tolerance test (ipGTT) in F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=7) mice. (**e**) Relative blood

glucose levels during intraperitoneal insulin tolerance test (ipITT) in F_0 LF (n=6), F_0 HF (n=6) and F_0

HF-LF (n=7) mice. (**f**) Random-fed blood glucose in F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=7)

mice. (**g**) Serum insulin levels as measured by ELISA in F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=7)

mice. (**h**) Serum leptin levels measured by ELISA in F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=7)

mice. (**i**) Relative organ weights in F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=7) mice. (**j**) Illustration

of organ acquisition and multi-modal omic methodologies (**k**) Principal component analysis (PCA) and

PC1-3 representation of n=600 polar metabolites in serum of F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF

(n=7) mice. (**l**) Correlation cofficients and ID of serum metabolites irreversibly (*left*, n=95) or reversibly

(*right*, n=118) altered in F_0 HF/F_0 HF-LF vs F_0 LF mice. Data in (**k,l**) were analyzed and plotted using

Metaboanalyst 6.0 (https://www.metaboanalyst.ca/MetaboAnalyst/). (**m**) Illustration of differential

metabolite abundances from (**k,l**) in the tricarbon acid cycle (TCA) pathway in serum of F_0 LF (n=6),

F_0 HF (n=6) and F_0 HF-LF (n=7) mice. One-way ANOVA (**d**,**f-i**) and two-way ANOVA (**b,c,e**) followed

by Tukey's multiple comparisons test were used for statistical analysis. Data are presented as mean ±

standard error (**b,c,e**) or individual values and superscript letters represent group comparisions (**a**,

p<0.05, F_0 LF vs F_0 HF; **b**, *p*<0.05, F_0 LF vs F_0 HF-LF; **c**, *p*<0.05, F_0 HF vs F_0 HF-LF).

Figure 2. Multimodal omics analysis reveals reversible mitochondrial gene aberrations in obese

adipose tissue. (**a**) Representative hematoxylin and eosin (H&E) stainings in eWAT of F_0 LF, F_0 HF

and F_0 HF-LF mice. (**b**) Size distribution of adipocytes area in eWAT of F_0 LF (n=6), F_0 HF (n=6) and

F_0 HF-LF (n=6) mice. (**c**) Multidimensional scaling (MDS) plots of transcriptomic changes in eWAT

as determined by m RNA-seq. Dots represent eWAT from individual animals, n=6 per group. (**d-g**)

Volcano plot of significantly up-(red) and down-regulated (blue) differentially expressed genes (DEG)

in eWAT from (**d**) F_0 HF vs F_0 LF, 3,867 upregulated and 2,337 downregulated DEG, (**e**) F_0 HF vs F_0

HF-LF, 2,858 upregulated and 1,986 downregulated DEG and (**f**) F_0 HF-LF vs F_0 LF, 459 upregulated

and 262 downregulated DEG. Plots depict log2 transformed fold-changes and log10- transformed false-

discovery rates (FDR) of expression changes. Significant DEG are defined as FDR≤0.05. (**g**) Gene Set

Enrichment Analysis (GSEA) and gene ontology (GO) enrichment of biological processes (GOBP) of

DEG shown in (**d-f**). (**h**) Heatmap depicting relative abundances of miRNAs in F_0 LF (n=6), F_0 HF

(n=6) and F_0 HF-LF (n=6) eWAT as determined by small RNA-seq. Boxes depict miRNAs that were

reported to positively (*brown*) or negatively (*red*) affect mitochondrial biogenesis and function. (**i**)

Abundance of positive (*miR-143-3p*, *miR-181a-2-3p*, *miR-378a-3p* and *miR-378*, brown) and negative

(*miR-532-5p*, red) miRNA regulators in F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=6) eWAT. One way

ANOVA (**b,i**) followed by Tukey's multiple comparisons test were used for statistical analysis. Data

are presented as individual values and superscript letters represent group comparisions (**a**, $p<0.05$, F_0

LF vs F_0 HF; **b**, $p<0.05$, F_0 LF vs F_0 HF-LF; **c**, $p<0.05$, F_0 HF vs F_0 HF-LF). (**j**) Schematic representation

of miRNA-mRNA differential expression networks using Cytoscape (https://cytoscape.org/). Shown

are predicted mRNA targets of miRNA from (**i**) as identified by AGO-HITS-CLIP in adipose tissue

from[56]. Node color depicts differential expression of predicted miRNA targets (F_0 HF (n=6) vs F_0 LF

(n=6)) and yellow highlights miRNA seed lengths (*left*, 7mer; *right*, 8mer). Edge color depits genic

localisation of the lontest miRNA seed.

Figure 3: Paternal obesity impairs glucose tolerance and instigates adipose dysfunction in F1

adipose tissue. (**a**) Body weight of F_1 [F_0 LF] (n=18), F_1 [F_0 HF] (n=25) and F_1 [F_0 HF-LF] (n=21). (**b,c**)

Blood glucose levels (**c**) and AUC of glucose levels during IPGTT in F_1 [F_0 LF] n=10), F_1 [F_0 HF]

(n=10) and F_1 [F_0 HF-LF] (n=10) mice. (**d**) Relative blood glucose levels during IPITT in F_1 [F_0 LF]

(n=10), F_1 [F_0 HF] (n=10) and F_1 [F_0 HF-LF] (n=9) mice. (**e**) Random-fed serum glucose levels in F_1

[F_0 LF] (n=18), F_1 [F_0 HF] (n=25) and F_1 [F_0 HF-LF] (n=21) mice. (**f**) Serum insulin levels measured by

ELISA in F_1 [F_0 LF] (n=6), F_1 [F_0 HF] (n=8) and F_1 [F_0 HF-LF] (n=8) mice. (**g**) Serum leptin levels

measured by ELISA in F_1 [F_0 LF] (n=7), F_1 [F_0 HF] (n=8), and F_1 of F_0 HF-LF (n=8) mice. (**h**) Body

composition measured by DXA in F_1 [F_0 LF] (n=11), F_1 [F_0 HF] (n=11) and F_1 [F_0 HF-LF] (n=11) mice.

(**i**) relative organ weights of F_1 [F_0 LF] (n=10), F_1 [F_0 HF] (n=10) and F_1 [F_0 HF-LF] (n=7). (**j,k**)

Representative H&E staining (**j**) and size distribution (**k**) of adipocytes in eWAT of F_1 [F_0 LF] (n=6),

F_1 [F_0 HF] (n=6) and F_1 [F_0 HF-LF] (n=6) mice. (**l**) GSEA-GOBP analysis of DEGs in eWAT of F_1 [F_0

LF] (n=4) F_1 [F_0 HF] (n=6) and F_1 [F_0 HF-LF] (n=6) mice. (**m,n**) Heat map and hierarchical clustering

differentially translated genes (**m**) and relative protein abundances (LFQ) of NDUFA6 and COX6B1

in eWAT of F_0 LF (n=6), F_0 HF (n=6), F_0 HF-LF (n=6), F_1 [F_0 LF] (n=6), F_1 [F_0 HF] (n=6), and F_1 of F_0

REV (n=6). One-way ANOVA (**c,g,i, k,n**) and two-way ANOVA (**b**) followed by Tukey's multiple

comparisons test were used for statistical analysis. Data are presented as mean ± standard error (**b,d**) or

individual values and superscript letters represent group comparisions (**a**, $p<0.05$, F_0 LF vs F_0 HF; **b**,

$p<0.05$, F_0 LF vs F_0 HF-LF; **c**, $p<0.05$, F_0 HF vs F_0 HF-LF).

Figure 4: Obesity induces spermatozoa and adipose miR-let-7 and blunts adipocyte mitochondrial

function via a miR-let-7-DICER1 axis. (**a,b**) Heatmap (**a**) and log10-transformed counts (**b**) of *miR-*

let7 isoforms in sperm from F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=6) mice. One-way ANOVA

followed by Tukey's multiple comparisons test was used for statistical analysis. Superscript letters

represent group comparisions and P values are indicated in the panel. (**c**) Heatmap representation of

shown *miR-let-7* isoforms in F_0 LF (n=5), F_0 HF (n=5), F_0 HF-LF (n=5), F_1 [F_0 LF] (n=6), F_1 [F_0 HF]

(n=5), and F_1 of F_0 REV (n=5) eWAT and spermatozoa. (**d**) Illustration of experimental workflow and

methodology to characterize the effects of *miR-Let-7d-5p* and *miR-Let-7e-5p* in 1°eWA from

C57BL/6N mice. (**e**) Absolute expression (Cq value) of *miR-let7-d-5p* and *miR-let7-e-5p* in 1°eWA

transfected with 100nM Negative Control, *miR-let7-d-5p* and *miR-let7-e-5p*. (**f,g**) Volcano plot of

significantly up- (red) and down-regulated (blue) DEG in 1°eWA transfected with 100nM of (**f**) *miR-*

let7-d-5p vs Negative Control, 1,301 upregulated and 977 downregulated DEGs (**g**) *miR-let7-e-5p* vs

Negative Control, 635 upregulated and 759 downregulated DEGs. (**h**) Schematic representation of *miR-*

let-7-mRNA DEN, depicted predicted *miR-Let-7* target mRNA repressed in F_0 HF (n=6) vs F_0 LF (n=6)

with seed lengths $\geq$7nt using adipose tissue AGO-HITS-CLIP[56]. Node color depicts differential

expression of *miR-let-7* gene targets and yellow highlights genes from Mitocarta3.0 [69]. Edge color

depits the genic localisation of the *miR-Let-7* seed. (**i,j**) Relative luciferase activity in HEK293 cells

after co-transfection of pmirGLO dual luciferase reporter assay (DLRA) constructs harbouring (**i**) wild-

type 3′-UTR fragments of indicated genes or (**j**) wild-type and mutated fragments of *Dicer1*, *Hif1an*,

Hk2 and *Slc25a24* transfected with 100nM of Negative Control, *miR-let-7d-5p* and *miR-let-7d-5p*

mimics. Data represent at least three biological replicates, each performed in three technical replicates.

One-way ANOVA (**j,k**) followed by Tukey's multiple comparisons test were used for statistical

analysis. Superscript letters represent group comparisions (**a**, $p<0.05$, Negative Control vs *miR-let-7d-*

5p; **b**, $p<0.05$, *miR-let-7e-5p* vs *miR-let-7e-5p*; **c**, $p<0.05$, Negative Control vs *miR-let-7e-5p*). (**k**)

Western blot analysis of 1°eWA transfected with 100nM of Negative Control, *miR-let-7d-5p* and *miR-*

let-7d-5p mimics performed with indicated antibodies. HSC70 served as loading control. (**l,m**) Oxygen

consumption rates (**l**, OCR) and extracellular acidification rates (**m**, ECAR) in 1°eWA transfected with

100nM of Negative Control A, *Dicer1* LNA1 and *Dicer1* LNA2 and stimulated with oligomycin (O),

FCCP (F) and Antimycin A plus Rotenone (A/R). Unpaired, two-tailed Student's t-tests were performed

at indicated timepoints (**a**, $p<0.05$, Negative Control vs *Dicer1* LNA1; **b**, $p<0.05$, *Dicer1* LNA1 vs

Dicer1 LNA2; **c**, $p<0.05$, Negative Control vs *Dicer1* LNA2).

Figure 5: Zygotic microinjection of obesity-associated *miR-let-7* causes F$_1$ glucose intolerance.

(**a**) llustration of experimental workflow for zygotic miRNA injection and phenotypic characterisation

of sired offsprings (F$_m$ mice). (**b**) Body weights of male F$_m$ mice sired from zygotes injected with

Negative Control (n=9), *miR-let-7d-5p* (n=14), *miR-let-7e-5p* (n=14) and *miR-699-5p* (n=12). (**c-e**),

Organ weights of eWAT (**c**) and liver (**d**) from male F$_m$ mice sired from Negative Control (n=7), *miR-*

let-7d-5p (n=6), *miR-let-7e-5p* (n=6) and *miR-699-5p* (n=7) treated zygotes. (**e**) Random-fed blood

glucose at indicated weeks of age in male F$_m$ mice sired from Negative Control (n=7), *miR-let-7d-5p*

(n=6), *miR-let-7e-5p* (n=6) and *miR-699-5p* (n=7) treated zygotes. One way ANOVA followed by

Tukey's multiple comparisons test were used for statistical analysis. Data are presented as mean ±

standard error. **, $p<0.01$, Negative Control vs *miR-let-7d-5p*, *miR-let-7e-5p* and *miR-699-5p*. (**f-h**),

Blood glucose levels during IPGTT in male F$_m$ mice sired from Negative Control (n=9), *miR-let-7d-5p*

(n=9), *miR-let-7e-5p* (n=9) and *miR-699-5p* (n=8) treated zygotes. Unpaired, two-tailed Student's t-

tests were performed at indicated timepoints and P values are indicated. (**i**) GSEA-GOBP analysis of

DEGs in eWAT of male F$_m$ mice sired from Negative Control (n=5), *miR-let-7d-5p* (n=6) and *miR-let-*

7e-5p (n=6) treated zygotes.

Figure 6: Obesity-associated *miR-let-7* rewires mitochondrial metabolism during embryogenesis.

(**a,b**) Correlation plot and regression analysis between differential BMI changes (ΔBMI) and expression

changes in human semen *MIR-LET-7D-5P* (**a**) and *MIR-LET-7E-5P* (**b**). (**c**) Illustration of experimental

and analytical workflow to study differential mRNA expression in 8-cell blastomeres obtained from

Negative Control, *miR-let-7d-5p*, *miR-let-7e-5p*and *miR-699-5p* injected zygotes. (**d**) Multidimensional

scaling (MDS) of single-cell RNAseq profiles in 8-cell stage blastomeres obtained from Negative

Control (n=22), *miR-let-7d-5p* (n=19), *miR-let-7e-5p* (n=20) and *miR-699-5p* (n=21) injected zygotes.

(**e**) GSEA-GOBP analysis of DEGs depicted in (**d**).

Supplementary figure legends:

Figure S1: Obesity and weight loss elicit reversible changes in serum metabolites associated with

mitochondrial function.

(a,b) Relative abundances of TCA intermediate metabolites (**a**) and carnithin moieties(**b**) as quantified

by LC-MS in serum of F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=7) mice. One-way ANOVA followed

by Tukey's multiple comparisons test were used for statistical analysis. P vales are indicated in each

panel.

Figure S2: Obesity and weight loss elicits reversible induces serum lipids linked to mitochondrial

dysfunction.

(**a,b**) Principal component analysis and PC1-2 representation (**a**) and heatmap illustration (**b**) of n=118

serum lipids from F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=7) mice. (**a,b**) were analyzed and plotted

using Metaboanalyst 6.0. (**c**) Relative abundances of ceramides species in serum of F_0 LF (n=6), F_0 HF

(n=6) and F_0 HF-LF (n=7) mice. One-way ANOVA followed by Tukey's multiple comparisons test

were used for statistical analysis and P vales are indicated in each panel.

Figure S3: Paternal obesity and weight loss cause irreversible gene alterations in F_0 liver.

(**a**) Representative histological (H&E), pan-leukocyte (anti-CD45) and anti-collagen/fibrosis (Sirius

Red) staining in livers of F_0 LF, F_0 HF and F_0 HF-LF mice. (**b-e**) Pathologist (SD) assessment of

macrovesicular (**b**) and microvesicular (**c**) steatosis (**d**) fibrosis and (**e**) lobular inflammation scores in

livers of F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=6) mice. A non-parametric Kruskal-Wallis test

followed by Dunn's multiple comparision test were used for statistical analysis and P vales are indicated

in each panel. (**f**) Multidimensional scaling (MDS) plot of F_0 liver. (**g-i**) Volcano plot of significantly

up-(red) and down-regulated (blue) differentially expressed genes (DEG) in liver from (**g**) F_0 HF vs F_0

LF, 3,495 upregulated and 1,639 downregulated DEG, (**h**) F_0 HF vs F_0 HF-LF, 2,858 upregulated and

1,986 downregulated DEG and (**i**) F_0 HF-LF vs F_0 LF, 459 upregulated and 262 downregulated DEG.

Plots depict log2 transformed fold-changes and log10- transformed false-discovery rates (FDR) of

expression changes. Significant DEG are defined as FDR≤0.05. (**j**) Gene Set Enrichment Analysis

(GSEA) and gene ontology (GO) enrichment of biological processes (GOBP) of DEG shown in (**g-i**).

Figure S4: Reversible mitochondrial alterations in obese F_0 adipose tissue.

(**a**) Multidimensional scaling (MDS) plots of transcriptomic changes in F_0 eWAT as determined by

RNA-seq. Dots represent eWAT from individual animals, n=6 per group. (**b**) Representation and colour

coding of mitochondrial electron transport chain complexes I-V (CI-V, *top*) and log-10 transformed

expression changes and significance of DEG (*bottom*) per CI-V in adipose tissue in eWAT of F_0 LF

(n=6), F_0 HF (n=6) and F_0 HF-LF (n=6) mice.

Figure S5: Reversible mitochondrial alterations in obese F_1 adipose tissue.

(**a**) Multidimensional scaling (MDS) plots of transcriptomic changes in F_1 eWAT as determined by

RNA-seq. Dots represent eWAT from F_1 [F_0 LF] (n=4), F_1 [F_0 HF] (n=6) and F_1 [F_0 HF-LF] (n=6) mice.

(**b**) Log-10 transformed expression changes and significance of DEG (*bottom*) per CI-V in adipose

tissue of eWAT of from F_1 [F_0 LF] (n=4), F_1 [F_0 HF] (n=6) and F_1 [F_0 HF-LF] (n=6) mice. (**c**) Relative

protein abundances (LFQ) of NDUFA8, NDUFS6, CPT1A and TOMM20 in eWAT of F_0 LF (n=6), F_0

HF (n=6), F_0 HF-LF (n=6), F_1 [F_0 LF] (n=6), F_1 [F_0 HF] (n=6), and F_1 of F_0 REV (n=6). One-way

ANOVA followed by Tukey's multiple comparisons test were used for statistical analysis. Data are

presented as mean ± standard error and P vales are indicated in each panel.

Figure S6: Paternal obesity and weight loss cause irreversible gene alterations in F_1 liver.

(**a**) Representative histological (H&E), pan-leukocyte (anti-CD45) and anti-collagen/fibrosis (Sirius

Red) staining in livers of F_1 [F_0 LF], F_1 [F_0 HF]and F_1 [F_0 HF-LF] mice. (**b-e**) Pathologist assessment of

macrovesicular (**b**) and microvesicular (**c**) steatosis, (**d**) lobular inflammation and (**e**) fibrosis scores in

livers of livers of F_1 [F_0 LF] (n=6), F_1 [F_0 HF] (n=6) and F_1 [F_0 HF-LF] (n=6) mice. A non-parametric

Kruskal-Wallis test followed by Dunn's multiple comparision test were used for statistical analysis and

P vales are indicated in each panel. (**f**) Multidimensional scaling (MDS) plot of F_1 liver. (**g-i**) Volcano

plot of up-(red) and down-regulated (blue) genes DEG in liver from (**g**) F_1 [F_0 HF] vs F_1 [F_0 LF] (**h**) F_1

[F_0 HF] vs F_1 [F_0 HF-LF] and (**i**) F_1 [F_0 HF-LF] vs F_1 [F_0 LF]. Plots depict log2 transformed fold-changes

and log10- transformed false-discovery rates (FDR) of expression changes. (**j**) Gene Set Enrichment

Analysis (GSEA) and gene ontology (GO) enrichment of biological processes (GOBP) of DEG shown

in (**g-i**).

Figure S7: Reversible alterations in F_1 adipose mitochondrial proteins and identification of cross-

generational miRNAs.

(**a**) Dot plot of enriched gene ontology terms in clusters I-III depicted of protein heatmap in **Fig.3m**.

(**b**) Heatmap depicting relative abundances of miRNAs in eWAT of F_1 [F_0 LF], F_1 [F_0 HF]and F_1 [F_0 HF-

LF] mice as determined by small RNA-seq. (**c,d**) Abundance of (**c**) *miR-143-3p*, *miR-181a-2-3p*, *miR-*

378, *miR-532-5p* and (**d**) *miR-let-7* isoforms in eWAT of F_1 [F_0 LF], F_1 [F_0 HF] and F_1 [F_0 HF-LF] mice.

An unpaired, two-tailed Student's t-test was used for statistical analysis in (**d**) and P values indicted.

Figure S8: Obesity-associated miR-let-7 impairs adipocyte mitochondrial function via silencing

of DICER1.

(**a**) Multidimensional scaling (MDS) plots of transcriptomic changes in 1°eWA transfected with

Negative Control (n=4), *miR-let-7d-5p* (n=4) and *miR-let-7e-5p* (n=4) mimetics as determined by

mRNA-seq. (**b**) GSEA-GOBP analysis of DEGs in 1°eWA transfected with Negative Control (n=4),

miR-let-7d-5p (n=4) and *miR-let-7e-5p* (n=4) mimics. (**c,d**) Western blot analysis (**c**) and densitometric

quantification (**d**) of indicated proteins in 1°eWA transfected with Negative Control, *miR-let-7d-5p* and

miR-let-7e-5p mimetics and analyzed using indicated antibodies. Heat shock protein (HSC) 70 antibody

was used as loading control and the blots shown are representative of three independent experiments.

Superscript letters represent group comparisions **a**, *p*<0.05 Negative Control vs *miR-let-7d-5p*; **b**,

p<0.05, *miR-let-7d-5p* vs *miR-let-7e-5p*; **c**, *p*<0.05, Negative Control vs *miR-let-7e-5p*). (**e**) *Dicer1*

mRNA levels in 1°eA after transfection with 100 nM Negative Control locked nucleic acids (LNA) and

two independent LNA inhibitors targeting *Dicer1* (Dicer1 LNA#1, Dicer1 LNA#2). Data represent

three independent experiments, with each experiment performed in three technical replicates. Unpaired,

two-tailed Student's T-tests were performed, and P values are indicated within the panel. (**f,g**) Western

blot analysis (**f**) and densitometric quantification (**g**) of DICER1 protein levels in 1°eWA transfected

after transfection with 100 nM Negative Control LNAs and two LNAs targeting *Dicer1*. Data represent

three independent experiments, with each experiment performed in three technical replicates. Unpaired,

two-tailed Student's T-tests were performed, and P values are indicated within the panel.

References:

1   Trerotola, M., Relli, V., Simeone, P. & Alberti, S. Epigenetic inheritance and the missing
989   heritability. *Human genomics* **9**, 17 (2015). https://doi.org:10.1186/s40246-015-0041-3

990   2   Bygren, L. O., Kaati, G. & Edvinsson, S. Longevity determined by paternal ancestors'
991   nutrition during their slow growth period. *Acta Biotheor* **49**, 53-59 (2001).
992   https://doi.org:10.1023/a:1010241825519

993   3   Pembrey, M. E. *et al.* Sex-specific, male-line transgenerational responses in humans.
994   *European journal of human genetics : EJHG* **14**, 159-166 (2006).
995   https://doi.org:10.1038/sj.ejhg.5201538

996   4   Lumey, L. H., Stein, A. D., Kahn, H. S. & Romijn, J. A. Lipid profiles in middle-aged men
997   and women after famine exposure during gestation: the Dutch Hunger Winter Families Study.
998   *Am J Clin Nutr* **89**, 1737-1743 (2009). https://doi.org:10.3945/ajcn.2008.27038

999   5   Carone, B. R. *et al.* Paternally induced transgenerational environmental reprogramming of
1000   metabolic gene expression in mammals. *Cell* **143**, 1084-1096 (2010).
1001   https://doi.org:10.1016/j.cell.2010.12.008

1002   6   Ng, S. F. *et al.* Paternal high-fat diet consumption induces common changes in the
1003   transcriptomes of retroperitoneal adipose and pancreatic islet tissues in female rat offspring.
1004   *FASEB journal : official publication of the Federation of American Societies for*
1005   *Experimental Biology* **28**, 1830-1841 (2014). https://doi.org:10.1096/fj.13-244046

1006   7   de Castro Barbosa, T. *et al.* High-fat diet reprograms the epigenome of rat spermatozoa and
1007   transgenerationally affects metabolism of the offspring. *Molecular metabolism* **5**, 184-197
1008   (2016). https://doi.org:10.1016/j.molmet.2015.12.002

1009   8   de Castro Barbosa, T., Alm, P. S., Krook, A., Barres, R. & Zierath, J. R. Paternal high-fat diet
1010   transgenerationally impacts hepatic immunometabolism. *FASEB journal : official publication*
1011   *of the Federation of American Societies for Experimental Biology* **33**, 6269-6280 (2019).
1012   https://doi.org:10.1096/fj.201801879RR

1013   9   Wang, Y. *et al.* Sperm microRNAs confer depression susceptibility to offspring. *Sci Adv* **7**
1014   (2021). https://doi.org:10.1126/sciadv.abd7605

1015   10   Dias, B. G. & Ressler, K. J. Parental olfactory experience influences behavior and neural
1016   structure in subsequent generations. *Nature neuroscience* **17**, 89-96 (2014).
1017   https://doi.org:10.1038/nn.3594

1018   11   Gapp, K. *et al.* Implication of sperm RNAs in transgenerational inheritance of the effects of
1019   early trauma in mice. *Nature neuroscience* **17**, 667-669 (2014).
1020   https://doi.org:10.1038/nn.3695

1021   12   Sun, W. *et al.* Cold-induced epigenetic programming of the sperm enhances brown adipose
1022   tissue activity in the offspring. *Nature medicine* **24**, 1372-1383 (2018).
1023   https://doi.org:10.1038/s41591-018-0102-y

Skvortsova, K., Iovino, N. & Bogdanovic, O. Functions and mechanisms of epigenetic inheritance in animals. *Nature reviews. Molecular cell biology* **19**, 774-790 (2018). https://doi.org:10.1038/s41580-018-0074-2

Rechavi, O., Minevich, G. & Hobert, O. Transgenerational inheritance of an acquired small RNA-based antiviral response in C. elegans. *Cell* **147**, 1248-1256 (2011). https://doi.org:10.1016/j.cell.2011.10.042

Fire, A. *et al.* Potent and specific genetic interference by double-stranded RNA in Caenorhabditis elegans. *Nature* **391**, 806-811 (1998). https://doi.org:10.1038/35888

Ashe, A. *et al.* piRNAs can trigger a multigenerational epigenetic memory in the germline of C. elegans. *Cell* **150**, 88-99 (2012). https://doi.org:10.1016/j.cell.2012.06.018

de Vanssay, A. *et al.* Paramutation in Drosophila linked to emergence of a piRNA-producing locus. *Nature* **490**, 112-115 (2012). https://doi.org:10.1038/nature11416

Rassoulzadegan, M. *et al.* RNA-mediated non-mendelian inheritance of an epigenetic change in the mouse. *Nature* **441**, 469-474 (2006). https://doi.org:10.1038/nature04674

Gross, N., Kropp, J. & Khatib, H. MicroRNA Signaling in Embryo Development. *Biology (Basel)* **6** (2017). https://doi.org:10.3390/biology6030034

Natt, D. *et al.* Human sperm displays rapid responses to diet. *PLoS Biol* **17**, e3000559 (2019). https://doi.org:10.1371/journal.pbio.3000559

Tomar, A. *et al.* Epigenetic inheritance of diet-induced and sperm-borne mitochondrial RNAs. *Nature* **630**, 720-727 (2024). https://doi.org:10.1038/s41586-024-07472-3

Grandjean, V. *et al.* RNA-mediated paternal heredity of diet-induced obesity and metabolic disorders. *Scientific reports* **5**, 18193 (2015). https://doi.org:10.1038/srep18193

Rodgers, A. B., Morgan, C. P., Leu, N. A. & Bale, T. L. Transgenerational epigenetic programming via sperm microRNA recapitulates effects of paternal stress. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 13699-13704 (2015). https://doi.org:10.1073/pnas.1508347112

Sharma, U. *et al.* Biogenesis and function of tRNA fragments during sperm maturation and fertilization in mammals. *Science* (2015). https://doi.org:10.1126/science.aad6780

Korhonen, H. M. *et al.* Dicer is required for haploid male germ cell differentiation in mice. *PloS one* **6**, e24821 (2011). https://doi.org:10.1371/journal.pone.0024821

Bernstein, E. *et al.* Dicer is essential for mouse development. *Nature genetics* **35**, 215-217 (2003). https://doi.org:10.1038/ng1253

Chen, Q. *et al.* Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic disorder. *Science* (2015). https://doi.org:10.1126/science.aad7977

Rodgers, A. B., Morgan, C. P., Bronson, S. L., Revello, S. & Bale, T. L. Paternal stress exposure alters sperm microRNA content and reprograms offspring HPA stress axis

regulation. *The Journal of neuroscience : the official journal of the Society for Neuroscience*

33, 9003-9012 (2013). https://doi.org:10.1523/JNEUROSCI.0914-13.2013

29   Cropley, J. E., Suter, C. M., Beckman, K. B. & Martin, D. I. Germ-line epigenetic

modification of the murine A vy allele by nutritional supplementation. *Proceedings of the*

National Academy of Sciences of the United States of America **103**, 17308-17312 (2006).

https://doi.org:10.1073/pnas.0607090103

30   Zhang, Y., Shi, J., Rassoulzadegan, M., Tuorto, F. & Chen, Q. Sperm RNA code programmes

the metabolic health of offspring. *Nature reviews. Endocrinology* **15**, 489-498 (2019).

https://doi.org:10.1038/s41574-019-0226-2

31   Sun, K., Kusminski, C. M. & Scherer, P. E. Adipose tissue remodeling and obesity. *The*

Journal of clinical investigation **121**, 2094-2101 (2011). https://doi.org:10.1172/JCI45887

32   Scherer, P. E. Adipose tissue: from lipid storage compartment to endocrine organ. *Diabetes*

55, 1537-1545 (2006). https://doi.org:10.2337/db06-0263

33   Lee, J. H. *et al.* The Role of Adipose Tissue Mitochondria: Regulation of Mitochondrial

Function for the Treatment of Metabolic Diseases. *Int J Mol Sci* **20** (2019).

https://doi.org:10.3390/ijms20194924

34   Czech, M. P. Mechanisms of insulin resistance related to white, beige, and brown adipocytes.

Molecular metabolism **34**, 27-42 (2020). https://doi.org:10.1016/j.molmet.2019.12.014

35   Kahn, C. R., Wang, G. & Lee, K. Y. Altered adipose tissue and adipocyte function in the

pathogenesis of metabolic syndrome. *The Journal of clinical investigation* **129**, 3990-4000

(2019). https://doi.org:10.1172/JCI129187

36   Bartelt, A. & Heeren, J. Adipose tissue browning and metabolic health. *Nature reviews.*

Endocrinology **10**, 24-36 (2014). https://doi.org:10.1038/nrendo.2013.204

37   Schottl, T., Kappler, L., Braun, K., Fromme, T. & Klingenspor, M. Limited mitochondrial

capacity of visceral versus subcutaneous white adipocytes in male C57BL/6N mice.

Endocrinology **156**, 923-933 (2015). https://doi.org:10.1210/en.2014-1689

38   Heinonen, S., Jokinen, R., Rissanen, A. & Pietilainen, K. H. White adipose tissue

mitochondrial metabolism in health and in obesity. *Obes Rev* **21**, e12958 (2020).

https://doi.org:10.1111/obr.12958

39   Kusminski, C. M. *et al.* MitoNEET-driven alterations in adipocyte mitochondrial activity

reveal a crucial adaptive process that preserves insulin sensitivity in obesity. *Nature medicine*

18, 1539-1549 (2012). https://doi.org:10.1038/nm.2899

40   Gonzalez-Franquesa, A. *et al.* Remission of obesity and insulin resistance is not sufficient to

restore mitochondrial homeostasis in visceral adipose tissue. *Redox Biol* **54**, 102353 (2022).

https://doi.org:10.1016/j.redox.2022.102353

41   Schottl, T., Kappler, L., Fromme, T. & Klingenspor, M. Limited OXPHOS capacity in white

adipocytes is a hallmark of obesity in laboratory mice irrespective of the glucose tolerance

status. *Molecular metabolism* **4**, 631-642 (2015).
https://doi.org:10.1016/j.molmet.2015.07.001

42 Soro-Arnaiz, I. *et al.* Role of Mitochondrial Complex IV in Age-Dependent Obesity. *Cell*
reports **16**, 2991-3002 (2016). https://doi.org:10.1016/j.celrep.2016.08.041

43 Wang, Y. *et al.* Metformin Improves Mitochondrial Respiratory Activity through Activation
of AMPK. *Cell reports* **29**, 1511-1523 e1515 (2019).
https://doi.org:10.1016/j.celrep.2019.09.070

44 Borralho, P. M., Rodrigues, C. M. & Steer, C. J. microRNAs in Mitochondria: An
Unexplored Niche. *Adv Exp Med Biol* **887**, 31-51 (2015). https://doi.org:10.1007/978-3-319-
22380-3_3

45 Bandiera, S. *et al.* Nuclear outsourcing of RNA interference components to human
mitochondria. *PloS one* **6**, e20746 (2011). https://doi.org:10.1371/journal.pone.0020746

46 Barrey, E. *et al.* Pre-microRNA and mature microRNA in human mitochondria. *PloS one* **6**,
e20220 (2011). https://doi.org:10.1371/journal.pone.0020220

47 Mori, M. A. *et al.* Role of MicroRNA Processing in Adipose Tissue in Stress Defense and
Longevity. *Cell metabolism* **16**, 336-347 (2012). https://doi.org:S1550-4131(12)00322-1 [pii]
10.1016/j.cmet.2012.07.017

48 Mori, M. A. *et al.* Altered miRNA processing disrupts brown/white adipocyte determination
and associates with lipodystrophy. *The Journal of clinical investigation* **124**, 3339-3351
(2014). https://doi.org:10.1172/JCI73468

49 Oliverio, M. *et al.* Dicer1-miR-328-Bace1 signalling controls brown adipose tissue
differentiation and function. *Nature cell biology* **18**, 328-336 (2016).
https://doi.org:10.1038/ncb3316

50 Turner, N. *et al.* A selective inhibitor of ceramide synthase 1 reveals a novel role in fat
metabolism. *Nature communications* **9**, 3165 (2018). https://doi.org:10.1038/s41467-018-
05613-7

51 Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. *Proceedings of the National Academy of*
Sciences of the United States of America **102**, 15545-15550 (2005).
https://doi.org:0506580102 [pii]
10.1073/pnas.0506580102

52 Brieno-Enriquez, M. A. *et al.* Exposure to endocrine disruptor induces transgenerational
epigenetic deregulation of microRNAs in primordial germ cells. *PloS one* **10**, e0124296
(2015). https://doi.org:10.1371/journal.pone.0124296

53 Zhu, H. *et al.* The Lin28/let-7 axis regulates glucose metabolism. *Cell* **147**, 81-94 (2011).
https://doi.org:10.1016/j.cell.2011.08.033

54 Frost, R. J. & Olson, E. N. Control of glucose homeostasis and insulin sensitivity by the Let-7
family of microRNAs. *Proceedings of the National Academy of Sciences of the United States*
of America **108**, 21075-21080 (2011). https://doi.org:10.1073/pnas.1118922109

55 Giroud, M. *et al.* Let-7i-5p represses brite adipocyte function in mice and humans. *Scientific*
reports **6**, 28613 (2016). https://doi.org:10.1038/srep28613

56 O'Connor, S. *et al.* AGO HITS-CLIP reveals distinct miRNA regulation of white and brown
adipose tissue identity. *Genes & development* (2021). https://doi.org:10.1101/gad.345447.120

57 Fullston, T. *et al.* Paternal obesity initiates metabolic disturbances in two generations of mice
with incomplete penetrance to the F2 generation and alters the transcriptional profile of testis
and sperm microRNA content. *FASEB journal : official publication of the Federation of*
American Societies for Experimental Biology **27**, 4226-4243 (2013).
https://doi.org:10.1096/fj.12-224048

58 Liu, W. *et al.* Expression of microRNA let-7 in cleavage embryos modulates cell fate
determination and formation of mouse blastocystsdagger. *Biol Reprod* **107**, 1452-1463
(2022). https://doi.org:10.1093/biolre/ioac181

59 Cheong, A. W. *et al.* MicroRNA Let-7a and dicer are important in the activation and
implantation of delayed implanting mouse embryos. *Human reproduction* **29**, 750-762
(2014). https://doi.org:10.1093/humrep/det462

60 Tobi, E. W. *et al.* DNA methylation signatures link prenatal famine exposure to growth and
metabolism. *Nature communications* **5**, 5592 (2014). https://doi.org:10.1038/ncomms6592

61 Schulz, L. C. The Dutch Hunger Winter and the developmental origins of health and disease.
Proceedings of the National Academy of Sciences of the United States of America **107**, 16757-
16758 (2010). https://doi.org:10.1073/pnas.1012911107

62 Jepsen, I. E. *et al.* Turning waiting time into treatment time: Weight reduction by a lifestyle
intervention programme for patients with obesity before fertility treatment. *Reproductive,*
Female and Child Health **2**, 133-142 (2023). https://doi.org:https://doi.org/10.1002/rfc2.46

63 Yuan, S. *et al.* Sperm-borne miRNAs and endo-siRNAs are important for fertilization and
preimplantation embryonic development. *Development* **143**, 635-647 (2016).
https://doi.org:10.1242/dev.131755

64 Suh, N. & Blelloch, R. Small RNAs in early mammalian development: from gametes to
gastrulation. *Development* **138**, 1653-1661 (2011). https://doi.org:10.1242/dev.056234

65 Pfisterer, U. *et al.* Identification of epilepsy-associated neuronal subtypes and gene expression
underlying epileptogenesis. *Nature communications* **11**, 5038 (2020).
https://doi.org:10.1038/s41467-020-18752-7

66 Popov, L. D. Mitochondrial biogenesis: An update. *J Cell Mol Med* **24**, 4892-4899 (2020).
https://doi.org:10.1111/jcmm.15194

67   Brandao, B. B. *et al.* Dynamic changes in DICER levels in adipose tissue control metabolic
adaptations to exercise. *Proceedings of the National Academy of Sciences of the United States*
of America **117**, 23932-23941 (2020). https://doi.org:10.1073/pnas.2011243117

68   Sullivan, R. & Saez, F. Epididymosomes, prostasomes, and liposomes: their roles in
mammalian male reproductive physiology. *Reproduction* **146**, R21-35 (2013).
https://doi.org:10.1530/REP-13-0058

69   Tan, K. *et al.* Downregulation of miR-199a-5p Disrupts the Developmental Potential of In
Vitro-Fertilized Mouse Blastocysts. *Biol Reprod* **95**, 54 (2016).
https://doi.org:10.1095/biolreprod.116.141051

70   Chambers, M. C. *et al.* A cross-platform toolkit for mass spectrometry and proteomics.
Nature biotechnology **30**, 918-920 (2012). https://doi.org:10.1038/nbt.2377

71   Kind, T. *et al.* LipidBlast in silico tandem mass spectrometry database for lipid identification.
Nature methods **10**, 755-758 (2013). https://doi.org:10.1038/nmeth.2551

72   Luan, H., Ji, F., Chen, Y. & Cai, Z. statTarget: A streamlined tool for signal drift correction
and interpretations of quantitative mass spectrometry-based omics data. *Anal Chim Acta*
1036, 66-72 (2018). https://doi.org:10.1016/j.aca.2018.08.002

73   Kleiner, D. E. *et al.* Design and validation of a histological scoring system for nonalcoholic
fatty liver disease. *Hepatology* **41**, 1313-1321 (2005). https://doi.org:10.1002/hep.20701

74   Rappsilber, J. & Mann, M. Analysis of the topology of protein complexes using cross-linking
and mass spectrometry. *CSH Protoc* **2007**, pdb prot4594 (2007).
https://doi.org:10.1101/pdb.prot4594

75   Tyanova, S., Mann, M. & Cox, J. MaxQuant for in-depth analysis of large SILAC datasets.
Methods in molecular biology **1188**, 351-364 (2014). https://doi.org:10.1007/978-1-4939-
1142-4_24

76   Tyanova, S. *et al.* The Perseus computational platform for comprehensive analysis of
(prote)omics data. *Nature methods* **13**, 731-740 (2016). https://doi.org:10.1038/nmeth.3901

77   Frankish, A. *et al.* GENCODE reference annotation for the human and mouse genomes.
Nucleic acids research **47**, D766-D773 (2019). https://doi.org:10.1093/nar/gky955

78   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).
https://doi.org:10.1093/bioinformatics/bts635

79   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).
https://doi.org:10.1093/bioinformatics/btt656

80   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140
(2010). https://doi.org:10.1093/bioinformatics/btp616

81   Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: Precision weights unlock linear model

analysis tools for RNA-seq read counts. *Genome biology* **15**, R29 (2014).

https://doi.org:10.1186/gb-2014-15-2-r29

82   Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.

Innovation (Camb) **2**, 100141 (2021). https://doi.org:10.1016/j.xinn.2021.100141

1210

Figure 1: Obesity and weight loss elicit reversible changes in glucose metabolism and mitochondrial metabolites

Figure 2: Multimodal omics analysis reveals reversible mitochondrial gene aberrations in obese adipose tissue

Figure 3: Paternal obesity impairs glucose tolerance and instigates adipose dysfunction in F$_1$ adipose tissue.

Figure 4: Obesity induces adipose tissue miR-let-7 and blunts mitochondrial function via a novel miR-let-7-DICER1 axis.

Figure 5: Zygotic microinjection of obesity-associated miR-let-7 causes glucose tolerance in sired mice.

Figure 6: Obesity-associated miR-let-7 rewires mitochondrial metabolism during embryogenesis

Figure S1: Obesity and weight loss elicits reversible changes in metabolites associated with mitochondrial function.

Figure S2: Obesity and weight loss elicits reversible changes in serum lipids controlling mitochondrial activity

Figure S3: Paternal obesity and weight loss cause irreversible gene alterations in F₀ liver

Figure S4: Reversible mitochondrial alterations in obese F_0 adipose tissue

Figure S5: Reversible mitochondrial alterations in F$_1$ adipose tissue

Figure S6: Paternal obesity and weight loss cause **irreversible** gene alterations in F₁ liver

Figure S7: Reversible alterations in F₁ adipose mitochondria and identification of cross-generational miRNAs

Figure S8: Obesity-associated miR-let-7 impairs adipocyte mitochondrial function via silencing of DICER1