Male obesity causes adipose mitochondrial dysfunction in F$_1$ progeny via a let-7-DICER axis.

Chien Huang[1,2#], Joo Hyun Park[1#], Ali Altıntaş [3], Natasa Stanic[1], Kristine Kyle de Leon[1], Signe Isacson[4], Panagiotis Kalogeropoulos[5], Hande Topel Batarlar[1], Rocio Valdebenito Malmros[1], Jesper Havelund[1], Bjørk Ditlev Marcher Larsen[1], Yen-Ting Chien[2], Wen-Chi Huang[2], Karolina Szczepanowska[6,7], Jan-Wilm Lackmann[6], Aleksandra Trifunovic[6], Eva Kildall Hejbøl[8], Sönke Detlefsen[8,9], Ida Engberg Jepsen[10], Stefanie Hansborg Kolstrup[11], Ricardo Laguna-Barraza[12], Javier Martin-Gonzalez[12], Konstantin Khodosevich[13], Nils J. Færgeman[1], Marcelo A. Mori[14-17], Marc R. Friedländer[5], Anita Oest[4], Romain Barrès[3,18] and Jan-Wilhelm Kornfeld[1,19*]

[1]*Department of Biochemistry and Molecular Biology (BMB), University of Southern Denmark (SDU), Odense, Denmark.* [2]*Laboratory of Animal Physiology, Department of Animal Science and Technology, National Taiwan University, Taipei, Taiwan.* [3]*Novo Nordisk Foundation Center for Basic Metabolic Research (CBMR), Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark.* [4]*Division of Cell and Neurobiology, Department of Biomedical and Clinical Sciences, Linköping University, Sweden.* [5]*Science for Life Laboratory, Department of Molecular Biosciences, The Wenner-Gren-Institute, Stockholm University, Sweden.* [6]*Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD) & Center for Molecular Medicine (CMMC), University of Cologne, Cologne, Germany.* [7]*IMol Polish Academy of Sciences, 02-247 Warsaw, Poland.* [8]*Department of Pathology, Odense University Hospital, Odense, Denmark.* [9]*Department of Clinical Research, Faculty of Health Science, University of Southern Denmark, Odense, Denmark.* [10] *The Fertility Clinic, Department of Obstetrics and Gynaecology, Zealand University Hospital, Koege, Denmark.* [11]*Biomedical Laboratory, Institute of Molecular Medicine, University of Southern Denmark, Odense, Denmark.* [12]*Core Facility for Transgenic Mice, Department of Experimental Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark.* [13]*Biotech Research & Innovation Centre (BRIC), University of Copenhagen, Copenhagen, Denmark.* [14]*Department of Biochemistry and Tissue Biology, Institute of Biology, Universidade Estadual de Campinas, Campinas, São Paulo, Brazil.* [15]*Program in Genetics and Molecular Biology, Institute of Biology, Universidade*

Estadual de Campinas, Campinas, São Paulo, Brazil. [16]Obesity and Comorbidities Research Center

(OCRC), Universidade Estadual de Campinas, Campinas, São Paulo, Brazil. [17]Experimental Medicine

Research Cluster (EMRC), Universidade Estadual de Campinas, Campinas, São Paulo, Brazil.

[18]Institut de Pharmacologie Moléculaire et Cellulaire, Université Côte d'Azur & Centre National pour

la Recherche Scientifique (CNRS), Valbonne, 06560, France. [19]Novo Nordisk Foundation Center for

Adipocyte Signalling, University of Southern Denmark, Odense, Denmark.

#These authors contributed equally to this work,

*Corresponding author

Abstract

We here describe that obesity and weight loss in male mice cause reversible abnormalities in glucose and

lipid metabolism, serum metabolomes and lipidomes as well as expression of microRNAs, mRNAs and

proteins controlling mitochondrial function in epididymal white adipose tissue. When mating obese male

mice with lean females, we observed reductions in expression and translation of genes encoding

mitochondrial respiratory components in (F_1) offspring that closely resemble those observed in the paternal

(F_0) generation. When mapping miRNA regulation across somatic organs (i.e., liver, adipose) and sperm and

$F_{0/1}$ generations, we found that obesity and weight loss reversibly affected miRNA levels, and that *let-7*

isoforms were induced in obese F_0 and F_1 adipose tissue and sperm of obese F_0 mice, eliciting qualitatively

similar responses in two adjacent tissues. Overexpressing *let-7* in adipocytes silenced DICER1, a miRNA

processing enzyme crucial for adipose adaptation to obesity as evidenced by deficiencies in mitochondrial

function following DICER1 loss in primary adipocytes. Also, microinjection of synthetic *let-7* mimetics at

physiological levels found in obese sperm into zygotes from lean mice elicited glucose intolerance and

impediments in adipose mitochondrial gene expression in mice sired from *let-7* microinjected zygotes,

phenocopying hereditary aspects of paternal obesity. When performing single-cell RNA-Seq of miRNA-

injected embryos, *let-7* impaired mitochondrial gene expression, suggesting altered oxidative metabolism

following zygotic *let-7* delivery. When studying miRNA alterations in human semen, lifestyle-induced

weight loss downregulated *hsa-let-7*, suggesting similar roles for human *let-7* in gametic epigenomes and

embryogenesis.

Introduction:

Intergenerational epigenetic inheritance of metabolic dysfunction

Genome-wide association studies have demonstrated that genetic (i.e., DNA) variants only explain 20% of obesity heritability[1]. Given pandemic rises in obesity in evolutionarily relatively short timescales, it was proposed that non-genetic or ,*epi-genetic*' processes, that are spurred by obesogenic lifestyles and other changes in environment exposures, might increase obesity susceptibility and contribute to disease incidences. This form of epigenetic inheritance, if convincingly demonstrated[2], is particularly intriguing for cases of paternal programming where affected fathers contribute only sperm as biological material to embryos (intergenerational inheritance) or no genetic/biological material at all (transgenerational inheritance). Epidemiological support for the existence of hereditary, paternally-acquired traits comes from historical records demonstrating that ample ancestral food supplies affects cardiometabolic traits up to two generations[3,4], showing that individuals carry increased risks of cardiometabolic dysfunction if their parents or grandparents were obese or were malnourished[5]. In line with correlative observations in human, mechanistic studies in isogenic (genetically identical) mouse strains like C57BL76 demonstrated that paternal obesity can impair metabolism in offsprings (F_1 generation and beyond) and, specific diet like low-protein[6] or high-fat diet feeding[7-9] and stress-related cues such as depression-like behavior[10], fear-conditioning[11] and psychological traumata[12] impart specific epigenetic behavioral and metabolic effects in offsprings. Recently, beneficial types of epigenetic programming in adipose tissue were also demonstrated where offsprings of mice exposed to cold temperatures display increased amounts of metabolically favourable brown adipose tissue [13], suggesting an unexpectedly high plasticity of the molecular and cellular mechanisms underlying intergenerational epigenetic responses.

The molecular basis for epigenetic inheritance of paternal obesity in male gametes

The molecular and cellular basis for epigenetic heredity remains largely enigmatic: Although changes in epigenetic marks like methylation of genomic DNA, histone post-translational modifications and alterations in small Noncoding RNAs (sncRNAs) correlate with hereditary processes in male gametes[14]. Although lifestyle/diet-induced changes to DNA methylation and histones modifications were reported,

the evidence for causatively implicating these in eliciting paternal response in progeny is rare for higher

mammalian species like rodents and humans. Other emerging instigators of intergenerational response

are found amongst the different types of gametic sncRNAs such as microRNAs (miRNAs): miRNAs in

gametes were initially identified and functionally dissected in nematodes and fruit flies[15-18], thus species

where sncRNAs-evoked gametic gene silencing processes are well-understood [19]. In contrast to DNA-

linked marks, sncRNAs are very mobile in nature and are subject to microvesicular trafficking between

cells and organs and secreted by somatic cell types in testes[20,21]. Thus, due to their high abundance of

miRNAs in sperm, their dynamic changes following diet exposures[22,23], their exchange between sperm

and oocytes[23] sperm-born sncRNAs have the unique potential for exchanging information about past

lifestyle choices to zygotes and F_1 progeny. Consistent with this, metabolic trait/disease-associated

miRNAs can affect intergenerational phenotypes and phenocopy F_0 lifestyles if microinjected [24,25] or

removed from zygotes[26]. Beyond controlling gametic epigenomes, miRNAs are also essential for germ

cell development processes such as spermatogenesis[27] and embryogenesis[28], and miRNA are exchanged

between zygotes and other cell types [12,25,29-31]. Taken together, it is plausible to hypothesize that obesity-

evoked changes in sperm miRNAs alter zygotic miRNA pools and embryonic development, instigating

epigenetic phenotypic traits and somatic gene expression in offsprings. As diverse paternal lifestyle

exposures ranging from inflammation[15], nutrition[29,31] to stress [12,30] alter gametic miRNA repertoires and

F_1 phenotypes in non-overlapping ways, the intriguing possibility exists that environmental experiences

are encoded in individual gametic sncRNA signature and that these sperm sncRNAs have the potential

to transmit complex lifestyle exposures given the high bandwidth of RNA-mediated effects[32].

Regulation of adipose tissue (mitochondrial) function by miRNAs

Adipose tissue is a multifunctional endocrine organ that coordinates tissue-level (autocrine, juxtacrine,

paracrine) and systemic (endocrine) responses by release of adipokines, lipids and inflammatory factors

and, shown recently, miRNA-containing microvesicles (exosomes)[33-35]. It is believed that adipose tissue

can safely store ingested nutrients as lipids, although chronic calorie excess causes insulin resistance[36],

aberrant gene expression in adipocytes[37], vascular remodelling and immune cell infiltration[33].

Mitochondria in adipose tissue exert important functions for tissue/energy homeostasis, particularly

during nutrient deprivation/excess[35]: Whereas brown adipose tissue (BAT) contains large numbers of

mitochondria to generate heat by dissipation of the electron transport chain (ETC) proton gradients[38],

inguinal (beige, iWAT) and epididymal (white, eWAT) adipose tissue protects from calorie overload

by removing nutrients from circulation and storage as energetic, indolent macromolecules such as

triglycerides. Mitochondrial function in WAT is impaired in obesity [39] and reduced mitochondrial

function in fat causes metabolic imbalance[40-43]. Mitochondrial DNA content, oxidative phosphorylation

(OxPhos), abundance and composition of ETC complexes are all functionally impaired in obesity and

morphological aberrations in mitochondrial structures are found in obesity[39,43] and aging[44]. Conversely,

treatment with the AMP-activated protein kinase (AMPK) agonist Metformin improves mitochondrial

function in adipose tissue, as do exercise or calorie restriction, and all these interventions correlate with

metabolic improvement[45]. Recently, miRNAs were identified as important regulators of mitochondrial

formation and function and known ‚mito-miRs' like *miR-143, miR-181a, miR-378* instigate, whereas

miR-532 which impairs mitochondrial function by (post-transcriptionally) silencing mitochondrial

translocation and mt-RNA homeostasis[46-48]. We and others showed that removing miRNAs from

adipocytes by conditionally ablating DICER1, a central component of the conserved miRNA processing

machinery, impairs adipogenesis, triggering lipodystrophy and metabolic dysfunction in mice[49-51].

Collectively, we report that obesity in male mice is linked to reversible miRNA, mRNA and proteins

alterations in adipose mitochondrial gene regulation that match those in adipose tissue of lean isogenic

F_1 progeny. Using integrative analyses of adipose, liver and sperm miRNA:mRNA gene networks, we

identify *let-7* as induced by obesity across cell types and generations and mechanistically implicate *let-*

7 driven suppression of DICER1 and concomitant declines in mitochondrial function in adipocytes in

inherited mouse phenotypes, thereby establishing sperm-born *let-7* as intergenerational driver of

mitochondrial dysfunction and glucose impairments in F_1 offspring.

Results

Paternal obesity elicits intergenerational impairments in adipose mitochondrial function

Whilst intergenerational effects of paternal (F_0) obesity on offspring (F_1) phenotypes and metabolic

health have been reported, the specific organ-level manifestations of this dysfunction in progeny, and

the precise somatic and germ cell-intrinsic molecular mechanisms in obese mice and their descendants,

remain surprisingly elusive. To investigate this using isogenic rodents (*Mus musculus*), we used diet-

induced obese (DIO) and weight-regressed C57BL/6N male mice, and fed 6-week-old animals with 1)

low-fat diet (LF) for 18 weeks (F_0 LF), 2) high-fat diet (HF) for 18 weeks (F_0 HF) or 3) HF for 9 weeks,

followed by 9 weeks of LF-driven weight loss (F_0 HF-LF, **Fig.1a**) to allow at least three cycles of

spermatogenesis. For each F_0 group, we mated eight randomly selected mice with unexposed lean

virgin, female C57BL/6N mice, and exposed both male and female F_1 progeny of each group, i.e., F_1

(F_0 LF, n=44 pups), F_1 (F_0 HF, n=52 pups) and F_1 (F_0 HF-LF, n=38 pups) mice to LF after weaning to

test for metabolic and metabolic sequelæ of F_0 diet exposure and removal. We did not additionally

challenge F_1 mice with HF given that effects of HF and epigenetic F_0 effects were reported as

cumulative with regards to F_1 phenotypes [52] (**Fig.1b**). Noteworthy, we detected no alterations in

fecundity, male-to-female pup ratio and litters sizes (**Fig.S1a,b**), suggesting that F_0 obesity did not

affect development and embryonic implantation rates nor indirectly affected maternal provisioning after

birth due to different litter sizes, two potential confounders.

When phenotyping F_0 and male F_1 animals for metabolic traits commonly associated with obesity and

type 2 diabetes (T2D), we found that HF increased and ensuing HF-LF feeding reduced body weight to

levels seen in F0 LF mice, whilst F_1 progeny were indistinguishable in terms of body weight (**Fig.1c**).

Intriguingly, male obesity impaired glucose tolerance (**Fig.1d, Fig.S2a**) and insulin sensitivity (**Fig.1e**)

not only in F_0 HF mice, but also in their F_1 (F_0 HF) offsprings without changes in circulating F_1 insulin

(**Fig.S2b**) and no alterations in liver and brown adipose tissue mass (not shown). In contrast, we found

that obesity increased iWAT (**Fig.S2c**) and eWAT (**Fig.1f**) masses and elevated serum levels of leptin,

an important adipokine reflecting adipocyte size and numbers, in F_0 and F_1 mice (**Fig.1g**). Adipocyte

size determination using haematoxylin/eosine (H/E) staining of formalin-fixated, paraffine-embedded

adipose sections demonstrated comparable shifts towards increased eWAT adipocyte sizes in F_0 and F_1

(**Fig.1h-j**), suggesting that paternal obesity might promote lipid accrual/lipolysis as well as metabolic

processes governing adipocyte health by similar molecular mechanisms in both generations and via this

mechanism contribute to glucose intolerance in F_0 and male F_1. When phenotyping female offsprings

of lean, obese and weight-regressed F_0 fathers, we observed no changes in F_1 weight (**Fig. S2d**), mild

glucose intolerance (**Fig.S2e,f**) and only trends towards changes in insulin sensitivity (**Fig.S2g**). In

support of overall mild impairments in female offsprings, random fed glycemia (**Fig.S2h**) and organ

weights (**Fig.S2i**) were also unchanged in female progeny. *Thus, male obesity and weight loss reversibly*

affect glucose tolerance in F_1, with metabolic effects predominantly observed in male progeny.

For comprehensive insights into molecular processes affected by paternal obesity and weight loss, and

those governing intergenerational effects, we performed bulk tissue mass spectrometry to determine

differentially abundant proteins in F_0 and F_1 eWAT. We observed 980/431 proteins increased in F_0 HF

compared to F_0 LF mice and in F_1 (F_0 HF) compared to F_1 (F_0 LF) mice. Conversely, 632/305 proteins

were less abundant comparing F_0 HF to F_0 LF mice (**Fig.S3a**) and F_1 (F_0 HF) compared to F_1 (F_0 LF)

mice (**Fig.S3b**). Using the STRING database (https://string-db.org/) followed by Gene Ontology (GO)

analysis of overrepresented biological processes (GOBP), we intriguingly saw that F_0 HF mice (**Fig.1k**)

and F_1 (F_0 HF) progeny (**Fig.1l**) shared the same downregulated, mitochondria-associated GOBPs such

as Oxphos, adenosine triphosphate (ATP) synthesis, and regulation of ETC subunits. Examples of

proteins similarly repressed in F_0 HF and F_1 (F_0 HF) eWAT included components of ETC Complex I

like NADH dehydrogenase 1 alpha subcomplex subunit 8 (NDUFA8) and NADH dehydrogenase iron-

sulfur protein 6 (NDUFS6), of Complex IV like Cytochrome C Oxidase Subunit 6B1 (COX6B1), of

mitochondrial membrane-intrinsic lipid carriers such as *Carnitine Palmitoyltransferase 1A* (CPT1A)

and of mitochondrial protein translocation such as *Translocase Of Outer Mitochondrial Membrane 20*

(TOMM20). In contrast, upregulated GOBP terms in F_0 HF mice (**Fig.S3a**) and their F_1 (F_0 HF) progeny

(**Fig.S3b**) included unrelated GOBP terms like protein translation, Golgi-mediated transport and

secretion as well as protein localisation. Importantly, abundance of metabolite- (**Fig.S4a-c**) and lipid-

based (**Fig.S4d,e**) biomarkers that can reflect systemic mitochondrial dysfunction, for instance tricarbon

cycle acid (TCA) intermediates like citrate, malate, fumarate, alpha-ketoglutarate and citrate, were

reduced by paternal obesity and restored by weight loss (**Fig.S4f**) and mitochondrial lipid carriers like

(acetylated) carnitine (**Fig.S4g**) were reduced. Furthermore, saturated, short-chain C:16:0/C:18:0

ceramides, i.e., lipid species reported to disrupt mitochondrial function and elicit mitochondrial ER

stress in obese adipocytes[53,54] were induced, whilst mito-protective long-chain C18:1/C24:1 ceramides

were reduced in F_0 HF mice (**Fig.S4h**). *Thus, male obesity downregulates proteins associated with*

mitochondrial processes in obese F_0 and F_1 offsprings that correlate with reduced levels of

mitochondrial function and lipid catabolism, likely reflecting a system-wide hereditary form of

mitochondrial dysfunction, also in adipose tissue.

Paternal obesity negatively affects expression of mitochondria-associated miRNA-mRNA networks

We next turned our attention to in detail study gene-regulatory processes at the messenger and miRNA

level that might underly the observed repression of mitochondria-associated proteins in F_0 and F_1

adipose tissue, and to this end performed mRNA-seq of eWAT and liver of F_0 and F_1 groups and

performed miRNA and mRNA differential expression analysis, specifically focussing on mitochondrial

processes. Comparing F_0 HF to F_0 LF eWAT using adjusted p-Values (pV_{adj}) ≤ 0.05 we detected yielded

3,265 up-regulated and 2,890 down-regulated genes (**Fig.2a**). When comparing F_1 (F_0 HF) to F_1 (F_0 LF)

mice we detected no significantly regulated genes when accounting for multiple testing (pV_{adj}). Yet

when specifically interrogating differential expression to reflect the more variable transcriptional effects

in F_1 mice, we found 766 up-regulated and 893 down-regulated genes using unadjusted p-Values (pV)

≤ 0.05 (**Fig.2b**). When performing gene set enrichment analysis (GSEA) of affected GOBP we found

upregulated processes linked to cell proliferation and immune activation that likely reflect adipocyte

hyperplasia and inflammation, thus cellular processes commonly associated with obesity, and GOBP

associated with germ cell and cilial function in F_1. Importantly, we confirmed our proteomic findings

and detected overrepresented GOBP related to lipid catabolism, ATP synthesis, OxPhos and ETC

downregulated in adipose tissue of F_0 HF mice (**Fig.2c**) and their F_1 offsprings (**Fig.2d**) and, when

interrogating expression of mRNAs encoding mitochondrial ETC I-V, we confirmed the repression of

nuclear-expressed mitochondrial constituents at the mRNA level (**Fig.2a,b**). Importantly, aberrant

mitochondrial effects were to large extends reversible by weight loss, as comparing F_0 HF to F_0 HF-LF

eWAT yielded 2,687 up-regulated and 2,237 down-regulated using pV_{adj} (**Fig.S5a**), whilst comparing

F_1 (F_0 HF) to F_1 (F_0 HF-LF) eWAT yielded 547 up-regulated and 732 down-regulated genes without

adjusting for multiple testing, (**Fig.S5b**) and GOBP restored by weight loss also fell into genes linked

to mitochondrial function (**Fig.S5c,d**).

Given that miRNAs are important (rheostatic) components in post-transcriptional gene network and

have previously been linked to mitochondrial maturation and function[55], albeit not in adipocytes, we

next wondered whether concerted down-regulation of mitochondrial programs in obese eWAT might

(partially) be instigated by altered expression of mito-miRs[1'] like *let-7[2]*, *miR-378[3]*, *miR-181[4]*, *miR-143[5]*

that functionally support mitochondrial formation and oxidative metabolism. Thus, when conducting

small RNA (sRNA)-seq to determine differential miRNA expression, we found that F_0 obesity repressed

mito-miRs that positively affect mitochondrial biogenesis and function, for instance *miR-143-3p[56]*, *miR-*

181a-2-3p[57,58], *miR-182[59]*, *miR-200[60]* and *miR-378a-3p[61]* in eWAT of F_0 HF (**Fig.2e**) and F_1 (F_0 HF,

Fig.2f) mice, whereas mitochondria-inactivating *miR-532[62]* was induced in F_0 HF (**Fig.6c**) and F_1 (F_0

HF, **Fig.6d**) eWAT, suggesting that miRNA dysregulation contributed to mitochondrial impediments

observed in obese F_0 eWAT. Integration of miRNA-mRNA NGS datasets with adipocyte-specific

miRNA-mRNA-Argonaute 2 interactomes using AGO-HITS-CLIP[6] revealed that *miR-378 and miR-*

143, despite being predicted, did not coinidce with downregulation of their mitochondrial protein targets

(**Fig.2g**), whilst *miR-532-5p* was both predicted to target and observed to coincide with mitochondrial

proteins repressed in F0 obesity, arguing for roles of *miR-532,* but not *miR-143/378* in driving

mitochondria dysfunction in F_0 HF eWAT (**Fig.2h**). In line with body weight loss-driven mRNA

transcriptional responses (**Fig.S6a,b**), miRNA alterations were highly dynamic, and weight loss

restored expression of mitochondria-promoting in F_0 HF (**Fig.S5e**) and F_1 (F_0 HF, **Fig.S5f**) mice. *Thus,*

F_0 obesity causes reversible repression of mitochondrial and lipid catabolic genes, and miRNA-mRNA

network analysis implicates obesity-regulated mito-miRs in this process.

The specificity of obesity-induced heritable epigenetic mitochondrial alterations to adipose tissue

We were in the following interested to address how confined the observed heritability of mitochondrial

dysfunction was i. as a cellular and metabolic process and to ii. adipose tissue and performed mRNA-

seq to assess the heredity of other obesity-associated processes such as chronic, low-grade inflammation

(meta-inflammation). Focussing on expression of gene markers of classically (M1) and alternatively

activated (M2) monocytes/macrophages, cell types that are linked to meta-inflammation-driven adipose

tissue dysfunction[33], we found that, as expected, mRNAs encoding M1 markers such as *Interleukin 1b*

(IL1b), *Monocyte chemoattractant protein-1 (Mcp1/Ccl2)* and *Tumor Necrosis Factor Alpha (TNF)*;

but also M2 markers like *Arginase 1 (Arg1)*, *Mannose Receptor C-Type 1 (Mrc1)*, *C-Type Lectin*

Domain Containing 10A (Clec10a) and *Resistin Like Beta (Retnla)* were induced in F_0 HF, but not F_1

eWAT (**Fig.S6e,f**), emphasising that a meta-inflammatory tone is not passed on from F_0 HF mice.

Given the dysfunction in a specialised cellular process like mitochondrial respiration in eWAT, we next

aimed to juxtapose adipose tissue with transcriptional effects in liver, an organ essential for glucose and

lipid homeostasis: Analysis of mRNAseq data using multidimensional scaling (MDS) plots showed that

F_0 LF segregated from F_0 HF and F_0 HF-LF, arguing for mostly irreversible effects of obesity in F_0 liver

(**Fig.S7a**) and modest effects of F_0 obesity on F_1 liver transcriptomes (**Fig.S7b**). When comparing F_0

HF to F_0 LF livers, we found 2,474 up- and 2,612 down-regulated genes with $pV_{adj} \leq 0.05$ (**Fig.S7c**),

whereas comparing F_1 (F_0 HF) to F_1 (F_0 LF) without adjusting for multiple testing ($pV \leq 0.05$) yielded

716up- and 546 down-regulated genes (**Fig.S7d**). Gene Set Enrichment analysis (GSEA)[63] of

differentially regulated mRNAs identified lipid metabolism and organelle formation as induced,

whereas repressed terms included lipoprotein trafficking, endoplasmic reticulum and ribosome, yet no

transcriptional evidence for altered mitochondrial processes in liver of obese mice and their F_1 progeny

(**Fig.S7e,f**), emphasising the specificity of mitochondrial alterations in obese fat.

The non-hereditary effects of paternal obesity were supported by histological evidence, and pathologist

assessment of H/E (morphology), Sirius Red (collagen/fibrosis) and anti-CD45 (leukocytes) staining

revealed important hallmarks of metabolic dysfunction-associated steatosis liver disease (MASLD) like

micro-/macro-vesicular steatosis, fibrosis and locular inflammation in F_0 HF mice, but not lean F_0 or F_1

descendants (**Fig.S8a-h**). Targeted analysis of expression changes in genes linked to hepatic lipid

metabolism and meta-inflammation such as *Adhesion G Protein-Coupled Receptor E1 (Adgre, F4/80),*

Apolipoprotein A4 (Apoa4), Apolipoprotein E (Apoe), Low Density Lipoprotein Receptor (Ldlr),

Peroxisome Proliferator Activated Receptor Gamma (Pparg), Cluster of Differentiation 36 (Cd36) and

Sterol Regulatory Element Binding Transcription Factor 1 (Srebf/Srebp1) confirmed that induction was

confined to F_0 HF livers (**Fig.S8i**). Furthermore, mitochondria-associated genes that were activated in

obesity, presumably counteracting lipid accumulation in MASLD, for instance *Carnithine*

Palmitoyltransferase 1A (Cpt1a), Carnithine O-Acetyltransferase (Crat), Acyl-Coa-Oxidase (Acox1)

and *Acetyl-CoA-Acetyltransferase (Acat1)* were induced in livers from F_0 HF, but not F_1, mice. *Thus,*

combinatorial analyses of transcriptional and histological changes argue against prominent effects of

paternal obesity onto F_1 gene expression and mitochondrial processes in hepatic cell types.

Paternal obesity causes qualitatively similar alterations in eWAT and sperm of obese male mice

Spermic sncRNA were reported to be subjected to dietary cues such as high-fat and -sugar diets in mice

and humans[22]; and it is therefore conceivable that HF feeding and weight loss alter spermic sncRNA

levels that can then, following oocyte fertilisation, be passed on to fertilised oocytes (zygotes) as shown

recently [23]. By changing sRNA composition in zygotes, sncRNAs thus have the unique ability to post-

transcriptionally affect decision points in embryogenesis, thereby contributing to the observed glucose

intolerance and mitochondrial protein repression observed in F_1 offsprings sired by F_0 HF bucks. A

particular aspect of sperm transcriptomes is the complex profile of sncRNAs involving miRNAs,

ribosomal RNAs (rRNAs), PIWI-interacting RNAs (piRNAs), (mitochondrial) transfer RNAs ((mt)-

tRNAs) and (mt)-tRNA fragments [26] that, as described by Tomar et al.[23], are affected by mitochondrial

dysfunction in obesity and, particularly relevant in this setting, are physically exchanged between sperm

and oocytes during fertilisation, a scenario also conceivable for other types of mitochondria-associated

sncRNAs, for example mito-miRs. To the map the effects of paternal obesity and weight loss on spermic

sncRNAs and to gain insights into sRNA-driven metabolic adaptations in F_0 HF sperm, we next isolated

motile spermatozoa using swim-up assays and charted differential spermic sncRNAs responses as

consequence of body weight gain and losses and found sperm harboured complex sncRNA pools of

thousands of (mt)-tRNAs, piRNAs, rRNAs and miRNAs (**Fig.3a-c**). In addition to effects elicited by

obesity and/or LF feeding at the single sncRNA level, we observed that (mt)-tRNAs and miRNAs as

entire RNA biotype were repressed by HF feeding. As a prior study from Tomar et. al[23] had shown

effects on mt-RNA in sperm after HF feeding, this was investigated further: Mt-tsRNA fragments were

found for nine mt-tRNAs (*mt-Tl1*, *mt-Tq*, *mt-Tm*, *mt-Ts1*, *mt-Tr*, *mt-Th*, *mt-Ts2*, *mt-Te*, *mt-Tp*), where

mean cpm above 10 was observed in *mt-Tm*, *mt-Ts1*, *mt-Ts2* and *mt-Th* (**Fig.9a**), but only 5' fragments

of *mt-Ts-1* were significantly changed in sperm from F_0 HF and F_0 HF-LF diet (**Fig.9b**). As miRNAs

are abundant in somatic cell types and sperm, harbour the possibility of horizontal transfer between

them, and exert gene-regulatory functions within both cellular compartments, we turned our attention

to miRNA regulation in sperm. We found that F_0 obesity and weight loss caused a coordinated and

inverse (i.e., downregulation and upregulation, respectively) effects on miRNome regulation (**Fig.3d,e**),

a finding previously observed by our labs [51] and others[49,50,64] for obese adipose tissues; in the latter case

at least partially due reduced abundance of DICER1. Intrigued by the repression of miRNAs in two

functionally distinct, yet anatomically adjacent tissues and cell types like eWAT and sperm, we asked

if differential miRNA responses were qualitative similar in both organ which, if present, would support

the possibility of physical miRNA exchange between eWAT adipocytes and sperm as shown for other

testicular cell types and sperm[21]: Indeed, upon comparison of lean, obese and weight-regressed sperm

and eWAT miRNAs, we found highly congruent responses across both compartments with activating

mito-miRNAs like *miR-378* and *miR-143* being down-regulated, whilst inactivating mito-miRs of the

let-7 family up-regulated in obese F_0 eWAT and sperm (**Fig.3f**). Importantly, when broadly mapping

the regulation of *let-7* family isoforms, we found mild, but synchronous inductions of *let-7* in F_0 eWAT

and sperm and F_1 eWAT (**Fig.3g-3i**). As increases of *let-7c* were reported in intergenerational rat studies

by us [8] and others[65] and because *let-7* impairs embryo implantation and glucose homeostasis when

overexpressed in mice [66,67], we focussed on *let-7* isoforms for functional follow-up studies in germ cells

and adipocytes. *Thus, paternal obesity triggers similar alterations in adipose tissue and sperm miRNAs*

and let-7 isoforms are consistently induced across cell types and generations.

Zygotic delivery of let-7 causes glucose intolerance and mitochondrial impairment in sired mice

Prompted by profound and reversible effects of male obesity and weight loss on sperm miRNomes and

let-7 in particular, as environmentally and artificially induced (e.g., elicited by miRNA microinjection)

alteration in zygote miRNA (pools) trigger abnormalities in progeny[25,68], and because embryonic

trajectories require maturation of pre- into mature miRNAs[69], we tested the functional relevance of *let-*

7 gain-of-function in one-cell embryos for F_1 phenotypes. We focussed on *let-7d-5p* and *let-7e-5p* given

that these exhibit the most concordant increases amongst *let-7* family members in F_0 and F_1 eWAT and

sperm (**Fig.3g-i**) and because *let-7* has before been connected to embryonic cell fate decisions[70,71]. To

test if zygotic delivery of *let-7* corresponding at levels seen in few obese spermatozoa triggers

phenotypical and metabolic changes in mice sired from *let-7* microinjected embryos (F_{MI} generation)

and whether these would resemble mice sired by F_0 HF fathers, we injected physiological amounts (50

fmol, i.e., *let-7* found in 5-10 spermatozoa) of *cel-mir-67* mimic as non-mammalian (*C.elegans*) control,

and mimics for *let-7d-5p* and *let-7e-5p* by microinjection into zygotes conceived from lean males and

females C57BL/6N mice using standard protocols[29] (**Fig.4a**). We injected 126/138/106 zygotes with

cel-mir-67, *let-7d-5p* and *let-7e-5p*, but observed no discernible difference in implantation success, litter

sizes and number of pregnancies in F_{MI} mice (**Fig.S10**), arguing against roles of *let-7* for zygote viability

and implantation. Next, we analysed at least 19 pups per group from 3-4 pregnancies but observed no

overt differences in postnatal body weight trajectories (**Fig.4b**). Despite similar weight, we detected

impaired glucose tolerance (**Fig.4c,d**), insulin resistance (**Fig.4e**) and elevated random-fed glycemia

(**Fig.4f**) in mice sired by *let-7* injected F_{MI} embryos, whilst eWAT and liver weights were

indistinguishable between F_{MI} groups (**Fig.4g**). Importantly, when performing mRNA-seq in eWAT of

F_{MI} mice, despite mild changes, we found that *let-7* down-regulated mitochondrial processes in F_{MI} mice

(**Fig.4h**). *Thus, zygotic delivery of let-7 at levels corresponding to few F_0 HF sperm cells, causes*

glucose intolerance and aberrant mitochondrial gene regulation in mice derived from these embryos.

Obesity-associated let-7 rewires mitochondrial metabolism during embryogenesis

As transfer of miRNA perturbation in sperm to oocytes can have deleterious consequences and give

rise to incompletely developed embryos [20,72,73], and as zygotes and blastocysts have the capacity to

differentiate into virtually all cell types, (early) embryogenesis represents an important window of

developmental plasticity that might affect F_{MI} phenotypes if perturbed by miRNA delivery, despite the

physiological levels of miRNA delivery and absence of embryonic lethality. Nonetheless, it is

conceivable that alterations in zygotic miRNAs still entails far-reaching consequences in sired mice due

to a combination of altered developmental, epigenetic and metabolic processes[73]. Based on the specific

mitochondrial alterations, but absence of increased *let-7* levels in F_{MI} eWAT, we next hypothesized that

let-7d-5p and *let-7e-5p* could carry out important roles in cell fate decisions during embryogenesis

which will indirectly contribute to the glucose intolerance in F_{MI} pups via altered embryonic

development. To test this experimentally, we repeated zygotic *let-7* injections but allowed miRNA-

injected embryos to divide thrice until the 8-cell blastomere stage. We dissociated embryos into single

blastomeres and performed single-cell RNA-Seq (scRNA-seq) using Smart-seq2[74] (**Fig.4i**).

Remarkably, we found that after three cell divisions, miRNA-injected blastomeres were still

transcriptionally distinct, reflecting the profound transcriptional and developmental consequences of

inducing *let-7* in zygotes (**Fig.4j**). When performing GSEA of *let-7*-dependent gene sets, we

corroborated that *let-7d-5p* repressed terms linked to mitochondrial processes like ATP synthesis, TCA

cycle and ETC (**Fig.4k**). Intergenerational programming in isogenic rodents has been described before,

yet it is elusive if these effects are translatable to species with highly different reproductive apparatuses,

genetic backgrounds, and timescales of obesity development like humans. Although epidemiological

evidence[4,75] suggests heritable effects of overfeeding and malnutrition in humans (e.g., the famous

'Dutch Hunger Winter'[76]), causal evidence for epigenetic inheritance is hard to separate from cultural

and ecological inheritance in humans[2]. As it is becoming appreciated that nutritional and metabolic cues

shape human sncRNA expression[22], we tested effects of weight change on semen *let-7* in a small cohort

of n=15 obese men (BMI, 39.49 ± 1.80) undergoing voluntary weight loss[77] and quantified *hsa-let-7*

using TaqMan qPCR. Intriguingly, we found that *hsa-let-7d-5p* and *hsa-let-73-5p* followed the trends

observed for weight regulation , suggesting similar correlations between metabolic improvement and

let-7 reductions in human semen (**Fig.4l,m**). *Thus, obesity-associated let-7d affects murine*

embryogenesis by impairing mitochondrial gene regulation and similar processes might be occurring

following weight loss in obese humans.

Adipocyte let-7 impairs mitochondrial function via post-transcriptional silencing of DICER1

Beyond F_0 HF sperm (**Fig.3g-i**), we found *let-7d-5p* and *let-7e-5p* upregulated in F_0/F_1 eWAT, and

AGO2-CLIP interactome analyses confirmed that *let-7* gene targets are enriched for genes encoding

mitochondrial proteins, that are downregulated by obesity in F_0/F_1 eWAT (**Fig.5a**). For this, and to add

to their previously ascribed role in fertilized oocytes, we aimed to dissect the transcriptional and

functional consequences of elevated *let-7* in adipocytes, with a specific focus on mitochondrial gene

regulation as highly sequence-related *let-7i* can impair commitment of mitochondria-rich adipocytes [78].

Yet, if roles for *let-7* in mitochondrial formation and function exist in mature adipocytes is unknown.

To test this, we delivered by reverse lipofection mimetics for *let-7d-5p* and *let-7e-5p* and *cel-mir-67*

into stromal vascular fraction-derived mature epididymal white adipocytes (1°eWA) to mimic obesity-

associated increased in *let-7* in eWAT. When interrogating the effects of *let-7* overexpression, we

observed that *let-7d-5p* elicited more pronounced effects (1,768 up-regulated, 1,387 down-regulated

genes, **Fig.5b**) than *let-7e-5p* (557 up-regulated, 893 down-regulated genes, **Fig.5c**) compared to *cel-*

mir-67, yet both *let-7* isoforms downregulated processes related to lipid metabolism, Insulin-AKT

(Protein Kinase B) signalling and carbohydrate metabolism (**Fig.S11**). Although not overrepresented at

the GOBP level, we found rate-limiting genes in glycolysis, TCA, and mitochondrial lipid import like

Carnithine Palmitoyltransferase 2 (*Cpt2*), Carnithine O-Acetyltransferase (*Crat*)*,* Hexokinase 2 (*Hk2*),

Isocitrate Dehydrogenase 2 (*Idh2*)*,* Pyruvate Carboxylase (*Pcx*), Pyruvate Dehydrogenase Kinase 2

(*Pdk2*) and mitochondrial ATP-magnesium/phosphate antiporter Solute Carrier Family 25 Member 24

(*Slc25a24*, **Fig.5b,c**; in blue) repressed upon *let-7* delivery. Importantly, we also detected pivotal genes

in miRNA maturation such as Ribonuclease III DICER1 *(Dicer1)* and Argonaute RISC Catalytic

Complex 2 (*Ago2,* **Fig.5b,c**; in red*)*. Next, we experimentally validated these predicted *let-7* targets by

dual luciferase reporter assays (DLRA)-based miRNA-mRNA 3'UTR interaction studies in HEK293

cells for genes linked to mitochondrial lipid and ATP uptake (*Slc25a24, Cpt2*), glycolysis (*Hk2*, *Pdk2*),

TCA activity (*Sdha*, **Fig.5d**) and miRNA processing (*Dicer1, Ago2*).

Dicer1 3'UTRs and exons harboured several independent *let-7* seeds of which, after mutation of

putative *let-7* responsive elements (MREs), two from exonic regions became refractory to *let-7*

repression, demonstrating direct negative effects of *let-7* on DICER1 translation (**Fig.5e**). We strived to validate *let-7* targets identified in HEK293 cells adipocytes, and delivered *let-7* and control mimics to 1°eWA and confirmed that *let-7* also repressed DICER1 in white adipocytes, yet caused only modest reductions in mitochondria-/TCA-/glycolysis-associated proteins upon overexpression (**Fig.5f-h**), prompting us to hypothesize that the chronic *let-7* induction seen in obese adipocytes might repress mitochondrial function rather via sustained DICER1 loss than by direct *let-7* effects on mitochondrial function; and that *Dicer1* inactivation might be a better proxy for mimicking the prolonged upregulation in *let-7d-5p* and *let-7e-5p* in obesity than acute overexpression of the miRNA.

To address if *Dicer1* silencing because of sustained *let-7* upregulation is sufficient to phenocopy *in vitro* the obesity-evoked impediments in mitochondrial function, we designed two independent *Dicer1*-targeting locked nucleic acid (LNA) inhibitors that repressed *Dicer1* at the mRNA and protein level, yet with varying efficiencies (**Fig.5i**) and interrogated mitochondrial respiration using Seahorse. We found that *Dicer1* knockdown reduced oxygen consumption rates (OCR) as proxy for mitochondrial respiration (**Fig.5j**) and extracellular acidification rates (ECAR) as proxy for glycolytic rates in 1°eWA (**Fig.5k**) supporting our hypothesis that obesity impairs mitochondrial respiration and gene regulation, at least partially, by post-transcriptional silencing of DICER1 via *let-7* (**Fig.5l**). *Thus, obesity-associated let-7 impairs mitochondrial oxidative respiration by silencing of DICER1 in mature adipocytes.*

Discussion:

High transcriptional plasticity of mitochondrial dysfunction in adipose tissue

In our study we found that hepatic expression changes inflicted by HFD feeding where to a large extend irreversible, whereas adipose tissue miRNA/mRNA transcriptomes and translatomes, and particularly mitochondrial OxPhos and TCA gene sets, reverted to a state *ante*. Similar studies have investigated the effects of reversible (i.e., diet- and exercise-evoked weight gain and loss) on hepatic and adipose transcriptomes: Gonzalez-Francesa et al. observed[42] that sustained diet-induced weight loss correlated with appreciable adipose transcriptional plasticity and mitochondrial gene expression, where short-term, exercise induced did not. Thus, durations of weight loss, and differences in control diet exposure might reflect the importance of nutritional and exposure factors during weight loss interventions on tissue responses in obesity. Despite not performing profiling of chromatic accessibility, transcriptional factor (TF) binding site enrichment analysis revealed that transcriptional activities of mitochondrial biogenesis factors like Nuclear Respiratory Factor 1 (*Nrf1*) and expression of transcriptional (co)-regulators such as Peroxisome proliferator-activated receptor gamma coactivator 1-alpha (*Ppargc1a*) and *Nrf1*[79] were repressed in obese fat (not shown). As Brandao et al. reported [80] that ablation of DICER1 in adipocytes repressed *Ppargc1a*/*Nrf1*, we can conceive a scenario where reduced DICER1 (for instance due to increased *let-7d/e-5p*) drives and/or synergizes with reduced *Ppargc1a*/*Nrf1* levels to impair mitochondrial biogenesis in hypercaloric adipocytes.

Rewiring of embryonal development by gametic miRNAs

In *C. elegans* and *D. melanogaster*, germline gene silencing by small Noncoding RNAs (sncRNA) is well-understood and similar processes are suggested for mammals [19]. miRNAs are not only critical for sperm cell formation [27] and embryogenesis [28], but can also be transferred from sperm to zygotes and it is plausible that oocyte-sperm fusion alters zygotic miRNA pools, affect embryonic development, and instructs phenotypic traits and somatic gene expression in offspring as observed in our intergenerational paradigm. In contrast to DNA-intrinsic epigenetic modifications like DNA methylation (DNAme) and histone post-translational modifications, miRNAs diffuse between and across cellular compartments, are highly abundant in most cells, and, crucially, can be exchanged between cells through extravesicular

transport [32]. As mature sperm is transcriptionally inert, and no mechanism exists how sperm senses

endocrine information and translates this into differential miRNA levels, sperm relies on 'decoding

somatic cell types' that possess these sentinel features and are capable of instructing sperm by transfer

of miRNAs and other sncRNAs like mitochondrial tRNA fragments that are exchanged between sperm

and oocyte at fertilisation [23]. Furthermore, epididymal cells, somatic epithelial cells of the extragonadal

male reproductive tracts, secrete *epididysomes*, microvesicles that fuse with nascent sperm [81], and as

we observed several miRNAs concordantly regulated in F_0 adipose tissue and sperm, horizontal miRNA

transfer from adipocytes to sperm, potentially via epididysomes is a plausible scenario that still lacks

experimental proof in our paradigm. Finally, altering miRNA levels like *miR-199a-5p* in zygotes by

microinjection is sufficient to reprogram embryonal metabolism [82], highlighting that miRNA alterations

can rewire embryonal development and metabolism which are still compatible with fetal development.

Methods:

Mice

All animal experiments were approved by the Ministry of Environment and Agriculture Denmark (Miljø- og Fødevarestyrelsen), the license no. 2018-15-0201-01544. All mice were housed under a 12-h light/ 12-h dark cycle in a temperature and humidity-controlled facility and had *ad libitum* access to diets and drinking water. 4-week-old C57BL/6J male mice (Janvier Labs) were fed with a chow diet (NIH-31, Zeigler Brothers Inc., 8% calories from fat) for 2 weeks of acclimatization and then were fed with either a low-fat diet (LF, D12450J, 10% kcal from fat, Research Diet) or a high-fat diet (HF, D12492, 60% kcal from fat, Research Diet) for 9 weeks. After 9 weeks of feeding, half of HF-fed mice were randomly assigned to the diet regression group which was given LFD. LFD is carefully chosen to match food source (animal vs plant) and micronutrient composition of HFD.

Mouse husbandry

All male virgin mice were maintained on the assigned diet until 21 weeks of age and mated with 10-week-old C57BL/6J female virgin mice (Janvier Labs). 8-week-old C57BL/6J female mice were fed with a chow diet for 2 weeks of acclimatization and then were fed with LF during breeding and lactation. Mating was set up for 5 days in a female's cage and vaginal plugs were checked every morning. All mice, including the F_0 HF group, were maintained on LF during breeding to remove confounding effects from maternal diets. Males were sent from breeding cages to the original cage to be fed with the assigned diet for additional 2 weeks and then sacrificed. The offspring from all groups of founders consumed only LF. Representative male offspring from six litters per group were included in all analyses.

Intraperitoneal glucose and insulin tolerance tests:

Mice were fasted for 6 h for glucose-tolerance tests. Fasting glucose levels were measured and then mice were intraperitoneally injected with 2 g/kg body weight of 20 % glucose solution (B. Braun Medical A/S). For intraperitoneal insulin-tolerance tests, random-fed glucose levels were measured, and mice were intraperitoneally injected with 0.75 IU/kg body weight of 100IU stock insulin (Novo

Nordisk). Blood glucose levels were measured at 15, 30, 60, 90, and 120 min after injecting glucose or

insulin using a CONTOUR®NEXT glucometer (Bayer). For ipGTT/ipITT, we omitted mice with

strongly delayed rises in glucose and no discernible drop in blood glucose, respectively, from analyses

as this was the consequence of mal injection outside the peritoneum.

Tissue collection:

Mice were sacrificed by carbon dioxide euthanasia. Livers, eWAT, iWAT and BAT were weighed,

snap-frozen in liquid nitrogen, and stored in -80 °C. Pieces from liver and eWAT were fixed in 4%

paraformaldehyde (PFA) for histology analysis. For serum collection, blood collected by the cardiac

puncture was centrifuged at 2,000 g for 15 min. The supernatant was collected and stored in -80 °C.

Sperm collection:

Mature sperm cells were collected from the cauda epididymis. Briefly, sperm cells are allowed to swim

out from the cauda epididymis and incubate in PBS at 37 °C for 15 min. Supernatant containing sperms

was passed through a 40-µm cell strainer and then incubated in a somatic cell lysis buffer (0.5 % Triton-

X and 0.1 % SDS) for 40 min on ice. Sperms were centrifuged at 600 g for 5 min and the pellets were

washed with PBS and pelted again at 600 g for 5 min. The supernatant was discarded, and the pellets

were snap-frozen in liquid nitrogen and stored at -80 °C.

Zygotic miRNA microinjection and embryo implantation:

Ovulation of prepubescent (4-week-old) C57BL/6J females was induced by intraperitoneal (IP)

injection of PMSG, 5 IU/female, followed by IP injection of hCG, 5IU/female, 47h later. After the

second injection, females were set in breeding with C57BL/6J stud males. Next morning, they were

monitored for formation of a vaginal plug (indicating mating has occurred during the night). Females

were euthanized and oviducts were dissected to harvest the cumuli containing zygotes and unfertilized

oocytes. Cumuli were disaggregated by incubation in a 10mg/ml solution of Hyaluronidase (Sigma-

Aldrich ref. H4272) in M2 medium (Sigma-Aldrich ref. M7167) at RT for 10 minutes. Zygotes were

sorted out, assessed by the presence of the second polar body and cultured in KSOM medium (Merck

Millipore ref. MR-106-D). Oocytes were discarded. Each miRNA was diluted at 10 ng/ul in

microinjection buffer (5 mM Tris, 0.1 mM EDTA, pH 7.4) as stock solution and kept at -80 °C. The

day of microinjection, 1,5 ng/ul solutions were prepared from the stock, and injected in the pronuclei

of the zygotes following standard microinjection procedures. miRIDIAN mimics were ordered from

Dharmacon: Inert miRNA control is based on *C.elegans cel-mir-67* which is not expressed in mouse or

human cells (5'-UCACAACCUCCUAGAAAGAGUAGA-3', cat. no. CN-001000-01-05), *mmu-let-*

7d-5p (cat. no. C-310486-07-0002) and *mmu-let-7e-5p* (cat. no. C-310507-07-0005). Injected zygotes

were cultured in KSOM until the microinjection session was completed and then cryopreserved by the

"Slow Freezing" Method (Manipulating the Mouse Embryo, a Laboratory Manual, fourth edition,

Chapter 16, Richard Behringer et al. Cold Spring Harbor Laboratory Press, 2014) and transported to

SDU at -150 °C for embryo transfer and recovery.

The frozen microinjected embryos were thawed at SDU and cultured in KSOM (Merck, MR-121-D)

for 40 minutes (Manipulating the Mouse Embryo, a Laboratory Manual, fourth edition, Chapter 16,

Richard Behringer et al. Cold Spring Harbor Laboratory Press, 2014). Viable embryos with intact zona

pellucida were selected for further transfer. These embryos were then washed in 10 drops of M2 and

surgically transferred to the oviducts of 0.5 dpc pseudo pregnant females. Pseudopregnancy was

induced in the females by mating them with a vasectomized male. Prior to implantation, the pseudo

pregnant females were anesthetized with ketamine (100 mg/kg) and xylazine (10 mg/kg) through

intraperitoneal injection, and the surgical embryo transfer procedure was performed following the

protocol described by Behringer et al. (Manipulating the Mouse Embryo, a Laboratory Manual, fourth

edition, Chapter 6, Richard Behringer et al. Cold Spring Harbor Laboratory Press, 2014). Postoperative

analgesic treatment with carprofen (5 mg/kg) was administered subcutaneously for 48 hours.

8-cell-morula blastomere disaggregation:

miRNAs were injected in the embryo at zygote stage as described above. After injection, embryos were

incubated in KSOM (KSOM-Embryomax advanced embryo medium, Merck Millipore #MD-101-D)

for 48-56 hours at 37 °C, in 5% CO_2. The culture medium was refreshed daily. In the afternoon of the

third day of culture, when the embryos had reached the 8-cell morula stage, they were incubated at RT

in Tyrod's solution (Sigma-Aldrich T1788) for up to 30 seconds until the zona pellucida was completely

dissolved. Afterwards the naked morulae were washed through three drops of M2 medium (Sigma-

Aldrich M7167) and finally incubated in KSOM without Ca^{2+} and Mg^{2+} at RT for 20 minutes. Then,

embryos were moved to a new drop of fresh M2 medium and there, pipetted up and down repeatedly to

facilitate the blastomere disaggregation. When single blastomeres were obtained, they were individually

transferred to single wells of a 96-well single-cell-sequencing plate from VWR (Cat. No. 82006-636)

with 2 ul of RNAse inhibitor mix (containing 0.12 ul Triton X-100 10 %, 0.1 ul RNAse inhibitor (Takara

(Cat. No. 2313B, 40 U/ul), 1.78 ul MilliQ water (RNAse-free)). Plates were them frozen at -80 °C until

the sequencing was performed.

Lipidomics and Metabolomics:

Serum Extraction:

100 µL of serum was mixed with 1000 µL of a 2:1 chloroform/methanol solution and 200 µL of water.

Four blank samples, each containing 100 µL of water, were processed in parallel. The mixtures were

shaken at 1000 rpm for 30 minutes at 4°C, followed by centrifugation at 16,000g for 10 minutes at 4

°C to separate the phases. The aqueous phase was re-extracted with 350 µL of an 86:14:1

chloroform/methanol/water solution. The organic phase from this re-extraction was combined with the

corresponding organic phase from the initial extraction. Aqueous fraction was dried using SpeedVac,

while the organic phase was dried under a stream of nitrogen. For subsequent analysis, the dried aqueous

fractions were reconstituted in 30 µL of 1 % formic acid for metabolomics. The organic phase was

reconstituted in 30 µL of lipidomic solvent A (5:1:4 isopropanol/methanol/water with 5 mM ammonium

acetate and 0.1 % acetic acid). A pooled quality control sample was prepared for each analytical setup

by combining 5 µL from each sample, excluding blanks.

Lipidomic analysis:

For **lipidomic** analysis, 2.5 µL of each sample was injected into an Agilent 1290 Infinity UPLC system

(Agilent Technologies, Santa Clara, CA, USA) equipped with a Zorbax Eclipse Plus C18 guard column

(2.1 × 5 mm, 1.8 µm) and an analytical column (2.1 × 150 mm, 1.8 µm), maintained at 50 °C. Analytes

were eluted at a flow rate of 400 µL/min using eluent A (water with 5 mM ammonium acetate and 0.1

% acetic acid) and eluent B (99:1 isopropanol/water with 5 mM ammonium acetate and 0.1 % acetic

acid). The gradient was as follows: 0 % B from 0 to 1 min, 0-25 % B from 1 to 1.5 min, 25-95 % B

from 1.5 to 12 min, 95 % B from 12 to 14 min, and 95-0 % B from 14 to 15 min, followed by a 3-

minute equilibration. For **metabolomics** analysis, 1.5 µL of each sample was injected into the same

UPLC-MS system, but with a ZORBAX Eclipse Plus C18 guard column (2.1 × 50 mm, 1.8 µm) and an

analytical column (2.1 × 150 mm, 1.8 µm) maintained at 40°C. Eluent A was 0.1 % formic acid, and

eluent B was 0.1 % formic acid in acetonitrile. The gradient was as follows: 3 % B from 0 to 1.5 min,

3-40 % B from 1.5 to 4.5 min, 40-95 % B from 4.5 to 7.5 min, 9 5% B from 7.5 to 10.1 min, and 95-3

% B from 10.1 to 10.5 min, followed by a 3.5-minute equilibration. Both analyses were performed on

a 6530B quadrupole time-of-flight mass spectrometer (Agilent Technologies, Santa Clara, CA, USA)

for mass spectrometric detection in positive and negative ion modes. MS1 mode settings included a

scanning range of 50-1700 m/z for lipidomic and 10-1050 m/z for metabolomics, with 2 scans/sec.

Internal calibration was performed using Hexakis (1H,1H,3H-tetrafluoropropoxy) phosphazene

delivered through a second needle in the ion source via an isocratic pump running at 20 µL/min. Iterative

data-dependent MS2 mode was used for fragmentation in both analyses with the following settings:

collision energy at 40 V (lipidomics) and 20-40 V (metabolomics), precursor threshold at 5000 counts,

and active exclusion after 2 spectra, with re-inclusion after 0.5 min.

Data Processing and Normalization:

All data was converted. mzML format using ProteoWizard[83]. Lipidomics data was annotated in MS-

Dial (v. 4.9) against the incorporated lipid-blast[84] with an MS1/MS2 mass tolerance of 0.005/0.01 Da

and a minimum identification score of 70 %. The annotated compounds were exported to PCDL

manager B.08.00 (Agilent Technologies) to create a database for area extraction in Profinder 10.0

(Agilent Technologies). Metabolomics data was processed in MzMine (v2.59) utilizing modules such

as ADAP chromatogram builder and deconvolution. Compounds were annotated at Metabolomics

Standards Initiative (MSI) levels 3 using the libraries of National Institute of Standards and Technology

2017 (NIST17) and MassBank of North America (MoNa). Features with signals less than 5 times those

in blanks or missing in more than 20 % of QC samples were removed and signals were QC corrected

using statTarget[85].

Data analysis:

For **metabolomics** pathway enrichment analysis (KEGG pathways of Mus Musculus) comparing F_0 LF

and F_0 HF serum (parameters 'Scatter plot', 'Hypergeometric test', 'relative-betweenness Centrality',

'Use all compounds in the selected pathway library') of differentially abundant polar metabolites (pV

≤ 0.1) were loaded into Metaboanalyst 6.0. For **lipidomics** and **metabolomics** principal component

analysis and heatmap representation of indicated no. of differentially abundant lipids and metabolites

were generated using ‚Statistical Feature' (one-factor option), Sample normalization = ‚Normalization

by sum', Data transformation = ‚Log transformation (base 10)' and Data scaling = ‚Mean Centering'.

Serum insulin and leptin ELISA:

Serum insulin and leptin levels were measured by mouse insulin ELISA kit (Mercodia, Uppsala,

Sweden) and mouse leptin ELISA kit (Crystal Chem, IL, USA) according to manufactures' protocol.

Histology and staining:

Liver and adipose tissues fixed with 4 % of paraformaldehyde were used for histology analyses.

Formalin-fixed, paraffin-embedded tissues were cut into 3 μm sections. Sections were stained by

haematoxylin and eosin (H&E) and Sirius Red by standard procedures at the Department of Pathology

at Odense University Hospital, Odense, Denmark. Immunohistochemistry for CD45 was performed on

Ventana BenchMark Ultra (Roche) with ready-to-use CD45 (LCA) (2B11 & PD7/26) Mouse

Monoclonal Antibody.

A liver pathologist (SD) assigned steatosis, lobular inflammation, ballooning, fibrosis stage, and portal

inflammation. Steatosis was graded as S0 (< 5% hepatocytes with large fat vacuoles), S1 (5-33%), S2

(>33-66%), and S3 (>66% hepatocytes with large fat vacuoles)[86]. Lobular inflammation (0-3) and

hepatocellular ballooning (0-2) were assessed according to the NAS-CRN Activity Score by Kleiner et

al.[86]. Also, portal inflammation was assessed, using a 4-tiered score by Kleine et al. Liver fibrosis stage

was assessed according to NAS-CRN for fibrosis; F-0, no fibrosis; F-1, perisinusoidal or periportal

fibrosis; F-2, perisinusoidal and portal/periportal fibrosis; F-3, bridging fibrosis; F-4, cirrhosis.

Measurement of adipocyte volumes:

The volume weighted mean volume of adipocytes was determined from the H&E sections with

newCAST stereology software (Visiopharm), autodisector; measurements on systematic, randomly

selected fields were performed on by at least 75 adipocyte profiles in each tissue.

Bulk proteomics from epididymal white adipose tissue:

Deep frozen fat pads (200 mg of epididymal adipose tissue) were powdered with mortar-pestle and

subjected to homogenization with ceramic beads (Percellys, 5500 rpm, 2 x 10 seconds, 30 seconds

break) in the buffer containing 50 mM HEPES pH 7.4, 1 % Triton X-100, 100 mM NaF, 10 mM sodium

orthovanadate, 10 mM EDTA, 0.2 % SDS, 100 mM NaCl. Tissue homogenates were further processed

to remove the nucleic acid traces (Diagenode SA; 10 min, cycle 30/30 seconds, at 4 °C). Floating fat

and tissue debris were removed from protein extracts by centrifugation at 20000 x g. 150 mg of proteins

were precipitated with 4x volume of ice-cold acetone and washed with ice-cold acetone twice. Air-dry

pellets were resuspended in 8M Urea/50 mM triethylammonium bicarbonate (TAEB) buffer

supplemented with a protease inhibitor cocktail (Sigma). 50 mg of the samples were incubated with 5

mM dithiothreitol at 25°C for 60 min, followed by treatment with 40 mM chloroacetamide for 30 min

(room temperature, light-protected). Lys-C protease (enzyme: substrate ratio of 1:75) was added for 4

hours at 25 °C. The TAEB-diluted sample (2M Urea final) was supplemented with trypsin (1:75 ratio)

followed by overnight digestion at 25 °C. Acidified peptides (1 % formic acid final) were purified on

the SDR-RP (C18) multi-stop-and-go-tip (Stage Tip)[87].

LC MS/MS Label-Free Quantitative Proteomic Analysis from epididymal white adipose tissue:

All samples were analyzed by the Proteomics Facility on a Q Exactive Plus Orbitrap mass spectrometer

coupled to an EASY nLC 1000 (Thermo Scientific). Peptides were loaded with solvent A (0.1 % formic

acid in water) onto an in-house packed analytical column (50 cm - 75 μm I.D., filled with 2.7 μm

Poroshell EC120 C18, Agilent) and separated with 150 min gradients. Peptides were

chromatographically separated at a constant flow rate of 250 nL/min using the following gradient: 4-6

% solvent B (0.1 % formic acid in 80 % acetonitrile) within 5.0 min, 6-23 % solvent B within 120.0

min, 23-54 % solvent B within 7.0 min, 54-85 % solvent B within 6.0 min, followed by washing and

column equilibration. The mass spectrometer was operated in data-dependent acquisition mode. The

MS1 survey scan was acquired from 300-1750 m/z at a resolution of 70,000. The top 10 most abundant

peptides were isolated within a 2.1 Th window and subjected to HCD fragmentation at a normalized

collision energy of 27 %. The AGC target was set to 5E5 charges, allowing a maximum injection time

of 60 ms. Product ions were detected in the Orbitrap at a resolution of 17,500. Precursors were

dynamically excluded for 20.0 s. All mass spectrometric raw data were processed with MaxQuant[88]

(version 1.5.3.8, using default parameters and searched against the canonical murine Uniprot reference

proteome (UP589, downloaded on 28/08/2020). Label-free quantification option was enabled as was

the match-between runs option between replicates. Follow-up analysis was done in Perseus[89] 1.6.15.

Hits from the decoy database, the contaminant list, and those only identified by modified peptides were

removed. Afterwards, results were filtered for data completeness in replicate groups and LFQ values

imputed using sigma downshift with standard settings. Finally, FDR-controlled T-tests between sample

groups were performed with s0 = 0.2. Proteins were annotated on GOBP, GOCC, KEGG and Pfam

terms. Afterwards, all one-way ANOVA (s= 0.2) significant candidates were Zscored and hierarchically

clustered using the Euclidean distance model. Resulting clusters were merged onto the whole data set

and a Fisher Exact Test calculated on cluster level with the whole dataset as background.

RNA isolation from mouse tissue and primary adipocytes:

Total RNAs from tissues, sperm, and primary adipocytes were isolated using the TRI reagent (Sigma)

followed by clean-up with RPE buffer (Qiagen, Germany). The quality of RNA was validated by the

Agilent RNA 6000 Nano-Kit in Agilent 2100 Bioanalyzer according to the manufacturer's protocol.

(Agilent Technologies, Waldbronn, Germany).

RNA sequencing:

Small RNA sequencing: (1) For sRNA-seq from F_0 sperm, library preparation and sRNA-seq was

performed at Cologne Center for Genomics, Germany. 50 ng of total RNA was used for library

preparation followed by size selection (Small RNA-Seq Library Prep Kit, Lexogen). The library was

sequenced in 1x50-bp single-end reads on NovaSeq6000 serie no. A00316. (2) Small RNA sequencing

from F_0 and F_1 eWAT was carried out after quality control, 125 ng of total RNA was used for library

preparation (Nextflex library kit) followed by sequencing by 1x50-bp single-end reads on Illumina

Novaseq 6000. *mRNA sequencing:* mRNA sequencing was performed in-house. After quality control,

1 μg of total RNA was used for library preparation (NEBNext RNA prep kit) followed by 2x50bp

paired-end sequencing on Illumina Novaseq 6000.

RNAseq data analysis:

Small RNA sequencing of spermatozoa: SncRNA-seq analysis of murine sperm was analyzed with

Seqpac ver. 1.2.0, where sncRNA biotypes were annotated with 1 mismatch against Ensembl protein

and non-coding RNA from GRCm39, miRbase, piRNA from piRBase ver. 3.0, tRNA from tRNAscan-

SE, rRNA from RNAcentral and mitochondria from GRCm39:MT. Biotypes were assessed based on

the following hierarchy: miRNAs, mitochondrial rRNAs, rRNAs, mitochondrial protein-coding

mRNAs, mitochondrial tRNAs, tRNAs, piRNAs, nuclear protein coding mRNAs, lncRNAs, and

miscRNAs. Baseline filtering of 1 read per sample and size of 15 to 75 nt long was performed.

Normalization was performed with counts per million (cpm).

Small RNA sequencing of somatic (adipose) tissue:

Adapter sequences were trimmed from raw miRNA-seq reads using Cutadapt. The processed reads

were mapped to the reference genome and aligned using the Bowtie algorithm within the miRDeep2

pipeline [90]. Normalization was performed using the Trimmed Mean of M-values (TMM) method to

account for sequencing depth and composition bias. Subsequent data analysis, including differential

expression analysis, was conducted in R[91] using respective Bioconductor packages[92].

mRNA sequencing:

Sequencing reads were aligned to the mouse reference genome from Gencode (vM25)[93] using STAR

aligner (v 2.7.9a)[94] and gene features were counted by using featureCountsv (2.0.3)[95]. The genes with

low expression were filtered out using *filterByExpr* function with default settings from edgeR

(v4.0.16)[96] R package. Differential expression analyses were performed using edgeR's *voomLmFit*

[97]unction with sample quality weights (sample.weights = TRUE) and linear model "0 + group", whereas

group represents the combination of all experimental groups of interest. The experimental conditions

and linear models used for each dataset are summarized below. Genes with adjusted p-values below

0.05 were considered as differentially expressed. Gene enrichment analyses were performed by

clusterProfiler (v4.10.1)[98] R package using both over-representation analysis (ORA) and gene-set

enrichment analysis (GSEA) using differentially expressed genes and log-fold-change (logFC) ranked

genes ($n_{permutation}$ = 1,000,000), respectively. Gene sets were retrieved from Gene Ontology (GO.db

v3.18.0) and Reactome (reactome.db, v1.86.2) databases.

Smart-seq2 mRNA sequencing of murine blastomeres:

Custom-made Smart-seq2 protocol was implemented as before[74]. Briefly, after thawing the plate, 1 µL

of dNTP mix (10 mM), 0.1 µL of oligo-dT (100 µM,

/5Biosg/AAGCAGTGGTATCAACGCAGAGTAC(T)30VN-3′) and 0.9 ul nuclease-free water was

added to each well and the lysis protocol was run when the samples were incubated at 72°C for 3 min

in thermocycler. Samples were then put on ice for at least 1 min. Next, 5.60 µL of reverse-transcription

mix containing 0.25 µL of RNase I, 2 µL of 5× first-strand buffer, 2 µL of betaine (5 M), 0.06 µL of

MgCl$_2$ (1 M), 0.5 µL of DTT (100 mM), 0.5 µL of SuperScript III (200 U/µL), and 0,29 ul of nuclease-

free water was added to each well and reverse transcription was carried out as follows: 42 °C-90:00, 42

°C-hold, 42 °C—12:20, 10 cycles (50 °C-2:00, 42 °C-2:00), 39 °C-12:00, 70 °C-15:00, 4 °C hold. After

the first 90 min of RT, 0.4 µL of 12,5 µM TSO (0.83 µM final concentration) was added to each reaction

at room temperature. The plate was resealed, centrifuged for 1 min at 700 g, and placed back on the

thermocycler to resume cycling from 42 °C-12:20. After reverse transcription, 15 µL of cDNA

enrichment PCR reaction mix containing 12.5 µL of 2× KAPA HiFi HotStart ReadyMix, 0.25 µL of

ISPCR primer (10 µM, /5Biosg/AAGCAGTGGTATCAACGCAGAGT-3′), and 2.25 µL of nuclease-

free water was added to each well and cDNA enrichment PCR was performed as follows for 21 cycles:

98 °C-3:00, 7 cycles (98 °C-0:20, 60 °C-4:00, and 72 °C-6:00), 7 cycles (98 °C-0:20, 64 °C-0:30, and

72 °C-6:00), and 7 cycles (98 °C-0:20, 67 °C-0:30, and 72 °C-7:00), 72 °C-10:00, 4 °C-hold. The cDNA

purification was performed using AMPure XP beads at a ratio of 1:1 (Beckmann). For elution, Qiagen

elution buffer (EB) was used. cDNA concentrations were measured using the Qubit 3.0 fluorometer

according to the manufacturer's protocol, and all cDNA samples were analyzed on Agilent Tape station

using the High Sensitivity DNA assay.

Dual-indexed Illumina Nextera XT sequencing libraries were prepared using 25 % of the recommended

reaction volumes of Nextera XT components. All Nextera XT libraries were prepared using 125 pg of

input cDNA for tagmentation, and were subsequently enriched for 12 PCR cycles, and purified using

AMPure XP beads (Beckmann) at a sample:bead ratio of 0.6:1. Library concentrations were measured

using Qubit 3.0 fluorometer, and the fragments were analyzed on the Agilent Tape station using the

High Sensitivity DNA assay. All libraries were individually normalized and diluted to a concentration

of pooled library to 5.4 ng/µL with average size ~515bp (16 nM). The libraries were sequenced on

Illumina NextSeq500 in a single-end 75-bp format (FC-404-2005, Illumina). Single-nuclei libraries

were sequenced at an average depth of 3 million reads.

Data availability – NGS:

(1) Illumina sRNA-seq and mRNA-seq datasets from F_0 and F_1 liver and eWAT (2) Illumina sRNA-

seq from F_0 sperm (3) Illumina mRNA-seq from offsprings sired from of miRNA-injected zygotes (F_m

/F_{MI}) eWAT (4) Illumina mRNA-seq from 1° eWA overexpressing *cel-mir-67* (negative control), *let-*

7d-5p and *let-7e-5p* mimics and (5) Smart-Seq2 single-cell mRNA-seq data from miRNA-injected 8-

cell embryos are available as Gene Expression Omnibus (GEO) SuperSeries GSE280278. The data are

available to the reviewers' discretion using **url**: https://tinyurl.com/3kdey4sk (**reviewer token**:

inqlkwqknhanxsn)

Data availability – Proteomics:

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via

the PRIDE partner repository with the dataset identifier PXD042800. The data are available to the

reviewer discretion: Username: reviewer_pxd042800@ebi.ac.uk, Password: QHj9fv33

Data availability – Metabolomics & lipidomics:

All metabolomic and lipidomic datasets are available at the Metabolomics Workbench[99] data repository

as project ID PR002238 (https://www.metabolomicsworkbench.org).

Male participants, sperm samples and analysis

The intervention study was approved by the Danish National Committee on Health Research Ethics

(SJ-769), the Danish Data Protection Agency (REG-048- 2019) and performed at the Fertility Clinic,

Zealand University Hospital, Denmark[77]. The semen samples were collected from obese patients

($\geq 30\,kg/m2$) via masturbation following two days of abstinence. The sperm analyses were performed

in the IVF laboratory in an authorized tissue establishment. The other semen samples were diluted with

MHM-C (Fujifilm) and frozen immediately for further analysis. The semen samples were collected via

masturbation following two days of abstinence, and the samples were analyzed a maximum of two to

three hours after production. After liquefaction, ejaculate volume and sperm concentration (total, motile

and progressive) were assessed using a microscope (Leica DM750). Sperm concentration was

determined with a CellVision counting chamber (CellVision Technology, Heerhugowaard, the

Netherlands). Total sperm count was calculated as the product between ejaculate volume and sperm

concentration. The sperm analyses were performed in the IVF laboratory in an authorized tissue

establishment.

Study design and approvals

This prospective intervention study was conducted between May 2020 and April 2022 at the Fertility

Clinic, Zealand University Hospital, Denmark. The study was approved by the Danish National

Committee on Health Research Ethics (SJ-769), the Danish Data Protection Agency (REG-048- 2019)

and registered on the Clinical Trial website (www. ClinicalTrials.gov identifier NCT04721938). All

participants provided written informed consent.

Primary adipocyte culture and transfection

Epididymal white adipose tissues (eWAT) from 6- to 8-week-old male C57BL/6J mice were removed,

minced, and digested with pre-warmed collagenase media (500 U/mL collagenase type 2 (LS004183,

Worthington) and 3.75U/mL DNase I (10104159001, Sigma). Isolated cells were seeded in 6-well

plates and grown in DMEM/Ham's F12 medium (21331, Vendor: Gibco), supplemented with 0.1 %

Biotin/D-Pantothenate (33 mM/17 mM), 1 % penicillin-streptomycin, and 20 % FBS. Until reaching

80 % confluency, primary cells were induced by induction medium with 1 μM rosiglitazone, 850 nM

insulin, 1 μM dexamethasone, 250 μM 3-isobutyl-1-methyl-xanthin (IBMX) and 10 % FBS.

Subsequently, the medium was changed every other day by a differentiation medium containing 1 μM

rosiglitazone, 850 nM insulin and 10 % FBS. 100 nM of Negative Control and *mmu-let-7-d/e-5p* mimics

or LNA oligos targeting *Dicer1* were reversely transfected using Lipofectamine RNAiMAX (Thermo

Fisher Scientific) at day 4 after induction. Completely differentiated cells were harvested at 7 days after

induction for further analysis.

Plasmid design and dual luciferase assay

For the generation of reporter constructs, the specific region of murine *Ago2*, *Dicer-1*, *Hif1an*, *Hk2*,

Cpt2, *Crat*, *Pcx*, *Idh2*, *Sdha*, *Sdhaf3*, *Pdk2*, *Slc25a22* and *Slc25a24* containing either wildtype (wt) or

mutated (mut) *let-7* binding sites was cloned behind the translation stop codon of the firefly-luciferase

gene in pmirGLO Dual-Luciferase vector using custom DNA oligos provided by IDT. Renilla luciferase

coding in the same plasmid was used as a control reporter for normalization purposes. To report the

miRNA activity in mammalian cells, HEK 293 cells were reversely transfected in 96-well plates with

60 ng DNA using the TransIT transfection reagent (mirusbio) for 3 days and forwardly transfected with

let-7d/e-5p mimics for 2 days. HEK-293 cells were cultured in DMEM medium (21013, Gibco)

supplemented with 5.6 mM L-glutamine, 1 % penicillin-streptomycin, and 10 % FBS. After 72-h vector

transfection and 48-h miRNA mimics transfection, dual luciferase reporter assays (Promega) were

performed. The relative activities of firefly luciferase indicating the interaction between miRNA and

specific region of targeted mRNA were normalized by the activities of Renilla luciferase.

TaqMan assay and general qPCR

To quantify the levels of mature miRNAs, TaqMan MicroRNA Reverse Transcription Kit (4366596,

Thermo Fisher), TaqMan MicroRNA Assays and TaqMan Universal Master Mix II (4440042, Thermo

Fisher) were applied according to manufacturer's protocols. Level of mature *hsa-miR-16* was used as

internal control in human samples. The relative expression of mature miRNA was calculated using a

comparative method ($2-\partial\partial$CT) according to the ABI Relative Quantification Method.

Determination of oxygen consumption rates and measurement of glycolytic activity:

Primary adipocytes from epididymal SVF (1°eWA) were seeded into Agilent Seahorse XFe96

Bioanalyzer microplates. Induction and differentiation were performed as described above. After four

days of differentiation, 16,000 per well mature adipocytes were re-seeded and transfected with LNA

oligos targeting *Dicer1* in DMEM/Ham's F-12 medium plus 10 % Fetal Bovine Serum, 1 % P/S, 0.1 %

Biotin and 0.1 % Pantothenic acid (Growth medium) at 37° C and 5 % CO_2. Seahorse measurement

was conducted at day 7 from differentiation. For each seahorse plate, the corresponding calibration plate

was prepared 24 hours prior to experiments using 200 µl XF Seahorse Calibrant Agilent per well.

Corresponding media were prepared before the experiment and consisted of Assay Medium

supplemented with 25 mM Glucose, 1 mM Glutamine, 2 mM Sodium Pyruvate, set to pH = 7.4 and

filtered sterile. The calibration plate possessed a cartridge having 4 pockets per well. Shortly before the

measurement pocket A was filled with 20 µl 20 µM Oligomycin, pocket B with 22 µl 20 µM FCCP and

pocket C with 25 µl 5 µM Antimycin A and Rotenone. One hour before measurements, plates were

washed with 1x PBS and changed to 180 µl Assay Medium in a non-CO_2 incubator at 37° C. Calibration

was started using calibration plates, measuring O_2 and pH LED Value/emission/Initial reference Delta

for each well. All measurements started with measuring basal values, followed by injection of

Oligomycin, FCCP, and Rotenone plus Antimycin A (MitoStressKit). Measurement parameters were:

Mix 3 min, wait 0 min, measure 3 min with each reagent's effect assessed within three consecutive

measurement cycles with a total duration of 18 min per reagent injection.

Protein extraction and western blot analysis

After being washed once by phosphate-buffered saline (PBS), the treated cells at harvest time were

lysed by RIPA buffer supplemented with cOmplete and PhosSTOP and quantified by Pierce BCA

Protein Assay Kit (23227, Thermo Fisher). Western blot analyses were carried out according to standard

protocols with primary antibodies against COXIV (Cell Signaling Technologies #4850, 1:1,1000

dilution), Cytochrome C (Cell Signaling Technologies #4280, 1:1,1000 dilution), HSP60 (Cell

Signaling Technologies #412165, 1:1,1000 dilution), PHB1 (Cell Signaling Technologies #2426,

1:1,1000 dilution), pyruvate dehydrogenase (Cell Signaling Technologies #3205, 1:1,1000 dilution),

SDHA (Cell Signaling Technologies #11998, 1:1,1000 dilution), SOD1 (Cell Signaling Technologies

#4266, 1:1,1000 dilution), VDAC (Cell Signaling Technologies #4661, 1:1,1000 dilution), HK2 (Cell

Signaling Technologies #2897, 1:1,1000 dilution), CPT2 (Abcam Ab181114, 1:1,1000 dilution) and

IDH2 (Cell Signaling Technologies #60322, 1:1,1000 dilution). Relative protein quantification was

normalized by HSC70 (Santa Cruz sc-7298, 1:1000 dilution).

Statistical analysis

Data were shown as mean ± standard error. One-way ANOVA and two-way ANOVA followed by

Tukey's multiple comparisons test were used for statistical analysis in *in vivo* experiments. Significant

differences between groups were noted in figure legends respectively.

Acknowledgements:

We thank the CECAD Proteomics Core Facility for support with proteomics analyses, the Laboratory

Manager, Stine Ravn, at The Zealand Fertility Clinic for handling and analysing the sperm samples.

We are thankful for technical support from Ronni Nielsen and specialists from the Functional Genomics

and Metabolism NGS team. We wish to thank Bente Møller and Irina Korshuniova from BRIC Single

Cell Genomics Core Facility for excellent help for Smart-seq2 experiments. Smart-seq2 work was partly

supported by CellX (The Danish Single Cell Examination Platform) funded by the Danish Research

Agency through the Danish national research infrastructure program (5229-0009B). JWK, JHP and

BDML received funding from the European Research Council Starting Grant TransGenRNA

(#675014). JWK, CH, and NS received funding from Sygeforsikring Denmark, University of Southern

Denmark and Danish Diabetes Academy that is funded by the Novo Nordisk Foundation

(NNF17SA0031406). JWK received support from the NNF Challenge (#33444) and Bioscience and

Basic Biomedicine Programs of the NNF (#28416). CH received grants from National Taiwan

University (113L7494) and the National Science and Technology, Taiwan (114-2314-B-002-055-).

MAM received support from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP)

and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and Coordenação de

Aperfeiçoamento de Pessoal de Nível Superior (CAPES). PK and MRF were supported by the

Vetenskapsrådet Consolidator Grant 2022-03953 ‚InSync'.

 Figure legends

Figure 1: Paternal obesity elicits heritable impairments in glucose metabolism and adipose

mitochondrial function. (a,b) Illustration of intergenerational *in vivo* cohorts depicting F_0 groups of

(a) paternal (F_0) obesity and weight-regression, male and female F_1 offspring mice **(b)** and analytical

workflows incl. RNA-seq for mRNA and sRNA, mass spectrometry for protein, and ultra-high pressure

liquid chromatography (UPLC) for metabolite and lipid differential regulation in F_0/F_1 epididymal white

adipose tissue (eWAT), liver, sperm and serum. **(c)** Body weights of F_0 LF (n=6), F_0 HF (n=13; n=6 for

15-24 weeks of age), F_0 HF-LF (n=7, *left*) and F_1 (F_0 LF, n=18), F_1 (F_0 HF, n=25) and F_1 (F_0 HF-LF,

n=21, *right*) male C57BL/6N mice. **(d)** Blood glucose during intraperitoneal glucose tolerance test in

F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=7, *left*) and F_1 (F_0 LF, n=10), F_1 (F_0 HF, n=10) and F_1 (F_0

HF-LF, n=10, *right*) male C57BL/6N mice. **(e)** Blood glucose during intraperitoneal insulin tolerance

test in F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=7, *left*) and F_1 (F_0 LF, n=9), F_1 (F_0 HF, n=10) and F_1

(F_0 HF-LF, n=10, *right*) male C57BL/6N mice. **(f)** Relative eWAT weights in F_0 LF (n=6), F_0 HF (n=6)

and F_0 HF-LF (n=7, *left*) and F_1 (F_0 LF, n=10), F_1 (F_0 HF, n=10) and F_1 (F_0 HF-LF, n=7, *right*) male

C57BL/6N mice. **(g)** Serum leptin levels measured by ELISA in F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-

LF (n=7, *left*) and F_1 (F_0 LF, n=8), F_1 (F_0 HF, n=8), and F_1 (F_0 HF-LF, n=8, *right*) male C57BL/6N mice.

(h) Fractional area size distribution of eWAT adipocytes in **(i)** F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF

(n=6, *left*) and F_1 (F_0 LF, n=6), F_1 (F_0 HF, n=6) and F_1 (F_0 HF-LF, n=6, *right*) male C57BL/6N mice.

(i,j) Representative haematoxylin and eosin staining of eWAT of **(i)** F_0 LF, F_0 HF and F_0 HF-LF and **(j)**

F_1 (F_0 LF), F_1 (F_0 HF), and F_1 (F_0 HF-LF) male C57BL/6N mice. **(k,l)** Visualisation of high-confidence

STRING interaction networks of proteins downregulated in **(k)** F_0 LF (n=6) vs F_0 HF (n=6) and **(l)** F_1

(F_0 LF, n=6), F_1 (F_0 HF, n=6) eWAT (*top*) and GOBP enrichment (*bottom*). Two-tailed Student's t-test

(c-e,h) for each timepoint and adipocyte sizes or one-way ANOVA followed by Tukey's multiple

comparisons **(f-h)** were used for statistical analysis. Data are presented as mean ± standard error with

individual values shown for n≤10. P-Values are indicated in the panel or represented as letters (**a**,

$p<0.05$, F_0 LF versus F_0 HF or F_1 (F_0 LF) versus F_1 (F_0 HF); **b**, $p<0.05$, F_0 HF versus F_0 HF-LF or F_1 (F_0

HF) versus F_1 (F_0 HF-LF); **c**, $p<0.05$, F_0 LF versus F_0 HF-LF or F_1 (F_0 LF) versus F_1 (F_0 HF-LF)).

Figure 2: Paternal obesity dysregulates miRNA:mRNA gene networks spurring intergenerational mitochondrial dysfunction. (a,b) Volcano plot of significantly up- (grey) and down-regulated (white) mRNAs in eWAT from **(a)** F_0 HF versus F_0 LF with 3,495 upregulated and 1,639 downregulated mRNAs and **(b)** F_1 (F_0 HF) versus F_1 (F_0 LF) with 1,284 up- and 702 down-regulated mRNAs. Plots show log10 transformed fold-changes (fc) and log10-transformed **(a)** adjusted or **(b)** non-adjusted p-Values of mRNA changes. Mitochondrial complex I-V genes are annotated in the plot. **(c,d)** Gene Set Enrichment Analysis and gene ontology enrichment of biological processes (GOBP) of differentially expressed mRNAs from **(a,b).** GOBP linked to mitochondrial respiration, ATP synthesis and Complex I assembly are marked with blue bars. **(e,f)** Volcano plot of significantly regulated miRNAs in eWAT from **(e)** F_0 HF versus F_0 LF with 154 up- and 193 down-regulated miRNAs and **(f)** F_1 (F_0 HF) versus F_1 (F_0 LF) with 63 up- and 87 downregulated miRNAs. Plots depict log10 transformed fc and log10-transformed p-Values of expression changes. Mitochondria-activating miRNAs are annotated in pink. **(g,h)** STRING interaction networks of **(g)** *miR-378/miR-143* and **(h)** *miR-532-5p* targets predicted by AGO2-CLIP (*left*) and network subset for eWAT proteins downregulated when comparing F_0 LF (n=6) versus F_0 HF (n=6) male C57BL/6N mice. Submodules containing mitochondrial proteins are marked by a red square and proteins marked by red dots are AGO2-CLIP predicted *let-7* targets and downregulated in F0 HF versus F_0 LF eWAT.

Figure 3: Obesity reversibly affects sperm sncRNomes and induces *let-7* in eWAT and sperm.
(a) Size distribution on nucleotide length of biotypes present in F_0 HF-LF sperm. Data presented is mean cpm per diet. **(b)** Size distribution on nucleotide length of biotypes present in F_0 HF sperm. Data presented is mean cpm per diet. **(c)** Size distribution on nucleotide length of biotypes present in F_0 LF sperm. Data presented is mean cpm per diet. **(d)** Fold change of F_0 HF vs F_0 LF sncRNA in sperm. Each points represents one sequence. **(e)** Fold change of F_0 HF-LF vs F_0 HF sncRNA in sperm. Each points represents one sequence. Dark blue= Mitochondrial tRNA, dark green=piRNA, green=ribosomal RNA, light green =long non-coding RNA, yellow=mitochondrial ribosomal RNA, orange= miRNA, dark orange=protein coding, red= miscellaneous RNA, dark red= tRNA, grey=unannotated. **(f)** Heatmap depicting scaled abundances of miRNAs in F_0 LF, F_0 HF and F_0 HF-LF (n=6/group) eWAT and sperm

as determined by sRNA-seq. **(g-i)** Abundance of *miR-143-3p*, *let-7d-5p* and in F_0 LF (n=6), F_0 HF (n=6)

and F_0 HF-LF (n=6) eWAT and sperm. One-way ANOVA followed by Tukey's multiple comparisons

(g-i) were used for statistical analysis. Data are presented as mean ± standard error with individual

values shown for n≤10. P-Values are indicated in the panel.

Figure 4: Zygotic delivery of *let-7* causes glucose intolerance in sired F_{MI} mice and impairs

embryonic mitochondria. (a) Illustration of experimental workflow for zygotic miRNA injection and

phenotypic characterisation of sired offsprings (F_{MI} mice). **(b)** Body weights of male F_{MI} mice sired

from with *cel-mir-67* (n=9), *let-7d-5p* (n=14) and *let-7e-5p* (n=14) injected zygotes. **(c,d)** Blood glucose

levels **(c)** and glucose AUC **(d)** during intraperitoneal glucose tolerance test in male F_{MI} mice sired from

cel-mir-67cel-mir-67 (n=7), *let-7d-5p* (n=10) and *let-7e-5p* (n=10) injected zygotes. **(e)** Blood glucose

levels during intraperitoneal insulin tolerance test in male F_{MI} mice sired from *cel-mir-67* (n=7), *let-7d-*

5p (n=9) and *let-7e-5p* (n=11) injected zygotes. **(f)** Random-fed blood glucose at indicated in male F_{MI}

mice sired from *cel-mir-67* (n=6), *let-7d-5p* (n=6) and *let-7e-5p* (n=6) injected zygotes **(g)** Relative

weights of indicated organs from male F_{MI} mice sired from *cel-mir-67* (n=7), *let-7d-5p* (n=6) and *let-*

7e-5p (n=6) injected zygotes. **(h)** GSEA-GOBP analysis of eWAT of male F_{MI} mice sired from *cel-mir-*

67 (n=5), *let-7d-5p* (n=6) and *let-7e-5p* (n=6) treated zygotes. GOBP linked to mitochondrial

respiration, ATP synthesis and Complex I assembly are marked with blue bars. **(i)** Illustration of

workflow to obtain single 8-cell stage blastomeres derived from zygotes injected with *cel-mir-67*, *let-*

7d-5p and *let-7e-5p* subjected to smart-seq2 single-cell sequencing. **(j)** Multidimensional scaling plots

of single-cell RNAseq profiles in 8-cell stage blastomeres obtained from *cel-mir-67* (n=22 blastomeres

derived from n=3 zygotes), *let-7d-5p* (n=19 cells blastomeres from n=3 zygotes) and *let-7e-5p* (n=20

blastomeres derived from n=4 zygotes). **(l,m)** Correlation plot and regression analysis between

differential BMI loss (ΔBMI) and expression changes (ΔCt) in human semen **(l)** *LET-7D-5P* and

(m)*LET-7E-5P.* Two-tailed Student's t-test **(c,e)** per timepoint, one-way ANOVA followed by Tukey's

multiple comparisons **(d,g)** or Pearson's regression analysis **(l,m)** were used for statistical analysis. Data

are presented as mean ± standard error with individual values shown for n≤10. P-Values are indicated

in the panel or represented as letters (**a**, *p<0.05*, *cel-mir-67* versus *let-7d-5p*, **b**, *p<0.05*, *cel-mir-67*

versus *let-7e-5p*).

Figure 5: Adipocyte *let-7* blunts mitochondrial function via a novel let-7-DICER1 axis

(a) Visualisation of high-confidence STRING interaction networks of *let-7* targets predicted by AGO2-

CLIP miRNA:mRNA interactions and subsetted for proteins downregulated comparing F_0 LF (n=6)

versus F_0 HF (n=6) in eWAT from male C57BL/6N mice. Submodules containing mitochondrial

proteins are marked by a red square and proteins marked by red dots are AGO2-CLIP predicted *let-7*

targets and downregulated in F0 HF versus F_0 LF eWAT. **(b,c)** Volcano plot of significantly up- (grey)

and down-regulated (white) genes in 1°eWA transfected with **(b)** *let-7-d-5p* versus *cel-mir-67*, yielding

974 up- and 1,295 down-regulated genes and **(c)** *let-7-e-5p* versus *cel-mir-67*, yielding 754 up- and 633

down-regulated genes. Genes connected to mitochondrial regulation are marked by blue, genes involved

in miRNA maturation by red dots. **(d)** Relative luciferase activity in HEK293 cells after transfection of

pmirGLO dual luciferase reporter assay (DLRA) constructs harbouring wild-type 3′-UTR or exon

containing *let-7* miRNA responsive elements (MRE) of indicated mitochondrial genes and **(e)** wild-

type and mutated *let-7* MREs in *Dicer1* transfected with 100 nM of *cel-mir-67*, *let-7d-5p* and *let-7d-5p*

mimics. Data represent n≥3 independent biological replicates, each performed in n=3 technical

replicates. **(f-h)** Western blot analysis of 1°eWA transfected with *cel-mir-67*, *let-7d-5p* and *let-7d-5p*

mimics performed using the indicated antibodies detecting **(f)** mitochondrial and **(g)** DICER1 proteins.

HSC70 served as loading control. **(h)** Densitometric quantification of (**f,g**) representing n=3 biological

replicates. **(i)** Western blot (*top*) and quantitative RT-PCR (*bottom*) quantification of DICER1 in

1°eWA transfected using two independent locked nucleic acid (LNA) inhibitors targeting *Dicer1*

mRNA. Data represent n=3 biological replicates, each performed in n=3 technical replicates. **(j,k)**

Oxygen consumption rates (**j,** OCR) and extracellular acidification rates (**k**, ECAR) in 1°eWA

transfected with 100 nM of *negative control LNA*, *Dicer1* LNA1 and *Dicer1* LNA2 and stimulated with

oligomycin (O), FCCP (F) and Antimycin A plus Rotenone (A/R). Experiments are representative of

n≥3 independent biological replicates. **(l)** Western blot analysis of eWAT from F_0 LF (n=4), F_0 HF

(n=4), F_0 HF-LF (n=4) male C57BL/6N mice performed using the indicated antibodies detecting

mitochondrial and DICER1 proteins. HSP90 served as loading control. **(m)** Densitometric

quantification of **(l)** representing n=4 biological replicates. **(n)** Illustration of proposed molecular

mechanism connecting obesity and elevated *let-7* in adipocytes to post-transcriptional silencing of

DICER1 and impaired oxidative phosphorylation and respiration. Two-tailed Student's t-test **(j,k)** for

each timepoint or one-way ANOVA **(d,e,h)** followed by Tukey's multiple comparisons test were used

for statistical analysis. Data are presented as mean ± standard error with individual values shown for

n≤10. P-Values are indicated in the panel or represented as letters (**a**, *p<0.05*, *cel-mir-67* versus *let-7d-*

5p; **b**, *p<0.05*, *cel-mir-67* versus *let-7e-5p*).

 Supplementary figure legends:

 Figure S1: Paternal obesity and weight loss do not affect reproductive fitness and litter sizes. (a)

 Total numbers, number of male and female offsprings, sex ratios and **(b)** litter sizes of pups conceived

 by (F_0 LF, n=8), (F_0 HF, n=8) and F_1 (F_0 HF-LF, n=8) bucks.

 Figure S2: Paternal obesity causes heritable alterations in glucose metabolism in male but not in

 female mice. (a) Blood glucose area under the curve (AUC) of glucose levels during intraperitoneal

 glucose tolerance test in F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=7, *left*) mice F_1 (F_0 LF, n=10), F_1

 (F_0 HF, n=10) and F_1 (F_0 HF-LF, n=10, *right*) mice. **(b)** Serum insulin levels as measured by ELISA in

 F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=7, *left*) and F_1 (F_0 LF, n=6), F_1 (F_0 HF, n=8) and F_1 (F_0 HF-

 LF, n=8) mice. **(c)** Relative inguinal white adipose tissue (iWAT) weights in F_0 LF (n=6), F_0 HF (n=6)

 and F_0 HF-LF (n=7, *left*) and F_1 (F_0 LF, n=10), F_1 (F_0 HF, n=10) and F_1 (F_0 HF-LF, n=7, *right*) mice. **(d)**

 Body weights of F_1 (F_0 LF, n=11), F_1 (F_0 HF, n=12) and F_1 (F_0 HF-LF, n=13) female C57BL/6N mice.

 (e) Blood glucose during intraperitoneal glucose tolerance test in F_1 (F_0 LF, n=8), F_1 (F_0 HF, n=10) and

 F_1 (F_0 HF-LF, n=10) female C57BL/6N mice. **(f)** AUC of glucose levels during intraperitoneal glucose

 tolerance test in F_1 (F_0 LF, n=8), F_1 (F_0 HF, n=10) and F_1 (F_0 HF-LF, n=10) female C57BL/6N mice. **(g)**

 Blood glucose during intraperitoneal insulin tolerance test in F_1 (F_0 LF, n=7), F_1 (F_0 HF, n=9) and F_1 (F_0

 HF-LF, n=9) female C57BL/6N mice. **(h)** Random-fed serum glucose levels in F_1 (F_0 LF, n=13), F_1 (F_0

 HF, n=12) and F_1 (F_0 HF-LF, n=13) female C57BL/6N mice. **(i)** Relative organ weights in indicated

 tissue from F_1 (F_0 LF, n=9), F_1 (F_0 HF, n=10) and F_1 (F_0 HF-LF, n=8) female C57BL/6N mice. Two-

 tailed Student's t-test **(d,e)** for individual timepoints or one-way ANOVA followed by Tukey's multiple

 comparisons **(a-c,f)** were used for statistical analysis. Data are presented as mean ± standard error with

 individual values shown for n≤10. p-Values are indicated in the panel or represented as letters (**a**,

 $p<0.05$, F_0 LF versus F_0 HF or F_1 (F_0 LF) versus F_1 (F_0 HF); **b**, $p<0.05$, F_0 HF versus F_0 HF-LF or F_1 (F_0

 HF) versus F_1 (F_0 HF-LF); **c**, $p<0.05$, F_0 LF versus F_0 HF-LF or F_1 (F_0 LF) versus F_1 (F_0 HF-LF)).

 Figure S3: Effects of paternal obesity on F_0 and F_1 eWAT proteomes. (a,b) Visualisation of high-

 confidence STRING interaction networks of proteins upregulated in **(a)** F_0 LF (n=6) vs F_0 HF (n=6) and

1054   **(b)** F_1 (F_0 LF, n=6), F_1 (F_0 HF, n=6) eWAT (*top*) and GOBP enrichment of differentially regulated

1055   proteins (*bottom*).

1056

Figure S4: Paternal obesity elicits alterations in circulating mitochondria-associated metabolites

and lipids. (a-c) Principal component analysis (PCA) representing PC1-2 **(a)** heatmap of the 74 most

differentially regulated serum metabolites **(c)** and (**b**) GOBP term enrichment obtained by integrating

differentially abundant serum metabolites and differentially expressed eWAT genes determined by

RNA-seq in eWAT of F_0 LF versus F_0 HF male C57L/6N mice and integrated using by Metaboanalyst

6.0. (**d,e**) PCA representing PC1-2 **(d)** and heatmap of the 100 most differentially regulated serum lipids

(e). Metabolites and lipids were determined by ultra-performance liquid chromatography in serum of

F_0 LF compared to F_0 HF (n=6 per group) male C57BL/6N mice. **(f-h)** Relative abundances of **(f)**

mitochondrial tricarbon acid intermediates **(g)** mitochondrial Carnitine lipid carriers **(h)** mitochondria-

inactivating short-chain and mitochondria-promoting long-chain ceramides in serum of F_0 LF (n=6), F_0

HF (n=6) and F_0 HF-LF (n=7) mice. One-way ANOVA followed by Tukey's multiple comparisons test

were used for statistical analysis and P vales are indicated in each panel.

1069

Figure S5: Paternal weight loss corrects obesity-associated mitochondrial dysfunction in eWAT.

(a,b) Volcano plot of significantly up- (grey) and down-regulated (white) mRNAs in eWAT from **(a)**

F_0 HF versus F_0 HF-LF with 2,779 up- and 1,741 down-regulated mRNAs and (**b**) F_1 (F_0 HF) versus F_1

(F_0 HF-F) with 1,118 up- and 550 down-regulated mRNAs. Plots show log10 transformed fold-changes

(fc) and log10-transformed **(a)** adjusted or **(b)** non-adjusted p-Values of mRNA changes. Mitochondrial

complex genes are annotated in the plot. **(c,d)** GSEA and GOBP enrichment of differentially expressed

mRNAs from **(a,b).** Terms linked to mitochondrial respiration, ATP synthesis and Complex I are

marked with blue bars. **(e,f)** Volcano plot of significantly regulated miRNAs in eWAT from **(e)** F_0 HF

versus F_0 HF-F with 147 up- and 148 down-regulated miRNAs and (**f**) F_1 (F_0 HF) versus F_1 (F_0 HF-LF)

with 92 up- and 37 downregulated miRNAs. Plots depict log10 transformed fc and p-Values of mRNA

changes. Mitochondria-activating miRNAs are annotated in pink.

1081

Figure S6: Paternal obesity does not affect F$_1$ adipose miRNA expression and meta-inflammation.
(a,b) Multidimensional scaling (MDS) plots and PC1/2 of mRNA changes in eWAT of **(a)** F$_0$ LF (n=6),
F$_0$ HF (n=6) and F$_0$ HF-LF (n=6) and **(b)** F$_1$ (F$_0$ HF, n=4), F$_1$ (F$_0$ HF, n=6) and F$_1$ (F$_0$ HF-LF, n=6) male
C57BL/6N mice determined by mRNA-seq. **(c,d)** Volcano plot of significantly regulated miRNAs in
eWAT from **(c)** F$_0$ HF versus F$_0$ LF with 154 up- and 193 down-regulated miRNAs and **(d)** F$_1$ (F$_0$ HF)
versus F$_1$ (F$_0$ LF) with 62 up- and 87 down-regulated miRNAs. Plots depict log10 transformed fc and
p-Values of mRNA changes. Mitochondria-inactivating miRNAs are annotated in light blue. **(e,f)**
Expression changes reflected by normalised mRNA-seq counts from **(e)** F$_0$ LF (n=6), F$_0$ HF (n=6) and
F$_0$ HF-LF (n=6) and **(f)** F$_1$ (F$_0$ HF, n=4), F$_1$ (F$_0$ HF, n=5) and F$_1$ (F$_0$ HF-LF, n=6) male C57BL/6N mice.
Genes represent markers of classically (M1) or alternatively (M2) activated macrophages/monocytes.
One-way ANOVA followed by Tukey's multiple comparisons **(e,f)** were used for statistical analysis.
Data are presented as mean ± standard error with individual values shown for n≤10. p-Values are
represented as letters (**a**, $p<0.05$, F$_0$ LF versus F$_0$ HF or F$_1$ (F$_0$ LF) versus F$_1$ (F$_0$ HF); **b**, $p<0.05$, F$_0$ HF
versus F$_0$ HF-LF or F$_1$ (F$_0$ HF) versus F$_1$ (F$_0$ HF-LF); **c**, $p<0.05$, F$_0$ LF versus F$_0$ HF-LF or F$_1$ (F$_0$ LF)
versus F$_1$ (F$_0$ HF-LF)).

Figure S7: Paternal obesity does not affect mitochondrial gene expression in F$_0$ and F$_1$ liver. (a,b)
MDS plots and PC1/2 of mRNA changes in liver of **(a)** F$_0$ LF (n=6), F$_0$ HF (n=6) and F$_0$ HF-LF (n=6)
and **(b)** F$_1$ (F$_0$ HF, n=4), F$_1$ (F$_0$ HF, n=5) and F$_1$ (F$_0$ HF-LF, n=6) male C57BL/6N mice determined by
mRNA-seq. **(c,d)** Volcano plot of significantly up-(grey) and down-regulated (white) genes in liver
from **(c)** F$_0$ HF versus F$_0$ LF, 3,312 up- and 1,597 down-regulated genes and **(d)** F$_1$ (F$_0$ HF) versus F$_1$ (F$_0$
LF), 457 upregulated and 1,041 downregulated genes. Plots depict log10 transformed fold-changes and
log10- transformed **(c)** adjusted or **(d)** non-adjusted p-Values of expression changes. **(e,f)** GSEA and
GOBP enrichment of biological processes expression changes shown in (**c,d**).

Figure S8: Paternal obesity and weight loss does not affect liver morphology and lipid metabolism
in F$_1$ progeny. (a-f) Representative **(a,b)** histological (haematoxylin/eosine staining cellular
structures), **(c,d)** fibrosis (Sirius Red staining collagen) and **(e,f)** pan-leukocyte (anti-CD45 stains

leukocytes) stained formalin-fixed paraffin embedded sections in livers of **(a,c,e)** F_0 LF, F_0 HF and F_0 HF-LF and **(b,d,f)** F_1 (F_0 LF), F_1 (F_0 HF) and F_1 (F_0 HF-LF) male C57BL/6N mice. **(g,h)** Pathologist assessment of macrovesicular and microvesicular steatosis, lobular inflammation and NAS fibrosis scores in livers of livers of F_0 LF, F_0 HF and F_0 HF-LF as well as in F_1 (F_0 LF), F_1 (F_0 HF) and F_1 (F_0 HF-LF, n=6 per group). **(i,j)** Expression changes reflected by normalised mRNA-seq counts from F_0 LF (n=6), F_0 HF (n=6) and F_0 HF-LF (n=6, *left*) and **(f)** F_1 (F_0 HF, n=4), F_1 (F_0 HF, n=5) and F_1 (F_0 HF-LF, n=5, *right*) male C57BL/6N mice. Data are shown for genes in liver **(i)** lipid and lipoprotein metabolism and **(j)** mitochondrial beta-oxidation. Data in **(i,j)** are presented as mean ± standard error with individual values shown for n≤10. p-Values are shown in the panels or represented as letters (**a**, *p*<0.05, F_0 LF versus F_0 HF or F_1 (F_0 LF) versus F_1 (F_0 HF); **b**, *p*<0.05, F_0 HF versus F_0 HF-LF or F_1 (F_0 HF) versus F_1 (F_0 HF-LF); **c**, *p*<0.05, F_0 LF versus F_0 HF-LF or F_1 (F_0 LF) versus F_1 (F_0 HF-LF)).

Figure S9: Paternal obesity and weight loss cause only modest changes in spermatozoa mitochondrial tRNA fragments. Fragments with a mean cpm above 10 shown. **(a)** Sequences mapped to mitochondrial tRNA, presented as mean cpm per diet. **(b)** Sum cpm per mitochondrial tsRNA fragment type. Each point represents one sample. (*p<0.05, One-way ANOVA with Tukey's multiple comparisons). n=6 per diet, pink=F_0 LF, beige= F_0 HF, blue= F_0 HF-LF.

Figure S10: Zygotic microinjection of obesity associated *let-7* does not affect embryonic viability and implantation.
Total numbers of zygotes microinjected with *cel-mir-67*, *let-7d-5p* and *let-7e-5p*, total numbers of pups born and relative survival rates of miRNA-proficient embryos.

Figure S11: Obesity-associated let-7 impairs adipocyte mitochondrial function via silencing of DICER1. GSEA-GOBP analysis of differentially abundant mRNAs in 8-cell blastomeres derived from *cel-mir-67* (n=22), *let-7d-5p* (n=19) and *let-7e-5p* (n=20) injected zygotes (corresponding **Fig.4j**).

<center>**References:**</center>

1138

1　Trerotola, M., Relli, V., Simeone, P. & Alberti, S. Epigenetic inheritance and the
missing heritability. *Human genomics* **9**, 17, doi:10.1186/s40246-015-0041-3 (2015).
2　Horsthemke, B. A critical view on transgenerational epigenetic inheritance in
humans. *Nature communications* **9**, 2973, doi:10.1038/s41467-018-05445-5 (2018).
3　Bygren, L. O., Kaati, G. & Edvinsson, S. Longevity determined by paternal ancestors'
nutrition during their slow growth period. *Acta Biotheor* **49**, 53-59,
doi:10.1023/a:1010241825519 (2001).
4　Pembrey, M. E. *et al.* Sex-specific, male-line transgenerational responses in humans.
European journal of human genetics : EJHG **14**, 159-166,
doi:10.1038/sj.ejhg.5201538 (2006).
5　Lumey, L. H., Stein, A. D., Kahn, H. S. & Romijn, J. A. Lipid profiles in middle-aged
men and women after famine exposure during gestation: the Dutch Hunger Winter
Families Study. *Am J Clin Nutr* **89**, 1737-1743, doi:10.3945/ajcn.2008.27038 (2009).
6　Carone, B. R. *et al.* Paternally induced transgenerational environmental
reprogramming of metabolic gene expression in mammals. *Cell* **143**, 1084-1096,
doi:10.1016/j.cell.2010.12.008 (2010).
7　Ng, S. F. *et al.* Paternal high-fat diet consumption induces common changes in the
transcriptomes of retroperitoneal adipose and pancreatic islet tissues in female rat
offspring. *FASEB journal : official publication of the Federation of American Societies*
for Experimental Biology **28**, 1830-1841, doi:10.1096/fj.13-244046 (2014).
8　de Castro Barbosa, T. *et al.* High-fat diet reprograms the epigenome of rat
spermatozoa and transgenerationally affects metabolism of the offspring. *Molecular*
metabolism **5**, 184-197, doi:10.1016/j.molmet.2015.12.002 (2016).
9　de Castro Barbosa, T., Alm, P. S., Krook, A., Barres, R. & Zierath, J. R. Paternal
high-fat diet transgenerationally impacts hepatic immunometabolism. *FASEB journal*
: official publication of the Federation of American Societies for Experimental Biology
33, 6269-6280, doi:10.1096/fj.201801879RR (2019).
10　Wang, Y. *et al.* Sperm microRNAs confer depression susceptibility to offspring. *Sci*
Adv **7**, doi:10.1126/sciadv.abd7605 (2021).
11　Dias, B. G. & Ressler, K. J. Parental olfactory experience influences behavior and
neural structure in subsequent generations. *Nature neuroscience* **17**, 89-96,
doi:10.1038/nn.3594 (2014).
12　Gapp, K. *et al.* Implication of sperm RNAs in transgenerational inheritance of the
effects of early trauma in mice. *Nature neuroscience* **17**, 667-669,
doi:10.1038/nn.3695 (2014).
13　Sun, W. *et al.* Cold-induced epigenetic programming of the sperm enhances brown
adipose tissue activity in the offspring. *Nature medicine* **24**, 1372-1383,
doi:10.1038/s41591-018-0102-y (2018).
14　Skvortsova, K., Iovino, N. & Bogdanovic, O. Functions and mechanisms of epigenetic
inheritance in animals. *Nature reviews. Molecular cell biology* **19**, 774-790,
doi:10.1038/s41580-018-0074-2 (2018).
15　Rechavi, O., Minevich, G. & Hobert, O. Transgenerational inheritance of an acquired
small RNA-based antiviral response in C. elegans. *Cell* **147**, 1248-1256,
doi:10.1016/j.cell.2011.10.042 (2011).
16　Fire, A. *et al.* Potent and specific genetic interference by double-stranded RNA in
Caenorhabditis elegans. *Nature* **391**, 806-811, doi:10.1038/35888 (1998).
17　Ashe, A. *et al.* piRNAs can trigger a multigenerational epigenetic memory in the
germline of C. elegans. *Cell* **150**, 88-99, doi:10.1016/j.cell.2012.06.018 (2012).
18　de Vanssay, A. *et al.* Paramutation in Drosophila linked to emergence of a piRNA-
producing locus. *Nature* **490**, 112-115, doi:10.1038/nature11416 (2012).
19　Rassoulzadegan, M. *et al.* RNA-mediated non-mendelian inheritance of an
epigenetic change in the mouse. *Nature* **441**, 469-474, doi:10.1038/nature04674
(2006).

20 Gross, N., Kropp, J. & Khatib, H. MicroRNA Signaling in Embryo Development.
Biology (Basel) **6**, doi:10.3390/biology6030034 (2017).
21 Sharma, U. *et al.* Small RNAs Are Trafficked from the Epididymis to Developing
Mammalian Sperm. *Developmental cell* **46**, 481-494 e486,
doi:10.1016/j.devcel.2018.06.023 (2018).
22 Natt, D. *et al.* Human sperm displays rapid responses to diet. *PLoS Biol* **17**,
e3000559, doi:10.1371/journal.pbio.3000559 (2019).
23 Tomar, A. *et al.* Epigenetic inheritance of diet-induced and sperm-borne
mitochondrial RNAs. *Nature* **630**, 720-727, doi:10.1038/s41586-024-07472-3 (2024).
24 Grandjean, V. *et al.* RNA-mediated paternal heredity of diet-induced obesity and
metabolic disorders. *Scientific reports* **5**, 18193, doi:10.1038/srep18193 (2015).
25 Rodgers, A. B., Morgan, C. P., Leu, N. A. & Bale, T. L. Transgenerational epigenetic
programming via sperm microRNA recapitulates effects of paternal stress.
Proceedings of the National Academy of Sciences of the United States of America
112, 13699-13704, doi:10.1073/pnas.1508347112 (2015).
26 Sharma, U. *et al.* Biogenesis and function of tRNA fragments during sperm
maturation and fertilization in mammals. *Science*, doi:10.1126/science.aad6780
(2015).
27 Korhonen, H. M. *et al.* Dicer is required for haploid male germ cell differentiation in
mice. *PloS one* **6**, e24821, doi:10.1371/journal.pone.0024821 (2011).
28 Bernstein, E. *et al.* Dicer is essential for mouse development. *Nature genetics* **35**,
215-217, doi:10.1038/ng1253 (2003).
29 Chen, Q. *et al.* Sperm tsRNAs contribute to intergenerational inheritance of an
acquired metabolic disorder. *Science*, doi:10.1126/science.aad7977 (2015).
30 Rodgers, A. B., Morgan, C. P., Bronson, S. L., Revello, S. & Bale, T. L. Paternal
stress exposure alters sperm microRNA content and reprograms offspring HPA
stress axis regulation. *The Journal of neuroscience : the official journal of the Society*
for Neuroscience **33**, 9003-9012, doi:10.1523/JNEUROSCI.0914-13.2013 (2013).
31 Cropley, J. E., Suter, C. M., Beckman, K. B. & Martin, D. I. Germ-line epigenetic
modification of the murine A vy allele by nutritional supplementation. *Proceedings of*
the National Academy of Sciences of the United States of America **103**, 17308-
17312, doi:10.1073/pnas.0607090103 (2006).
32 Zhang, Y., Shi, J., Rassoulzadegan, M., Tuorto, F. & Chen, Q. Sperm RNA code
programmes the metabolic health of offspring. *Nature reviews. Endocrinology* **15**,
489-498, doi:10.1038/s41574-019-0226-2 (2019).
33 Sun, K., Kusminski, C. M. & Scherer, P. E. Adipose tissue remodeling and obesity.
The Journal of clinical investigation **121**, 2094-2101, doi:10.1172/JCI45887 (2011).
34 Scherer, P. E. Adipose tissue: from lipid storage compartment to endocrine organ.
Diabetes **55**, 1537-1545, doi:10.2337/db06-0263 (2006).
35 Lee, J. H. *et al.* The Role of Adipose Tissue Mitochondria: Regulation of
Mitochondrial Function for the Treatment of Metabolic Diseases. *Int J Mol Sci* **20**,
doi:10.3390/ijms20194924 (2019).
36 Czech, M. P. Mechanisms of insulin resistance related to white, beige, and brown
adipocytes. *Molecular metabolism* **34**, 27-42, doi:10.1016/j.molmet.2019.12.014
(2020).
37 Kahn, C. R., Wang, G. & Lee, K. Y. Altered adipose tissue and adipocyte function in
the pathogenesis of metabolic syndrome. *The Journal of clinical investigation* **129**,
3990-4000, doi:10.1172/JCI129187 (2019).
38 Bartelt, A. & Heeren, J. Adipose tissue browning and metabolic health. *Nature*
reviews. Endocrinology **10**, 24-36, doi:10.1038/nrendo.2013.204 (2014).
39 Schottl, T., Kappler, L., Braun, K., Fromme, T. & Klingenspor, M. Limited
mitochondrial capacity of visceral versus subcutaneous white adipocytes in male
C57BL/6N mice. *Endocrinology* **156**, 923-933, doi:10.1210/en.2014-1689 (2015).

40   Heinonen, S., Jokinen, R., Rissanen, A. & Pietilainen, K. H. White adipose tissue
mitochondrial metabolism in health and in obesity. *Obes Rev* **21**, e12958,
doi:10.1111/obr.12958 (2020).
41   Kusminski, C. M. *et al.* MitoNEET-driven alterations in adipocyte mitochondrial
activity reveal a crucial adaptive process that preserves insulin sensitivity in obesity.
Nature medicine **18**, 1539-1549, doi:10.1038/nm.2899 (2012).
42   Gonzalez-Franquesa, A. *et al.* Remission of obesity and insulin resistance is not
sufficient to restore mitochondrial homeostasis in visceral adipose tissue. *Redox Biol*
54, 102353, doi:10.1016/j.redox.2022.102353 (2022).
43   Schottl, T., Kappler, L., Fromme, T. & Klingenspor, M. Limited OXPHOS capacity in
white adipocytes is a hallmark of obesity in laboratory mice irrespective of the
glucose tolerance status. *Molecular metabolism* **4**, 631-642,
doi:10.1016/j.molmet.2015.07.001 (2015).
44   Soro-Arnaiz, I. *et al.* Role of Mitochondrial Complex IV in Age-Dependent Obesity.
Cell reports **16**, 2991-3002, doi:10.1016/j.celrep.2016.08.041 (2016).
45   Wang, Y. *et al.* Metformin Improves Mitochondrial Respiratory Activity through
Activation of AMPK. *Cell reports* **29**, 1511-1523 e1515,
doi:10.1016/j.celrep.2019.09.070 (2019).
46   Borralho, P. M., Rodrigues, C. M. & Steer, C. J. microRNAs in Mitochondria: An
Unexplored Niche. *Adv Exp Med Biol* **887**, 31-51, doi:10.1007/978-3-319-22380-3_3
(2015).
47   Bandiera, S. *et al.* Nuclear outsourcing of RNA interference components to human
mitochondria. *PloS one* **6**, e20746, doi:10.1371/journal.pone.0020746 (2011).
48   Barrey, E. *et al.* Pre-microRNA and mature microRNA in human mitochondria. *PloS*
one **6**, e20220, doi:10.1371/journal.pone.0020220 (2011).
49   Mori, M. A. *et al.* Role of MicroRNA Processing in Adipose Tissue in Stress Defense
and Longevity. *Cell metabolism* **16**, 336-347, doi:S1550-4131(12)00322-1 [pii]
10.1016/j.cmet.2012.07.017 (2012).
50   Mori, M. A. *et al.* Altered miRNA processing disrupts brown/white adipocyte
determination and associates with lipodystrophy. *The Journal of clinical investigation*
124, 3339-3351, doi:10.1172/JCI73468 (2014).
51   Oliverio, M. *et al.* Dicer1-miR-328-Bace1 signalling controls brown adipose tissue
differentiation and function. *Nature cell biology* **18**, 328-336, doi:10.1038/ncb3316
(2016).
52   Huypens, P. *et al.* Epigenetic germline inheritance of diet-induced obesity and insulin
resistance. *Nature genetics* **48**, 497-499, doi:10.1038/ng.3527 (2016).
53   Turpin, S. M. *et al.* Obesity-induced CerS6-dependent C16:0 ceramide production
promotes weight gain and glucose intolerance. *Cell metabolism* **20**, 678-686,
doi:10.1016/j.cmet.2014.08.002 (2014).
54   Hammerschmidt, P. *et al.* CerS6-dependent ceramide synthesis in hypothalamic
neurons promotes ER/mitochondrial stress and impairs glucose homeostasis in
obese mice. *Nature communications* **14**, 7824, doi:10.1038/s41467-023-42595-7
(2023).
55   Cui, Q., Yu, Z., Purisima, E. O. & Wang, E. Principles of microRNA regulation of a
human cellular signaling network. *Mol Syst Biol* **2**, 46, doi:10.1038/msb4100089
(2006).
56   Hong, H. *et al.* MicroRNA-143 promotes cardiac ischemia-mediated mitochondrial
impairment by the inhibition of protein kinase Cepsilon. *Basic Res Cardiol* **112**, 60,
doi:10.1007/s00395-017-0649-7 (2017).
57   Das, S. *et al.* miR-181c regulates the mitochondrial genome, bioenergetics, and
propensity for heart failure in vivo. *PloS one* **9**, e96820,
doi:10.1371/journal.pone.0096820 (2014).
58   Roman, B. *et al.* Nuclear-mitochondrial communication involving miR-181c plays an
important role in cardiac dysfunction during obesity. *J Mol Cell Cardiol* **144**, 87-96,
doi:10.1016/j.yjmcc.2020.05.009 (2020).

59   Alhadidi, Q. M. *et al.* MiR-182 Inhibition Protects Against Experimental Stroke in vivo
and Mitigates Astrocyte Injury and Inflammation in vitro via Modulation of Cortactin
Activity. *Neurochemical research* **47**, 3682-3696, doi:10.1007/s11064-022-03718-6
(2022).
60   Lee, H. *et al.* microRNA-200a-3p enhances mitochondrial elongation by targeting
mitochondrial fission factor. *BMB reports* **50**, 214-219,
doi:10.5483/bmbrep.2017.50.4.006 (2017).
61   Carrer, M. *et al.* Control of mitochondrial metabolism and systemic energy
homeostasis by microRNAs 378 and 378. *Proceedings of the National Academy of*
Sciences of the United States of America **109**, 15330-15335, doi:1207605109 [pii]
10.1073/pnas.1207605109 (2012).
62   Wang, J. X. *et al.* MicroRNA-532-3p regulates mitochondrial fission through targeting
apoptosis repressor with caspase recruitment domain in doxorubicin cardiotoxicity.
Cell Death Dis **6**, e1677, doi:10.1038/cddis.2015.41 (2015).
63   Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach
for interpreting genome-wide expression profiles. *Proceedings of the National*
Academy of Sciences of the United States of America **102**, 15545-15550,
doi:0506580102 [pii]
10.1073/pnas.0506580102 (2005).
64   Madsen, S. *et al.* Rapid downregulation of DICER is a hallmark of adipose tissue
upon high-fat diet feeding. *Molecular and cellular endocrinology* **595**, 112413,
doi:10.1016/j.mce.2024.112413 (2025).
65   Brieno-Enriquez, M. A. *et al.* Exposure to endocrine disruptor induces
transgenerational epigenetic deregulation of microRNAs in primordial germ cells.
PloS one **10**, e0124296, doi:10.1371/journal.pone.0124296 (2015).
66   Zhu, H. *et al.* The Lin28/let-7 axis regulates glucose metabolism. *Cell* **147**, 81-94,
doi:10.1016/j.cell.2011.08.033 (2011).
67   Frost, R. J. & Olson, E. N. Control of glucose homeostasis and insulin sensitivity by
the Let-7 family of microRNAs. *Proceedings of the National Academy of Sciences of*
the United States of America **108**, 21075-21080, doi:10.1073/pnas.1118922109
(2011).
68   Fullston, T. *et al.* Paternal obesity initiates metabolic disturbances in two generations
of mice with incomplete penetrance to the F2 generation and alters the transcriptional
profile of testis and sperm microRNA content. *FASEB journal : official publication of*
the Federation of American Societies for Experimental Biology **27**, 4226-4243,
doi:10.1096/fj.12-224048 (2013).
69   Cheong, A. W. *et al.* MicroRNA Let-7a and dicer are important in the activation and
implantation of delayed implanting mouse embryos. *Human reproduction* **29**, 750-
762, doi:10.1093/humrep/det462 (2014).
70   Liu, W. *et al.* Expression of microRNA let-7 in cleavage embryos modulates cell fate
determination and formation of mouse blastocystsdagger. *Biol Reprod* **107**, 1452-
1463, doi:10.1093/biolre/ioac181 (2022).
71   Kolenda, T., Przybyla, W., Teresiak, A., Mackiewicz, A. & Lamperska, K. M. The
mystery of let-7d - a small RNA with great power. *Contemp Oncol (Pozn)* **18**, 293-
301, doi:10.5114/wo.2014.44467 (2014).
72   Yuan, S. *et al.* Sperm-borne miRNAs and endo-siRNAs are important for fertilization
and preimplantation embryonic development. *Development* **143**, 635-647,
doi:10.1242/dev.131755 (2016).
73   Suh, N. & Blelloch, R. Small RNAs in early mammalian development: from gametes
to gastrulation. *Development* **138**, 1653-1661, doi:10.1242/dev.056234 (2011).
74   Pfisterer, U. *et al.* Identification of epilepsy-associated neuronal subtypes and gene
expression underlying epileptogenesis. *Nature communications* **11**, 5038,
doi:10.1038/s41467-020-18752-7 (2020).

75 Tobi, E. W. *et al.* DNA methylation signatures link prenatal famine exposure to
growth and metabolism. *Nature communications* **5**, 5592, doi:10.1038/ncomms6592
(2014).
76 Schulz, L. C. The Dutch Hunger Winter and the developmental origins of health and
disease. *Proceedings of the National Academy of Sciences of the United States of*
America **107**, 16757-16758, doi:10.1073/pnas.1012911107 (2010).
77 Jepsen, I. E. *et al.* Turning waiting time into treatment time: Weight reduction by a
lifestyle intervention programme for patients with obesity before fertility treatment.
Reproductive, Female and Child Health **2**, 133-142,
doi:https://doi.org/10.1002/rfc2.46 (2023).
78 Giroud, M. *et al.* Let-7i-5p represses brite adipocyte function in mice and humans.
Scientific reports **6**, 28613, doi:10.1038/srep28613 (2016).
79 Popov, L. D. Mitochondrial biogenesis: An update. *J Cell Mol Med* **24**, 4892-4899,
doi:10.1111/jcmm.15194 (2020).
80 Brandao, B. B. *et al.* Dynamic changes in DICER levels in adipose tissue control
metabolic adaptations to exercise. *Proceedings of the National Academy of Sciences*
of the United States of America **117**, 23932-23941, doi:10.1073/pnas.2011243117
(2020).
81 Sullivan, R. & Saez, F. Epididymosomes, prostasomes, and liposomes: their roles in
mammalian male reproductive physiology. *Reproduction* **146**, R21-35,
doi:10.1530/REP-13-0058 (2013).
82 Tan, K. *et al.* Downregulation of miR-199a-5p Disrupts the Developmental Potential
of In Vitro-Fertilized Mouse Blastocysts. *Biol Reprod* **95**, 54,
doi:10.1095/biolreprod.116.141051 (2016).
83 Chambers, M. C. *et al.* A cross-platform toolkit for mass spectrometry and
proteomics. *Nature biotechnology* **30**, 918-920, doi:10.1038/nbt.2377 (2012).
84 Kind, T. *et al.* LipidBlast in silico tandem mass spectrometry database for lipid
identification. *Nature methods* **10**, 755-758, doi:10.1038/nmeth.2551 (2013).
85 Luan, H., Ji, F., Chen, Y. & Cai, Z. statTarget: A streamlined tool for signal drift
correction and interpretations of quantitative mass spectrometry-based omics data.
Anal Chim Acta **1036**, 66-72, doi:10.1016/j.aca.2018.08.002 (2018).
86 Kleiner, D. E. *et al.* Design and validation of a histological scoring system for
nonalcoholic fatty liver disease. *Hepatology* **41**, 1313-1321, doi:10.1002/hep.20701
(2005).
87 Rappsilber, J. & Mann, M. Analysis of the topology of protein complexes using cross-
linking and mass spectrometry. *CSH Protoc* **2007**, pdb prot4594,
doi:10.1101/pdb.prot4594 (2007).
88 Tyanova, S., Mann, M. & Cox, J. MaxQuant for in-depth analysis of large SILAC
datasets. *Methods in molecular biology* **1188**, 351-364, doi:10.1007/978-1-4939-
1142-4_24 (2014).
89 Tyanova, S. *et al.* The Perseus computational platform for comprehensive analysis of
(prote)omics data. *Nature methods* **13**, 731-740, doi:10.1038/nmeth.3901 (2016).
90 Friedlander, M. R., Mackowiak, S. D., Li, N., Chen, W. & Rajewsky, N. miRDeep2
accurately identifies known and hundreds of novel microRNA genes in seven animal
clades. *Nucleic acids research* **40**, 37-52, doi:10.1093/nar/gkr688 (2012).
91 Chan, B. K. C. Data Analysis Using R Programming. *Adv Exp Med Biol* **1082**, 47-
122, doi:10.1007/978-3-319-93791-5_2 (2018).
92 Reimers, M. & Carey, V. J. Bioconductor: an open source framework for
bioinformatics and computational biology. *Methods Enzymol* **411**, 119-134,
doi:10.1016/S0076-6879(06)11008-3 (2006).
93 Frankish, A. *et al.* GENCODE reference annotation for the human and mouse
genomes. *Nucleic acids research* **47**, D766-D773, doi:10.1093/nar/gky955 (2019).
94 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
doi:10.1093/bioinformatics/bts635 (2013).

95 Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930,
doi:10.1093/bioinformatics/btt656 (2014).
96 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**,
139-140, doi:10.1093/bioinformatics/btp616 (2010).
97 Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: Precision weights unlock linear
model analysis tools for RNA-seq read counts. *Genome biology* **15**, R29,
doi:10.1186/gb-2014-15-2-r29 (2014).
98 Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics
data. *Innovation (Camb)* **2**, 100141, doi:10.1016/j.xinn.2021.100141 (2021).
99 Sud, M. *et al.* Metabolomics Workbench: An international repository for metabolomics
data and metadata, metabolite standards, protocols, tutorials and training, and
analysis tools. *Nucleic acids research* **44**, D463-470, doi:10.1093/nar/gkv1042
(2016).
1422

Figure 1: Paternal obesity elicits heritable impairments in glucose metabolism and adipose mitochondrial function

Figure 2: Paternal obesity dysregulates miRNA:mRNA gene networks spurring intergenerational mitochondrial dysfunction

Figure 3: Paternal obesity reversibly affects sperm sncRNomes and induces *let7 isoforms* in eWAT and sperm.

Figure 4: Zygotic delivery of *let-7d/e-5p* causes glucose intolerance in sired F$_{MI}$ mice and impairs embryonic mitochondria.

Figure 5: Adipocyte let-7 blunts mitochondrial function via a novel miR-let-7-DICER1 axis.

Figure S1: Paternal obesity and weight loss do not affect reproductive fitness and litter sizes

a

b

Figure S2: Paternal obesity causes heritable alterations in glucose metabolism in male but not in female mice

Figure S3: Effects of paternal obesity on F$_0$ and F$_1$ eWAT proteomes

Figure S4: Paternal obesity elicits alterations in circulating mitochondria-associated metabolites and lipids

Figure S5: Paternal weight loss corrects obesity-associated mitochondrial dysfunction in eWAT.

Figure S6: Paternal obesity does not affect F_1 adipose miRNA expression and meta-inflammation

Figure S7: Paternal obesity does not affect mitochondrial gene expression in F_0 and F_1 liver

Figure S8: Paternal obesity and weight loss do not affect liver morphology and lipid metabolism in F_1 progeny

Figure S9: Paternal obesity and weight loss cause only modest changes in spermatozooa mitochondrial tRNA fragments

Figure S10: Zygotic microinjection of obesity associated miR-let7 does not affect embryonic viability and implantation