1 ORIGINAL RESEARCH

2 O'Connor et al

3 Differential effects of conventional transcranial direct

4 current stimulation (tDCS) and high-definition

5 transcranial direct current stimulation (HD-tDCS) of

6 the cerebellum on offset analgesia

Niamh O'Connor[1,3*]

Hannah Ashe[1*]

Max Wragan[1,3*]

Ruairí O'Flaherty[1*]

Eoin Deevy-Gray[1]

Alice G. Witney [1,2,3]

[1] Trinity College Institute of Neuroscience, Trinity College Dublin, Ireland

[2] Trinity Centre for Biomedical Engineering, Trinity College Dublin, Ireland

[3] Discipline of Physiology, School of Medicine, Trinity College Dublin, Ireland

*These authors contributed equally to this work

Correspondence:

Department of Physiology,

2nd Floor Trinity Biomedical Sciences Institute,

152-160 Pearse Street,

Dublin D2

Tel +353 1 896 2080

Email : Dr Alice Witney awitney@tcd.ie

Abstract

Background : Offset analgesia (OA) describes the large decrease in perceived pain in

response to a minor decrease in applied painful thermal stimulus. Here non-invasive brain

stimulation (NIBS) is used to modulate the cerebellum, since the cerebellum is known to

signal sensory prediction errors and is implicated in pain processing.

Methods : An OA protocol individualized to heat pain threshold (HPT) was applied via TSA-II

(Medoc, Israel). NIBS interventions were applied prior to OA. Cathodal cerebellar transcranial

direct current stimulation (tDCS) and high-definition (4X1) transcranial direct current

stimulation (HD-tDCS) were applied to 46 healthy participants within a sham controlled

repeated measures design to examine whether diffuse or focal stimulation differentially

modulates OA.

Results : OA induced hypoalgesia was robust, with 90% of responses showing a drop in

perceived pain (δVAS) > 10 following the 1°C fall in temperature. This OA response was

augmented following a protocol with sham and focal cathodal cerebellar stimulation on four

OA parameters (OA $_{latency}$, VAS $_{minimum}$, VAS $_{mean}$ and VAS $_{2nd\ max}$) relative to pre-stimulation.

This effect was differential to the protocol with sham and conventional tDCS where two OA

metrics altered (OA $_{duration}$, VAS $_{2nd\ max}$).

Conclusion : OA enhancement via cathodal cerebellar NIBS may involve both a placebo

effect and sustaining a noxious sensory prediction error. Understanding how the cerebellum is

involved in OA could enhance therapies for pain patients.

Introduction

Offset analgesia (OA) is a pain modulation phenomenon where an individual experiences a disproportionately large decrease in pain sensation in response to a small decrease in experimentally applied painful thermal stimulus [14; 16]. The transient unexpected decrease in applied pain leads to the experience of analgesia and hypoalgesia in 70-90% of healthy participants[16]. This OA response is thought to be due endogenous pain regulation systems with research demonstrating that abnormal or absent OA responses in chronic pain patients [1; 34; 36; 45]. Therefore, the OA protocol has potential to provide valuable insights into the function of endogenous pain regulation systems. These pathways are obvious therapeutic targets, so there is clinical utility in understanding the mechanisms underlying the strong analgesic effect that occurs following the unexpected drop in nociceptive thermal stimuli.

The cerebellum is known to contribute to pain modulation, but the precise role in pain processing has not been established [21; 31; 42]. The cerebellum has been widely studied in computational neuroscience due to the potential to map its geometric structure to theoretical models of learning, where a mismatch between predicted and actual sensory signals can provide an error correction mechanism that drives adaptation[29]. Further this learning process can be mapped to the cellular architecture of the cerebellum[18]. The cerebellar cortex has a modular organization with repeated microzones with a repeated simple neuronal arrangement. In each microzone, climbing fibres from the inferior olive project to Purkinje cells which then output to cerebellar and vestibular nuclei. Frameworks for learning suggest that synaptic plasticity, particularly long-term depression (LTD) drive error correction and motor learning. These ideas have led to the concept of the cerebellum involvement in the learning and updating an 'internal model', with this framework having broader applications than motor control[52]. Offset analgesia could be explained via an internal model, with the mismatch between actual and expected nociceptive stimuli leading to a re-calibration of the pain perceived via the same error correction mechanism.

Non-invasive brain stimulation (NIBS) is a means of modulating neuronal activity and has

been found effective for pain modulation[3; 10; 11; 19; 22; 30]. Currently, the greatest

efficacy for pain relief is associated with high frequency repetitive transcranial magnetic

stimulation (rTMS) of primary motor cortex (M1) and anodal transcranial direct current

stimulation (tDCS) of primary motor cortex (M1) [12; 20; 24]. Anodal M1 HD-tDCS has been

found to modulate pain thresholds [50] with increased focality. These forms of M1 NIBS lead

to increased corticomotoneuronal and corticocortical excitability with long term potentiation

(LTP) like mechanisms proposed to underlie the synaptic plasticity. In contrast, cathodal

tDCS is thought to reduce corticomotoneuronal excitability and induce LTD like changes.

Within experimental studies of the cerebellum, both excitatory and inhibitory cerebellar

stimulation have been shown to lead to modulation of pain thresholds in animal studies,

dependent on the region stimulated. Further, the higher folding, increased neuronal density

and wider diversity of cell types of the cerebellum relative to M1 add complexity to designing

appropriate cerebellar NIBS protocols and may necessitate focal stimulation. Some evidence

has shown that cathodal tDCS may inhibit motor adaptation [17; 51]with the proposal that this

can be explained via an inhibition of the error correction learning. If applied to pain, inhibition

of updating the mismatch between actual and expected nociception could be predicted to

sustain OA induced analgesia. Therefore, this study applies inhibitory cathodal NIBS to the

cerebellum, comparing diffuse tDCS with the more focal HD-tDCS.

Material and methods

The study protocol was approved by the Faculty of Health Science Research Ethics

Committee, Trinity College Dublin, Ireland and conducted in accordance with the Declaration

of Helsinki (2013) with pre-registration on the Open Science Framework. Participant eligibility

was evaluated with a medical questionnaire, which excluded participants with neurological

and psychiatric conditions. Power calculations determined that at least 44 participants were

required. 46 participants (22 females, mean age 22.35 years) completed both experiments in

the study. A sham-controlled repeated measures design was used, with tDCS and HD-tDCS

completed in two sessions, separated by at least one week, to ensure no overlap in NIBS

effects between experimental sessions. The order of the two NIBS interventions was counter-

balanced between participants. Sham stimulation enables placebo NIBS effects on OA to be

examined. Sham stimulation occurred prior to active stimulation in all participants in each

session to avoid after-effects.

Offset analgesia protocol

A TSA-II Neurosensory Analyzer, Medoc, Israel [28] applied heat via the 16x16 mm peltier

thermode to the participants right forearm, secured with Velcro©. A computerized visual

analogue scale (CoVAS, Medoc, Israel) [27] continuously recorded participants' pain report at

9 Hz [55] with the displacement of a marker between two endpoints of 'no pain' and 'most

intense pain imaginable' corresponding to values between 0 and 100. Training on CoVAS

occurred prior to testing.

Heat pain threshold (HPT) were determined to individualize the subsequent OA protocol [2;

32; 40; 43; 44]. HPTs were determined by increasing the temperature of the thermode

applied to the right forearm at a rate of 1.5°C/s from a baseline of 32-35°C until pain was

reported. HPT trials were repeated three times, with 10 s rest intervals between trials.

The OA protocol comprised of an OA trial and a control trial, presented in random order. To

ensure data accuracy, 60 s of rest followed data collection trials in addition to slight thermode

repositioning on the forearm to prevent sensitization [54]. On an OA trial (Figure 1), three

sequential temperatures (T1-3) were utilised: the initial stimulus temperature (T1) is the

participant's HPT, ramped from the baseline temperature representing normal skin

temperature (32-35°C) at a rate of 1.5℃/s. T1 (HPT) was held constant for 7 - 10 seconds.

The duration of T1 varied to decrease participants' expectation of a temperature change at

temperature 2 (T2). The T2 phase elevates the temperature a single degree to HPT + 1℃ for

5 s, and the Temperature 3 (T3) falls to HPT for 30 s before returning to baseline skin

temperature at a rate of 6℃/s. (Figure 1). The transition from T2 to T3 is where OA is

experienced, the characteristic disproportionate decrease in pain sensation relative to the

associated temperature decrease. This OA protocol differs from some previous OA protocols

[14; 54] because the T3 phase was extended to characterize the duration of OA, previous

studies show ongoing OA at the termination of T3 [2; 14; 46].

Control trials in the OA protocol consisted of the same T1 temperature as the OA trials

(HPT), similarly ramped from baseline skin temperature at a rate of 1.5℃/s. However, in

contrast, T1 was sustained for 30s, before ramping down to baseline skin temperature at rate

of 6℃/s.

Transcranial direct current stimulation montages

Conventional cerebellar tDCS was applied using a neuroConn DC stimulator (NeuroConn

GmBH, Germany) with sponge electrodes (surface area 25 cm^2) soaked in 0.9% saline. The

stimulating electrode was positioned at the mid-point between the inion and the maximum

convexity of the right ear with the return electrode on the right buccinator[15; 17; 23].

Cerebellar HD-tDCS was applied using the neuroConn DC stimulator connected to a Soterix

4x1 HD-tDCS adaptor (Model 4x1-C2; Soterix Medical Inc., New York, USA). A 4X1

electrode montage using sintered Ag/AgCl electrodes (EasyCap, GmbH, Germany) filled with

conductive gel (Electro-Gel, Brainmaster Inc., USA) were secured with an electrode cap

(EasyCap, GmbH, Germany). The central electrode was placed at the mid-point of the inion

and the maximum convexity of the right ear, with the four return electrodes spaced 3 cm

radially around the central electrode. Impedance values were measured for all electrodes

prior to usage to ensure good contact quality [49; 50]. Active NIBS applied current of 2mA of

tDCS for 20 minutes for both NIBS protocols. Sham stimulation applied 2mA for 30s to elicit a

tingling sensation, with no current applied in remainder of the 20 minute protocol [13; 48].

Current flow modelling is critical to NIBS protocol design[9; 38]. The cerebellar tDCS and HD-

tDCS protocols were simulated (HD-Explore Version 2.1, Soterix Medical, New York, US) to

characterise the current distribution with the given electrode montage and current level of

2mA as applied experimentally. All modelling was performed on a 93 standard adult male

head model sourced from the Soterix model library. Electrode positioning was based on the

10/20 EEG system. For conventional tDCS modelling, O10, PO10, EX2, and EX6

represented the cathode sponge electrode and EX14, EX16, and EX18 represented the

anodal sponge electrode (Figure 2a). For HD-tDCS modelling, PO10 represented the

cathode and PO4, P10, EXZ, and EX4 represented the anodes (Figure 2b).

OA and control trial response characterization

OA and Control response metrics were extracted using a custom written MATLAB script

(Version 9.0, The MathWorks Inc., MA, USA). Control trial metrics were mean (VAS $_{mean}$),

minimum (VAS $_{min}$), maximum (VAS$_{max}$) and standard deviation (VAS$_{sd}$) of the VAS score

while HPT was applied over a 30s period (T1).

CoVAS profiles were averaged dependent on neurostimulation (pre-stimulation, sham, active

stimulation) from the onset T3 for 22 s to qualitatively compare the impact of NIBS on the OA

response, which is defined as the VAS ratings made during T3. The OA response metrics

used in the study (Table 1, Figure 1) included both magnitude (Figure 1a) and temporal

(Figure 1b) features with metrics selected to be consistent with previous studies, VAS $_{min}$ [2;

6; 34], δVAS [1; 6; 34] and VAS $_{mean}$ [6]. Further, in this study, a second maximum VAS was

included since the OA protocol extended the T3 phase. Some OA metrics could not be

extracted on all trials due to features of the OA response, for example there could be no VAS

$_{2nd\ max}$ or OA $_{rate\ up}$ metric if the OA continued throughout T3.

Statistical analysis

Statistical analysis was performed in SPSS (IBM SPSS Statistics, Version 27; IBM Corp).

Descriptive statistics were collected for all OA metrics across conditions (pre-stimulation OA;

sham stimulation OA; active stimulation OA). Within day ICC on OA metrics are consistently

high in previous research [35] and pilot work in the laboratory. Between day ICC comparing

OA metrics at pre-stimulation baseline on each session were moderate or low. Therefore, the

tDCS and HD-tDCS experiments were considered separately via two 3X1 repeated

measures ANOVA for both control and OA trial metrics. Mauchly's test of sphericity was

performed, and when sphericity could not be assumed the Huynh-Feldt estimate was applied.

Post-hoc pairwise comparisons were made with Bonferroni corrections for multiple

comparisons.

Results

On control trials where T1 was held at HPT for 30s, the VAS $_{mean}$ was 30.07 (SD 18.95), VAS

$_{max}$ of 42.31 (SD 22.41), VAS $_{min}$15.96 (SD 18.67) and VAS $_{sd}$ was 28.11 (SD 9.06). As

expected, there were no consistent periods of analgesia observed in this VAS response to

sustained HPT.

In contrast, the OA trials successfully consistently induced analgesia, with only 3% of trials

not showing a decrease in VAS at the T2-T3 transition. 90% of trials showed a fall greater

than 10 VAS, with most trials eliciting a much larger decrease in pain perception.

The fall of temperature in T3 to HPT gave hypoalgesia with a minimum VAS (VAS $_{min}$) at the

nadir of the OA response of 12.71 (SD 15.50). The mean δVAS in the current study was

slightly lower at 38.13 (SD, 22.39) relative to scores found by previous studies [1; 32; 33].

The T3 VAS $_{mean}$ 30.35 (SD 23.61) for T3 phase was lower than studies that do not

individualize OA protocols to HPTs [6]. The OA $_{rate\ down}$ at the T2-T3 transition a mean of 8.45

VAS/s (SD 8.45) was much steeper than OA $_{rate\ up}$, during T3, (1.85 VAS/s, SD 1.71),

revealing the rapidity of OA at the transition and the sustained hypoalgesia that demonstrates

the necessity of extending T3 in the protocol. Taken together the protocol used in this study

robustly generated OA and provided baseline values consistent with earlier studies in the

area [1; 26; 32].

Modelling cerebellar NIBS

Current flow in the conventional tDCS (Figure 2a) is more diffuse than HD-tDCS (Figure 2b).

Highest flow intensities with tDCS are near the active electrode, at the cerebellar cortical

level in the posterior lobe. Medial structures also receive strong current and spread to the

temporal lobe and frontal lobe. Current reached the upper and lower brainstem. For HD-

tDCS, current flow was more restricted, with high field intensities in the posterior and anterior

cerebellar lobe and posterior aspects of the temporal lobe. Very little current reaches the

contralateral cerebellar hemisphere or brainstem.

The effect of cerebellar NIBS on OA

There were no significant differences in the control trial metrics (VAS $_{mean}$, VAS $_{min}$, VAS$_{max}$ and VAS$_{sd}$) dependent on stimulation for either tDCS or HD-tDCS. However, there was a differential impact of NIBS on the OA response (Figure 3).

The average impact of active conventional cerebellar tDCS stimulation relative to pre-stimulation and sham can be seen in Figure (3a).

The effect of sham conventional tDCS and cerebellar cathodal tDCS on OA

The stimulation protocol led to two significant changes in OA metrics with moderate effect sizes. The alteration in OA $_{duration}$ was found to be significant $F(2,86)=4.35$, p =.037, partial η^2 = .07. Post-hoc pairwise comparisons with a Bonferroni adjustment indicated there was no significance difference between OA $_{duration}$ at baseline and following sham stimulation, p=.18. However, OA $_{duration}$ was significantly longer after active tDCS stimulation (11.17 s, SD 7.97) relative to pre-stimulation (8.11 s, SD 6.75), p=.048. Interestingly, there was also no significant difference between OA $_{duration}$ after sham and after active neurostimulation, p=1. OA $_{duration}$ represents the length of maximal OA; from the T2-T3 transition to the point at which pain rating begins to rise again (Figure 1b, Table 1, metric 8).

Additionally, there was an effect of stimulation of VAS $_{2nd\ max}$, $F(2,90)=5.40$,p=.006, partial η^2 = .11. Post-hoc pairwise comparisons with a Bonferroni adjustment showed there was a significant difference between mean VAS $_{2nd\ max}$ at baseline (37.35, SD 26.94) and following sham stimulation, (29.75, SD 25.32) p=.009 additionally VAS $_{2nd\ max}$ was significantly lower after active tDCS stimulation, (29.52, SD 25.33), p=.028. However, the reduction in VAS $_{2nd\ max}$ also occurred after sham stimulation, with no significant difference between VAS $_{2nd\ max}$ and after active stimulation, p=1, suggestive of a placebo effect. Statistical analysis revealed there was no effect of cerebellar tDCS neurostimulation on δVAS, VAS $_{min}$, OA $_{rate\ down}$, VAS $_{mean}$, OA $_{rate\ up}$ or OA l$_{atency}$ metrics of the OA response.

The effect of sham HD-tDCS and cathodal cerebellar HD-tDCS on OA

Qualitatively, the effect of active cathodal cerebellar HD-tDCS relative to sham is to enhance the OA effect (Figure 3b). Statistical analysis revealed differences on four OA metrics with medium effect sizes. The effect of neurostimulation on δVAS, that is the drop in VAS at the T2-T3 transition was found to be significant $F(1.64,76.09)=3.72$, $p =.036$, partial $\eta^2 = .08$. Post-hoc pairwise comparisons with a Bonferroni adjustment indicated there was no significance difference between mean δVAS at baseline (32.70, SD 22.63) and following sham stimulation, (38.02, SD 23.51), $p=.46$. However, δVAS was significantly greater after active stimulation, (39.80, SD=21.79), $p=.007$. There was also no significant difference between δVAS after sham and after active neurostimulation, $p=.96$.

The effect of neurostimulation on VAS $_{rate\ down}$, that is the rate of OA-induced hypoalgesia, was found to be significant $F(1.70,71.18)=3.36$, $p =.048$, partial $\eta^2 = .07$. Post-hoc pairwise comparisons with a Bonferroni adjustment indicated there was no significance difference between mean VAS $_{rate\ down}$ at baseline (6.59 VAS/s, SD 4.75) and following sham stimulation, (8.91 VAS/s, SD=8.38), $p=.55$. However, VAS $_{rate\ down}$ was significantly faster after active tDCS stimulation, (8.31 VAS/s, SD 5.37) $p=.021$. There was also no significant difference between VAS $_{min}$ after sham and after active neurostimulation, $p=.86$.

The effect of neurostimulation on VAS $_{min}$, the VAS at the nadir of the OA response was found to be significant $F(2,90)=4.96$, $p =.01$, partial $\eta^2 = .1$. Post-hoc pairwise comparisons with a Bonferroni adjustment indicated there was no significance difference between mean VAS_{min} at baseline (16.17, SD=20.25) and following sham stimulation, (10.08, SD 11.62), $p=.12$. However, mean VAS_{min} was significantly reduced after active tDCS stimulation, (9.79, SD 11.62), $p=.018$. There was also no significant difference between VAS_{min} after sham and after active neurostimulation, $p=1$.

There was an effect of stimulation of VAS $_{mean}$, $F(2,90)=4.91$, $p =.009$, partial $\eta^2 = .98$. Post-hoc pairwise comparisons with a Bonferroni adjustment indicated there was no significance difference between mean VAS_{mean} at baseline (32.04, SD 23.81) and following sham stimulation, $p=.055$. However, VAS $_{mean}$ was significantly lower after active tDCS stimulation,

(26.02, SD 21.75) p=.023. There was also no significant difference between VAS $_{mean}$ after

sham (27.79, SD 18.99), and after active neurostimulation, p=1.

The effect of neurostimulation on $VAS_{2nd\ max}$ was found to be significant F(2,90)=3.56, p

=.033, partial η^2 = .07 indicating pain does not rise as much even though OA is reduced

relative to the start of T3. Post-hoc pairwise comparisons with a Bonferroni adjustment

indicated there was no significance difference between mean $VAS_{2nd\ max}$ at baseline (32.04,

SD 23.81) and following sham stimulation, (27.29, SD 18.99), p=.19.   However, $VAS_{2nd\ max}$

was significantly reduced after active tDCS stimulation, (26.02, SD 21.75), p=.038. There was

also no significant difference between $VAS_{2nd\ max}$ after sham and after active neurostimulation,

p=1.

There were no effect of the NIBS protocol on OA $_{duration}$, OA $_{latency}$ or OA $_{rate\ up}$.

282 Discussion

Offset analgesia, the disproportionate decrease in perceived pain that follows a slight

decrease in applied nociceptive stimuli, is a robust experimental pain protocol[14]. One

previously untested explanation for offset analgesia is that it is experienced due to sensory

error signaling in the cerebellum. Error signaling in the cerebellum drives sensorimotor

adaptation and motor learning in altering environments[52]. If this concept were applied to

pain, a reduction in adaptation might prolong the analgesia evoked by the transient sensory

mismatch in an OA trial. To address this, the study applied inhibitory NIBS to the cerebellum,

to examine whether cathodal tDCS may reduce adaptation to the sensory mismatch implicit

within the OA protocol and thus sustain analgesia. This prediction of the study was

supported, yet also it was further revealed that placebo effects could contribute to modulation

of OA.

The study first designed an individualized offset analgesia protocol based on prior

studies[16]. Consistent with earlier OA studies [14], the protocol provided a robust

methodology for inducing hypoalgesia, with participants consistently experiencing a profound

reduction in the magnitude of pain experienced during the protocol. One aim of this study

was quantifying the OA temporal pain profile with metrics that could characterize both

magnitude and duration of the experienced pain. The increased duration of an OA trial in this

study protocol is an important modification to earlier protocols as it assists differentiation and

characterization of OA pain profiles; the essential basis for determining the effects of NIBS on

OA.

Focal, HD-tDCS was found to augment the effects of OA pain profiles in four of the metrics

with medium effect sizes with conventional cerebellar tDCS augmenting the OA pain profile

on two metrics relative to pre-stimulation. However, in this study the interpretation of NIBS

effects on the OA response is likely to summative to the occurrence of repeated OA protocols

and perhaps also to sham NIBS providing a placebo effect. Qualitatively there appears to be

a trend towards increased OA after sham (Figure 3), and one metric in the conventional

cerebellar tDCS experiment showed a significant difference post sham, suggestive that a

placebo effect is present. However, no other OA metrics post-sham reach significance in

either experiment. Therefore, it is possible that the effect of repeated OA with sham NIBS

primes the OA response, with subsequent active NIBS intervention augmenting the effect,

producing the overall significant changes to the OA pain profiles. Previous studies have

investigated the impact of repeated OA, with mixed findings[5; 16]. Importantly, the

differential impact of the two active NIBS interventions on OA is observable both qualitatively

and statistically, indicating that interpretation must include consideration of current flow and

the focality of current applied and not be solely a placebo effect.

Analgesic enhancement in OA can be explained by several different pathways, and

cerebellar stimulation via NIBS can contribute to this enhancement. The exact mechanisms

of NIBS on the cerebellum are less well established than cortical regions [48], partly due to

the increased diversity of cell types in cerebellar cortex, though it is thought to modulate

Purkinje cells. There are multiple possible consequences to Purkinje cell modulation, since

these are the sole output cells from cerebellar cortex. First, cathodal cerebellar tDCS may

decrease inhibitory effects of Purkinje cells and so increases deep cerebellar nuclei activity,

to indirectly reduce inhibition of M1 [12]. Numerous studies link analgesia with primary motor

cortex excitability and the cerebellum has a known role in modulating corticomotoneuronal

activity[37]. This pathway forms the cerebello-thalamo-cortical pathway, a major output

pathway of the cerebellum. Alternatively, increased caudate putamen activity is associated

with cathodal and anodal cerebellar tDCS [24]. The caudate is a known region for pain

suppression [53] and is critically involved in the reward pathways. The hypoalgesia

experienced after the transient decrease in supra-threshold pain, that is characteristic of the

OA protocol, is known to activate reward pathways. Additionally, cerebellar stimulation may

impact on OA via connectivity with known pain modulation pathways including the PAG and

dorsolateral prefrontal cortex (DLPFC)[4].

This study showed in a healthy young population, cathodal cerebellar NIBS modulated the

OA response relative to pre-stimulation consistent with disruption of cerebellar adaptation.

This is notable since the OA response in a healthy population such as the one in this study

can be subject to ceiling effects. Future work would apply the study protocol to an elderly and

chronic pain patient population, both of whom have deficits in OA. However, studies in

healthy, young adults are first necessary to ensure a robust protocol, minimizing individual

differences that would impact on the OA response. The differences in effects due to focality

of stimulation are perhaps unsurprising given that animal studies show both pro and anti-

nociceptive effects of the cerebellum[7; 41] and due to the relatively small target regions

[47]. This suggests that future work should use high-definition montages, alongside current

flow modeling. Cerebellar NIBS protocols and electrode montages are developing [8; 25; 48]

but there is increasing evidence that the choice of optimal electrode montage and 'dosage' is

critical for an optimal effect [39].

Overall OA is a robustly observed form of hypoalgesia and better understanding of the

underlying mechanism could open therapeutic options. This study supports further

investigation of cerebellar involvement in OA.

Author contributions

354 NOC and AW designed the experimental protocol with EDG involved in pilot testing; NOC,

355 HA, ROF collected the experimental data, MW and AW collated the data, MW and AW

356 performed statistical and data analysis. NOC, HA, ROF, EDG contributed to initial drafts. MW

357 and AW wrote the manuscript.

Disclosure

The author reports no conflicts of interest in this work. A proportion of this work was presented as an abstract at EFIC, Valencia 2019

References

[1] Alter BJ, Aung MS, Strigo IA, Fields HL. Onset hyperanalgesia and offset analgesia: transient increases or decrease of noxious thermal stimulus intensity robustly modulate subsequent perceived pain intensity. PLoS One 2020;15(e0231124).

[2] Alter BJ, Santosa H, Nguyen QH, Huppert TJ, Wasan AD. Offset analgesia is associated with opposing modulation of medial versus dorsolateral prefrontal cortex activations: a functional near infrared spectroscopy study. Molecular Pain 2022;18.

[3] Antal A, Brepohl N, Poreisz C, Boros K, Csifcsak G, Paulus W. Transcranial direct current stimulation over somatosensory cortex decreases experimentally induced acute pain perception. Clinical Journal of Pain 2008;24:56-63.

[4] Cacciola A, Bertino S, Basile GA, Di Maruo D, Calamuneri A, Chillemi G, Duca A, Bruschetta D, Flace P, Favalaro A, Calabro RS, Anastasi G, Milardi D. Mapping the structural connectivity between the periaqueductal gray and the cerebellum in humans. Brain Structure Function 2019;224:2153-2165.

[5] Derbyshire SWG, Osborn J. Enhancement of offset analgesia during sequential testing. European Journal of Pain 2008;12:980-989.

[6] Derbyshire SWG, Osborn J. Offset analgesia is mediated by activation in the region of the periaqueductal grey and rostral ventromedial medulla. Neuroimage 2009;47:1002-1006.

[7] Dey PK, Ray AK. Anterior cerebellum as a site for morphine analgesia and post-stimulation analgesia. Indian J Physiol Pharmacol;26:3-12.

[8] Ferrucci R, Cortese F, Priori A. Cerebellar tDCS: how to do it. The Cerebellum 2015;14:27-30.

[9] Fiocchi S, Chiaramello E, Gazzola V, Suttrup J, Ravazzani P, Parazzini M. Assessment of the capability to target cerebellar sub-regions with high-definition transcranial direct current stimulation (HD-tDCS) over the cerebellum. IEEE 3rd international forum on research and technologies for society and industry (RTSI) 2017:1-6.

[10] Fregni F, Boggio PS, Lima MC, Ferreira MJL, Wagner T, Rigonatti SP, Castro AW, Souza DR, Riberto M, Freedman SD, Nitsche MA, Pascual-Leone A. A sham-controlled, phase II trial of transcranial direct current stimulation for the treatment of central pain in traumatic spinal cord injury. Pain 2006;122:197-209.

[11] Fregni F, Gimenes R, Valle AC, Ferreira MJL, Rocha RR, Natalle L, Bravo R, Rigonatti SP, Freedman SD, Nitsche MA, Pascual-Leone A, Boggio PS. A randomised, sham-controlled, proof-of-principle study of transcranial direct current stimulation for the treatment of pain in fibromyalgia. Arthritis and Rheumatism 2006;54:3988-3998.

[12] Galea JM, Jayaram G, Ajagbe L, Celnik P. Modulation of cerebellar excitability by polarity-specific noninvasive direct current stimulation. J Neurosci 2009;29:9115-9122.

[13] Gandiga PC, Hummel FC, Cohen LG. Transcranial DC stimulation (tDCS): a tool for double blind sham-controlled clinical studies in brain stimulation. Clin Neurophysiol 2006;117:845-850.

[14] Grill JD, Coghill RC. Transient analgesia evoked by noxious stimulus offset. J Neurophysiol 2002;87:2205-2208.

[15] Hamada M, Strigaro G, Murase N, Sadnicka A, Galea JM, Edwards MJ, Rothwell JC. Cerebellar modulation of human associative plasticity. Journal of Physiology 2012;590:2365-2374.

[16] Hermans L, Calders P, Van Oosterwijck J, Verschelde E, Bertel E, Meeus M. An
overview of offset analgesia and the comparison with conditioned pain modulation: a
systematic literature review. Pain Physician 2016;19:307-326.
[17] Herzfeld DJ, Pastor D, Haith AM, Rossetti Y, Shadmehr R, O'Shea J. Contributions of the
cerebellum and the motor cortex to acquisition and retention of motor memories.
Neuroimage 2014;98:147-158.
[18] Ito M. Neurophysiological aspect of the cerebellar control system. International Journal of
Neurology 1970;7:162-176.
[19] Kong Q, Li T, Reddy S, Hodges S, Kong J. Brain stimulation targets for chronic pain:
insights from meta-analysis, functional connectivity and literature review.
Neurotherapeutics 2024;21(e00297).
[20] Lefaucher JP, Antal A, Ahdab R, Ciampi de Andrade D, Fregni F, Khedr EM, Nitsche M,
Paulus W. The use of repetitive transcranial magnetic stimulation (rTMS) and
transcranial direct current stimulation (tDCS) to relieve pain. Brain Stimulation
2008;1:337-344.
[21] Li CN, Keay KA, Henderson LA, Mychasiuk R. Re-examining the mysterious role of the
cerebellum in pain. J Neurosci 2024;44(17):1-18.
[22] Lima MC, Fregni F. Motor cortex stimulation for chronic pain: systematic review and
meta-analysis of the literature. Neurology 2008;70:2329-2337.
[23] Macher K, Bohringer A, Villringer A, Pleger B. Cerebellar-parietal connections underpin
phonological storage. Journal of Neuroscience 2014;34:5029-5037.
[24] Maldonado T, Jackson TB, Bernard JA. Anodal cerebellar stimulation increases cortical
activation: evidence for cerebellar scaffolding of cortical processing. Human Brain
Mapping 2023;44(4):1666-1682.
[25] Manto M, Argyropolos GPD, Bocci T, Celnik PA, Corben LA, Guidetti M, Koch G, Priori A,
Rothwell JC, Sadnicka A, Spampinato D, Ugawa Y, Wessel MJ, Ferrucci R.
Consensus paper: novel directions and next steps of non-invasive brain stimulation of
the cerebellum in health and disease. The Cerebellum 2022;21:1092-1122.
[26] Martucci KT, Eisenach JC, Tong C, Coghill RC. Opiod-independent mechanisms
supporting offset analgesia and temporal sharpening of nociceptive information. Pain
2012;153:1232-1243.
[27] Medoc L. CoVAS (apparatus and software), 1996.
[28] Medoc L. TSA-II Neurosensory Analyzer (apparatus and software), 1999.
[29] Miall RC, Wolpert DM. Forward models for physiological motor control. Neural Netw
1996;9(8):1265-1279.
[30] Mordillo Mateos L, Dileone M, Soto-Leon V, Brocalero-Camacho A, Perez-Borrego YA,
Onate-Figuerez A, Aguilar JAM, Oliviero A. Effects of transcranial direct current
stimulation on temperature and pain perception. Scientific Reports 2017;7.
[31] Moulton EA, Schmahmann JD, Becerra L, Borsook D. The cerebellum and pain: passive
integrator or active participator? Brain Res Rev 2010;65(1):14-27.
[32] Naugle KM, Cruz-Almeida Y, Fillingim RB, Riley JL. Offset analgesia is reduced in older
adults. Pain 2013;154:2381-2387.
[33] Niesters M, Dahan A, Swartjes M, Noppers I, Fillingim RB, Aarts L, Sarton EY. Effect of
ketamine on endogenous pain modulation in healthy volunteers. Pain
2011;152(3):656-663.
[34] Niesters M, Hoitsma E, Sarton E, Aarts L, Dahan A. Offset analgesia in neuropathic pain
patients and effect of treatment with morphine and ketamine. Anesthesiology
2011;115:1063-1071.
[35] Nilsson M, Nissen TD, Graversen C, Gazerani P, Drewes AM, Brock C. Offset analgesia:
a reproducibility study. Scandinavian Journal of Pain 2012;3:192.
[36] Oudejans LCJ, Smit JH, van Velzen M, Dahan A, Niesters M. The influence of offset
analgesia on the onset and offset of pain in patients with fibromyalgia. Pain
2015;156:2521-2527.
[37] Oulad Ben Taib N, Manto M. Trains of epidural DC stimulation of the cerebellum tune
corticomotor excitability. Neural Plasticity 2013:1-12.
[38] Parazzini M, Rossi E, Ferrucci R, Fiocchi S, Liorni I, Priori A, Ravazzani P.
Computational model of cerebellar transcranial direct current stimulation. Conf Proc
Annu Int Conf IEEE Eng Med Biol Soc 2013;IEEE Eng Med Biol Soc Annu Conf:237-
240.

[39] Peterchev AV, Wagner TA, Miranda PC, Nitsche MA, Paulus W, Lisanby SH, Pascual-
Leone A, Bikson M. Fundamentals of transcranial electric and magnetic stimulation
dose: definition, selection and reporting practices. Brain Stimulation 2012;5(4):435-
453.
[40] Shulman J, Zurakowski D, Keysor J, Jervis K, Sethna NF. Offset analgesia identifies
impaired endogenous pain modulation in pediatric chronic pain disorders. Pain
2020;161(2852).
[41] Siegel P, Wepsic JG. Alteration of nociception by stimulation of cerebellar structures in
the monkey. Physiol Behav 1974;13:189-194.
[42] Strick PL, Dum RP, Fiez JA. Cerebellum and nonmotor function. Annual Review of
Neuroscience 2009;32:413-434.
[43] Szikszay TM, Adamczyk WM, Carvalho GF, May A, Luedtke K. Offset analgesia:
somatotopic endogenous pain modulation in migraine. Pain 2020;161(3):557-564.
[44] Szikszay TM, Adamczyk WM, Carvalho GF, May A, Luedtke K. Offset analgesia is
increased intra-orally. Journal of Oral Rehabilitation 2022;49(10):993-1001.
[45] Szikszay TM, Adamczyk WM, Luedtke K. The magnitude of offset analgesia as a
measure of endogenous pain modulation in healthy participants and patients with
chronic pain: a systematic review and meta-analysis. The Clinical Journal of Pain
2019;35:189-204.
[46] Szikszay TM, Melz N, von Glasenapp B, Adamczyk WM, Luedtke K. Effects of
stimulation area and temperature rates on offset analgesia. Pain Reports
2022;7(e1043).
[47] Tomlinson SP, Davis NJ, Bracewell RM. Brain stimulation studies of non-motor cerebellar
function: A systematic review. Eurosci biobehav Rev 2013;37(5):766-789.
[48] Van Dun K, Bodranghien FCAA, Marien P, Manto MU. tDCS of the cerebellum: where do
we stand in 2016? Technical issues and critical review of the literature Frontiers in
Human Neuroscience 2016;10(199).
[49] Villamar MF, Volz MS, Bikson M, Datta A, Dasilva AF, Fregni F. Techniques and
considerations in the use of 4X1 ring high-definition transcranial direct current
stimulation (HD-tDCS). J Vis Exp 2013;e50309.
[50] Villamar MF, Wivatvongvana P, Patumanond J, Bikson M, Truong DQ, Datta A, Fregni F.
Focal modulation of the primary motor cortex in fibromyalgia using 4X1 ring high
definition transcranial direct current stimulation (HD-tDCS): immediate and delayed
analgesic effects of cathodal and anodal stimulation. Journal of Pain 2013;14:371-
383.
[51] Weightman M, Brittain J-S, Miall RC. Direct and indirect effects of cathodal cerebellar
TDCS on visuomotor adaption of hand and arm movements. Scientific Reports
2021;11(4464).
[52] Wolpert DM, Miall RC, Kawato M. Internal models in the cerebellum. TRENDS COGN
SCI 1998;2(9):338-347.
[53] Wunderlich AP, Klug R, Stuber G, Landwehrmeyer B, Weber F, Feund W. Caudate
nucleus and insular activation during a pain suppression paradigm comparing thermal
and electrical stimulation The Open Neuroimaging journal 2011;5:1-8.
[54] Yelle MD, Rogers JM, Coghill RC. Offset analgesia: a temporal contrast mechanisms for
nociceptive information. Pain 2008;134:174-186.
[55] Zhang K-L, Yuan H, Wu F-F, Pu X-Y, Liu B-Z, Li Z, Liu H, Yang Y, Wang Y-Y. Analgesia
effect of noninvasive brain stimulation for neuropathic pain patients: a systematic
review. Pain Therapy 2021;10:315-332.
514

Figure Legends

Figure 1: Offset analgesia protocol showing extracted OA metric definitions for magnitude (A) 1: VAS_{min}, 2: VAS_{mean}, 3: δVAS_{OA} and temporal (B) 5: OA rate $_{down}$, 6: OA $_{rate\ up}$, 7: OA $_{latency}$ and 8: 0A $_{duration}$ of the OA response. See Table 1 for OA metric definitions.

Figure 2: Current flow modelling: 3D representation of current flow (upper panel), current flow through coronal, sagittal and axial slices (lower panel) A) Cerebellar tDCS montage: High intensity in the cerebellum, temporal lobe, frontal lobe and brainstem. B) Cerebellar HD-tDCS montage restricted current to the right cerebellar hemisphere and the posterior occipital lobe, with peak field intensity in the posterior cerebellum.

Figure 3: Average of OA response during T3 period dependent on cerebellar NIBS stimulation for a) tDCS protocol and b) HD-tDCS protocol. Pre-stimulation (red); Sham-stimulation (blue); Active-stimulation (green). Shading represents standard error of the mean.

Table Legends

Table 1: OA response metric definitions for magnitude and temporal features (see Figure 1 for graphical representation)

Table 2: OA pain profile metric means

Metric Label	Offset Analgesia Metric	Description
1	VAS $_{min}$	Lowest reported VAS in t3
2	VAS $_{mean}$	Mean VAS in t3
3	δVAS $_{OA}$	Magnitude of OA: the change in VAS at t2-t3
4	VAS $_{2nd\ Max}$	Maximum VAS in t3 following hypoanalgesia
5	OA $_{rate\ down}$	Rate of decrease in VAS (δVAS) due to OA
6	OA $_{rate\ up}$	Rate of increase from VAS $_{min}$ to VAS $_{2nd\ Max}$
7	OA $_{latency}$	Time taken for maximum VAS score at the start of t3 to drop to VAS $_{min}$ (the OA induced hypoanalgesia)
8	OA $_{duration}$	Time taken for VAS $_{min}$ to rise again following hypoanalgesia

Table 1

stimulation	condition	VASmin	VASmean	δVAS	VAS (2nd max)	OA Rate down	OA Rate Up	OA Latency	OA duration
tDCS	Pre	14.74 (16.67)	26.77 (20.62)	32.70 (20.26)	27.79 (18.98)	6.59 (4.75)	1.41 (1.40)	6.09 (3.31)	8.11 (6.75)
	Sham	10.08 (11.71)	23.51 (17.62)	40.19 (23.76)	29.52 (25.33)	9.80 (10.07)	2.10 (1.78)	5.93 (3.34)	10.48 (8.59)
	Active	14.29 (16.37)	25.23 (18.17)	40.06 (22.63)	29.75* (23.80)	9.08 (9.81)	1.96 (1.45)	6.20 (3.50)	11.17* (7.97)
HD-tDCS	Pre	16.17 (20.25)	28.77 (19.59)	37.80 (22.40)	37.35 (26.94)	8.04 (7.65)	1.87 (1.76)	6.23 (3.35)	9.80 (6.75)
	Sham	11.35 (14.56)	21.49 (13.49)	38.02 (23.51)	27.79 (18.99)	9.80 (10.08)	1.81 (1.87)	5.53 (2.97)	8.90 (6.84)
	Active	9.79* (11.62)	20.91* (15.19)	39.75 (21.79)	26.02* (21.7)	8.31* (5.37)	1.96 (1.90)	5.69 (2.92)	10.62 (7.69)

Table 2

Figure 1

A

tDCS

Coronal Slice

Saggital Slice

Axial Slice

L   R

F   B

L   R

Field Intensity V/m

0.15

B

HD-tDCS

Coronal Slice

Saggital Slice

Axial Slice

L   R

F   B

L   R

Field Intensity V/m

0.15

Differential effect of neurostimulation on the OA response