1   ORIGINAL RESEARCH

2   O'Connor et al

3   Differential effects of conventional transcranial direct

4   current stimulation (tDCS) and high-definition

5   transcranial direct current stimulation (HD-tDCS) of

6   the cerebellum on offset analgesia

Niamh O'Connor[1,3*]

Hannah Ashe[1*]

Max Wragan[1,3*]

Ruairí O'Flaherty[1*]

Eoin Deevy-Gray[1]

Alice G. Witney [1,2,3]

[1] Trinity College Institute of Neuroscience, Trinity College Dublin, Ireland

[2]Trinity Centre for Biomedical Engineering, Trinity College Dublin, Ireland

[3]Discipline of Physiology, School of Medicine, Trinity College Dublin, Ireland

*These authors contributed equally to this work

Correspondence:

Department of Physiology,

2nd Floor Trinity Biomedical Sciences Institute,

152-160 Pearse Street,

Dublin D2

Tel +353 1 896 2080

Email : Dr Alice Witney awitney@tcd.ie

Abstract

Background : Endogenous analgesic systems in the brain modulate pain perception. Offset analgesia (OA) describes the large decrease in perceived pain in response to a minor decrease in applied painful thermal stimulus. Here non-invasive brain stimulation (NIBS) of the cerebellum is used to probe OA.

Methods : An OA protocol individualized to heat pain threshold (HPT) was applied via TSA-II (Medoc, Israel). Heat was ramped to HPT, with a transient temperature increase of HPT+1°C. NIBS interventions were applied prior to OA in 46 participants within a sham controlled repeated measures design. Cathodal cerebellar transcranial direct current stimulation (tDCS) and high-definition (4X1) transcranial direct current stimulation (HD-tDCS) were applied in separate experimental sessions to examine whether diffuse (tDCS) or focal (HD-tDCS) stimulation differentially modulates OA.

Results : OA induced hypoanalgesia was robust, with 90% of responses showing a substantial drop in perceived pain (δVAS) following the 1°C fall in temperature, with an average VAS decrease of 38 in response to the 1°C fall in temperature. Cathodal cerebellar HD-tDCS enhances the analgesic impact of OA on four OA parameters (OA $_{latency}$, VAS $_{minimum}$, VAS $_{mean}$ and VAS $_{2nd\ max}$) relative to pre-stimulation. Conventional tDCS modulates two OA metrics relative to pre-stimulation (OA $_{duration}$, VAS $_{2nd\ max}$) with an increase in OA $_{duration}$ following sham tDCS.

Conclusion : There is a differential influence of conventional and high-definition cerebellar NIBS on OA. This is suggestive of cerebellar modulation of OA and highlights the importance of electrode montage in delineating the influence of the cerebellum in pain processing.

Introduction

Endogenous pain modulatory pathways, that alter the pain experienced in response to nociceptive stimuli, are obvious therapeutic targets. However, these pathways are not fully elucidated. Offset analgesia (OA) is a pain modulation phenomenon where an individual experiences a disproportionately large decrease in pain sensation in response to a small decrease in experimentally applied thermal stimulus [20]. The hypoanalgesia evoked by OA experimental protocols has potential to provide valuable insights into the function of endogenous pain regulation systems. Previous studies have demonstrated abnormal OA responses in chronic pain patients relative to healthy participants, including neuropathic pain and osteoarthritis [1; 41; 43; 57] thus validating the potential clinical utility of this protocol. This study aims to further characterize the OA response and examine whether non-invasive brain stimulation (NIBS) of the cerebellum modulates these pathways.

The cerebellum contributes to pain modulation, but the precise role has not been elucidated [27; 38; 54]. fMRI pain studies report cerebellar activity in response to nociceptive stimuli, with cerebellar lobules IV-VI and crus I consistently showing increased activity during nociceptive stimulation [12; 36]. Thermal nociceptive stimuli increase activity in cerebellar lobules III-VIII [4; 37; 38]. Injury typically leads to increases in cerebellar activity ipsilateral to injury, whereas thermal and chemical nociception can cause bilateral increases in cerebellar activity [24; 27; 47]. Cerebellar lesions in humans are associated with increased pain sensation in response to mechanical and thermal stimulus, suggesting the cerebellum modulates pain via connectivity with sensorimotor, limbic, executive and reward regions [48]. A NIBS technique, repetitive transcranial magnetic stimulation (rTMS) of the cerebellum has been found to modulate pain thresholds with mechanisms thought to be related to peripheral nerve effects rather than the pathway to motor cortex [68]. Animal models show evidence of both anti-nociceptive and pro-nociceptive role of the cerebellum [49; 50] on pain thresholds that is thought to be due to the involvement of the anterior and posterior cerebellar cortex respectively.

Imaging studies show the cerebellum increases in activity during an OA protocol relative to tonic pain, with close correlation demonstrated with the periacqueductal gray (PAG) [22]. The

cerebellum's connectivity between the cerebellum and PAG are highly indicative of the

cerebellum's important role in mediating pain perception [38]. Further, patients with cerebellar

lesions demonstrated poor OA responses relative to age-matched healthy participants[48].

NIBS is effective for pain modulation[3; 16; 17; 25; 28; 35]. Currently, the greatest efficacy is

associated with high frequency repetitive transcranial magnetic stimulation (rTMS) of primary

motor cortex (M1) and anodal transcranial direct current stimulation (tDCS) of primary motor

cortex (M1) [18; 26; 30]. Anodal M1 HD-tDCS has been found to modulate pain thresholds

[62] with increased focality. The cerebellum is currently a relatively unexplored target for pain

modulation. Cerebellar NIBS protocols and electrode montages are developing [14; 31; 60]

but there is increasing evidence that the choice of optimal electrode montage and 'dosage' is

critical for an optimal effect [46]. Cerebellar NIBS may aid understanding of endogenous pain

modulation [25; 63]. A recent study has suggested that anodal and cathodal tDCS enhance

and inhibit endogenous pain pathways [53] respectively with small effect sizes. Given the

substantial evidence behind cerebellar involvement in OA, further research into cerebellar

NIBS protocols is appropriate to further understand this finding, as consideration of

stimulation polarity effects should be treated cautiously[7]. Folding of the brain region

underlying stimulation will likely result in the direction of current flow on neurons being difficult

to predict [3] and perhaps distort or diminish effect. This presents a particular challenge to

NIBS of the cerebellum, due to its complex cytoarchitecture [60] . Additionally the cerebellum

contains the highest density of neurons in the brain, making interpretation of diffuse current

application as occurs in conventional tDCS particularly challenging [64]. Therefore, cerebellar

NIBS may benefit from the more focused HD-tDCS method for optimizing behavioural effects.

This study compares two forms of transcranial direct current stimulation (tDCS) of the

cerebellum on OA, conventional tDCS and the more focal HD-tDCS on the OA response.

Material and methods

106 The study protocol was approved by the Faculty of Health Science Research Ethics

107 Committee, Trinity College Dublin, Ireland and conducted in accordance with the Declaration

108 of Helsinki (2013). Participant eligibility was evaluated with a medical questionnaire, which

109 excluded participants with neurological and psychiatric conditions. 46 participants (22

females, mean age 22.35 years) completed all experiments in the study. A sham-controlled repeated measures design was used, with tDCS and HD-tDCS completed in two sessions, separated by at least one week, to ensure no overlap in NIBS effects between experimental sessions.

Offset analgesia protocol

A TSA-II Neurosensory Analyzer, Medoc, Israel [34] applied heat via the 16x16 mm peltier thermode to the participants right forearm, secured with Velcro©. A computerized visual analogue scale (CoVAS, Medoc, Israel) [33] continuously recorded participants' pain report at 9 Hz [67] with the displacement of a marker between two endpoints of 'no pain' and 'most intense pain imaginable' corresponding to values between 0 and 100. Trained using the CoVAS occurred prior to testing.

Heat pain threshold (HPT) individualized the subsequent OA protocol [2; 39; 51; 55; 56]. HPTs were determined by increasing the temperature of the thermode applied to the right forearm at a rate of 1.5°C/s from a baseline of 32-35°C until pain was reported. HPT trials were repeated three times, with 10 s rest intervals between trials.

The OA protocol comprised of an OA trial and a control trial, presented in random order. To ensure data accuracy, 60 s of rest followed data collection trials in addition to slight thermode repositioning on the forearm to prevent sensitization [66]. On an OA trial (Figure 1), three sequential temperatures (T1-3) were utilised: the initial stimulus temperature (T1) is the participant's HPT, ramped from the baseline temperature representing normal skin temperature (32-35°C) at a rate of 1.5℃/s. T1 (HPT) was held constant for 7 - 10 seconds. The duration of T1 varied to decrease participants' expectation of a temperature change at temperature 2 (T2). The T2 phase elevates the temperature a single degree to HPT + 1℃ for 5 s, and the Temperature 3 (T3) falls to HPT for 30 s before returning to baseline skin temperature at a rate of 6℃/s. (Figure 1). The transition from T2 to T3 is where OA is experienced, the characteristic disproportionate decrease in pain sensation relative to the associated temperature decrease. This OA protocol differs from some previous OA protocols [20; 66] because the T3 phase was extended to characterize the duration of OA, previous studies show ongoing OA at the termination of T3 [2; 20; 58].

139   Control trials consisted of the same T1 temperature as the OA trials (HPT), similarly ramped

140   from baseline skin temperature at a rate of 1.5℃/s. However, in contrast, T1 was sustained

141   for 30s, before ramping down to baseline skin temperature at rate of 6℃/s.

Transcranial direct current stimulation montages

Conventional cerebellar tDCS was applied using a neuroConn DC stimulator (NeuroConn

GmBH, Germany) with sponge electrodes (surface area 25 cm^2) soaked in 0.9% saline. The

stimulating electrode was positioned at the mid-point between the inion and the maximum

convexity of the right ear with the return electrode on the right buccinator[21; 23; 29].

Cerebellar HD-tDCS was applied using the neuroConn DC stimulator connected to a Soterix

4x1 HD-tDCS adaptor (Model 4x1-C2; Soterix Medical Inc., New York, USA).   A 4X1

electrode montage using sintered Ag/AgCl electrodes (EasyCap, GmbH, Germany) filled with

conductive gel (Electro-Gel, Brainmaster Inc., USA) were secured with an electrode cap

(EasyCap, GmbH, Germany). The central electrode was placed at the mid-point of the inion

and the maximum convexity of the right ear, with the four return electrodes spaced 3 cm

radially around the central electrode. Impedance values were measured for all electrodes

prior to usage to ensure good contact quality [61; 62]. Active NIBS applied current of 2mA of

tDCS for 20 minutes for both tDCS protocols. Sham stimulation applied 2mA for 30s to elicit a

tingling sensation, with no current applied in remainder of the 20 minute protocol [19; 60].

Current flow modelling is critical to NIBS protocol design[15; 45]. The cerebellar tDCS and

HD-tDCS protocols were simulated (HD-Explore Version 2.1, Soterix Medical, New York, US)

to characterise the current distribution with the given electrode montage and current level of

2mA as applied experimentally. All modelling was performed on a 93 standard adult male

head model sourced from the Soterix model library. Electrode positioning was based on the

10/20 EEG system. For conventional tDCS modelling, O10, PO10, EX2, and EX6

represented the cathode sponge electrode and EX14, EX16, and EX18 represented the

anodal sponge electrode (Figure 2a). For HD-tDCS modelling, PO10 represented the

cathode and PO4, P10, EXZ, and EX4 represented the anodes (Figure 2b).

OA and control trial response characterization

OA and Control response metrics were extracted using a custom written MATLAB script

(Version 9.0, The MathWorks Inc., MA, USA). Control trial metrics were mean (VAS $_{mean}$),

minimum (VAS $_{min}$), maximum (VAS$_{max}$) and standard deviation (VAS$_{sd}$) of the VAS score

while HPT was applied over a 30s period (T1).

CoVAS profiles were averaged dependent on neurostimulation (pre-stimulation, sham, active

stimulation) from the onset T3 for 22 s to qualitatively compare the impact of NIBS on the OA

response, which is defined as the VAS ratings made during T3. The OA response metrics

used in the study (Table 1, Figure 1) included both magnitude (Figure 1a) and temporal

(Figure 1b) features with metrics selected to be consistent with previous studies, VAS $_{min}$ [2;

10; 41], δVAS [1; 10; 41] and VAS $_{mean}$ [10]. Further, in this study, a second maximum VAS

was included since the OA protocol extended the T3 phase. Some OA metrics could not be

extracted on all trials due to features of the OA response, for example there could be no VAS

$_{2nd\ max}$ or OA $_{rate\ up}$ metric if the OA induced hypoanalgesia continued throughout T3.

Statistical analysis

Statistical analysis was performed in SPSS (IBM SPSS Statistics, Version 27; IBM Corp).

Descriptive statistics were collected for all OA metrics across conditions (pre-stimulation OA;

sham stimulation OA; active stimulation OA). Within day ICC on OA metrics are consistently

high in previous research [42] and pilot work in the laboratory. Between day ICC comparing

OA metrics at pre-stimulation baseline on each session were moderate or low. Therefore, the

tDCS and HD-tDCS experiments were considered separately via two 3X1 repeated

measures ANOVA for both control and OA trial metrics. Mauchly's test of sphericity was

performed, and when sphericity could not be assumed the Huynh-Feldt estimate was applied.

Post-hoc pairwise comparisons were made with Bonferroni corrections for multiple

comparisons.

Results

On control trials where T1 was held at HPT for 30s, the VAS_{mean} was 30.07 (SD 18.95), VAS_{max} of 42.31 (SD 22.41), VAS_{min} 15.96 (SD 18.67) and VAS_{sd} was 28.11 (SD 9.06). As expected, there were no consistent periods of hypoanalgesia observed in this VAS response to sustained HPT.

In contrast, the OA protocol successfully consistently induced hypoanalgesia, with only 3% of trials not showing a decrease in VAS at the T2-T3 transition. 90% of trials showed a fall greater than 10 VAS, with most trials eliciting a much larger decrease in pain perception.

The fall of temperature in T3 to HPT gave hypoanalgesia with a minimum VAS (VAS_{min}) at the nadir of the OA response of 12.71 (SD 15.50). The mean δVAS in the current study was slightly lower at 38.13 (SD, 22.39) relative to scores found by previous studies [1; 39; 40]. The T3 VAS_{mean} 30.35 (SD 23.61) for T3 phase was lower than studies that do not individualize OA protocols to HPTs [10]. The $OA_{rate\ down}$ at the T2-T3 transition a mean of 8.45 VAS/s (SD 8.45) was much steeper than $OA_{rate\ up}$, during T3, (1.85 VAS/s, SD 1.71), revealing the rapidity of OA at the transition and the sustained hypoanalgesia that demonstrates the necessity of extending T3 in the protocol. Taken together the protocol used in this study robustly generated OA and provided baseline values consistent with earlier studies in the area [1; 32; 39].

Modelling cerebellar NIBS

Current flow in the conventional tDCS (Figure 2a) is more diffuse than HD-tDCS (Figure 2b). Highest flow intensities with tDCS are near the active electrode, at the cerebellar cortical level in the posterior lobe. Medial structures also receive strong current and spread to the temporal lobe and frontal lobe. Current reached the upper and lower brainstem. For HD-tDCS, current flow was more restricted, with high field intensities in the posterior and anterior cerebellar lobe and posterior aspects of the temporal lobe. Very little current reaches the contralateral cerebellar hemisphere or brainstem.

The effect of cerebellar NIBS on OA

There were no significant differences in the control trial metrics (VAS $_{mean}$, VAS $_{min}$, VAS$_{max}$

and VAS$_{sd}$) dependent on stimulation for either tDCS or HD-tDCS. However, there was a

differential impact of NIBS on the OA response (Figure 3).

The average impact of active conventional cerebellar tDCS stimulation relative to pre-

stimulation and sham can be seen in Figure (3a).

Statistical analysis revealed there was no effect of cerebellar tDCS neurostimulation on

δVAS, VAS $_{min}$, OA $_{rate\ down}$, VAS $_{mean}$, OA $_{rate\ up}$ or OA l$_{atency}$ metrics of the OA response.

However, the effect of neurostimulation on OA $_{duration}$ was found to be significant $F_{(2,86)}=4.35$,

p =.037, partial η^2 = .07. Post-hoc pairwise comparisons with a Bonferroni adjustment

indicated there was no significance difference between OA $_{duration}$ at baseline and following

sham stimulation, p=.18.   However, OA $_{duration}$ was significantly longer after active tDCS

stimulation (11.17 s, SD 7.97) relative to pre-stimulation (8.11 s, SD 6.75), p=.048.

Interestingly, there was also no significant difference between OA $_{duration}$ after sham and after

active neurostimulation, p=1. OA $_{duration}$ represents the length of maximal OA; from the T2-T3

transition to the point at which pain rating begins to rise again (Figure 1b, Table 1, metric 8).

Additionally, there was an effect of stimulation of VAS $_{2nd\ max}$, $F_{(2,90)}=5.40$,p=.006, partial η^2 =

.11. Post-hoc pairwise comparisons with a Bonferroni adjustment showed there was a

significant difference between mean VAS $_{2nd\ max}$ at baseline (37.35, SD 26.94) and following

sham stimulation, (29.75, SD 25.32) p=.009 additionally VAS $_{2nd\ max}$ was significantly lower

after active tDCS stimulation, (29.52, SD 25.33), p=.028. However, the reduction in VAS $_{2nd}$

$_{max}$ also occurred after sham stimulation, with no significant difference between VAS $_{2nd\ max}$

and after active stimulation, p=1, suggestive of a placebo effect.

Second, the effect of the more focal 4 X 1 electrode montage was considered. Qualitatively,

the effect of active cathodal cerebellar HD-tDCS relative to sham is to enhance the OA effect

(Figure 3b). Statistical analysis revealed no effect of cerebellar HD-tDCS neurostimulation

was found on OA $_{duration}$, OA $_{latency}$ or OA $_{rate\ up.}$ However, there were differences on four OA

metrics. The effect of neurostimulation on δVAS, that is the drop in VAS at the T2-T3

transition was found to be significant $F_{(1.64,76.09)}=3.72$, p =.036, partial η^2 = .08. Post-hoc

pairwise comparisons with a Bonferroni adjustment indicated there was no significance difference between mean δVAS at baseline (32.70, SD 22.63) and following sham stimulation, (38.02, SD 23.51), p=.46. However, δVAS was significantly greater after active tDCS stimulation, (39.80, SD=21.79), p=.007. There was also no significant difference between δVAS after sham and after active neurostimulation, p=.96.

The effect of neurostimulation on VAS $_{rate\ down}$, that is the rate of OA-induced hypoanalgesia, was found to be significant $F_{(1.70, 71.18)}=3.36$, p =.048, partial η^2 = .07. Post-hoc pairwise comparisons with a Bonferroni adjustment indicated there was no significance difference between mean VAS $_{rate\ down}$ at baseline (6.59 VAS/s, SD 4.75) and following sham stimulation, (8.91 VAS/s, SD=8.38), p=.55. However, VAS $_{rate\ down}$ was significantly faster after active tDCS stimulation, (8.31 VAS/s, SD 5.37) p=..021. There was also no significant difference between VAS $_{min}$ after sham and after active neurostimulation, p=.86.

The effect of neurostimulation on VAS $_{min}$, the VAS at the nadir of the OA response was found to be significant $F_{(2,90)}=4.96$, p =.01, partial η^2 = .1. Post-hoc pairwise comparisons with a Bonferroni adjustment indicated there was no significance difference between mean VAS$_{min}$ at baseline (16.17, SD=20.25) and following sham stimulation, (10.08, SD 11.62), p=.12. However, mean VAS$_{min}$ was significantly reduced after active tDCS stimulation, (9.79, SD 11.62), p=.018. There was also no significant difference between VAS$_{min}$ after sham and after active neurostimulation, p=1.

There was an effect of stimulation of VAS $_{mean}$, $F_{(2,90)}=4.91$, p =.009, partial η^2 = .98. Post-hoc pairwise comparisons with a Bonferroni adjustment indicated there was no significance difference between mean VAS$_{mean}$ at baseline (32.04, SD 23.81) and following sham stimulation, p=.055. However, VAS $_{mean}$ was significantly lower after active tDCS stimulation, (26.02, SD 21.75) p=.023. There was also no significant difference between VAS $_{mean}$ after sham (27.79, SD 18.99), and after active neurostimulation, p=1.

The effect of neurostimulation on VAS$_{2nd\ max}$ was found to be significant $F_{(2,90)}=3.56$, p =.033, partial η^2 = .07 indicating pain does not rise as much even though OA is reduced relative to the start of T3. Post-hoc pairwise comparisons with a Bonferroni adjustment indicated there was no significance difference between mean VAS$_{2nd\ max}$ at baseline (32.04, SD 23.81) and following sham stimulation, (27.29, SD 18.99), p=.19. However, VAS$_{2nd\ max}$

was significantly reduced after active tDCS stimulation, (26.02, SD 21.75), p=.038. There was also no significant difference between $VAS_{2nd\ max}$ after sham and after active neurostimulation, p=1.

Discussion

The individualized offset analgesia protocol provided a robust methodology for inducing hypoanalgesia. Given the recent advances in NIBS as interventions for pain modulation [13], NIBS of the cerebellum is potentially an important means of further elucidating this role and potentially augmenting hypoanalgiesa. A recent study has indicated anodal tDCS of the cerebellum marginally enhances endogenous pain modulation whereas cathodal tDCS suppresses it, with small effect sizes [53]. However, recent research on the optimization of NIBS suggest it is critical to evaluate differing electrode montages and current dosage, with these parameters paramount for the specific effect and yet gaining less attention than target area and polarity[6; 7; 46].

Here cathodal cerebellar HD-tDCS was found to augment the effects of OA relative to pre-stimulation with VAS pain ratings in T3 both lower at the nadir and demonstrating more sustained hyponanalgesia. Conventional cerebellar tDCS increased OA on two metrics relative to pre-stimulation, with an increase in the initial hypoanalgesia experienced. Qualitatively there appears to be a trend towards increased OA after sham (Figure 3), and one metric in the conventional cerebellar tDCS experiment showed a significant difference post sham, suggestive of a placebo effect. However, no other OA metrics showed an effect of sham in either experiment. Further, the differential impact of the NIBS interventions on OA is observable both qualitatively and statistically, indicating that interpretation must include consideration of current flow and the focality of current applied. Additionally, these differential effects indicate that there is not simply an effect of sequential testing of an OA protocol, which has been found in some studies to enhance the analgesia experienced[9]. The study supports increased focality and attention to current flow to different cerebellar regions as critical for further understanding the cerebellum's role in OA. Cerebellar stimulation presents a challenge for NIBS due to the relatively small target regions [59] that may optimally enhance OA. Animal studies show both pro and anti-nociceptive effects of the cerebellum[11;

52], therefore exploration of different high-definition tDCS montages will enhance understanding further in future studies. In parallel, the exact mechanisms of NIBS on the cerebellum are less well established than cortical regions [60] therefore increased understanding of mechanisms from animal studies alongside current flow modelling would further benefit NIBS design [8].

Cerebellar tDCS mediating analgesic enhancement can be explained by several different pathways. First, cathodal cerebellar tDCS may decrease inhibitory effects of Purkinje cells and so increases deep cerebellar nuclei activity to indirectly reduce inhibition of M1 [18]. Numerous studies link analgesia with primary motor cortex excitability and the cerebellum has a known role in modulating corticomotoneuronal activity[44]. Alternatively, increased caudate putamen activity is associated with cathodal and anodal cerebellar tDCS [30]. The caudate is a known region for pain suppression [65] and is critically involved in the reward pathways. The hypoanalgesia experienced after the transient decrease in supra-threshold pain, that is characteristic of the OA protocol, is known to activate reward pathways. Additionally, cerebellar stimulation may impact on OA via connectivity with known pain modulation pathways including the PAG and dorsolateral prefrontal cortex (DLPFC)[5].

To conclude, offset analgesia is a robustly observed form of hypoanalgesia and better understanding of the underlying mechanism could open therapeutic options. The role of the cerebellum in pain processing is further strengthened by this study. High-definition cerebellar tDCS shows promise as a novel neuromodulation protocol to enhance endogenous pain modulation pathways. Further investigation should occur of the optimal electrode montage, current level, and duration for cerebellar stimulation. Such 'dosing' effects of neuromodulation are currently under-explored yet likely critical for maximizing NIBS effects, as well as improving mechanistic understanding.

Author contributions

NOC and AW designed the experimental protocol with EDG involved in pilot testing; NOC, HA, ROF collected the experimental data, MW and AW collated the data, MW and AW

336   performed statistical and data analysis. NOC, HA, ROF, EDG contributed to initial drafts. MW

337   and AW wrote the manuscript.

Disclosure

339   The author reports no conflicts of interest in this work.   A proportion of this work was

340   presented as an abstract at EFIC, Valencia 2019

References

[1] Alter BJ, Aung MS, Strigo IA, Fields HL. Onset hyperanalgesia and offset analgesia:
transient increases or decrease of noxious thermal stimulus intensity robustly
modulate subsequent perceived pain intensity. PLoS One 2020;15(e0231124).
[2] Alter BJ, Santosa H, Nguyen QH, Huppert TJ, Wasan AD. Offset analgesia is associated
with opposing modulation of medial versus dorsolateral prefrontal cortex activations:
a functional near infrared spectroscopy study. Molecular Pain 2022;18.
[3] Antal A, Brepohl N, Poreisz C, Boros K, Csifcsak G, Paulus W. Transcranial direct current
stimulation over somatosensory cortex decreases experimentally induced acute pain
perception. Clinical Journal of Pain 2008;24:56-63.
[4] Borsook D, Moulton EA, Tully S, Schmahmann JD, Becerra L. Human cerebellar
responses to brush and heat stimuli in healthy and neuropathic pain subjects.
Cerebellum 2008;7(3):252-272.
[5] Cacciola A, Bertino S, Basile GA, Di Maruo D, Calamuneri A, Chillemi G, Duca A,
Bruschetta D, Flace P, Favalaro A, Calabro RS, Anastasi G, Milardi D. Mapping the
structural connectivity between the periaqueductal gray and the cerebellum in
humans. Brain Structure Function 2019;224:2153-2165.
[6] Caulfield KA, Badran BW, DeVries WH, Summers PM, Kofmehl E, Li X, Borckardt JJ,
Bikson M, George MS. Transcranial electrical stimulation motor threshold can
estimate individualised tDCS dosage from reverse calculation electric-field modeling.
Brain Stimulation 2020;13:961-969.
[7] Caulfield KA, George MS. Optimized APPS-tDCS electrode position, size, and distance
doubles the on-target stimulation magnitude in 3000 electric field models. Scientific
Reports 2022;12.
[8] Dayan E, Censor N, Buch ER, Sandrini M, Cohen LG. Noninvasive brain stimulation: from
physiology to network dynamics and back. Nat Neurosci 2013;16:838-844.
[9] Derbyshire SWG, Osborn J. Enhancement of offset analgesia during sequential testing.
European Journal of Pain 2008;12:980-989.
[10] Derbyshire SWG, Osborn J. Offset analgesia is mediated by activation in the region of
the periaqueductal grey and rostral ventromedial medulla. Neuroimage
2009;47:1002-1006.
[11] Dey PK, Ray AK. Anterior cerebellum as a site for morphine analgesia and post-
stimulation analgesia. Indian J Physiol Pharmacol;26:3-12.
[12] Diano M, D'Agata F, Cauda F, Costa T, Geda E, Sacco K, Duca S, Torta DM, Germiniani
GC. Cerebellar clustering and functional connectivity during pain processing.
Cerebellum 2015.
[13] Fenton BW, Palmieri PA, Boggio PS, Fanning J, Fregni F. A preliminary study of
transcranial direct current stimulation for the treatment of refactory chronic pelvic
pain. Brain Stimulation 2009;2(2):103-107.
[14] Ferrucci R, Cortese F, Priori A. Cerebellar tDCS: how to do it. The Cerebellum
2015;14:27-30.
[15] Fiocchi S, Chiaramello E, Gazzola V, Suttrup J, Ravazzani P, Parazzini M. Assessment
of the capability to target cerebellar sub-regions with high-definition transcranial direct

current stimulation (HD-tDCS) over the cerebellum. IEEE 3rd international forum on
research and technologies for society and industry (RTSI) 2017:1-6.
[16] Fregni F, Boggio PS, Lima MC, Ferreira MJL, Wagner T, Rigonatti SP, Castro AW,
Souza DR, Riberto M, Freedman SD, Nitsche MA, Pascual-Leone A. A sham-
controlled, phase II trial of transcranial direct current stimulation for the treatment of
central pain in traumatic spinal cord injury. Pain 2006;122:197-209.
[17] Fregni F, Gimenes R, Valle AC, Ferreira MJL, Rocha RR, Natalle L, Bravo R, Rigonatti
SP, Freedman SD, Nitsche MA, Pascual-Leone A, Boggio PS. A randomised, sham-
controlled, proof-of-principle study of transcranial direct current stimulation for the
treatment of pain in fibromyalgia. Arthritis and Rheumatism 2006;54:3988-3998.
[18] Galea JM, Jayaram G, Ajagbe L, Celnik P. Modulation of cerebellar excitability by
polarity-specific noninvasive direct current stimulation. J Neurosci 2009;29:9115-
9122.
[19] Gandiga PC, Hummel FC, Cohen LG. Transcranial DC stimulation (tDCS): a tool for
double blind sham-controlled clinical studies in brain stimulation. Clin Neurophysiol
2006;117:845-850.
[20] Grill JD, Coghill RC. Transient analgesia evoked by noxious stimulus offset. J
Neurophysiol 2002;87:2205-2208.
[21] Hamada M, Strigaro G, Murase N, Sadnicka A, Galea JM, Edwards MJ, Rothwell JC.
Cerebellar modulation of human associative plasticity. Journal of Physiology
2012;590:2365-2374.
[22] Hermans L, Calders P, Van Oosterwijck J, Verschelde E, Bertel E, Meeus M. An
overview of offset analgesia and the comparison with conditioned pain modulation: a
systematic literature review. Pain Physician 2016;19:307-326.
[23] Herzfeld DJ, Pastor D, Haith AM, Rossetti Y, Shadmehr R, O'Shea J. Contributions of the
cerebellum and the motor cortex to acquisition and retention of motor memories.
Neuroimage 2014;98:147-158.
[24] Keltner JR, Furst A, Fan C, Redfern R, Inglis B, Fields HL. Isolating the modulatory effect
of expectation on pain transmission: a functional magnetic resonance imaging study.
J Neurosci 2006;26(16):4437-4443.
[25] Kong Q, Li T, Reddy S, Hodges S, Kong J. Brain stimulation targets for chronic pain:
insights from meta-analysis, functional connectivity and literature review.
Neurotherapeutics 2024;21(e00297).
[26] Lefaucher JP, Antal A, Ahdab R, Ciampi de Andrade D, Fregni F, Khedr EM, Nitsche M,
Paulus W. The use of repetitive transcranial magnetic stimulation (rTMS) and
transcranial direct current stimulation (tDCS) to relieve pain. Brain Stimulation
2008;1:337-344.
[27] Li CN, Keay KA, Henderson LA, Mychasiuk R. Re-examining the mysterious role of the
cerebellum in pain. J Neurosci 2024;44(17):1-18.
[28] Lima MC, Fregni F. Motor cortex stimulation for chronic pain: systematic review and
meta-analysis of the literature. Neurology 2008;70:2329-2337.
[29] Macher K, Bohringer A, Villringer A, Pleger B. Cerebellar-parietal connections underpin
phonological storage. Journal of Neuroscience 2014;34:5029-5037.
[30] Maldonado T, Jackson TB, Bernard JA. Anodal cerebellar stimulation increases cortical
activation: evidence for cerebellar scaffolding of cortical processing. Human Brain
Mapping 2023;44(4):1666-1682.
[31] Manto M, Argyropolos GPD, Bocci T, Celnik PA, Corben LA, Guidetti M, Koch G, Priori A,
Rothwell JC, Sadnicka A, Spampinato D, Ugawa Y, Wessel MJ, Ferrucci R.
Consensus paper: novel directions and next steps of non-invasive brain stimulation of
the cerebellum in health and disease. The Cerebellum 2022;21:1092-1122.
[32] Martucci KT, Eisenach JC, Tong C, Coghill RC. Opiod-independent mechanisms
supporting offset analgesia and temporal sharpening of nociceptive information. Pain
2012;153:1232-1243.
[33] Medoc L. CoVAS (apparatus and software), 1996.
[34] Medoc L. TSA-II Neurosensory Analyzer (apparatus and software), 1999.
[35] Mordillo Mateos L, Dileone M, Soto-Leon V, Brocalero-Camacho A, Perez-Borrego YA,
Onate-Figuerez A, Aguilar JAM, Oliviero A. Effects of transcranial direct current
stimulation on temperature and pain perception. Scientific Reports 2017;7.

[36] Morton D, Jones A, Sandhu J. Brain imaging of pain: state of the art. Journal of Pain Research 2016;9:613-624.

[37] Moulton EA, Elman I, Pendse G, Schmahmann J, Becerra L, Borsook D. Aversion-related circuitry in the cerebellum: responses to noxious heat and unpleasant images. J Neurosci 2011;31:3795-3804.

[38] Moulton EA, Schmahmann JD, Becerra L, Borsook D. The cerebellum and pain: passive integrator or active participator? Brain Res Rev 2010;65(1):14-27.

[39] Naugle KM, Cruz-Almeida Y, Fillingim RB, Riley JL. Offset analgesia is reduced in older adults. Pain 2013;154:2381-2387.

[40] Niesters M, Dahan A, Swartjes M, Noppers I, Fillingim RB, Aarts L, Sarton EY. Effect of ketamine on endogenous pain modulation in healthy volunteers. Pain 2011;152(3):656-663.

[41] Niesters M, Hoitsma E, Sarton E, Aarts L, Dahan A. Offset analgesia in neuropathic pain patients and effect of treatment with morphine and ketamine. Anesthesiology 2011;115:1063-1071.

[42] Nilsson M, Nissen TD, Graversen C, Gazerani P, Drewes AM, Brock C. Offset analgesia: a reproducibility study. Scandinavian Journal of Pain 2012;3:192.

[43] Oudejans LCJ, Smit JH, van Velzen M, Dahan A, Niesters M. The influence of offset analgesia on the onset and offset of pain in patients with fibromyalgia. Pain 2015;156:2521-2527.

[44] Oulad Ben Taib N, Manto M. Trains of epidural DC stimulation of the cerebellum tune corticomotor excitability. Neural Plasticity 2013:1-12.

[45] Parazzini M, Rossi E, Ferrucci R, Fiocchi S, Liorni I, Priori A, Ravazzani P. Computational model of cerebellar transcranial direct current stimulation. Conf Proc Annu Int Conf IEEE Eng Med Biol Soc 2013;IEEE Eng Med Biol Soc Annu Conf:237-240.

[46] Peterchev AV, Wagner TA, Miranda PC, Nitsche MA, Paulus W, Lisanby SH, Pascual-Leone A, Bikson M. Fundamentals of transcranial electric and magnetic stimulation dose: definition, selection and reporting practices. Brain Stimulation 2012;5(4):435-453.

[47] Ploghaus A, Tracey I, Clare S, Gati JS, Rawlins NP, Matthews PM. Learning about pain: The neural substrate of the prediction error for aversive events. PNAS 2000;97:16.

[48] Ruscheweyh R, Kuhnel M, Filippopulos F, Blum B, Eggert T, Straube A. Altered experimental pain perception after cerebellar infarction. Pain 2014;155:1303-1312.

[49] Saab CY, Kawasaki M, Al-Chaer ED, Willis WD. Cerebellar cortical stimulation increase spinal visceral nociceptive responses. J Neurophysiol 2001;85:2359-2363.

[50] Saab CY, Wilis WD. The cerebellum: organization, functions and its role in nociception. Brain Res Brain Res Rev 2003;42:85-95.

[51] Shulman J, Zurakowski D, Keysor J, Jervis K, Sethna NF. Offset analgesia identifies impaired endogenous pain modulation in pediatric chronic pain disorders. Pain 2020;161(2852).

[52] Siegel P, Wepsic JG. Alteration of nociception by stimulation of cerebellar structures in the monkey. Physiol Behav 1974;13:189-194.

[53] Stachendeder R, Alt L, Straube A, Ruscheweyh R. Effects of transcranial direct current stimulation (ct-DCS) of the cerebellum on pain perception and endogenous pain modulation: a randomized, moncentric, double-bllind, sham-contolled crossover study. The Cerebellum 2022.

[54] Strick PL, Dum RP, Fiez JA. Cerebellum and nonmotor function. Annual Review of Neuroscience 2009;32:413-434.

[55] Szikszay TM, Adamczyk WM, Carvalho GF, May A, Luedtke K. Offset analgesia: somatotopic endogenous pain modulation in migraine. Pain 2020;161(3):557-564.

[56] Szikszay TM, Adamczyk WM, Carvalho GF, May A, Luedtke K. Offset analgesia is increased intra-orally. Journal of Oral Rehabilitation 2022;49(10):993-1001.

[57] Szikszay TM, Adamczyk WM, Luedtke K. The magnitude of offset analgesia as a measure of endogenous pain modulation in healthy participants and patients with chronic pain: a systematic review and meta-analysis. The Clinical Journal of Pain 2019;35:189-204.

[58] Szikszay TM, Melz N, von Glasenapp B, Adamczyk WM, Luedtke K. Effects of
stimulation area and temperature rates on offset analgesia. Pain Reports
2022;7(e1043).
[59] Tomlinson SP, Davis NJ, Bracewell RM. Brain stimulation studies of non-motor cerebellar
function: A systematic review. Eurosci biobehav Rev 2013;37(5):766-789.
[60] Van Dun K, Bodranghien FCAA, Marien P, Manto MU. tDCS of the cerebellum: where do
we stand in 2016? Technical issues and critical review of the literature Frontiers in
Human Neuroscience 2016;10(199).
[61] Villamar MF, Volz MS, Bikson M, Datta A, Dasilva AF, Fregni F. Techniques and
considerations in the use of 4X1 ring high-definition transcranial direct current
stimulation (HD-tDCS). J Vis Exp 2013;e50309.
[62] Villamar MF, Wivatvongvana P, Patumanond J, Bikson M, Truong DQ, Datta A, Fregni F.
Focal modulation of the primary motor cortex in fibromyalgia using 4X1 ring high
definition transcranial direct current stimulation (HD-tDCS): immediate and delayed
analgesic effects of cathodal and anodal stimulation. Journal of Pain 2013;14:371-
383.
[63] Witney AG. From mechanisms to analgesia: towards the use of non-invasive
neuromodulation for pain relief in the clinic. In: D Larrivee, SM Rayegani, editors.
Neurostimulation and Neuromodulation in Contemporary Therapeutic Practice.
London: Intech Open, 2020. pp. 191-208.
[64] Woods AJ, Antal A, Bikson M, Boggio PS, Brunoni AR, Celnik P, Cohen LG, Fregni F,
Herrmann CS, Kappenman ES, Knotkova H, Liebetanz D, Miniussi C, Miranda PC,
Paulaus W, Priori A, Reato D, Stagg CJ, Wenderoth N, Nitsche MA. A technical guide
to tDCS and related non-invasive brain stimulation tools. Clincial Neurophysiology
2016;127:1031-1048.
[65] Wunderlich AP, Klug R, Stuber G, Landwehrmeyer B, Weber F, Feund W. Caudate
nucleus and insular activation during a pain suppression paradigm comparing thermal
and electrical stimulation The Open Neuroimaging journal 2011;5:1-8.
[66] Yelle MD, Rogers JM, Coghill RC. Offset analgesia: a temporal contrast mechanisms for
nociceptive information. Pain 2008;134:174-186.
[67] Zhang K-L, Yuan H, Wu F-F, Pu X-Y, Liu B-Z, Li Z, Liu H, Yang Y, Wang Y-Y. Analgesia
effect of noninvasive brain stimulation for neuropathic pain patients: a systematic
review. Pain Therapy 2021;10:315-332.
[68] Zunhammer M, Busch V, Griesbach F, Landgrebe M, Hajak G, Langguth B. rTMS over
the cerebellum modulates temperature detection and pain thresholds through
peripheral mechanisms. Brain Stimulation 2011;4:210-217.
535
536

Figure Legends

Figure 1: Offset analgesia protocol showing extracted OA metric definitions for magnitude (A)

1: VAS $_{min}$, 2: VAS $_{mean}$, 3: δVAS $_{OA}$ and temporal (B) 5: OA rate $_{down}$, 6: OA $_{rate\ up}$, 7: OA $_{latency}$

and 8: 0A $_{duration}$ of the OA response. See Table 1 for OA metric definitions.

Figure 2: Current flow modelling: 3D representation of current flow (upper panel), current flow

through coronal, sagittal and axial slices (lower panel) A) Cerebellar tDCS montage: High

intensity in the cerebellum, temporal lobe, frontal lobe and brainstem. B) Cerebellar HD-tDCS

montage restricted current to the right cerebellar hemisphere and the posterior occipital lobe,

with peak field intensity in the posterior cerebellum.

546 **Figure 3:** Average of OA response during T3 period dependent on cerebellar NIBS

547 stimulation for a) tDCS protocol and b) HD-tDCS protocol. Pre-stimulation (red); Sham-

548 stimulation (blue); Active-stimulation (green). Shading represents standard error of the mean.

549 Table Legends

550 **Table 1:** OA response metric definitions for magnitude and temporal features (see Figure 1

551 for graphical representation)

554 **Table 1:**

Metric Label	Offset Analgesia Metric	Description
1	VAS_{min}	Lowest reported VAS in t3
2	VAS_{mean}	Mean VAS in t3
3	δVAS_{OA}	Magnitude of OA: the change in VAS at t2-t3
4	$VAS_{2nd\ Max}$	Maximum VAS in t3 following hypoanalgesia
5	$OA_{rate\ down}$	Rate of decrease in VAS (δVAS) due to OA
6	$OA_{rate\ up}$	Rate of increase from VAS_{min} to $VAS_{2nd\ Max}$
7	$OA_{latency}$	Time taken for maximum VAS score at the start of t3 to drop to VAS_{min} (the OA induced hypoanalgesia)
8	$OA_{duration}$	Time taken for VAS_{min} to rise again following hypoanalgesia

Figure 1

A tDCS

Coronal Slice   Saggital Slice   Axial Slice

Field Intensity V/m   0.15   0

L   R   F   B   L   R

B HD-tDCS

Coronal Slice   Saggital Slice   Axial Slice

Field Intensity V/m   0.15   0

L   R   F   B   L   R

Differential effect of neurostimulation on the OA response

A **tDCS, pre/sham/active**

T3

HPT ⁰C

Co-VAS (1-100)

Time(s)

B **HD-tDCS, pre/sham/active**

T3

HPT ⁰C

Co-VAS (1-100)

Time(s)