Reduced gravity and muon flux absence affect *Caenorhabditis elegans* life history traits and viral infection

Ana Villena-Giménez,[1] Victoria G. Castiglioni[1], Juan C. Muñoz-Sánchez,[1,2], Esmeralda G. Legarda[1], Rubén González[3], and Santiago F. Elena[1,4]

[1]Instituto de Biología Integrativa de Sistemas (CSIC-UV), Paterna, 46980 Valencia, Spain

[2]Department of Theoretical Physics, Universitat de València, Burjassot, 46100 Valencia, Spain

[3]Institute Pasteur, 75015 Paris, France

[4]Santa Fe Institute, Santa Fe, NM 87501, USA

Correspondence: santiago.elena@csic.es

ORCIDs:

A.V-G.: 0009-0000-3564-4293

V.G.C.: 0000-0001-9884-2537

J.C.M-S.: 0000-0002-5570-4983

E.G.L.: 0000-0002-8560-8407

R.G.: 0000-0001-9588-1127

S.F.E.: 0000-0001-8249-5593

ABSTRACT Environmental conditions fundamentally shape host-pathogen interactions, yet how multiple extreme abiotic stressors combine to influence infection outcomes remains poorly understood. Organisms evolved under specific gravitational and radiation regimes; deviations from these conditions —whether in extreme terrestrial environments or beyond Earth— may alter immune function and pathogen replication. In this study, we investigated the effects of reduced gravity and lowered muon flux on Orsay virus infection in the nematode *Caenorhabditis elegans*. We employed a fully factorial experimental design, examining how each factor, alone and in combination, influences physiological traits and viral load. While below-background radiation radically affected viral accumulation dynamics, reduced gravity had a minor effect. Both factors significantly impacted reproduction and morphology, with some effects magnified by viral infection. These results reveal how even partial modifications of Earth-like gravity and radiation levels can alter pathogen–host interactions. By integrating experimental observations with mathematical modeling, we show that these environmental stressors primarily affect prezygotic reproductive processes and modulate viral replication through distinct and sometimes antagonistic mechanisms. Although this work does not encompass the full complexity of space environments, where cosmic radiation includes high-energy protons and heavy ions, it provides insight into how adjustable models of reduced gravity and radiation can advance our understanding of biological adaptation beyond standard terrestrial conditions.

IMPORTANCE Understanding how extreme environmental conditions affect host–pathogen interactions is critical both for safeguarding biological systems during spaceflight and for exploring fundamental principles of stress biology. This study demonstrates that reduced gravity and diminished muon radiation flux can significantly alter viral infection dynamics and host physiology in *Caenorhabditis elegans*. By integrating experimental data with mathematical modeling, we reveal that these abiotic stresses impact prezygotic reproductive processes and modulate viral replication in distinct and sometimes antagonistic ways. Our findings suggest that even partial deviations from Earth-like conditions can reshape infection outcomes and developmental trajectories, highlighting the need for deeper mechanistic insights into biological adaptation beyond terrestrial norms. These results have implications for space biosciences, evolutionary virology, radiation hormesis theory, and the design of countermeasures to preserve organismal health in extreme or non-terrestrial habitats.

KEYWORDS environment-dependent virulence, host-virus interactions, mathematical modeling, off-Earth stresses, Orsay virus, radiation hormesis

Environmental conditions fundamentally shape host-microbe interactions, yet our understanding of how extreme environments alter these relationships remains limited. Organisms evolved under specific gravitational and radiation regimes, and deviations from these conditions may reveal fundamental constraints on physiological and immune function. Such deviations occur in off-Earth environments where organisms may experience a variety of unusual stresses, including reduced gravitational forces (microgravity, µG) and substantial changes in radiation exposure. Understanding how biological systems respond to such perturbations is essential for comprehending life's responses to conditions beyond Earth: whether encountered during space exploration, in extreme environments, or throughout evolutionary history.

Off-Earth environments can affect the organism's overall health and immunity. For instance, within relatively short-duration orbital missions (10 - 15 days), blood samples from astronauts showed altered levels of leukocyte distribution and cytokine production and a decrease in the function of Epstein-Barr virus specific CD8[+] T cells (1). During longer missions (6 months), immune alterations persisted, as evidenced by reduced T-cell function (2), impairment of natural killer cells function (3) and elevated levels of salivary antimicrobial proteins (4). Indeed, the reactivation of latent herpesviruses is well documented in astronauts (4 - 6): Epstein-Barr virus, varicella-zoster virus and cytomegalovirus infections increased in frequency, duration and amplitude the longer the duration of space missions (5, 7). These observations raise fundamental questions about how gravitational and radiation changes modulate host-pathogen dynamics.

Radiation composition in off-Earth settings differs substantially from that on Earth's surface. One of the radiation components that differs between space and Earth's surface is the muon, a charged elementary particle produced when pions decay following cosmic ray interactions with atmospheric molecules. Muons are the main component of cosmic radiation at sea level, with about 600 muons crossing a human body every minute (8). As an important source of radiation for living beings, cells have likely developed mechanisms that buffer muon's detrimental effects (9, 10). This observation is relevant to radiation hormesis, in which organisms may benefit from low-level radiation exposure but experience detrimental effects when doses deviate significantly in either direction (11). Understanding biological responses to conditions with dramatically reduced muon flux can therefore reveal whether organisms require baseline radiation exposure for optimal function, a key prediction of hormesis theory.

Caenorhabditis elegans has long served as a valuable model for studying diverse stresses that can be challenging to dissect in more complex animals. *C. elegans* is a well-characterized metazoan that has been widely employed as a model organism (12). It has many advantages: easy visualization, fast development, a complete characterized cell lineage and genome, and an estimated genetic homology of ~80% of the protein-coding genome with humans (13). *C. elegans* is exposed to fluctuating environmental conditions in its natural environment, such as heat, cold, food scarcity, oxygen levels, and osmotic perturbations. While repeated mild heat-shock

treatments extend *C. elegans* lifespan (14), moderate treatments impair fecundity (15) and embryogenesis (16), and severe treatments decreases longevity (17). These heat-shock phenotypic alterations are synchronized with a strong change in the transcriptional response (18). Exposure to other abiotic stresses, such as hypertonic environments induce transcriptional responses that mimic infection response to a variety of pathogens (19) and heat shocked animals have increased survival upon exposure to *Staphylococcus aureus* (20). In contrast, animals fed with pathogenic bacteria (21) or infected with virus (22) show an increased resistance to heat shock and ubiquitin ligases involved in the intracellular pathogen response (IPR) promote thermotolerance in order to cope with proteotoxic stress, unveiling the close link between biotic and abiotic stresses (23). These findings demonstrate that stress responses and immune pathways are deeply interconnected, raising the question of how non-standard gravitational and radiation environments might modulate infection outcomes.

C. elegans has long been used as a model in immunity. It can regulate immune responses recognizing pathogens, influencing the body's development, and controlling avoidance behavior (24). Although there are significant differences with the immune system of mammals, some mechanisms used to limit pathogenesis show remarkable phylogenetic conservation, such as uridylation and RNAi pathways (25, 26). Since the discovery of Orsay virus (OrV), a lot of effort has been put into the characterization of virus-host interaction dynamics. OrV is a bipartite positive-sense single-stranded RNA virus related to the *Nodaviridae* family that infects *C. elegans* intestinal cells (27). This natural host-pathogen system provides an ideal model for studying how environmental stressors influence real viral infection dynamics, in contrast to artificial infection models. Viral infection produces mild intestinal symptoms such as changes in cytoplasm viscosity, nuclear degeneration and cellular fusion (27), while mild effects in progeny production where only observed in a more susceptible strain (26). Links between viral infection and abiotic stresses have been described in the nematode, with heat stress reducing its susceptibility to OrV (28) and, conversely, OrV-infected animals showing an increased tolerance to heat-shock (22).

The effect of lack of muon radiation flux, hereafter referred as below background radiation (BBR), and of µG on *C. elegans* and other model organisms has been explored before. Van Voorhies *et al*. (29) carried out phenotypic and transcriptomic studies in *C. elegans* at the underground laboratory at the Waste Isolation Pilot Plant (New Mexico, USA). Animals had faster larval growth rates, higher egg laying rates and a different gene expression profile, both at 72 h after being at BBR and after 10 generations. Moreover, a reduction in fertility has been observed in *Drosophila melanogaster* flies reared at the Gran Sasso Underground laboratory (Italy) (30). Studies in simulated µG conditions in *C. elegans* have reported no reproductive and lifespan changes (31), increased intestinal barrier permeability (32) and induction of oxidative stress and dysregulated antioxidant machinery (33).

In this study, we have explored the role of two environmental stressors on the outcome of a viral infection. To do so, we have studied *C. elegans*/OrV pathosystem under a factorial combination of two abiotic factors: gravity intensity (standard and μG), using a Random Positioning Machine (RPM), and muon radiation flux (standard and BBR), achieved by conducting experiments in the Canfranc Underground Laboratory (LSC; Estación de Canfranc, Huesca, Spain). We acclimatized the nematodes to the abiotic stresses and performed the experiments in the third generation of acclimatized animals. This enabled us to investigate long-term effects of μG and BBR in the nematode since it is known that in *C. elegans* responses to stress are transmitted intergenerationally (32). We have measured different fertility traits as a proxy for the animal's fitness, and body length and morphology to identify developmental impairments. We also assessed intestinal barrier permeability to determine the functional state of the intestine under these stresses, since OrV infects intestinal cells and intestinal collagens are involved in preventing viral entry (34). As a proxy for viral fitness under these different stress conditions, we quantified OrV genomic RNA2 accumulation. Finally, the data was fitted to a fertility model and different parameters associated with prezygotic (early oocyte development, maturation and fertilization rates) and postzygotic (abortion rate) effects estimated. By combining phenotypic observations with mathematical modeling, we reveal how altered gravity and radiation interact to reshape host-pathogen dynamics.

RESULTS

The impairment of intestinal barrier by μG is antagonized by BBR

Intestinal integrity is critical for overall health (35), as it is key to nutrient absorption and serves as a primary defense against enteric pathogens, including OrV. Previous work has shown that various abiotic stresses can alter intestinal permeability, with μG causing a significant increase (31). Accordingly, we sought to validate our simulated microgravity conditions by determining whether they recapitulate this known phenotype. Additionally, we examined how BBR might counterbalance or compound these effects. To measure gut permeability, we employed a dye-based assay in which nematodes are exposed to a blue dye. An intact intestinal barrier restricts the dye to the lumen; conversely, compromised integrity leads to dye leakage into surrounding tissues (Fig. 1A). Firstly, we observed a large and significant overall increase in permeability of animals grown in μG conditions (χ^2 = 15.882, 1 d.f., P < 0.001, η_P^2 = 0.187), as previously reported. Conversely, BBR conditions were associated to a moderate but still significant reduction in permeability (χ^2 = 8.387, 1 d.f., P = 0.004, η_P^2 = 0.099). No significant interaction between the two factors has been found (χ^2 = 0.461, 1 d.f., P = 0.497), suggesting that permeability alterations are a consequence of overlapping mechanisms (Fig. 1B). These results

167 validate our experimental system and demonstrate that µG and BBR exert opposing effects on
168 intestinal physiology.

FIG 1 (A) Representative animals in standard (non-permeable) and µG conditions (impaired permeability). Nematodes showing blue staining outside the intestinal tube were scored as permeable whether the salt diffused throughout the body out. (B) Intestinal permeability of animals under standard gravity ($n = 30$), µG ($n = 28$), BBR ($n = 34$) and µG plus BBR ($n = 24$) conditions. Synchronized L1 animals were grown at 20 ºC during 48 h and then fed with a solution containing erioglaucine disodium for 3 h. Error bars represent ±1 SEM.

Viral accumulation dynamics are affected by gravity and radiation intensity

Having confirmed that our simulated conditions produced the expected phenotypic changes in *C.*
elegans (*e.g.*, increased intestinal permeability under reduced gravity), we next examined whether
viral infections remained successful under these modified environments. Our laboratory has
previously demonstrated that, under standard laboratory conditions (*i.e.*, Earth-like gravity and
muon radiation flux), OrV infection in *C. elegans* follows two waves of viral load from L1 larvae
to sexually mature adults (36). A first peak is reached approximately 12 hours post-inoculation
(hpi) in L1 larvae, which is followed by a sharp decline as antiviral responses are activated by the

developing animals, reaching a minimum approximately 18 – 20 hpi (L2 larvae). From there on,

a second wider peak is reached that last for at least 18 additional hours (L3 larvae and L4),

afterwards, viral load is low, yet viral replication continues at low levels (36). In this work, we

observed that under our standard reference conditions, OrV likewise exhibited these two distinct

waves, thus confirming the reproducibility of its infection dynamics (Fig. 2, upper left panel). We

then asked whether reducing gravity and muon flux would alter either the timing or amplitude of

these phases.

Gravity intensity has a net significant effect on viral load across time points (χ^2 = 2255.942, 1

d.f., P < 0.001; η_P^2 = 0.058). The mean viral load under μG conditions was (5.78

$\pm 0.06) \cdot 10^4$,·significantly lower than the $(1.02 \pm 0.01) \cdot 10^5$ observed under standard gravity,

representing a 1.8-fold reduction over 44 hpi (Fig. 2, upper right panel: green line *vs* gray

reference; *post hoc* test, P < 0.001). This effect is time-dependent (χ^2 = 1415.456, 19 d.f., P <

0.001; η_P^2 = 0.037), primarily reducing the height of the first peak of viral replication. Muon

radiation flux also has a significant impact on viral replication (χ^2 = 1593.430, 1 d.f., P < 0.001;

η_P^2 = 0.041). Under BBR conditions, the mean viral load reached $(6.36 \pm 0.07) \cdot 10^5$, a 6.3-fold

increase compared to normal muon flux (Fig. 2, lower left panel: green line *vs* gray reference).

The effect of BBR strongly altered the temporal dynamics of OrV accumulation (χ^2 = 3103.138,

19 d.f., P < 0.001; η_P^2 = 0.080), eliminating the biphasic replication pattern: a single peak emerges

around 12 hpi, remains stable until 36 hpi, and then gradually declines.

These environmental effects were not independent. A significant interaction was detected

between gravity and radiation conditions (χ^2 = 1636.075, 1 d.f., P < 0.001; η_P^2 = 0.042), which

was also time-dependent (χ^2 = 1934.595, 19 d.f., P < 0.001; η_P^2 = 0.050). The combined stress of

μG and BBR produces a distinctive pattern resembling the BBR conditions but with a reduction

in viral load similar to the one observed in μG —approximately 1.8-fold lower than the control

throughout the entire developmental period (Fig. 2, lower-right panel: green line *vs* gray

reference; *post hoc* test, P < 0.001).

In summary, μG exerts a negative effect on viral replication, BBR enhances it while altering

infection dynamics, and their combination results in a compounded reduction of viral

accumulation.

FIG 2 Log-fold change of OrV accumulation, relative to the expression of housekeeping gene *cdc-42*, evaluated in inoculated animals reared under the indicated combination of gravity intensity and muon radiation flux conditions. Different green tones represent independent biological replicates (pools of ~300 animals per time point). The solid green lines represent the median trajectory. For the sake of comparison, the gray solid lines represent the OrV dynamics observed in standard laboratory conditions (as in the left upper panel).

Fecundity traits are affected by reduced gravity and radiation

210
OrV infections are typically described as having a mild impact on *C. elegans* fitness (37). To
determine whether this impact might be altered by our environmental treatments, we performed
time-series measurements of three fecundity-related traits —number of viable progeny (*A*),
number of unfertilized eggs (*U*), and number of nonviable embryos (*N*)— in four conditions:
standard environment, μG, BBR, and the combination of μG plus BBR (Fig. S1). These
measurements were done both in non-inoculated animals (control; blue lines and circles) and in
OrV-inoculated ones (OrV; green lines and triangles). All these data were fitted to the generalized
linear mixed model (GLMM) described in the Materials and Methods section. The results from
these analyses are shown in Table 1. For a detailed description of the effect of the different
stresses on *A*, *U* and *N*, we refer to the supplementary text in Supplementary Materials online.
Here, we will focus on the combination of these three traits into a more eco-evolutionary relevant
trait, the per-day reproductive success defined as $R(t) = A(t)/[A(t) + N(t) + U(t)]$ (Fig. 3). Focusing

first on the effects of the environmental conditions in non-inoculated animals, we found that differences in gravity intensity had a significant effect, though of small magnitude (Table 1; $P < 0.001$, $\eta_P^2 = 0.022$), with animals reared at standard gravity having an average R of 0.950 ±0.005 that was reduced to 0.927 ±0.004 in μG (Fig. 3, upper left *vs* upper right panels, blue). The effect of muon radiation flux was conditional to the value of gravity intensity (Table 1; $P = 0.002$, $\eta_P^2 = 0.016$). In μG, R was not affected by differences in muon radiation flux (*post hoc* test, $P = 0.691$) (Fig. 3, upper left *vs* lower right panels, blue). However, in standard gravity conditions the minimization of muon radiation flux was associated with a reduction in average R from 0.962 ±0.007 to 0.939 ±0.006 (*post hoc* test, $P < 0.001$) (Fig. 3, upper left *vs* lower left panels, blue).

FIG 3 Per day reproductive success (R) of OrV-inoculated (green) and non-inoculated (blue) animals under different combinations of environmental stresses. Color triangles and circles rep   ribbon. Nu   r BBR, an   ns, $n = $ 12

R was also negatively affected by infection itself (Table 1; $P < 0.001$, $\eta_P^2 = 0.074$), with

inoculated animals having grand mean *R* of 0.917 ±0.004 and non-inoculated animals of 0.961

±0.005, an effect of moderate magnitude. The effect of inoculation was mediated by gravity

intensity (Table 1; $P < 0.001$, $\eta_P^2 = 0.024$). The mean *R* of inoculated animals exposed to μG

(0.894 ±0.005) was significantly lower than in standard gravity (0.941 ±0.006; *post hoc* test, $P <$

0.001) (Fig. 3, upper left *vs* upper right panels, difference between blue and green). The effect of

inoculation was also mediated by the muon radiation flux (Table 1; $P = 0.004$, $\eta_P^2 = 0.014$). In

this case, *R* of inoculated animals in BBR conditions (0.910 ±0.006) was lower than for animals

inoculated in standard muon radiation flux (0.949 ±0.006; *post hoc* test, $P = 0.020$) (Fig. 3, upper

left *vs* lower left panels, difference between blue and green). Finally, the effect of inoculation

significantly depended on the interaction between the two environmental conditions (Table 1; *P*

< 0.001, $\eta_P^2 = 0.018$). The largest effect on *R* was observed for inoculated animals grown under

μG and standard muon radiation flux (0.866 ±0.007) (Fig. 3, upper left *vs* lower right panels,

difference between blue and green), while the smallest effect was seen for non-inoculated animals

under the same combination of factors (0.97 ±0.01) (green, upper left *vs* lower right panels in Fig.

3).

In conclusion, both environmental stressors impacted reproductive success, though their

effects varied across traits. μG always reduced reproductive success. BBR reduced reproductive

success in standard gravity, and its effects were smaller or absent in μG. In standard conditions,

OrV negatively impacted the animals' reproductive success, especially by increasing the number

of unfertilized eggs. These effects are worsened by the combination of μG and BBR, indicating

that these stresses likely affect different pathways of the infection response.

The observed effects of BBR and viral infection are likely prezygotic

The above statistical descriptions and analyses highlight the effect of the three experimental

factors in the data. However, this sort of analyses fails to provide a mechanistic explanation for

the observed differences. To dig into possible mechanisms, we developed a simple model that

describes the processes from the maturation of germline precursors up to the hatching of eggs to

liberate L1 larvae. The model is described by the set of six ordinary differential equations given

by Eqs. 1 – 6 in Materials and Methods. The model includes five parameters to describe different

processes. Fitting the data shown in Fig. 1 into Eqs. 1 – 6 allows to disentangle whether observed

differences in *A*, *N* and *U* under different environments and infection status can be attributable to

pre- or post-zygotic effects. Table 2 shows the fitting estimates for each condition (see Fig. S2

for illustration of the best fittings). Table S1 shows the results of the three-ways ANOVA model

fitted to the parameter estimates. Gravity intensity itself had no significant effect on any of the

six model parameters. Muon radiation flux had significant effects, of large magnitude, in the rates

of germline precursors differentiation into mature oocytes (δ) as well as in the rate of oocyte migration from the gonads into the spermathecae (η). Animals reared in BBR conditions experienced a reduction in both parameters, in the case of δ from 0.65 ±0.03 d^{-1} to 0.45 ±0.02 d^{-1} (*post hoc* test $P < 0.001$); in the case of η from 0.92 ±0.06 d^{-1} to 0.69 ±0.06 d^{-1} (*post hoc* test $P = 0.008$).

Infection status had a significant effect of large magnitude in δ and η (Table S1; $P < 0.001$, $\eta_P^2 \geq 0.209$ in both cases) and significant but of moderate magnitude in the initial number of germline precursors (G_0) and in spermatocyte to germline precursors initial ratio (α) (Table S1; $P \leq 0.038$, $\eta_P^2 \leq 0.107$ in both cases). Inoculated animals show a reduction in δ from 0.62 ±0.03 d^{-1} to 0.47 ±0.02 d^{-1} (*post hoc* test $P < 0.001$), an increase in η from 0.65 ±0.06 d^{-1} to 0.96 ±0.06 d^{-1} (*post hoc* test $P < 0.001$), a reduction in α from 0.92 ±0.02 to 0.86 ±0.02 (*post hoc* test $P = 0.016$), and an increase in G_0 from 270 ±14 to 309 ±12 (*post hoc* test $P = 0.038$).

Table S1 also shows that some of the effects involved interactions between the three environmental factors. Firstly, δ was affected by gravity intensity in a manner dependent on the muon radiation flux (significant interaction gravity by radiation term; $P = 0.006$, $\eta_P^2 = 0.138$): in standard muon radiation flux and standard gravity $\delta = 0.72$ ±0.04 d^{-1}, but reduced to $\delta = 0.58$ ±0.03 d^{-1} in μG (19.7% reduction; *post hoc* test $P = 0.003$); in BBR conditions and standard gravity $\delta = 0.42$ ±0.03 h^{-1}, statistically undistinguishable from $\delta = 0.47$ ±0.03 d^{-1} in μG (*post hoc* test $P = 0.107$). Secondly, the magnitude of the effect of muon radiation flux on δ and η also depended on the infection status (Table S1; $P \leq 0.001$, $\eta_P^2 \geq 0.188$ in both cases). In the case of δ, non-inoculated animals reared in standard muon radiation flux conditions had $\delta = 0.78$ ±0.04 d^{-1}, while at BBR conditions, $\delta = 0.46$ ±0.03 d^{-1}, representing a significant reduction (*post hoc* test $P < 0.001$); inoculated animals grown at standard muon radiation flux conditions had $\delta = 0.51$ ±0.03 d^{-1}, while at BBR conditions, $\delta = 0.43$ ±0.03 d^{-1}, representing a significant reduction (*post hoc* test $P = 0.036$). In other words, the magnitude of the effect on δ due to muon radiation flux was larger in non-inoculated animals than in inoculated ones (41.6% *vs*. 15.56% reductions). In the case of η, non-inoculated animals reared in standard muon radiation flux had $\eta = 0.6$ ±0.1 d^{-1}, while at BBR conditions, $\eta = 0.68$ ±0.08 d^{-1}, values that were not significantly different (*post hoc* test $P = 0.315$); inoculated animals grown at standard muon radiation flux conditions had $\eta = 1.22$ ±0.08 d^{-1}, while at BBR conditions, $\eta = 0.70$ ±0.08 d^{-1}, representing a significant reduction (*post hoc* test $P < 0.001$). In other words, the magnitude of the effect on η due to muon radiation flux was larger in inoculated animals than in non-inoculated ones (44.0% reduction *vs*. 9.8% nonsignificant increase).

In conclusion, we found that the three environmental factors had significant effects in parameters related to prezygotic processes, with the magnitude and sign of the observed changes

305   depending on the precise combination of factors. In terms of wider effects, OrV infection affected

306   more developmental processes than either muon radiation flux or gravity intensity.

Synergistic effects of µG, BBR and infection in developmental rates

To better understand the effects of these environmental factors in another key

developmental/health metric, we next focused on larval development (Fig. 4A). There was a

significant overall effect of gravitational differences yet of small magnitude ($\chi^2 = 45.734$, 1 d.f.,

$P < 0.001$, $\eta_P^2 = 0.005$), with animals raised in µG being larger than those raised under standard

conditions (mean values across time points: 678 ±4 µm vs 640 ±4 µm) (Fig. 4A, upper left vs

upper right panels, blue). Furthermore, the difference is size increased with time post-hatching

($\chi^2 = 48.942$, 3 d.f., $P < 0.001$; $\eta_P^2 = 0.006$). Muon radiation flux also had a significant overall

effect of small magnitude ($\chi^2 = 28.179$, 1 d.f., $P < 0.001$, $\eta_P^2 = 0.003$), with nematodes exposed

to BBR conditions being larger than those in standard conditions (674 ±4 µm vs 644 ±4 µm) (Fig.

4A, upper left vs lower right panels, blue). The effect magnitude was also time-dependent ($\chi^2 =$

17.747, 3 d.f., $P < 0.001$; $\eta_P^2 = 0.002$). Inoculation had a significant impact as well ($\chi^2 = 20.630$,

1 d.f., $P < 0.001$, $\eta_P^2 = 0.002$), with inoculated animals being smaller than their non-inoculated

counterparts (647 ±4 µm vs 672 ±4 µm). The effect magnitude varied over time, generally

increasing ($\chi^2 = 14.983$, 3 d.f., $P = 0.002$; $\eta_P^2 = 0.002$), though not uniformly across all conditions.

The magnitude of these effects depended on the pairwise interactions of the three factors (χ^2

≥ 10.424, 1 d.f., $P \leq 0.001$, $\eta_P^2 \leq 0.003$ in all combinations) and varied over time ($\chi^2 \geq 12.269$, 3

d.f., $P \leq 0.007$, $\eta_P^2 \leq 0.002$ in all combinations). Focusing first in the interaction between gravity

intensity and muon radiation flux, animals grown in µG under BBR conditions were, on average,

larger than those grown only in µG (702 ±6 µm vs 654 ±5 µm; $post\ hoc$ test, $P < 0.001$) (Fig. 4A,

upper right vs lower right panels, blue) and larger than those grown only under BBR conditions

(702 ±6 µm vs 646 ±6 µm; $post\ hoc$ test, $P < 0.001$) (Fig. 4A , lower left vs lower right panels,

blue lines). Now looking at the interaction between gravity intensity and inoculation, we found

that under standard gravity intensity, inoculated animals were on average smaller than non-

inoculated ones (618 ±6 µm vs 671 ±4 µm; $post\ hoc$ test, $P < 0.001$) (Fig. 4A, upper left panel,

differences between blue and green), while inoculation had no significant effect on growth under

µG conditions (675 ±6 µm vs 673 ±6 µm; $post\ hoc$ test, $P = 0.602$) (Fig. 4A, upper right panel,

differences between blue and green). Finally, exploring the interaction between muon radiation

flux and inoculation, we observed that inoculation had no net effect on animals grown under BBR

conditions (675 ±6 µm vs 673 ±6 µm; $post\ hoc$ test, $P = 1$) (Fig. 4A, lower left panel, differences

between blue and green). Three-way interactions were not statistically significant either

independently ($\chi^2 = 0.225$, 1 d.f., $P = 0.636$) or over developmental time ($\chi^2 = 2.034$, 3 d.f., $P =$

0.565), indicating that the effect of infection was independent on the interaction between the two

abiotic stresses (Fig. 4A, lower right panel, differences between blue and green).

FIG 4 Growth and development of non-inoculated and OrV-inoculated animals under different gravity and radiation conditions. (A) Starting from plates with 400 individuals, photographs were taken at 12, 24, 36, and 48 hours after hatching to assess the effect of the stress conditions in larval development. Solid lines represent the mean and ribbons represent ±1 SEM (number of replicates ranged from 19 to 130). (B) Percentage of males quantified 48 h after hatching under standard ($n = 1606$), μG ($n = 733$), BBR ($n = 3490$), and BBR plus μG ($n = 1640$). Error bars represent ±1 SEM. (C) Animal grown in standard conditions. (D) Individual with bulges

around the body and short length. Extracted from the population of 24 h in BBR conditions; scale bar of 100 µm. (E) Individual defective in molting. Extracted from the population 48 h in BBR plus µG; scale bar of 100 µm.

Populations of *C. elegans* are typically dominated by hermaphrodites, with males arising
infrequently under standard conditions, only once per 1606 individuals in our experiments.
Although rare, males can be biologically significant by enabling outcrossing, which increases
genetic diversity and may facilitate adaptation under stressful conditions (38). Notably, the
frequency of males increased when nematodes were subjected to reduced gravity or BBR. For
instance, we found four males in a population of 733 under µG, 15 out of 3490 under BBR alone,
and five out of 1640 under combined µG and BBR (Fig. 4B). While neither gravity intensity (χ^2
= 2.606, 1 d.f., P = 0.106) nor muon ration flux (χ^2 = 1.431, 1 d.f., P = 0.232) alone significantly
affected the number of males produced, their combination led to a ~5-fold increase in male
frequency compared to standard conditions, a statistically significant yet small effect (χ^2 = 5.315,
1 d.f., P = 0.021, η_P^2 = 0.001).

In addition, under BBR conditions we observed dumpy animals, characterized by a shorter
and stouter body morphology due to defects in structural genes like collagens (3 in a population
of 1843 individuals), as well as other phenotypes that deviate from the standard (Fig. 4C), which
included increased head-length proportion, body bulges (Fig. 4D) and molting defects (Fig. 4E).
Neither of these phenotypes were observed in standard conditions (Binomial test, P < 0.001). A
possible explanation is that lack of radiation might affect genes that are involved in molting, such
as collagens (39).

In conclusion, OrV inoculation mitigates the effect of the two abiotic stresses on
developmental rates, the two abiotic stresses result in increased male frequency and BBR also
causes diverse morphological abnormalities.

DISCUSSION

In this study, we sought to investigate the independent and combined effects of two environmental
stressors (reduced gravity and absence of muon radiation flux) on a biological system comprising
C. elegans and its natural viral pathogen OrV. Organisms evolved under specific gravitational
and radiation conditions at Earth's surface; deviations from these parameters provide
opportunities to understand the constraints these factors impose on physiology and immunity.
Previous work suggested that even partial modifications of Earth-like gravity and radiation levels
could influence immune responses and pathogen–host interactions (7, 40, 41). Consequently, we
focused primarily on reduced muon flux (the major component of cosmic radiation at sea level)

in an underground laboratory, and on simulated reduced gravity using an RPM. Although space

environments include additional stressors such as heavy-ion radiation and solar particle events

(42), our results demonstrate how extreme environments can fundamentally alter host-pathogen

interactions.

We first validated that our reduced-gravity simulation recapitulated previously reported

phenotypes, such as increased intestinal permeability under microgravity (32). Although agar

plate-based RPM experiments do not fully mimic true weightlessness, and surface tension can

interfere with microgravity simulation fidelity (43), we still observed a significant compromise

in intestinal integrity under μG. Furthermore, RPM microgravity simulation in agar plates has

been proved to induce changes in *C. elegans* physiology (31), while liquid culture in a RPM

produces a shear stress that should be acknowledged (44). It is also worth-mentioning that liquid

culture in *C. elegans* has an impact in gene expression (45) and growth (46), which represents a

disadvantage when considering other microgravity simulation systems. Conversely, culturing

nematodes under BBR (*i.e.*, drastically reduced muon flux) partially offset μG-induced intestinal

permeability, but their combination worsens reproductive success, suggesting that microgravity

and BBR elicit overlapping but distinct physiological responses.

Our results confirm that μG and BBR represent stressful environments that affect responses to

viral infections and development (1 - 7, 29). Regarding development and fecundity, the small but

significant increase in male frequency in the combination of the two stresses suggests that off-

Earth conditions may disrupt cellular homeostasis, potentially affecting chromosomal disjunction

as observed with other environmental stresses (47). Increased male frequency could have major

population-level consequences, as increased mating increases genetic diversity through allele

shuffling, bolstering adaptation to stressful conditions (38, 48). Animals grown under μG or BBR

conditions were larger, with their combination producing further increases. These μG results are

in disagreement with previous works, which reported decreased body length in simulated

microgravity (33) and in spaceflight (49), likely due to our experimental design: animals were

preadapted to μG during two generations, grew on solid medium, and experienced different

exposure timing and duration without additional stresses like bleaching synchronization.

Regarding viral infection, OrV-inoculated animals in standard conditions were shorter than non-

inoculated, while it had no effect under other stresses. This confirms early observations of OrV

infection effects on development (50) and consistent with observations in adult nematodes

expressing human hepatitis delta viral antigens (51). Regarding viral infection dynamics, μG had

a small effect in the shape of the virus accumulation curve, reducing the accumulated viral load

over 44 hpi. BBR had major effects in OrV accumulation dynamics, erasing the distinction

between both viral production phases and resulting in a large reduction in accumulated OrV.

Combining both stresses fully changed temporal viral production shape, resembling BBR patterns

but further reducing accumulated viral load. These observations suggest that (*i*) simulated µG conditions may cause minor nematode immune system impairment and (*ii*) BBR-induced stress on nucleic acids stability impacts OrV replication through unknown mechanisms. Our results are consistent with spaceflight-induced immune alterations and herpesvirus reactivation in astronauts (3, 5 - 7, 52).

The molecular mechanisms underlying these effects likely involve multiple conserved pathways. Short-term exposures to µG cause small, reversible *C. elegans* transcriptomic alterations, while prolonged expositions result in larger transgenerational reversible effects (31, 53, 54). These alterations affect longevity-regulating insulin/IGF-1 (55) and sphingolipid signaling pathways (56), neuronal function and/or cellular metabolism mediated by the *daf-16*/FOXO signaling pathway (57), and misregulation of responses to oxidative stress and the antioxidant defense system (33). Some of these pathways are also linked to OrV infection (55), which generates reactive oxygen species (ROS) as part of the immune response, creating oxidative stress (56). Failure to produce ROS antiviral responses would result in enhanced virus accumulation. Our fertility modeling approach revealed effects appear largely attributable to differences in prezygotic parameters, such as oocyte maturation and fertilization rates. This aligns with findings showing altered major sperm protein genes (MSP) regulation in BBR condition (29). MSPs are involved in sperm motility and amoeboid-like crawling (57) and are critical for oocyte and egg development, ovulation, spermathecal valve dilation, and parthenogenesis (58, 59). Given MSP functional diversity, altered *msp* gene expression under BBR conditions likely reflects in our model's prezygotic parameters.

Our study is amongst the first exploring biological implications of muon radiation flux absence on viral infection. While most research focused on cosmic radiation exposure on natural background radiation, the reduction of secondary cosmic rays, including muons, presents an underexplored variable (29). No study has explored radiation absence effects on viral infection progression, despite potential contributions to viral activation in astronauts. Muon radiation has been present during organisms' evolution as a result organisms have developed mechanisms directed to cope with the effects of this type of radiation in the integrity of their genetic material integrity and physiological processes. The negative impact of lack of muon radiation in reproductive success aligns with the emerging theory of radiation as an hormetic agent (11), where doses below and above natural background may harm organisms.

Combined µG and BBR impact creates a singular environment affecting viral infection dynamics. These off-Earth stresses are not isolated but interact to modulate viral reactivation or replication. The immune dysregulation caused by µG, when compounded by the altered radiation environment, demonstrates how environmental factors integrate to produce emergent biological properties not predictable from individual treatments. The systems-level interactions we observed

(where reduced gravity and radiation absence create overlapping yet distinct physiological
responses) highlight the complex dependencies organisms have evolved under Earth's
environmental conditions.

In conclusion, evidence presented here highlights the intricate interplay between reduced
gravity, diminished muon flux and viral infection. Even introducing a couple of off-Earth
conditions (microgravity and below-background radiation) substantially reshape host physiology
and pathogen replication. Our findings suggest that lowered gravity may impair certain *C.*
elegans immune function aspects, allowing modest viral load increases, while the lack of muon
flux alters viral accumulation dynamics more pronouncedly. These stressors create an
environment in which viral activity and host fitness are influenced by overlapping yet distinct
mechanisms. For organisms evolved in Earth's gravitational field and background radiation,
deviations in these parameters (whether in orbital stations, underground conditions, or on other
planetary bodies) can induce notable biological effects. Further research involving a wider range
of radiation types and additional parameters will help clarify the mechanisms behind these
observations and guide strategies to preserve health under non-terrestrial conditions.

MATERIAL AND METHODS

C. elegans strains and culturing

C. elegans was cultured and maintained at 20 °C on nematode growth media (NGM) agar plates
seeded with *Escherichia coli* OP50. ERT54 (*jyIs8[pals-5p::GFP + myo-2p::mCherry]X*), a
transgenic strain with a genetic wild-type (Bristol N2) background that expresses GFP in response
to intracellular infection (60) was used for all experiments. The more susceptible strain SFE2
(*drh-1[ok3495]IV;mjIs228*) was used to produce OrV stocks.

In order to obtain synchronized animal populations, plates with embryos were carefully
washed with M9 buffer to remove larvae and adults but leaving the embryos behind. Plates were
washed again using M9 buffer after 1 h to collect larvae hatched within that time span and
transferred to seeded NGM plates.

A variable number of plates of 500 synchronized L1 animals were used to assess male
frequency. Plates were stored at 20 °C and, two days after hatching, males and total population
were counted using a Zeiss Stemi305 stereomicroscope.

Experimental design and locations

Four different environmental conditions have been tested in this work: standard conditions, μG,
BBR, and the combination of μG plus BBR. Experiments conducted in standard and microgravity
conditions were performed in the Evolutionary Systems Virology laboratory in the Institute for

Integrative Systems Biology (Paterna, Spain). The experiments that included BBR conditions
were carried out in the LSC in three different periods from 2022 to 2024 due to technical issues.

μG was simulated using an RPM (Yuri Gravity GmbH) set to the zero-gravity mode. Values
of G-force were monitored through the RPM software and maintained at approximately 0.001 –
0.002 g throughout the experiments. The RPM was placed inside an incubator at 20 ºC. Animals
used in experiments of μG conditions were acclimated along two generations, with studies done
on the third generation and its progeny. Plates were sealed with parafilm to maintain their
humidity.

BBR experiments were performed in the LSC LAB2400. The measured integrated muon
radiation flux in this facility is ~5×10^{-3} m^{-2} s^{-1} (61). For comparison, the muon radiation flux at
sea level in the Northern hemisphere is about 150 m^{-2} s^{-1} (62). Thermoluminescent dosimeters
sensitive to various sources of radiation are placed at different locations of the underground
laboratory. In 2022 they showed and average dose rate of 0.71 ±0.03 mSv per year in the
underground facilities, compared with the 1.36 ±0.03 mSv per year of the above ground laboratory
(63). Radon radiation from the mountain was reduced to 1 mBq m^{-3} thanks to a good ventilation
system and a Radon Abatement System, a value substantially lower than the 200 – 280 Bq m^{-3}
measured at the surface laboratory (64). Nematodes used in BBR experiments were acclimated
to this condition during two generations prior to experiments.

Fertility assay

L4 larvae were picked individually from a synchronized population and put in separate fresh
plates. Animals were transferred to fresh plates every 24 h until egg laying stopped.
Synchronized animal populations were inoculated after hatching. The number of nonviable (N)
and unfertilized eggs (U) was counted 24 h after the adult was removed from the plate to ensure
all eggs had hatched. Viable progeny (A) was counted 48 h after the adult was removed from the
plate. Reproductive success, R, was then computed as $R = A/(A + N + U)$.

For OrV infection experiments, a population of animals was inoculated one hour after hatching
with the OrV stock described below.

Viral stock preparation, virus quantification and inoculation procedure

For OrV (strain JUv1580_vlc) stock preparation, SFE2 animals were inoculated as previously
described (65). In short, animals were allowed to grow for 5 days and then resuspended in M9
(0.22 M KH$_2$PO$_4$, 0.42 M Na$_2$HPO$_4$, 0.85 M NaCl, 0.001 M MgSO$_4$), let stand for 15 min at room
temperature, vortexed, and centrifuged for 2 min at 400 g. The supernatant was centrifuged twice
at 21,000 g for 5 min and then passed through a 0.2 μm filter. RNA of the resulting viral stock
was extracted using the Viral RNA Isolation kit (NZYTech). The concentration of viral RNA was

then determined by RT-qPCR using a standard curve, and normalized across different stocks (details below). Primers used for RT-qPCRs can be found in Table S2.

For the standard curve, cDNA of JUv1580_vlc was obtained using AccuScript High-Fidelity Reverse Transcriptase (Agilent) and reverse primers at the 3' end of the genome. Approximately 1000 bp of the 3' end of RNA2 were amplified using forward primers containing 20 bp coding the T7 promoter and DreamTaq DNA Polymerase (Thermo Fisher). The PCR products were gel purified using MSB Spin PCRapace (Invitek Molecular) and an *in vitro* transcription was performed using T7 Polymerase (Merck). The remaining DNA was then degraded using DNAse I (Life Technologies). RNA concentration was determined by NanoDrop (Thermo Fisher) and the number of molecules per μL was determined using the online tool EndMemo RNA Copy Number Calculator (https://www.endmemo.com/bio/dnacopynum.php). Primers used for the standard curve can be found in Table S2.

For inoculation experiments, synchronized populations were inoculated by pipetting 60 μL of viral stock on top of the bacterial lawn containing the animals. The normalized inoculum contained $2.6 \cdot 10^7$ copies of OrV RNA2/μL. The efficiency of this viral stock (measured as animals showing activation of the *pals-5p::GFP* reporter at 48 hpi) was 72 ±3 % (mean ±1 SEM, n = 5 plates with 44 – 48 animals per plate).

RNA extractions and RT-qPCRs

Sample preparation and RNA extractions were performed as previously described (65). Synchronized populations of 300 inoculated and control animals were collected at the designated times (from 6 hpi to 44 hpi every 2 h) with PBS-0.05% Tween. Samples of inoculated animals were performed in triplicates. Samples were centrifuged for 2 min at 1350 rpm and the supernatant was discarded. Another two wash steps were performed before flash-freezing the samples in liquid N_2. 500 μL of Trizol (Invitrogen) were added to the nematode pellet and the pellet was disrupted by following five cycles of freeze-thawing and five cycles of 30 seconds of vortex followed by 30 seconds of rest. 100 μL of chloroform were then added and the tubes were shaken for 15 s and let rest for 2 min. Samples were centrifuged for 15 min at 11,000 g at 4 °C and the top layer containing the RNA was then mixed with the same volume of 100% ethanol. The sample was then loaded into RNA Clean & Concentrator columns (Zymo Research) and the rest of the protocol was followed according to manufacturer instructions.

RT-qPCRs were performed using Power SYBR Green PCR Master Mix (Applied Biosystems) on an ABI StepOne Plus Real-time PCR System (Applied Biosystems). 10 ng of total RNA were loaded and a comparison with the endogenous gene *cdc-42* was used for relative OrV quantifications. Primers used for RT-qPCRs can be found in Table S2.

Body size measurements and analysis of abnormal phenotypes

Synchronized animals were collected at 12, 24, 36, and 48 h, plates were washed with M9 buffer to collect nematodes and another M9 wash was made to remove bacteria from the media. Animals were mounted in a 4% agarose pad with 20% 100 mM azide and photographed using a Leica Thunder DMi8 microscope with DFC9000 GTC sCMOS camera and objective HC PL APO 40×/0.95 CORR PH2 (Leica Microsystems). Body length was measured using the segmented line tool in ImageJ (66).

The photographs employed to measure body size were also used to describe abnormal phenotypes.

Intestinal permeability assay

The intestinal permeability assay was performed as previously described with minor modifications (67). Synchronized nematodes were washed after 48 h with S buffer (6.5 mM K_2HPO_4, 43.5 mM KH_2PO_4, 100 mM NaCl) and suspended in a 1:1 solution of 1 mL of M9 buffer mixed with 0.05 g of erioglaucine disodium salt (Thermo Scientific) and 1 mL of *E. coli* OP50 suspended in S buffer, and gently rocked for 3 h.

Animals were washed carefully three times with 10 mL of M9 buffer. Animals were mounted on microscopic slides on top of a 4% agarose pad with 20% 100 mM azide. Photographs were taken using a Leica MZ10F stereomicroscope with Flexacam C3 camera and with a Leica DMi1 microscope with Flexacam C1 camera. Quantifications were randomized by assigning random names to photographs to avoid observer bias. A positive control was prepared to help us to classify the different phenotypes. Positive control was prepared by following the same assay except that animals were grown in NGM plates with 0.8 mM H_2O_2 during 48 h, as previously described (68). Four-hundred nematodes were used in each condition.

A model of *C. elegans* fecundity and parameter estimation

Fig. 5 describes a simple fecundity model for sexually mature adults at L4 developmental stage, in which a fixed number of spermatocyte cells (*S*) have already been produced and stored in the spermathecae. Germline precursor cells (*G*) differentiate into mature oocytes (*O*) at rate δ_G. This rate is assumed to be constant $\delta_G = \delta$ as far as germline cells exist ($G(t) > 0$) and zero otherwise. These oocytes migrate from the distal to the proximal gonads at rate η until reaching the spermathecae where, if spermatocytes are available, are fecundated with an efficiency function ψ that depends on the amount of *S* present in the spermatheca at time *t*: $\psi(S) = (1 - \gamma)S/(\gamma + S) \in [0,1]$, where γ is a dimensionless parameter which modulates the value of the efficiency, resulting in fertilized and unfertilized (*U*) eggs. After passing through the uterus, the eggs are deposited in the agar, where, if they have been fecundated, a fraction μ will develop into L1 larvae (*i.e*, the

592 viable progeny, A). A fraction $1 - \mu$ will stop embryonic development, resulting in nonviable

593 embryos (N). N and U eggs will simply remain in the plates and are easily distinguishable.

Parameter	Description
δ_G	Rate of germline precursor cells differentiation
η	Rate of oocytes migration
Ψ	Fertilization efficiency
μ	Fraction of fertilized eggs viability
α	Spermatocytes to germline initial ratio

Variable	Description
G	Germline precursor cells
O	Oocytes
S	Spermatocytes
U	Unfertilized eggs
A	Viable progeny
N	Non-viable eggs

FIG 5 Parameters, variables and schematic representation of the fecundity model described by Eq. 1 – 6.

596 These processes can be written according to the following set of ordinary differential

597 equations,

$$\frac{dG}{dt} = -\delta_G, \qquad \text{Eq. 1}$$

$$\frac{dO}{dt} = \delta_G - \eta O, \qquad \text{Eq. 2}$$

$$\frac{dS}{dt} = -\eta \psi O, \qquad \text{Eq. 3}$$

$$\frac{dA}{dt} = \eta \mu \psi O, \qquad \text{Eq. 4}$$

$$\frac{dN}{dt} = \eta(1 - \mu)\psi O, \text{and} \qquad \text{Eq. 5}$$

$$\frac{dU}{dt} = \eta(1 - \psi)O, \qquad \text{Eq. 6}$$

being A, N, and U are the three experimental observables (Eqs. 4 – 6). All variables were

normalized by the initial number of non-differentiated germinal cells G_0. Model parameters δ, η

and γ represent prezygotic stages, while μ represents a postzygotic event. Furthermore, we define

the dimensionless parameter $\alpha = S_0/G_0$, which for a hermaphrodite in which the spermatocytes

will be the limiting factor for asexual reproduction will be $\alpha < 1$.

We used a genetic algorithm to estimate the vector of parameters (δ, η, μ, γ, α, G_0) better

fitting the experimental data for every condition. The size of the population of vectors of

parameters was fixed at 600, and the number of generations was set to 10^4. The exploration of

the parameter space was refined through successive generations. The cost function ($\mathcal{F}$) to

minimize was the logarithm of the square sum of discrepancies:

$$\mathcal{F} = \log \sum_{i=1}^{T} \left[\left(A_i - \hat{A}_i \right)^2 + \left(N_i - \hat{N}_i \right)^2 + \left(U_i - \hat{U}_i \right)^2 \right]. \qquad \text{Eq. 7}$$

Here, $\{t_i, A_i, N_i, U_i\}_{i=1}^{T}$ represent the experimental data obtained for T different times and $\hat{A}_i$, $\hat{N}_i$

and $\hat{U}_i$ denote the estimated values for the different species obtained with the mathematical model

for a given set of parameters.

Within each generation, the 5% of the parameters with the lowest $\mathcal{F}$ values were designated as

the elite population, remaining constant for the next generation. The next-best 80% underwent

parameter crossover, while the remaining 15%, representing the least favorable parameters,

experienced random mutations. The parameter space was subdivided into a logarithmic scale for

parameters δ, η and γ to ensure a more balanced exploration. The model was fitted to every

fertility assay data, resulting in a vector of estimated parameters per experimental replicate and

condition.

Statistical analyses

Data were fitted to generalized linear mixed models (GLMM) with gravity intensity (standard *vs*

μG), radiation level (standard *vs* BBR) and infection status (non-inoculated *vs* inoculated)

incorporated as fixed orthogonal factors, whereas individual animals (subjects) and days-post

inoculation (repeated measures per subject) were both modeled as random factors. In the case of

counts (A, N and U), a Poisson distribution with a log-link function was used; in the case of R and

body size, a Normal distribution with identity-link function was used.

Intestinal permeability and the frequency of males per population data were fitted to logistic

regression models using generalized linear models (GLM) with gravity intensity and radiation

level incorporated as fixed orthogonal factors and assuming a Binomial distribution and a logit-

link function.

Viral load data were fitted to a GLMM with gravity intensity and radiation level incorporated

as fixed orthogonal factors, whereas replicate plates (subjects) and hpi (repeated measures per

subject) were both modeled as random factors, a Gamma distribution with a log-link function was

used.

Estimated parameters of the model described by Eqs. 1 – 6 were independently fitted to

ANOVA models with gravity intensity, radiation level and infection status incorporated as fixed

orthogonal factors. Errors were assumed to be Normal.

In all cases, pairwise *post hoc* comparisons were done with the sequential Bonferroni method.

The magnitude of the effects was calculated using the η_P^2 statistic, *i.e.*, the proportion of variance

in the dependent variable that can be attributed to each model factor after accounting for the
variance explained by other factors in the model. Typically, values of $0.06 \leq \eta_P^2 < 0.15$ are
considered as medium size effects and $\eta_P^2 \geq 0.15$ are as large magnitude effects.

These analyses were done with SPSS version 29.0.2.0 (IBM Corporation, Armonk, NY).

643
644

ACKNOWLEDGEMENTS
We thank Francisca de la Iglesia (I2SysBio) and Rebecca Hernández-Antolín (LSC) for excellent
technical support. We are deeply grateful to Carlos Peña-Garay (LSC) for his commitment and
dedication to research in subterranean biology, as well as his constant material and emotional
support. Many computations were performed on the HPC cluster Garnatxa at I2SysBio (CSIC-
UV).

FUNDING
This study was supported by ESA contract 4000135960/21/NL/GLC/my, grant PID2022-
136912NB-I00 funded by MCIN/AEI/10.13039/501100011033 and by "ERDF a way of making
Europe", grant CIPROM/2022/59 funded by Generalitat Valenciana, and grant PB-02-22 from
ICTS Laboratorio Subterráneo de Canfranc to S.F.E. V.G.C. was supported by grant FJC2021-
047264-I funded by MCIN/AEI/10.13039/501100011033 and by NextGenerationEU/PRTR.
E.G.L. was supported by grant FPU21/00410 funded by MCIN/AEI/10.13039/501100011033 and
by "ESF invest in your future". J.C.M-S. was supported by grant ACIF/2021/296 from
Generalitat Valenciana. The funders played no role in study design, data collection, analysis and
interpretation of data, or the writing of this manuscript.

AUTHOR CONTRIBUTIONS
A.V.G.: conceptualization, data curation, formal analysis, investigation, methodology, validation,
writing-original draft, writing-review and editing; V.G.C.: conceptualization, investigation,
methodology, supervision, validation, writing-review and editing; J.C.M.S.: conceptualization,
formal analysis, investigation, methodology, software, writing-review and editing; E.G.L.:
investigation, writing-review and editing; R.G.: conceptualization, supervision, writing-review
and editing; S.F.E.: conceptualization, data curation, formal analysis, funding acquisition, project
administration, supervision, writing-original draft, writing-review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work
performed therein.

DATA AVAILABILITY

The datasets generated during the current study are available in the Zenodo repository, https://doi.org/10.5281/zenodo.13869555.

ADDITIONAL FILES

Supplementary materials are available online at https://git.csic.es/sfelenalab/spaceworms/-/blob/070697e9dc66061d671094f6906a4a18f25b5e6b/Supplementary_Materials.pdf.

REFERENCES

1. Crucian B *et al*. 2013. Immune system dysregulation occurs during short duration spaceflight on board the space shuttle. J Clin Immunol 33:456–465.

2. Crucian B *et al*. 2015. Alterations in adaptive immunity persist during long-duration spaceflight. npj Microgravity **1**:15013.

3. Bigley AB *et al.* 2019. NK cell function is impaired during long-duration spaceflight. J Appl Physiol 126:842–853.

4. Agha NH *et al*. 2020. Salivary antimicrobial proteins and stress biomarkers are elevated during a 6-month mission to the International Space Station. J Appl Physiol 128:264–275.

5. Rooney BVB, Crucian BE, Pierson DL, Laudenslager ML, Mehta SK. 2019. Herpes virus reactivation in astronauts during spaceflight and its application on Earth. Front Microbiol 10:16.

6. Mehta SK *et al*. 2014. Multiple latent viruses reactivate in astronauts during Space Shuttle missions. Brain Behavior Immunity 41:210–217.

7. Mehta SK *et al*. 2017. Latent virus reactivation in astronauts on the international space station. npj Microgravity 3:11.

8. Bonomi G, Checchia P, D'Errico M, Pagano D, Saracino G. 2020. Applications of cosmic-ray muons. Prog Particle Nucl Phys 112:103768.

9. Zhao L *et al.* 2020. The determinant of DNA repair pathway choices in ionizing radiation-induced DNA double-strand breaks. BMC Res Int 2020:4834965.

10. Ceccaldi R, Rondinelli B, D'Andrea AD. 2016. Repair pathway choices and consequences at the double-strand break. Trends Cell Biol **26**:52–64.

11. Baldwin J, Grantham V. 2015. Radiation hormesis: historical and current perspectives. J Nucl Med Technol **43**:242–246.

12. Brenner S. 1974. The genetics of *Caenorhabditis elegans*. Genetics 77:71–94.

13. Kim W, Underwood RS, Greenwald I, Shaye DD. 2018. OrthoList 2: A New comparative genomic analysis of human and *Caenorhabditis elegans* genes. Genetics 210:445–461.

14. Olsen A, Vantipalli MC, Lithgow GJ. 2006. Lifespan extension of *Caenorhabditis elegans* following repeated mild hormetic heat treatments. Biogerontology 7:221–230.

15. Aprison EZ, Ruvinsky I. 2014. Balanced trade-offs between alternative strategies shape the
response of *C. elegans* reproduction to chronic heat stress. PLoS ONE 9:e105513.

16. Begasse ML, Leaver M, Vazquez F, Grill SW, Hyman AA. 2015. Temperature dependence of
cell division timing accounts for a shift in the thermal limits of *C. elegans* and *C. briggsae*.
Cell Rep 10:647–653.

17. Rodriguez M, Snoek LB, Riksen JAG. Bevers RP, Kammenga JE. 2012. Genetic variation for
stress-response hormesis in *C. elegans* lifespan. Exp Gerontol 47:581–587.

18. Jovic K *et al*. 2017. Temporal dynamics of gene expression in heat-stressed *Caenorhabditis*
elegans. PLoS ONE 12:e0189445.

19. Rohlfing AK, Miteva Y, Hannenhalli S, Lamitina T. 2010. Genetic and physiological activation
of osmosensitive gene expression mimics transcriptional signatures of pathogen infection in
C. elegans. PLoS ONE 5:e9010.

20. Prithika U, Deepa V, Balamurugan K. 2016. External induction of heat shock stimulates the
immune response and longevity of *Caenorhabditis elegans* towards pathogen exposure. Innate
Immunity 22:466–478.

21. Leroy M, Mosser T, Manière X, Alvarez D,. Matic I. 2012. Pathogen-induced *Caenorhabditis*
elegans developmental plasticity has a hormetic effect on the resistance to biotic and abiotic
stresses. BMC Evol Biol 12:187.

22. Castiglioni VG, Elena SF. 2024. Orsay virus infection increases *Caenorhabditis elegans*
resistance to heat-shock. Biol Lett 20:20240278.

23. Panek J *et al*. 2020. A cullin-RING ubiquitin ligase promotes thermotolerance as part of the
intracellular pathogen response in *Caenorhabditis elegans*. Proc Natl Acad Sci USA
117:7950–7960.

24. De Souza EA, Thompson MA, Taylor RC. 2023. Olfactory chemosensation extends lifespan
through TGF-β signaling and UPR activation. Nat Aging 3:938–947.

25. Le Pen J *et al*. 2018. Terminal uridylyltransferases target RNA viruses as part of the innate
immune system. Nat Struct Mol Biol 25:778–786.

26. Ashe A *et al*. 2013. A deletion polymorphism in the *Caenorhabditis elegans* RIG-I homolog
disables viral RNA dicing and antiviral immunity. eLife 2:e00994.

27. Félix MA *et al*. 2011. Natural and experimental infection of *Caenorhabditis* nematodes by
novel viruses related to nodaviruses. PLoS Biol 9:e1000586.

28. Huang Y. *et al*. 2021. Heat stress reduces the susceptibility of *Caenorhabditis elegans* to Orsay
virus infection. Genes 12:1161.

29. Van Voorhies WA, Castillo HA, Thawng CN, Smith GB. 2020. The phenotypic and
transcriptomic response of the *Caenorhabditis elegans* nematode to background and below-
background radiation levels. Front Public Health 8:581796.

30. Morciano P. *et al*. 2018. Effects of reduced natural background radiation on *Drosophila*

melanogaster growth and development as revealed by the FLYINGLOW program. J Cell Physiol 233:23–29.

31. Tee LF. *et al.* 2017. Effects of simulated microgravity on gene expression and biological phenotypes of a single generation *Caenorhabditis elegans* cultured on 2 different media. Life Sci Space Res 15:11–17.

32. Liu H, Guo D, Kong Y, Rui Q, Wang D. 2019. Damage on functional state of intestinal barrier by microgravity stress in nematode *Caenorhabditis elegans*. Ecotox Environ Safety 183:109554.

33. Zhao L, Rui Q, Wang D. 2017. Molecular basis for oxidative stress induced by simulated microgravity in nematode *Caenorhabditis elegans*. Sci Total Environ 607–608:1381–1390.

34. Zhou Y, Chen H, Zhong W, Tao YJ. 2024. Collagen and actin network mediate antiviral immunity against Orsay virus in *C. elegans* intestinal cells. PLoS Pathog 20:e1011366.

35. Martel J. *et al.* 2022. Gut barrier disruption and chronic disease. Trends Endocrinol Metabol 33:247–265.

36. Castiglioni VG. *et al.* 2024. Story of an infection: viral dynamics and host responses in the *Caenorhabditis elegans*-Orsay virus pathosystem. Sci Adv 10:eadn5945.

37. Félix MA, Wang D. 2019. Natural viruses of *Caenorhabditis* nematodes. Annu Rev Genet 53:313–326.

38. Morran LT, Cappy BJ, Anderson JL, Phillips PC. 2009. Sexual partners for the stressed: facultative outcrossing in the self-fertilizing nematode *Caenorhabditis elegans*. Evolution 63:1473–1482.

39. Johnstone IL. 2000. Cuticle collagen genes. Expression in *Caenorhabditis elegans*. Trends Genet 16:21–27.

40. Hicks J, Olson M, Mitchell C, Juran CM, Paul AM. 2023. The impact of microgravity on immunological states. Immunohorizons 7:670–682.

41. Shearer WT, Zhang S, Reuben JM, Lee BN, Butel JS. 2005. Effects of radiation and latent virus on immune responses in a space flight model. J Allergy Clin Immunol 115:1297–1303.

42. Chancellor JC. *et al.* 2018. Limitations in predicting the space radiation health risk for exploration astronauts. npj Microgravity 4:8.

43. Adenle AA, Johnsen B, Szewczyk NJ. 2009. Review of the results from the International *C. elegans* first experiment (ICE-FIRST). Adv Space Res 44:210–2016.

44. Wuest SL, Stern P, Casartelli E, Egli M. 2017. Fluid dynamics appearing during simulated microgravity using random positioning machines. PLoS ONE 12:e0170826.

45. Çelen I, Doh JH, Sabanayagam CR. 2018. Effect of liquid cultivation on gene expression and phenotype of *C. elegans*. BMC Genomics 19:562.

46. Lev I, Bril R, Liu Y, Ceré LI, Rechavi O. 2019. Inter-generational consequences for growing *Caenorhabditis elegans* in liquid. Phil Trans R Soc B 374:20180125.

47. Goswami PC, Roti Roti JL, Hunt CR. 1996. The cell cycle-couple expression of topoisomerase
IIα during S phase is regulated by mRNA stability and is disrupted by heat shock or ionizing
radiation. Mol Cell Biol 16:1500–1508.

48. Slowinski SP. *et al*. 2024. Outcrossing in *Caenorhabditis elegans* increases in response to food
limitation. Ecol Evol 14:e11166.

49. Higashibata A. *et al*. 2016. Microgravity elicits reproducible alterations in cytoskeletal and
metabolic gene and protein expression in space-flown *Caenorhabditis elegans*. npj
Microgravity 2:15022.

50. Melero I. *et al.* 2025. The interplay between *Caenorhabditis elegans* larval development and
Orsay virus infection. bioRxiv doi: 10.1101/2025.05.23.655711

51. Lee LW, Chang TY, Lo HW, Lo SJ. 2011. Hepatitis D antigens cause growth retardation and
brood-size reduction in *C. elegans*. Front Biosci 3:380–390.

52. Lv H. *et al*. 2023. Microgravity and immune cells. J R Soc Interface 20:20220869.

53. Willis CRG. *et al*. 2020. Comparative transcriptomics identifies neuronal and metabolic
adaptations to hypergravity and microgravity in *Caenorhabditis elegans*. iScience 23:101734.

54. Çelen I, Jayasinghe A, Doh JH, Sabanayagam CR. 2023. Transcriptomic signature of the
simulated microgravity response in *Caenorhabditis elegans* and comparison to spaceflight
experiments. Cells 12:270.

55. Duxbury EML. *et al*. 2024. Reduced insulin/IGF-1 signaling upregulates two anti-viral
immune pathways, decreases viral load and increases survival under viral infection in *C.
elegans*. GeroScience 46:5767–5780.

56. Van der Hoeven R, McCallum KC, Cruz MR, Garsin DA. 2011. Ce-Duox1/BLI-3 generated
reactive oxygen species trigger protective SKN-1 activity via p39 MAPK signaling during
infection in *C. elegans*. PLoS Pathog 7:e1002453.

57. L'Hernault WE 1997. Spermatogenesis. In: Riddle, D. L. (ed). *C. elegans II*. Cold Spring
Harbor, NY: Cold Spring Harbor Laboratory Press. pp. 271-94.

58. McGovern M, Yu L, Kosinski M, Greenstein D, Savage-Dunn C. 2007. A role for sperm in
regulation of egg-laying in the nematode *C. elegans*. BMC Develop Biol 7:41.

59. Kuwabara PE. 2003. The multifaceted *C. elegans* major sperm protein: an ephrin signaling
antagonist in oocyte maturation. Genes Develop 17:155–161.

60. Bakowski MA. *et al*. 2014. Ubiquitin-mediated response to microsporidia and virus infection
in *C. elegans*. PLoS Pathog 10:e1004200.

61. Trzaska WH, *et al*. 2019. Cosmic-ray muon flux at Canfranc Underground Laboratory. Eur
Phys J C 79:721.

62. Workman RL. *et al*. 2022. The review of particle physics. Prog Theor Exp Phys 2022:083C01.

63. Hernández-Antolín R, Cid-Barrio L, Peña-Garay C. 2024. Canfranc biology platform:
exploring life in cosmic silence. Front Phys 12:1397799.

64. Pérez-Pérez J. *et al.* 2022. Radon mitigation applications at the Laboratorio Subterráneo de
Canfranc (LSC). Universe 8:112.

65. Castiglioni VG. *et al*. 2024. Experimental evolution of an RNA virus in *Caenorhabditis*
elegans. Infect Genet Evol 123:105623.

66. Schindelin J. *et al*. 2012. Fiji: an open-source platform for biological-image analysis. Nat Meth
9:676-682.

67. Kissoyan KAB. *et al.* 2019. Natural *C. elegans* microbiota protects against infection via
production of a cyclic lipopeptide of the viscosin group. Curr Biol 29:1030-1037.e5.

68. Balaguer F. *et al*. 2023. *Bifidobacterium animalis* subsp. *lactis* BPL1[TM] and its lipoteichoic
acid modulate longevity and improve age/stress-related behaviors in *Caenorhabditis elegans*.
Antioxidants 12:2107.

TABLE 1. Results of the fecundity data fitting to the GLMM testing for the effect of gravity intensity, muon radiation flux and OrV infection. Significant effects of small magnitude are highlighted in light gray, while significant effects of medium and large magnitude are highlighted in dark gray.

Source of variation	d.f.	Unfertilized eggs (U)			Nonviable embryos (N)			Progeny (A)			Reproductive success (R)		
		F	P	η_P^2	F	P	η_P^2	F	P	η_P^2	F	P	η_P^2
Model	7, 592	3.113	0.003	0.036	11.702	< 0.001	0.122	13.129	< 0.001	0.134	19.737	< 0.001	0.189
Gravity intensity	1, 592	1.923	0.166	0.003	3.825	0.051	0.006	22.538	< 0.001	0.037	13.221	< 0.001	0.022
Muon radiation flux	1, 592	1.122	0.290	0.002	5.010	0.026	0.008	0.410	0.522	0.001	0.425	0.515	0.001
Infection status	1, 592	3.864	0.050	0.006	9.244	0.002	0.015	19.131	< 0.001	0.031	47.457	< 0.001	0.074
Gravity × Radiation	1, 592	1.866	0.172	0.003	23.054	< 0.001	0.037	21.024	< 0.001	0.034	9.599	0.002	0.016
Gravity × Infection	1, 592	1.968	0.161	0.003	12.585	< 0.001	0.021	13.209	< 0.001	0.022	14.810	< 0.001	0.024
Radiation × Infection	1, 592	1.624	0.203	0.003	15.129	< 0.001	0.025	1.097	0.295	0.002	8.321	0.004	0.014
Gravity × Radiation × Infection	1, 592	2.020	0.156	0.003	0.259	0.611	0.000	6.689	0.010	0.011	10.964	< 0.001	0.018

TABLE 2. Estimation of model parameters and initial condition. Values represent the mean of the replicates (±1 SD). See Fig. S2 for the fitting to the experimental data.

Gravity intensity	Muon radiation flux	Infection status	G_0	α	$\delta\,(\text{d}^{-1})$	$\eta\,(\text{d}^{-1})$	γ	μ
Standard	Standard	non-inoculated	252.773 ±32.217	0.936 ±0.011	0.891 ±0.215	0.584 ±0.235	0.015 ±0.011	0.989 ±0.003
Standard	Standard	inoculated	303.236 ±113.438	0.912 ±0.129	0.543 ±0.126	1.199 ±0.497	0.004 ±0.005	0.988 ±0.005
μG	Standard	non-inoculated	277.392 ±21.302	0.938 ±0.049	0.667 ±0.166	0.659 ±0.103	0.004 ±0.003	0.986 ±0.014
μG	Standard	inoculated	292.661 ±84.967	0.824 ±0.100	0.486 ±0.138	1.242 ±0.329	0.029 ±0.036	0.970 ±0.028
Standard	BBR	non-inoculated	293.401 ±27.615	0.924 ±0.068	0.416 ±0.072	0.676 ±0.220	0.013 ±0.020	0.974 ±0.008
Standard	BBR	inoculated	356.747 ±66.461	0.844 ±0.105	0.417 ±0.075	0.764 ±0.282	0.019 ±0.019	0.983 ±0.012
μG	BBR	non-inoculated	257.598 ±65.897	0.888 ±0.095	0.495 ±0.084	0.689 ±0.438	0.012 ±0.018	0.978 ±0.015
μG	BBR	inoculated	283.136 ±57.662	0.853 ±0.103	0.451 ±0.128	0.641 ±0.197	0.028 ±0.033	0.975 ±0.014

G_0: Germline precursor cells at $t = 0$; α: germline precursor cells to spermatocytes initial ratio; δ: rate of germline precursor cells differentiation into mature oocytes; η: rate of oocytes migration from the distal gonads to the spermathecae; γ: scale parameter of the fecundation efficiency function; μ: fraction of a fecundated eggs developing into an L1 larvae ($1 - \mu$ would be the fraction of a fecundated eggs which abort its development).