A Targeted Reference Database for Improved Analysis

of Environmental 16S rRNA Oxford Nanopore

Sequencing Data

Melcy Philip[1], Tonje Nilsen[1] Sanna Majaneva[2], Ragnhild Pettersen[2], Morten Stokkan[3], Jessica

Louise Ray[4], Nigel Keeley[5], Knut Rudi[1], Lars-Gustav Snipen[1]

[1]Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life

Sciences, Ås (formerly known as viken), Norway

[2]Akvaplan-niva, Framsenteret, P.O. Box 6606, Stakkevollan, 9296 Tromsø, Norway

[3]STIM department Bergen, Thormøhlens gate 55, 5006 Bergen, Norway

[4]Aqua Kompetanse AS, Storlavika 7, 7770 Flatanger, Norway

[5]Institute of Marine Research, Tromsø, Norway

Corresponding author: Melcy Philip, Faculty of Chemistry, Biotechnology and Food Science,

Norwegian University of Life Sciences, Christian Magnus Falsens vei 18, 1433, Ås (formerly

known as viken), Norway (philip.melcy@nmbu.no and melcy.p.j@gmail.com)

Running Title: Targeted Database for Enhanced 16S ONT Analysis

Abstract

The Oxford Nanopore Technologies (ONT) sequencing platform is compact and efficient, making it suitable for rapid biodiversity assessments in remote areas. Despite its long reads, ONT has a higher error rate compared to other platforms, necessitating high-quality reference databases for accurate taxonomic assignments. However, the absence of targeted databases for underexplored habitats, such as the seafloor, limits ONT's broader applicability for exploratory analysis.

To address this, we have developed a strategy to build environmentally-targeted databases to enhance 16S rRNA gene(16S) analysis using ONT. We start by using Illumina short-read data to create a database of full-length or near full-length 16S sequences from seafloor samples. Initially, amplicons are mapped to the SILVA database, with matches added to our database. Unmatched amplicons are reconstructed using METASEED and Barrnap methodologies with amplicon and metagenome data. Finally, if the previous strategies did not succeed, we included the short-read sequences in the database. This resulted in AQUAeD-DB, which contains 14 545 16S sequences clustered at 95% identity. Comparative database analysis and relative abundance correlation studies reveal that AQUAeD-DB provides consistent results for both Illumina and Nanopore read assignments (median correlation coefficient: 0.50), whereas a standard database showed a substantially weaker correlation. These findings also emphasis its potential to recognize both high and low-abundance taxa, which could be key indicators in environmental studies. This work highlights the necessity of targeted databases for environmental analysis, especially for ONT-based studies, and lays foundations for future extension of the database.

42   **Keywords:** 16S rRNA, metagenomics, seafloor sediment, database, Nanopore, ONT, Illumina,

43   biodiversity assessment

Introduction

The Earth's surface is primarily covered by oceans, yet we know little about the seafloor, or

its microbiota (Canals et al., 2021). Microorganisms play a key role in biogeochemical cycling

in the global ocean thus the potential importance of microbiota to healthy ecosystem function

cannot be overstated (Madsen, 2011). Many aspects of our understanding of soft-sediment

microbiota is based on short read amplicon sequencing of the 16S rRNA gene V3 and V4

regions (Ghate et al., 2021; Hoshino et al., 2020; Yang et al., 2022). Advancements in long read

sequencing, including Oxford Nanopore Technology (ONT), have enabled rapid analysis of

environmental microbiota (Zorz et al., 2023). However, the relatively high error rate of ONT

sequencing technology remains a persistent challenge, rendering ONT strategies dependent

on reference databases (Amarasinghe et al., 2020). For effective and robust use of ONT

technology that generates reads spanning nearly the entire 16S gene, databases of nearly full-

length 16S rRNA gene sequences with comprehensive representation of habitat-relevant taxa

are needed.

The most widely used 16S databases are the curated Greengenes and SILVA databases, as well

as the non-curated NCBI 16S rRNA database (Federhen, 2011; McDonald et al., 2024; Yilmaz

et al., 2013) . These databases are limited by coverage despite the large numbers of reference

sequences that they house. For instance, the 16S component of the GenBank database

consists of 43% chimeric sequences, the presence of which may confound accurate taxonomic

assignments (DeSantis et al., 2006). Furthermore, despite the use of quality control tools,

public 16S reference databases still contain 0.2% to 2.5% misidentified species, leading to

taxonomic misassignments (Kozlov et al., 2016). In addition, there is a major bias towards

characterization of human-associated and highly abundant bacteria, while most of

environmental microbial diversity remains uncharacterized (Yang & Iwasaki, 2014). In this

context, we fail to notice potentially low-abundant microorganisms, such as nitrifying

bacteria, anammox bacteria and methanotrophs, that may be crucial for reflecting the health

of the environment (Marlow et al., 2014; Patil et al., 2021; Wu et al., 2019). Benthic microbes,

in particular, are underrepresented in databases and our understanding of them is constrained

by very few studies (Tatsuhiko Hoshinoa, 2020). The current scenario thus emphasises the

need for a specialised database for seafloor microorganisms, since environmental microbial

communities are often missing from current databases.

Studies reveal that employing targeted databases can improve the accuracy and coverage of

amplicon-based microbial profiling (Dueholm et al., 2022; Meola et al., 2019; Overgaard et al.,

2022). Although ONT sequencing technology is advancing, there are only few bioinformatics

tools for ONT 16S profiling (Ciuffreda et al., 2021; Santos et al., 2020; Sissel et al., 2015) apart

from the workflow offered by Oxford Nanopore. There are several ongoing efforts to improve

taxonomic assignments within ONT-driven studies, with the EMU tool showing promise as an

extensively used approach (Curry et al., 2022; Rodríguez-Pérez et al., 2020; Zorz et al., 2023).

However, all these tools, require a good reference database for improved taxonomic

assignment. This underlines the necessity of an environment-specific or specialized reference

databases and the associated potential benefits. (F. Escapa et al., 2020; Henderson et al.,

2019; Nierychlo et al., 2020; Ritari et al., 2015). Studies have focused on achieving high

resolution using Nanopore, given its longer reads covering most of the 16S gene, whereas our

study aims to capture the broadest possible diversity from environmental samples (Benítez-

Páez et al., 2016; Lemoinne et al., 2023).

The aim of this work was to build a targeted 16S database for seafloor sediment environments

to facilitate full length 16S (v2-v9) ONT sequencing for microbial profiling and prediction of

ecosystem status. We have called this database "AQUAeD-DB" in acknowledgment of the

research project "AQUAeD" which provided environmental samples for sequencing analysis

(see Acknowledgements). Our strategy (Fig. 1) makes use of short-read amplicon data in

combination with existing databases, as well as whole genome shotgun sequencing in

combination with our recently developed method METASEED (Philip et al., 2024). We also

assess and present the effectiveness of this targeted database in comparison to the existing

tools and databases for taxonomic assignment of 16S ONT amplicons.

Fig. 1 The16S database generation. a) The seafloor samples used in this study were collected

from various sites along the Norwegian coast. b) The samples for database generation were

sequenced using Illumina, providing shotgun and 16S rRNA amplicon reads, c) Various

methods are used to collect or rebuild longer 16S rRNA sequences, which are then graded

into different quality levels, from Q1 (best) to Q5 (poorest). These sequences are added to our

reference database e) and used to for taxonomic assignment of ONT sequences amplified

106 from   geographically   proximal   seafloor   sediment   material.

Materials and Methods

Data

In this study, we used total of 182 marine sediment samples collected along a north-south

gradient along the Norwegian coast as mentioned previously (Nilsen et al., 2024; Pettersen et

al., 2022). The data was randomly divided into two subsets: subset A consisted of 94 samples

and was used for the database construction; subset B consisted of 88 samples and was used

for database comparison.

Subset A was subject to Illumina sequencing, both 16S amplicon and whole genome

(metagenome) sequencing. Sediment sample processing was carried out as described in

(Pettersen et al., 2022). In brief, sediment DNA was extracted on a KingFisher Flex automated

liquid handling platform using the MagAttract PowerSoil DNA KF Kit (Qiagen). The

amplification of the V3-V4 region of the 16S rRNA gene was done using primers 341F and

806R (Yu et al., 2005). Individual amplicon PCR products were dual-indexed using different

combinations of Illumina forward and reverse primers with unique barcode tags. The library

pools were sequenced a MiSeq v3 platform at the Norwegian Sequencing Centre in Oslo.

Shotgun sequencing was performed on an Illumina Novaseq 6000 platform at Novogene, UK

facility using the protocol stated in (Gourlé et al., 2018).

Subset B samples were subjected to 16S amplicon sequencing using both Illumina and ONT

sequencing platforms. Library preparation for Illumina sequencing of Subset B samples was

performed as described above. ONT sequencing libraries were prepared using the Oxford

Nanopore Sequencing kit (SQK-LSK114). The first amplification was performed using modified

16S forward and reverse primers (Genetic Analysis, Norway) with the following PCR program:

95°C for 15 min, 30 cycles of 95°C for 30 sec, 55°C for 30 sec and 72°C for 80 sec. Amplicons

were purified with 1:1 Ampure XP paramagnetic beads (Beckman Coulter) before indexing

PCR. Between 10 and 60 ng of purified full-length 16S amplicons was added to the PCR

mixture, in addition to 0.2 µM of a unique PCR barcode (ONT cat.nr. EXP-PBC096). The

barcoding amplification was done using the following program: 95°C for 15 min, 12 cycles of

95°C for 30 sec, 62°C for 15 min and 65°C for 2 min, followed by 65°C for 10 min. Barcoded

PCR products were concentration-normalized and pooled in equimolar ratios, the library from

which was purified with 1:1 Ampure XP beads before sequencing on a MinION flow cell

(Eurofins Genomics, Germany).

Amplicon data processing

The 16S amplicon sequencing data from subset A were processed with VSEARCH (Rognes et

al., 2016) for clustering into Operational Taxonomic Units (OTUs) using a 95% identity

threshold. ONT reads are longer than Illumina reads but still have error levels making it

difficult to go much above a 95% identity resolution for the 16S sequences. Because the subset

A Illumina sequence data will be used to generate reference sequences for error-prone ONT

sequences in subset B, a lower 95% resolution was chosen as opposed to a higher value or a

denoising approach. In addition, a recent study of very high diversity environments like

seafloor sediments demonstrated unstable compositional profiles when using high-resolution

denoising of illumina reads(Nilsen et al., 2024).

The Illumina paired-end reads from subset B were merged and de-replicated by VSEARCH, but

not clustered. Only reads with at least five copies per sample were used, since these are most

likely without sequencing error. Nanopore reads from the same subset B samples were filtered

based on their length, which ranged between 800 – 1 400 bp.

SILVA alignments

The OTU represents the sequences of around 420 - 440 bases, which corresponds to the

lengths of the amplicons we obtained by the primers used. First, all OTU representative

sequences were aligned against the 2 152 651 16S sequences in the SILVA 138.1 (Boratyn et

al., 2013). The SILVA sequences have a median length of 1 403 bases and range in length from

900 to 4 000 bases. OTUs from subset A achieving a best match with ≥ 95% identity over ≥

95% of their length were marked as 'known', i.e. resembling something already known from

the SILVA database. The full-length corresponding matching SILVA sequences were then

retrieved as representative for' known 'OTUs (Fig. 1, box Q1).

It should be noted that many of the seed sequences used for reconstruction (see Section:

Reconstructing 16S using METASEED) derive from the same' known 'OTUs, obfuscating the

need for sequence reconstruction as matching SILVA sequences are already known. The

reconstruction was, however, still performed in order to test how closely the reconstructed

sequence could be matched to the corresponding SILVA sequence.

Reconstructing 16S using METASEED

OTUs were used as 'seed sequences' within the METASEED workflow to reconstruct a longer

section of the 16S rRNA gene from shotgun metagenome reads. METASEED collects reads

from the shotgun metagenome data, aligns them to the seed sequences and combines

assignment results with abundance information. For each seed, a reconstructed sequence is

reported if the sequence is ≥ 500 bp in length and the alignment occurs with ≥99% identity to

the corresponding seed.

Reconstructed sequences were subsequently aligned to the SILVA database, using the same

thresholds as above. As mentioned before (see Section: SILVA alignments), the 'known 'seeds

could now be checked to verify whether the seed and its reconstructed sequence matched

the same SILVA sequence, or at least the same taxon. The most interesting reconstructed

sequences derive from cases where the SILVA database generated no matches to the seed nor

to its reconstructed sequence. These reconstructed sequences were then added to our

collection as new 16S sequences, representing new taxa relative to the SILVA collection (Fig.

1, boxes Q2 and Q3).

Shotgun data assembly

The shotgun metagenome data from the 94 samples in subset A were filtered and merged

using BBduk and BBmerge (Bushnell, 2014) from BBmap v39.01. The assembly was done using

metaspades v3.15.5 (Nurk et al., 2017). From the assembled contigs we used Barrnap v0.9

(https://github.com/tseemann/barrnap) to extract 16S sequences from the contigs using

length threshold ≥ 800 bases. As 16S sequences extracted from different samples may be

identical, all 16S sequences were clustered using VSEARCH v2.22.1, as described below, to

obtain a resolution of 95% identity (Fig. 1, box Q4).

Finally, we also included 16S centroids that didn't give any match to the above sequences.

These unmatched 16S sequences represent the lowest quality part of the reference database

(Fig. 1, box Q5).

Taxonomic assignments

Taxonomic assignments of all database sequences was obtained using the SINTAX algorithm

(Edgar, 2016) implemented in the VSEARCH software, where we used the GSR database and

the NCBI taxonomy (Molano et al., 2024). This means sequences were assigned to all ranks to

species, with confidence scores at each rank.

Phylogenetic placement of METASEEED and Barrnap derived sequences

Phylogenetic placement of the METASEED- and Barrnap- derived sequences were determined

since these are 'new '16S variants. First, all database sequences (excluding short amplicons)

were analysed with `cmsearch` in the `infernal` software v1.1.4 (Eddy, 2002) and the

secondary structure model for bacterial 16S downloaded from the `Rfam` database (Kalvari et

al., 2021). Sequences with an overall score (bitscore/length of the sequence) below 0.9 were

filtered out. A multiple alignment of the resulting sequences was created using `cmalign` and

computed evolutionary distances using Felsenstein-81 model (Felsenstein, 1981) and the `ape`

R package (Paradis & Schliep, 2019). These distances were then used in a hierarchical

clustering with average linkage and the sequences were split into 64 clusters. The clusters with

at least 5 members were subject to a Fisher exact test for enrichment of sequences originating

from either METASEED or Barrnap, i.e. sequences not previously found in SILVA. We

considered a cluster to be enriched with new variants, if the odds-ratio is greater than 1 and

the p-value is less than 0.05. In these clusters there are some 16S variants where the taxonomy

is known to phylum rank, and we list the dominating phyla in the enriched clusters based on

this.

Database comparisons

As previously stated, one of the main motivations behind this work is to improve the database

that will be used to analyse amplicon data from Oxford Nanopore Technology. EMU (Curry et

al., 2022) is a standard software for profiling such data and we wanted to compare our

database to the default EMU database. We first used VSEARCH alignments to assess the

overlap between our generated database and the default EMU database. Matches that failed

to show at least 95% identity were not considered.

Next, we wanted to test how the Nanopore reads from the 90 subset B sediment samples

would be profiled using the EMU software in combination with its default database (EMU-DB)

as well as our custom database (AQUAeD-DB). In addition, we also tested the profiling of

Illumina de-replicated reads using standard VSEARCH alignments, i.e. pairwise alignment

between read and database sequences, and assigning to the best, an identity threshold of

95% was used. The read count matrices from the Illumina VSEARCH and Nanopore EMU tool

assignments for the AQUAeD-DB and EMU-DB databases were used to compute relative

abundance. Prior to the correlation analysis, missing database sequences from the Illumina

read count were added to the Nanopore read count matrix, and vice versa, with a count of

zero for each sample. This was done to resolve disparities in matrix lengths.

Results

Addition   of   SILVA   sequences   to   the   database

The full-length sequences were first collected from the SILVA database, by matching them

against the16S rRNA amplicon representative sequences, generated by conventional short-

read data processing. Using a 95% identity threshold, our 16S rRNA pipeline generated 14,393

OTUs from the subset A data. Each OTU has a representative sequence consisting of 400 - 440

bases in accord with the length of the V3-V4 amplicons. These sequences were then aligned

against the SILVA database using VSEARCH. In total 5 278 of OTUs showed at least 95% identity

matches with some of the SILVA sequences (identity distribution shown in Fig. 2). This

represents 84% - 96% of the total OTU abundance in these 94 samples. The 4 921 distinct

SILVA sequences matched to the OTU sequences were collected and filtered using `infernal`

`cmsearch,` which resulted in 4 838 sequences. The SILVA sequences were added to the

database as full-length representatives of these OTUs and they belong to the highest quality

(Table 1, Q1) sequences in the database.

Fig. 2 Quality of OTU identifications. Frequency distribution of OTU sequences (y-axis) with

varying degrees of sequence similarity (x-axis) to SILVA database sequences. The green

columns in indicate the number of OTUs with a SILVA match while the grey columns indicate

no relevant match to SILVA database.

Reconstructing 16S rRNA genes using METASEED

All 14 393 OTUs were passed to the METASEED together with shotgun metagenome data in order to re-construct near full-length 16S rRNA from the shorter amplicon sequences. The screening process for the METASEED results is elaborated in Fig. S1. This analysis resulted in 3 525 16S rRNA sequences, of which 3 392 were ≥ 500 bases in length. Of these, 2 947 had at least 99% sequence identity to their seeds, and were considered re-constructed 16S sequences.

From the 2 947 reconstructed 16S sequences, 1 176 had no match to SILVA sequences already collected (≥ 95% identity). Reconstructed 16S rRNA sequences with a minimum length of 800 bases were regarded as high quality, while sequences 500 - 800 bases in length were considered low quality. This resulted in 421 high quality and 755 low quality reconstructed sequences. The total of 1 176 METASEED reconstructed sequences were filtered using `cmsearch`, which led to the addition of 717 reconstructed sequences to the database. The Q2 and Q3 category in Table 1 summarize these results. The increase in the captured relative abundance when including Q1+Q2+Q3 sequences versus only Q1 (Fig. S2).

Addition of 16S rRNA from contigs using Barrnap

We also looked for 16S rRNA genes in the assembled contigs from the metagenome data using the `barrnap` software. From the 94 samples, we found in total 9 839 short or longer sequences recognized as 16S rRNA. The number of sequences was reduced to 1 149 when applying a length threshold of at least 800 bp. Finally, since the same 16S rRNA sequence may

have been assembled and extracted from different samples, these sequences were clustered

using a 95% identity threshold, which reduced the number of sequences to 518. Again, we

used the `cmsearch` filtering, resulting in 320 high quality near full-length 16S rRNA

sequences, which were added to database. The total length of assembled contigs was greater

than 32 000 megabasepairs, implying that roughly one full-length high-quality 16S rRNA was

found for every 100 mega base-pairs of genomic sequence (Table 1, Q4).

Apart from the sequences collected from the above-mentioned sources, we also included 8

670 OTU sequences that did not match to any sequences added to the database from SILVA,

METASEED, and Barrnap when using a 95% identity threshold (Table 1, Q5). Including all

categories (Table 1), the final size of the AQUAeD database was 14 545 16S sequences.

Taxonomic assignments

All sequences in the AQUAeD database were classified using `SINTAX` (Fig. 3), with the

exception of sequences belonging to the Q1 category, which were copied from the SILVA

database with taxonomic assignments.

Fig. 3 The bars depict the fraction of classified sequences (Q2 - Q5) in the AQUAeD Database

using SINTAX (bootstrap value 0.8 and 0.5) and the GSR databases. The x-axis shows various

taxonomic ranks and y-axis shows the fraction of classified sequences.

Enrichment of 'new '16S variants

Sequences in categories Q2, Q3 and Q4 (Table 1) are 'new' 16S rRNA gene variants in the sense

that they are not included in the SILVA database. We wanted to see how these distribute when

clustered among the sequences in Q1 as well. The multiple sequence alignment of these 5

875 sequences was followed by the evolutionary distance calculation, cluster generation and

using the Fisher test for enrichment of the 'new variants'. From 64 clusters we found that 17

clusters were enriched by 'new' variants. The other members of these enriched clusters

predominantly consisted of phyla Planctomycetota, Myxococcota, and Verrucomicrobiota

(Table 2).

Database comparison

EMU software is the standard tool used for taxonomic profiling of Nanopore 16S reads. It comes with a default database (EMU-DB), consisting of 49 243 sequences. From a comparison between the EMU-DB and AQUAeD-DB, we observed that 18 584 (38%) EMU database sequences had a match to 509 (3.5 %) AQUAeD database sequences when using a 95% identity threshold.

The 88 subset B samples resulted in a total of 973 758 dereplicated Illumina reads, corresponding to 51 397 distinct sequences. The alignment of these reads (95% identity) to the EMU-DB resulted in a match for 3% of the database sequences, comprising 52% of the total Illumina reads. Alignment with the AQUAeD-DB yielded a match to 23% of the database sequences, covering 99% of the total Illumina reads. The Nanopore amplicon reads from the 88 subset B samples having length between 800 and 1 400 bases resulted in **4 972 056 reads**. The analysis of Nanopore reads resulted in 3 967 AQUAeD database sequences, accounting for approximately 27% of the database, while EMU-DB matched to 2 560 database sequences, accounting for approximately 5% of the database.

Fig. 4 The Database comparison. A) Illumina results for taxonomic assignments using VSEARCH. B, Nanopore results for taxonomic assignments using EMU. B) the percentage reads assigned exceeds 100% due to the fuzzy assignment by EMU which resulted in a total 'count' of assigned reads that was larger than the number of reads. Pie charts represent reads and database sequences are represented by bars. Shaded/striped regions in each pie chart or bar indicate the fraction of reads assigned or the number of database sequences to which they were assigned, respectively. Detailed information about alignments can be found in Table S1.

A comparison of read count matrices revealed that 82% of the EMU-DB sequences in the

Illumina read count matrix were missing in the corresponding Nanopore read count matrix,

while 17% of the sequences in the Nanopore matrix were the missing from Illumina matrix.

For the AQUAeD-DB read count matrices, the overlap was notably better, with 45% of Illumina

sequences and 13% of Nanopore sequences missing from their counterparts, indicating a

lower level of missing database sequences. Correlation analysis demonstrated high positive

correlation between read count matrices for the AQUAeD-DB, whereas the EMU-DB shows

very weak negative correlation (Fig. 5). The relative mean abundance of Illumina and

Nanopore results for AQUAeD-DB demonstrate a significant positive correlation (Fig. 6).

Fig. 5 Correlation between Illumina and ONT read count matrices. Boxplots indicate

correlation coefficients for read count matrix similarity when using the AQUAeD or EMU

database.

Fig. 6 Relative mean abundance of Illumina versus Nanopore sequences for subset B samples.

The scatterplot depicts the log of relative mean abundance of Nanopore (y-axis) versus the

log of relative abundance of Illumina(x-axis) of AQUAeD-DB database sequences across

samples. Cases where either Illumina or Nanopore values were zero are omitted.

Discussion

The motivation for building the AQUAeD-DB was to enhance the use of Nanopore data in the

study of seafloor sediments. Thus, we wanted to see if the use of AQUAeD-DB has some

benefit compared to a standard database. Since the EMU tool is a popular choice for

Nanopore 16S data processing, we used its default database (EMU-DB) in our comparison. In

addition to using EMU for assigning the reads, we also used VSEARCH, which is typically used

for Illumina data, just for comparison. The VSEARCH alignment directly between the AQUAeD-

DB and EMU-DB reveals that 3.5% of the AQUAeD database could represent 38% of the EMU

database, implying that the AQUAeD-DB, despite being smaller in the number of sequences,

covers an order of magnitude more seafloor diversity compared to EMU-DB.

The AQUAeD-DB comprises of 14 545 16S sequences that were collected from seafloor

sediments (Fig. 1). The Illumina amplicon sequencing of subset A seafloor samples yielded 14

393 centroid sequences when clustered at the coarse level of 95% identity. Approximately

40% of these sequences matched the SILVA database sequences, even with relatively liberal

identity threshold of 95% (Fig. 2). However, this set of matching SILVA sequences, the Q1,

makes up the backbone of our AQUAeD-DB. To add more longer sequences to the database,

we used our METASEED tool (Philip et al., 2024) to extend the amplicon sequences, making

use of the Illumina metagenome data from the same samples (subset A). This resulted in the

Q2 and Q3 part of the database. The total abundance covered by the database sequences in

each sample did not increase appreciably by adding these sequences, but the increase was

larger for the exceptionally high diversity samples (high Shannon index of 5-7, Fig. S2). This

typically allowed us to obtain 16S sequences for the low-abundance and ecologically

significant organisms from phyla like Planctomycetota (table 2). In the Q4 part of the database,

we tried to collect long 16S directly from metagenome assembled reads. The fact that we only

collect one such sequence per 100 megabases of assembled contigs indicates how difficult it

is to obtain such information from the assembly of short read metagenome data. The

limitations of metagenome assembly, specifically the challenges in retrieving 16S sequences,

have been previously addressed (Hiseni et al., 2022; Mise & Iwasaki, 2022; Shaiber & Eren,

2019).

The AQUAeD-DB can be roughly split into two parts: The 'known 'sequences (Q1) and the 'new

variants '(Q2-Q5, Fig 3). The fractions of taxonomic assignments were generally low, especially

for the Q3-Q5 sequences. This emphasizes that new sequence variants in the AQUAeD

database (Q2-Q5) have poor representation in a public database, like the GSR used here.

Considering the poor taxonomic information, we wanted to see if phylogenetic distances

between these new variants, and the Q1 sequences indicate a grouping of these new

sequences, or if they are scattered around in the phylogeny. By clustering based on

phylogenetic distances, we found several clusters where these new variants are over-

represented, which include significant marine microorganisms from the phyla

Planctomycetota, Myxococcota, and Verrucomicrobiota (Dell'Anno et al., 2021; Freitas et al.,

2012; Wu et al., 2024)

A clear difference was evident when we did the assignment with both VSEARCH, which is

normally used for Illumina data, and EMU, which is the standard choice for Nanopore (Fig 4).

The VSEARCH tool aligned the high-precision Illumina reads, with EMU-DB assigning almost

half of the reads and spreading them out among only 1 474 of the nearly 50 000 database

sequences. For AQUAeD-DB, almost all reads are assigned to 3 301 database sequences. The

Nanopore reads showed different results compared to the VSEARCH assignment of Illumina

data, especially in terms of database coverage. One of the major contributing factors can be

the significantly higher noise levels compared to Illumina data; this leads to reads being

distributed across a much larger number of sequences in both databases (Magi et al., 2017;

Santos et al., 2020). Another difference lies in the fact that the EMU tool assigns more than

100% of the reads, which likely indicates an artifact of the fuzzy assignment process

(Szoboszlay et al., 2023). From VSEARCH+Illumina and EMU+Nanopore, we see how more

organisms are being given read counts in the AQUAeD-DB compared to the EMU-DB. This is

the effect of AQUAeD-DB containing more relevant taxa for seafloor data, while EMU-DB is a

generic collection and probably more relevant for use in medical research.

The previously stated percentage of missing database sequences in the read count matrices

for Illumina and Nanopore indicate that the EMU-DB analysis results exhibit greater disparities

compared to the AQUAeD-DB results. A clear positive correlation between Illumina and

Nanopore relative abundance were revealed in AQUAeD-DB whereas EMU-DB shows very

inconsistent and poor results (Fig 5). From these results it is evident that AQUAeD-DB results

in more reliable and consistent assignments for both Illumina and ONT analysis. These findings

were supported by the broad range of diversity, especially for the low- abundance taxa that

the AQUAeD-DB database was able to cover (Fig 6). These observations once again underline

the benefits of AQUAeD-DB and emphasize the potential implementation for Nanopore based

environmental studies.

These findings indicated that AQUAeD-DB have illustrated a significant contribution in

environmental analysis using ONT sequencing data. The database enrichment analysis implies

that significant marine microbiota is represented in newly developed AQUAeD database,

comprising both existing sequences and novel sequences from the aforementioned strategies.

This optimizes taxonomical assignment and provides potential for more explicit

environmental context. The study emphasizes the potential, and necessity of such databases

in facilitating the implementation of the cutting-edge technologies, thus enabling the

comprehensive environmental assessments.

Conclusion

The newly developed AQUAeD-DB database is a specialized database for 16S rRNA gene

sequences from highly diverse and complex environmental samples. The primary objective of

AQUAeD-DB was to provide a better reference database for identification of Nanopore

sequences targeting microbial communities in North Atlantic seafloor sediments. Based on

results of the enrichment study and database comparison, this database is substantially

diverse, and efficiently covers an expanded range of microorganisms that are poorly

represented in other databases. This study was designed as a pilot project with the potential

for future scalability of the database to incorporate additional 16S rRNA gene sequences. This

expansion will be rooted in ongoing efforts to improve environmental surveys and monitoring

using microbial profiles.

Acknowledgements

433   We express our gratitude to the fish farmers that provided samples for this research. We

434   would also like to thank the employees of STIM, Aqua Kompetanse, and Akvaplan-NIVA that

435   performed the sampling.

Data availability

The AQUAED-DB and taxonomy details are publicly available (https://arken.nmbu.no/~pmelcy/share/AQUAED-DB/). All raw sequences used in this study have been deposited in the NCBI BioProject database. Illumina 16S rRNA sequences from subset A and subset B is a part of the project with the accession code PRJNA1128851. The raw shotgun data of subset A has been uploaded with the accession code PRJNA1128851. The raw Nanopore sequencing data for subset B has been uploaded with the accession code PRJNA1153974. The metadata files for all datasets are available in https://arken.nmbu.no/~pmelcy/share/AQUAED-DB/metadata/

Author contributions

Knut Rudi, Lars Snipen and Melcy Philip conceived the study; Jessica Louise Ray, Morten Stokkan, Ragnhild Pettersen and Sanna Majaneva design the collection of sediment samples; Tonje Nilsen performed all lab experiments; Lars Snipen and Melcy Philip performed all analyses, with conceptual input from Knut Rudi; Knut Rudi, Lars Snipen and Melcy Philip contributed to interpretation of results; Knut Rudi, Lars Snipen and Melcy Philip drafted the manuscript. All author(s) read and commented on the manuscript.

Funding

456   This study is a part of the project: "AQUAeD", funded by The Research Council of Norway

457   (Project Number: 320076).

Competing interests

At the time of manuscript writing, Sanna Majaneva, Ragnhild Pettersen were employed by

company Akvaplan-niva AS. Jessica Louise Ray was employed by company Aqua Kompetanse

AS and Morten Stokkan was employed by company STIM AS. The remaining authors declare

that the research was conducted in the absence of any commercial or financial relationships

that could be construed as a potential conflict of interest.

Table 1: The AQUAeD-DB contains sequences of 5 quality categories, from the best (Q1) to the poorest (Q5)

Category descriptions are provided in Materials and Methods. Lengths are shown in basepairs (bp).

Category	Origin	Number of sequences	Minimum length	Median length	Maximum length
Q1	SILVA	4838	900	1461	1668
Q2	High Quality METASEED	311	800	918	1522
Q3	Low Quality METASEED	406	501	688.5	799
Q4	16S from contigs	320	800	924	1577
Q5	Amplicons	8670	288	426	438

469   **Table 2:** Enrichment of 'new' 16S variants. Results of Fisher tests for 17 'new variants 'enriched clusters when
470   k=64.

Odds ratio	Cluster size	New variants fraction	Dominant phylum
7.06	20	0.6	Planctomycetota
7.01	5	0.6	Desulfobacterota,Proteobacteria
6.56	12	0.58	Patescibacteria
4.015	13	0.46	Patescibacteria
3.97	35	0.46	Spirochaetota
3.27	39	0.41	Verrucomicrobiota
3.01	23	0.39	Spirochaetota
2.71	30	0.37	Verrucomicrobiota
2.59	62	0.35	Chloroflexi
2.58	114	0.35	Verrucomicrobiota
2.55	153	0.35	Planctomycetota
2.54	37	0.35	Firmicutes
2.34	30	0.33	Firmicutes
2.34	27	0.33	Fibrobacterota
2.13	61	0.31	Planctomycetota
2.07	72	0.31	Verrucomicrobiota
1.61	241	0.25	Myxococcota

References

Amarasinghe, S. L., Su, S., Dong, X., Zappia, L., Ritchie, M. E., & Gouil, Q. (2020). Opportunities and
challenges in long-read sequencing data analysis. *Genome Biology*, *21*(1), 30.
https://doi.org/10.1186/s13059-020-1935-5
Benítez-Páez, A., Portune, K. J., & Sanz, Y. (2016). Species-level resolution of 16S rRNA gene amplicons
sequenced through the MinION™ portable nanopore sequencer. *GigaScience*, *5*(1).
https://doi.org/10.1186/s13742-016-0111-z
Boratyn, G. M., Camacho, C., Cooper, P. S., Coulouris, G., Fong, A., Ma, N., Madden, T. L., Matten, W.
T., McGinnis, S. D., Merezhuk, Y., Raytselis, Y., Sayers, E. W., Tao, T., Ye, J., & Zaretskaya, I. (2013).
BLAST: a more efficient report with usability improvements. *Nucleic Acids Research*, *41*(W1),
W29-W33. https://doi.org/10.1093/nar/gkt282
Bushnell, B. (2014). *BBMap: a fast, accurate, splice-aware aligner*.
Canals, M., Pham, C. K., Bergmann, M., Gutow, L., Hanke, G., van Sebille, E., Angiolillo, M., Buhl-
Mortensen, L., Cau, A., Ioakeimidis, C., Kammann, U., Lundsten, L., Papatheodorou, G., Purser,
A., Sanchez-Vidal, A., Schulz, M., Vinci, M., Chiba, S., Galgani, F.,…Giorgetti, A. (2021). The
quest for seafloor macrolitter: a critical review of background knowledge, current methods
and future prospects. *Environmental Research Letters*, *16*(2), 023001.
https://doi.org/10.1088/1748-9326/abc6d4
Curry, K. D., Wang, Q., Nute, M. G., Tyshaieva, A., Reeves, E., Soriano, S., Wu, Q., Graeber, E., Finzer, P.,
Mendling, W., Savidge, T., Villapol, S., Dilthey, A., & Treangen, T. J. (2022). Emu: species-level
microbial community profiling of full-length 16S rRNA Oxford Nanopore sequencing data.
Nature Methods, *19*(7), 845-853. https://doi.org/10.1038/s41592-022-01520-4
Dell'Anno, F., Rastelli, E., Tangherlini, M., Corinaldesi, C., Sansone, C., Brunet, C., Balzano, S., Ianora, A.,
Musco, L., Montereali, M. R., & Dell'Anno, A. (2021). Highly Contaminated Marine Sediments
Can Host Rare Bacterial Taxa Potentially Useful for Bioremediation [Original Research].
Frontiers in Microbiology, *12*. https://doi.org/10.3389/fmicb.2021.584850
DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., Huber, T., Dalevi, D., Hu, P.,
& Andersen, G. L. (2006). Greengenes, a chimera-checked 16S rRNA gene database and
workbench compatible with ARB. *Applied and Environmental Microbiology*, *72*(7), 5069-5072.
Dueholm, M. K. D., Nierychlo, M., Andersen, K. S., Rudkjøbing, V., Knutsson, S., Arriaga, S., Bakke, R.,
Boon, N., Bux, F., Christensson, M., Chua, A. S. M., Curtis, T. P., Cytryn, E., Erijman, L.,
Etchebehere, C., Fatta-Kassinos, D., Frigon, D., Garcia-Chaves, M. C., Gu, A. Z.,…Mi, D. A. S. G.
C. (2022). MiDAS 4: A global catalogue of full-length 16S rRNA gene sequences and taxonomy
for studies of bacterial communities in wastewater treatment plants. *Nature Communications*,
13(1), 1908. https://doi.org/10.1038/s41467-022-29438-7
Eddy, S. R. (2002). A memory-efficient dynamic programming algorithm for optimal alignment of a
sequence to an RNA secondary structure. *BMC Bioinformatics*, *3*(1), 18.
https://doi.org/10.1186/1471-2105-3-18
Edgar, R. (2016). SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS sequences. In:
bioRxiv.
F. Escapa, I., Huang, Y., Chen, T., Lin, M., Kokaras, A., Dewhirst, F. E., & Lemon, K. P. (2020). Construction
of habitat-specific training sets to achieve species-level assignment in 16S rRNA gene datasets.
Microbiome, *8*(1), 65. https://doi.org/10.1186/s40168-020-00841-w
Federhen, S. (2011). The NCBI Taxonomy database. *Nucleic Acids Research*, *40*(D1), D136-D143.
https://doi.org/10.1093/nar/gkr1178

Felsenstein, J. (1981). Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol*, *17*(6), 368-376. https://doi.org/10.1007/bf01734359

Freitas, S., Hatosy, S., Fuhrman, J. A., Huse, S. M., Welch, D. B., Sogin, M. L., & Martiny, A. C. (2012). Global distribution and diversity of marine Verrucomicrobia. *Isme j*, *6*(8), 1499-1505. https://doi.org/10.1038/ismej.2012.3

Ghate, S. D., Shastry, R. P., Arun, A. B., & Rekha, P. D. (2021). Unraveling the bacterial community composition across aquatic sediments in the Southwestern coast of India by employing high-throughput 16S rRNA gene sequencing. *Regional Studies in Marine Science*, *46*, 101890. https://doi.org/https://doi.org/10.1016/j.rsma.2021.101890

Henderson, G., Yilmaz, P., Kumar, S., Forster, R. J., Kelly, W. J., Leahy, S. C., Guan, L. L., & Janssen, P. H. (2019). Improved taxonomic assignment of rumen bacterial 16S rRNA sequences using a revised SILVA taxonomic framework. *PeerJ*, *7*, e6496. https://doi.org/10.7717/peerj.6496

Hiseni, P., Snipen, L., Wilson, R. C., Furu, K., & Rudi, K. (2022). Questioning the Quality of 16S rRNA Gene Sequences Derived From Human Gut Metagenome-Assembled Genomes [Opinion]. *Frontiers in Microbiology*, *12*. https://doi.org/10.3389/fmicb.2021.822301

Hoshino, T., Doi, H., Uramoto, G. I., Wörmer, L., Adhikari, R. R., Xiao, N., Morono, Y., D'Hondt, S., Hinrichs, K. U., & Inagaki, F. (2020). Global diversity of microbial communities in marine sediment. *Proc Natl Acad Sci U S A*, *117*(44), 27587-27597. https://doi.org/10.1073/pnas.1919139117

Kalvari, I., Nawrocki, E. P., Ontiveros-Palacios, N., Argasinska, J., Lamkiewicz, K., Marz, M., Griffiths-Jones, S., Toffano-Nioche, C., Gautheret, D., & Weinberg, Z. (2021). Rfam 14: expanded coverage of metagenomic, viral and microRNA families. *Nucleic Acids Research*, *49*(D1), D192-D200.

Kozlov, A. M., Zhang, J., Yilmaz, P., Glöckner, F. O., & Stamatakis, A. (2016). Phylogeny-aware identification and correction of taxonomically mislabeled sequences. *Nucleic Acids Research*, *44*(11), 5022-5033.

Lemoinne, A., Dirberg, G., Georges, M., & Robinet, T. (2023). Fine-scale congruence in bacterial community structure from marine sediments sequenced by short-reads on Illumina and long-reads on Nanopore.

Madsen, E. L. (2011). Microorganisms and their roles in fundamental biogeochemical cycles. *Current Opinion in Biotechnology*, *22*(3), 456-464. https://doi.org/https://doi.org/10.1016/j.copbio.2011.01.008

Magi, A., Semeraro, R., Mingrino, A., Giusti, B., & D'Aurizio, R. (2017). Nanopore sequencing data analysis: state of the art, applications and challenges. *Briefings in Bioinformatics*, *19*(6), 1256-1272. https://doi.org/10.1093/bib/bbx062

Marlow, J., Steele, J., Case, D., Connon, S., Levin, L., & Orphan, V. (2014). Microbial abundance and diversity patterns associated with sediments and carbonates from the methane seep environments of Hydrate Ridge, OR. *Frontiers in Marine Science*, *1*. https://doi.org/10.3389/fmars.2014.00044

McDonald, D., Jiang, Y., Balaban, M., Cantrell, K., Zhu, Q., Gonzalez, A., Morton, J. T., Nicolaou, G., Parks, D. H., Karst, S. M., Albertsen, M., Hugenholtz, P., DeSantis, T., Song, S. J., Bartko, A., Havulinna, A. S., Jousilahti, P., Cheng, S., Inouye, M.,...Knight, R. (2024). Author Correction: Greengenes2 unifies microbial data in a single reference tree. *Nature Biotechnology*, *42*(5), 813-813. https://doi.org/10.1038/s41587-023-02026-w

Meola, M., Rifa, E., Shani, N., Delbès, C., Berthoud, H., & Chassard, C. (2019). DAIRYdb: a manually curated reference database for improved taxonomy annotation of 16S rRNA gene sequences from dairy products. *BMC Genomics*, *20*(1), 560. https://doi.org/10.1186/s12864-019-5914-8

Mise, K., & Iwasaki, W. (2022). Unexpected absence of ribosomal protein genes from metagenome-assembled genomes. *ISME Communications*, *2*(1). https://doi.org/10.1038/s43705-022-00204-6

Molano, L.-A. G., Vega-Abellaneda, S., & Manichanh, C. (2024). GSR-DB: a manually curated and optimized taxonomical database for 16S rRNA amplicon analysis. *mSystems*, *9*(2), e00950-00923. https://doi.org/doi:10.1128/msystems.00950-23

Nierychlo, M., Andersen, K. S., Xu, Y., Green, N., Jiang, C., Albertsen, M., Dueholm, M. S., & Nielsen, P. H. (2020). MiDAS 3: An ecosystem-specific reference database, taxonomy and knowledge platform for activated sludge and anaerobic digesters reveals species-level microbiome composition of activated sludge. *Water Research*, *182*, 115955. https://doi.org/https://doi.org/10.1016/j.watres.2020.115955

Nilsen, T., Snipen, L.-G., Angell, I. L., Keeley, N. B., Majaneva, S., Pettersen, R., & Rudi, K. (2024). Swarm and UNOISE outperform DADA2 and Deblur for denoising high-diversity marine seafloor samples. *ISME Communications*. https://doi.org/10.1093/ismeco/ycae071

Nurk, S., Meleshko, D., Korobeynikov, A., & Pevzner, P. A. (2017). metaSPAdes: a new versatile metagenomic assembler. *Genome research*, *27*(5), 824-834.

Overgaard, C. K., Tao, K., Zhang, S., Christensen, B. T., Blahovska, Z., Radutoiu, S., Kelly, S., & Dueholm, M. K. D. (2022). Application of ecosystem-specific reference databases for increased taxonomic resolution in soil microbial profiling. *Front Microbiol*, *13*, 942396. https://doi.org/10.3389/fmicb.2022.942396

Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*, *35*(3), 526-528.

Patil, P. K., Baskaran, V., Vinay, T. N., Avunje, S., Leo-Antony, M., Shekhar, M. S., Alavandi, S. V., & Vijayan, K. K. (2021). Abundance, community structure and diversity of nitrifying bacterial enrichments from low and high saline brackishwater environments. *Letters in Applied Microbiology*, *73*(1), 96-106. https://doi.org/10.1111/lam.13480

Pettersen, R., Ormaasen, I., Angell, I. L., Keeley, N. B., Lindseth, A., Snipen, L., & Rudi, K. (2022). Bimodal distribution of seafloor microbiota diversity and function are associated with marine aquaculture. *Marine Genomics*, *66*, 100991. https://doi.org/https://doi.org/10.1016/j.margen.2022.100991

Philip, M., Rudi, K., Ormaasen, I., Angell, I. L., Pettersen, R., Keeley, N. B., & Snipen, L.-G. (2024). METASEED: a novel approach to full-length 16S rRNA gene reconstruction from short read data. *BMC Bioinformatics*, *25*(1), 237. https://doi.org/10.1186/s12859-024-05837-z

Ritari, J., Salojärvi, J., Lahti, L., & de Vos, W. M. (2015). Improved taxonomic assignment of human intestinal 16S rRNA sequences by a dedicated reference database. *BMC Genomics*, *16*(1), 1056. https://doi.org/10.1186/s12864-015-2265-y

Rodríguez-Pérez, H., Ciuffreda, L., & Flores, C. (2020). NanoCLUST: a species-level analysis of 16S rRNA nanopore sequencing data. *Bioinformatics*, *37*(11), 1600-1601. https://doi.org/10.1093/bioinformatics/btaa900

Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: a versatile open source tool for metagenomics. *PeerJ*, *4*, e2584. https://doi.org/10.7717/peerj.2584

Santos, A., van Aerle, R., Barrientos, L., & Martinez-Urtaza, J. (2020). Computational methods for 16S metabarcoding studies using Nanopore sequencing data. *Computational and Structural Biotechnology Journal*, *18*, 296-305. https://doi.org/https://doi.org/10.1016/j.csbj.2020.01.005

Shaiber, A., & Eren, A. M. (2019). Composite Metagenome-Assembled Genomes Reduce the Quality of Public Genome Repositories. *mBio*, *10*(3), 10.1128/mbio.00725-00719. https://doi.org/doi:10.1128/mbio.00725-19

Szoboszlay, M., Schramm, L., Pinzauti, D., Scerri, J., Sandionigi, A., & Biazzo, M. (2023). Nanopore Is Preferable over Illumina for 16S Amplicon Sequencing of the Gut Microbiota When Species-Level Taxonomic Classification, Accurate Estimation of Richness, or Focus on Rare Taxa Is Required. *Microorganisms*, *11*(3), 804. https://www.mdpi.com/2076-2607/11/3/804

Tatsuhiko Hoshinoa, H. D., Go-Ichiro Uramotoa, Lars Wörmerc, Rishi R. Adhikari, Nan Xiaoa,e,, Yuki
Moronoa, Steven D'Hondtf, Kai-Uwe Hinrichsc , Fumio Inagaki. (2020). Global diversity of
microbial communities in

marine sediment.
Wu, J., Hong, Y., Chang, X., Jiao, L., Li, Y., Liu, X., Xie, H., & Gu, J.-D. (2019). Unexpectedly high diversity
of anammox bacteria detected in deep-sea surface sediments of the South China Sea. *FEMS*
Microbiology Ecology, *95*(3). https://doi.org/10.1093/femsec/fiz013
Wu, X.-T., He, Y.-Q., Qiu, M., Li, G.-X., Ou, X.-B., Wu, Y.-B., & Bao, P. (2024). Adaptive Changes in Bacterial
and Archaeal Community Structures and Functions Occur under Primordial Conditions. *ACS*
Earth   and   Space   Chemistry,   *8*(7),   1470-1479.
https://doi.org/10.1021/acsearthspacechem.4c00089
Yang, C.-c., & Iwasaki, W. (2014). MetaMetaDB: A Database and Analytic System for Investigating
Microbial   Habitability.   *PLOS   ONE*,   *9*(1),   e87126.
https://doi.org/10.1371/journal.pone.0087126
Yang, N., Tian, C., Lv, Y., Hou, J., Yang, Z., Xiao, X., & Zhang, Y. (2022). Novel primers for 16S rRNA gene-
based archaeal and bacterial community analysis in oceanic trench sediments. *Applied*
Microbiology and Biotechnology, *106*(7), 2795-2809. https://doi.org/10.1007/s00253-022-
11893-3
Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J., Ludwig, W.,
& Glöckner, F. O. (2013). The SILVA and "All-species Living Tree Project (LTP)" taxonomic
frameworks.   *Nucleic   Acids   Research*,   *42*(D1),   D643-D648.
https://doi.org/10.1093/nar/gkt1209
Yu, Y., Lee, C., Kim, J., & Hwang, S. (2005). Group-specific primer and probe sets to detect
methanogenic communities using quantitative real-time polymerase chain reaction.
Biotechnology   and   Bioengineering,   *89*(6),   670-679.
https://doi.org/https://doi.org/10.1002/bit.20347
Zorz, J., Li, C., Chakraborty, A., Gittins, D. A., Surcon, T., Morrison, N., Bennett, R., MacDonald, A., &
Hubert, C. R. J. (2023). SituSeq: an offline protocol for rapid and remote Nanopore 16S rRNA
amplicon   sequence   analysis.   *ISME   Communications*,   *3*(1),   33.
https://doi.org/10.1038/s43705-023-00239-3