A Targeted Reference Database for Improved Analysis of Environmental 16S rRNA Oxford Nanopore Sequencing Data

Melcy Philip[1], Tonje Nilsen[1] Sanna Majaneva[2], Ragnhild Pettersen[2], Morten Stokkan[3], Jessica Louise Ray[4], Nigel Keeley[5], Knut Rudi[1], Lars-Gustav Snipen[1]

[1]Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences, Ås (formerly known as viken), Norway

[2]Akvaplan-niva, Framsenteret, P.O. Box 6606, Stakkevollan, 9296 Tromsø, Norway

[3]STIM department Bergen, Thormøhlens gate 55, 5006 Bergen, Norway

[4]Aqua Kompetanse AS, Storlavika 7, 7770 Flatanger, Norway

[5]Institute of Marine Research, Tromsø, Norway

Corresponding author: Melcy Philip, Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences, Christian Magnus Falsens vei 18, 1433, Ås (formerly known as viken), Norway (philip.melcy@nmbu.no and melcy.p.j@gmail.com)

Running Title: Targeted Database for Enhanced 16S ONT Analysis

Abstract

The Oxford Nanopore Technologies (ONT) sequencing platform is compact and efficient, making it suitable for rapid biodiversity assessments in remote areas. Despite its long reads, ONT has a higher error rate compared to other platforms, necessitating high-quality reference databases for accurate taxonomic assignments. However, the absence of targeted databases for underexplored habitats, such as the seafloor, limits ONT's broader applicability for exploratory analysis.

To address this, we have developed a strategy to build environmentally-targeted databases to enhance 16S rRNA gene(16S) analysis using ONT. We start by using Illumina short-read data to create a database of full-length or near full-length 16S sequences from seafloor samples. Initially, amplicons are mapped to the SILVA database, with matches added to our database. Unmatched amplicons are reconstructed using METASEED and Barrnap methodologies with amplicon and metagenome data. Finally, if the previous strategies did not succeed, we included the short-read sequences in the database. This resulted in AQUAeD-DB, which contains 14 545 16S sequences clustered at 95% identity. Comparative database analysis and relative abundance correlation studies reveal that AQUAeD-DB provides consistent results for both Illumina and Nanopore read assignments (median correlation coefficient: 0.50), whereas a standard database showed a substantially weaker correlation. These findings also emphasis its potential to recognize both high and low-abundance taxa, which could be key indicators in environmental studies. This work highlights the necessity of targeted databases for environmental analysis, especially for ONT-based studies, and lays foundations for future extension of the database.

42 **Keywords:** 16S rRNA, metagenomics, seafloor sediment, database, Nanopore, ONT, Illumina,

43 biodiversity assessment

Introduction

The Earth's surface is primarily covered by oceans, yet we know little about the seafloor, or

its microbiota (Canals et al., 2021). Microorganisms play a key role in biogeochemical cycling

in the global ocean thus the potential importance of microbiota to healthy ecosystem

function cannot be overstated (Madsen, 2011). Many aspects of our understanding of soft-

sediment microbiota is based on short read amplicon sequencing of the 16S rRNA gene V3

and V4 regions (Ghate et al., 2021; Hoshino et al., 2020; Yang et al., 2022). Advancements in

long read sequencing, including Oxford Nanopore Technology (ONT), have enabled rapid

analysis of environmental microbiota (Zorz et al., 2023). However, the relatively high error

rate of ONT sequencing technology remains a persistent challenge, rendering ONT strategies

dependent on reference databases (Amarasinghe et al., 2020). For effective and robust use

of ONT technology that generates reads spanning nearly the entire 16S gene, databases of

nearly full-length 16S rRNA gene sequences with comprehensive representation of habitat-

relevant taxa are needed.

The most widely used 16S databases are the curated Greengenes and SILVA databases, as

well as the non-curated NCBI 16S rRNA database (Federhen, 2011; McDonald et al., 2024;

Yilmaz et al., 2013) . These databases are limited by coverage despite the large numbers of

reference sequences that they house. For instance, the 16S component of the GenBank

database consists of 43% chimeric sequences, the presence of which may confound accurate

taxonomic assignments (DeSantis et al., 2006). Furthermore, despite the use of quality

control tools, public 16S reference databases still contain 0.2% to 2.5% misidentified species,

leading to taxonomic misassignments (Kozlov et al., 2016). In addition, there is a major bias

towards characterization of human-associated and highly abundant bacteria, while most of

environmental microbial diversity remains uncharacterized (Yang & Iwasaki, 2014). In this

context, we fail to notice potentially low-abundant microorganisms, such as nitrifying

bacteria, anammox bacteria and methanotrophs, that may be crucial for reflecting the

health of the environment (Marlow et al., 2014; Patil et al., 2021; Wu et al., 2019). Benthic

microbes, in particular, are underrepresented in databases and our understanding of them is

constrained by very few studies (Tatsuhiko Hoshinoa, 2020). The current scenario thus

emphasises the need for a specialised database for seafloor microorganisms, since

environmental microbial communities are often missing from current databases.

Studies reveal that employing targeted databases can improve the accuracy and coverage of

amplicon-based microbial profiling (Dueholm et al., 2022; Meola et al., 2019; Overgaard et

al., 2022). Although ONT sequencing technology is advancing, there are only few

bioinformatics tools for ONT 16S profiling (Ciuffreda et al., 2021; Santos et al., 2020; Sissel et

al., 2015) apart from the workflow offered by Oxford Nanopore. There are several ongoing

efforts to improve taxonomic assignments within ONT-driven studies, with the EMU tool

showing promise as an extensively used approach (Curry et al., 2022; Rodríguez-Pérez et al.,

2020; Zorz et al., 2023). However, all these tools, require a good reference database for

improved taxonomic assignment. This underlines the necessity of an environment-specific or

specialized reference databases and the associated potential benefits. (F. Escapa et al., 2020;

Henderson et al., 2019; Nierychlo et al., 2020; Ritari et al., 2015). Studies have focused on

achieving high resolution using Nanopore, given its longer reads covering most of the 16S

gene, whereas our study aims to capture the broadest possible diversity from environmental

samples (Benítez-Páez et al., 2016; Lemoinne et al., 2023).

The aim of this work was to build a targeted 16S database for seafloor sediment

environments to facilitate full length 16S (v2-v9) ONT sequencing for microbial profiling and

prediction of ecosystem status. We have called this database "AQUAeD-DB" in

acknowledgment of the research project "AQUAeD" which provided environmental samples

for sequencing analysis (see Acknowledgements). Our strategy (Fig. 1) makes use of short-

read amplicon data in combination with existing databases, as well as whole genome

shotgun sequencing in combination with our recently developed method METASEED (Philip

et al., 2024). We also assess and present the effectiveness of this targeted database in

comparison to the existing tools and databases for taxonomic assignment of 16S ONT

amplicons.

Fig. 1 The16S database generation. a) The seafloor samples used in this study were collected from various sites along the Norwegian coast. b) The samples for database generation were sequenced using Illumina, providing shotgun and 16S rRNA amplicon reads, c) Various methods are used to collect or rebuild longer 16S rRNA sequences, which are then graded into different quality levels, from Q1 (best) to Q5 (poorest). These sequences are added to our reference database e) and used to for taxonomic assignment of ONT sequences

107 amplified   from   geographically   proximal   seafloor   sediment   material.

Materials and Methods

Data

In this study, we used total of 182 marine sediment samples collected along a north-south

gradient along the Norwegian coast as mentioned previously (Nilsen et al., 2024; Pettersen

et al., 2022). The data was randomly divided into two subsets: subset A consisted of 94

samples and was used for the database construction; subset B consisted of 88 samples and

was used for database comparison.

Subset A was subject to Illumina sequencing, both 16S amplicon and whole genome

(metagenome) sequencing. Sediment sample processing was carried out as described in

(Pettersen et al., 2022). In brief, sediment DNA was extracted on a KingFisher Flex

automated liquid handling platform using the MagAttract PowerSoil DNA KF Kit (Qiagen).

The amplification of the V3-V4 region of the 16S rRNA gene was done using primers 341F

and 806R (Yu et al., 2005). Individual amplicon PCR products were dual-indexed using

different combinations of Illumina forward and reverse primers with unique barcode tags.

The library pools were sequenced a MiSeq v3 platform at the Norwegian Sequencing Centre

in Oslo. Shotgun sequencing was performed on an Illumina Novaseq 6000 platform at

Novogene, UK facility using the protocol stated in (Gourlé et al., 2018).

Subset B samples were subjected to 16S amplicon sequencing using both Illumina and ONT

sequencing platforms. Library preparation for Illumina sequencing of Subset B samples was

performed as described above. ONT sequencing libraries were prepared using the Oxford

Nanopore Sequencing kit (SQK-LSK114). The first amplification was performed using

modified 16S forward and reverse primers (Genetic Analysis, Norway) with the following

PCR program: 95°C for 15 min, 30 cycles of 95°C for 30 sec, 55°C for 30 sec and 72°C for 80

sec. Amplicons were purified with 1:1 Ampure XP paramagnetic beads (Beckman Coulter)

before indexing PCR. Between 10 and 60 ng of purified full-length 16S amplicons was added

to the PCR mixture, in addition to 0.2 µM of a unique PCR barcode (ONT cat.nr. EXP-

PBC096). The barcoding amplification was done using the following program: 95°C for 15

min, 12 cycles of 95°C for 30 sec, 62°C for 15 min and 65°C for 2 min, followed by 65°C for

10 min. Barcoded PCR products were concentration-normalized and pooled in equimolar

ratios, the library from which was purified with 1:1 Ampure XP beads before sequencing on

a MinION flow cell (Eurofins Genomics, Germany).

Amplicon data processing

The 16S amplicon sequencing data from subset A were processed with VSEARCH (Rognes et

al., 2016) for clustering into Operational Taxonomic Units (OTUs) using a 95% identity

threshold. ONT reads are longer than Illumina reads but still have error levels making it

difficult to go much above a 95% identity resolution for the 16S sequences. Because the

subset A Illumina sequence data will be used to generate reference sequences for error-

prone ONT sequences in subset B, a lower 95% resolution was chosen as opposed to a

higher value or a denoising approach. In addition, a recent study of very high diversity

environments like seafloor sediments demonstrated unstable compositional profiles when

using high-resolution denoising of illumina reads(Nilsen et al., 2024).

The Illumina paired-end reads from subset B were merged and de-replicated by VSEARCH,

but not clustered. Only reads with at least five copies per sample were used, since these are

most likely without sequencing error. Nanopore reads from the same subset B samples were

filtered based on their length, which ranged between 800 – 1 400 bp.

SILVA alignments

The OTU represents the sequences of around 420 - 440 bases, which corresponds to the

lengths of the amplicons we obtained by the primers used. First, all OTU representative

sequences were aligned against the 2 152 651 16S sequences in the SILVA 138.1 (Boratyn et

al., 2013). The SILVA sequences have a median length of 1 403 bases and range in length

from 900 to 4 000 bases. OTUs from subset A achieving a best match with ≥ 95% identity

over ≥ 95% of their length were marked as 'known', i.e. resembling something already

known from the SILVA database. The full-length corresponding matching SILVA sequences

were then retrieved as representative for' known 'OTUs (Fig. 1, box Q1).

It should be noted that many of the seed sequences used for reconstruction (see Section:

Reconstructing 16S using METASEED) derive from the same' known 'OTUs, obfuscating the

need for sequence reconstruction as matching SILVA sequences are already known. The

reconstruction was, however, still performed in order to test how closely the reconstructed

sequence could be matched to the corresponding SILVA sequence.

Reconstructing 16S using METASEED

OTUs were used as 'seed sequences' within the METASEED workflow to reconstruct a longer

section of the 16S rRNA gene from shotgun metagenome reads. METASEED collects reads

from the shotgun metagenome data, aligns them to the seed sequences and combines

assignment results with abundance information. For each seed, a reconstructed sequence is

reported if the sequence is ≥ 500 bp in length and the alignment occurs with ≥99% identity

to the corresponding seed.

Reconstructed sequences were subsequently aligned to the SILVA database, using the same

thresholds as above. As mentioned before (see Section: SILVA alignments), the 'known'

seeds could now be checked to verify whether the seed and its reconstructed sequence

matched the same SILVA sequence, or at least the same taxon. The most interesting

reconstructed sequences derive from cases where the SILVA database generated no matches

to the seed nor to its reconstructed sequence. These reconstructed sequences were then

added to our collection as new 16S sequences, representing new taxa relative to the SILVA

collection (Fig. 1, boxes Q2 and Q3).

Shotgun data assembly

The shotgun metagenome data from the 94 samples in subset A were filtered and merged

using BBduk and BBmerge (Bushnell, 2014) from BBmap v39.01. The assembly was done

using metaspades v3.15.5 (Nurk et al., 2017). From the assembled contigs we used Barrnap

v0.9 (https://github.com/tseemann/barrnap) to extract 16S sequences from the contigs

using length threshold ≥ 800 bases. As 16S sequences extracted from different samples may

be identical, all 16S sequences were clustered using VSEARCH v2.22.1, as described below,

to obtain a resolution of 95% identity (Fig. 1, box Q4).

Finally, we also included 16S centroids that didn't give any match to the above sequences.

These unmatched 16S sequences represent the lowest quality part of the reference

database (Fig. 1, box Q5).

Taxonomic assignments

Taxonomic assignments of all database sequences was obtained using the SINTAX algorithm

(Edgar, 2016) implemented in the VSEARCH software, where we used the GSR database and

the NCBI taxonomy (Molano et al., 2024). This means sequences were assigned to all ranks

to species, with confidence scores at each rank.

Phylogenetic placement of METASEEED and Barrnap derived sequences

Phylogenetic placement of the METASEED- and Barrnap- derived sequences were

determined since these are 'new '16S variants. First, all database sequences (excluding short

amplicons) were analysed with cmsearch in the infernal software v1.1.4 (Eddy, 2002)

and the secondary structure model for bacterial 16S downloaded from the Rfam database

(Kalvari et al., 2021). Sequences with an overall score (bitscore/length of the sequence)

below 0.9 were filtered out. A multiple alignment of the resulting sequences was created

using cmalign and computed evolutionary distances using Felsenstein-81 model

(Felsenstein, 1981) and the ape R package (Paradis & Schliep, 2019). These distances were

then used in a hierarchical clustering with average linkage and the sequences were split into

64 clusters. The clusters with at least 5 members were subject to a Fisher exact test for

enrichment of sequences originating from either METASEED or Barrnap, i.e. sequences not

previously found in SILVA. We considered a cluster to be enriched with new variants, if the

odds-ratio is greater than 1 and the p-value is less than 0.05. In these clusters there are

some 16S variants where the taxonomy is known to phylum rank, and we list the dominating

phyla in the enriched clusters based on this.

Database comparisons

As previously stated, one of the main motivations behind this work is to improve the

database that will be used to analyse amplicon data from Oxford Nanopore Technology.

EMU (Curry et al., 2022) is a standard software for profiling such data and we wanted to

compare our database to the default EMU database. We first used VSEARCH alignments to

assess the overlap between our generated database and the default EMU database.

Matches that failed to show at least 95% identity were not considered.

Next, we wanted to test how the Nanopore reads from the 90 subset B sediment samples

would be profiled using the EMU software in combination with its default database (EMU-

DB) as well as our custom database (AQUAeD-DB). In addition, we also tested the profiling of

Illumina de-replicated reads using standard VSEARCH alignments, i.e. pairwise alignment

between read and database sequences, and assigning to the best, an identity threshold of

95% was used. The read count matrices from the Illumina VSEARCH and Nanopore EMU tool

assignments for the AQUAeD-DB and EMU-DB databases were used to compute relative

abundance. Prior to the correlation analysis, missing database sequences from the Illumina

read count were added to the Nanopore read count matrix, and vice versa, with a count of

zero for each sample. This was done to resolve disparities in matrix lengths.

Results

Addition   of   SILVA   sequences   to   the   database

The full-length sequences were first collected from the SILVA database, by matching them

against the16S rRNA amplicon representative sequences, generated by conventional short-

read data processing. Using a 95% identity threshold, our 16S rRNA pipeline generated

14,393 OTUs from the subset A data. Each OTU has a representative sequence consisting of

400 - 440 bases in accord with the length of the V3-V4 amplicons. These sequences were

then aligned against the SILVA database using VSEARCH. In total 5 278 of OTUs showed at

least 95% identity matches with some of the SILVA sequences (identity distribution shown in

Fig. 2). This represents 84% - 96% of the total OTU abundance in these 94 samples. The 4

921 distinct SILVA sequences matched to the OTU sequences were collected and filtered

using `infernal cmsearch,` which resulted in 4 838 sequences. The SILVA sequences

were added to the database as full-length representatives of these OTUs and they belong to

the highest quality (Table 1, Q1) sequences in the database.

Fig. 2 Quality of OTU identifications. Frequency distribution of OTU sequences (y-axis) with

varying degrees of sequence similarity (x-axis) to SILVA database sequences. The green

columns in indicate the number of OTUs with a SILVA match while the grey columns indicate

no relevant match to SILVA database.

Reconstructing 16S rRNA genes using METASEED

All 14 393 OTUs were passed to the METASEED together with shotgun metagenome data in

order to re-construct near full-length 16S rRNA from the shorter amplicon sequences. The

screening process for the METASEED results is elaborated in Fig. S1. This analysis resulted in

3 525 16S rRNA sequences, of which 3 392 were ≥ 500 bases in length. Of these, 2 947 had

at least 99% sequence identity to their seeds, and were considered re-constructed 16S

sequences.

From the 2 947 reconstructed 16S sequences, 1 176 had no match to SILVA sequences

already collected (≥ 95% identity). Reconstructed 16S rRNA sequences with a minimum

length of 800 bases were regarded as high quality, while sequences 500 - 800 bases in length

were considered low quality. This resulted in 421 high quality and 755 low quality

reconstructed sequences. The total of 1 176 METASEED reconstructed sequences were

filtered using `cmsearch`, which led to the addition of 717 reconstructed sequences to the

database. The Q2 and Q3 category in Table 1 summarize these results. The increase in the

captured relative abundance when including Q1+Q2+Q3 sequences versus only Q1 (Fig. S2).

Addition of 16S rRNA from contigs using Barrnap

We also looked for 16S rRNA genes in the assembled contigs from the metagenome data

using the `barrnap` software. From the 94 samples, we found in total 9 839 short or longer

sequences recognized as 16S rRNA. The number of sequences was reduced to 1 149 when

applying a length threshold of at least 800 bp. Finally, since the same 16S rRNA sequence

may have been assembled and extracted from different samples, these sequences were

clustered using a 95% identity threshold, which reduced the number of sequences to 518.

Again, we used the `cmsearch` filtering, resulting in 320 high quality near full-length 16S

rRNA sequences, which were added to database. The total length of assembled contigs was

greater than 32 000 megabasepairs, implying that roughly one full-length high-quality 16S

rRNA was found for every 100 mega base-pairs of genomic sequence (Table 1, Q4).

Apart from the sequences collected from the above-mentioned sources, we also included 8

670 OTU sequences that did not match to any sequences added to the database from SILVA,

METASEED, and Barrnap when using a 95% identity threshold (Table 1, Q5). Including all

categories (Table 1), the final size of the AQUAeD database was 14 545 16S sequences.

Taxonomic assignments

All sequences in the AQUAeD database were classified using `SINTAX` (Fig. 3), with the

exception of sequences belonging to the Q1 category, which were copied from the SILVA

database with taxonomic assignments.

Fig. 3 The bars depict the fraction of classified sequences (Q2 - Q5) in the AQUAeD Database

using SINTAX (bootstrap value 0.8 and 0.5) and the GSR databases. The x-axis shows various

taxonomic ranks and y-axis shows the fraction of classified sequences.

Enrichment of new 16S variants

Sequences in categories Q2, Q3 and Q4 (Table 1) are new' 16S rRNA gene variants in the

sense that they are not included in the SILVA database. We wanted to see how these

distribute when clustered among the sequences in Q1 as well. The multiple sequence

alignment of these 5 875 sequences was followed by the evolutionary distance calculation,

cluster generation and using the Fisher test for enrichment of the new variants . From 64

clusters we found that 17 clusters were enriched by new' variants. The other members of

these enriched clusters predominantly consisted of phyla Planctomycetota, Myxococcota,

and Verrucomicrobiota (Table 2).

Database comparison

EMU software is the standard tool used for taxonomic profiling of Nanopore 16S reads. It

comes with a default database (EMU-DB), consisting of 49 243 sequences. From a

comparison between the EMU-DB and AQUAeD-DB, we observed that 18 584 (38%) EMU

database sequences had a match to 509 (3.5 %) AQUAeD database sequences when using a

95% identity threshold.

The 88 subset B samples resulted in a total of 973 758 dereplicated Illumina reads,

corresponding to 51 397 distinct sequences. The alignment of these reads (95% identity) to

the EMU-DB resulted in a match for 3% of the database sequences, comprising 52% of the

total Illumina reads. Alignment with the AQUAeD-DB yielded a match to 23% of the

database sequences, covering 99% of the total Illumina reads. The Nanopore amplicon reads

from the 88 subset B samples having length between 800 and 1 400 bases resulted in 4 972

056 reads. The analysis of Nanopore reads resulted in 3 967 AQUAeD database sequences,

accounting for approximately 27% of the database, while EMU-DB matched to 2 560

database sequences, accounting for approximately 5% of the database.

Fig. 4 The Database comparison. A) Illumina results for taxonomic assignments using

VSEARCH. B, Nanopore results for taxonomic assignments using EMU. B) the percentage

reads assigned exceeds 100% due to the fuzzy assignment by EMU which resulted in a total

'count 'of assigned reads that was larger than the number of reads. Pie charts represent

reads and database sequences are represented by bars. Shaded/striped regions in each pie

chart or bar indicate the fraction of reads assigned or the number of database sequences to

which they were assigned, respectively. Detailed information about alignments can be found

in Table S1.

A comparison of read count matrices revealed that 82% of the EMU-DB sequences in the

Illumina read count matrix were missing in the corresponding Nanopore read count matrix,

while 17% of the sequences in the Nanopore matrix were the missing from Illumina matrix.

For the AQUAeD-DB read count matrices, the overlap was notably better, with 45% of

Illumina sequences and 13% of Nanopore sequences missing from their counterparts,

indicating a lower level of missing database sequences. Correlation analysis demonstrated

high positive correlation between read count matrices for the AQUAeD-DB, whereas the

EMU-DB shows very weak negative correlation (Fig. 5). The relative mean abundance of

Illumina and Nanopore results for AQUAeD-DB   demonstrate a significant positive

correlation (Fig. 6).

Fig. 5 Correlation between Illumina and ONT read count matrices. Boxplots indicate

correlation coefficients for read count matrix similarity when using the AQUAeD or EMU

database.

Fig. 6 Relative mean abundance of Illumina versus Nanopore sequences for subset B

345 samples. The scatterplot depicts the log of relative mean abundance of Nanopore (y-axis)

346 versus the log of relative abundance of Illumina(x-axis) of AQUAeD-DB database sequences

347 across samples. Cases where either Illumina or Nanopore values were zero are omitted.

Discussion

The motivation for building the AQUAeD-DB was to enhance the use of Nanopore data in

the study of seafloor sediments. Thus, we wanted to see if the use of AQUAeD-DB has some

benefit compared to a standard database. Since the EMU tool is a popular choice for

Nanopore 16S data processing, we used its default database (EMU-DB) in our comparison. In

addition to using EMU for assigning the reads, we also used VSEARCH, which is typically used

for Illumina data, just for comparison. The VSEARCH alignment directly between the

AQUAeD-DB and EMU-DB reveals that 3.5% of the AQUAeD database could represent 38%

of the EMU database, implying that the AQUAeD-DB, despite being smaller in the number of

sequences, covers an order of magnitude more seafloor diversity compared to EMU-DB.

The AQUAeD-DB comprises of 14 545 16S sequences that were collected from seafloor

sediments (Fig. 1). The Illumina amplicon sequencing of subset A seafloor samples yielded

14 393 centroid sequences when clustered at the coarse level of 95% identity.

Approximately 40% of these sequences matched the SILVA database sequences, even with

relatively liberal identity threshold of 95% (Fig. 2). However, this set of matching SILVA

sequences, the Q1, makes up the backbone of our AQUAeD-DB. To add more longer

sequences to the database, we used our METASEED tool (Philip et al., 2024) to extend the

amplicon sequences, making use of the Illumina metagenome data from the same samples

(subset A). This resulted in the Q2 and Q3 part of the database. The total abundance

covered by the database sequences in each sample did not increase appreciably by adding

these sequences, but the increase was larger for the exceptionally high diversity samples

(high Shannon index of 5-7, Fig. S2). This typically allowed us to obtain 16S sequences for the

low-abundance and ecologically significant organisms from phyla like Planctomycetota (table

2). In the Q4 part of the database, we tried to collect long 16S directly from metagenome

assembled reads. The fact that we only collect one such sequence per 100 megabases of

assembled contigs indicates how difficult it is to obtain such information from the assembly

of short read metagenome data. The limitations of metagenome assembly, specifically the

challenges in retrieving 16S sequences, have been previously addressed (Hiseni et al., 2022;

Mise & Iwasaki, 2022; Shaiber & Eren, 2019).

The AQUAeD-DB can be roughly split into two parts: The 'known 'sequences (Q1) and the

'new variants '(Q2-Q5, Fig 3). The fractions of taxonomic assignments were generally low,

especially for the Q3-Q5 sequences. This emphasizes that new sequence variants in the

AQUAeD database (Q2-Q5) have poor representation in a public database, like the GSR used

here. Considering the poor taxonomic information, we wanted to see if phylogenetic

distances between these new variants, and the Q1 sequences indicate a grouping of these

new sequences, or if they are scattered around in the phylogeny. By clustering based on

phylogenetic distances, we found several clusters where these new variants are over-

represented, which include significant marine microorganisms from the phyla

Planctomycetota, Myxococcota, and Verrucomicrobiota (Dell'Anno et al., 2021; Freitas et al.,

2012; Wu et al., 2024)

A clear difference was evident when we did the assignment with both VSEARCH, which is

normally used for Illumina data, and EMU, which is the standard choice for Nanopore (Fig 4).

The VSEARCH tool aligned the high-precision Illumina reads, with EMU-DB assigning almost

half of the reads and spreading them out among only 1 474 of the nearly 50 000 database

sequences. For AQUAeD-DB, almost all reads are assigned to 3 301 database sequences. The

Nanopore reads showed different results compared to the VSEARCH assignment of Illumina

data, especially in terms of database coverage. One of the major contributing factors can be

the significantly higher noise levels compared to Illumina data; this leads to reads being

distributed across a much larger number of sequences in both databases (Magi et al., 2017;

Santos et al., 2020). Another difference lies in the fact that the EMU tool assigns more than

100% of the reads, which likely indicates an artifact of the fuzzy assignment process

(Szoboszlay et al., 2023). From VSEARCH+Illumina and EMU+Nanopore, we see how more

organisms are being given read counts in the AQUAeD-DB compared to the EMU-DB. This is

the effect of AQUAeD-DB containing more relevant taxa for seafloor data, while EMU-DB is a

generic collection and probably more relevant for use in medical research.

The previously stated percentage of missing database sequences in the read count matrices

for Illumina and Nanopore indicate that the EMU-DB analysis results exhibit greater

disparities compared to the AQUAeD-DB results. A clear positive correlation between

Illumina and Nanopore relative abundance were revealed in AQUAeD-DB whereas EMU-DB

shows very inconsistent and poor results (Fig 5). From these results it is evident that

AQUAeD-DB results in more reliable and consistent assignments for both Illumina and ONT

analysis. These findings were supported by the broad range of diversity, especially for the

low- abundance taxa that the AQUAeD-DB database was able to cover (Fig 6). These

observations once again underline the benefits of AQUAeD-DB and emphasize the potential

implementation for Nanopore based environmental studies.

These findings indicated that AQUAeD-DB have illustrated a significant contribution in

environmental analysis using ONT sequencing data. The database enrichment analysis

implies that significant marine microbiota is represented in newly developed AQUAeD

database, comprising both existing sequences and novel sequences from the

aforementioned strategies. This optimizes taxonomical assignment and provides potential

for more explicit environmental context. The study emphasizes the potential, and necessity

of such databases in facilitating the implementation of the cutting-edge technologies, thus

enabling the comprehensive environmental assessments.

Conclusion

The newly developed AQUAeD-DB database is a specialized database for 16S rRNA gene

sequences from highly diverse and complex environmental samples. The primary objective

of AQUAeD-DB was to provide a better reference database for identification of Nanopore

sequences targeting microbial communities in North Atlantic seafloor sediments. Based on

results of the enrichment study and database comparison, this database is substantially

diverse, and efficiently covers an expanded range of microorganisms that are poorly

represented in other databases. This study was designed as a pilot project with the potential

for future scalability of the database to incorporate additional 16S rRNA gene sequences.

This expansion will be rooted in ongoing efforts to improve environmental surveys and

monitoring using microbial profiles.

Acknowledgements

436 We express our gratitude to the fish farmers that provided samples for this research. We

437 would also like to thank the employees of STIM, Aqua Kompetanse, and Akvaplan-NIVA that

438 performed the sampling.

Data availability

The   AQUAED-DB   and   taxonomy   details   are   publicly   available

(https://arken.nmbu.no/~pmelcy/share/AQUAED-DB/). All raw sequences used in this study

have been deposited in the NCBI BioProject database. Illumina 16S rRNA sequences from

subset A and subset B is a part of the project with the accession code PRJNA1128851. The

raw shotgun data of subset A has been uploaded with the accession code PRJNA1128851.

The raw Nanopore sequencing data for subset B has been uploaded with the accession code

PRJNA1153974. The metadata files for all datasets are available in

https://arken.nmbu.no/~pmelcy/share/AQUAED-DB/metadata/

Author contributions

Knut Rudi, Lars Snipen and Melcy Philip conceived the study; Jessica Louise Ray, Morten

Stokkan, Ragnhild Pettersen and Sanna Majaneva design the collection of sediment samples;

Tonje Nilsen performed all lab experiments; Lars Snipen and Melcy Philip performed all

analyses, with conceptual input from Knut Rudi; Knut Rudi, Lars Snipen and Melcy Philip

contributed to interpretation of results; Knut Rudi, Lars Snipen and Melcy Philip drafted the

manuscript. All author(s) read and commented on the manuscript.

Funding

459 This study is a part of the project: "AQUAeD", funded by The Research Council of Norway

460 (Project Number: 320076).

Competing interests

At the time of manuscript writing, Sanna Majaneva, Ragnhild Pettersen were employed by

company Akvaplan-niva AS. Jessica Louise Ray was employed by company Aqua Kompetanse

AS and Morten Stokkan was employed by company STIM AS. The remaining authors declare

that the research was conducted in the absence of any commercial or financial relationships

that could be construed as a potential conflict of interest.

Table 1: The AQUAeD-DB contains sequences of 5 quality categories, from the best (Q1) to the poorest (Q5)

Category descriptions are provided in Materials and Methods. Lengths are shown in basepairs (bp).

Category	Origin	Number of sequences	Minimum length	Median length	Maximum length
Q1	SILVA	4838	900	1461	1668
Q2	High Quality METASEED	311	800	918	1522
Q3	Low Quality METASEED	406	501	688.5	799
Q4	16S from contigs	320	800	924	1577
Q5	Amplicons	8670	288	426	438

472 **Table 2:** Enrichment of 'new' 16S variants. Results of Fisher tests for 17 'new variants 'enriched clusters when
473 k=64.

Odds ratio	Cluster size	New variants fraction	Dominant phylum
7.06	20	0.6	Planctomycetota
7.01	5	0.6	Desulfobacterota,Proteobacteria

6.56	12	0.58	Patescibacteria
4.015	13	0.46	Patescibacteria
3.97	35	0.46	Spirochaetota
3.27	39	0.41	Verrucomicrobiota
3.01	23	0.39	Spirochaetota
2.71	30	0.37	Verrucomicrobiota
2.59	62	0.35	Chloroflexi
2.58	114	0.35	Verrucomicrobiota
2.55	153	0.35	Planctomycetota
2.54	37	0.35	Firmicutes
2.34	30	0.33	Firmicutes
2.34	27	0.33	Fibrobacterota
2.13	61	0.31	Planctomycetota
2.07	72	0.31	Verrucomicrobiota
1.61	241	0.25	Myxococcota

477 **Supplementary data**

Fig. S1 The Fig. depicts the workflow of the screening process in METASEED analysis

Fig. S2 The Fig. depicts the sum of relative OTU abundance (y-axis) in each of the ninety-four

samples covered by the SILVA set only (orange dots) and the increase obtained by adding the

reconstructed sequences as well (green dots). The x-axis shows Shannon diversity for each

sample. The distance between a brown and green dot shows the increase in relative

abundance as we add the Q1 and Q2 sequences to the collection.

Table S1: The table presents the VSEARCH alignment results of Illumina dereplicated reads and EMU

assignment results of Nanopore reads against EMU database and AQUAeD database.

	Total	**AQUAeD (14545)**	**EMU (49243)**
Distinct Illumina Sequences	51397	3301	1474
Corresponding Illumina Reads	973758	966477	358760
Illumina Reads Percent	100	99	52
Number of total reads and unique database sequence matches (Nanopore)	49 72 056	3967	2560
Fraction of database sequences (Nanopore)	100	27	5

References

493   Amarasinghe, S. L., Su, S., Dong, X., Zappia, L., Ritchie, M. E., & Gouil, Q. (2020). Opportunities and
494   challenges in long-read sequencing data analysis. *Genome Biology*, *21*(1), 30.
495   https://doi.org/10.1186/s13059-020-1935-5
496   Benítez-Páez, A., Portune, K. J., & Sanz, Y. (2016). Species-level resolution of 16S rRNA gene
497   amplicons sequenced through the MinION™ portable nanopore sequencer. *GigaScience*,
498   *5*(1). https://doi.org/10.1186/s13742-016-0111-z
499   Boratyn, G. M., Camacho, C., Cooper, P. S., Coulouris, G., Fong, A., Ma, N., Madden, T. L., Matten, W.
500   T., McGinnis, S. D., Merezhuk, Y., Raytselis, Y., Sayers, E. W., Tao, T., Ye, J., & Zaretskaya, I.

(2013). BLAST: a more efficient report with usability improvements. *Nucleic Acids Research*,
41(W1), W29-W33. https://doi.org/10.1093/nar/gkt282
Bushnell, B. (2014). *BBMap: a fast, accurate, splice-aware aligner*.
Canals, M., Pham, C. K., Bergmann, M., Gutow, L., Hanke, G., van Sebille, E., Angiolillo, M., Buhl-
Mortensen, L., Cau, A., Ioakeimidis, C., Kammann, U., Lundsten, L., Papatheodorou, G.,
Purser, A., Sanchez-Vidal, A., Schulz, M., Vinci, M., Chiba, S., Galgani, F.,...Giorgetti, A. (2021).
The quest for seafloor macrolitter: a critical review of background knowledge, current
methods and future prospects. *Environmental Research Letters*, *16*(2), 023001.
https://doi.org/10.1088/1748-9326/abc6d4
Curry, K. D., Wang, Q., Nute, M. G., Tyshaieva, A., Reeves, E., Soriano, S., Wu, Q., Graeber, E., Finzer,
P., Mendling, W., Savidge, T., Villapol, S., Dilthey, A., & Treangen, T. J. (2022). Emu: species-
level microbial community profiling of full-length 16S rRNA Oxford Nanopore sequencing
data. *Nature Methods*, *19*(7), 845-853. https://doi.org/10.1038/s41592-022-01520-4
Dell'Anno, F., Rastelli, E., Tangherlini, M., Corinaldesi, C., Sansone, C., Brunet, C., Balzano, S., Ianora,
A., Musco, L., Montereali, M. R., & Dell'Anno, A. (2021). Highly Contaminated Marine
Sediments Can Host Rare Bacterial Taxa Potentially Useful for Bioremediation [Original
Research]. *Frontiers in Microbiology*, *12*. https://doi.org/10.3389/fmicb.2021.584850
DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., Huber, T., Dalevi, D., Hu,
P., & Andersen, G. L. (2006). Greengenes, a chimera-checked 16S rRNA gene database and
workbench compatible with ARB. *Applied and Environmental Microbiology*, *72*(7), 5069-
5072.
Dueholm, M. K. D., Nierychlo, M., Andersen, K. S., Rudkjøbing, V., Knutsson, S., Arriaga, S., Bakke, R.,
Boon, N., Bux, F., Christensson, M., Chua, A. S. M., Curtis, T. P., Cytryn, E., Erijman, L.,
Etchebehere, C., Fatta-Kassinos, D., Frigon, D., Garcia-Chaves, M. C., Gu, A. Z.,...Mi, D. A. S. G.
C. (2022). MiDAS 4: A global catalogue of full-length 16S rRNA gene sequences and taxonomy
for studies of bacterial communities in wastewater treatment plants. *Nature
Communications*, *13*(1), 1908. https://doi.org/10.1038/s41467-022-29438-7
Eddy, S. R. (2002). A memory-efficient dynamic programming algorithm for optimal alignment of a
sequence to an RNA secondary structure. *BMC Bioinformatics*, *3*(1), 18.
https://doi.org/10.1186/1471-2105-3-18
Edgar, R. (2016). SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS sequences. In:
bioRxiv.
F. Escapa, I., Huang, Y., Chen, T., Lin, M., Kokaras, A., Dewhirst, F. E., & Lemon, K. P. (2020).
Construction of habitat-specific training sets to achieve species-level assignment in 16S rRNA
gene datasets. *Microbiome*, *8*(1), 65. https://doi.org/10.1186/s40168-020-00841-w
Federhen, S. (2011). The NCBI Taxonomy database. *Nucleic Acids Research*, *40*(D1), D136-D143.
https://doi.org/10.1093/nar/gkr1178
Felsenstein, J. (1981). Evolutionary trees from DNA sequences: a maximum likelihood approach. *J
Mol Evol*, *17*(6), 368-376. https://doi.org/10.1007/bf01734359
Freitas, S., Hatosy, S., Fuhrman, J. A., Huse, S. M., Welch, D. B., Sogin, M. L., & Martiny, A. C. (2012).
Global distribution and diversity of marine Verrucomicrobia. *Isme j*, *6*(8), 1499-1505.
https://doi.org/10.1038/ismej.2012.3
Ghate, S. D., Shastry, R. P., Arun, A. B., & Rekha, P. D. (2021). Unraveling the bacterial community
composition across aquatic sediments in the Southwestern coast of India by employing high-
throughput 16S rRNA gene sequencing. *Regional Studies in Marine Science*, *46*, 101890.
https://doi.org/https://doi.org/10.1016/j.rsma.2021.101890
Henderson, G., Yilmaz, P., Kumar, S., Forster, R. J., Kelly, W. J., Leahy, S. C., Guan, L. L., & Janssen, P. H.
(2019). Improved taxonomic assignment of rumen bacterial 16S rRNA sequences using a
revised SILVA taxonomic framework. *PeerJ*, *7*, e6496. https://doi.org/10.7717/peerj.6496

Hiseni, P., Snipen, L., Wilson, R. C., Furu, K., & Rudi, K. (2022). Questioning the Quality of 16S rRNA
Gene Sequences Derived From Human Gut Metagenome-Assembled Genomes [Opinion].
Frontiers in Microbiology, *12*. https://doi.org/10.3389/fmicb.2021.822301
Hoshino, T., Doi, H., Uramoto, G. I., Wörmer, L., Adhikari, R. R., Xiao, N., Morono, Y., D'Hondt, S.,
Hinrichs, K. U., & Inagaki, F. (2020). Global diversity of microbial communities in marine
sediment. *Proc Natl Acad Sci U S A*, *117*(44), 27587-27597.
https://doi.org/10.1073/pnas.1919139117
Kalvari, I., Nawrocki, E. P., Ontiveros-Palacios, N., Argasinska, J., Lamkiewicz, K., Marz, M., Griffiths-
Jones, S., Toffano-Nioche, C., Gautheret, D., & Weinberg, Z. (2021). Rfam 14: expanded
coverage of metagenomic, viral and microRNA families. *Nucleic Acids Research*, *49*(D1),
D192-D200.
Kozlov, A. M., Zhang, J., Yilmaz, P., Glöckner, F. O., & Stamatakis, A. (2016). Phylogeny-aware
identification and correction of taxonomically mislabeled sequences. *Nucleic Acids Research*,
44(11), 5022-5033.
Lemoinne, A., Dirberg, G., Georges, M., & Robinet, T. (2023). Fine-scale congruence in bacterial
community structure from marine sediments sequenced by short-reads on Illumina and
long-reads on Nanopore.
Madsen, E. L. (2011). Microorganisms and their roles in fundamental biogeochemical cycles. *Current*
Opinion in Biotechnology, *22*(3), 456-464.
https://doi.org/https://doi.org/10.1016/j.copbio.2011.01.008
Magi, A., Semeraro, R., Mingrino, A., Giusti, B., & D'Aurizio, R. (2017). Nanopore sequencing data
analysis: state of the art, applications and challenges. *Briefings in Bioinformatics*, *19*(6),
1256-1272. https://doi.org/10.1093/bib/bbx062
Marlow, J., Steele, J., Case, D., Connon, S., Levin, L., & Orphan, V. (2014). Microbial abundance and
diversity patterns associated with sediments and carbonates from the methane seep
environments of Hydrate Ridge, OR. *Frontiers in Marine Science*, *1*.
https://doi.org/10.3389/fmars.2014.00044
McDonald, D., Jiang, Y., Balaban, M., Cantrell, K., Zhu, Q., Gonzalez, A., Morton, J. T., Nicolaou, G.,
Parks, D. H., Karst, S. M., Albertsen, M., Hugenholtz, P., DeSantis, T., Song, S. J., Bartko, A.,
Havulinna, A. S., Jousilahti, P., Cheng, S., Inouye, M.,…Knight, R. (2024). Author Correction:
Greengenes2 unifies microbial data in a single reference tree. *Nature Biotechnology*, *42*(5),
813-813. https://doi.org/10.1038/s41587-023-02026-w
Meola, M., Rifa, E., Shani, N., Delbès, C., Berthoud, H., & Chassard, C. (2019). DAIRYdb: a manually
curated reference database for improved taxonomy annotation of 16S rRNA gene sequences
from dairy products. *BMC Genomics*, *20*(1), 560. https://doi.org/10.1186/s12864-019-5914-8
Mise, K., & Iwasaki, W. (2022). Unexpected absence of ribosomal protein genes from metagenome-
assembled genomes. *ISME Communications*, *2*(1). https://doi.org/10.1038/s43705-022-
00204-6
Molano, L.-A. G., Vega-Abellaneda, S., & Manichanh, C. (2024). GSR-DB: a manually curated and
optimized taxonomical database for 16S rRNA amplicon analysis. *mSystems*, *9*(2), e00950-
00923. https://doi.org/doi:10.1128/msystems.00950-23
Nierychlo, M., Andersen, K. S., Xu, Y., Green, N., Jiang, C., Albertsen, M., Dueholm, M. S., & Nielsen, P.
H. (2020). MiDAS 3: An ecosystem-specific reference database, taxonomy and knowledge
platform for activated sludge and anaerobic digesters reveals species-level microbiome
composition of activated sludge. *Water Research*, *182*, 115955.
https://doi.org/https://doi.org/10.1016/j.watres.2020.115955
Nilsen, T., Snipen, L.-G., Angell, I. L., Keeley, N. B., Majaneva, S., Pettersen, R., & Rudi, K. (2024).
Swarm and UNOISE outperform DADA2 and Deblur for denoising high-diversity marine
seafloor samples. *ISME Communications*. https://doi.org/10.1093/ismeco/ycae071
Nurk, S., Meleshko, D., Korobeynikov, A., & Pevzner, P. A. (2017). metaSPAdes: a new versatile
metagenomic assembler. *Genome research*, *27*(5), 824-834.

601 Overgaard, C. K., Tao, K., Zhang, S., Christensen, B. T., Blahovska, Z., Radutoiu, S., Kelly, S., &
602   Dueholm, M. K. D. (2022). Application of ecosystem-specific reference databases for
603   increased taxonomic resolution in soil microbial profiling. *Front Microbiol*, *13*, 942396.
604   https://doi.org/10.3389/fmicb.2022.942396

605 Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary
606   analyses in R. *Bioinformatics*, *35*(3), 526-528.

607 Patil, P. K., Baskaran, V., Vinay, T. N., Avunje, S., Leo-Antony, M., Shekhar, M. S., Alavandi, S. V., &
608   Vijayan, K. K. (2021). Abundance, community structure and diversity of nitrifying bacterial
609   enrichments from low and high saline brackishwater environments. *Letters in Applied*
610   *Microbiology*, *73*(1), 96-106. https://doi.org/10.1111/lam.13480

611 Pettersen, R., Ormaasen, I., Angell, I. L., Keeley, N. B., Lindseth, A., Snipen, L., & Rudi, K. (2022).
612   Bimodal distribution of seafloor microbiota diversity and function are associated with marine
613   aquaculture. *Marine Genomics*, *66*, 100991.
614   https://doi.org/https://doi.org/10.1016/j.margen.2022.100991

615 Philip, M., Rudi, K., Ormaasen, I., Angell, I. L., Pettersen, R., Keeley, N. B., & Snipen, L.-G. (2024).
616   METASEED: a novel approach to full-length 16S rRNA gene reconstruction from short read
617   data. *BMC Bioinformatics*, *25*(1), 237. https://doi.org/10.1186/s12859-024-05837-z

618 Ritari, J., Salojärvi, J., Lahti, L., & de Vos, W. M. (2015). Improved taxonomic assignment of human
619   intestinal 16S rRNA sequences by a dedicated reference database. *BMC Genomics*, *16*(1),
620   1056. https://doi.org/10.1186/s12864-015-2265-y

621 Rodríguez-Pérez, H., Ciuffreda, L., & Flores, C. (2020). NanoCLUST: a species-level analysis of 16S
622   rRNA nanopore sequencing data. *Bioinformatics*, *37*(11), 1600-1601.
623   https://doi.org/10.1093/bioinformatics/btaa900

624 Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: a versatile open source tool
625   for metagenomics. *PeerJ*, *4*, e2584. https://doi.org/10.7717/peerj.2584

626 Santos, A., van Aerle, R., Barrientos, L., & Martinez-Urtaza, J. (2020). Computational methods for 16S
627   metabarcoding studies using Nanopore sequencing data. *Computational and Structural*
628   *Biotechnology Journal*, *18*, 296-305.
629   https://doi.org/https://doi.org/10.1016/j.csbj.2020.01.005

630 Shaiber, A., & Eren, A. M. (2019). Composite Metagenome-Assembled Genomes Reduce the Quality
631   of Public Genome Repositories. *mBio*, *10*(3), 10.1128/mbio.00725-00719.
632   https://doi.org/doi:10.1128/mbio.00725-19

633 Szoboszlay, M., Schramm, L., Pinzauti, D., Scerri, J., Sandionigi, A., & Biazzo, M. (2023). Nanopore Is
634   Preferable over Illumina for 16S Amplicon Sequencing of the Gut Microbiota When Species-
635   Level Taxonomic Classification, Accurate Estimation of Richness, or Focus on Rare Taxa Is
636   Required. *Microorganisms*, *11*(3), 804. https://www.mdpi.com/2076-2607/11/3/804

637 Tatsuhiko Hoshinoa, H. D., Go-Ichiro Uramotoa, Lars Wörmerc, Rishi R. Adhikari, Nan Xiaoa,e,, Yuki
638   Moronoa, Steven D'Hondtf, Kai-Uwe Hinrichsc , Fumio Inagaki. (2020). Global diversity of
639   microbial communities in

640 marine sediment.

641 Wu, J., Hong, Y., Chang, X., Jiao, L., Li, Y., Liu, X., Xie, H., & Gu, J.-D. (2019). Unexpectedly high
642   diversity of anammox bacteria detected in deep-sea surface sediments of the South China
643   Sea. *FEMS Microbiology Ecology*, *95*(3). https://doi.org/10.1093/femsec/fiz013

644 Wu, X.-T., He, Y.-Q., Qiu, M., Li, G.-X., Ou, X.-B., Wu, Y.-B., & Bao, P. (2024). Adaptive Changes in
645   Bacterial and Archaeal Community Structures and Functions Occur under Primordial
646   Conditions. *ACS Earth and Space Chemistry*, *8*(7), 1470-1479.
647   https://doi.org/10.1021/acsearthspacechem.4c00089

648 Yang, C.-c., & Iwasaki, W. (2014). MetaMetaDB: A Database and Analytic System for Investigating
649   Microbial Habitability. *PLOS ONE*, *9*(1), e87126.
650   https://doi.org/10.1371/journal.pone.0087126

Yang, N., Tian, C., Lv, Y., Hou, J., Yang, Z., Xiao, X., & Zhang, Y. (2022). Novel primers for 16S rRNA
gene-based archaeal and bacterial community analysis in oceanic trench sediments. *Applied*
Microbiology and Biotechnology, 106(7), 2795-2809. https://doi.org/10.1007/s00253-022-
11893-3
Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J., Ludwig,
W., & Glöckner, F. O. (2013). The SILVA and "All-species Living Tree Project (LTP)" taxonomic
frameworks. *Nucleic Acids Research, 42*(D1), D643-D648.
https://doi.org/10.1093/nar/gkt1209
Yu, Y., Lee, C., Kim, J., & Hwang, S. (2005). Group-specific primer and probe sets to detect
methanogenic communities using quantitative real-time polymerase chain reaction.
Biotechnology and Bioengineering, 89(6), 670-679.
https://doi.org/https://doi.org/10.1002/bit.20347
Zorz, J., Li, C., Chakraborty, A., Gittins, D. A., Surcon, T., Morrison, N., Bennett, R., MacDonald, A., &
Hubert, C. R. J. (2023). SituSeq: an offline protocol for rapid and remote Nanopore 16S rRNA
amplicon sequence analysis. *ISME Communications, 3*(1), 33.
https://doi.org/10.1038/s43705-023-00239-3
667