A Targeted Reference Database for Improved Analysis of Environmental 16S rRNA Oxford Nanopore Sequencing Data

Melcy Philip[1], Tonje Nilsen[1] Sanna Majaneva[2], Ragnhild Pettersen[2], Morten Stokkan[3], Jessica Louise Ray[4], Nigel Keeley[5], Knut Rudi[1], Lars-Gustav Snipen[1]

[1]Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences, Ås (formerly known as viken), Norway

[2]Akvaplan-niva, Framsenteret, P.O. Box 6606, Stakkevollan, 9296 Tromsø, Norway

[3]STIM department Bergen, Thormøhlens gate 55, 5006 Bergen, Norway

[4]Aqua Kompetanse AS, Storlavika 7, 7770 Flatanger, Norway

[5]Institute of Marine Research, Tromsø, Norway

Corresponding author: Melcy Philip, Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences, Christian Magnus Falsens vei 18, 1433, Ås (formerly known as viken), Norway (philip.melcy@nmbu.no)

Knut Rudi: knut.rudi@nmbu.no,

Tonje Nilsen: tonje-n2@online.no,

Sanna Majaneva: sam@akvaplan.niva.no,

Ragnhild Pettersen: rap@akvaplan.niva.no,

Morten Stokkan: morten.stokkan@stim.no

Jessica Louise Ray: jessica@aqua-kompetanse.no

Nigel B Keeley: nigel.brian.keeley@hi.no,

Lars-Gustav Snipen: lars.snipen@nmbu.no

Abstract

The Oxford Nanopore Technologies (ONT) sequencing platform is compact and efficient, making it suitable for rapid biodiversity assessments in remote areas. Despite its long reads, ONT has a higher error rate compared to other platforms, necessitating high-quality reference databases for accurate taxonomic assignments. However, the absence of targeted databases for underexplored habitats, such as the seafloor, limits ONT's broader applicability for exploratory analysis.

To address this, we propose an approach for building environmentally-targeted databases to improve 16S rRNA gene (16S) analysis using Oxford Nanopore Technologies (ONT), using seafloor sediment samples from the Norwegian coast as an example. We started by using Illumina short-read data to create a database of full-length or near full-length 16S sequences from seafloor samples. Initially, amplicons are mapped to the SILVA database, with matches added to our database. Unmatched amplicons are reconstructed using METASEED and Barrnap methodologies with amplicon and metagenome data. Finally, if the previous strategies did not succeed, we included the short-read sequences in the database. This resulted in AQUAeD-DB, which contains 14 545 16S sequences clustered at 95% identity. Comparative database analysis reveal that AQUAeD-DB provides consistent results for both Illumina and Nanopore read assignments (median correlation coefficient: 0.50), whereas a standard database showed a substantially weaker correlation. These findings also emphasize its potential to recognize both high and low-abundance taxa, which could be key indicators in environmental studies. This work highlights the necessity of targeted databases for environmental analysis, especially for ONT-based studies, and lays foundations for future extension of the database.

46   **Keywords:** 16S rRNA, metagenomics, seafloor sediment, database, Nanopore, ONT, Illumina,

47   biodiversity assessment

49   Introduction

The Earth's surface is primarily covered by oceans, yet we know little about the seafloor, or

its microbiota (Carrasco De La Cruz, 2021; Thurber et al., 2014). Microorganisms play a key

role in biogeochemical cycling in the global ocean thus the potential importance of

microbiota to healthy ecosystem function cannot be overstated (Madsen, 2011). Many

aspects of our understanding of soft-sediment microbiota is based on short read amplicon

sequencing of the 16S rRNA gene V3 and V4 regions (Ghate et al., 2021; Hoshino et al.,

2020; Yang et al., 2022). Advancements in long read sequencing, including Oxford Nanopore

Technology (ONT), have enabled rapid analysis of environmental microbiota (Zorz et al.,

2023). However, the relatively high error rate of ONT sequencing technology remains a

persistent challenge, rendering ONT strategies dependent on reference databases

(Amarasinghe et al., 2020). For effective and robust use of ONT technology that generates

reads spanning nearly the entire 16S gene, databases of nearly full-length 16S rRNA gene

sequences with comprehensive representation of habitat-relevant taxa are needed.

The most widely used and recognized 16S databases are the curated Greengenes and SILVA

databases, as well as the non-curated NCBI 16S rRNA database (Federhen, 2011; McDonald

et al., 2024; Yilmaz et al., 2013). However, these databases cover only a limited range of

environment-associated or marine-related microorganisms, despite housing a large number

of reference sequences. Moreover, recent studies have revealed other underlying issues

such as taxonomic gaps, partial annotation and redundancy (Cabezas et al., 2024). For

instance, the 16S component of the GenBank database consists of 43% chimeric sequences,

the presence of which may confound accurate taxonomic assignments (DeSantis et al., 2006;

Frenkel-Morgenstern et al., 2012). Furthermore, despite the use of quality control tools,

public 16S reference databases still contain 0.2% to 2.5% misidentified species, leading to

taxonomic misassignments (Kozlov et al., 2016). In addition, there is a major bias towards

characterization of human-associated and highly abundant bacteria, while most of

environmental microbial diversity remains uncharacterized (Yang & Iwasaki, 2014). In this

context, we fail to notice potentially low-abundant microorganisms, such as nitrifying

bacteria, anammox bacteria and methanotrophs, that may be crucial for reflecting the

health of the environment (Marlow et al., 2014; Patil et al., 2021; Wu et al., 2019). Benthic

microbes, in particular, are underrepresented in databases and our understanding of them is

constrained by very few studies (Tatsuhiko Hoshinoa, 2020). The current scenario thus

emphasizes the need for a specialized database for seafloor microorganisms, since

environmental microbial communities are often missing from current databases.

Studies reveal that employing targeted databases can improve the accuracy and coverage of

amplicon-based microbial profiling (Dueholm et al., 2022; Meola et al., 2019; Overgaard et

al., 2022). Although ONT sequencing technology is advancing, there are only few

bioinformatics tools for ONT 16S profiling (Ciuffreda et al., 2021; Santos et al., 2020; Sissel et

al., 2015) apart from the workflow offered by Oxford Nanopore. There are several ongoing

efforts to improve taxonomic assignments within ONT-driven studies, with the EMU tool

showing promise as an extensively used approach (Curry et al., 2022; Rodríguez-Pérez et al.,

2020; Zorz et al., 2023). However, all these tools, require a good reference database for

improved taxonomic assignment. This underlines the necessity of an environment-specific or

specialized reference database and the associated potential benefits. (F. Escapa et al., 2020;

Henderson et al., 2019; Nierychlo et al., 2020; Ritari et al., 2015). Studies have focused on

achieving high resolution using Nanopore, given its longer reads covering most of the 16S

gene, whereas our study aims to capture the broadest possible diversity from environmental

samples (Benítez-Páez et al., 2016; Lemoinne et al., 2023).

The aim of this work is to present an approach for developing a targeted 16S database

specifically designed for seafloor sediment or similar environments. This database facilitates

microbial profiling and ecosystem status prediction using full-length 16S (V2-V9) ONT

sequencing . We have called this database "AQUAeD-DB" in acknowledgment of the

research project "AQUAeD" which provided environmental samples for sequencing analysis

(see Acknowledgements). Our strategy (Fig. 1) makes use of short-read amplicon data in

combination with existing databases, as well as whole genome shotgun sequencing in

combination with our recently developed method METASEED (Philip et al., 2024). We also

assess and present the effectiveness of this targeted database in comparison to the existing

tools and databases for taxonomic assignment of 16S ONT amplicons. As an example, we

demonstrate this methodology using sediment samples from the Norwegian coast, with

plans to apply it in the monitoring of seafloor sediments to assess the impacts of

aquaculture sites across Norway. Additionally, this approach is adaptable for broader and

global studies in the future.

Fig. 1 The16S database generation. a) The seafloor samples used in this study were collected

from various sites along the Norwegian coast. b) The samples for database generation were

sequenced using Illumina, providing shotgun and 16S rRNA amplicon reads, c) Various

methods are used to collect or rebuild longer 16S rRNA sequences, which are then graded

into different quality levels, from Q1 (best) to Q5 (poorest). These sequences are added to

our reference database e) and used to for taxonomic assignment of ONT sequences

118 amplified   from   geographically   proximal   seafloor   sediment   material.

Materials and Methods

Data

In this study, we used total of 182 marine sediment samples collected along a north-south gradient along the Norwegian coast as mentioned previously (Nilsen et al., 2024; Pettersen et al., 2022). The sample information and collection details can be found in this study (Nilsen et al., 2024). The data was randomly divided into two subsets: subset A consisted of 94 samples and was used for the database construction; subset B consisted of 88 samples and was used for database comparison.

Subset A was subject to Illumina sequencing, both 16S amplicon and whole genome (metagenome) sequencing. Sediment sample processing was carried out as described in (Pettersen et al., 2022). In brief, sediment DNA was extracted on a KingFisher Flex automated liquid handling platform using the MagAttract PowerSoil DNA KF Kit (Qiagen). The amplification of the V3-V4 region of the 16S rRNA gene was done using primers 341F and 806R (Yu et al., 2005). Individual amplicon PCR products were dual-indexed using different combinations of Illumina forward and reverse primers with unique barcode tags. The library pools were sequenced a MiSeq v3 platform at the Norwegian Sequencing Centre in Oslo. Shotgun sequencing was performed on an Illumina Novaseq 6000 platform at Novogene, UK facility using the protocol stated in (Gourlé et al., 2018).

Subset B samples were subjected to 16S amplicon sequencing using both Illumina and ONT sequencing platforms. Library preparation for Illumina sequencing of Subset B samples was

performed as described above. ONT sequencing libraries were prepared using the Oxford

Nanopore Sequencing kit (SQK-LSK114-XL). The first amplification was performed using

modified 16S forward and reverse primers (Genetic Analysis, Norway) with the following PCR

program: 95°C for 15 min, 30 cycles of 95°C for 30 sec, 55°C for 30 sec and 72°C for 80 sec.

Amplicons were purified with 1:1 Ampure XP paramagnetic beads (Beckman Coulter) before

indexing PCR. Between 10 and 60 ng of purified full-length 16S amplicons was added to the

PCR mixture, in addition to 0.2 µM of a unique PCR barcode (ONT cat.nr. EXP-PBC096). The

barcoding amplification was done using the following program: 95°C for 15 min, 12 cycles of

95°C for 30 sec, 62°C for 15 min and 65°C for 2 min, followed by 65°C for 10 min. Barcoded

PCR products were concentration-normalized and pooled in equimolar ratios, the library

from which was purified with 1:1 Ampure XP beads before sequencing on a MinION flow

cell. The samples were sequenced at Eurofins Genomics, Germany.

Amplicon data processing

The 16S amplicon sequencing data from subset A were processed with VSEARCH (Rognes et

al., 2016) for clustering into Operational Taxonomic Units (OTUs) using a 95% identity

threshold. ONT reads are longer than Illumina reads but still have error levels making it

difficult to go much above a 95% identity resolution for the 16S sequences. Because the

subset Illumina sequence data will be used to generate reference sequences for error-prone

ONT sequences in subset B, a lower 95% resolution was chosen as opposed to a higher value

or a denoising approach. In addition, a recent study of very high diversity environments like

seafloor sediments demonstrated unstable compositional profiles when using high-

resolution denoising of illumina reads (Nilsen et al., 2024).

The Illumina paired-end reads from subset B were merged and de-replicated by VSEARCH,

but not clustered. Only reads with at least five copies per sample were used, since these are

most likely without sequencing error. Nanopore reads from the same subset B samples were

filtered based on their length, which ranged between $800 - 1\,400$ bp.

SILVA alignments

The OTU represents the sequences of around 420 - 440 bases, which corresponds to the

lengths of the amplicons we obtained by the primers used. First, all OTU representative

sequences were aligned against the 2 152 651 16S sequences in the SILVA 138.1 (Boratyn et

al., 2013). The SILVA sequences have a median length of 1 403 bases and range in length

from 900 to 4 000 bases. OTUs from subset A achieving a best match with ≥ 95% identity

over ≥ 95% of their length were marked as 'known', i.e. resembling something already

known from the SILVA database. The full-length corresponding matching SILVA sequences

were then retrieved as representative for' known 'OTUs (Fig. 1, box Q1).

It should be noted that many of the seed sequences used for reconstruction (see Section:

Reconstructing 16S using METASEED) derive from the same' known 'OTUs, obfuscating the

need for sequence reconstruction as matching SILVA sequences are already known. The

reconstruction was, however, still performed in order to test how closely the reconstructed

sequence could be matched to the corresponding SILVA sequence.

Reconstructing 16S using METASEED

OTUs were used as 'seed sequences' within the METASEED workflow to reconstruct a longer

section of the 16S rRNA gene from shotgun metagenome reads. METASEED collects reads

from the shotgun metagenome data, aligns them to the seed sequences and combines

assignment results with abundance information. For each seed, a reconstructed sequence is

reported if the sequence is ≥ 500 bp in length and the alignment occurs with ≥99% identity

to the corresponding seed. Detailed information on our METASEED approach can be referred

from the recent publication (Philip et al., 2024).

Reconstructed sequences were subsequently aligned to the SILVA database, using the same

thresholds as above. As mentioned before (see Section: SILVA alignments), the 'known'

seeds could now be checked to verify whether the seed and its reconstructed sequence

matched the same SILVA sequence, or at least the same taxon. The most interesting

reconstructed sequences derive from cases where the SILVA database generated no matches

to the seed nor to its reconstructed sequence. These reconstructed sequences were then

added to our collection as new 16S sequences, representing new taxa relative to the SILVA

collection (Fig. 1, boxes Q2 and Q3).

Shotgun data assembly

The shotgun metagenome data from the 94 samples in subset A were filtered and merged

using BBduk and BBmerge (Bushnell, 2014) from BBmap v39.01. BBDuk parameters included

k-mer size 23 (k=23), hamming distance 1 (h_dist=1), 3' end trimming quality 20

(Q_trim=20), and read discarding for average quality below 20 (Q_discard=20). No trimming

was applied to the 5' end (trim_left=0), bases beyond position 150 were trimmed

(trim_right=150), and reads shorter than 30 bases were discarded (min_len=30). The

assembly was done using metaspades v3.15.5 , executed within a Singularity container (Nurk

et al., 2017). Inputs included paired-end, merged, and unpaired reads, with assembly

parameters configured for memory allocation, thread usage, and temporary file handling.

The `--only-assembler` option was used to bypass read error correction.From the

assembled contigs we used Barrnap v0.9 (https://github.com/tseemann/barrnap) to extract

16S sequences from the contigs using length threshold ≥ 800 bases. As 16S sequences

extracted from different samples may be identical, all 16S sequences were clustered using

VSEARCH v2.22.1, as described below, to obtain a resolution of 95% identity (Fig. 1, box Q4).

Finally, we also included 16S centroids that didn't give any match to the above sequences.

These unmatched 16S sequences represent the lowest quality part of the reference

database (Fig. 1, box Q5).

Taxonomic assignments

Taxonomic assignments of all database sequences was obtained using the SINTAX algorithm

(Edgar, 2016) implemented in the VSEARCH software, where we used the GSR database and

the NCBI taxonomy (Molano et al., 2024). This means sequences were assigned to all ranks

to species, with confidence scores at each rank.

Phylogenetic placement of METASEEED and Barrnap derived sequences

Phylogenetic placement of the METASEED- and Barrnap- derived sequences were

determined since these are 'new' 16S variants. First, all database sequences (excluding short

amplicons) were aligned and filtered using `cmsearch` in the `infernal` software v1.1.4

(Eddy, 2002) and secondary structure model for bacterial 16S downloaded from the `Rfam`

database (Kalvari et al., 2021). This level of filtration, utilizing `cmsearch`, was applied to

eliminate potential low-quality sequences from the database and short amplicons were

excluded to minimize potential noise in this analysis. Sequences with an overall score

(bitscore/length of the sequence) below 0.9 were filtered out. A multiple alignment of the

resulting sequences was created using `cmalign` and computed evolutionary distances

using Felsenstein-81 model (Felsenstein, 1981) and the ape R package (Paradis & Schliep,

2019). These distances were then used in a hierarchical clustering with average linkage and

the sequences were split into 64 clusters. The clusters with at least 5 members were subject

to a Fisher exact test for enrichment of sequences originating from either METASEED or

Barrnap, i.e. sequences not previously found in SILVA. We considered a cluster to be

enriched with new variants, if the odds-ratio is greater than 1 and the p-value is less than

0.05. In these clusters there are some 16S variants where the taxonomy is known to phylum

rank, and the dominating phyla in the enriched clusters are mentioned in the results.

Database comparisons

As previously stated, one of the main motivations behind this work is to improve the

database that will be used to analyses amplicon data from Oxford Nanopore Technology.

EMU (Curry et al., 2022) is a standard software for profiling such data and we wanted to

compare our database to the default EMU database. We first used VSEARCH alignments with

default settings to assess the overlap between our generated database and the default EMU

database. Matches that failed to show at least 95% identity were not considered.

Next, we wanted to test how the Nanopore reads from the 90 subset B sediment samples

would be profiled using the EMU software in combination with its default database (EMU-

DB) as well as our custom database (AQUAeD-DB) using default settings. In addition, we also

tested the profiling of Illumina de-replicated reads using standard VSEARCH alignments, i.e.

pairwise alignment between read and database sequences, and assigning to the best, an

identity threshold of 95% was used. The EMU output was processed using an R script to

convert it into the read count matrix format. For VSEARCH, the read count matrix is provided

if specified in the command. The read count matrices from the Illumina VSEARCH and

Nanopore EMU tool assignments for the AQUAeD-DB and EMU-DB databases were used to

compute relative abundance. Prior to the correlation analysis, missing database sequences

from the Illumina read count were added to the Nanopore read count matrix, and vice versa,

with a count of zero for each sample. This was done to resolve disparities in matrix lengths.

Results

Addition   of   SILVA   sequences   to   the   database

The full-length sequences were first collected from the SILVA database, by matching them

against the16S rRNA amplicon representative sequences, generated by conventional short-

read data processing. Using a 95% identity threshold, our 16S rRNA pipeline generated

14,393 OTUs from the subset A data. Each OTU has a representative sequence consisting of

400 - 440 bases in accord with the length of the V3-V4 amplicons. These sequences were

then aligned against the SILVA database using VSEARCH. In total 5 278 of OTUs showed at

least 95% identity matches with some of the SILVA sequences (identity distribution shown in

Fig. 2). This represents 84% - 96% of the total OTU abundance in these 94 samples. The 4

921 distinct SILVA sequences matched to the OTU sequences were collected and filtered

using `infernal cmsearch,` which resulted in 4 838 sequences. The SILVA sequences

were added to the database as full-length representatives of these OTUs and they belong to

the highest quality (Table 1, Q1) sequences in the database.

Fig. 2 Quality of OTU identifications. Frequency distribution of OTU sequences (y-axis) with

varying degrees of sequence similarity (x-axis) to SILVA database sequences. The green

columns indicate the number of OTUs with a SILVA match while the grey columns indicate no

relevant match to SILVA database.

Reconstructing 16S rRNA genes using METASEED

All 14 393 OTUs were passed to the METASEED together with shotgun metagenome data in

order to re-construct near full-length 16S rRNA from the shorter amplicon sequences. The

screening process for the METASEED results is elaborated in Fig. S1. This analysis resulted in

3 525 16S rRNA sequences, of which 3 392 were ≥ 500 bases in length. Of these, 2 947 had

at least 99% sequence identity to their seeds, and were considered re-constructed 16S

sequences.

From the 2 947 reconstructed 16S sequences, 1 176 had no match to SILVA sequences

already collected (≥ 95% identity). Reconstructed 16S rRNA sequences with a minimum

length of 800 bases were regarded as high quality, while sequences 500 - 800 bases in length

were considered low quality. This resulted in 421 high quality and 755 low quality

reconstructed sequences. The total of 1 176 METASEED reconstructed sequences were

filtered using `cmsearch`, which led to the addition of 717 reconstructed sequences to the

database. The Q2 and Q3 category in Table 1 summarize these results. Fig.S2 shows the

increase in captured relative abundance when Q1+Q2+Q3 sequences are included,

compared to using only Q1.

Addition of 16S rRNA from contigs using Barrnap

We also looked for 16S rRNA genes in the assembled contigs from the metagenome data

using the `barrnap` software. From the 94 samples, we found in total 9 839 short or longer

sequences recognized as 16S rRNA. The number of sequences was reduced to 1 149 when

applying a length threshold of at least 800 bp. Finally, since the same 16S rRNA sequence

may have been assembled and extracted from different samples, these sequences were

clustered using a 95% identity threshold, which reduced the number of sequences to 518.

Again, we used the `cmsearch` filtering, resulting in 320 high quality near full-length 16S

rRNA sequences, which were added to database. The total length of assembled contigs was

greater than 32 000 megabasepairs, implying that roughly one full-length high-quality 16S

rRNA was found for every 100 mega base-pairs of genomic sequence (Table 1, Q4).

Apart from the sequences collected from the above-mentioned sources, we also included

OTU sequences that did not match to any sequences added to the database from SILVA,

METASEED, and Barrnap when using a 95% identity threshold. This 8 670 OTUs accounts for

the Q5 category(Table 1). Including all categories (Table 1), the final size of the AQUAeD

database was 14 545 16S sequences.

Taxonomic assignments

All sequences in the AQUAeD database were classified using SINTAX (Fig. 3), with the

exception of sequences belonging to the Q1 category, which were collected from the SILVA

database with taxonomic assignments.

Fig. 3 The bars depict the fraction of classified sequences (Q2 - Q5) in the AQUAeD Database

using SINTAX (bootstrap value 0.8 and 0.5) and the GSR databases. The x-axis shows various

taxonomic ranks, and y-axis shows the fraction of classified sequences.

Enrichment of 'new '16S variants

Sequences in categories Q2, Q3 and Q4 (Table 1) are 'new' 16S rRNA gene variants in the sense that they are not included in the SILVA database. We wanted to see how these distribute when clustered among the sequences in Q1 as well. The multiple sequence alignment of these 5 875 sequences was followed by the evolutionary distance calculation, cluster generation and using the Fisher test for enrichment of the 'new variants'. From 64 clusters we found that 17 clusters were enriched by 'new' variants. The other members of these enriched clusters predominantly consisted of phyla Planctomycetota, Myxococcota, and Verrucomicrobiota (Table 2).

Database comparison

EMU software is the standard tool used for taxonomic profiling of Nanopore 16S reads. It comes with a default database (EMU-DB), consisting of 49 243 sequences. From a comparison between the EMU-DB and AQUAeD-DB, we observed that 18 584 (38%) EMU database sequences had a match to 509 (3.5 %) AQUAeD database sequences when using a 95% identity threshold.

The 88 subset B samples resulted in a total of 973 758 dereplicated Illumina reads, corresponding to 51 397 distinct sequences. The alignment of these reads (95% identity) to the EMU-DB resulted in a match for 3% of the database sequences, comprising 52% of the total Illumina reads. Alignment with the AQUAeD-DB yielded a match to 23% of the database sequences, covering 99% of the total Illumina reads. The Nanopore amplicon reads from the 88 subset B samples having length between 800 and 1 400 bases resulted in 4 972 056 reads. The analysis of Nanopore reads resulted in 3 967 AQUAeD database sequences,

accounting for approximately 27% of the database, while the Nanopore reads matched to

2560 EMU-DB sequences, accounting for approximately 5% of the database.

Fig. 4 The Database comparison. A) Illumina results for taxonomic assignments using

VSEARCH. B, Nanopore results for taxonomic assignments using EMU. B) the percentage

reads assigned exceeds 100% due to the fuzzy assignment by EMU which resulted in a total

'count 'of assigned reads that was larger than the number of reads. Pie charts represent

reads and database sequences are represented by bars. Shaded/striped regions in each pie

chart or bar indicate the fraction of reads assigned or the number of database sequences to

which they were assigned, respectively. Detailed information about alignments can be found

in Table S1.

A comparison of read count matrices revealed that 82% of the EMU-DB sequences in the

Illumina read count matrix were missing in the corresponding Nanopore read count matrix,

while 17% of the sequences in the Nanopore matrix were the missing from Illumina matrix.

For the AQUAeD-DB read count matrices, the overlap was notably better, with 45% of

Illumina sequences and 13% of Nanopore sequences missing from their counterparts,

indicating a lower level of missing database sequences. Correlation analysis demonstrated

high positive correlation between read count matrices for the AQUAeD-DB, whereas the

EMU-DB shows very weak negative correlation (Fig. 5). The relative mean abundance of

Illumina and Nanopore results for AQUAeD-DB demonstrate a significant positive correlation

(Fig. 6).

Fig. 5 Correlation between Illumina and ONT read count matrices. Boxplots indicate

correlation coefficients for read count matrix similarity when using the AQUAeD or EMU

database.

Fig. 6 Relative mean abundance of Illumina versus Nanopore sequences for subset B

370   samples. The scatterplot depicts the log of relative mean abundance of Nanopore (y-axis)

371   versus the log of relative abundance of Illumina(x-axis) of AQUAeD-DB database sequences

372   across samples. Cases where either Illumina or Nanopore values were zero are omitted.

Discussion

The motivation for building the AQUAeD-DB was to enhance the use of Nanopore data in

the study of seafloor sediments. Thus, we wanted to see if the use of AQUAeD-DB has some

benefit compared to a standard database. Since the EMU tool is a popular choice for

Nanopore 16S data processing, we used its default database (EMU-DB) in our comparison. In

addition to using EMU for assigning the reads, we also used VSEARCH, which is typically used

for Illumina data, just for comparison. The VSEARCH alignment directly between the

AQUAeD-DB and EMU-DB reveals that 3.5% of the AQUAeD database could represent 38%

of the EMU database, implying that the AQUAeD-DB, despite being smaller in the number of

sequences, AQUAeD-DB covers an order of magnitude more seafloor diversity compared to

EMU-DB. This difference could also be due to the EMU-DB containing more sequence

variants from the same taxa, specifically from human-associated environments, which may

be less relevant to seafloor ecosystems. This underscores the importance of utilizing

environment-specific databases in monitoring studies.

The AQUAeD-DB comprises of 14 545 16S sequences that were collected from seafloor

sediments (Fig. 1). The Illumina amplicon sequencing of subset A seafloor samples yielded

14 393 centroid sequences when clustered at the coarse level of 95% identity.

Approximately 40% of these sequences matched the SILVA database sequences, even with

relatively liberal identity threshold of 95% (Fig. 2). However, this set of matching SILVA

sequences, the Q1, makes up the backbone of our AQUAeD-DB. To add more longer

sequences to the database, we used our METASEED tool (Philip et al., 2024) to extend the

amplicon sequences, making use of the Illumina metagenome data from the same samples

(subset A). This resulted in the Q2 and Q3 part of the database. The total abundance

covered by the database sequences in each sample did not increase appreciably by adding

these sequences, but the increase was larger for the exceptionally high diversity samples

(high Shannon index of 5-7, Fig. S2). This typically allowed us to obtain 16S sequences for the

low-abundance and ecologically significant organisms from phyla like Planctomycetota (table

2). In the Q4 part of the database, we tried to collect long 16S directly from metagenome

assembled reads. The fact that we only collect one such sequence per 100 megabases of

assembled contigs indicates how difficult it is to obtain such information from the assembly

of short read metagenome data. The limitations of metagenome assembly, specifically the

challenges in retrieving 16S sequences, have been previously addressed (Hiseni et al., 2022;

Mise & Iwasaki, 2022; Shaiber & Eren, 2019).

The AQUAeD-DB can be roughly split into two parts: The 'known 'sequences (Q1) and the

'new variants '(Q2-Q5, Fig 3). The fractions of taxonomic assignments were generally low,

especially for the Q3-Q5 sequences. This emphasizes that new sequence variants in the

AQUAeD database (Q2-Q5) have poor representation in a public database, like the GSR used

here. Considering the poor taxonomic information, we wanted to see if phylogenetic

distances between these new variants, and the Q1 sequences indicate a grouping of these

new sequences, or if they are scattered around in the phylogeny. By clustering based on

phylogenetic distances, we found several clusters where these new variants are over-

represented, which include significant marine microorganisms from the phyla

Planctomycetota, Myxococcota, and Verrucomicrobiota (Dell'Anno et al., 2021; Freitas et al.,

2012; Wu et al., 2024)

A clear difference was evident when we did the assignment with both VSEARCH, which is

normally used for Illumina data, and EMU, which is the standard choice for Nanopore (Fig 4).

The VSEARCH tool aligned the high-precision Illumina reads, with EMU-DB assigning almost

half of the reads and spreading them out among only 1 474 of the nearly 50 000 database

sequences. For AQUAeD-DB, almost all reads are assigned to 3 301 database sequences. The

Nanopore reads showed different results compared to the VSEARCH assignment of Illumina

data, especially in terms of database coverage. One of the major contributing factors can be

the significantly higher noise levels compared to Illumina data; this leads to reads being

distributed across a much larger number of sequences in both databases (Magi et al., 2017;

Santos et al., 2020). Another difference lies in the fact that the EMU tool uses Expectation-

Maximization (EM) algorithm and results in non-integer read counts for each taxa/database

sequence, while result in the assignment more than 100% of the reads as we sume the

readcounts. This likely indicates an artifact of the fuzzy assignment process (Szoboszlay et

al., 2023). From VSEARCH+Illumina and EMU+Nanopore, we see how more organisms are

being given read counts in the AQUAeD-DB compared to the EMU-DB. This is the effect of

AQUAeD-DB containing more relevant taxa for seafloor data, while EMU-DB is a generic

collection and probably more relevant for use in medical research.

The previously stated percentage of missing database sequences in the read count matrices

for Illumina and Nanopore indicate that the EMU-DB analysis results exhibit greater

disparities compared to the AQUAeD-DB results. The discrepancies here can be due to the

different primers, possible primer bias, differences in the 16S rRNA regions used for

Nanopore vs Illumina sequencing (Abellan-Schneyder et al., 2021; Wang et al., 2021; Zhang

et al., 2024). However, clear positive correlation between Illumina and Nanopore relative

abundance were revealed in AQUAeD-DB whereas EMU-DB shows very inconsistent and

poor results (Fig 5). From these results it is evident that AQUAeD-DB results in more reliable

and consistent assignments for both Illumina and ONT analysis. These findings were

supported by the broad range of diversity, especially for the low- abundance taxa that the

AQUAeD-DB database was able to cover (Fig 6). These observations once again underline the

benefits of AQUAeD-DB and emphasize the potential implementation for Nanopore based

environmental studies.

These findings indicated that AQUAeD-DB have illustrated a significant contribution in

environmental analysis using ONT sequencing data. The database enrichment analysis

implies that significant marine microbiota is represented in newly developed AQUAeD

database, comprising both existing sequences and novel sequences from the

aforementioned strategies. This optimizes taxonomical assignment and provides potential

for more explicit environmental context. The study emphasizes the potential, and necessity

of such databases in facilitating the implementation of the cutting-edge technologies, thus

enabling the comprehensive environmental assessments.

Conclusion

The newly developed AQUAeD-DB database is a specialized database for 16S rRNA gene

sequences from highly diverse and complex environmental samples. The primary objective

of AQUAeD-DB was to provide a better reference database for identification of Nanopore

sequences targeting microbial communities in North Atlantic seafloor sediments. Based on

results of the enrichment study and database comparison, this database is substantially

diverse, and efficiently covers an expanded range of microorganisms that are poorly

represented in other databases. This study was designed as a pilot project with the potential

for future scalability of the database to incorporate additional 16S rRNA gene sequences.

This expansion will be rooted in ongoing efforts to improve environmental surveys and

monitoring using microbial profiles.

Acknowledgements

470   We express our gratitude to the fish farmers that provided samples for this research. We

471   would also like to thank the employees of STIM, Aqua Kompetanse, and Akvaplan-NIVA that

472   performed the sampling.

Data availability

The   AQUAED-DB   and   taxonomy   details   are   publicly   available

(https://arken.nmbu.no/~pmelcy/share/AQUAED-DB/). All raw sequences used in this study

have been deposited in the NCBI BioProject database. Illumina 16S rRNA sequences from

subset A and subset B is a part of the project with the accession code PRJNA1128851. The

raw shotgun data of subset A has been uploaded with the accession code PRJNA1128851.

The raw Nanopore sequencing data for subset B has been uploaded with the accession code

PRJNA1153974.   The   metadata   files   for   all   datasets   are   available   in

https://arken.nmbu.no/~pmelcy/share/AQUAED-DB/metadata/

Author contributions

Knut Rudi, Lars Snipen and Melcy Philip conceived the study; Jessica Louise Ray, Morten

Stokkan, Ragnhild Pettersen and Sanna Majaneva design the collection of sediment samples;

Tonje Nilsen performed all lab experiments; Lars Snipen and Melcy Philip performed all

analyses, with conceptual input from Knut Rudi; Knut Rudi, Lars Snipen and Melcy Philip

contributed to interpretation of results; Knut Rudi, Lars Snipen and Melcy Philip drafted the

manuscript. All author(s) read and commented on the manuscript.

Funding

493 This study is a part of the project: "AQUAeD", funded by The Research Council of Norway

494 (Project Number: 320076).

Competing interests

At the time of manuscript writing, Sanna Majaneva, Ragnhild Pettersen were employed by

company Akvaplan-niva AS. Jessica Louise Ray was employed by company Aqua Kompetanse

AS and Morten Stokkan was employed by company STIM AS. The remaining authors declare

that the research was conducted in the absence of any commercial or financial relationships

that could be construed as a potential conflict of interest.

Table 1: The AQUAeD-DB contains sequences of 5 quality categories, from the best (Q1) to the poorest (Q5)

Category descriptions are provided in Materials and Methods. Lengths are shown in basepairs (bp).

Category	Origin	Number of sequences	Minimum length	Median length	Maximum length
Q1	SILVA	4838	900	1461	1668
Q2	High Quality METASEED	311	800	918	1522
Q3	Low Quality METASEED	406	501	688.5	799
Q4	16S from contigs	320	800	924	1577

Q5	Amplicons	8670	288	426	438

506 **Table 2:** Enrichment of 'new' 16S variants. Results of Fisher tests for 17 'new variants 'enriched clusters when
507 k=64.

Odds ratio	Cluster size	New variants fraction	Dominant phylum
7.06	20	0.6	Planctomycetota
7.01	5	0.6	Desulfobacterota,Proteobacteria
6.56	12	0.58	Patescibacteria
4.015	13	0.46	Patescibacteria
3.97	35	0.46	Spirochaetota
3.27	39	0.41	Verrucomicrobiota
3.01	23	0.39	Spirochaetota
2.71	30	0.37	Verrucomicrobiota
2.59	62	0.35	Chloroflexi
2.58	114	0.35	Verrucomicrobiota
2.55	153	0.35	Planctomycetota
2.54	37	0.35	Firmicutes
2.34	30	0.33	Firmicutes
2.34	27	0.33	Fibrobacterota
2.13	61	0.31	Planctomycetota
2.07	72	0.31	Verrucomicrobiota

1.61	241	0.25	Myxococcota

References

Abellan-Schneyder, I., Matchado, M. S., Reitmeier, S., Sommer, A., Sewald, Z., Baumbach, J., List, M., & Neuhaus, K. (2021). Primer, Pipelines, Parameters: Issues in 16S rRNA Gene Sequencing. *mSphere*, *6*(1), 10.1128/msphere.01202-01220. https://doi.org/doi:10.1128/msphere.01202-20

Amarasinghe, S. L., Su, S., Dong, X., Zappia, L., Ritchie, M. E., & Gouil, Q. (2020). Opportunities and challenges in long-read sequencing data analysis. *Genome Biology*, *21*(1), 30. https://doi.org/10.1186/s13059-020-1935-5

Benítez-Páez, A., Portune, K. J., & Sanz, Y. (2016). Species-level resolution of 16S rRNA gene amplicons sequenced through the MinION™ portable nanopore sequencer. *GigaScience*, *5*(1). https://doi.org/10.1186/s13742-016-0111-z

Boratyn, G. M., Camacho, C., Cooper, P. S., Coulouris, G., Fong, A., Ma, N., Madden, T. L., Matten, W. T., McGinnis, S. D., Merezhuk, Y., Raytselis, Y., Sayers, E. W., Tao, T., Ye, J., & Zaretskaya, I. (2013). BLAST: a more efficient report with usability improvements. *Nucleic Acids Research*, *41*(W1), W29-W33. https://doi.org/10.1093/nar/gkt282

Bushnell, B. (2014). *BBMap: a fast, accurate, splice-aware aligner*.

Cabezas, M. P., Fonseca, N. A., & Muñoz-Mérida, A. (2024). MIMt: a curated 16S rRNA reference database with less redundancy and higher accuracy at species-level identification. *Environmental Microbiome*, *19*(1), 88. https://doi.org/10.1186/s40793-024-00634-w

Carrasco De La Cruz, P. M. (2021). The Knowledge Status of Coastal and Marine Ecosystem Services - Challenges, Limitations and Lessons Learned From the Application of the Ecosystem Services Approach in Management [Review]. *Frontiers in Marine Science*, *8*. https://doi.org/10.3389/fmars.2021.684770

Curry, K. D., Wang, Q., Nute, M. G., Tyshaieva, A., Reeves, E., Soriano, S., Wu, Q., Graeber, E., Finzer, P., Mendling, W., Savidge, T., Villapol, S., Dilthey, A., & Treangen, T. J. (2022). Emu: species-level microbial community profiling of full-length 16S rRNA Oxford Nanopore sequencing data. *Nature Methods*, *19*(7), 845-853. https://doi.org/10.1038/s41592-022-01520-4

Dell'Anno, F., Rastelli, E., Tangherlini, M., Corinaldesi, C., Sansone, C., Brunet, C., Balzano, S., Ianora, A., Musco, L., Montereali, M. R., & Dell'Anno, A. (2021). Highly Contaminated Marine Sediments Can Host Rare Bacterial Taxa Potentially Useful for Bioremediation [Original Research]. *Frontiers in Microbiology*, *12*. https://doi.org/10.3389/fmicb.2021.584850

DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., Huber, T., Dalevi, D., Hu, P., & Andersen, G. L. (2006). Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Applied and Environmental Microbiology*, *72*(7), 5069-5072.

Dueholm, M. K. D., Nierychlo, M., Andersen, K. S., Rudkjøbing, V., Knutsson, S., Arriaga, S., Bakke, R., Boon, N., Bux, F., Christensson, M., Chua, A. S. M., Curtis, T. P., Cytryn, E., Erijman, L., Etchebehere, C., Fatta-Kassinos, D., Frigon, D., Garcia-Chaves, M. C., Gu, A. Z.,...Mi, D. A. S. G. C. (2022). MiDAS 4: A global catalogue of full-length 16S rRNA gene sequences and taxonomy for studies of bacterial communities in wastewater treatment plants. *Nature Communications*, *13*(1), 1908. https://doi.org/10.1038/s41467-022-29438-7

Eddy, S. R. (2002). A memory-efficient dynamic programming algorithm for optimal alignment of a
sequence to an RNA secondary structure. *BMC Bioinformatics*, *3*(1), 18.
https://doi.org/10.1186/1471-2105-3-18
Edgar, R. (2016). SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS sequences. In:
bioRxiv.
F. Escapa, I., Huang, Y., Chen, T., Lin, M., Kokaras, A., Dewhirst, F. E., & Lemon, K. P. (2020).
Construction of habitat-specific training sets to achieve species-level assignment in 16S rRNA
gene datasets. *Microbiome*, *8*(1), 65. https://doi.org/10.1186/s40168-020-00841-w
Federhen, S. (2011). The NCBI Taxonomy database. *Nucleic Acids Research*, *40*(D1), D136-D143.
https://doi.org/10.1093/nar/gkr1178
Felsenstein, J. (1981). Evolutionary trees from DNA sequences: a maximum likelihood approach. *J
Mol Evol*, *17*(6), 368-376. https://doi.org/10.1007/bf01734359
Freitas, S., Hatosy, S., Fuhrman, J. A., Huse, S. M., Welch, D. B., Sogin, M. L., & Martiny, A. C. (2012).
Global distribution and diversity of marine Verrucomicrobia. *Isme j*, *6*(8), 1499-1505.
https://doi.org/10.1038/ismej.2012.3
Frenkel-Morgenstern, M., Gorohovski, A., Lacroix, V., Rogers, M., Ibanez, K., Boullosa, C., Andres
Leon, E., Ben-Hur, A., & Valencia, A. (2012). ChiTaRS: a database of human, mouse and fruit
fly chimeric transcripts and RNA-sequencing data. *Nucleic Acids Research*, *41*(D1), D142-
D151. https://doi.org/10.1093/nar/gks1041
Ghate, S. D., Shastry, R. P., Arun, A. B., & Rekha, P. D. (2021). Unraveling the bacterial community
composition across aquatic sediments in the Southwestern coast of India by employing high-
throughput 16S rRNA gene sequencing. *Regional Studies in Marine Science*, *46*, 101890.
https://doi.org/https://doi.org/10.1016/j.rsma.2021.101890
Henderson, G., Yilmaz, P., Kumar, S., Forster, R. J., Kelly, W. J., Leahy, S. C., Guan, L. L., & Janssen, P. H.
(2019). Improved taxonomic assignment of rumen bacterial 16S rRNA sequences using a
revised SILVA taxonomic framework. *PeerJ*, *7*, e6496. https://doi.org/10.7717/peerj.6496
Hiseni, P., Snipen, L., Wilson, R. C., Furu, K., & Rudi, K. (2022). Questioning the Quality of 16S rRNA
Gene Sequences Derived From Human Gut Metagenome-Assembled Genomes [Opinion].
Frontiers in Microbiology, *12*. https://doi.org/10.3389/fmicb.2021.822301
Hoshino, T., Doi, H., Uramoto, G. I., Wörmer, L., Adhikari, R. R., Xiao, N., Morono, Y., D'Hondt, S.,
Hinrichs, K. U., & Inagaki, F. (2020). Global diversity of microbial communities in marine
sediment. *Proc Natl Acad Sci U S A*, *117*(44), 27587-27597.
https://doi.org/10.1073/pnas.1919139117
Kalvari, I., Nawrocki, E. P., Ontiveros-Palacios, N., Argasinska, J., Lamkiewicz, K., Marz, M., Griffiths-
Jones, S., Toffano-Nioche, C., Gautheret, D., & Weinberg, Z. (2021). Rfam 14: expanded
coverage of metagenomic, viral and microRNA families. *Nucleic Acids Research*, *49*(D1),
D192-D200.
Kozlov, A. M., Zhang, J., Yilmaz, P., Glöckner, F. O., & Stamatakis, A. (2016). Phylogeny-aware
identification and correction of taxonomically mislabeled sequences. *Nucleic Acids Research*,
44(11), 5022-5033.
Lemoinne, A., Dirberg, G., Georges, M., & Robinet, T. (2023). Fine-scale congruence in bacterial
community structure from marine sediments sequenced by short-reads on Illumina and
long-reads on Nanopore.
Madsen, E. L. (2011). Microorganisms and their roles in fundamental biogeochemical cycles. *Current
Opinion in Biotechnology*, *22*(3), 456-464.
https://doi.org/https://doi.org/10.1016/j.copbio.2011.01.008
Magi, A., Semeraro, R., Mingrino, A., Giusti, B., & D'Aurizio, R. (2017). Nanopore sequencing data
analysis: state of the art, applications and challenges. *Briefings in Bioinformatics*, *19*(6),
1256-1272. https://doi.org/10.1093/bib/bbx062
Marlow, J., Steele, J., Case, D., Connon, S., Levin, L., & Orphan, V. (2014). Microbial abundance and
diversity patterns associated with sediments and carbonates from the methane seep

environments of Hydrate Ridge, OR. *Frontiers in Marine Science*, *1*.
https://doi.org/10.3389/fmars.2014.00044

McDonald, D., Jiang, Y., Balaban, M., Cantrell, K., Zhu, Q., Gonzalez, A., Morton, J. T., Nicolaou, G.,
Parks, D. H., Karst, S. M., Albertsen, M., Hugenholtz, P., DeSantis, T., Song, S. J., Bartko, A.,
Havulinna, A. S., Jousilahti, P., Cheng, S., Inouye, M.,…Knight, R. (2024). Author Correction:
Greengenes2 unifies microbial data in a single reference tree. *Nature Biotechnology*, *42*(5),
813-813. https://doi.org/10.1038/s41587-023-02026-w

Meola, M., Rifa, E., Shani, N., Delbès, C., Berthoud, H., & Chassard, C. (2019). DAIRYdb: a manually
curated reference database for improved taxonomy annotation of 16S rRNA gene sequences
from dairy products. *BMC Genomics*, *20*(1), 560. https://doi.org/10.1186/s12864-019-5914-8

Mise, K., & Iwasaki, W. (2022). Unexpected absence of ribosomal protein genes from metagenome-
assembled genomes. *ISME Communications*, *2*(1). https://doi.org/10.1038/s43705-022-
00204-6

Molano, L.-A. G., Vega-Abellaneda, S., & Manichanh, C. (2024). GSR-DB: a manually curated and
optimized taxonomical database for 16S rRNA amplicon analysis. *mSystems*, *9*(2), e00950-
00923. https://doi.org/doi:10.1128/msystems.00950-23

Nierychlo, M., Andersen, K. S., Xu, Y., Green, N., Jiang, C., Albertsen, M., Dueholm, M. S., & Nielsen, P.
H. (2020). MiDAS 3: An ecosystem-specific reference database, taxonomy and knowledge
platform for activated sludge and anaerobic digesters reveals species-level microbiome
composition of activated sludge. *Water Research*, *182*, 115955.
https://doi.org/https://doi.org/10.1016/j.watres.2020.115955

Nilsen, T., Snipen, L.-G., Angell, I. L., Keeley, N. B., Majaneva, S., Pettersen, R., & Rudi, K. (2024).
Swarm and UNOISE outperform DADA2 and Deblur for denoising high-diversity marine
seafloor samples. *ISME Communications*. https://doi.org/10.1093/ismeco/ycae071

Nurk, S., Meleshko, D., Korobeynikov, A., & Pevzner, P. A. (2017). metaSPAdes: a new versatile
metagenomic assembler. *Genome research*, *27*(5), 824-834.

Overgaard, C. K., Tao, K., Zhang, S., Christensen, B. T., Blahovska, Z., Radutoiu, S., Kelly, S., &
Dueholm, M. K. D. (2022). Application of ecosystem-specific reference databases for
increased taxonomic resolution in soil microbial profiling. *Front Microbiol*, *13*, 942396.
https://doi.org/10.3389/fmicb.2022.942396

Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. *Bioinformatics*, *35*(3), 526-528.

Patil, P. K., Baskaran, V., Vinay, T. N., Avunje, S., Leo-Antony, M., Shekhar, M. S., Alavandi, S. V., &
Vijayan, K. K. (2021). Abundance, community structure and diversity of nitrifying bacterial
enrichments from low and high saline brackishwater environments. *Letters in Applied*
Microbiology, *73*(1), 96-106. https://doi.org/10.1111/lam.13480

Pettersen, R., Ormaasen, I., Angell, I. L., Keeley, N. B., Lindseth, A., Snipen, L., & Rudi, K. (2022).
Bimodal distribution of seafloor microbiota diversity and function are associated with marine
aquaculture. *Marine Genomics*, *66*, 100991.
https://doi.org/https://doi.org/10.1016/j.margen.2022.100991

Philip, M., Rudi, K., Ormaasen, I., Angell, I. L., Pettersen, R., Keeley, N. B., & Snipen, L.-G. (2024).
METASEED: a novel approach to full-length 16S rRNA gene reconstruction from short read
data. *BMC Bioinformatics*, *25*(1), 237. https://doi.org/10.1186/s12859-024-05837-z

Ritari, J., Salojärvi, J., Lahti, L., & de Vos, W. M. (2015). Improved taxonomic assignment of human
intestinal 16S rRNA sequences by a dedicated reference database. *BMC Genomics*, *16*(1),
1056. https://doi.org/10.1186/s12864-015-2265-y

Rodríguez-Pérez, H., Ciuffreda, L., & Flores, C. (2020). NanoCLUST: a species-level analysis of 16S
rRNA nanopore sequencing data. *Bioinformatics*, *37*(11), 1600-1601.
https://doi.org/10.1093/bioinformatics/btaa900

Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: a versatile open source tool
for metagenomics. *PeerJ*, *4*, e2584. https://doi.org/10.7717/peerj.2584

Santos, A., van Aerle, R., Barrientos, L., & Martinez-Urtaza, J. (2020). Computational methods for 16S
metabarcoding studies using Nanopore sequencing data. *Computational and Structural*
Biotechnology   *Journal*,   *18*,   296-305.
https://doi.org/https://doi.org/10.1016/j.csbj.2020.01.005
Shaiber, A., & Eren, A. M. (2019). Composite Metagenome-Assembled Genomes Reduce the Quality
of   Public   Genome   Repositories.   *mBio*,   *10*(3),   10.1128/mbio.00725-00719.
https://doi.org/doi:10.1128/mbio.00725-19
Szoboszlay, M., Schramm, L., Pinzauti, D., Scerri, J., Sandionigi, A., & Biazzo, M. (2023). Nanopore Is
Preferable over Illumina for 16S Amplicon Sequencing of the Gut Microbiota When Species-
Level Taxonomic Classification, Accurate Estimation of Richness, or Focus on Rare Taxa Is
Required. *Microorganisms*, *11*(3), 804. https://www.mdpi.com/2076-2607/11/3/804
Tatsuhiko Hoshinoa, H. D., Go-Ichiro Uramotoa, Lars Wörmerc, Rishi R. Adhikari, Nan Xiaoa,e,, Yuki
Moronoa, Steven D'Hondtf, Kai-Uwe Hinrichsc , Fumio Inagaki. (2020). Global diversity of
microbial communities in

marine sediment.
Thurber, A. R., Sweetman, A. K., Narayanaswamy, B. E., Jones, D. O. B., Ingels, J., & Hansman, R. L.
(2014). Ecosystem function and services provided by the deep sea. *Biogeosciences*, *11*(14),
3941-3963. https://doi.org/10.5194/bg-11-3941-2014
Wang, Y., Zhao, Y., Bollas, A., Wang, Y., & Au, K. F. (2021). Nanopore sequencing technology,
bioinformatics   and   applications.   *Nature   Biotechnology*,   *39*(11),   1348-1365.
https://doi.org/10.1038/s41587-021-01108-x
Wu, J., Hong, Y., Chang, X., Jiao, L., Li, Y., Liu, X., Xie, H., & Gu, J.-D. (2019). Unexpectedly high
diversity of anammox bacteria detected in deep-sea surface sediments of the South China
Sea. *FEMS Microbiology Ecology*, *95*(3). https://doi.org/10.1093/femsec/fiz013
Wu, X.-T., He, Y.-Q., Qiu, M., Li, G.-X., Ou, X.-B., Wu, Y.-B., & Bao, P. (2024). Adaptive Changes in
Bacterial and Archaeal Community Structures and Functions Occur under Primordial
Conditions.   *ACS   Earth   and   Space   Chemistry*,   *8*(7),   1470-1479.
https://doi.org/10.1021/acsearthspacechem.4c00089
Yang, C.-c., & Iwasaki, W. (2014). MetaMetaDB: A Database and Analytic System for Investigating
Microbial   Habitability.   *PLOS   ONE*,   *9*(1),   e87126.
https://doi.org/10.1371/journal.pone.0087126
Yang, N., Tian, C., Lv, Y., Hou, J., Yang, Z., Xiao, X., & Zhang, Y. (2022). Novel primers for 16S rRNA
gene-based archaeal and bacterial community analysis in oceanic trench sediments. *Applied*
Microbiology and Biotechnology, *106*(7), 2795-2809. https://doi.org/10.1007/s00253-022-
11893-3
Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J., Ludwig,
W., & Glöckner, F. O. (2013). The SILVA and "All-species Living Tree Project (LTP)" taxonomic
frameworks.   *Nucleic   Acids   Research*,   *42*(D1),   D643-D648.
https://doi.org/10.1093/nar/gkt1209
Yu, Y., Lee, C., Kim, J., & Hwang, S. (2005). Group-specific primer and probe sets to detect
methanogenic communities using quantitative real-time polymerase chain reaction.
Biotechnology   and   Bioengineering,   *89*(6),   670-679.
https://doi.org/https://doi.org/10.1002/bit.20347
Zhang, T., Li, H., Jiang, M., Hou, H., Gao, Y., Li, Y., Wang, F., Wang, J., Peng, K., & Liu, Y.-X. (2024).
Nanopore sequencing: flourishing in its teenage years. *Journal of Genetics and Genomics*.
https://doi.org/https://doi.org/10.1016/j.jgg.2024.09.007
Zorz, J., Li, C., Chakraborty, A., Gittins, D. A., Surcon, T., Morrison, N., Bennett, R., MacDonald, A., &
Hubert, C. R. J. (2023). SituSeq: an offline protocol for rapid and remote Nanopore 16S rRNA
amplicon   sequence   analysis.   *ISME   Communications*,   *3*(1),   33.
https://doi.org/10.1038/s43705-023-00239-3