Disruption of the Novel Nested Gene Aff3ir Mediates Disturbed Flow-Induced Atherosclerosis in Mice

Shuo He[1#], Lei Huang[2,3#], Zhuozheng Chen[1], Ze Yuan[1], Yue Zhao[2,4], Lingfang Zeng[4], Yi Zhu[1*], Jinlong He[1*]*

[1]Tianjin Key Laboratory of Metabolic Diseases, Province and Ministry Co-sponsored Collaborative Innovation Center for Medical Epigenetics, Department of Physiology and Pathophysiology, Tianjin Medical University, Tianjin, 300070, China;

[2]Department of Heart Center, The Third Central Hospital of Tianjin, 83 Jintang Road, Hedong District, Tianjin 300170, China;

[3]Nankai University Affiliated Third Center Hospital, Tianjin 300170, China;

[4]School of Cardiovascular and Metabolic Medicine and Sciences, King's College London British Heart Foundation Centre of Excellence, Faculty of Life Sciences and Medicine, King's College London, London SE5 9NU, United Kingdom.

Running title: AFF3ir-ORF2 exerts anti-atherosclerosis roles

[#]These authors contributed equally to this study.

*Correspondence:

Jinlong He, Ph.D.

Department of Physiology and Pathophysiology, Tianjin Medical University; Qixiangtai Rd 22[nd], Tianjin, 300070, China.

Tel: +86-22-83336591; E-mail: hejinlong@tmu.edu.cn

or

Yi Zhu, M.D., Ph.D.

Department of Physiology and Pathophysiology, Tianjin Medical University; Qixiangtai Rd 22[nd], Tianjin, 300070, China.

Tel: +86-22-83336591; E-mail: zhuyi@tmu.edu.cn

or

Lingfang Zeng, Ph.D.

School of Cardiovascular and Metabolic Medicine and Sciences, Faculty of Life Science and

Medicine, King's College London, London SE5 9NU, United Kingdom. Tel: +44 7848

85295; E-mail: lingfang.zeng@kcl.ac.uk

Keywords: Shear stress, Atherosclerosis, Nested gene, IRF5

Abstract

Disturbed shear stress-induced endothelial atherogenic responses are pivotal in the initiation

and progression of atherosclerosis, contributing to the uneven distribution of atherosclerotic

lesions. This study investigates the role of Aff3ir-ORF2, a novel nested gene variant, in

disturbed flow-induced endothelial cell activation and atherosclerosis. We demonstrate that

disturbed shear stress significantly reduces Aff3ir-ORF2 expression in athero-prone regions.

Using three distinct mouse models with manipulated AFF3ir-ORF2 expression, we

demonstrate that AFF3ir-ORF2 exerts potent anti-inflammatory and anti-atherosclerotic

effects in ApoE$^{-/-}$ mice. RNA sequencing revealed that interferon regulatory factor 5 (IRF5), a

key regulator of inflammatory processes, mediates inflammatory responses associated with

AFF3ir-ORF2 deficiency. AFF3ir-ORF2 interacts with IRF5, promoting its retention in the

cytoplasm, thereby inhibiting the IRF5-dependent inflammatory pathways. Notably, IRF5

knockdown in AFF3ir-ORF2 deficient mice almost completely rescues the aggravated

atherosclerotic phenotype. Moreover, endothelial-specific AFF3ir-ORF2 supplementation

using the CRISPR/Cas9 system significantly ameliorated endothelial activation and

atherosclerosis. These findings elucidate a novel role for AFF3ir-ORF2 in mitigating

endothelial inflammation and atherosclerosis by acting as an inhibitor of IRF5, highlighting

its potential as a valuable therapeutic approach for treating atherosclerosis.

Introduction

Atherosclerosis, characterized by the formation of fibrofatty lesions in the arterial wall, is a leading cause of morbidity and mortality worldwide, contributing to most myocardial infarctions and many strokes (Herrington, Lacey, Sherliker, Armitage, & Lewington, 2016; Libby et al., 2019). The activation of vascular endothelial cells (ECs), induced by various chemical and mechanical stimuli, such as lipopolysaccharide and shear stress, is an initial step in the development of atherosclerosis (Davignon & Ganz, 2004). Consequently, atherosclerotic lesions preferentially develop at the branches and curvatures of the arterial tree, where blood flow is disturbed (Davis, Earley, Li, & Chien, 2023). For decades, researchers have been interested in exploring the mechanisms underlying mechanotransduction during endothelial activation caused by disturbed flow (Davis et al., 2023). Several mechanosensitive proteins, such as YAP/TAZ (B. Li et al., 2019), Annexin A2 (Zhang et al., 2020), BMP4 (Sorescu et al., 2004), and NAD(P)H oxidase (Hwang et al., 2003; Jo, Song, & Mowbray, 2006), have been identified as key regulators of disturbed shear stress in ECs and have been implicated in the progression of atherosclerosis. Emerging evidence has revealed that pharmacological or genetic inhibition of endothelial YAP activation ameliorates the progression of atherosclerotic plaques in mice (B. Li et al., 2019; Wang et al., 2016; Yang et al., 2021), indicating that targeting disturbed flow-induced endothelial activation could be a promising therapeutic strategy for atherosclerosis. However, the precise mechanisms by which the disturbed flow exerts detrimental effects remain unclear.

The interferon regulatory factor (IRF) family of transcription factors, comprising nine members (IRF1-IRF9) in mammals, is primarily characterized by its role in mediating antiviral responses and type I interferon production (Sato, Taniguchi, & Tanaka, 2001). Although these members share a conserved DNA-binding domain in their N-terminal region that recognizes similar DNA sequences, IRF5 plays a central role in inflammation (Almuttaqi & Udalova, 2019; Takaoka et al., 2005). IRF5 mediates the production of proinflammatory cytokines, including IL-12b and IL-23a, and promotes the expression of inflammatory genes (Cai, Yao, & Li, 2017; Saliba et al., 2014; Weiss, Blazek, Byrne, Perocheau, & Udalova, 2013). It promotes inflammatory responses in various immune cells, including macrophages (Seneviratne et al., 2017), neutrophils (Weiss et al., 2015), and B cells (Savitsky, Yanai, Tamura, Taniguchi, & Honda, 2010). Global or myeloid-specific knockouts of IRF5 have been shown to exert anti-atherosclerotic effects (Leipner et al., 2021; Seneviratne et al., 2017). Despite the established importance of IRF5 in immune cells, its restrictively regulatory mechanism and role in shear stress-induced endothelial activation remain unknown.

We recently reported that a novel protein-coding nested gene, *Aff3ir*, contributes to
endothelial maintenance by promoting the differentiation of vascular stem/progenitor cells
(SPCs) into ECs (Zhao Y et al., 2024). AFF3ir-ORF2, encoded by the *Aff3ir* transcript variant
2, is predominantly expressed in the EC layer of the mouse aorta (Zhao Y et al., 2024).
Notably, our recent study indicated that the overexpression of AFF3ir-ORF2 could enhance
laminar flow-induced mRNA levels of essential EC markers in SPCs (Zhao Y et al., 2024),
suggesting that AFF3ir-ORF2 may be a novel mechanotransduction protein in ECs. However,
the regulation of Aff3ir-ORF2 under disturbed flow and its role in atherosclerosis remain
unclear.

In this study, we aimed to elucidate the mechanism by which disturbed blood flow induces
endothelial activation and atherosclerosis. Our study showed that disrupted Aff3ir-ORF2
expression in athero-prone regions led to inflammatory responses and development of
atherosclerosis. Aff3ir-ORF2, the expression of which is reduced by disturbed shear stress,
exerts critical anti-inflammatory effects by binding to IRF5 and mitigating disturbed shear
stress-induced IRF5 activation. Additionally, we demonstrated that endothelial-specific
supplementation with Aff3ir-ORF2 significantly ameliorated disturbed flow-induced
endothelial activation and the development of atherosclerotic plaques, highlighting its
promising therapeutic potential for the treatment of atherosclerosis.

Results

Disturbed shear stress reduces the expression of AFF3ir-ORF2.

Our recent study showed the active participation of *Aff3ir* in EC differentiation from vascular SPCs induced by laminar shear stress (Zhao Y et al., 2024), suggesting the potential involvement of this novel protein-encoding nested gene in mediating hemodynamic stimulation. To further elucidate the functional role of *Aff3ir* and its encoded proteins in disturbed shear stress-induced EC activation, we examined the expression of *AFF3* and *AFF3ir* in segments of the mouse aorta. We found that the mRNA level of *AFF3ir*, but not that of its parent gene *AFF3*, was significantly lower in the aortic arch, an area exposed to disturbed shear stress, than in the thoracic aorta, which was exposed to steady unidirectional shear stress (B. Li et al., 2019) (Figure 1A). *Aff3ir* transcript variants can generate two proteins (Zhao Y et al., 2024), therefore, we measured the protein levels of AFF3ir-ORF1 and AFF3ir-ORF2. While AFF3ir-ORF1 and AFF3 showed comparable expression levels in the aortic arch and thoracic aorta of mice, the expression of AFF3ir-ORF2 showed an 87% reduction in the aortic arch compared to the thoracic aorta (Figure 1B, C), suggesting that AFF3ir-ORF2 may be a novel mechanosensitive protein that responds to disturbed shear stress. Enface immunofluorescence staining confirmed a marked reduction in AFF3ir-ORF2 expression in the aortic arch compared to that in the thoracic aorta (Figure 1D, E). Moreover, we observed the expression of AFF3ir-ORF2 in longitudinal sections of the mouse aorta (B. Li et al., 2019). We found that the expression of AFF3ir-ORF2, but not AFF3, was notably downregulated in athero-prone regions (the inner curvature of the aortic arch and bifurcation of the carotid artery) compared to that in the protective region in the outer curvature of the aortic arch (Figure 1F).

Next, we explored the impact of disturbed shear stress on AFF3ir-ORF2 expression *in vitro*. Mouse embryonic fibroblasts (MEFs) exhibit responses consistent with those of ECs (Chen et al., 2010; Wen et al., 2013), therefore, we investigated AFF3ir-ORF2 expression in MEFs from WT mice exposed to static or disturbed flow (0.5 ± 4 dyn/cm^2, 1 Hz). Consistent with our *in vivo* findings, while disturbed shear stress increased the expression of VCAM-1, a critical inflammatory marker of ECs (Nakashima, Raines, Plump, Breslow, & Ross, 1998), it significantly reduced both the protein and mRNA levels of AFF3ir-ORF2 (Figure 1G–I). The expression of AFF3 in response to the disturbed flow was minimally affected at both the mRNA and protein levels (Figure 1G–I). These results collectively demonstrate that disturbed shear stress induces a reduction in AFF3ir-ORF2 expression both *in vivo* and *in vitro*.

AFF3ir-ORF2 ameliorates disturbed shear stress-induced inflammation and atherosclerosis

Disturbed shear stress-induced atherogenic responses are initial events in atherosclerotic plaque formation (Davis et al., 2023). To elucidate the regulatory role of AFF3ir-ORF2 in disturbed shear stress-induced inflammation, we overexpressed AFF3ir-ORF2 in MEFs. AFF3ir-ORF2 overexpression attenuated ICAM-1 expression induced by disturbed shear stress at both the protein and mRNA levels (Figure 2A–C). Additionally, AFF3ir-ORF2 overexpression blunted the disturbed shear stress-induced expression of several other inflammatory genes, including VCAM-1, IL-6, and IL-1β (Figure 2C).

Given the anti-inflammatory effects of AFF3ir-ORF2, we speculated that it may ameliorate disturbed shear stress-induced inflammation and atherosclerosis *in vivo*. ApoE knockout (ApoE$^{-/-}$) mice were subjected to partial ligation surgery to induce disturbed flow in the left carotid arteries (LCAs). The endothelium of the LCAs was intravascularly infected with adenovirus (Ad-Scramble or Ad-AFF3ir-ORF2) prior to surgery (Nam et al., 2009; Zhang et al., 2020). Enface immunofluorescence staining confirmed the successful AFF3ir-ORF2 overexpression in the left carotid artery (Figure S1A). Mice infected with Ad-AFF3ir-ORF2 exhibited a significant decrease in lesion area in the LCAs compared to those infected with Ad-Scramble (23±17% vs 63±14%) (Figure 2D–F). Overexpression of AFF3ir-ORF2 also attenuated disturbed flow-induced inflammatory responses, as evidenced by decreased VCAM-1 expression in the endothelium of LCAs (Figure 2G, H). These findings suggested that AFF3ir-ORF2 ameliorates shear stress-induced inflammation and atherosclerosis.

AFF3ir-ORF2 deficiency aggravates inflammation and atherosclerotic lesions in ApoE$^{-/-}$ mice

To explore the effects of AFF3ir-ORF2 on inflammation and atherosclerosis, we generated AFF3ir-ORF2 global knockout (AFF3ir-ORF2$^{-/-}$) mice. No obvious phenotypic abnormalities were observed in AFF3ir-ORF2$^{-/-}$ mice up to 20 weeks of age and monitoring was discontinued thereafter. Additionally, AFF3ir-ORF2 deficiency did not alter systolic blood pressure, diastolic blood pressure, or mean arterial pressure (Figure S2A), suggesting that AFF3ir-ORF2 is dispensable for physiological blood pressure maintenance. We then isolated MEFs from WT and AFF3ir-ORF2$^{-/-}$ mice. RT-PCR analysis confirmed the deficiency of AFF3ir-ORF2 in AFF3ir-ORF2$^{-/-}$ MEFs (Figure S2B). Under disturbed flow stimulation, AFF3ir-ORF2 deficient MEFs displayed higher expression of inflammatory genes, including *ICAM-1, VCAM-1,* and *IL-1b*, compared to those in WT MEFs (Figure S2C).

Next, we crossed AFF3ir-ORF2$^{-/-}$ mice with ApoE$^{-/-}$ mice to generate double-knockout (ApoE$^{-/-}$AFF3ir-ORF2$^{-/-}$) mice. Eight-week-old ApoE$^{-/-}$ and ApoE$^{-/-}$AFF3ir-ORF2$^{-/-}$ mice were fed a high-fat diet for 12 weeks to induce atherosclerosis (Figure 3A). *En face* Oil-Red O staining indicated that AFF3ir-ORF2 deficiency accelerated the development of atherosclerosis in the entire aorta (Figure 3B, C). Furthermore, AFF3ir-ORF2 deletion increased the lesion area and lipid deposition in the aortic roots of ApoE$^{-/-}$ mice without altering the collagen fiber content (Figure 3D, E). Similar results were observed in distributing arteries (LCAs) (Figure 3F, G). Given that the expression of adhesion proteins, such as VCAM-1 in ECs is crucial for monocyte infiltration into plaques (Kobiyama & Ley, 2018), we assessed VCAM-1 expression in the aortic roots of these mice. We found that AFF3ir-ORF2 deletion increased VCAM-1 expression in the aortic roots of ApoE$^{-/-}$ mice (Figure 3H, I), indicating that the atherogenic effects of AFF3ir-ORF2 deletion may result from endothelial inflammation. Additionally, there were no significant differences between the two groups in body weight or triglyceride, total cholesterol, LDL cholesterol, and HDL cholesterol levels (Figure S3A, B), indicating that the atherogenic effect of AFF3ir-ORF2 silencing is unlikely to be related to lipid metabolism. Taken together, these results indicate that AFF3ir-ORF2 deficiency aggravates inflammation and atherosclerotic lesions in mice.

AFF3ir-ORF2 mitigates disturbed shear stress-induced inflammation by interacting with IRF5 and retaining it within the cytosol.

To explore the mechanism by which AFF3ir-ORF2 mitigates atherogenesis, we performed RNA sequencing (RNA-seq) on MEFs from WT and AFF3ir-ORF2$^{-/-}$ mice. We identified 1,167 upregulated and 310 downregulated genes in the AFF3ir-ORF2$^{-/-}$ group, with a criterion of 1.5-fold change and P<0.05 (Figure 4A, B). Differentially expressed genes were subjected to bioinformatics enrichment analysis using Gene Ontology (GO) databases. GO analysis showed that these genes were mainly enriched in processes, including leukocyte cell-cell adhesion, regulation of cell−cell adhesion, and leukocyte activation involved in immune response (Figure 4C), which is highly consistent with the phenotypes observed in AFF3ir-ORF2$^{-/-}$ mice. To further identify the upstream transcriptional regulators of these genes, we predicted upstream transcription factors using the ChEA3 database (Keenan et al., 2019). The top 20 transcription factors obtained from the ChEA3 database were mapped to the atherosclerotic disease-related gene list in the Disgenet database (Pinero et al., 2017). Interferon regulatory factor 5 (IRF5) and IRF8 were identified as key upstream regulators (Figure 4D).

IRF5 and IRF8, which are members of the same family of transcription factors originally implicated in interferon production, have been identified as critical regulators of the inflammatory response and contribute to the pathogenesis of various inflammatory diseases (Almuttaqi & Udalova, 2019; Salem, Salem, & Gros, 2020). However, their potential roles in disturbed shear stress-induced inflammation remain unclear. We speculated that AFF3ir-ORF2 interacts with IRF5 and/or IRF8. Coimmunoprecipitation assays indicated that endogenous AFF3ir-ORF2 could bind to both IRF5 and IRF8 (Figure 4E). To determine which transcription factor that mediates the inflammatory effects of AFF3ir-ORF2 deficiency, we silenced IRF5 and IRF8 in WT and AFF3ir-ORF2$^{-/-}$ MEFs exposed to disturbed flow. Notably, silencing IRF5, but not IRF8, blunted the upregulation of inflammatory genes, including *ICAM-1, VCAM-1, IL-6,* and *IL-1β* (Figure 4F), suggesting that IRF5 was the predominant factor mediating the anti-inflammatory effects of AFF3ir-ORF2 in the context of disturbed shear stress. In addition, neither IRF5 nor IRF8 expression was regulated by AFF3ir-ORF2 (Figure 4F). Consistently, we found that IRF5 silencing significantly inhibited the upregulation of ICAM-1 protein levels induced by AFF3ir-ORF2 deficiency under disturbed shear stress (Figure 4G, H). Given that the transcriptional activity of IRF5 depends on its nuclear translocation (Lv et al., 2024), we next explored whether AFF3ir-ORF2 affects the subcellular localization of IRF5. Subcellular fractionation assays indicated that IRF5 was predominantly localized in the cytoplasm under static conditions, but exhibited obvious nuclear localization when exposed to disturbed shear stress (Figure 4I, J). While the total expression of IRF5 was barely affected by AFF3ir-ORF2 deficiency or overexpression, nuclear localization of IRF5 increased with AFF3ir-ORF2 deficiency (Figure 4I, J). To further ascertain the role of AFF3ir-ORF2 in regulating the transcriptional activity of IRF5, we performed a luciferase reporter assay (Qiao, Lv, Qiao, Wang, & Miao, 2022). AFF3ir-ORF2 overexpression significantly decreased the transcriptional activity of IRF5 (Figure 4K). In summary, these results suggested that AFF3ir-ORF2 acts as an endogenous inhibitor of IRF5 and exerts anti-inflammatory effects by retaining IRF5 in the cytosol.

IRF5 knockdown prevents the aggravation of atherosclerosis induced by ORF2 deficiency.

Next, we investigated the role of IRF5 in disturbed flow-induced atherosclerosis *in vivo* and whether it mediates the atherogenic phenotype associated with AFF3ir-ORF2 deficiency. ApoE$^{-/-}$ and ApoE$^{-/-}$AFF3ir-ORF2$^{-/-}$ mice were subjected to partial ligation surgery in the LCAs and intravascularly infected with lentiviruses expressing either IRF5-specific shRNA

(lenti-shIRF5) or Scramble shRNA (lenti-shScramble). En-face immunofluorescence staining confirmed successful IRF5 deletion in the left carotid artery (Figure S4A). After a 4-week high-fat diet challenge, IRF5 deletion resulted in an approximately 60% reduction in plaque area in the LCAs of ApoE$^{-/-}$ mice (Figure 5A, B). In addition, IRF5 deletion attenuated endothelial activation, as evidenced by reduced VCAM-1 expression in the endothelium of LCAs (Figure 5C, D). Notably, although ApoE$^{-/-}$AFF3ir-ORF2$^{-/-}$ mice exhibited an increased plaque area in the LCAs compared to ApoE$^{-/-}$ mice, IRF5 deletion almost completely abolished these differences, reducing both the plaque area and VCAM-1 expression in the endothelium of the LCAs (Figure 5A–D). These findings provide in vivo evidence that AFF3ir-ORF2 deficiency-induced atherosclerosis is mediated by endothelial IRF5.

Endothelial-specific AFF3ir-ORF2 supplementation ameliorates EC activation and atherosclerosis in mice.

Given the significant anti-inflammatory effects of AFF3ir-ORF2 on endothelial activation and atherosclerosis, we explored the potential use of gene therapy targeting AFF3ir-ORF2 to treat atherosclerosis. Endothelial-specific AFF3ir-ORF2 overexpression was achieved using an EC-enhanced AAV-mediated CRISPR/Cas9 genome-editing system controlled by an EC-specific ICAM2 promoter as we previously reported (Z. Li et al., 2024; Swiech et al., 2015). ApoE$^{-/-}$ mice infected with AAV-ICAM2-Control or AAV-ICAM2- AFF3ir-ORF2 were fed a high-fat diet for 12 weeks (Figure 6A). En-face immunofluorescence staining confirmed successful AFF3ir-ORF2 overexpression in ECs (Figure S5A, B). Endothelial-specific AFF3ir-ORF2 overexpression had a minimal effect on triglycerides, total cholesterol, LDL cholesterol, and HDL cholesterol levels in the plasma of mice (Figure S5C). However, compared to the negative control, endothelial-specific AFF3ir-ORF2 overexpression significantly reduced the Oil-red O-positive lesion area in the whole aortas of ApoE$^{-/-}$ mice (19±5% vs 54±8%) (Figure 6B, C). Moreover, endothelial-specific AFF3ir-ORF2 overexpression reduced the lesion area and lipid deposition in the aortic roots of ApoE$^{-/-}$ mice without altering the collagen fiber content (Figure 6D, E). In addition, AFF3ir-ORF2 overexpression effectively suppressed VCAM-1 expression in the endothelium of the aortic roots of ApoE$^{-/-}$ mice (Figure 6F, G). Collectively, these results suggest that supplementation with AFF3ir-ORF2 was effective in preventing atherosclerosis development.

Discussion

Endothelial activation is a critical initial event in the development of atherosclerosis, and emerging evidence suggests that targeting disturbed shear stress-induced endothelial activation is a promising therapeutic strategy. In the present study, we elucidated the role of the novel nested gene-encoded protein, AFF3ir-ORF2, in sensing disturbed shear stress. Moreover, we demonstrated that AFF3ir-ORF2 acts as an endogenous inhibitor of IRF5, a key regulator of the inflammatory response, thereby exerting potent anti-inflammatory and anti-atherogenic effects (Figure 7).

Using three mouse models (global AFF3ir-ORF2 knockout, locally AFF3ir-ORF2 endothelial expression, and endothelial-specific AFF3ir-ORF2 overexpression), we demonstrated that AFF3ir-ORF2 exerted potent anti-inflammatory and anti-atherosclerosis effects in ApoE$^{-/-}$ mice. Notably, while AFF3ir-ORF2 knockout increased VCAM-1 expression in the endothelium and enlarged the plaque area in the aortic roots, it had a minimal effect on collagen deposition within the plaques. This discrepancy may be attributed to the differential expression patterns of IRF5 across various cell types (Roberts, Collado, & Barnes, 2024). Phenotypically modulated vascular smooth muscle cells (VSMCs) within the fibrous cap produce extracellular matrix molecules critical for plaque composition and stabilization (Bennett, Sinha, & Owens, 2016). A previous study has shown minimal colocalization between IRF5 and the VSMC marker, α-smooth muscle actin, in aortic root lesions of ApoE$^{-/-}$ mice (Seneviratne et al., 2017), indicating a relatively low IRF5 expression level in VSMCs. Consequently, AFF3ir-ORF2 likely exerts its anti-inflammatory effects primarily through endothelial IRF5. Consistently, endothelial-specific AFF3ir-ORF2 overexpression reduced the aortic plaque area, but had minimal effects on collagen deposition. Our findings establish the potent anti-atherosclerotic role of AFF3ir-ORF2 in early and advanced atherosclerosis mouse models. However, given the multiple critical roles of ECs throughout the initiation and progression of atherosclerosis, further investigations are needed to explore the potential role of AFF3ir-ORF2 in other atherosclerotic processes, including endothelial-to-mesenchymal transition, plaque rupture, and atherothrombotic occlusion. Additionally, we found that disturbed shear stress transcriptional downregulated the expression of AFF3ir. However, the protein levels of AFF3ir-ORF2, but not those of AFF3ir-ORF1, were reduced by disturbed shear stress. Since both AFF3ir-ORF1 and AFF3ir-ORF2 are derived from AFF3ir, different translation mechanisms may be involved in the production of AFF3ir-ORF1/2.

In addition to ECs, various other cell types, particularly immune cells, play crucial roles in the

progression of atherosclerotic plaques (Wolf & Ley, 2019). Macrophage polarization and inflammatory responses accelerate plaque development, leading to an increase in necrotic core and vulnerable plaques (Krausgruber et al., 2011). Elevated IRF5 expression and nuclear localization have been observed in macrophages within plaques of ApoE$^{-/-}$ mice (Seneviratne et al., 2017). IRF5 has been demonstrated to drive macrophages towards a pro-inflammatory state, thereby affecting plaque stability (Seneviratne et al., 2017). Global or myeloid cell-specific deletion of IRF5 stabilizes atherosclerotic plaques by suppressing the inflammatory phenotypes of macrophages (Leipner et al., 2021; Seneviratne et al., 2017). Despite these findings, the role of IRF5 in shear stress-induced endothelial activation remains largely unknown. Our study provides evidence that disturbed shear stress is sufficient to induce IRF5 nuclear translocation and activation in ECs. Furthermore, IRF5 knockdown in ECs significantly reduced disturbed flow-induced plaque formation in LCAs. These findings suggest that targeting endothelial IRF5 may be an effective strategy for combating the early stages of atherosclerosis.

Given the key role of IRF5 in mediating inflammatory responses, it has been considered as an attractive therapeutic target, and various strategies have been developed to study and modulate its function (Almuttaqi & Udalova, 2019). For example, nanoparticle-delivered siRNA targeting IRF5 in macrophages promotes inflammation resolution, improves infarct healing, and attenuates post-myocardial infarction remodeling (Courties et al., 2014). Additionally, manipulating IRF5 protein levels through the E3 ubiquitin ligase, TRIM21, has been explored as a strategy for modulating its activity (Lazzari et al., 2014). Given the crucial physiological role of IRF5, strategies aimed at suppressing its pathophysiological activation without altering basal levels may offer additional benefits. Our study introduces a novel approach to inhibit IRF5 activation. We found that the novel nested gene-encoded protein, AFF3ir-ORF2, interacts with IRF5, leading to cytoplasmic retention and inactivation under disturbed shear stress conditions. Importantly, endothelium-specific supplementation with AFF3ir-ORF2 effectively attenuated endothelial activation and reduced the atherosclerotic plaque area in ApoE$^{-/-}$ mice, suggesting that targeting endothelial IRF5 activation with AFF3ir-ORF2 holds promise for the treatment of atherosclerosis. Furthermore, as emerging studies have highlighted the substantial contributions of IRF5 to autoimmune diseases (Graham et al., 2006), neuropathic pain (Masuda et al., 2014), obesity (Dalmas et al., 2015), and hepatic fibrosis (Alzaid et al., 2016), future research should investigate whether AFF3ir-ORF2 has beneficial effects in these contexts.

Conclusion

340 In conclusion, this study provides novel evidence that the disruption of AFF3ir-ORF2
341 expression under disturbed flow promotes endothelial inflammatory responses and
342 atherosclerosis. AFF3ir-ORF2 serves as an endogenous inhibitor of IRF5 by binding to IRF5
343 and preventing its nuclear translocation. Supplementation with endothelial AFF3ir-ORF2 may
344 be a promising therapeutic strategy for treating atherosclerosis.

Materials and methods

Animals

C57BL/6 and Apolipoprotein E-null (ApoE$^{-/-}$) mice were purchased from the Experimental Animal Centre of Military Medical Science Academy (Beijing, China). AFF3ir-ORF2-null (AFF3ir-ORF2$^{-/-}$) mice were acquired from Dr. Lingfang Zeng's Laboratory at King's College London (Zhao Y et al., 2024). To generate ApoE and AFF3ir-ORF2 double knockout mice (ApoE$^{-/-}$AFF3ir-ORF2$^{-/-}$), ApoE$^{-/-}$ mice were crossed with AFF3ir-ORF2$^{-/-}$ mice. The animals were maintained at 21 ± 1°C under a 12-hour light/dark cycle (lights on at 07:00, lights off at 19:00) with ad libitum access to water and standard chow unless specified otherwise. This study adhered to the *Guide for the Care and Use of Laboratory Animals* of the US National Institutes of Health (NIH Publication No. 85-23, revised 2011). All study protocols were approved by the Institutional Animal Care and Use Committee of Tianjin Medical University.

Carotid artery partial ligation surgery

The surgery was performed as we previously described (Yang et al., 2021). Briefly, mice were anesthetized with isoflurane (2-3%). A ventral midline incision (4-5 mm) was made in the neck and the left carotid artery was exposed through blunt dissection of subcutaneous fat and muscle tissue. The left external carotid, internal carotid, and occipital arteries were ligated with a 6-0 silk suture, leaving the superior thyroid artery intact. For adenovirus and lentivirus infection studies, adenovirus (Ad-ORF2 or Ad-Scramble) or lentivirus (lenti-shRNA-IRF5 or lenti-shRNA-Scramble) was introduced into the lumen of the left carotid artery and kept inside for 40 min. After infection, the adenovirus or lentivirus was released, and blood flow to the common carotid artery was restored. Mice were fed with a high-fat diet (TD88137, ENVIGO, USA) immediately after surgery and continued for 4 weeks.

Endothelial AFF3ir-ORF2 overexpression in mice

Endothelial-specific adeno-associated virus (AAV)-mediated CRISPR/Cas9 shuttle plasmid was constructed by Cell & Gene Therapy (Shanghai, China) as previously reported (Wang et al., 2016). ApoE$^{-/-}$ mice received a single tail vein injection of recombinant AAV containing an endothelial-specific human ICAM-2 promoter driving AFF3ir-ORF2 overexpression (AAV-AFF3ir-ORF2) or a control empty vector (AAV-Scramble), with a dose of 1×10^{11} viral genomes in a 200 μL volume of sterile PBS. Subsequently, the mice were fed with a high-fat diet (TD88137, ENVIGO, USA) for 3 months.

Oil-Red O staining for atherosclerotic plaques in mouse aorta

The ApoE$^{-/-}$, ApoE$^{-/-}$AFF3ir-ORF2$^{-/-}$, and EC-specific AFF3ir-ORF2 overexpression mice were anaesthetized by inhalation of 2% isoflurane and euthanized by cervical dislocation. The aortas were dissected in 1×PBS and opened to expose the atherosclerotic plaques. After fixation in 4% formaldehyde for 1 hat 4 °C, the tissues were rinsed in water for 10 min, followed by 60% isopropanol. The aortas were then stained with Oil-Red O for 30 min with gentle shaking, rinsed again in 60% isopropanol, and subsequently rinsed in water three times. The samples were mounted on wax with the endothelial surface facing upwards. Images were captured using an HP Scanjet G4050. Plaque areas were quantified using NIH ImageJ software by calculating the plaque area relative to the total vascular area.

Immunofluorescence staining

MEFs slides or frozen sections were fixed in 4% paraformaldehyde for 30 min, then permeabilized in 0.1% Triton X-100 (in PBS) and blocked with 1% bovine serum albumin for 30 min at room temperature. Sections were incubated overnight at 4°C with primary antibodies (1:100). The vWF (Cat. No. ab11713) and VE-Cadherin (Cat. No. ab33168) antibodies were obtained from Abcam (Cambridge, UK). VCAM-1 (Cat. No. sc-13160) antibody was from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Following primary antibody incubation, sections were treated with Alexa Fluor 488- or Alexa Fluor 594- conjugated secondary antibodies (1:200, Thermo Fisher Scientific, Grand Island, NY, USA) at room temperature for 1 hour. Slides were then mounted with DAPI-containing mounting medium. Antibody specificity and target staining authenticity were verified using negative controls. Immunofluorescence micrographs were acquired using a Leica confocal laser scanning microscope. Representative images were randomly selected from each group.

Histological analysis of atherosclerotic lesions

Harvested carotid arteries and cross-sections of aortic roots were fixed in 4% paraformaldehyde and embedded in optimal cutting temperature compound (OCT). OCT-embedded tissues were sectioned at a thickness of 7 μm. Slides were immersed in 1×PBS for 5 min to remove OCT, and subsequently stained with Oil-Red O, haematoxylin and eosin (HE), and Masson's trichrome stain to assess lipid accumulation, lesion area, and collagen deposition, respectively (B. Li et al., 2019). Images were acquired using microscopy.

Quantification of plasma lipid levels

Blood samples were obtained via cardiac puncture, rinsed with heparin, and collected in 1.5

mL Eppendorf tubes. Total plasma cholesterol, triglycerides, LDL cholesterol, and HDL cholesterol levels were measured enzymatically using an automated clinical chemistry analyser kit (Biosino Biotech, Beijing, China).

Cell culture, transfection, and shear stress experiments

Mouse Embryonic Fibroblasts (MEFs) were obtained and cultured as previously described (Ferreira & Hein, 2023). Cell passages 4 to 7 were used in all experiments. MEFs and HEK293 cells were cultured in the Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% FBS, penicillin (100 U/mL), and streptomycin (100 μg/mL). Cells were incubated at 37°C in a humidified environment containing 5% CO_2 and grown to 70-80% confluence before treatment.

Small interfering RNA against IRF5 or IRF8 were synthesized from General Biosystems (Hefei, China). The sequences of siRNAs are shown in the Table S1. The MEFs were passaged to 6-well plated and transfected with 20 nmol/L siRNA per well using the Lipofectamine RNA iMAX Reagent (Invitrogen, Carlsbad, CA, USA).

For flow experiments, confluent monolayers of MEFs were seeded onto glass slides, and a parallel plate flow system was used to launch oscillatory flow (0.5 ± 4 dyn/cm^2). The flow system was enclosed in a chamber (Frangos, Eskin, McIntire, & Ives, 1985; Fu et al., 2011).

Adenovirus and lentivirus production and infection

AFF3ir-ORF2 sequences were inserted into the GV138 vector (CMV-MCS-3FLAG) to generate recombinant adenovirus (Ad-AFF3ir-ORF2). The short hairpin RNA (shRNA) sequences targeting mouse IRF5 were 5'-GGGACAACACCATCTTCAAGG-3', 5'GGTTGCTGCTGGAGATGTTCT-3', and 5'-GCCTAGAGCAGTTTCTCAATG-3'. The control shRNA was 5'-GCGTGATCTTCACCGACAAGA-3'. These shRNAs were constructed and cloned into pLV-U6-shRNA-CMV-EGFP to generate recombinant lentivirus (lenti-shRNA-IRF5 or lenti-shCtrl). MEFs were infected with adenovirus or lentivirus at a multiplicity of infection (MOI) of 10, with no detectable cellular toxicity observed.

Western blot analysis

Whole-cell lysates were prepared in a lysis buffer containing a complete protease inhibitor cocktail, PhosSTOP, and PMSF (Roche, Mannheim, Germany). Cytoplasmic and nuclear proteins were extracted from wild-type and AFF3ir-ORF2$^{-/-}$ MEFs using a protein extraction kits (Invent Biotechnologies, SC-003, Beijing, China). Protein were separated by SDS-PAGE and transferred to nitrocellulose membranes (Cat. No. 10600001; GE Healthcare; Chicago, IL,

USA). The membranes were incubated with primary antibodies. IRF5 (Cat. No. 96527), IRF8

(Cat. No. 98344), and Flag (Cat. No. 14793) antibodies were from Cell Signaling Technology

(Danvers, MA, USA). ICAM-1 (Cat. No. ab222736) antibodies were from Abcam

(Cambridge, UK). AFF3ir-ORF2 (Cat. No. C0302HL300-4) and AFF3ir-ORF1 (Cat. No.

C0302HL300) antibodies were from GenScript (NJ, USA). GAPDH (Cat. No. 60004-1-Ig)

antibody was from Proteintech (Wuhan, China). AFF3 (Cat. No. PA5-68961) antibody was

from Thermo Fisher Scientific (Waltham, MA, USA).

After incubation with horseradish peroxidase-conjugated secondary antibodies, the proteins

were visualized using enhanced chemiluminescence reagents in a ChemiScope3600 Mini

chemiluminescence imaging system (Clinx Science Instruments; Shanghai, China). Protein

levels were quantified by measuring integrated density with NIH Image J software

(https://imagej.nih.gov/ij/), using GAPDH as a loading control for normalization.

Co-immunoprecipitation

Whole-cell lysates were prepared by lysing cells in a 1% NP-40 lysis buffer containing 50

mM Tris-HCl, 1% Nonidet-P40, 0.1% SDS, and 150 mM NaCl, supplemented with a

complete protease inhibitor cocktail (Cat. No. 04693132001; Roche, Indianapolis, IN, USA),

a phosphatase inhibitor (PhosSTOP; Cat. No. 04906845001; Roche), and PMSF (Cat. No.

IP0280; Solarbio Life Sciences; Beijing, China). Samples were incubated on ice for 30

minutes, then centrifuged at 12,000 g for 10 minutes, and the supernatant was transferred to a

new tube. Protein concentrations were determined using the BCA Protein Assay Kit (Thermo

Fisher Scientific, Grand Island, NY, USA).

For immunoprecipitation, 1000 μg of protein was incubated with specific antibodies at 4 ℃

for 12 hour with constant rotation. Subsequently, 50 μL of 50% Protein A/G PLUS-Agarose

beads was added, and the incubation continued for an additional 2 hr. Beads were washed five

times with the lysis buffer and collected by centrifugation at 12,000 g for 2 min at 4℃. After

the final wash, the supernatant was removed and discarded. Precipitated proteins were eluted

by resuspending the beads in 2×SDS-PAGE loading buffer and boiling for 5 min. The eluates

from immunoprecipitation were subjected to Western blot analysis.

Total RNA extraction and real-time quantitative PCR analysis

Total RNA was extracted from cells using an RNA extraction kits (Transgen Biotech, ER501-

01, Beijing, China). Reverse transcription was performed with a reverse transcription kit

(Thermo Fisher Scientific, Grand Island, NY, USA). Quantitative PCR was conducted using

SYBR Select (Thermo Fisher Scientific) according to the manufacturer's protocol, with GAPDH serving as the internal control. The primers for quantitative real-time PCR are listed in Supplementary Table 2.

Luciferase reporter assay

The IRF5-binding motif and the full-length ZNF217 promoter were ligated into pGl3-based plasmids (Genechem, Shanghai, China), as previously described (Qiao et al., 2022). HEK293T cells were seeded into 24-well plates and grown to 70-80% confluency. Cells were transfected with the firefly luciferase reporter plasmid containing the IRF5-responsive ZNF217 promoter along with a β-galactosidase reporter plasmid (Promega, Madison, WI, USA) for 24 hour. Subsequently, cells were then infected with adenovirus (Ad-ORF2 or Ad-Scramble) for an additional 24 hour. Relative luciferase activity was measured using a luciferase assay and normalized to β-galactosidase activity as determined by the β-Galactosidase Enzyme Assay System (Promega, Madison, WI, USA).

RNA-sequencing (RNA-seq)

RNA-seq was performed as we previously described (Z. Li et al., 2024). Wild-type (WT) and AFF3ir-ORF2$^{-/-}$ MEFs were harvested, and RNA was extracted using the MagicPure Total RNA Kit (TransGen, Beijing, China). Whole transcriptome RNA-seq analysis were conducted by the Beijing Genomics Institute (BGI). Paired-end sequencing in 150-bp length was performed using the DNBSEQ-G400 platform. Raw data was filtrated using SOAPnuke (v1.5.6). Differential gene expression analysis, with thresholds set at $P < 0.05$ and fold change ≥ 1.5, was performed via the BGI website (http://omiscribe.bgi.com). Pathway enrichment analysis was carried out using DAVID tools.

Statistical analysis

Statistics analyses were performed using GraphPad Prism 8.0. No sample outliers were excluded. At least six independent experiments were performed for all biochemical experiments and the representative images were shown. Unpaired Student's t test (2-tailed), 1-way ANOVA or 2-way ANOVA with Bonferroni multiple comparison post hoc test were used for analyses, as appropriate. Sample size, statistical method, and statistical significance are specified in Figures and Figure Legends. Levels of probabilities less than 0.05 were regarded as significant.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Acknowledgments

This work was supported by National Natural Science Foundation of China Grants (82330012, 82127808, 82270516, and 82070451), British Heart Foundation (FS-15/74/31669), Tianjin Science and Technology Program of China [21JCYBJC01590], and Research Project of Tianjin Municipal Health Commission [2023013].

Disclosures

None.

References:

Almuttaqi, H., & Udalova, I. A. (2019). Advances and challenges in targeting IRF5, a key regulator of

inflammation. *FEBS J, 286*(9), 1624-1637. doi:10.1111/febs.14654

Alzaid, F., Lagadec, F., Albuquerque, M., Ballaire, R., Orliaguet, L., Hainault, I., . . . Venteclef, N.

(2016). IRF5 governs liver macrophage activation that promotes hepatic fibrosis in mice and

humans. *JCI Insight, 1*(20), e88689. doi:10.1172/jci.insight.88689

Bennett, M. R., Sinha, S., & Owens, G. K. (2016). Vascular Smooth Muscle Cells in Atherosclerosis.

Circ Res, 118(4), 692-702. doi:10.1161/CIRCRESAHA.115.306361

Cai, H., Yao, Z., & Li, W. (2017). IRF-5 accelerates leukocyte adhesion to endothelial cells in

ischemia-reperfusion injury through regulating the transcription of VCAM-1. *Biochem Biophys*

Res Commun, 492(2), 192-198. doi:10.1016/j.bbrc.2017.08.044

Chen, Z., Peng, I. C., Cui, X., Li, Y. S., Chien, S., & Shyy, J. Y. (2010). Shear stress, SIRT1, and

vascular homeostasis. *Proc Natl Acad Sci U S A, 107*(22), 10268-10273.

doi:10.1073/pnas.1003833107

Courties, G., Heidt, T., Sebas, M., Iwamoto, Y., Jeon, D., Truelove, J., . . . Nahrendorf, M. (2014). In

vivo silencing of the transcription factor IRF5 reprograms the macrophage phenotype and

improves infarct healing. *J Am Coll Cardiol, 63*(15), 1556-1566.

doi:10.1016/j.jacc.2013.11.023

Dalmas, E., Toubal, A., Alzaid, F., Blazek, K., Eames, H. L., Lebozec, K., . . . Venteclef, N. (2015). Irf5

deficiency in macrophages promotes beneficial adipose tissue expansion and insulin

sensitivity during obesity. *Nat Med, 21*(6), 610-618. doi:10.1038/nm.3829

Davignon, J., & Ganz, P. (2004). Role of endothelial dysfunction in atherosclerosis. *Circulation, 109*(23

Suppl 1), III27-32. doi:10.1161/01.CIR.0000131515.03336.f8

Davis, M. J., Earley, S., Li, Y. S., & Chien, S. (2023). Vascular mechanotransduction. *Physiological*

Reviews, 103(2), 1247-1421. doi:10.1152/physrev.00053.2021

Ferreira, R., & Hein, N. (2023). Purification of Mouse Embryonic Fibroblasts (MEFs). *Curr Protoc,*

3(11), e921. doi:10.1002/cpz1.921

Frangos, J. A., Eskin, S. G., McIntire, L. V., & Ives, C. L. (1985). Flow effects on prostacyclin

production by cultured human endothelial cells. *Science, 227*(4693), 1477-1479.

doi:10.1126/science.3883488

Fu, Y., Hou, Y., Fu, C., Gu, M., Li, C., Kong, W., . . . Zhu, Y. (2011). A novel mechanism of

gamma/delta T-lymphocyte and endothelial activation by shear stress: the role of ecto-ATP

synthase beta chain. *Circ Res, 108*(4), 410-417. doi:10.1161/CIRCRESAHA.110.230151

Graham, R. R., Kozyrev, S. V., Baechler, E. C., Reddy, M. V., Plenge, R. M., Bauer, J. W., . . .

Alarcon-Riquelme, M. E. (2006). A common haplotype of interferon regulatory factor 5 (IRF5)

regulates splicing and expression and is associated with increased risk of systemic lupus

erythematosus. *Nat Genet, 38*(5), 550-555. doi:10.1038/ng1782

Herrington, W., Lacey, B., Sherliker, P., Armitage, J., & Lewington, S. (2016). Epidemiology of

Atherosclerosis and the Potential to Reduce the Global Burden of Atherothrombotic Disease.

Circ Res, 118(4), 535-546. doi:10.1161/CIRCRESAHA.115.307611

Hwang, J., Saha, A., Boo, Y. C., Sorescu, G. P., McNally, J. S., Holland, S. M., . . . Jo, H. (2003).
Oscillatory shear stress stimulates endothelial production of O2- from p47phox-dependent
NAD(P)H oxidases, leading to monocyte adhesion. *Journal of Biological Chemistry, 278*(47),
47291-47298. doi:10.1074/jbc.M305150200

Jo, H., Song, H., & Mowbray, A. (2006). Role of NADPH oxidases in disturbed flow- and BMP4-
induced inflammation and atherosclerosis. *Antioxid Redox Signal, 8*(9-10), 1609-1619.
doi:10.1089/ars.2006.8.1609

Keenan, A. B., Torre, D., Lachmann, A., Leong, A. K., Wojciechowicz, M. L., Utti, V., . . . Ma'ayan, A.
(2019). ChEA3: transcription factor enrichment analysis by orthogonal omics integration.
Nucleic Acids Res, 47(W1), W212-W224. doi:10.1093/nar/gkz446

Kobiyama, K., & Ley, K. (2018). Atherosclerosis. *Circ Res, 123*(10), 1118-1120.
doi:10.1161/circresaha.118.313816

Krausgruber, T., Blazek, K., Smallie, T., Alzabin, S., Lockstone, H., Sahgal, N., . . . Udalova, I. A.
(2011). IRF5 promotes inflammatory macrophage polarization and TH1-TH17 responses. *Nat
Immunol, 12*(3), 231-238. doi:10.1038/ni.1990

Lazzari, E., Korczeniewska, J., Ni Gabhann, J., Smith, S., Barnes, B. J., & Jefferies, C. A. (2014).
TRIpartite motif 21 (TRIM21) differentially regulates the stability of interferon regulatory factor
5 (IRF5) isoforms. *PLoS One, 9*(8), e103609. doi:10.1371/journal.pone.0103609

Leipner, J., Dederichs, T. S., von Ehr, A., Rauterberg, S., Ehlert, C., Merz, J., . . . Hardtner, C. (2021).
Myeloid cell-specific Irf5 deficiency stabilizes atherosclerotic plaques in Apoe(-/-) mice. *Mol
Metab, 53*, 101250. doi:10.1016/j.molmet.2021.101250

Li, B., He, J., Lv, H., Liu, Y., Lv, X., Zhang, C., . . . Ai, D. (2019). c-Abl regulates YAPY357
phosphorylation to activate endothelial atherogenic responses to disturbed flow. *J Clin Invest,
129*(3), 1167-1179. doi:10.1172/JCI122440

Li, Z., Zhang, J., Ma, Z., Zhao, G., He, X., Yu, X., . . . He, J. (2024). Endothelial YAP Mediates
Hyperglycemia-Induced Platelet Hyperactivity and Arterial Thrombosis. *Arterioscler Thromb
Vasc Biol, 44*(1), 254-270. doi:10.1161/ATVBAHA.123.319835

Libby, P., Buring, J. E., Badimon, L., Hansson, G. K., Deanfield, J., Bittencourt, M. S., . . . Lewis, E. F.
(2019). Atherosclerosis. *Nat Rev Dis Primers, 5*(1), 56. doi:10.1038/s41572-019-0106-z

Lv, J. J., Wang, H., Zhang, C., Zhang, T. J., Wei, H. L., Liu, Z. K., . . . Bian, H. (2024). CD147 Sparks
Atherosclerosis by Driving M1 Phenotype and Impairing Efferocytosis. *Circ Res, 134*(2), 165-
185. doi:10.1161/circresaha.123.323223

Masuda, T., Iwamoto, S., Yoshinaga, R., Tozaki-Saitoh, H., Nishiyama, A., Mak, T. W., . . . Inoue, K.
(2014). Transcription factor IRF5 drives P2X4R+-reactive microglia gating neuropathic pain.
Nat Commun, 5, 3771. doi:10.1038/ncomms4771

Nakashima, Y., Raines, E. W., Plump, A. S., Breslow, J. L., & Ross, R. (1998). Upregulation of VCAM-
1 and ICAM-1 at atherosclerosis-prone sites on the endothelium in the ApoE-deficient mouse.
Arterioscler Thromb Vasc Biol, 18(5), 842-851. doi:10.1161/01.atv.18.5.842

Nam, D., Ni, C. W., Rezvan, A., Suo, J., Budzyn, K., Llanos, A., . . . Jo, H. (2009). Partial carotid
ligation is a model of acutely induced disturbed flow, leading to rapid endothelial dysfunction

and atherosclerosis. *Am J Physiol Heart Circ Physiol, 297*(4), H1535-1543. doi:10.1152/ajpheart.00510.2009

Pinero, J., Bravo, A., Queralt-Rosinach, N., Gutierrez-Sacristan, A., Deu-Pons, J., Centeno, E., . . . Furlong, L. I. (2017). DisGeNET: a comprehensive platform integrating information on human disease-associated genes and variants. *Nucleic Acids Res, 45*(D1), D833-D839. doi:10.1093/nar/gkw943

Qiao, X., Lv, S., Qiao, Y., Wang, F., & Miao, L. (2022). Interferon regulatory factor 5-induced upregulation of zinc-finger protein 217 promotes pancreatic carcinoma progression. *Mol Med Rep, 25*(5). doi:10.3892/mmr.2022.12705

Roberts, B. K., Collado, G., & Barnes, B. J. (2024). Role of interferon regulatory factor 5 (IRF5) in tumor progression: Prognostic and therapeutic potential. *Biochim Biophys Acta Rev Cancer, 1879*(1), 189061. doi:10.1016/j.bbcan.2023.189061

Salem, S., Salem, D., & Gros, P. (2020). Role of IRF8 in immune cells functions, protection against infections, and susceptibility to inflammatory diseases. *Hum Genet, 139*(6-7), 707-721. doi:10.1007/s00439-020-02154-2

Saliba, D. G., Heger, A., Eames, H. L., Oikonomopoulos, S., Teixeira, A., Blazek, K., . . . Udalova, I. A. (2014). IRF5:RelA interaction targets inflammatory genes in macrophages. *Cell Rep, 8*(5), 1308-1317. doi:10.1016/j.celrep.2014.07.034

Sato, M., Taniguchi, T., & Tanaka, N. (2001). The interferon system and interferon regulatory factor transcription factors -- studies from gene knockout mice. *Cytokine Growth Factor Rev, 12*(2-3), 133-142. doi:10.1016/s1359-6101(00)00032-0

Savitsky, D. A., Yanai, H., Tamura, T., Taniguchi, T., & Honda, K. (2010). Contribution of IRF5 in B cells to the development of murine SLE-like disease through its transcriptional control of the IgG2a locus. *Proc Natl Acad Sci U S A, 107*(22), 10154-10159. doi:10.1073/pnas.1005599107

Seneviratne, A. N., Edsfeldt, A., Cole, J. E., Kassiteridi, C., Swart, M., Park, I., . . . Monaco, C. (2017). Interferon Regulatory Factor 5 Controls Necrotic Core Formation in Atherosclerotic Lesions by Impairing Efferocytosis. *Circulation, 136*(12), 1140-1154. doi:10.1161/CIRCULATIONAHA.117.027844

Sorescu, G. P., Song, H., Tressel, S. L., Hwang, J., Dikalov, S., Smith, D. A., . . . Jo, H. (2004). Bone morphogenic protein 4 produced in endothelial cells by oscillatory shear stress induces monocyte adhesion by stimulating reactive oxygen species production from a nox1-based NADPH oxidase. *Circ Res, 95*(8), 773-779. doi:10.1161/01.RES.0000145728.22878.45

Swiech, L., Heidenreich, M., Banerjee, A., Habib, N., Li, Y., Trombetta, J., . . . Zhang, F. (2015). In vivo interrogation of gene function in the mammalian brain using CRISPR-Cas9. *Nat Biotechnol, 33*(1), 102-106. doi:10.1038/nbt.3055

Takaoka, A., Yanai, H., Kondo, S., Duncan, G., Negishi, H., Mizutani, T., . . . Taniguchi, T. (2005). Integral role of IRF-5 in the gene induction programme activated by Toll-like receptors. *Nature, 434*(7030), 243-249. doi:10.1038/nature03308

Wang, L., Luo, J. Y., Li, B., Tian, X. Y., Chen, L. J., Huang, Y., . . . Huang, Y. (2016). Integrin-YAP/TAZ-JNK cascade mediates atheroprotective effect of unidirectional shear flow. *Nature,*

540(7634), 579-582. doi:10.1038/nature20602

Weiss, M., Blazek, K., Byrne, A. J., Perocheau, D. P., & Udalova, I. A. (2013). IRF5 is a specific

marker of inflammatory macrophages in vivo. *Mediators Inflamm, 2013*, 245804.

doi:10.1155/2013/245804

Weiss, M., Byrne, A. J., Blazek, K., Saliba, D. G., Pease, J. E., Perocheau, D., . . . Udalova, I. A.

(2015). IRF5 controls both acute and chronic inflammation. *Proc Natl Acad Sci U S A, 112*(35),

11001-11006. doi:10.1073/pnas.1506254112

Wen, L., Chen, Z., Zhang, F., Cui, X., Sun, W., Geary, G. G., . . . Shyy, J. Y. (2013). Ca2+/calmodulin-

dependent protein kinase kinase beta phosphorylation of Sirtuin 1 in endothelium is

atheroprotective. *Proc Natl Acad Sci U S A, 110*(26), E2420-2427.

doi:10.1073/pnas.1309354110

Wolf, D., & Ley, K. (2019). Immunity and Inflammation in Atherosclerosis. *Circ Res, 124*(2), 315-327.

doi:10.1161/CIRCRESAHA.118.313591

Yang, Y., Ma, Q., Li, Z., Wang, H., Zhang, C., Liu, Y., . . . He, J. (2021). Harmine alleviates

atherogenesis by inhibiting disturbed flow-mediated endothelial activation via protein tyrosine

phosphatase PTPN14 and YAP. *Br J Pharmacol, 178*(7), 1524-1540. doi:10.1111/bph.15378

Zhang, C., Zhou, T., Chen, Z., Yan, M., Li, B., Lv, H., . . . Ai, D. (2020). Coupling of Integrin alpha5 to

Annexin A2 by Flow Drives Endothelial Activation. *Circ Res, 127*(8), 1074-1090.

doi:10.1161/CIRCRESAHA.120.316857

Zhao Y, Ehteramyan M, Li Y, Bai X, Huang L, Gao Y, . . . Zeng L. (2024). A novel nested gene Aff3ir

participates in vascular remodeling by enhancing endothelial cell differentiation in mice. *Genes*

& Diseases. doi:10.1016/j.gendis.2024.101339.

Figures and Figure legends

Figure 1. Disturbed shear stress reduces the expression of AFF3ir-ORF2 in vivo and in vitro.

A–F, Aortas were isolated from the C57BL/6 mice. **A,** RT-PCR analysis of the mRNA levels

of *AFF3ir-ORF2* and AF4/FMR2 family member 3 (*AFF3*) in the thoracic aorta (TA) and

aortic arch (AA) of C57BL/6 mice. Data are presented as mean ± SEM (n=6 mice per group).

*$P<0.05$, unpaired two-tailed *t*-test. **B, C,** Western blot analysis of the expression of the

indicated proteins in the TA and AA of mice. Protein levels were normalized to those of

GAPDH, and the relative expression values were relative to those of the TA group. Data are

presented as mean ± SEM (n=6 mice per group). *$P<0.05$, unpaired two-tailed *t*-test. **D, E,**

En-face immunofluorescence staining of AFF3ir-ORF2, VE-cadherin, and DAPI, and

quantification of AFF3ir-ORF2 expression in AA and TA. Scale bar, 20 μm. The

immunofluorescence intensity of AFF3ir-ORF2 was normalized to that of DAPI, and the

relative expression of the values was relative to that of the TA group. Data are presented as

mean ± SEM (n=6 mice per group). *$P< 0.05$, unpaired two-tailed *t*-test. **F,** Representative

immunofluorescent staining for von Willebrand factor (vWF), AFF3ir-ORF2, and AFF3 in

longitudinal sections of mouse aortas. n=6 mice per group. Scale bar, 25 μm. Inner, inner

curvature of the AA; outer, outer curvature of the AA; BIF, Bifurcation. **G, H,** Mouse

embryonic fibroblasts (MEFs) isolated from the embryo of C57BL/6 mice were subjected to

static (ST) or oscillatory shear stress (OSS, 0.5 ± 4 dyn/cm^2, 1 Hz) for indicated time. Western

blot analysis of the indicated proteins. Protein levels were normalized to GAPDH and the

relative expression values were relative to that of the ST group. Data are mean ± SEM (n = 6

independent experiments). *$P< 0.05$, one-way ANOVA with Tukey post-test. **I,** MEFs were

subjected to ST or OSS treatment for 6 h. RT-PCR analysis of the mRNA levels of *AFF3ir-*

ORF2, *AFF3,* intercellular adhesion molecule 1 (*ICAM*-1), and vascular cell adhesion

molecule 1 (*VCAM-1*) in MEFs. Data are mean ± SEM (n = 6 independent experiments). *$P<$

0.05, unpaired two-tailed *t*-test.

Figure 2. AFF3ir-ORF2 overexpression alleviates disturbed flow-induced inflammation and atherosclerosis.

A–C, Mouse embryonic fibroblasts (MEFs) isolated from C57BL/6 mice were infected with indicated adenoviruses (Ad-Scramble or Ad-Aff3ir-ORF2) for 48 h and then exposed to static (ST) or oscillatory shear stress (OSS, 0.5 ± 4 dyn/cm^2, 1 Hz) for another 6 h. **A, B,** Western blot analysis of the indicated proteins in MEFs and quantification of their relative expression of proteins are shown. The protein levels were normalized to GAPDH and the relative

expression of the values is relative to MEFs infected with Ad-Scramble and treated with ST.

Data are presented as mean ± SEM (n = 6 independent experiments). *$P<$ 0.05, two-way

ANOVA with Tukey post-test. **C,** RT-PCR analysis of mRNA levels of *AFF3, AFF3ir,*

VCAM-1, ICAM-1, interleukin-6 (*IL-6*), and interleukin-1 beta (*IL-1β*). The relative

expression of the values is relative to MEFs infected with Ad-Scramble and treated with ST.

Data are presented as mean ± SEM (n = 6 independent experiments). *$P<$ 0.05, two-way

ANOVA with Tukey post-test. **D–H,** Eight-week-old male ApoE$^{-/-}$ mice were subjected to

partial ligation of the carotid artery along with 10 μL of adenovirus suspension at 1×10^8

transducing units (TU)/mL was instilled into the left carotid artery (LCA). The mice were then

fed high-fat diet for 4 weeks. **D,** Arterial tissues were isolated to examine the atherosclerotic

lesions. Scale bar, 2 mm. **E, F,** LCAs were sectioned for hematoxylin and eosin staining and

quantification of the lesion area. Scale bar, 25 μm. Data are presented as mean ± SEM (n=6

mice per group). *$P<$ 0.05, unpaired two-tailed *t*-test. **G**, **H,** Immunofluorescence staining for

vWF, VCAM-1, and DAPI in the LCA, and quantification of the relative fluorescent intensity

of VCAM-1. The immunofluorescence intensity of VCAM-1 was normalized to DAPI, and

the relative expression of the values were relative to that of the Ad-Scramble group. Scale bar,

50 μm. Data are presented as mean ± SEM (n=6 mice per group). *$P<$ 0.05, unpaired two-

tailed *t*-test.

Figure 3. AFF3ir-ORF2 deletion aggravates inflammation and atherosclerotic lesions in

ApoE$^{-/-}$ mice.

Eight-week-old male ApoE$^{-/-}$ and ApoE$^{-/-}$AFF3ir-ORF2$^{-/-}$ mice were fed a high-fat diet for 12

weeks. Arterial tissues and aortic roots were isolated to examine atherosclerotic lesions. **A,**

Schematic of experimental strategy. **B,** Representative images of en face Oil-Red O staining

of the aortas. Scale bar, 4 mm. **C,** Quantification of the plaque area in the aortas. Data are

presented as mean ± SEM (n=12 mice per group). *P< 0.05, unpaired two-tailed t-test. **D,** Oil-

Red O, hematoxylin and eosin (HE), and Masson staining of the aortic roots. Scale bars, 500

μm. **E,** Quantification of plaque size, Oil-Red O-positive area, and collagen fiber content in

aortic root sections. Data are presented as mean ± SEM (n=6 mice per group). *P< 0.05,

unpaired two-tailed t-test. **F**, LCAs were sectioned and stained with Oil-Red O, HE, and

Masson's trichrome. Scale bars, 500 μm. **G,** Quantification of plaque size, Oil-Red O-positive

area, and collagen fiber content in the LCA sections. Data are presented as mean ± SEM (n=6

mice per group). *P< 0.05, unpaired two-tailed t-test. **H**: Representative immunofluorescence

images of vWF, VCAM-1, and DAPI in the aortic roots. Scale bar, 500 μm. **I**, Quantification

of the relative fluorescence intensity of VCAM-1. The immunofluorescence intensity of

VCAM-1 was normalized to that of DAPI, and the relative expression of the values were

relative to that of the ApoE$^{-/-}$ group. Data are presented as mean ± SEM (n=6 mice per group).

*P< 0.05, unpaired two-tailed t-test.

Figure 4. AFF3ir-ORF2 mitigates disturbed shear stress-induced inflammation by interacting with IRF5 and retaining it within the cytosol.

A–D, Mouse embryonic fibroblasts (MEFs) were isolated from wild-type (WT) and AFF3ir-ORF2$^{-/-}$ mice. **A**, Heat map showing mRNA profiles of WT and AFF3ir-ORF2$^{-/-}$ MEFs (n=3). **B**, Volcano map displays the differentially expressed genes between the groups. **C,** Gene

Ontology enrichment pathway analysis of the differentially expressed genes. **D**, Venn diagrams of the top 20 transcription factors from the ChEA3 and DisGENET analysis related to atherosclerosis. **E,** Immunoprecipitation performed using antibodies against AFF3ir-ORF2, IRF5, and IRF8. n = three independent experiments. **F–H,** WT and AFF3ir-ORF2$^{-/-}$ MEFs were subjected to silence of Control (siNC), IRF5 (siIRF5), or IRF8 (siIRF8) with siRNAs for 24 h, followed by exposure to Static (ST) or oscillatory shear stress (OSS, 0.5 ± 4 dyn/cm^2, 1 Hz) for another 6 h. **F**, RT-PCR analysis of the mRNA levels of *VCAM-1, ICAM-1, IL-6,* and *IL-1b*. The relative expression of the values is relative to WT MEFs transfected with siNC and treated with ST. Data are mean $\pm$ SEM (n = 6 independent experiments). *$P < 0.05$, two-way ANOVA with Tukey post-test. **G, H**, Representative western blots of IRF5 and ICAM-1 expression. Data are mean $\pm$ SEM (n = 5 independent experiments). *$P < 0.05$, two-way ANOVA with Tukey post-test. **I, J**, WT, and AFF3ir-ORF2$^{-/-}$ MEFs were exposed to ST or OSS for 6 h. Nuclear and cytoplasmic proteins were extracted from the cells. Representative western blots of the indicated proteins and quantification of IRF5 expression in nucleus are shown. The expression of these proteins was relative to the level of nuclear IRF5 in ST-treated WT MEFs. Data are mean $\pm$ SEM (n = 6 independent experiments). *$P < 0.05$, two-way ANOVA with Tukey post-test. **K,** HEK293 cells were transfected with the firefly luciferase reporter plasmid containing the IRF5-responsive ZNF217 promoter along with a β-galactosidase reporter plasmid for 24 hours. Cells were infected with the indicated adenoviruses (Ad-Scramble or Ad-Aff3ir-ORF2) for 24 h. Promoter activity was measured using luciferase, which was normalized to β-gal. Data are mean $\pm$ SEM (n = 6 independent experiments). *$P < 0.05$, two-way ANOVA with Tukey post-test.

Figure 5. IRF5 knockdown prevents the aggravation of atherosclerosis in AFF3ir-ORF2 deficienct mice.

Eight-week-old male ApoE$^{-/-}$ mice were subjected to partial ligation of the left carotid artery (LCA) along with 10 μL of lentivirus suspension at 1×10^8 transducing units (TU)/mL was instilled into the LCA. The mice were then fed a high-fat diet for 4 weeks. **A,** Arterial tissues were isolated to examine the atherosclerotic lesions. Scale bar, 2 mm. **B,** LCAs were

sectioned for hematoxylin and eosin staining and quantification of the lesion area. Scale bar,

25 μm. Data are mean ± SEM (n=6 mice per group). *$P < 0.05$, two-way ANOVA with Tukey

post-test. **C, D,** Immunofluorescence staining for vWF, VCAM-1, and DAPI in the LCAs and

quantification of the relative fluorescent intensity of VCAM-1. Scale bar, 50 μm. The

immunofluorescence intensity of VCAM-1 was normalized to DAPI, and the relative

expression of the values were relative to that of the group of ApoE$^{-/-}$ mice infected with Ad-

scramble. Data are presented as mean ± SEM (n=6 mice per group). *$P < 0.05$, two-way

ANOVA with Tukey post-test.

Figure 6. Endothelial-specific AFF3ir-ORF2 supplementation alleviates EC activation and atherosclerosis in ApoE$^{-/-}$ mice.

Eight-week old ApoE$^{-/-}$ male mice were infused with the indicated adeno-associated virus (AAV) and then fed a high-fat diet for 12 weeks. **A,** Schematic of the experimental strategy. **B**, Representative images of enface Oil-Red O staining of the aortas. Scale bar, 4 mm. **C,** Quantification of the plaque area in the entire aortas. Data are presented as mean ± SEM (n=6 mice per group). *$P< 0.05$, unpaired two-tailed t-test. **D,** Hematoxylin and eosin (HE), Oil-Red O, and Masson staining of the aortic roots. Scale bars, 500 μm. **E,** Quantification of

plaque size, Oil-Red O-positive area, and collagen fiber content in aortic root sections. Data

are presented as mean ± SEM (n=6 mice per group). *$P< 0.05$, unpaired two-tailed *t*-test. **F**,

Representative immunofluorescence image of vWF, VCAM-1, and DAPI in the aortic roots.

Scale bar, 500 μm. **G**, Quantification of the relative fluorescent intensity of VCAM-1. The

immunofluorescence intensity of VCAM-1 was normalized to that of DAPI, and the relative

expression of the values were relative to that of the AAV-Scramble group. Data are presented

as mean ± SEM (n=6 mice per group). *$P< 0.05$, unpaired two-tailed *t*-test.

Figure 7. Schematic illustration of the AFF3ir-ORF2/IRF5 cascade in disturbed flow-induced endothelial activation and atherosclerosis.

Disturbed flow induced a down-regulation of AFF3ir-ORF2, which could interact with IRF5 and promote the latter's retention in the cytoplasm, thereby boosting IRF5-dependent inflammatory pathways in endothelial cells and leading to atherogenesis.