The G3BP Stress-Granule Proteins Reinforce the Translation Program of the Integrated

Stress Response

Jarrett Smith[1,2,3] & David P. Bartel[1,2,3]

[1]Howard Hughes Medical Institute and [2]Whitehead Institute for Biomedical Research, 9 Cambridge Center, Cambridge, MA 02142, USA

[3]Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

Correspondence: dbartel@wi.mit.edu

ABSTRACT

When mammalian cells are exposed to extracellular stress, they coordinate the condensation of stress granules (SGs) through the action of key nucleating proteins G3BP1 and G3BP2 (G3BPs) and, simultaneously, undergo a massive reduction in translation.[1-5] Although SGs and G3BPs have been linked to this translation response, their overall impact has been unclear. Here, we investigate the longstanding question of how, and indeed whether, G3BPs and SGs shape the stress translation response. We find that SGs are enriched for mRNAs that are resistant to the stress-induced translation shutdown. Although the accurate recruitment of these stress-resistant mRNAs does require the context of stress, a combination of optogenetic tools and spike-normalized ribosome profiling demonstrates that G3BPs and SGs are necessary and sufficient to both help prioritize the translation of their enriched mRNAs and help suppress cytosolic translation. Together these results support a model in which G3BPs and SGs reinforce the stress translation program by prioritizing the translation of their resident mRNAs.

INTRODUCTION

When cells are exposed to environmental stresses, including oxidative stress, high temperature, harmful chemicals, and viral infection, the cells initiate the integrated stress response (ISR).[5-8] This response integrates signals from different stress sensors and ultimately results in reduced global translation, through phosphorylation of a critical translation initiation factor eif2α.[5-9] At the same time, select mRNAs continue to be translated or even undergo enhanced translation to help cope with and recover from the stress, whereas mRNAs whose products are energetically expensive for the cell are especially sensitive to down regulation.[6,8,10-13] During this dramatic alteration of the translation state, cytoplasmic puncta known as stress granules (SGs) form through the process of liquid-liquid phase separation.[2-5,10,14-16] These granules are non-membrane-bound collections of RNA-binding proteins, mRNAs, and translation machinery.[1,5,15,17-19] Although SGs contain a diverse proteome, they require the RNA-binding proteins G3BP1 and G3BP2 (G3BPs) for their formation.[1,2] Both SGs and G3BPs have been hypothesized to play roles in translation,[1,20-23] but their precise impacts on translation during the stress response are not well understood.

Because they are highly enriched for translation machinery and form concurrently with translation shutdown, SGs were originally hypothesized to be required for the dramatic translation shutdown observed during the ISR.[20,24,25] However, more recent studies indicate that SGs are not required for the massive reduction in translation that occurs during the ISR; G3BP KO cell lines, which are devoid of microscopically visible stress granules, still undergo the eif2α-dependent translation reduction associated with the ISR.[1,26]

Because the formation of SGs is enhanced by polysome collapse, and SGs are enriched in small ribosomal subunits but not large ribosomal subunits, SGs are also proposed to simply be sites of devoid of translation.[5,6,14,20,27,28] However, single-molecule analyses show that translating mRNAs are not forbidden inside of SGs,[29] and some report that translation inside of SGs might not be a rare event compared to translation in the cytosol.[30]

G3BPs, in addition to being required for formation of canonical SGs,[1,2] are also sufficient to nucleate SG assembly in the absence of stress, either by overexpression or by optogenetic manipulation.[31,32] These proteins also play roles in unstressed cells, including translational

upregulation of target mRNAs in fly cells and rescue of ribosomal subunits from lysosomal
degradation.[21,23]
Despite their many connections to translation and the stress response, the overall impact
that G3BPs and SGs have on translation during stress or, indeed, whether they have any impact
on translation during stress is not fully understood.[14,33-35] Previous attempts to determine the
effects of G3BPs or SGs on translation have relied either on bulk measurements of translation
with limited sensitivity and specificity, such as S^{35} labeling of nascent peptides, or have been
restricted to observations of single genes and reporters.[1,21,30] Here, we use spike-normalized
ribosome profiling and optogenetic methods to measure the impact of G3BPs and SGs on
translation with and without stress. These results support a model in which G3BPs and SGs
reinforce the ISR translation program by prioritizing the translation of their most enriched
transcripts.
78

RESULTS
Ribo-spike Enables Observation of Absolute Translation Changes during the ISR
A variety of cellular stresses induce the ISR, resulting in eif2α phosphorylation and a drastic
global reduction in translation. Such global changes to translation have been challenging to
observe by ribosome profiling because ribosome profiling typically reports the relative
translation efficiencies (TEs) in each sample with no information on the absolute differences
between samples.[36] In other high-throughput approaches, spiking known amounts of
orthogonal standards into to each starting sample enables quantification of absolute
differences. However, spiking in short RNAs resembling ribosome-protected fragments (RPFs)
would not be appropriate for samples analyzed by ribosome profiling, as these samples are
treated with RNAse to create RPFs, which would destroy the unprotected spike molecules.
Normalizing to endogenous mitochondrial RPFs can bypass this problem.[36,37] However, an
assumption underlying this approach is that mitochondrial translation is unaffected by the
experimental condition—an assumption not always expected to be valid, especially if cells are
subjected to stress.
To use ribosome profiling to measure absolute changes to TE occurring as a consequence of
stress, we developed a ribosome profiling spike-in (ribo-spike) (Fig 1A). Firefly luciferase mRNA
was translated in vitro, using a rabbit-reticulocyte translation system, and while the reaction
was undergoing maximal translation, it was treated with cycloheximide to block ribosome
translocation and flash frozen for storage. A defined amount of this reaction containing
ribosomes translating the orthogonal firefly luciferase mRNA was then spiked into each sample
as the ribo-spike. To confirm that these ribo-spikes functioned as intended, we added them in
equal quantities to a dilution series of cellular lysates and performed ribosome profiling on
these lysates. For both RPF and mRNA measurements, concentrations of the dilution series
calculated based on the spike-ins closely matched the known dilutions (Fig S1A).
To apply the ribo-spike in the context of the ISR, we stressed cells with 500 μM sodium
arsenite, which causes oxidative stress, for 1 h and compared them to unstressed cells using
ribosome profiling with our ribo-spike. This dose of sodium arsenite activates the ISR, leading to
eif2α phosphorylation and substantial global reduction in translation.[1,20] By normalizing to the
spike-in TE across samples we were able to detect and measure absolute changes in TE for

mRNAs of >10,000 individual genes. Aggregating these measurements revealed a drastic (20-
fold) global reduction in TE, consistent with previous bulk measurements (Fig 1B and
S1B,C).[1,38,39] Interestingly, TEs of mitochondrial mRNAs increased in the presence of sodium
arsenite (Fig 1B), validating our concern that the TEs of mitochondrial mRNAs might not remain
constant under stress conditions.

Against the backdrop of a drastic global reduction of translation during stress, some mRNAs
retained or even enhanced their translation in response to stress. Among them were mRNAs of
several stress-responsive genes previously reported to maintain their translation during the ISR,
including *ATF4*, *ATF5*, *CEBPα*, and *DDIT3/CHOP* (Fig 1B). Many of these mRNAs are known to
contain upstream open reading frames (uORFs), which can help regulate their translation during
stress.[40] Indeed, regulation by uORFs is proposed to help shape the general translation response
to stress.[6,8,11,12,41,42] Supporting this proposal, among the mRNAs with retained or enhanced
translation, we observed a strong bias in those reported to contain one or more functional uORF
(Fig S1D).[43]

In yeast, newly transcribed mRNAs tend to be preferentially translated during stress.[44,45] To
assess this possibility in mammalian cells, we compared the translation change in response to
stress to mRNA half-lives, reasoning that mRNAs with short half-lives would be predominantly
composed of newly synthesized transcripts. Consistent with the observations made in yeast, we
observed a moderate negative correlation between the translation change of an mRNA upon
stress and its estimated half-life (Fig S1E; R_S = –0.22).[46]

At the other extreme, some mRNAs were especially sensitive to the translation shutdown,
including *CSDE1* and *PCBP2*, consistent with a previous study.[42] Interestingly, other
hypersensitive mRNAs encoded proteins involved in energetically expensive processes, such as
vesicle trafficking (*COPS3*, *COPS6*, *CLTC*, *SCFD1*), which are strongly down-regulated during stress
(Fig 1B).[47]

To measure absolute TE changes in another stress context, we performed analogous
experiments in cells exposed to 45°C heat shock (Fig 1C and S1B,C). The results resembled those
observed for arsenite stress. First, normalizing to the ribo-spike again revealed a large reduction
in global translation, although the median reduction of 8-fold was somewhat less than the 20-
fold reduction caused by arsenite stress. Second, mitochondrially translated mRNAs again had
higher TE in stressed cells, indicating that mitochondrial translation might generally be altered
in the context of global cellular stress. Third, mRNAs with retained or enhanced TE again
included those of previously identified stress-responsive, uORF-containing genes (*ATF4*, *ATF5*,
CEBPα, *DDIT3/CHOP*, and *BiP/GPR78*), and short half-life transcripts (Fig 1C and S1F,G). Fourth,
mRNAs hypersensitive to heat shock translation shut down again included *CSDE1*, *PCBP2*, and
CLTC, as well as the mRNA for the vesicle trafficking gene *VSP4A* (Fig 1C). Despite this overlap
and an overall correlation between the changes observed with these two different stressors
(Spearman *R*, R_S = 0.63), the translation changes were not identical, and many genes were called
as being differentially regulated in one stress but not the other (Fig S1H–J). This observation
concurred with previous reports that the extent to which specific mRNAs are up- or
downregulated can depend on the particular type and dosage of stress.[7]

We consider the combination of widespread translation shutdown, the retained or
increased translation of select transcripts, and the hypersensitive downregulated translation of
others to be the ISR translation program. The ability to observe these ISR translation programs

Figure 1. Ribo-spike enables observation of absolute translation changes during ISR. (A) Schematic representation of the ribo-spike. The ribo-spike consists of polysomes formed on an orthogonal mRNA sequence, in this case firefly luciferase mRNA (*fLuc*) translated in a rabbit reticulocyte lysate. A defined amount of the ribo-spike sample is added to each experimental sample, ultimately generating orthogonal RPFs that enable normalization between samples. **(B)** The ISR translation program induced upon arsenite stress. Plotted for each mRNA is the $\log_2$ fold-change (FC) in TE observed in HCT116 cells responding to 500 µM sodium arsenite (for 1 h) as a function of the mRNA expression. The red dashed line indicates unchanged TE, as determined by the ribo-spike. Points for stress-enhanced, stress-resistant, and stress-hypersensitive mRNAs are indicated by colors, with exemplar mRNAs listed for each category. Stress response categories were determined by DESeq2 ($\log_2$ FC > 1, adjusted *p* value < 0.05). Mitochondrially translated mRNAs are indicated in orange. *n* indicates number of unique mRNAs. **(C)** The ISR translation program induced upon heat shock (45°C for 25 min); otherwise, as in (A).

both under arsenite stress and heat shock illustrates how using ribo-spike enables accurate and

precise monitoring of absolute translation changes between samples and emphasizes the utility

of a truly orthogonal spike-in for studies monitoring global changes to translation.

G3BPs Reinforce the ISR Translation Program

Having developed a method to globally monitor absolute TE changes, we wanted to use it to

determine the effects of G3BPs and SGs on the cellular stress response. To do so, we also

needed to be able deplete cells of G3BPs and SGs. Therefore, we created a cell line in which

G3BPs could be inducibly depleted using an auxin-inducible degron (AID).[48,49] In HCT116 cells,
we tagged both alleles of both endogenous G3BP1 and endogenous G3BP2 with the AID and
RFP (AID::RFP::G3BPs). Adding indole-3-acetic acid (IAA, a form of auxin) induced rapid and
efficient depletion of both G3BPs, with approximately 90% of each protein being depleted
within 3 h, while imparting minimal effects on the stability of another SG-resident protein
PABPC1 (Fig 2A–C).
To determine whether our AID-mediated depletion of G3BPs was sufficient to prevent
formation of SGs, we labeled endogenous PABPC1, a canonical SG marker, with GFP and
assessed SG formation after subjecting cells to oxidative stress by treatment with sodium
arsenite. As expected, depletion of G3BPs prior to stress largely prevented formation of SGs, as
marked by PABPC1, with < 3% of cells still able to form visible SGs (Fig 2C,D). To examine if this
result was generalizable across different stresses, we repeated this experiment using heat shock
instead of arsenite stress. Under this different stress, results were similar, with < 5% of cells able
to form visible SGs (Fig 2E,F). These results were consistent with those of previous experiments
that use long-term knockdown or knockout of G3BPs to show that these proteins are necessary
nucleators of SG assembly.[1,2]
Having generated these degron lines, we set out to determine the impact of G3BPs and SGs
on translation during stress, using ribosome profiling with the ribo-spike to monitor translation
differences in cells subjected to arsenite stress either with or without prior depletion of G3BPs,
and thus with or without the ability to form SGs (Fig 2G). Compared to G3BP-depleted cells,
normal cells underwent a somewhat stronger inhibition of global translation (1.4-fold, Fig S2A).
This modest effect was consistent with reports that G3BP is largely dispensable for the global
reduction in translation caused by the ISR.[1]
Although the effect was relatively small, we were nonetheless intrigued to find that G3BPs
and perhaps SGs did seem to contribute, if only modestly, to the massive reduction in
translation that occurs during the ISR. To further examine this contribution of G3BPs and
potentially SGs to the ISR translation program, we compared the translation changes caused by
presence of G3BPs to those caused by activation of the ISR. We observed a clear positive
correlation (R_S = 0.33) between the mRNAs regulated by G3BPs and those regulated by the ISR
(Fig 2H and S2B). This implied that G3BPs, and perhaps SGs, tend to upregulate the same
mRNAs that the ISR upregulates and tend to downregulate the same mRNAs that the ISR
downregulates, thereby reinforcing the ISR translation program.
Given that the ISR translation program for arsenite-treated cells and heat-shocked cells were
distinct, we examined whether G3BPs also reinforced the heat shock ISR translation program.
When comparing the G3BP-dependent translational changes observed during heat shock to the
heat-shock-induced ISR translation changes, the results resembled those observed for arsenite
stress. Although the G3BP-dependent reduction to translation was negligible, especially when
compared to the much larger reduction caused by the ISR (1.06- and 8-fold, respectively; Fig 1C
and S2C), we again observed a positive correlation (R_S = 0.43) between changes of mRNAs
regulated by G3BPs and those regulated by the ISR (Fig 2J and S2D). Taken together, these
results indicated that G3BPs, and perhaps SGs, reinforce the ISR translation program across
multiple stresses.
To examine whether the factors we observed impacting mRNA translation during stress
(presence of uORFs and short half-life) also impacted the effect of G3BP on translation, we

compared both uORF presence and estimated mRNA half-lives to G3BP-dependent translation
in cells treated with arsenite or heat shock. We observed that mRNAs containing one or more
functional uORF were preferentially translated in the presence of heat stress. However this
trend did not hold in arsenite-stressed cells (Fig S2E,F), indicating that, although uORFs may play
a role in G3BP-dependent translation during stress, it appears to be a minor or stress-specific
role. Furthermore, we observed no correlation between the half-life of an mRNA and its G3BP-
dependent translation (Fig S2G,H), indicating that, although newly synthesized mRNAs might be
translationally favored, their regulation appears to be stress-granule independent.
To determine whether the G3BP-dependent translation effects that we observed were a
constitutive or stress-specific function of G3BPs— potentially through formation of SGs, we
performed identical ribosome profiling experiments, comparing cells containing G3BPs to cells
depleted for G3BPs but this time in the absence of stress. This allowed us to quantify
transcriptome-wide G3BP-dependent translation changes in the absence of stress. When
comparing G3BP-dependent translation changes in unstressed cells to the changes caused by
activation of the ISR, unstressed G3BP-dependent translation changes had no positive
correlation to the translation changes of either the arsenite-induced ISR or the heat-shock-
induced ISR (R_S = −0.033 and −0.12 respectively; Fig 2I, K). Together these results argued that
G3BPs reinforce the ISR translation program in a stress-dependent manner and were consistent
with a model in which this is a function of G3BP through its nucleation of SGs.
Because G3BPs are required for SG formation, we could not say whether the effects we
observed upon depleting G3BPs were a consequence of losing SGs or whether they were a
consequence of losing some other G3BP function. We sought to disambiguate G3BP and SG
function by targeting CAPRIN1 or UBAP2L, two other proteins reported to be required for SG
formation,[1,19,50,51] reasoning that observing the same effects when SGs were depleted by
removing a different protein would support the idea that the effects were a consequence of
removing SGs. We first targeted CAPRIN1, a G3BP interactor that colocalizes to SGs, using the
AID system and assessed SG formation by following G3BP and eIF4G localization. No reduction
in SG formation was detected (Fig S2I – M). We then targeted UBAP2L, another SG-localized
protein, and assessed granule formation by following PABPC1 localization. Again, no significant
reduction in SG formation was detected (Fig S2N – Q). These results concurred with findings
that in many cell types G3BPs are uniquely required for SG formation and illustrated the
difficulties in genetically disentangling the functions of SGs from those of their required G3BP
components.[2]
238
SGs are Enriched for mRNAs that are Favored by the ISR Translation Program
After difficulty using this genetic approach to define the role of SGs, we turned to other
approaches. One attractive hypothesis was that mRNAs that are differentially regulated by the
ISR translation program are also differentially localized to SGs and that this localization to SGs
determines the effect that G3BPs have on translation of those mRNAs during stress. To assess
the relationships between regulation by the ISR translation program, regulation by G3BPs, and
SG localization, we purified stress granule cores. Using an established protocol,[18,52] cores were
purified from HCT116 cells treated with 500 µM sodium arsenite for one hour, and both the SG-
associated and total-cytoplasmic transcripts were sequenced. Analysis of these datasets
identified 488 transcripts that were significantly enriched in SG cores and 453 that were

Figure 2. G3BPs reinforce the ISR translation program. (A) Depletion of endogenous G3BPs fused to AID. Immunoblots probed for the indicated proteins show specific depletion of AID fusion proteins after treatment with 500 μM IAA for 3 h. **(B)** Quantification of protein depletion in (A), normalizing to levels of GAPDH. Points show values for two biological replicates. **(C)** Prevention of SG formation during arsenite stress. HCT116 cells were either treated with 500 μM IAA to deplete G3BPs (right) or not treated (left) prior to arsenite stress (500 μM sodium arsenite for 1 h). Images show either G3BPs in red or SG marker protein PABPC1 in green (scale bar, 10 μm). **(D)** Quantification of SG formation in (C). Points show values for three biological replicates (***, $p \leq 0.001$; Welch's two-sample t-test). **(E, F)** Prevention of SG formation during heat shock (45°C for 25 min); otherwise, as in (C, D). **(G)** Schematic representation of G3BP-depletion experiment. Ribosome profiling was performed on stressed HCT116 cells to determine the effects of stress on translation, G3BP-depleted unstressed HCT116 cells to determine the effect of G3BPs on translation, and G3BP-depleted stressed HCT116 cells to determine the effects of G3BPs/SGs on translation. The effects of stress were then compared to the effects of G3BPs and SGs. **(H)** Relationship between G3BP-dependent translation during arsenite stress and the ISR translation

(H) Relationship between G3BP-dependent translation during arsenite stress and the ISR translation program. The plots indicate the $\log_2$ FC in TE due to presence of G3BPs in cells treated with arsenite as a function of the $\log_2$ FC in TE due to arsenite. Points for stress-enhanced, stress-resistant, and stress-hypersensitive mRNAs are indicated by colors. *n* indicates number of unique mRNAs. **(I)** Relationship between G3BP-dependent translation in the absence of stress and the ISR translation program; otherwise, as in (H). **(J)** Relationship between G3BP-dependent translation during heat shock and the heat-shock ISR translation program; otherwise, as in (H). **(K)** Relationship between G3BP-dependent translation in the absence of stress and the ISR translation program; otherwise, as in (H).

250
significantly depleted (Fig 3A, >2-fold change, adjusted *p* value <0.05).
Overall, our SG enrichments observed in HCT116 cells resembled those previously observed
in U2OS cells (Fig S3A, R_S = 0.50),[18] especially when considering the differences between the
two cell types, as reflected in their overall total-mRNA expression profiles (Fig S3B, R_S = 0.66).
Moreover, as found in U2OS cells,[18] longer mRNAs and mRNAs that were poorly translated prior
to stress tended to be enriched in SGs purified from HCT116 cells (Fig S3C,D). These similarities
with the results previously obtained with U2OS cells indicated that we had successfully purified
SG cores.
We next examined the relationship between differential regulation by the ISR translation
program and SG localization and found that mRNAs that retained or improved their translation
during stress tended to be more enriched in SGs, whereas mRNAs that were especially sensitive
to translation shutdown tended to be depleted (Fig 3B). To evaluate whether this relationship
was simply a consequence of SGs being enriched for transcripts that were poorly translated
prior to stress, which might have experienced a smaller decline upon translation shutdown, we
also examined the relationship between TE during stress and SG enrichment and found that SGs
were enriched for mRNAs that were translated relatively well during stress (Fig S3E).
Furthermore, analyses of published measurements of SG-enrichment in U2OS cells treated with
sodium arsenite[18], stress-dependent RNA-granule enrichment in NIH 3T3 cells treated with the
potent ER stressor thapsigargin [53], and stress-dependent G3BP proximity labeling in HEK293T
cells treated with sodium arsenite[54] yielded similar, albeit sometimes weaker, trends (Fig S3F –
J). Together, these results supported a model in which mRNAs that are favored by the ISR
translation response preferentially localize to SGs and suggest that this model might be
generalizable across multiple stresses and cell types.
274
SG-Enriched mRNAs are Favored by G3BPs During Stress
To assess whether SG localization might influence regulation by G3BPs during stress, we
examined the relationship between G3BP dependence of translation during arsenite treatment
and SG enrichment. Surprisingly, mRNAs enriched in SGs of our HCT116 cells tended to have
higher TE in the presence of G3BPs during stress, whereas mRNAs depleted from SGs tended to
be translationally repressed in the presence of G3BPs during stress (Fig 3C). Repeating this
analysis with SG enrichment data from arsenite-treated U2OS cells[18] yielded similar but weaker
relationships (Fig S3K). Likewise, an analysis comparing our G3BP-dependent translation during
heat shock to previous measurements of stress-dependent RNA-granule enrichment in heat-
shocked NIH 3T3 cells[53] showed that mRNAs targeted to the SG-enriched fraction tended to
have higher TEs in the presence of G3BPs (Fig S3L). Moreover, comparison of our G3BP-

dependent translation during arsenite stress to SG-enrichment in arsenite-stressed HEK293T
cells previously measured by proximity labeling[54] yielded analogous results (Fig S3M). Thus, in a
variety of stress and cell-type contexts, the G3BPs favor the translation of SG-enriched mRNAs
or inhibit the translation of SG-depleted mRNAs. Taken together, the results of our SG-
purification experiments suggested a model in which those transcripts that are most strongly
upregulated by the ISR translation program localize to SGs, leading to further upregulation by
G3BPs.

Figure 3. Translation of SG-enriched mRNAs is favored by G3BPs. (A) mRNAs enriched and depleted in SGs. Plotted for each mRNA is its abundance in the SG fraction as a function of its abundance in the cytoplasmic fraction. Points for SG-enriched and SG-depleted mRNAs, as determined by DESeq2 ($\log_2$ FC > 1, adjusted p value < 0.05), are indicated in red and blue, respectively, with their numbers in parentheses. n indicates total number of unique mRNAs. **(B)** Relationship between SG enrichment and ISR translation. Plotted are the SG enrichments of mRNAs in each of three ISR translation categories: stress-enhanced/resistant, bulk behavior, and stress-hypersensitive (line, median; notch, 95% confidence interval; box, quartiles; whiskers, 1.5 x IQR); ***, $p \leq 0.001$; Welch's two-sample t-test). **(C)** Relationship between G3BP-dependent translation and SG enrichment. The plots indicate the $\log_2$ FC in TE due to presence of G3BPs in stressed cells for mRNAs in each of three SG-enrichment categories: SG-enriched, SG-depleted, and neither; otherwise as in (B). All stresses were 500 µM sodium arsenite for 1 h.

SG-like OptoGranules Drive Global Reduction in Translation without a Full ISR

294
Our global analyses of the effects of G3BPs on translation showed that G3BPs, while not
required to establish the ISR translation program, do reinforce it. To determine whether
formation of SGs was responsible for these additional changes to translation during the ISR and
whether SGs were indeed sufficient to drive these translation changes in the absence of
exogenous stress, we set out to find a system in which we could form ectopic SGs and assess
their effect on translation in the absence of exogenous stress.
Ubiquitin-associated protein 2-like (UBAP2L) is reported to be sufficient to drive ectopic SG
formation in the absence of exogenous stress.[55] Accordingly, in an attempt to nucleate ectopic
SG formation, we used the PiggyBac system to integrate multiple copies of doxycycline-

inducible, RFP-tagged UBAP2L (RFP::UBAP2L) into HCT116 cells. As expected, doxycycline
induction caused UBAP2L puncta to form in most cells. However, these puncta were smaller and
more numerous than typical SGs, and did not colocalize with the canonical SG marker eIF4G,
which indicated that they were not authentic SGs (Fig S4A,B).
Overexpression of G3BP1 is also reported to be sufficient to drive ectopic SG formation in
the absence of exogenous stress.[31] When we overexpressed GFP-tagged G3BP1 using the
PiggyBac system, G3BP puncta formed with similar size and morphology as stress granules.
Moreover, these puncta were positive for the canonical SG marker, eIF4G, as expected if these
puncta were authentic SGs. However, these SGs formed in only ~20% of cells overexpressing
G3BP1 (Fig S4C,D).
To form ectopic SGs in a higher fraction of cells, we turned to the CRY2 optogenetic system,
in which the light-dependent cryptochrome 2 (CRY2) photolyase homology region (PHR)
oligomerization domain (hereafter referred to simply as CRY2) is fused to G3BP1, enabling
spontaneous SG-like condensates to be nucleated in response to blue light.[32] These
OptoGranules contain many known markers for authentic SGs.[32] In our adaptation of this
method, we removed the NTF2L domain at the N-terminus of G3BP1 and replaced it with the
CRY2 domain tagged with GFP (GFP::CRY2::G3BP$_{\Delta N}$) (Fig 4A). This construct was placed under a
doxycycline-inducible promoter and integrated into cells using the PiggyBac system. Even in the
absence of blue light, inducing the expression of GFP::CRY2::G3BP$_{\Delta N}$ caused formation of
ectopic puncta in ~20% of cells, which resembled the percentage observed when
overexpressing GFP::G3BP1. Nevertheless, exposing these cells to 488 nm blue light from an
LED array enhanced puncta formation, such that they were observed in ~80% of cells within 3 h.
Removal of the blue light caused disassembly of puncta, reaching the basal 20% of cells within 2
h, which showed that for most, if not all, of the granules, formation was reversible (Fig 4A, S4E).
We next examined whether these granules had SG markers and behavior. Consistent with
previous findings,[32] our light-induced granules were positive for canonical SG markers eIF4G and
CAPRIN1 (Fig 4B and S4F). FRAP experiments showed that these granules recovered from
photobleaching on a time scale similar to that of canonical SGs, indicating that these granules
shared similar dynamics (Fig S4G). Together, these results showed that our OptoGranules (OGs)
were dynamic, reversible, SG-like condensates formed in the absence of exogenous stress.
To determine whether formation of these OGs was sufficient to drive translation changes
even in the absence of exogenous stress, cells expressing GFP::CRY2::G3BP$_{\Delta N}$ were exposed to
blue light for three hours, allowing OGs to form in most cells, and then subjected to ribosome
profiling using our ribo-spike. To control for the effects of exposing cells to blue light, ribosome
profiling was also performed on cells that were exposed to blue light but were not induced to
express GFP::CRY2::G3BP$_{\Delta N}$ and therefore formed no OGs.
If formation of OGs were sufficient to drive ISR-like translation changes, then we would
predict that analysis of our ribosome-profiling data would yield the following results. First, as
with the ISR translation program, cells with OGs would be predicted to undergo a global
reduction in TE compared to cells only exposed to blue light. Second, these translation changes
would be predicted to positively correlate with those observed in the ISR, with similar genes
retaining their translation in the face of widespread decreases in TE, and with similar genes
being especially sensitive to that translation shutdown. Third, OG-induced translation changes
would be predicted to correlate positively with the G3BP-dependent translation during stress.

Fourth, transcripts enriched in SGs would be predicted to also be resistant to any translation
shutdown caused by the formation of these SG-like condensates.
When TEs in OG-induced cells were compared to those in blue-light controls, a global
reduction in translation was observed. This reduction was modest (1.6-fold) compared to the
drops in translation caused by the ISR translation program (20-fold for 1 h of 500 μM sodium
arsenite stress and 8-fold for 25 min of 45°C heat shock) (Fig 4C). This modest reduction was
consistent with the results of the AID-depletion experiments that indicted that G3BPs, which are
essential for SG formation, were responsible for only a small fraction of the translation
shutdown observed during the ISR.
Although this observation was consistent with the first of our four predictions, analyses of
the TE results failed to confirm any of the other three predictions regarding the effects of OG
induction on translation. OG translation changes did not correlate with either the ISR translation
program or G3BP-dependent translation changes in stressed cells (Fig 4D,E, R_S = 0.019 and −
0.054, respectively). Additionally, although mRNAs depleted in SGs tended to be more sensitive
to the translation shutdown caused by formation of OGs, mRNAs that were enriched in SGs did
not perform any better than those that were neither enriched nor depleted (Fig 4F). Considered
together, these results argued that spontaneous formation of the SG-like OGs, while sufficient
to drive a modest reduction in global translation, was not sufficient to specify the ISR translation
program.
367
OptoGranules Require Cellular Stress to Recruit an SG Transcriptome
One caveat of our predictions of TE changes that might be observed upon OG formation was
that we assumed that the spontaneously formed OGs contained the same transcriptome as SGs
and therefore would have the opportunity to regulate those transcripts in the same manner.
This assumption is consistent with the recent proposal that G3BP is the central scaffold of a
complex protein-RNA interaction network. According to this model, G3BPs nucleate SG
formation by switching to a higher valency conformation when triggered by increased levels of
free cytosolic RNA.[2-4] This model suggests that simply increasing the valency of G3BPs, as in the
case of the CRY2 system, might be sufficient to establish that same complex interaction
network, even in the absence of stress. Interestingly, this prediction is also consistent with the
observation that the established determinants of the SG transcriptome (transcript length and TE
prior to stress) are parameters that are independent of stress and is further supported by the
observation that OGs are positive for many canonical SG markers (Fig S4E).[18,32] However,
prominent paradigms of SG formation state that the SG transcriptome is largely determined by
the transcriptome RNA-binding landscape (the landscape created by proteins bound to the
transcriptome), meaning that mRNAs that are most available to self-associate or be stably
bound by SG assembly factors and remodeled into messenger ribonucleoprotein (mRNP)
complexes are the most likely to be incorporated into SGs.[2,14,56,57] Indeed, stress can
dramatically shift the RNA content of other biological condensates such as P-bodies and P
granules.[58,59] Thus, because cellular stress dramatically remodels the RNA-binding landscape,
SGs formed in the presence of stress and OGs formed in the absence of stress might recruit
different sets of mRNAs, despite their general compositional, morphological, and biophysical
similarities.
391

Figure 4. SG-like OGs Can Drive a Global Reduction in Translation. (A) Schematic representation of the OG system. HCT116 cells expressing endogenous G3BP containing an NTF2L dimerization domain were edited to also express doxycycline (dox)-inducible ectopic G3BP in which the NTF2L domain had been replaced with a fusion of GFP and the light-inducible CRY2 dimerization domain (GFP::CRY2::G3BP$_{\Delta N}$). After doxycycline induction, a minority of cells formed ectopic granules. After exposure to blue light (488 nm), most cells formed ectopic granules. **(B)** SG marker localization to OGs. HCT116 cells expressing wild-type GFP::G3BP1 were either stressed with 500 μM sodium arsenite for 1 h (top) or treated with 1 μM doxycycline to induce expression of GFP::CRY2::G3BP$_{\Delta N}$ and exposed to blue light for 3 h (bottom). Fluorescent or immunostained proteins are as indicated (scale bars, 10 μm). White boxes highlight insets that are expanded on the right.

(C) Comparison of TE in OG-induced cells to that in blue-light control cells. Plotted are the TE distributions in HCT116 cells in which GFP::CRY2::G3BP$_{\Delta N}$ was expressed and exposed to blue light to form OGs (right) and in control cells expressing only wild-type G3BP exposed to blue light, in which no OGs were formed (left) (line, median; notch, 95% confidence interval; box, quartiles; whiskers, 1.5 x IQR). Significance was determined using ribo-spike values as in Figure S1B (*, $p \leq 0.05$; Welch's two-sample t-test). **(D)** Comparison of OG and ISR translation programs. Plotted is TE change caused by induction of OGs as a function of TE change caused by arsenite. *n* indicates total number of unique mRNAs. **(E)** Comparison of OG and G3BP-dependent translation programs. Plotted is TE change caused by OGs as a function of TE change caused by presence of G3BPs in arsenite-treated cells; otherwise, as in (C). **(F)** Comparison of OG translation program and SG enrichment. The plots indicate the log$_2$ FC in TE caused by induction of OGs for mRNAs in each of three SG enrichment categories: SG-enriched, SG-depleted, and neither (line, median; notch, 95% confidence interval; box, quartiles; whiskers, 1.5 x IQR); ns, not significant; ***, $p \leq 0.001$; Welch's two-sample t-test).

To test this hypothesis, we set out to purify OGs formed in either the absence or the presence
of stress. To purify OGs formed in the absence of stress, we used the same approach as in our
previous ribosome-profiling experiment. To purify OGs formed in the presence of stress, we
combined our OG and AID systems. We integrated the doxycycline-inducible GFP::CRY2::G3BP$_{\Delta N}$
construct into cells expressing endogenously edited AID::RFP::G3BPs and doxycycline-inducible
OsTIR1, the E3 ligase responsible for the turnover of the AID. Upon treatment with doxycycline
and IAA, these cells degraded their endogenous G3BPs and replaced them with
GFP::CRY2::G3BP$_{\Delta N}$, which rendered them incapable of forming SGs in the absence of blue light,
even when treated with 500 µM sodium arsenite for 1 h. However, when these stressed cells
were exposed to blue light, they readily formed granules (Fig 5A,B). We refer to these OGs
formed in the presence of stress as StressOptoGranules (SOGs).
We then purified and sequenced the transcripts associated with both OGs and SOGs. As
controls, we also examined real SGs purified from each cell line, i.e., from the OG line subjected
to stress in the absence of blue light, and from the SOG line subjected to stress in the absence
of both IAA and blue light (Fig 5C,D). We then examined all pairwise comparisons of granule
enrichments, including our original SG enrichment data acquired from the parental line as an
additional control. When we compared across control SGs we saw that results for all of our
controls were similar to each other, with R_S values ranging from 0.72–0.75 (Fig 5E, Fig 5SA).
However, when comparing results of control SGs to those of OGs formed in the absence of
stress, substantially weaker correlation was observed, with R_S values ranging from 0.35–0.45
(Fig 5E, Fig 5SA). To determine if this difference was due to lack of cellular stress as opposed to
some intrinsic bias of OG formation, we compared results of our control SGs to those of SOGs
formed in the presence of stress. We found that transcripts enriched in SOGs were much more
similar, and in fact nearly indistinguishable from those in SG controls, with R_S values ranging
from 0.69–0.70 (Fig 5E, Fig 5SA). These differences in granule composition were not driven by
differences in gene expression, as the cytoplasmic mRNA levels correlated quite well between
these groups, with R_S values ranging from 0.84–0.96 (Fig 5F, Fig 5SB). Taken together, our results
indicated that establishing the SG transcriptome requires not only the interaction network
established by high valency G3BPs but also the RNA-binding landscape established during
cellular stress.
Equipped with data on the enrichments within OGs, we were able to re-evaluate our
prediction that granule-enriched transcripts would have favored translation during OG

formation. As expected, transcripts enriched in OGs better retained their translation, whereas
transcripts that were depleted from OGs were most sensitive to the OG-induced translation
shutdown (Fig 5G). This observation was consistent with a model in which enrichment in SGs
favors translation, while depletion from SGs favors repression. Taken together, our OG
ribosome-profiling and purification results supported a model in which SGs are sufficient to
drive a modest decrease in global translation and to prioritize the translation of their enriched
mRNAs while downregulating translation of mRNAs for which they are depleted. However, SGs
require stress to induce an ISR-like translation program, presumably because the sorting of SG
mRNAs is dictated by the cellular RNA-binding landscape, which is drastically remodeled upon
activation of the ISR.
435
Tethering to SGs Imparts Resistance to ISR Translation Shutdown
Our findings suggested that localization to an SG grants an mRNA prioritized translation during
the ISR translation shutdown, allowing the mRNA to better retain its TE. To test this model, we
examined whether tethering reporters to SGs influenced their translation. The reporters each
encoded nanoluciferase (nLuc) fused to the *E. coli* dihydrofolate reductase (ecDHFR)
destabilizing domain, which causes rapid turnover of its protein fusions, ensuring reporting on
recently translated nLuc.[60] The 3'UTR of each reporter mRNA also included an array of 24
bacteriophage MS2 hairpins, which bind MS2 coat protein (MCP).[30,61]
G3BPs strongly partition into SGs[1,31,62], and we found that in HCT116 cells almost half of our
endogenous G3BP1 fusion protein (45.9%) was inside of SGs after 90 minutes of arsenite stress,
a fraction that exceeded estimates in other cell lines (Fig S6A,B).[62] Considering this high
partitioning coefficient, we chose endogenous G3BP1 as our SG-tethering protein. Using the
PiggyBac system, reporters were stably integrated into HCT116 cells in which all endogenous
G3BP1 alleles were tagged with either GFP (GFP::G3BP1) or MCP and GFP (MCP::GFP::G3BP1)
(Fig 6A). We also explored the λN–BoxB aptamer system for tethering reporters to G3BP,[63,64] but
in contrast to our results when expressing MCP::GFP::G3BP (Fig S6C), we observed perturbed
G3BP localization and SG formation in cells expressing λN::GFP::G3BP.
To assess the effects of SG tethering for a variety of 5' and 3' UTRs, we created four different
reporters using 5' and 3' UTRs from four mRNAs that had diverse behaviors in response to stress
and to G3BPs, as measured by ribosome profiling (Table S1). Single-molecule fluorescence in
situ hybridization (smFISH) with probes targeting the *nLuc* sequence indicated that the reporters
were recruited to SGs at a basal level in cells expressing only GFP::G3BP, with an average of 30%
of molecules localized to SGs across the four reporters (Fig 6B – D). In cells expressing
MCP::GFP::G3BP, the reporters were more robustly recruited to SGs, with an average of 85% of
reporter molecules localized to SGs (Fig 6B – D), which was a higher fraction recruited to SGs
than that observed for 99% of endogenous mRNAs.[18]
Next, we measured how the translation of these reporters responded to arsenite stress.
Cells were treated with 500 µM sodium arsenite for 90 min, and changes to protein production
were monitored by measuring luciferase signal from these cells compared to that from
unstressed control cells, and normalizing to total-protein levels. In addition, changes to mRNA
levels were monitored by reverse transcription and quantitative PCR (RT-qPCR). Using these
changes in translation and mRNA, changes in TE were calculated for each reporter. For
untethered reporters, large decreases in TE were observed upon induction of stress, with a 25-

Figure 5. OGs require stress to recruit the SG transcriptome. (A) Replacement of G3BP with GFP::CRY2::G3BP$_{\Delta N}$ to induce SOGs. In the left panel, HCT116 cells expressing endogenously tagged AID::RFP::G3BPs were either untreated (left) or treated with 1 μM doxycycline and 500 μM IAA for 18 h (right), leading to turnover of endogenous G3BP shown in red and expression of GFP::CRY2::G3BP$_{\Delta N}$ shown in green. In the middle panel, cells were treated as in left panel, but were subsequently stressed with 500 μM sodium arsenite for 1 h in the absence of blue light. In the right panel, cells were treated as in the middle panel, but were also exposed to blue light, beginning 30 min after start of the arsenite treatment (scale bars, 10 μm). White boxes highlight insets that are expanded on the right. **(B)** Quantification of granule formation in (A). Error bars report standard deviation for three biological replicates (*** $p \leq 0.001$, ns, not significant, Welch's two-sample t-test). **(C)** Schematic representation of granule purification experiment. Three cell lines were used: the parental HCT116 line expressing endogenous G3BPs, an OG HCT116 line expressing endogenous G3BPs and doxycycline-inducible GFP::CRY2::G3BP$_{\Delta N}$, and an SOG line expressing endogenous G3BP fused to AID and RFP (RFP::AID::G3BP), doxycycline-inducible OsTIR1, and doxycycline-inducible GFP::CRY2::G3BP$_{\Delta N}$. Control SGs were purified from each cell line by treating them with 500 μM sodium arsenite for 1 h. Additionally, OGs were purified from the OG line treated with doxycycline for 18 h, followed by 3 h of blue light, and SOGs were purified from the SOG line treated with both doxycycline and IAA for 18 h followed by exposure to arsenite stress for 30 min in the absence of blue light and then an additional 30 min of stress in the presence of blue light. **(D)** mRNAs enriched and depleted in the indicated types of granules; otherwise, as in Fig. 3A. **(E)** Comparisons of granule enrichments. The heatmap depicts the pairwise correlations (R_S values) observed between the enrichments of granules from (Fig. 3A and 5D). **(F)** Comparisons of cytoplasmic transcriptomes of cells used to determine granule enrichments (E). **(G)** Comparison of OG-dependent translation and OG enrichment. Plotted are distributions of TE changes caused by induction of OGs for mRNAs in each of three OG enrichment categories; otherwise, as in Fig 4E.

fold average drop in TE observed across the four reporters (Fig 6E,F). This drop was consistent
with that expected for the translation of an mRNA during the ISR. Indeed, it aligned closely with
the 20-fold average decrease in TE observed by ribosome profiling, indicating that our assay
accurately reported on typical translation changes observed during stress. However, this
uniform drop in translation also indicated that several of our reporters did not behave like their
endogenous counterparts, some of which continue to be efficiently translated under stress
conditions (Table S1), which implied that the 5' and 3' UTRs of these mRNAs were not sufficient
to impart their endogenous translation responses to stress.
We proceeded to examine what happens to the translation of these reporters when
tethered to SGs in the MCP::GFP::G3BP1 cell line. When tethered to SGs, each of our four
reporters retained its translation significantly better than when untethered, with an average
drop in TE of 2-fold when tethered, compared to 25-fold when untethered (Fig 6E,F). To control
for the possibility that this might have been the result of recruiting a generic protein to the 3'
UTRs of the reporters, we repeated the experiment in the untethered GFP::G3BP cell lines but
this time also expressing MCP::BFP at levels similar to those of MCP::GFP::G3BP (Fig 6E,F, Fig
S6E). The effects of MCP::BFP co-expression were minor compared to those of MCP::GFP::G3BP
(Fig 6E,F).
To determine whether this tethering effect was a stress-specific, we examined the effect in
unstressed cells and stressed cells separately. Although all reporters showed effects of tethering
in arsenite-treated cells, no substantial differences between tethered and untethered controls
were observed in unstressed cells (Fig S6F,G), arguing that tethering to SGs via G3BP reinforces
translation in a stress-dependent manner and consistent with a model in which this is due to SG
recruitment.
To further distinguish between SG and G3BP function, we examined whether the effect of
tethering was specific to G3BPs. To do this, we measured the TEs of reporters tethered to SGs
expressing MCP fused to CAPRIN1 (MCP::RFP::CAPRIN1). Although we were able to recover cell

Figure 6. Tethering to SGs via G3BP1 imparts resistance to ISR translation shutdown. (A) Schematic representation of SG tethering experiment. The 5' and 3' UTRs of an mRNA of interest were appended to a nanoluciferase reporter (NanoLuc) that contained 24x repeats of the MS2 hairpin. The reporter was then expressed in each of three different HCT116-derived cell lines: a line expressing endogenous G3BP1 tagged with GFP (GFP::G3BP), a line expressing endogenous G3BP1 tagged with the MS2 coat protein (MCP) fused to GFP (MCP::GFP::G3BP), and a line expressing endogenous G3BP1 tagged with GFP and expressing MCP fused to ectopic BFP (MCP::BFP). **(B)** Tethering a luciferase reporter to SGs via G3BP1. HCT116 cells expressing a NanoLuc reporter bearing 5' and 3' UTRs from *CNOT10* and endogenous G3BP1 tagged with either GFP (top) or MCP::GFP (bottom) were stressed with 500 μM sodium arsenite for 90 min. Images show localization of G3BP in green and smFISH of NanoLuc reporter molecules in magenta (scale bars, 1 μm). White boxes highlight insets that are expanded on the right. **(C)** SG tethering efficiency. Percent of cytoplasmic reporter molecules localized to SGs in untethered cells (green) and to SGs in tethered cells (pink) are shown for reporters with 5' and 3' UTRs of the indicated genes. Points show values for at least three biological replicates (** $p \leq 0.01$, *** $p \leq 0.001$, Welch's two-sample t-test). The average SG tethering efficiency for the four different reporters was 33 for untethered cells and 86 for tethered cells, with standard deviations of 12% and 9%, respectively.

(D) G3BP1-tethered ISR translation response. Plotted for each reporter is the change in TE caused by arsenite stress (500 µM sodium arsenite for 90 min) in untethered cells (green), untethered cells expressing MCP::BFP (blue), or tethered cells (pink) (ns, not significant). The average $\log_2$ FC in TE for the ISR translation response was –5.05 for untethered cells, –4.15 for MCP::BFP expressing cells, and –0.98 for tethered cells; otherwise, as in (C).

using another MCP-tagged SG protein. Accordingly, reporters were integrated into a cell line
lines for only three of the four reporters, for all three, recruitment to SGs robust, resembling
that observed with G3BP tethering (Fig S6H,I). To measure the effects of this alternative
tethering strategy, we expressed both the reporters and doxycycline-inducible, MCP tagged
CAPRIN1 in our AID::G3BP cell line. With this combination, we could assess the effects of
tethering the reporter to SGs via CAPRIN1 and test whether those affects were dependent on
the presence of G3BPs and SGs using the AID degron system.
To induce MCP-tagged CAPRIN1 without depleting the AID-tagged G3BPs, we treated cells with
both 1 µg/ml doxycycline and 200 µM auxinole, a small-molecule inhibitor of OsTIR1 activity.
Robust expression of MCP-tagged CAPRIN1 was observed, with minimal inhibition of SG
formation due to G3BP depletion (Fig S6J – L). Under these conditions, our reporters, tethered
to SGs via CAPRIN1, significantly retained translation following arsenite, compared to
uninduced, untethered cells, which showed that this effect was not specific to tethering via
G3BPs. We note that the effect size observed with CAPRIN1 tethering was smaller than that
observed which G3BP tethering, which might be attributable to the smaller translation
inhibition observed upon stress, potentially due to differences in cell line background.
Furthermore, when cells were instead treated with both 1 µg/ml doxycycline and 500 µM IAA to
degrade G3BPs and prevent SG formation, this retention of translation was abolished (Fig S6M),
which indicated that this effect of tethering required the presence of G3BPs. Taken together, the
results of these tethering experiments indicated that tethering the 3'UTR of an mRNA to either
G3BP1 or CAPRIN1 caused increased recruitment to SGs and imparted resistance to the global
reduction in translation due to the ISR—consistent with a model in which localization to SGs
grants an mRNA prioritized translation during stress.
521
522   **DISCUSSION**
523   SGs were once thought to be required for the global repression of translation observed during
524   the ISR.[5,15,20,24,25] This idea fell out of favor as many studies showed that cells incapable of
525   forming SGs still mount an ISR translation response.[1,5,26] Since then, additional evidence has led
526
to the hypothesis that SGs, far from being required for the ISR translation response, have no
function and are the unavoidable, incidental byproducts of increased RNA availability due to
translation repression.[34,35] Here, we propose a model in which SGs are neither required for the
substantial changes in translation induced during the ISR nor incidental condensates that have
no impact on the stress translation response, but instead measurably amplify and reinforce the
ISR translation program, thereby leading to a widespread and statistically significant, albeit
relatively modest, impact on global translation.
This widespread reinforcement of the ISR is consistent with several observations of SG
biology. SGs recruit a wide array of mRNAs and proteins, constituting a significant redistribution

of the cytoplasm during stress.[14,17,18] At the same time, SGs contain only ~10% of the
cytoplasmic mRNAs and protein molecules,[18,53] which implies that they would be less likely to
have dramatic impacts on cellular outcomes from stress and more likely to play more subtle
tuning roles on expression of many different genes.
Indeed, a recent study investigating the impact of SGs during viral infection reports that,
although SGs are not required for the innate immune response to viral infection or double-
stranded RNA (dsRNA), they play a role in tuning that response.[51] However, instead of
amplifying the stress response as we saw during arsenite and heat stress, SGs are reported to
act as shock absorbers in response to viral infection, limiting the extent of the ISR transcriptional
response.[51] The difference in orientation of these potential tuning functions might be explained
by the difference in the nature of their stressors and composition of their SGs. Different stresses
lead not only to different stress responses but also to distinct stress granules with distinct
compositions.[19,39] Thus, viral infection or treatment with dsRNA may result in SGs with very
different composition and biology. Indeed, the granules formed in response to dsRNA tend to be
smaller than those formed by arsenite or heat shock and tend to only weakly recruit PABPC1
and polyadenylated mRNAs, with some of the RNAs most enriched in canonical SGs such as
AHNAK and NORAD not present in dsRNA SGs.[65] Furthermore, whereas canonical SGs require
the phosphorylation of eIF2α, granules formed in response to dsRNA do not require the dsRNA-
dependent eIF2α kinase PKR for their formation.[51,65] Moreover, the response to viral infection
might differ between different cells or viruses, in that the shock-absorber function is not
reported to apply in other conditions designed to induce viral stress.[66] All of this is further
complicated by the presence of RNAse L bodies (RLBs) which also form in response to viral
infection and share markers with SGs, such as G3BPs.[67] Thus, although our results seem likely to
be generalizable across canonical SGs that require eIF2α phosphorylation, granules formed
under viral infection appear to have different effects.
In presenting a model of SGs functioning to reinforce the ISR translation program, we
propose that, in addition to repressing translation of some mRNAs, SGs also help to rescue
translation of others. SGs have previously been hypothesized to play strictly repressive roles in
translation for a number of reasons. First, SGs form in conditions of translation repression and
are nucleated by the availability of translationally silent mRNAs, implying that the mRNAs
localized to them are not translating.[25,68] Second, although SGs are enriched for small ribosomal
subunits and eukaryotic initiation factors, they are not enriched for large ribosomal
subunits.[20,28] However, neither of these observations preclude SGs from imparting pro-
translation effects. First, the mRNAs in SGs are in dynamic exchange with the cytosol and
therefore could transit out to the cytosol after their translation was initiated, which would
maintain enrichment for translationally silent mRNAs inside the SGs.[29,30,38] Indeed, translation
initiation inside the SG might even lead to expulsion of the mRNA, as the presence of even a
single ribosome can prevent an mRNA from localizing to an SG.[69] Second, although 40S and 60S
ribosomal subunits in SGs are at different stoichiometries than they are in the cytosol, 60S
subunits are not depleted from SGs; their availability is equal to that in the cytosol.[20,28] Indeed,
a recent study following the translation of *ATF4* reporter molecules during stress reports that
translation in the SG is neither impossible nor a rare event.[30] Thus, these prior observations
leave open the possibility that SGs could be sites that lead to privileged translation for a subset
of mRNAs.

Although this model of mRNAs being preferentially initiated or translated while localized to
the SG is perhaps the simplest interpretation of our data, it is not the only interpretation.
Another possibility is that while transiting through an SG, mRNAs are licensed to be more
efficiently translated, and then, having acquired this licensing, they go on to be translated
independent of their localization in the SG or cytosol (Fig 7). Elaborating on this possibility,
perhaps an mRNA is translated at the same efficiency inside or outside of the SG, but in either
case its TE might be proportional to its dwell time inside of the SG and thus proportional to its
likelihood of acquiring the license to be more efficiently translated.

This license to translate could be acquired through the binding of either translation
machinery or a set of pro-translation RNA-binding proteins. For example, loading of a
preinitiation complex (PIC) assembled from the 40S ribosome subunit and initiation factors,
which are known to be enriched in SGs,[20,28] could poise a transcript to be more quickly initiated
whether in the SG or in the cytosol.[5,14] Similarly, UBAP2L, which enhances the translation of its
associated mRNAs, is highly enriched in SGs.[50,70] Perhaps preferential binding of UBAP2L to an
SG mRNA could lead to more efficient translation even after the mRNA has diffused out of the
SG. Additional interesting candidates for translational licensing are the G3BPs themselves. The
G3BPs have been observed to regulate translation and translation machinery.[1,21-23,71,72] G3BPs
also exhibit exceptionally high partitioning into SGs, with almost half of G3BP1 localized to SGs
in our cell lines.[63] If G3BPs act to license privileged translation, their strong partitioning to SGs
might lead to preferential translation of SG-resident transcripts (Fig 7).

This proposed increased mRNA licensing in SGs is consistent with the idea that SGs are
formed by non-translating RNAs and with the observation that, although an *ATF4* reporter is
translated in the SG, it is translated no more efficiently in the SG than in the cytosol.[30] Indeed,
we found that *ATF4* mRNA was neither enriched in SGs nor influenced by the presence of SGs,
which indicated that other transcripts might better represent mRNAs influenced by SGs.

Lastly, it is important to acknowledge the difficulty of distinguishing the function of SGs from
the function of the SG-nucleating G3BPs. One approach for inferring the function of SGs is to
consider phenotypes consistently observed after different proteins reported to be required for
SG formation are depleted. However, many of these proteins, such as the G3BPs, CAPRIN1, and
UBAP2L, are thought to form complexes with each other even outside of the context of SGs,
meaning that any phenotypes shared between these interacting proteins could be due to the
function of complexes that exist independent of SGs.[19,34,35] Moreover, we and others found that
the ability to prevent SG formation through depletion of non-G3BP nucleators appears to be
stress or cell-type specific, which limits what can be learned regarding general SG function by
depleting these factors.

Although we have not conclusively distinguished between G3BP function and SG function,
five lines of reasoning support the proposal that the ISR translation program is reinforced by SGs
rather than by only G3BPs. First, in both our ribosome profiling experiments and our tethered
reporter experiments, we found that G3BPs regulate these mRNAs only in the context of stress.
Second, although our SG purifications did rely on the IP of G3BPs, this IP was performed
specifically on the cellular fraction that pellets after centrifugation at 18,000*g*, which removes
most cytosolic G3BP.[17,18,52,73] Thus, our SG transcriptomes should be substantially enriched for
the content of G3BP-positive SGs and not simply the G3BP-RNA interactome. Furthermore, we
were able to identify similar trends in data from SG-enriched fractions that do not require IP of

G3BPs.[53] Third, we found that formation of OptoGranules, which manipulates only localization
of G3BPs, was sufficient to trigger a translation response, and that the content of the
OptoGranule was what determined which mRNAs were regulated. Fourth, we found that
tethering reporter mRNAs to SGs was sufficient to impart resistance to stress induced
translation shut down, whether tethered by G3BP1 or CAPRIN1. Although G3BPs and CAPRIN1
are thought to be binding partners[1], this result indicates that this effect does not require direct
recruitment via G3BPs. Fifth, in distinguishing between SG- and G3BP-centric models, we should
not dismiss the possibility that both G3BPs and SGs contribute to this translation reinforcement.
If G3BPs are a licensing factor for improved translation during stress, and are also strongly
partitioned to SGs, perhaps the effects we observe are direct effects of G3BP binding to mRNA,
and that recruitment to SGs is a molecular strategy to increase interaction with G3BPs during
stress, therefor reinforcing translation of a subset of privileged mRNAs. Although these
considerations do not conclusively distinguish between SG and G3BP function, they do support
the proposal that SGs are a functional condensate responsible for reinforcing the ISR translation
program.

Figure 7. A Plausible Model of SG Translation Control. This model reconciles the previous observation that mRNAs devoid of ribosomes are preferentially recruited to SGs with our observation that mRNAs that are translationally enhanced during the ISR are enriched in SGs, further enhancing their translation. In this model, 1) ISR-enhanced transcripts are preferentially recruited to SGs between cycles of translation, when they lack ribosomes. This preferential recruitment might be the result of various factors, including transcript length, presence of relevant RBP-binding motifs, or translation state prior to stress. 2) As a consequence of being recruited to SGs, these transcripts are licensed for translation. One possible mechanism of licensing (depicted here) is the recruitment of G3BPs, which are highly enriched in SGs. 3) After this licensing, this model is agnostic to the location of the translation that follows. The transcript might diffuse out to the cytosol, or could remain inside the SG. Once these transcripts complete their translation, they would be available to, once again, be preferentially recruited to (or retained in) the granule.

Methods

Cell culture

All cells were cultured at 37°C with 5% CO_2 using McCoy's 5A medium supplemented with 10% fetal bovine serum (FBS) and 2mM L-glutamine. HCT116 (CCL-247 from ATCC) were obtained from the Young lab.

Cell-line construction

Cas9 mediated editing. Fusion proteins were introduced into the endogenous locus of genes using the Cas9 genome editing system. The 5' and 3' homology arms (each 200 – 700 bp) flanking an insertion site were either amplified from genomic DNA of the parental cell line or from gblock gene fragments (IDT) cloned into a rescue template plasmid, flanking the fusion protein sequence. This plasmid, along with pX330 plasmid expressing a Cas9 guide RNA, was reverse transfected into the parental cell line using lipofectamine 3000 transfection reagent in a 6-well dish at 100% confluency. After 48 h, these cells of each well were expanded to a 10 cm dish and cultured for an additional 24 h. Cells with strong signal from the fusion protein fluorescence were single-cell sorted into 96-well plates using flow cytometry. Colonies derived from single cells were grown, expanded, and screened by PCR genotyping and western blotting. Unless otherwise stated, all endogenously edited cell lines were edited homozygously.

PiggyBac editing. Exogenous genes were introduced into cells using the PiggyBac transposon system. The exogenous gene was either amplified from genomic DNA of the parental cell line or from gblock gene fragments (IDT) and cloned into a plasmid expressing the PiggyBac inverted-repeat sequences along with a selection marker for either hygromycin, puromycin, or blasticidin. This plasmid, along with the super PiggyBac transposase expression vector (System Biosciences #PB210PA-1), was reverse transfected into the parental cell line using lipofectamine 3000 transfection reagent (Thermo #L3000015) in a 6-well dish at 100% confluency. After 48 h, cells of each well were expanded to a 10 cm dish and successfully edited cells were selected for using the appropriate antibiotic. For cases in which a fluorescent protein had been transposed into the genome, cells expressing the protein at the appropriate level were single-cell sorted into 96-well plates by flow cytometry. Colonies derived from single cells were grown, expanded, and screened by either microscopy, western blotting, or both.

Ribo-spike production

In vitro transcription. Ribo-spike mRNA was generated using mMessage mMachine kit (Thermo # AM1344) following the standard T7 protocol. DNA transcription template was generated by PCR of a plasmid template encoding the firefly luciferase coding region (*fLuc*) preceded by the T7 promotor and followed by an encoded poly(A) tail of 30 nt length. The resulting PCR product was purified using phenol–chloroform extraction followed by ethanol precipitation and was size selected and purified from an agarose gel using the "freeze and squeeze" method in which the gel slice was frozen at −80°C for 30 min, followed by centrifugation at 21,000g for 5 min.[74] The eluted DNA was then concentrated by an additional ethanol precipitation and added to the mMessage mMachine in vitro transcription reaction, incubated for at 37°C for 2 h, and then treated with TURBO DNAse for 15 min at 37°C. RNA from the in vitro transcription reaction was recovered by passing the reaction over a tris-buffered micro bio-spin p30 gel column (Bio-Rad

#7326223) centrifuging at 1000*g* for 4 min. The resulting flowthrough was then purified by
phenol–chloroform extraction and ethanol precipitation, and then resuspended in water. RNA
was examined by running an aliquot on a 4% urea–polyacrylamide denaturing gel and staining
with Sybr Gold (Thermo #S11494).
In vitro translation. The *fLuc* mRNA was in vitro translated using a rabbit reticulocyte lysate
system (Promega #L4151) following the standard non-nuclease treated protocol, with the
exception of using a shorter incubation time to maximize ribosome occupancy. The reaction was
assembled by combining 0.06 µM *fLuc* mRNA with reticulocyte lysate, 0.02 mM complete amino
acid mixture, 0.8 units/µl RNAsein ribonuclease inhibitor, 0.01 M creatine phosphate, 0.05
mg/ml creatine phosphokinase, 2 mM DTT, 0.05 mg/ml tRNA, 80 mM potassium acetate, and
0.5 mM magnesium acetate in 23 117 µl reactions. The reactions were incubated at 30°C for 15
min, cooled on ice and treated with 0.1 mg/ml cycloheximide to block ribosome translocation.
Reactions were then combined, mixed, aliquoted, snap frozen in liquid nitrogen, and stored at –
80°C.
709
IAA-induced protein depletion
HCT116 cells were engineered using the PiggyBac system to express multiple copies of
doxycycline-inducible *OsTIR1*. These cells were then also engineered using Cas9 to expressed
AID-tagged fusion proteins edited at their endogenous loci. Cells expressing fusion proteins
were first doxycycline-induced for 4 h and were then treated with 500 µM IAA (Gold Bio #I-110-
25) to induce rapid (< 3 h) depletion of the AID-fusion protein. Depletion was confirmed by
microscopy, western blot, or both. When confirming by western, as was the case with G3BP1,
G3BP2, and CAPRIN1 proteins, antibodies raised directly against the target proteins were used
to enable detection of untagged alleles, if present. Where indicated, non-depleted controls
were treated with 200 µM auxinole (Aobious #AOB8812) to minimize background protein
degradation.[75] Unless otherwise stated, all other non-depleted control cells were treated only
with ethanol, the solvent used for IAA. Previous work has shown that addition of IAA alone does
not affect translation.[76]
723
724 **Stress conditions**
725 For oxidative stress, unless otherwise stated, cells were treated with media containing 500 µM
726 sodium arsenite (Sigma #93289) for 1 h. For heat shock, media was separately heated to 45°C.
727 Media was removed from cells previously grown in a 37°C, rapidly replaced with 45°C media,
728 and then cells were quickly placed in a 45°C incubator for 25 min.
729
730 **Doxycycline induction**
731 For all doxycycline inductions, cells were treated with 1 µg/ml doxycycline (Clonetech Takara Bio
732 #631311) for indicated amount of time.
733
Tethered reporter TE experiments
To measure changes in TE for the nanoluciferase reporters (NanoLuc), cells expressing a reporter
containing selected 5' and 3' UTRs, NanoLuc fused to an ecDHFR domain and 24x MS2 stem
loops, were harvested using trypsin, centrifuged at 300*g* for 5 min, aspirated, washed with cold
PBS, lysed in buffer containing 10 mM Tris HCl pH 7.5, 5 mM MgCl$_2$, 100 mM KCl, 1% triton, 1x

cOmplete mini tablet (EDTA free), and 0.02 U/μl SUPERase-In RNase inhibitor, and depleted of
nuclei by centrifuging at 1300g for 10 min at 4°C. Nuclear-depleted supernatant was then
transferred to a new tube and flash frozen using liquid nitrogen and stored at –80°C. Cells were
then assayed for total protein using bicinchoninic acid, luciferase production, and reporter
mRNA using RT-quantitative PCR (RT-qPCR) comparing to the *GAPDH* mRNA, which did not
change in abundance during treatment with sodium arsenite (Fig S6C). Change in translation
efficiency was then calculated as
746

747
$$\Delta \text{TE}_{\text{MCP/GFP}} = \frac{\Delta \text{NanoLuc signal}_{\text{MCP/GFP}}}{\Delta \text{mRNA}_{\text{MCP/GFP}}} = \frac{\dfrac{\text{NanoLuc signal/Total Protein}_{\text{MCP}}}{\text{NanoLuc signal/Total Protein}_{\text{GFP}}}}{\dfrac{\textit{NanoLuc} \text{ mRNA}/\textit{GAPDH} \text{ mRNA}_{\text{MCP}}}{\textit{NanoLuc} \text{ mRNA}/\textit{GAPDH} \text{ mRNA}_{\text{GFP}}}}$$

748
where MCP indicates the reporter being expressed in a cell line expressing MCP::BFP or
MCP::GFP::G3BP1, and GFP indicates the reporter being expressed in a cell line expressing
GFP::G3BP1.
Luciferase Assays. Luciferase assays were conducted using the standard protocol of the
Luciferase Reporter Assay System (Promega #E1910). Cell lysates were analyzed in 48-well
microplate format at room temperature using a GloMax microplate reader and luminometer
device (Promega #GM3000).
RT-qPCR. RNA was extracted using TRI reagent (sigma #93289), precipitated using ethanol and
linear acrylamide (thermo #J67830-XF) and resuspended in water. RT reactions were performed
using a Quantitect RT kit following kit protocols. Genomic DNA was first eliminated using 7x
gDNA wipeout buffer and incubating at 42°C for 2–5 min. Reactions were then placed on ice, RT
reagents were added, and then RT was performed at 42°C for 30 min. RT enzyme was heat
inactivated by incubating at 95°C for 3 min. cDNA product was then diluted 1:2 in water and
stored at –20°C. qPCR was then performed using the PowerUP SYBR Green qPCR master mix
and primers targeting either the *NanoLuc* or *GAPDH* sequences. Each sample was run in
technical triplicate PCR using a QuantStudio 6 Pro RT-PCR System (Thermo).
765
766 **Blue-light LED treatment**
767 Cells were exposed to blue light for indicated amounts of time using a 12 inch square LED Grow
768 Light System with 225 x 14W Blue LEDs (HQRP #884667106091218) mounted inside of their cell
769 culture incubator.
770
771 **Fluorescence recovery after photobleaching**
772 Cells were bleached using a blue light (488 nm) laser, followed by green channel imaging every
773 second for 3 min. Imaging was performed using an Andor Revolution Spinning Disk Confocal
774 with FRAPPA system. FRAP Analysis was performed using Fiji (ImageJ).
775
Immunofluoresence and live imaging
Cells were grown in plastic Nunc Cell Culture Treated Multidishes (Thermo #140675). For live
imaging, cells were aspirated of their cell growth media and replaced with PBS. For
immunofluorescence, cells were fixed using 3.7% formaldehyde in PBS at room temperature for

15 min, washed with PBS for 5 min three times, blocked using normal goat serum diluted in PBS
with 0.3% triton for at least 1 h, then treated with primary antibody diluted in 1% normal goat
serum, 0.3% triton, and PBS. Primary antibody treatments were performed overnight at 4°C.
Cells were then washed in PBS for 5 min three times and treated with secondary antibody using
same dilution buffer for 1–2 h at room temperature. Cells were washed a final time with PBS
and mounted using mounting media containing DAPI (VECTASHIELD #H-2000-10). Imaging was
performed on a Nikon Ti automated inverted microscope with incubation enclosure. Images
were analyzed using Fiji (Image J).
788
smFISH
For smFISH, cells were plated in glass bottom, 10 mm, 24 well plates (MatTek Corporation
#P24G-1.5-10-F). Cells were prepared following the standard Stellaris RNA FISH protocol for
adherent cells (Biosearch Technologies). Cells were fixed using 3.7% formaldehyde in PBS at
room temperature for 10 min. Cells were then permeabilized using 70% ethanol for at least 1 h
at 4°C. Cells were washed with Wash buffer A and incubated at room temperature for 5 min.
Wash buffer A was aspirated and cells were treated with approximately 30 μl of 125 nM FISH
probe diluted in hybridization buffer. The plate was sealed with parafilm and stored in dark
humid chamber overnight at 37°C. Cells were then washed with wash buffer A for 30 min at
37°C , then washed with wash buffer B for 5 min and mounted with mounting media containing
DAPI (VECTASHIELD #H-2000-10). smFISH imaging was performed using a Zeiss LSM 980 with
Airyscan 2 Laser Scanning Confocal imaging system with Zeiss AxioObserver motorized inverted
microscope stand, DIC optics, and a motorized XYZ stage. Images were analyzed using IMARIS.
802
Western Blots
Samples used for western blots were lysed by boiling in NuPAGE LDS Sample Buffer (Thermo #
NP0007) diluted to 1x concentration and supplemented with 1 mM DTT. Samples were run on
NuPAGE 4 to 12% Bis-Tris 1.0 mm miniprotein gels (Thermo #NP0321BOX), then transferred to
0.2μM PVDF transfer membranes (Thermo #22860). All protein gels and transfers were run
using a mini gel tank (Thermo #A25977) with protein gels run at 200V for 80 min and transfers
run at 30V for 60 min. Transferred membranes were blocked in PBS with 1% Triton and 5% milk
for at least 1 h at room temperature before they were transferred to primary antibody diluted in
the same buffer at 4°C. Primary antibody was applied overnight followed by at least three 5 min
washes with the same blocking buffer, secondary antibody diluted in the same buffer for 30 min
at room temperature, and three 5 min washes in the same buffer. Westerns were imaged using
the Odyssey CLx imaging system by LI-COR. All western blot quantifications were done using
Fiji/ImageJ software.
816
Ribosome profiling and matched RNA-seq
HCT116 cells were cultured in a 15 cm dish, washed twice with cold PBS, scraped into lysis
buffer containing 10 mM Tris HCl pH 7.5, 5 mM $MgCl_2$, 100 mM KCl, 1% triton, 2 mM DTT, 1x
cOmplete mini tablet (EDTA free), 0.3 U/μl RNasin Plus RNase inhibitor, and 1:60 dilution of
ribo-spike. Cells were lysed by passing through 26G ½ inch needle at least 7 times. Lysates were
depleted of nuclei by centrifugation at 1300*g* for 10 min at 4°C, divided into aliquots for ribo-

seq and matched RNA libraries, and snap frozen using liquid nitrogen. Ribosome profiling and
matched RNA seq libraries were then prepared according to previously established protocol,[77] a
detailed version of which is available at http://bartellab.wi.mit.edu/ protocols.html. Sequencing
was performed on an Illumina HiSeq 2500 using 50 cycles. Only reads mapping to ORFs
of GRCh38-annotated genes, fluorescent protein-coding genes, the OsTIR1 gene, or the *fLuc*
gene were used. Reads mapping to the first 50 nt of each ORF were excluded. Reads mapping to
heme-related genes from the rabbit genome (Broad/oryCun2) were also excluded. TE changes
were calculated and count depth normalization was performed for using DESeq2.[78]
831

Isolation of RNA from HCT116 cells and SG cores for RNA sequencing
Isolation and sequencing of SG cores was adapted from an established protocol.[18,52] HCT116
cells were cultured in a 24.5 cm square dish, washed with PBS, scraped into PBS, centrifuged at
1500g for 3 min, aspirated to remove liquid, and snap frozen in liquid nitrogen. Frozen cell
pellets were thawed in lysis buffer (50 mM Tris HCl pH 7.4, 2 mM MgOAc, 10 mM $MgCl_2$, 100
mM KOAc, 0.1% NP40, 0.5 mM DTT, 1x cOmplete mini tablet (EDTA free), and 1 U/µl RNasin Plus
RNase inhibitor). Cells were lysed by passing through 26G ½ inch needle at least seven times.
Lysates were depleted of their nuclei by centrifugation at 1000g for 5 min at 4°C. 5% of this
nuclear-depleted sample was aliquoted and snap frozen in liquid nitrogen and stored at −80°C
as the cytoplasmic fraction. The remaining 95% was used to isolate the SG fraction. Sample was
centrifuged at 18,000g for 20 min at 4°C. Pellet was resuspended in 1 ml lysis buffer and
allowed to turn end-over-end for at least 10 min at 4°C. Once resuspended, sample was
centrifuged again at 18,000g for 20 min at 4°C. The pellet was resuspended in 300 µl lysis buffer
and centrifuged at 850g for 2 min at 4°C. The supernatant was taken and incubated with 6 µg of
G3BP2 antibody (Bethyl Laboratories #A302-040A) overnight at 4°C, rotating end-over-end.
Sample was centrifuged at 18,000g for 20 min at 4°C to remove unbound antibody, and the
pellet was resuspended in 500 µl lysis buffer. 625 µl of M-280 sheep, anti-rabbit IgG dynabeads
(Thermo # 11204D) were then added to the sample and incubated for at least 3 h at 4°C,
rotating end-over-end. Beads were washed with buffer containing 20 mM Tris HCl pH 8.0 and 50
mM NaCl and eluted using a high-salt buffer (10mM Tris HCl pH 7.5, 1M NaCl, 5% 1,6
hexanediol, 5mM $MgCl_2$, 100mM KCl, 1% triton, 2mM DTT, 1x cOmplete mini tablet (EDTA free),
and 0.3 U/µl RNasin Plus RNase inhibitor), turning end-over-end for at least 4 h. Eluate was snap
frozen using liquid nitrogen and stored at −80°C as the SG fraction. RNA-sequencing libraries
were then prepared from the cytoplasmic and SG fractions in parallel, following the protocol
used for the RNA-sequencing libraries that matched the ribosome profiling. SG enrichment
values were calculated using DESeq2.[78]
858

Statistical analysis and comparison with published datasets
860 Plots and statistical analyses were performed using R. Statistical tests and parameters are shown
861 in figure legends. Additional data were obtained from publicly available datasets associated with
862 published work[18,43,46,53,54] and re-analyzed using R.
863

Figure S1. Heat shock and oxidative stress illicit similar but distinct ISR translation programs.
(A) Quantification of ribo-spike. Cytoplasmic extract from HCT116 cells was serially diluted to
achieve the indicated relative concentrations, and then a constant amount of ribo-spike was
added to each sample before ribosome profiling and RNA-seq analyses. Plotted are the relative
concentrations of total mRNA (dashed line) and RPFs (solid line) as calculated from the 1000
most highly expressed genes, using no-spike controls (blue) and ribo-spike normalized samples
(red) as a function of the actual relative concentrations. **(B)** Relative TE measurements before
adjusting for ribo-spike. Plotted with circles are the average $\log_2$ TE values (line, median; notch,
95% confidence interval; box, quartiles; whiskers, 1.5 x IQR) calculated for mRNAs from either
untreated cells (black, three replicates), cells treated with 500 µM sodium arsenite for 1 h
(green, four replicates), or cells treated with heat shock at 45°C for 25 min (pink, 2 replicates).
Red Xs denote the ribo-spike TE values (one for each replicate). Statistical significance was
calculated based on ribo-spike values alone (***, $p \leq 0.001$; Welch's two-sample t-test). **(C)** TE
measurements of panel B after adjusting for ribo-spike; otherwise, as in (B). Note that statistical
significance is determined in panel B, using ribo-spike values prior to adjustment (***, $p \leq$
0.001). **(D)** Preferential translation of mRNAs containing uORFs during arsenite stress. The plots
indicate $\log_2$ FC in TE caused by arsenite for mRNAs containing at least one uORF and those
containing no uORFs, according to annotations from Chen et al. (2020)[43] (line, median; notch,
95% confidence interval; box, quartiles; whiskers, 1.5 x IQR); ***, $p \leq 0.001$; Welch's two-
sample t-test). **(E)** Relationship between translation during arsenite stress and mRNA half-lives.
The plots indicate the $\log_2$ FC in TE due to arsenite as a function of the mean-normalized half-
life (PC1) of mRNAs as calculated from consensus of multiple human cell data sets by Agarwal et
al. (2022).[46] n indicates number of unique mRNAs. **(F)** Preferential translation of mRNAs
containing uORFs during heat shock. Plots indicate $\log_2$ FC in TE caused by treatment with 45°C
heat shock; otherwise, as in (D). **(G)** Relationship between translation during heat shock and
mRNA half-lives. The plots indicate the $\log_2$ FC in TE due to heat shock as a function of the
mean-normalized half-life of mRNAs; otherwise, as in (E). **(H – J)** Comparison of arsenite stress
and heat shock ISR translation programs. Plotted for each mRNA are changes in TE caused by
arsenite as a function of changes in TE caused by heat shock. Points for mRNAs classified as
either stress-enhanced, stress-resistant, or stress-hypersensitive mRNAs in either arsenite stress
(D), heat shock (E), or both (F) are colored. n indicates number of genes; R_s indicates Spearman
coefficient.

901
902

Figure S2. Neither CAPRIN1 nor UBAP2L are required for SG formation in HCT116 cells. (A) The ISR translation response to arsenite stress, either with or without G3BPs. The plots indicate the $\log_2$ FC in TE caused by arsenite stress in either untreated HCT116 cells or cells depleted of their G3BPs by addition of 500 μM IAA (line, median; notch, 95% confidence interval; box, quartiles; whiskers, 1.5 x IQR). Significance was determined using ribo-spike values, as in Figure S1B (* $p \leq$ 0.05, ***, $p \leq$ 0.001; Welch's two-sample t-test). **(B)** Comparison of G3BP-dependent translation and ISR translation in HCT116 cells treated with arsenite stress. The plots indicate the $\log_2$ FC in TE due to presence of G3BPs in stressed cells in each of three ISR translation categories: stress-enhanced/resistant, bulk behavior, and stress-hypersensitive (line, median; notch, 95% confidence interval; box, quartiles; whiskers, 1.5 x IQR; ***, $p \leq$ 0.001; Welch's two-sample t-test). **(C)** The ISR translation response to heat shock, either with or without G3BPs. ns, not significant; otherwise, as in (A). **(D)** Comparison of G3BP-dependent translation and ISR translation in HCT116 cells treated with heat-shock stress; otherwise, as in (B). **(E)** G3BP-dependent translation of mRNAs containing uORFs during arsenite stress. The plots indicate the $\log_2$ FC in TE due to presence of G3BPs in arsenite-stressed cells for mRNAs containing at least one uORF and those containing no uORFs, according to annotations from Chen et al. (2020)[43] (line, median; notch, 95% confidence interval; box, quartiles; whiskers, 1.5 x IQR); ***, $p \leq$ 0.001; Welch's two-sample t-test). **(F)** G3BP-dependent translation of mRNAs containing uORFs during heat stress. The plots indicate the $\log_2$ FC in TE due to presence of G3BPs in heat-stressed cells for mRNAs containing at least one uORF and those containing no uORFs; otherwise as in (E). **(G)** Relationship between G3BP-dependent translation during arsenite stress and mRNA half-lives. The plots indicate the $\log_2$ FC in TE due to presence of G3BPs in arsenite-stressed cells as a function of the mean-normalized half-life (PC1) of mRNAs as calculated from consensus of multiple human cell data sets by Agarwal et al. (2022).[46] n indicates number of unique mRNAs. **(H)** Relationship between G3BP-dependent translation during heat stress and mRNA half-lives. The plots indicate the $\log_2$ FC in TE due to presence of G3BPs in heat-stressed cells as a function of the mean-normalized half-life of mRNAs; otherwise, as in (G). **(I)** Depletion of endogenous CAPRIN1 fused to AID. Immunoblots probed for the indicated proteins show specific depletion of AID fusion protein after treatment with 500 μM IAA for 3 h. **(J)** Quantification of protein depletion in (I), normalizing to levels of GAPDH. Points show values for three biological replicates. **(K)** Colocalization of CAPRIN1 in SGs. Quantification of colocalization was performed using Mander's coefficient between G3BP1 and indicated SG marker, comparing to a control image from an irrelevant field of cells (***, $p \leq$ 0.001; Welch's two-sample t-test). **(L)** SG formation despite CAPRIN1 depletion. HCT116 cells were either treated with 500 μM IAA to deplete CAPRIN1 (right) or not treated (left) prior to arsenite stress (500 μM sodium arsenite for 1 h). Images show either CAPRIN1 in red, SG marker protein G3BP1 in green, or SG marker protein eIF4G in magenta (scale bars, 10 μm). **(M)** Quantification of SG formation in (L). Points show values for three biological replicates (ns, not significant). **(N)** Depletion of endogenous UBAP2L fused to AID. Quantification was based on AID::RFP::UBAP2L and PABPC1::GFP fluorescence in homozygously tagged HCT116 cells (images not shown). Points show values for three biological replicates. **(O)** Colocalization of UBAP2L in SGs. Quantification of colocalization was performed using Mander's coefficient between PABPC1 and UBAP2L, comparing to a control image from an irrelevant field of cells (***, $p \leq$ 0.001; Welch's two-sample t-test). **(P)** SG formation despite UBAP2L depletion. HCT116 cells were either treated with 500 μM IAA to

947   deplete UBAP2L (right) or not treated (left) prior to arsenite stress (500 µM sodium arsenite for
948   1 h). Images show either UBAP2L in red or SG marker protein PABPC1 in green; otherwise, as in
949   (L). **(Q)** Quantification of SG formation in (P); otherwise, as in (M).

Figure S3. Conclusions regarding SG-enrichment are consistent across multiple cell types and stresses. (A) Comparison of SG enrichments. Shown are SG enrichments calculated in Khong et al. (2017)[18] plotted as a function of SG enrichments calculated in this study. *n* indicates total number of unique mRNAs. **(B)** Comparison of mRNA levels for the cells used in (A). Shown are relative mRNA abundances in U2OS cells, as measured by Khong et al. (2017)[18], plotted as a function of relative abundance in HCT116 cells, as measured in this study. Otherwise as in (A). **(C – E)** SG enrichment determinants. Shown is the relationship between either transcript length (C), unstressed TE (D), or stressed TE (E) and SG enrichment; otherwise as in Fig 3C (* $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$). **(F)** Comparison of SG enrichment in arsenite-stressed U2OS cells as measured by Khong et al. (2017)[18] and ISR translation in arsenite stressed HCT116 cells; otherwise, as in Fig 3B. **(G)** Comparison of arsenite-stressed TE in HCT116 cells and SG enrichment in arsenite-stressed U2OS cells as measured by Khong et al. (2017)[18]; otherwise, as in (E). **(H)** Comparison of Stress-Dependent RNP Granule (RG) enrichment and ISR translation in ER-stressed NIH 3T3 cells as measured by Namkoong et al. (2018);[53] otherwise, as in (F). **(I)** Comparison of G3BP-depdenent proximity labeling in unstressed HEK293T cells as measured by Ren et al. (2023)[54] and ISR translation in arsenite stressed HCT116 cells. ns, not significant; otherwise, as in (F). **(J)** Comparison of G3BP-depdenent proximity labeling in arsenite-stressed HEK293T cells as measured by Ren et al. (2023)[54] and ISR translation in arsenite-stressed HCT116 cells; otherwise, as in (I). **(K)** Comparison of G3BP-dependent translation in HCT116 cells and SG enrichment in U2OS cells as measured by Khong et al. (2017)[18]. All stresses were 500 µM sodium arsenite for 1 h; otherwise, as in (E). **(L)** Comparison of G3BP-dependent translation in HCT116 cells and RG enrichment in NIH 3T3 cells as measured by Namkoong et al. (2018).[53] All stresses were heat shock; otherwise, as in (I). **(M)** Comparison of G3BP-dependent translation in HCT116 cells and G3BP-depdenent proximity labeling in stressed HEK293T cells as measured by Ren et al. (2023)[54]. All stresses were 500 µM sodium arsenite for 1 h; otherwise, as in (K).

Figure S4. OGs are SG-like condensates formed in the absence of stress. (A) No induction of ectopic SGs caused by UBAP2L overexpression. Unstressed U2OS cells were either treated with 1 μM doxycycline (right) or not treated (left) to induce overexpression of UBAP2L. Images show either UBAP2L in red or SG marker protein eIF4G in magenta (scale bars, 10 μm). White boxes highlight insets that are expanded on the right. **(B)** Colocalization of UBAP2L puncta with SG marker. Quantification of colocalization was performed using Mander's coefficient between UBAP2L and eIF4G as an SG marker, comparing to a control image from an irrelevant field of cells (***, $p \leq 0.001$; Welch's two-sample t-test)**. (C)** Induction of ectopic SGs caused by G3BP overexpression. Unstressed HCT116 cells were either treated with 1 μM doxycycline (right) or not treated (left) to induce overexpression of G3BP1. Images show either G3BP1 in green or SG marker protein eIF4G in magenta; otherwise, as in (A). **(D)** Quantification of ectopic granule formation in (C). Points indicate values for three biological replicates. **(E)** Kinetics of OG formation and dissolution. Plotted over time is the percent of cells with visible OGs. Cells were exposed to blue light starting at 0 min and removed from blue light at 360 min. Error bars report standard deviation for six biological replicates shown as points. **(F)** Colocalization of SG markers in OGs. Quantification of colocalization was performed using Mander's coefficient between GFP::CRY2::G3BP$_{\Delta N}$ and indicated SG marker, comparing to a control image from an irrelevant field of cells (***, $p \leq 0.001$; Welch's two-sample t-test)**. (G)** FRAP of OGs. Plotted over time is the fluorescence intensity of GFP-labeled granules after photobleaching. SG (green) and OG (blue) recoveries are shown over a 180 s time course. Photobleaching occurred at 3 s. Error bars report standard deviation for at least six granules from at least three cells.

1076 **Figure S5. OGs require stress to recruit the SG transcriptome. (A)** Expanded comparisons of
1077 granule enrichment, showing the distributions used to calculate the R_S values of Fig 5E. n
1078 indicates total number of unique mRNAs. **(B)** Expanded comparisons of cell line transcriptomes,
1079 showing the distributions used to calculate the R_S values of Fig 5F.

Figure S6. Tethering to SGs via CAPRIN1 imparts G3BP-dependent resistance to ISR translation
shutdown. (A) G3BP1 partitioning to SGs. HCT116 cells expressing G3BP1 endogenously tagged
with GFP were stressed with 500 μM sodium arsenite for 90 min. Images show localization of
G3BP in green. White outlines show cell boundaries as determined by Imaris software based on
cytosolic fluorescence. Purple outlines show SG boundaries as determined by Imaris software
(scale bars, 1 μm). **(B)** Quantification of G3BP1 partitioning to SGs. Points show the amount of
fluorescence from SGs as a percent of total fluorescence, for eight independent replicates each
containing at least three cells. **(C)** Effects of tagging G3BP with λN and MCP. HCT116 cells
expressing endogenous G3BP tagged with the indicated GFP fusion protein were either
unstressed (top) or treated with 500 μM sodium arsenite for 1 h (bottom) to assess SG
formation (scale bar, 10 μm). White boxes highlight insets that are expanded on the right. **(D)**
mRNA changes during stress. Plotted for each mRNA is its abundance in arsenite-stressed
HCT116 cells (500 μM sodium arsenite for 1 h) as a function of its abundance in unstressed
HCT116 cells. The *GAPDH* mRNA, which was used as a control for RT-qPCR, is highlighted in
green. The red dashed line indicates unchanged abundance. *n* indicates total number of unique
mRNAs. **(E)** Quantification of MCP expression. Plot indicates relative MCP expression as
determined by RT-qPCR in HCT116 cell lines expressing either endogenously tagged GFP::G3BP
(untethered), GFP::G3BP and MCP::BFP (MCP::BFP), or MCP::GFP::G3BP (tethered). The points
show values of at least three biological replicates. **(F)** Unstressed SG-tethered translation
response. Plotted for each reporter is the unstressed $\log_2$ TE in either untethered cells (green),
untethered cells expressing MCP::BFP (blue), or tethered cells (pink). **(G)** Stressed SG-tethered
translation response. Plotted for each reporter is the stressed $\log_2$ TE; otherwise, as in (F). **(H)**
Tethering a luciferase reporter to SGs via CAPRIN1. HCT116 cells expressing a NanoLuc reporter
bearing 5' and 3' UTRs from *CNOT10* and MCP::RFP::CAPRIN1 were stressed with 500 μM
sodium arsenite for 90 min. Images show localization of CAPRIN1 in red and smFISH of NanoLuc
reporter molecules in yellow (scale bars, 1 μm). White boxes highlight insets that are expanded
on the right. **(I)** SG tethering efficiency. Percent of cytoplasmic reporter molecules localized to
SGs in untethered cells (replotted from Figure 6C in green) and to SGs in CAPRIN1-tethered cells
(pink) are shown for reporters with 5' and 3' UTRs of the indicated genes. Points show values for
at least three biological replicates (** $p \le 0.01$, *** $p \le 0.001$, Welch's two-sample t-test). The
average SG tethering efficiency for the four different reporters was 30% for untethered cells and
89% for tethered cells, with standard deviations of 11% and 4%, respectively. **(J)** Cells expressing
CAPRIN1 tether while depleted of G3BPs. HCT116 cells expressing endogenously tagged
AID::RFP::G3BPs were either untreated (left), treated with 1 μM doxycycline and 200 μM
auxinole for 10 h, leading to retention of G3BPs and expression of MCP::GFP::CAPRIN1 (middle),
or treated with 1 μM doxycycline and 500 μM IAA for 10 h, leading to turnover of endogenous
G3BPs and expression of MCP::GFP::CAPRIN1 (right). Otherwise as in (C). **(K)** Assessment of SG
formation in cells expressing CAPRIN1 tether while depleted of G3BPs. Cells were treated as in
(J) and then subsequently stressed with 500 μM sodium arsenite for 90 min; otherwise, as in (J).
(L) Quantification of granule formation in (K). Error bars report standard deviation for three
biological replicates (*** $p \le 0.001$, ns, not significant, Welch's two-sample t-test). **(M)**
CAPRIN1-tethered ISR translation response. Plotted for each reporter is the change in TE caused
by arsenite stress (500 μM sodium arsenite for 90 min) in untethered cells (green), tethered
cells expressing G3BPs (pink), or tethered cells depleted of G3BPs (red) (*** $p \le 0.001$, ** $p \le$

0.01, ns, not significant, Welch's two-sample t-test). The average $\log_2$ FC in TE for the ISR
translation response was −2.24 for untethered cells, −1.12 for CAPRIN1-tethered cells
expressing G3BPs, and −2.09 for CAPRIN1-tethered cells depleted of G3BPs.

Table S1

RNA	Endogenous TE change log$_2$(stressed/unstressed)	Reporter TE change log$_2$(stressed/unstressed)
CNOT10	−4.44	−5.49
DIMT1	−1.82	−4.42
DDIT4L	4.18	−5.362
RB1	−2.24	−3.24

1206 **Table S1. Luciferase reporters do not recapitulate endogenous TE behavior.** Listed for each
1207 mRNA are endogenous changes in TE due to presence of arsenite stress (500 µM sodium
1208 arsenite for 1 h), and reporter changes in TE due to presence of arsenite stress (500 µM sodium
1209 arsenite for 1.5 h).
1210
1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1　Kedersha, N. *et al.* G3BP-Caprin1-USP10 complexes mediate stress granule condensation
and associate with 40S subunits. *J Cell Biol* **212**, 845-860, doi:10.1083/jcb.201508028
(2016).
2　Yang, P. *et al.* G3BP1 Is a Tunable Switch that Triggers Phase Separation to Assemble
Stress Granules. *Cell* **181**, 325-345 e328, doi:10.1016/j.cell.2020.03.046 (2020).
3　Sanders, D. W. *et al.* Competing Protein-RNA Interaction Networks Control Multiphase
Intracellular Organization. *Cell* **181**, 306-324 e328, doi:10.1016/j.cell.2020.03.050 (2020).
4　Guillen-Boixet, J. *et al.* RNA-Induced Conformational Switching and Clustering of G3BP
Drive Stress Granule Assembly by Condensation. *Cell* **181**, 346-361 e317,
doi:10.1016/j.cell.2020.03.049 (2020).
5　Buchan, J. R. & Parker, R. Eukaryotic stress granules: the ins and outs of translation. *Mol*
Cell **36**, 932-941, doi:10.1016/j.molcel.2009.11.020 (2009).
6　Advani, V. M. & Ivanov, P. Translational Control under Stress: Reshaping the Translatome.
Bioessays **41**, e1900009, doi:10.1002/bies.201900009 (2019).
7　Pakos-Zebrucka, K. *et al.* The integrated stress response. *EMBO Rep* **17**, 1374-1395,
doi:10.15252/embr.201642195 (2016).
8　Harding, H. P. *et al.* Regulated translation initiation controls stress-induced gene
expression in mammalian cells. *Mol Cell* **6**, 1099-1108, doi:10.1016/s1097-
2765(00)00108-8 (2000).
9　Harding, H. P., Zhang, Y. & Ron, D. Protein translation and folding are coupled by an
endoplasmic-reticulum-resident kinase. *Nature* **397**, 271-274, doi:10.1038/16729
(1999).
10　Yamasaki, S. & Anderson, P. Reprogramming mRNA translation during stress. *Curr Opin*
Cell Biol **20**, 222-226, doi:10.1016/j.ceb.2008.01.013 (2008).
11　Starck, S. R. *et al.* Translation from the 5' untranslated region shapes the integrated
stress response. *Science* **351**, aad3867, doi:10.1126/science.aad3867 (2016).
12　Sidrauski, C., McGeachy, A. M., Ingolia, N. T. & Walter, P. The small molecule ISRIB
reverses the effects of eIF2alpha phosphorylation on translation and stress granule
assembly. *Elife* **4**, doi:10.7554/eLife.05033 (2015).
13　Vattem, K. M. & Wek, R. C. Reinitiation involving upstream ORFs regulates ATF4 mRNA
translation in mammalian cells. *Proc Natl Acad Sci U S A* **101**, 11269-11274,
doi:10.1073/pnas.0400541101 (2004).
14　Protter, D. S. W. & Parker, R. Principles and Properties of Stress Granules. *Trends Cell Biol*
26, 668-679, doi:10.1016/j.tcb.2016.05.004 (2016).
15　Kedersha, N. L., Gupta, M., Li, W., Miller, I. & Anderson, P. RNA-binding Proteins TIA-1
and TIAR Link the Phosphorylation of eIF-2 to the Assembly of Mammalian Stress
Granules. *The Journal of Cell Biology* **147**, 1431-1441 (1999).
16　Molliex, A. *et al.* Phase separation by low complexity domains promotes stress granule
assembly and drives pathological fibrillization. *Cell* **163**, 123-133,
doi:10.1016/j.cell.2015.09.015 (2015).
17　Jain, S. *et al.* ATPase-Modulated Stress Granules Contain a Diverse Proteome and
Substructure. *Cell* **164**, 487-498, doi:10.1016/j.cell.2015.12.038 (2016).

18   Khong, A. *et al.* The Stress Granule Transcriptome Reveals Principles of mRNA
1273   Accumulation in Stress Granules. *Mol Cell* **68**, 808-820 e805,
1274   doi:10.1016/j.molcel.2017.10.015 (2017).
1275   19   Markmiller, S. *et al.* Context-Dependent and Disease-Specific Diversity in Protein
1276   Interactions within Stress Granules. *Cell* **172**, 590-604 e513,
1277   doi:10.1016/j.cell.2017.12.032 (2018).
1278   20   Kedersha, N. *et al.* Evidence That Ternary Complex (eIF2-GTP-tRNAiMet)-Deficient
1279   Preinitiation Complexes Are Core Constituents of Mammalian Stress Granules. *Molecular*
1280   *Biology of the Cell* **13**, 195-210, doi:10.1091/mbc.01-05-0221 (2002).
1281   21   Laver, J. D. *et al.* The RNA-Binding Protein Rasputin/G3BP Enhances the Stability and
1282   Translation of Its Target mRNAs. *Cell Rep* **30**, 3353-3367 e3357,
1283   doi:10.1016/j.celrep.2020.02.066 (2020).
1284   22   Alam, U. & Kennedy, D. Rasputin a decade on and more promiscuous than ever? A
1285   review of G3BPs. *Biochim Biophys Acta Mol Cell Res* **1866**, 360-370,
1286   doi:10.1016/j.bbamcr.2018.09.001 (2019).
1287   23   Meyer, C., Garzia, A., Morozov, P., Molina, H. & Tuschl, T. The G3BP1-Family-USP10
1288   Deubiquitinase Complex Rescues Ubiquitinated 40S Subunits of Ribosomes Stalled in
1289   Translation from Lysosomal Degradation. *Mol Cell* **77**, 1193-1205 e1195,
1290   doi:10.1016/j.molcel.2019.12.024 (2020).
1291   24   Anderson, P. & Kedersha, N. RNA granules: post-transcriptional and epigenetic
1292   modulators of gene expression. *Nat Rev Mol Cell Biol* **10**, 430-436, doi:10.1038/nrm2694
1293   (2009).
1294   25   Kedersha, N. *et al.* Dynamic shuttling of TIA-1 accompanies the recruitment of mRNA to
1295   mammalian stress granules. *J Cell Biol* **151**, 1257-1268, doi:10.1083/jcb.151.6.1257
1296   (2000).
1297   26   Kwon, S., Zhang, Y. & Matthias, P. The deacetylase HDAC6 is a novel critical component of
1298   stress granules involved in the stress response. *Genes Dev* **21**, 3381-3394,
1299   doi:10.1101/gad.461107 (2007).
1300   27   Youn, J. Y. *et al.* Properties of Stress Granule and P-Body Proteomes. *Mol Cell* **76**, 286-
1301   294, doi:10.1016/j.molcel.2019.09.014 (2019).
1302   28   Kimball, S. R., Horetsky, R. L., Ron, D., Jefferson, L. S. & Harding, H. P. Mammalian stress
1303   granules represent sites of accumulation of stalled translation initiation complexes. *Am J*
1304   *Physiol Cell Physiol* **284**, C273-284, doi:10.1152/ajpcell.00314.2002 (2003).
1305   29   Moon, S. L. *et al.* Multicolour single-molecule tracking of mRNA interactions with RNP
1306   granules. *Nat Cell Biol* **21**, 162-168, doi:10.1038/s41556-018-0263-4 (2019).
1307   30   Mateju, D. *et al.* Single-Molecule Imaging Reveals Translation of mRNAs Localized to
1308   Stress Granules. *Cell* **183**, 1801-1812 e1813, doi:10.1016/j.cell.2020.11.010 (2020).
1309   31   Tourriere, H. *et al.* The RasGAP-associated endoribonuclease G3BP assembles stress
1310   granules. *J Cell Biol* **160**, 823-831, doi:10.1083/jcb.200212128 (2003).
1311   32   Zhang, P. *et al.* Chronic optogenetic induction of stress granules is cytotoxic and reveals
1312   the evolution of ALS-FTD pathology. *Elife* **8**, doi:10.7554/eLife.39578 (2019).
1313   33   Baymiller, M. & Moon, S. L. Stress Granules as Causes and Consequences of Translation
1314   Suppression. *Antioxid Redox Signal* **39**, 390-409, doi:10.1089/ars.2022.0164 (2023).

34   Putnam, A., Thomas, L. & Seydoux, G. RNA granules: functional compartments or
incidental condensates? *Genes Dev* **37**, 354-376, doi:10.1101/gad.350518.123 (2023).
35   Mateju, D. & Chao, J. A. Stress granules: regulators or by-products? *FEBS J* **289**, 363-373,
doi:10.1111/febs.15821 (2022).
36   Ingolia, N. T. Ribosome Footprint Profiling of Translation throughout the Genome. *Cell*
165, 22-33, doi:10.1016/j.cell.2016.02.066 (2016).
37   Iwasaki, S., Floor, S. N. & Ingolia, N. T. Rocaglates convert DEAD-box protein eIF4A into a
sequence-selective translational repressor. *Nature* **534**, 558-561,
doi:10.1038/nature17978 (2016).
38   Wilbertz, J. H. *et al.* Single-Molecule Imaging of mRNA Localization and Regulation
during the Integrated Stress Response. *Mol Cell* **73**, 946-958 e947,
doi:10.1016/j.molcel.2018.12.006 (2019).
39   Aulas, A. *et al.* Stress-specific differences in assembly and composition of stress granules
and related foci. *J Cell Sci* **130**, 927-937, doi:10.1242/jcs.199240 (2017).
40   Sonenberg, N. & Hinnebusch, A. G. Regulation of translation initiation in eukaryotes:
mechanisms and biological targets. *Cell* **136**, 731-745, doi:10.1016/j.cell.2009.01.042
(2009).
41   Holcik, M. & Sonenberg, N. Translational control in stress and apoptosis. *Nat Rev Mol*
Cell Biol **6**, 318-327, doi:10.1038/nrm1618 (2005).
42   Andreev, D. E. *et al.* Translation of 5' leaders is pervasive in genes resistant to eIF2
repression. *Elife* **4**, e03971, doi:10.7554/eLife.03971 (2015).
43   Chen, J., Brunner, A.-D., Nunez, J. K. & Weissman, J. Pervasive functional translation of
noncanonical human open reading frames. *Science* **367**, 1135-1140,
doi:10.1126/science.aay0262 (2020).
44   Glauninger, H. *et al.* Transcriptome-wide mRNA condensation precedes stress granule
formation and excludes stress-induced transcripts. *bioRxiv*,
doi:10.1101/2024.04.15.589678 (2024).
45   Mostafa Zedan, A. P. S., Stephanie Heinrich, Pablo Aurelio Gómez García, Sarah Khawaja,
Karsten Weis. Newly synthesized mRNA selectively escapes translational repression
following acute stress. *bioRxiv*, doi:https://doi.org/10.1101/2024.04.14.589419 (2024).
46   Agarwal, V. & Kelley, D. R. The genetic and biochemical determinants of mRNA
degradation rates in mammals. *Genome Biology* **23**, 245, doi:10.1186/s13059-022-
02811-x (2022).
47   Pizzinga, M. *et al.* The cell stress response: extreme times call for post-transcriptional
measures. *Wiley Interdiscip Rev RNA* **11**, e1578, doi:10.1002/wrna.1578 (2020).
48   Natsume, T., Kiyomitsu, T., Saga, Y. & Kanemaki, M. T. Rapid Protein Depletion in Human
Cells by Auxin-Inducible Degron Tagging with Short Homology Donors. *Cell Rep* **15**, 210-
218, doi:10.1016/j.celrep.2016.03.001 (2016).
49   Nishimura, K., Fukagawa, T., Takisawa, H., Kakimoto, T. & Kanemaki, M. An auxin-based
degron system for the rapid depletion of proteins in nonplant cells. *Nat Methods* **6**, 917-
922, doi:10.1038/nmeth.1401 (2009).
50   Cirillo, L. *et al.* UBAP2L Forms Distinct Cores that Act in Nucleating Stress Granules
Upstream of G3BP1. *Curr Biol* **30**, 698-707 e696, doi:10.1016/j.cub.2019.12.020 (2020).

51   Paget, M. *et al.* Stress granules are shock absorbers that prevent excessive innate
immune responses to dsRNA. *Mol Cell* **83**, 1180-1196 e1188,
doi:10.1016/j.molcel.2023.03.010 (2023).

52   Khong, A., Jain, S., Matheny, T., Wheeler, J. R. & Parker, R. Isolation of mammalian stress
granule cores for RNA-Seq analysis. *Methods* **137**, 49-54,
doi:10.1016/j.ymeth.2017.11.012 (2018).

53   Namkoong, S., Ho, A., Woo, Y. M., Kwak, H. & Lee, J. H. Systematic Characterization of
Stress-Induced RNA Granulation. *Mol Cell* **70**, 175-187 e178,
doi:10.1016/j.molcel.2018.02.025 (2018).

54   Ren, Z., Tang, W., Peng, L. & Zou, P. Profiling stress-triggered RNA condensation with
photocatalytic proximity labeling. *Nat Commun* **14**, 7390, doi:10.1038/s41467-023-
43194-2 (2023).

55   Huang, C. *et al.* UBAP2L arginine methylation by PRMT1 modulates stress granule
assembly. *Cell Death Differ* **27**, 227-241, doi:10.1038/s41418-019-0350-5 (2020).

56   Panas, M. D., Ivanov, P. & Anderson, P. Mechanistic insights into mammalian stress
granule dynamics. *J Cell Biol* **215**, 313-323, doi:10.1083/jcb.201609081 (2016).

57   Ivanov, P., Kedersha, N. & Anderson, P. Stress Granules and Processing Bodies in
Translational Control. *Cold Spring Harb Perspect Biol* **11**,
doi:10.1101/cshperspect.a032813 (2019).

58   Matheny, T., Rao, B. S. & Parker, R. Transcriptome-Wide Comparison of Stress Granules
and P-Bodies Reveals that Translation Plays a Major Role in RNA Partitioning. *Mol Cell*
Biol **39**, doi:10.1128/MCB.00313-19 (2019).

59   Lee, C. S. *et al.* Recruitment of mRNAs to P granules by condensation with intrinsically-
disordered proteins. *Elife* **9**, doi:10.7554/eLife.52896 (2020).

60   Ramadurgum, P. & Hulleman, J. D. Protocol for Designing Small-Molecule-Regulated
Destabilizing Domains for In Vitro Use. *STAR Protoc* **1**, doi:10.1016/j.xpro.2020.100069
(2020).

61   Bertrand, E. *et al.* Localization of ASH1 mRNA particles in living yeast. *Mol Cell* **2**, 437-
445, doi:10.1016/s1097-2765(00)80143-4 (1998).

62   Wheeler, J. R., Jain, S., Khong, A. & Parker, R. Isolation of yeast and mammalian stress
granule cores. *Methods* **126**, 12-17, doi:10.1016/j.ymeth.2017.04.020 (2017).

63   Matheny, T., Van Treeck, B., Huynh, T. N. & Parker, R. RNA partitioning into stress
granules is based on the summation of multiple interactions. *RNA* **27**, 174-189,
doi:10.1261/rna.078204.120 (2021).

64   Khong, A., Matheny, T., Huynh, T. N., Babl, V. & Parker, R. Limited effects of m(6)A
modification on mRNA partitioning into stress granules. *Nat Commun* **13**, 3735,
doi:10.1038/s41467-022-31358-5 (2022).

65   Burke, J. M., Moon, S. L., Matheny, T. & Parker, R. RNase L Reprograms Translation by
Widespread mRNA Turnover Escaped by Antiviral mRNAs. *Mol Cell* **75**, 1203-1217 e1205,
doi:10.1016/j.molcel.2019.07.029 (2019).

66   Burke, J. M., Ratnayake, O. C., Watkins, J. M., Perera, R. & Parker, R. G3BP1-dependent
condensation of translationally inactive viral RNAs antagonizes infection. *Sci Adv* **10**,
eadk8152, doi:10.1126/sciadv.adk8152 (2024).

67 Burke, J. M., Lester, E. T., Tauber, D. & Parker, R. RNase L promotes the formation of
unique ribonucleoprotein granules distinct from stress granules. *J Biol Chem* **295**, 1426-
1438, doi:10.1074/jbc.RA119.011638 (2020).
68 Buchan, J. R., Muhlrad, D. & Parker, R. P bodies promote stress granule assembly in
Saccharomyces cerevisiae. *J Cell Biol* **183**, 441-455, doi:10.1083/jcb.200807043 (2008).
69 Fedorovskiy, A. G. *et al.* A Solitary Stalled 80S Ribosome Prevents mRNA Recruitment to
Stress Granules. *Biochemistry (Mosc)* **88**, 1786-1799, doi:10.1134/S000629792311010X
(2023).
70 Luo, E. C. *et al.* Large-scale tethered function assays identify factors that regulate mRNA
stability and translation. *Nat Struct Mol Biol* **27**, 989-1000, doi:10.1038/s41594-020-
0477-6 (2020).
71 Bidet, K., Dadlani, D. & Garcia-Blanco, M. A. G3BP1, G3BP2 and CAPRIN1 are required for
translation of interferon stimulated mRNAs and are targeted by a dengue virus non-
coding RNA. *PLoS Pathog* **10**, e1004242, doi:10.1371/journal.ppat.1004242 (2014).
72 Ortega, A. D., Willers, I. M., Sala, S. & Cuezva, J. M. Human G3BP1 interacts with beta-F1-
ATPase mRNA and inhibits its translation. *J Cell Sci* **123**, 2685-2696,
doi:10.1242/jcs.065920 (2010).
73 Wheeler, J. R., Matheny, T., Jain, S., Abrisch, R. & Parker, R. Distinct stages in stress
granule assembly and disassembly. *Elife* **5**, doi:10.7554/eLife.18413 (2016).
74 Downey, N. Extraction of DNA from agarose gels. *Methods Mol Biol* **235**, 137-139,
doi:10.1385/1-59259-409-3:137 (2003).
75 Yesbolatova, A., Natsume, T., Hayashi, K. I. & Kanemaki, M. T. Generation of conditional
auxin-inducible degron (AID) cells and tight control of degron-fused proteins using the
degradation inhibitor auxinole. *Methods* **164-165**, 73-80,
doi:10.1016/j.ymeth.2019.04.010 (2019).
76 Xiang, K. & Bartel, D. P. The molecular basis of coupling between poly(A)-tail length and
translational efficiency. *Elife* **10**, doi:10.7554/eLife.66493 (2021).
77 Subtelny, A. O., Eichhorn, S. W., Chen, G. R., Sive, H. & Bartel, D. P. Poly(A)-tail profiling
reveals an embryonic switch in translational control. *Nature* **508**, 66-71,
doi:10.1038/nature13007 (2014).
78 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8
(2014).
1434