ATAD2 mediates chromatin-bound histone chaperone turnover

Ariadni Liakopoulou[1], Fayçal Boussouar[1#], Daniel Perazza[1], Sophie Barral[1], Emeline Lambert[1],

Tao Wang[1], Florent Chuffart[1], Ekaterina Bourova-Flin[1], Charlyne Gard[2], Denis Puthier[2], Sophie

Rousseaux[1], Christophe Arnoult[1], André Verdel[1], Saadi Khochbin[1#]

1- Université Grenoble Alpes, INSERM U1209, CNRS UMR 5309, Institute for Advanced

Biosciences, 38000, Grenoble, France

2- Aix Marseille Université, INSERM, UMR U1090, TAGC, Marseille, France

Correspondence:

Saadi Khochbin: saadi.khochbin@univ-grenoble-alpes.fr

Fayçal Boussouar: faycal.boussouar@univ-grenoble-alpes.fr

Abstract

ATAD2, a conserved protein which is predominantly expressed in embryonic stem (ES) cells

and spermatogenic cells, emerges as a crucial regulator of chromatin plasticity. Our previous

parallel studies conducted in both ES cells and *S. pombe* highlighted the fundamental role of

ATAD2 in facilitating chromatin-bound histone chaperone turnover. Focusing on

spermatogenesis, we demonstrate here that ATAD2 regulates the HIRA-dependent

localization of H3.3 on the genome and influences H3.3-mediated gene transcription.

Moreover, by modulating histone eviction and the assembly of protamines, ATAD2 ensures

proper chromatin condensation and genome packaging in mature sperm. Disruption of *Atad2*

function in mice leads to abnormal genome organization in mature spermatozoa. Together,

these findings establish a previously overlooked level of chromatin dynamic regulation,

governed by ATAD2-controlled histone chaperones binding to chromatin, which defines the

balance between histone deposition and removal.

Introduction

Spermatogenesis involves unique and dramatic chromatin remodelling and genome reorganization events that take place in preparation for a shift from the universal nucleosome-based eukaryotic genome organization into a unique nucleoprotamine-based structure (Gaucher et al. 2010; Bao and Bedford 2016; Le Blevec et al. 2020; Okada 2022). During the last decade, important information on the molecular basis of the nucleosome-to-nucleoprotamine transition during the postmeiotic phases of spermatogenesis became available. Several testis-specific H2A, H2B and H3 histone variants replace the canonical histones, locally or globally, at different stages, starting early during the commitment of cells into meiotic and post-meiotic differentiation (Hoghoughi et al. 2018). Most of this nucleosome dismantling and assembly occurs in the absence of DNA replication. In other words, the replication-independent nucleosome assembly pathways, such as the well-documented HIRA-dependent H3.3 assembly, are expected to be critical in ensuring the necessary chromatin remodeling in these cells. Finally, hints to the functions of transition proteins (TPs), and their functional relationship with protamines (PRMs) were proposed based on the study of a specific late-expressing H2A variant, H2A.L.2 (Barral et al. 2017b). In the proposed model we suggest that TPs do not directly replace histones, but rather could be loaded on open H2A.L.2-containing nucleosomes to control PRM assembly. Nucleosome-to-nucleoprotamine transition also includes a TP-dependent pre-PRM2 processing, histone displacement by PRMs and the structuration of the nucleoprotamine-based genome organization in the mature spermatozoa (Barral et al. 2017b; Rezaei-Gazik et al. 2022). In mice, two protamines, PRM1 and PRM2, replace histones. PRM2 is initially synthesized as a precursor protein, which undergoes sequential proteolytic processing at its N-terminal region during its incorporation, ultimately producing the mature, fully processed PRM2 (Arevalo et al. 2022).

Taking into account this unique dynamic of post-meiotic cells' nucleosomes, it is expected that these cells massively express generic factors, or those predominantly expressed in spermatogenic cells that act on chromatin, including histone acetylating machineries, acetylated histone readers and histone variants and their chaperones, all of which mediate histone and non-histone protein exchanges. More specifically, the replication-independent nature of nucleosome assembly, both in cells undergoing meiosis and in post-meiotic cells,

strongly suggests the involvement of the chaperone / histone substrate couple, HIRA and

H3.3.

An important part of this replication-independent histone assembly and transitions in the

genome organizations occurs in the context of a massive histone H4 hyperacetylation in the

haploid spermatogenic cells, spermatids, mediated by NUT-p300/CBP histone

acetyltransferase complex (Shiota et al. 2018; Rousseaux et al. 2022). Hyperacetylation of

histone H4, especially, the double acetylation on H4K5K8 in these cells, leads to the binding

of H4 by the first bromodomain of BRDT and to histone replacement (Moriniere et al. 2009;

Gaucher et al. 2012; Goudarzi et al. 2016; Shiota et al. 2018). Additionally, Histone H4

hyperacetylation, especially at K5, could also be recognized by another bromodomain-

containing factor of unknown function in spermatogenic cells, ATAD2 (Caron et al. 2010;

Morozumi et al. 2016).

ATAD2 is a highly conserved bromodomain-containing factor that also harbours an AAA

ATPase domain (Boussouar et al. 2013; Cattaneo et al. 2014). Our early work showed that

Atad2 is predominantly expressed in ES cells, as well as in spermatogenic cells, and is

aberrantly activated in virtually all solid cancers (Caron et al. 2010). Although ATAD2's function

in cancer prompted many investigations (Liu et al. 2022), its function in its physiological

context of expression has remained largely overlooked. We reported the first investigations

of ATAD2's function in ES cells (Morozumi et al. 2016). These studies in ES cells showed that

ATAD2 is recruited into acetylated transcriptionally active chromatin and is critical to keep the

chromatin of ES cells highly dynamic, as illustrated by an ATAD2-dependent high rate of

histone exchange (Morozumi et al. 2016). Subsequent studies conducted by our laboratory, in

ES cells and in *S. pombe*, deciphered the molecular basis of this activity of ATAD2, which relies

on its capacity to ensure the turnover of the chromatin-bound histone chaperone, HIRA and

FACT (Wang et al. 2021). In addition to ES cells, *Atad2* is also highly expressed in male germ

cells, pointing to its potential role in the regulation of HIRA -dependent H3.3 assembly in these

cells, especially, in post-meiotic cells. Indeed, previous published works have highlighted the

important function of H3.3 assembly in meiotic and post-meiotic male genome programming

(van der Heijden et al. 2007; Yuen et al. 2014; Tang et al. 2015; Fontaine et al. 2022). We

therefore hypothesized that ATAD2-regulated HIRA-H3.3 activity would also be a critical

player in the post-meiotic male genome reorganization, as HIRA has been shown to be

essential post-fertilization (Lin et al. 2014) and during oogenesis (Nashun et al. 2015).

Using a lacZ tagged *Atad2* knockout (KO) mouse model, we could show that *Atad2*'s

expression is enhanced in post-meiotic male germ cells, ensuring a proper turnover of histone

H3.3, including on the sex chromosomes. More remarkably, our data demonstrate that ATAD2

controls the appropriate expression of a series of genes whose activity is known to be H3.3-

dependent in post-meiotic cells. Additionally, in the absence of ATAD2, perturbed nucleosome

structural organization leads to a delayed histone-to-PRM-replacement, leading to a defective

global final male genome compaction. Although this defect in mature spermatozoon genome

compaction has minimal effect on male fertility during natural reproduction, it impacts the

success rate of *in vitro* fertilization using spermatozoa from *Atad2* KO mice pointing to a more

fragile genome in these cells.

In summary, our work revealed a novel mechanism for the regulation of chromatin plasticity,

involving the control of histone chaperone actions, particularly that of HIRA, controlling the

proper male germ cell transcriptional regulation, genome compaction and optimal male

fertility.

Results

Atad2 is predominantly expressed in haploid male germ cells

Atad2 is normally expressed in embryonic stem (ES) and spermatogenic cells and remains

silent in the majority of adult tissues (Caron et al. 2010; Morozumi et al. 2016). Therefore, in

addition to ES cells, male germ cells constitute the second cell type where the molecular basis

of the physiological ATAD2 activity could be investigated.

To decipher the function of ATAD2 during spermatogenesis, we used a mouse model

harboring an insertion of a *lacZ* reporter cassette between exons 11 and 12 that enables

splicing and fusion of the *lacZ* transgene with exon 11 of the long and short forms of *Atad2*

(**Fig.1A**) and production of a non-functional ATAD2-lacZ fusion protein (**Fig. 1B**), referred to as

Atad2 KO in this study. Furthermore, we crossed these mice with transgenic mice expressing

ubiquitously Cre recombinase under *CMV* promoter, in order to remove the *neo* cassette and

to delete *Atad2's* exon 12 (**Fig. 1B**). In addition to the long somatic form of ATAD2, ATAD2L,

the mouse testis encodes a short isoform with predominant expression in spermatogenic cells

that we previously named ATAD2S (Caron et al. 2010). The combined LacZ fusion and *Cre*

recombinase-dependent deletion of *Atad2's* exon 12 make both ATAD2L and ATAD2S

indetectable with our homemade antibody, while as expected, a reduced level of both ATAD2

isoforms in Atad2 heterozygous cells can be observed (**Fig. 1C**).

The expression of *lacZ* under the control of the *Atad2* promoter allows one to visualize *Atad2*

gene activity in a stage-specific manner. X-gal staining of seminiferous tubule sections from

Atad2 KO mice showed a remarkably high expression of *Atad2* in post-meiotic spermatogenic

cells (**Fig. 1D).** It is noteworthy that, as expected, no staining was observed in the wild-type

tubules.

ATAD2 controls HIRA accumulation

In our previous work performed in both *S. pombe* and mouse ES cells, we demonstrated the

crucial role for ATAD2 in HIRA-dependent histone dosage as well as in HIRA-chromatin

interaction. Among other discoveries, this work showed that in the absence of ATAD2, HIRA

becomes trapped on chromatin leading to its accumulation in ES cells, as well as in a variety

of cancer cell lines (Wang et al. 2021). Because of this very conserved action of ATAD2 in

controlling the HIRA activity, we reasoned that the ATAD2-dependent HIRA activity should be

particularly active during spermatogenic cell differentiation, especially, in post-meiotic cells,

when ATAD2 expression is highly induced (**Fig. 1D**). To test if the absence of ATAD2 could lead

to the accumulation of HIRA, as we previously reported in other systems, we quantified HIRA

in spermatogenic cells from both wild-type and *Atad2* KO mice. These investigations revealed

an accumulation of HIRA protein in *Atad2* KO spermatogenic cells (**Fig. 2A**), consistent with its

increased association with chromatin (Wang et al. 2021). We have also monitored HIRA

accumulation as a function of spermatogenic cells stages. Due to the strong activation of

Atad2 after meiosis, in spermatids (**Fig.1D**), the ATAD2-dependent accumulation of HIRA was

expected to mostly occur in post-meiotic cells. Accordingly, we also compared HIRA

accumulation in fractionated spermatogenic cell extracts. This experiment demonstrated that,

in the absence of functional ATAD2, the accumulation of HIRA primarily occurs in round

spermatids (**Fig. 2B**). This finding is consistent with the post-meiotic activation of Atad2 and

its role in regulating the turnover of chromatin-bound HIRA.

ATAD2 controls HIRA-dependent H3.3 localization

We speculated that an increased presence of HIRA on chromatin in the absence of ATAD2,

should also be associated with increased amounts of deposited histones. Indeed, we have

demonstrated previously in *S. pombe* that histone gene inactivation could rescue the growth

arrest due to the absence of *Atad2/Abo1.* In parallel, we found that in ES cells, in the absence

of ATAD2, an increased nucleosome assembly occurs, even on the so-called nucleosome-free

region (NFR) (Wang et al. 2021).

Interestingly, visualization of H3.3, the major target of HIRA, by immunofluorescence, in *Atad2*

KO round spermatids, displayed a diffuse, intense staining (**Fig. 3**), which is normally

concentrated on the sex chromosomes (van der Heijden et al. 2007; Fontaine et al. 2022) in

the wild-type cells. Indeed, as expected and previously reported, in the wild-type round

spermatids, H3.3 was mostly observed in a single domain, close to the unique chromocenter

in these cells (**Fig. 3**, left panels). In the absence of ATAD2, in agreement with an increased

residence time of HIRA on chromatin in general and a reduced H3.3 turnover, an increased

level of H3.3 could be observed on the sex chromosome (adjacent to the chromocenter), and

more interestingly, on all over the genome (**Fig. 3**, right panels).

General chromatin accessibility and transcriptional regulation by ATAD2

H3.3 is known to be dynamically incorporated on a group of active genes, especially around

their transcriptional start sites (TSS) (de Dieuleveult et al. 2016). We therefore predicted that

the observed increased H3.3 residence time in the absence of ATAD2 would decrease TSS

regions' accessibility specially those of highly active genes. To verify this prediction, we

purified round spermatid fractions from wild-type and *Atad2 KO* mice testes and measured

the general chromatin accessibility following an ATAC-seq approach. Chromatin accessibility

was then assessed around the TSS of genes, which were categorized into four groups based

on the quartile of their mean expression from three independent transcriptomes of wild-type

round spermatids (**Fig. 4A**), from the top 25% most highly expressed genes (group 1) to the

25% least expressed genes (group 4). As predicted, the absence of ATAD2 was associated with

a significant decrease in TSS region accessibility of the most active genes (**Fig. 4B and 4C**),

supporting a role for ATAD2 in maintaining chromatin dynamics at the most highly active

chromatin regions.

To visualize the genes more sensitive to ATAD2-mediated chromatin remodeling that affect

transcription, we monitored whole genome transcriptional activity during the first

spermatogenic wave at day 20 post-partum (PP), when meiosis is complete, and in the first post-meiotic cells that appear on 22-, 24- and 26-days PP.

In order to definitively confirm our working hypothesis on the ATAD2-HIRA-H3.3 functional interaction, we took advantage of published H3.3-dependent transcriptomic data (Fontaine et al. 2022). Indeed, the authors of this publication identified a series of genes that are mis-regulated in cells with inactivated *H3.3b* gene. Because of the post-meiotic major activation of *Atad2*, we specifically focussed on the gene lists that show a H3.3-dependent deregulated expression in post-meiotic round spermatids.

In these transcriptomes, we specifically followed the expression of the published lists of genes that are downregulated or upregulated in the absence of H3.3 (Fontaine et al. 2022). Very interestingly, we observed that a significant number of genes (37%) that were reported to be down-regulated in the absence of H3.3 (H3.3-activated genes), are upregulated in the absence of ATAD2, particularly at day 26 PP, when post-meiotic cells are present in the testes (**Fig. 5A**, **Fig. S1)**. Gene Set Enrichment Analysis (GSEA) applied to genes that are differentially expressed between wild-type and *Atad2* KO cells at day 26 PP, confirmed that the "H3.3-activated" gene set (corresponding to the published list of H3.3-activated genes) are clearly enriched in *Atad2* KO cells (**Fig. 5A**, right panel).

This regulation can also be observed when we considered the published list of genes that were up-regulated in the absence of H3.3 (H3.3-repressed genes). In this case, the absence of ATAD2 led to the enhancement of the down-regulation of these genes (37%). In other words, the absence of ATAD2 led to an enhancement of H3.3-dependent gene repression (**Fig. 5B, Fig. S1**).

Fontaine and colleagues also reported that sex chromosome-linked genes that normally undergo Meiotic Sex Chromosome Inactivation (MSCI), were up-regulated in the absence of H3.3, indicating that H3.3 is required for their repression. We also investigated these genes and showed that they present an enhancement of repression in the absence of ATAD2 (51%) (**Fig. 5C, Fig. S1**)**.**

These analyses clearly indicate that the absence of ATAD2 enhances H3.3 functions in agreement with increased residence time of H3.3 in nucleosomes. Genes that require H3.3 to be repressed or activated present an enhanced repression or activation, respectively, in the absence of ATAD2.

Overall, these investigations remarkably confirmed a role for ATAD2 in regulating H3.3
function.

ATAD2 is required for controlled histone eviction and PRM assembly

ATAD2's involvement in HIRA-chromatin interaction mobility and its role in H3.3 assembly in
post-meiotic cells, along with its similar function in the control of other histone chaperones
like FACT (Wang et al. 2021), led us to investigate whether ATAD2 participates in controlling
the histone-to-protamine replacement that occurs later in these cells. The histone eviction
process involves a genome-wide histone H4 hyperacetylation (Shiota et al. 2018) and a TP-
mediated PRMs assembly, which directs histone eviction and the final packaging of the male
genome (Barral et al. 2017a). To test the involvement of ATAD2 in spermatogenic cell
chromatin reorganization, we sought to co-detect testis-specific H2B variant, TH2B
(Montellier et al. 2013), along with transition protein 1 (TP1) and protamine 1 (PRM1). TH2B
is first expressed at the commitment of spermatogenic cells into meiotic divisions and
progressively becomes the major H2B species of spermatogenic cell chromatin in the
subsequent stages (Montellier et al. 2013). As expected, the accumulation of TP1 and PRM1
in wild-type cells is associated with the eviction of the majority of histones, reflected by the
mutually exclusive labelling of TH2B and TP1/PRM1 in distinct cells (**Fig. 6A** and **6B**, *Atad2*
wild-type mice). In contrast, in *Atad2* KO cells, TH2B and TP1/PRM1 were co-detected in a
significant number of cells (**Fig. 6A** and **6B**), suggesting that the process of histone eviction is
disturbed. However, in *Atad2* KO cells, the initial delay in histone removal is transient, as
spermatids with high levels of transition protein TP1 and PRM1 eventually appear (**Fig. 6**).
These observations strongly suggest that, in the absence of ATAD2, the process of histone
eviction is also slowed down or delayed, allowing for the simultaneous detection of histones
and histone-replacing proteins in the same cells.

Given that NUT-CBP/p300-dependent H4 hyperacetylation, along with the expression of
H2A.L.2, TPs, and PRMs, is essential for histone removal, we also investigated whether the
observed delay in histone removal is due to defective histone acetylation, particularly at H4K5,
as well as disrupted cell type-specific expression and localization of H2A.L.2, TPs, and PRMs.
Immunohistochemical detection of H4K5ac, H2A.L.2, TP2, PRM1, and PRM2 revealed a similar

pattern and cell type-specific expression of the investigated proteins when comparing *Atad2* KO and wild-type post-meiotic cells. (**Fig. S2**).

These data support the idea that the perturbed modified flexibility of histone/TP/PRM chaperones does not grossly affect the other molecular events involved in the histone-to-PRM replacement.

Perturbed pre-PRM2 processing in the absence of ATAD2

We have shown previously that the H2A.L.2-dependent process of incorporation of TPs into nucleosomes is required for the proper processing of pre-PRM2. The abnormal overlapping staining of histones, TP and protamines observed by immunofluorescence in *Atad2* KO cells, suggested that TPs may not effectively mediate the pre-PRM processing. To test this hypothesis, we took advantage of our specific pre-PRM2 antibody (Rezaei-Gazik et al. 2022) to investigate pre-PRM2 maturation and assembly. **Figure 7A** shows that although pre-PRM2 is present in both wild-type and *Atad2* KO elongating spermatids, in the absence of ATAD2, pre-PRM2 tends to form dense aggregates as well as fragmented genomic domains. This phenomenon is also associated with an increased accumulation of pre-PRM2 protein level in testis extracts of *Atad2* KO spermatogenic cells compared to wild-type cells as revealed by ELISA and immunoblotting (**Fig. 7B, Fig. S3**).

These data provide strong evidence that ATAD2 is required to streamline the activity of histone chaperones as well as other factors involved in histone exchange and TP/PRM assembly.

Defective mature spermatozoa genome organization and male fertility *in vitro*

We show that although the *Atad2* KO post-meiotic cells' defective chromatin organization does not affect the testis histology (**Fig. S2** and **data not shown**), it impairs testis weight and sperm count, which are significantly reduced in *Atad2* KO mice (**Fig. S4**)).

As expected, the impact of ATAD2 on HIRA function and TP/PRM assembly also has an impact on proper sperm genome compaction. Indeed, mature sperm genome decompaction test revealed a significant difference in the ability of the genome to resist genome decompaction conditions between wild-type *Atad2* and *Atad2* KO spermatozoa (**Fig. 8A; Fig. S5**). Given that spermatozoa with defective genome compaction are known to have poor success rate in *in vitro* fertilization (Bashiri et al. 2021), we expected that spermatozoa from *Atad2* KO mice

would perform worse in *in vitro* fertilization assays compared to wild-type spermatozoa. **Figure 8B** shows that this is, in fact, the case. However, surprisingly, fertility tests carried out on sexually mature adult males revealed that *Atad2* KO male mice are fertile, with only a statistically non-significant effect observed on their ability to foster pups (**Fig. 8C**), indicating that *in vivo* fertilization could buffer a relatively large series of sperm genome compaction defaults.

Discussion

Taking advantage of the very conserved structural features of ATAD2 and relying on the power of yeast genetics, we previously discovered that ATAD2's function is mostly to control major histone chaperone functions. Using a screen of the Abo1/ATAD2-dependent arrest of *S. pombe* cell growth, we found that impressively, 11 of the 14-growth arrest suppressor strains that we isolated, impacted histone chaperones or histones themselves, with 8 distinct inactivating mutations in the histone chaperone HIRA complex. Furthermore, investigation of ATAD2-depleted mouse ES cells showed that HIRA accumulates and remains bound to nucleosomes, leading to nucleosome overloading around active gene TSSs and on the normally nucleosome-free regions (NFRs) of these genes.

Here, we showed that *Atad2* is highly expressed in post-meiotic spermatogenic cells and that in its absence, as in ES cells and in different cancer cells, HIRA accumulates. Our new data suggest that in the absence of ATAD2 activity, HIRA residency time on chromatin increases leading to decreased turnover and overloading of its substrate, the histone variant H3.3 (**Fig. 9**). This replication-independent histone has been first shown to assemble on the transcriptionally inactive meiotic cell sex chromosomes by Peter de Boer's laboratory (van der Heijden et al. 2007). Subsequent studies showed that H3.3 remained enriched on the sex chromosomes also in early post-meiotic cells, usually localized close to the unique large chromocenter in these cells (Fontaine et al. 2022).

In agreement with the proposed role for ATAD2 in the control of HIRA-dependent dynamics of H3.3, we observed a more intense labelling of H3.3 on the sex bodies in round spermatids and impressively, an increase in the genomic background detection of H3.3 in these cells. The role of ATAD2 in the HIRA/H3.3 functions is further supported by comparing the reported H3.3-dependent transcriptome with the ATAD2-dependent transcriptome identified here.

Indeed, in the absence of ATAD2, we observed an enhancement of H3.3-dependent gene

regulatory events (activation or repression), in agreement with an increased H3.3 presence

and hence function, in the chromatin of *Atad2* KO post-meiotic cells.

These investigations on ATAD2's role in post-meiotic spermatogenic cells built upon our

previous findings in *S. pombe* and ES cells, uncover an additional level of regulation of histone

turnover, which relies on the modulation of histone chaperone–chromatin interactions.

Previously, A. Groth's laboratory discovered that the histone chaperone DNAJC9 recruits the

ATPase HSP70 to provide the necessary energy to release the chaperone from chromatin

(Hammond et al. 2021). Considering our data on the role of the ATPase ATAD2 with respect

to HIRA and that of HSP70 with respect to DNAJC9, it is tempting to propose that ATP energy

is required for the proper function of histone chaperones in controlling histone turnover.

Interestingly, very recently, Song laboratory demonstrated that the *S. pombe* Abo1/ATAD2, is

required for ATP-dependent dissociation of the histone chaperone FACT from chromatin,

particularly at TSSs (Jang et al. 2024). In line with this conclusion, ChIP-seq mapping of FACT

in ATAD2-depleted ES cells revealed an accumulation of nucleosome-bound FACT at the TSSs

of active genes (Wang et al. 2021), supporting the idea that ATAD2 not only regulates the

dynamics of histone-bound HIRA but also that of other histone chaperones such as FACT.

In addition to histones, many chromatin-bound regulatory factors also need to dynamically

interact with chromatin and hence ATAD2 could also be involved in mediating the dynamic

interactions of such chromatin-bound factors. In agreement with this hypothesis, using

proteomics of ATAD2-bound nucleosomes in ES cells, we previously co-purified various

chromatin bound factors (Morozumi et al. 2016). Other AAA ATPase factors, such as p97/VCP,

have also been implicated in the remodeling of chromatin-bound factors. Indeed, the AAA

ATPase factor, p97/VCP, has been shown to remove H3 variant CENP-A (van den Berg et al.

2023), the chromatin trapped PARP1 (Krastev et al. 2022), trapped Ku70/80 (van den Boom et

al. 2016), DNA damage sensors, DDB2 and XPC (Puumalainen et al. 2014), RNA pol II (Lafon et

al. 2015), and other chromatin interacting factors. Similarly, ATAD2 AAA ATPase very likely

acts on other chromatin-bound factors than histone chaperones. However, in contrast to

p97/VCP, and due to its bromodomain, ATAD2 would preferentially act on H4K5 acetylated

chromatin regions and would specifically ensure the dynamics of H4 acetylated chromatin-

bound factors. Based on the data presented here, we can also include, a role for ATAD2 to

streamline steps involving the H2A.L.2-dependent recruitment of TPs and protamine assembly

and the final histone displacement in spermatids. The chaperones responsible for TP/PRM

assembly are not yet identified, but it is possible that ATAD2 plays a role in dissociating these

chaperones from their substrate proteins after the assembly process. That is why in our *Atad2*

KO spermatids, we observed an overlap between TP/PRM and histones. This could be due to

the delayed removal of chaperones bound to TPs and PRMs and the inability of pre-PRM2 to

be processed and hence of PRMs to efficiently replace histones. Since histone replacement

still occurs in the absence of ATAD2, it suggests that other AAA ATPases, such as p97/VCP,

may compensate for its function. However, under these conditions, although most of the

histone replacement takes place, our functional tests such as *in vitro* chromatin decompaction

and *in vitro* fertilization, strongly suggest that mature spermatozoa genome organization is

not fully complete and the spermatozoa nuclei remain fragile and sensitive to external

assaults.

In summary, the functional analysis of ATAD2 during its physiological expression in the post-

meiotic phases of spermatogenesis supports our previous conclusions regarding its role in ES

cells and its conserved activity in *S. pombe*. Together, these studies reveal a novel layer of

regulation in histone turnover and chromatin dynamics, based on the control of histone-

bound histone chaperone release.

Materials and Methods

Animal care and breading

Mice were housed at the Grenoble High Technology Animal Facility (PHTA). Mice were euthanized following a procedure approved by the official ethic committee of the University Grenoble Alpes (COMETH, C2EA-12). The investigators directly involved in care and breading of mice had an official animal-handling authorization obtained after 2 weeks of intensive training and a final formal evaluation.

Generation of *Atad2*-KO mice

Atad2 KO mice were obtained from the International Knockout Mouse Consortium, the Welcome Trust Sanger Institute, UK. The 'knockout-first' allele (tm1a) contains an IRES: *lacZ* trapping cassette and a floxed human β-*actin* promoter-driven *neo* cassette inserted into the intron 12 of the *Atad2* gene (Skarnes et al. 2011). We created $Atad2^{\Delta Neo\text{-}Exon\,12}$ mice by crossing mice bearing the *tm1a* allele with mice expressing Cre recombinase under the control of a ubiquitous human promoter CMV. This cross results in an $Atad2^{LacZ\Delta Neo\text{-}Exon12}$ knockout mouse with the first 11 exons of *Atad2* being fused to β-gal, leading to a non-functional ATAD2 protein depleted of its bromodomain and AAA ATPase domain.

For *Atad2* KO mice genotyping, multiplex PCR was used to detect LacZ cassette producing a 393-bp WT band and a 293-bp mutant band using primers *Atad2*_84231_F, TATCCAACAAGCCTGAGCCC, *Atad2*_84231_R, CAACTGGAGCTGGGTCTTCC and Cas-R, TCGTGGTATCGTTATGCGCC.

Fertility test

Fertility tests were carried out with sexually mature 3-month-old males. Five WT males and five *Atad2* KO males were crossed with two C57BL6 females each, during 3 months, and the litter sizes were monitored during this period.

***In vitro* Fertilization**

Eggs were obtained from 6- to 7-week-old superovulated C57/BL6J females, stimulated by a first injection of 7.5 UI of PMSG (Pregnant Mare Serum Gonadotropin), followed 48 hours later by a second injection of 7.5 UI of HCG (Human Chorionic Gonadotropin). Fourteen hours later,

females were euthanized by cervical dislocation and cumulus-oocytes complexes (COCs)
released from the ampulla were collected in 500 μL M2 medium. Spermatozoa from wild type
and *Atad2* KO males were harvested by dilaceration of the cauda epididymitis and allowed to
swim in 1 mL M2 medium for 10 min at 37°C. Two hundred microliters of sperm were
transferred into 800 μL M16 /2% BSA and capacitated for 80 min at 37°C /5% CO_2. Finally,
300,000 sperm were simultaneously added to the COCs and incubated in M16 medium at 37°C
/5% CO_2. After 4 hours of incubation, unbound spermatozoa were removed by 3 successive
washes with 500 μL of M16. Twenty-four hours after fertilization, unfertilized eggs and two-
cell embryos (as an indication of successful fertilization), were scored.
405

Protein extraction and Immunoblotting
Proteins were extracted in 8M Urea, sonicated at 150J with a probe sonicator, mixed with 4x
Laemli loading buffer and heated for 5 minutes at 95 °C. Protein samples were loaded on 4-
12% Bis-Tris gels and migrated at 150 V for 1h30. The transfer step was realized using a
nitrocellulose membrane with the Bio-Rad Transfer Turbo machine. Nitrocellulose membrane
was blocked in 5% milk PBS-Tween 0.1% solution, washed 3 times for 10 minutes with PBS-
Tween 0.1% solution and incubated overnight at 4°C with specific antibodies listed in Table
S1. Membranes were washed 3 times for 10 minutes with PBS-Tween 0.1% solution and were
incubated with HRP secondary antibodies at 1/10000 dilution for 1h at RT under agitation.
Revelation was performed using ECL substrates and visualization was performed with the
Vilber Fusion FX Chemiluminescence imaging system (Vilber).
417

ELISA
Urea protein extracts were diluted in PBS and used to coat 96-well round-bottom plates,
followed by overnight incubation at 4 °C. Coating solution was removed and wells were
washed 3 times with washing buffer (BSA 1%, PBS-T 0,1%). Wells were blocked with blocking
buffer (5% BSA in PBS-T 0,1%) for 2 hours at room temperature under agitation. Blocking
buffer was removed and wells were washed 2 times with washing buffer. Primary antibodies
for pre-PRM2 (1:1000) and tubulin (1:5000) were diluted into blocking buffer, pipetted into
their corresponding wells and incubated overnight at 4°C under mild agitation. Wells were
washed 4 times with washing buffer and subsequent incubation with secondary HRP
antibodies, diluted 1:5000 in blocking buffer, was performed for 1 hour at room temperature

under mild agitation. Wells were washed 4 times with washing buffer, and TMB ELISA
Substrate was added to wells and incubated until the color developed, at which time Stop
Solution was added to the wells to stop the reaction. The signal was quantified using a plate
reader at 450nm.
432

X-Gal staining

Male mice were euthanized by cervical dislocation. Testes were dissected out, drilled in 3
points and incubated overnight at 4℃ under agitation in fixative solution (0.5% glutaraldehyde,
2 mM $MgCl_2$ and 5 mM EGTA in PBS). After incubation until equilibration in an equilibration
buffer (30% sucrose, 2 mM $MgCl_2$ and 5mM EGTA in PBS), testes were frozen individually in
OCT on dry ice. Frozen blocks were sectioned at 10 µm slices on Superfrost^{TM} slides and let
dry overnight. Slides were fixed with fixation buffer (0.2% glutaraldehyde, 100 mM $MgCl_2$ and
5 mM EGTA in PBS) for 10 min, washed with washing buffer (PBS, 0.02% NP-40, 0.01% Sodium
deoxycholate and 2 mM $MgCl_2$) for 5min and incubated three times in 50% Ethanol for 5 min
each. Then the slides were stained with staining solution (2 mM $MgCl_2$, 0.02% NP-40, 0.01%Na
Doc, 5 mM potassium ferricyanide, 10 mM potassium ferrocyanide and 0.5 mg/ml X-gal) in a
wet chamber for several hours at 37℃. The slides were then washed with washing buffer for
5 min and with distilled water, and a counterstaining was performed with hematoxylin. Finally,
the slides were washed with running tap water for several minutes and mounted with Dako
fluorescent mounting medium.
448

Immunofluorescence staining of germ cells

Staged seminiferous tubules were prepared as detailed in Gaucher et al. 2012. In order to
permeabilize the cells, the slides were placed in 0.5% saponin, 0.2% Triton X-100, and 1 × PBS
at RT for 15 min and washed for 5 min with PBS. The preparations were then incubated in 5%
milk, 0.2% PBS-Tween blocking buffer, at RT for 30 min under agitation. The primary
antibodies were diluted in 1% dry milk, 0.2% PBS-Tween buffer. The slides were incubated
overnight with this diluted solution of the primary antibodies in a humidified chamber at 4℃
and then washed three times for 5 min each in the antibodies' dilution buffer. The secondary
antibodies were diluted at 1:500 in the same buffer and incubated in a humidified chamber
for 30 min at 37℃, and then washed as for the primary antibodies. The DNA was
counterstained with Hoechst, and the slides were mounted using Dako fluorescent mounting

medium.

Sperm head decompaction test

Spermatozoa from wild type and *Atad2* KO males were harvested by dilaceration of the cauda

epididymis and allowed to swim in 1 mL M2 medium for 10 min at 37°C. The sperm-containing

suspension was transferred into a 1.5 ml Eppendorf tube and centrifuged at 3000 rpm for 8

min in 4°C. The sperm pellets were resuspended in PBS and dropped onto glass slides to

spread and dry at RT. The decompaction mix (50 mM DTT, 400 IU/ml heparin and 0.2% Triton

in PBS) was added to each dried sperm drop for 2 min and the decompaction was stopped by

placing the slides in 4% PFA for 15 min. Slides were washed with PBS for 5 minutes, stained

with DAPI, and mounted using Dako fluorescent mounting medium. Fluorescent images were

captured by a Zeiss inverted microscope under a 63X numeric aperture oil-immersion lens

(Carl Zeiss). The size of spermatozoa heads was calculated with ImageJ software.

Spermatogenic cell fraction purification

Spermatogenic cell fraction purification was performed as previously described (Shiota et al.

2018).

ATAC-seq and data analysis

Library preparation

Frozen nuclei from four biologically independent purified spermatid fractions were

centrifuged at 3000 g for 5 minutes and then washed in a 1% BSA-PBS solution. Libraries were

then prepared from 100,000 nuclei using the ATAC-seq Kit (Catalog No. 53150) from Active

Motif, following the manufacturer's protocol (version B9). After quality control (QC) with

Qubit and Fragment Analyzer, libraries were subsequently purified using AMPure XP

(Beckman Coulter, Catalog No. A63881) beads with a double size selection of 100-800 bp

fragments. After a second round of library QC (Qubit and Fragment Analyzer) we performed a

paired-end sequencing on a NextSeq 2000 device using a 200-cycle P2 flow cell.

Trimming

The raw fastq files were processed by 5' end trimming, keeping 30bp-length fragments, using fastx_trimmer [http://hannonlab.cshl.edu/fastx_toolkit/. Accessed 28 Feb. 2022.], with options -l 30 -Q33.

Alignment

The trimmed fastq files were aligned on the UCSC Mus_musculus mm10 genome using the Bowtie2 aligner (Langmead and Salzberg 2012), with options –end-to-end, –no-mixed, –no-discordant.

Normalization

The BigWig files (.bw) containing normalized integrated aligned read count signals were obtained from by normalizing and smoothing bam files using bamCoverage (from deepTools suite (Ramirez et al. 2014) with options: –binSize 4 –minMappingQuality 30 –normalizeUsing RPM.

Heatmap

The normalized ChIP signal were converted into a 10bp bin matrix of the signal 500b upstream and 1500b downstream protein-coding genes' TSS, using computeMatrix (from deepTools suite (Ramirez et al. 2014), with options reference-point -R mm10_pc.bed –referencePoint TSS –binSize 10 –beforeRegionStartLength 500 –afterRegionStartLength 1500

Heatmaps were generated using plotHeatmap (deepTools suite (Ramirez et al. 2014).

RNA-seq data processing

Wild-type (WT) and *Atad2* knockout (KO) homozygous males were sacrificed by cervical dislocation on days 20-, 22-, 24-, and 26- postpartum (PP), after which their testes were harvested. RNA was extracted using TRIzol™ reagent, followed by DNA digestion with DNase I. The resulting RNA samples were used for RNA sequencing. For both WT and *Atad2* KO mice, the biological replicates included 3 samples for D20, 4 samples for D22, 2 samples for D24 and 2 samples for D26.

The raw sequencing data (FASTQ files) were aligned to the UCSC mm10 mouse genome using

the STAR software (version 2.7.11b) (Dobin et al. 2013) to generate BAM files. Raw read counts

were extracted from the BAM files using HTSeq (version 2.0.5) (Anders et al. 2015) with the

following parameters: -t exon, -f bam, -r pos, --stranded=reverse, -m intersection-strict, and -

-nonunique none.

Normalization of read counts was performed using the DESeq2 R package (version 1.22.2)

(Anders and Huber 2010; Love et al. 2014). Counts were transformed into DESeq2-normalized

values and subsequently log-transformed using the formula log2(1 + DESeq2-normalized

counts). Differential expression analysis was conducted with the R package "SARTools" (Varet

et al. 2016).

Purified RNA from fractionated round spermatids were also sequenced to establish gene

transcriptional group quartile shown in Fig. 4. RNA purification and sequencing were

performed as described above.

Quantification and statistical analysis

Statistical analysis was performed with R and GraphPad Prism 8. The results of fertility tests

were analyzed by Student t-test. Pairwise Student t-test was used to analyze testis weights

and spermatozoa counts of same-aged mice and ratio pairwise Student t-test was used for the

IVF efficiency results. The results of sperm decompaction test were analyzed by two-way

ANOVA and Tukey's HSD post two-way ANOVA test was used to determine the significance

between groups. *P-values* < 0.05 were considered statistically significant: *$p < 0.05$, **$p < 0.01$,

***$p < 0.001$.

Gene Set Enrichment Analysis (GSEA)

Differential expression analysis and the corresponding GSEA analysis (Mootha et al. 2003;

Subramanian et al. 2005) were performed to compare *Atad2* KO versus *Atad2* wild-type

samples at days 20, 22, 24, and 26 PP. For the GSEA analysis, we used the Python package

"gseapy" (Fang et al. 2023) available at https://www.gsea-msigdb.org/gsea.

The GSEA was carried out on three custom gene sets. The first gene set, referred to as "sex

chromosome-linked genes", corresponds to genes located on the X and Y chromosomes. The

remaining two gene sets include genes identified as upregulated or downregulated in the

absence of histone H3.3 by Fontaine and colleagues (Fontaine et al. 2022). These gene sets

were designated as "post-meiotic H3.3-activated genes" and "post-meiotic H3.3-repressed

genes", respectively.

Data availability

The transcriptomic data generated in this study, corresponding to testes collected at different

days post-partum, have been deposited on GEO under the accession N°. GSE277943.

To review GEO accession GSE277943 go to :

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE277943

Enter token wrszesqgjfkxxmx into the box.

Transcriptomic data from fractionated round spermatids (three biological replicates, Fig. 4A)

were deposited on GEO accession GSE284749. To review GEO accession GSE277943 go to:

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE284749.

Enter token uzmvcaygpnydrgt into the box

ATAC-seq data generated in this study were deposited on GEO under the accession N°

GSE283879. To review GEO accession GSE283879, go to :

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE283879

Enter token ozazwsmoxjofzkb into the box

Manuscript preparation

Parts of the text have been refined, with the help of free online tools (ChatGPT and DeepL).

COMPETING INTEREST STATEMENT

The authors declare that they have no conflict of interest.

ACKNOWLEDGEMENTS

This work was supported by ANR EpiSperm 4 (ANR-19-CE12- 0014), ANR Episperm 5 (ANR-23-

CE12-0028), ANR NME-CoA (ANR-23-CE14-0050) and the Cancer ITMO [Multi-Organization

Thematic Institute of the French Alliance for Life Sciences and Health (AVIESAN)] MIC program

to SK and SR. AL is a recipient of a 4th year PhD fellowship by Foundation pour la Recherche

Médicale (FRM). High throughput sequencing was performed at the TGML Platform,

supported by grants from Inserm, GIS IBiSA, Aix-Marseille Université. We thank Dr. Alexandra
Varga for X-gal staining of testis sections. The authors thank Dr Kelly Matmati, Stroke Program
Director, Rochester General Hospital, Connecticut, USA, for her critical reading of the
manuscript.
580

AUTHOR CONTRIBUTIONS

AL: experimentation, data interpretation, writing and preparation of the manuscript, FB:
supervision, experimentation, data interpretation, writing and preparation of the manuscript,
management of acquisition of *Atad2* KO mouse model. SB: experimentation and the maintain
of *Atad2* mouse model, EL and CA *in vitro* fertilization essays. FC and EBF transcriptomic data
analyses. DePu and CG: transcriptomic data generation, SR: founding acquisition and
transcriptomic analysis. DaPe and AV: experimentation and data generation and analyses. SK:
conception, founding acquisition and supervision, and coordination of the whole project and
writing of the manuscript.

  Figure legends

  Liakopoulou_Fig1

Figure 1: *Atad2* is highly expressed in post-meiotic male germ cells.

(**A**) The exon/intron organization of *Atad2* gene and the two ATAD2-encoding transcriptional units (encoding for ATAD2-long and ATAD2-short) are indicated (Dyer et al. 2025). The red arrows indicate the position of insertion of the *Lac-Z-Lox-Neo-Lox* cassette.

(**B**) Gene targeting strategy for the generation of the *Atad2* KO allele is represented. Exon number and relative position, selection gene cassettes, LoxP sites and Frt sites are indicated.

Crossing mice bearing this construct with *CMV-Cre* mice resulted in the generation of the *Atad2 LacZ*$^{\Delta Neo\text{-}Exon12}$ allele, which can be verified through PCR amplification of the genomic DNA (shown on the ethidium bromide-stained gel). The expected amplified bands for each genotype allowing the detection of both wild-type and KO alleles of *Atad2* are indicated. E = Exon.

(**C**) Total protein extracts from wild-type (WT), heterozygous (HET) and *Atad2* knockout (KO) mice testes were probed with anti-ATAD2 and anti-H3 antibodies as indicated. Four different mice were used for this experiment.

(**D**) Seminiferous tubules sections from wild-type and *Atad2* KO mice were stained with X-gal to visualize *lacZ* gene expression and β -galactosidase activity under the endogenous *Atad2* gene promoter. Testes from three different wild-type and *Atad2* KO mice were used to generate the represented sections. Scale bar: 200 µm

Figure 2: *Atad2* gene inactivation leads to the accumulation of HIRA

(**A**) Total protein extracts from the testes of five wild-type (WT) and five *Atad2* KO mice were used to detect HIRA and H3 by immunoblotting. The panel shows a representative immunoblotting after HIRA detection. The histogram in the right panel shows the quantification of HIRA immunoblotting signal normalized to H3. The mean ± standard deviation was 0.86± 0.21 for wild type and 1.15± 0.06 for *Atad2* KO samples. The p value was 0.02 for unpaired Student t-test.

(**B**) Protein extracts from fractionated spermatogenic cells from wild-type (*Atad2* WT) and knockout (*Atad2* KO) mice (pool of testes from 3 individuals per genotype) were used to detect HIRA and H3 by immunoblotting.

Figure 3: Enhanced H3.3 detection in *Atad2* KO post-meiotic cells

A specific antibody against H3.3 was used to detect H3.3 in round spermatid cells from wild-
type (*Atad2* WT) and knockout (*Atad2* KO*)* mice. Round spermatids are recognizable by their
distinctive Hoechst - bright chromocenter (upper panels). The lower panels represent a higher
magnification of selected cells that are displayed at lower magnification in the upper panels.
The images show H3.3 labelling and Hoechst staining alone or the merged images of these
staining as indicated. The nuclear boundaries defined by Hoechst staining, are outlined by
white lines.

Fig. 4: ATAD2 controls active gene TSS accessibility

A) Gene selection according to expression in wild-type round spermatids. The plot displays the mean expression levels (y-axis, labeled as "mean expression level") versus the corresponding standard deviations (x-axis), both calculated from three independent biological replicates of isolated round spermatids. The standard deviation reflects the variability of gene expression across biological replicates. Genes were grouped into four categories (Group1: blue, Group 2: cyan, Group 3: green, Group 4: orange) according to the quartile of their mean expression. For Group 4, all genes have no detectable expression, resulting in a mean expression of zero and a standard deviation of zero; consequently, the 5611 genes in this group are represented by a single overlapping point (red open circle) at the origin.

B) Heatmap on ATAC-seq signal. The heatmap shows the ATAC-seq signal of each group (Group 1 in blue, Group 2 in cyan, Group 3 in green and Group 4 in orange) in rows, for 4 *Atad2* WT and 4 *Atad2* KO samples in column.

C) Profiles according to selected genes. The mean profiles are displayed together with their variability (± 2 s.e.m.) across the four replicates for both *Atad2* WT (blue) and *Atad2* KO (red). For groups 1, 2, and 3, the envelopes of the curves remain clearly separated around the peak, indicating a consistent difference in signal between the two conditions. In contrast, group 4 does not present a strong signal and, accordingly, no marked difference is observed between *Atad2* WT and KO in this group.

Figure 5: ATAD2 enhances H3.3 gene regulatory functions.

(A) Expression profiles of post-meiotic H3.3-activated genes. The heatmap (left panel) displays the normalized expression levels of genes identified by Fontaine and colleagues as upregulated in the absence of histone H3.3 (Fontaine et al. 2022) for *Atad2* WT (WT) and *Atad2* KO (KO) samples at days 20, 22, 24, and 26 PP (D20 to D26). The colour scale represents

the z-score of log-transformed DESeq2-normalized counts. The middle panel box plots display, pooled, normalized expression levels, aggregated across replicates and genes, for each condition (WT and KO) and each time point (D20 to D26). Statistical significance between WT and KO conditions was determined using a two-sided t-test, with p-values indicated as follows: * for p-value<0.05, ** for p-value<0.01 and *** for p-value<0.001. The right panel shows the results of gene set enrichment analysis (GSEA), which assesses whether predefined groups of genes show statistically significant differences between conditions. Here, the post-meiotic H3.3-activated genes set, identified by Fontaine et al. (2022), is significantly enriched in *Atad2* KO compared with WT samples at day 26 (p < 0.05, FDR < 0.25). Coloured vertical bars indicate the "leading edge" genes (i.e., those contributing most to the enrichment signal), located before the point of maximum enrichment score. **(B)** As shown in (A) but for the "post-meiotic H3.3-repressed genes" gene set. **(C)** As shown in (A) but for the " sex chromosome-linked genes " gene set.

A

B

Figure 6: ATAD2 is required for efficient histone-to-protamine replacement.

(**A**) Spermatogenic cell preparations from wild-type and *Atad2* KO mice were stained with

antibodies against TH2B (in green) and transition protein 1 (TP1) or protamine 1 (PRM1) (in

red). A yellow signal indicates the co-detection of TH2B along with TP1 or PRM1. The indicated

fields are shown at higher magnification in panel "B". Scale bar: 10 μm.

(**B**) Fields shown in panel "A" are ordered to illustrate the transition from the histone

associated genome to its packaging by protamines, considering both wild-type and *Atad2* KO

spermatogenic cells.

Figure 7: ATAD2 is required for proper pre-protamine 2 processing and protamine assembly

(**A**) A specific antibody raised against the processed part of pre-PRM2 (Rezaei-Gazik et al. 2022) was used to detect pre-PRM2 expression and localization in tubular elongating spermatid cells. The lower panels represent the magnification of selected cells.

(**B**) The amount of pre-PRM2 was detected using soluble extracts from testis of wild-type and *Atad2* KO mice using ELISA. Histograms show the normalized values (pre-PRM2/tubulin) of 4 measurements of independent sedimentation samples of Elongating/Condensing fractions. Mean ± Standard Deviation is 1.6 ± 0.2 and 2.1 ± 0.1 for wild type and *Atad2* KO samples, respectively. The p-value obtained using an unpaired Student's *t*-test was 0.0158.

Figure 8: ATAD2s' function in sperm genome compaction and fertility parameters

(**A**) Epididymal sperm cells were isolated by the swim-up method from wild-type or *Atad2* KO

mice and underwent a decompaction test and the sizes of the sperm heads were measured

and represented as box plots (*n* = 75 sperm heads / box plot). The value ranges (mean) were

10.4 ~ 19.3 (16.0) for wild type sperms without decompaction; 11.4 ~ 20.1 (16.4) for *Atad2* KO

sperms without decompaction; 33.4 ~ 61.8 (48.5) for wild type sperms after decompaction;

45.0 ~ 78.9 (62.6) *Atad2* KO sperms after decompaction. Tukey's HSD post two-way ANOVA

test was used. ****$p < 0.001$.

(**B**) Pools of oocytes from C57BL6 females were obtained and used for *in vitro* fertilization (IVF)

with spermatozoa from WT or *Atad2* KO males (the experiment was repeated 4 times

independently). The fertilization success rate is indicated as a percentage of oocytes giving

rise to stage 2 embryos. Mean ± Standard Deviation is 25.5 ± 14 and 12 ± 5.2 for wild type and

Atad2 KO experiments respectively and the p = 0.002 for ratio paired Student t-test.

(**C**) Five wild-type and five *Atad2* KO male mice were each mated with two C57BL6 females,

the resulting litter sizes were recorded and represented as box plots (*n* = 28 and 30,

respectively). The average numbers of pups are respectively 8.0 ± 2.1 and 7.1 ± 2.5 and the *p-*

value of Student t-test = 0.1.

 Liakopoulou_Fig9

Figure 9: A model summarizing the role of ATAD2 in regulating chromatin-bound HIRA dynamics (along with other histone and non-histone protein chaperones) during the post-meiotic stages of mouse spermatogenesis. Following the deposition of H3.3 by HIRA onto chromatin, ATAD2 facilitates the release of HIRA. In the absence of ATAD2, both HIRA and the newly incorporated H3.3 persist longer in nucleosomes. The scheme was created using BioRender.

REFERENCES

Anders S, Huber W. 2010. Differential expression analysis for sequence count data. *Genome*
Biol **11**: R106.
Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-throughput
sequencing data. *Bioinformatics* **31**: 166-169.
Arevalo L, Esther Merges G, Schneider S, Schorle H. 2022. Protamines: lessons learned from
mouse models. *Reproduction* **164**: R57-R74.
Bao J, Bedford MT. 2016. Epigenetic regulation of the histone-to-protamine transition during
spermiogenesis. *Reproduction* **151**: R55-70.
Barral S, Morozumi Y, Hoghoughi N, Rousseaux S, Khochbin S. 2017a. [The mystery of histone
disappearance during spermatogenesis]. *Med Sci (Paris)* **33**: 588-590.
Barral S, Morozumi Y, Tanaka H, Montellier E, Govin J, de Dieuleveult M, Charbonnier G, Coute
Y, Puthier D, Buchou T et al. 2017b. Histone Variant H2A.L.2 Guides Transition Protein-
Dependent Protamine Assembly in Male Germ Cells. *Mol Cell* **66**: 89-101 e108.
Bashiri Z, Amidi F, Amiri I, Zandieh Z, Maki CB, Mohammadi F, Amiri S, Koruji M. 2021. Male
Factors: the Role of Sperm in Preimplantation Embryo Quality. *Reprod Sci* **28**: 1788-
1811.
Boussouar F, Jamshidikia M, Morozumi Y, Rousseaux S, Khochbin S. 2013. Malignant genome
reprogramming by ATAD2. *Biochim Biophys Acta* **1829**: 1010-1014.
Caron C, Lestrat C, Marsal S, Escoffier E, Curtet S, Virolle V, Barbry P, Debernardi A, Brambilla
C, Brambilla E et al. 2010. Functional characterization of ATAD2 as a new cancer/testis
factor and a predictor of poor prognosis in breast and lung cancers. *Oncogene* **29**:
5171-5181.
Cattaneo M, Morozumi Y, Perazza D, Boussouar F, Jamshidikia M, Rousseaux S, Verdel A,
Khochbin S. 2014. Lessons from yeast on emerging roles of the ATAD2 protein family
in gene regulation and genome organization. *Mol Cells* **37**: 851-856.
de Dieuleveult M, Yen K, Hmitou I, Depaux A, Boussouar F, Bou Dargham D, Jounier S,
Humbertclaude H, Ribierre F, Baulard C et al. 2016. Genome-wide nucleosome
specificity and function of chromatin remodellers in ES cells. *Nature* **530**: 113-116.
Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.
2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15-21.
Dyer SC, Austine-Orimoloye O, Azov AG, Barba M, Barnes I, Barrera-Enriquez VP, Becker A,
Bennett R, Beracochea M, Berry A et al. 2025. Ensembl 2025. *Nucleic Acids Res* **53**:
D948-D957.
Fang Z, Liu X, Peltz G. 2023. GSEApy: a comprehensive package for performing gene set
enrichment analysis in Python. *Bioinformatics* **39**.
Fontaine E, Papin C, Martinez G, Le Gras S, Nahed RA, Hery P, Buchou T, Ouararhni K, Favier
B, Gautier T et al. 2022. Dual role of histone variant H3.3B in spermatogenesis: positive
regulation of piRNA transcription and implication in X-chromosome inactivation.
Nucleic Acids Res **50**: 7350-7366.
Gaucher J, Reynoird N, Montellier E, Boussouar F, Rousseaux S, Khochbin S. 2010. From
meiosis to postmeiotic events: the secrets of histone disappearance. *FEBS J* **277**: 599-
604.
Gaucher JF, Mauge C, Didry D, Guichard B, Renault L, Carlier MF. 2012. Interactions of isolated
C-terminal fragments of neural Wiskott-Aldrich syndrome protein (N-WASP) with actin
and Arp2/3 complex. *J Biol Chem* **287**: 34646-34659.

Goudarzi A, Zhang D, Huang H, Barral S, Kwon OK, Qi S, Tang Z, Buchou T, Vitte AL, He T et al.
2016. Dynamic Competing Histone H4 K5K8 Acetylation and Butyrylation Are
Hallmarks of Highly Active Gene Promoters. *Mol Cell* **62**: 169-180.
Hammond CM, Bao H, Hendriks IA, Carraro M, Garcia-Nieto A, Liu Y, Reveron-Gomez N, Spanos
C, Chen L, Rappsilber J et al. 2021. DNAJC9 integrates heat shock molecular chaperones
into the histone chaperone network. *Mol Cell* **81**: 2533-2548 e2539.
Hoghoughi N, Barral S, Vargas A, Rousseaux S, Khochbin S. 2018. Histone variants: essential
actors in male genome programming. *J Biochem* **163**: 97-103.
Jang J, Kang Y, Zofall M, Cho C, Grewal S, Lee JY. 2024. Abo1 ATPase facilitates the dissociation
of FACT from chromatin. *bioRxiv*.
Krastev DB, Li S, Sun Y, Wicks AJ, Hoslett G, Weekes D, Badder LM, Knight EG, Marlow R, Pardo
MC et al. 2022. The ubiquitin-dependent ATPase p97 removes cytotoxic trapped
PARP1 from chromatin. *Nat Cell Biol* **24**: 62-73.
Lafon A, Taranum S, Pietrocola F, Dingli F, Loew D, Brahma S, Bartholomew B, Papamichos-
Chronakis M. 2015. INO80 Chromatin Remodeler Facilitates Release of RNA
Polymerase II from Chromatin for Ubiquitin-Mediated Proteasomal Degradation. *Mol
Cell* **60**: 784-796.
Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**:
357-359.
Le Blevec E, Muronova J, Ray PF, Arnoult C. 2020. Paternal epigenetics: Mammalian sperm
provide much more than DNA at fertilization. *Mol Cell Endocrinol* **518**: 110964.
Lin CJ, Koh FM, Wong P, Conti M, Ramalho-Santos M. 2014. Hira-mediated H3.3 incorporation
is required for DNA replication and ribosomal RNA transcription in the mouse zygote.
Dev Cell **30**: 268-279.
Liu H, Wen Q, Yan S, Zeng W, Zou Y, Liu Q, Zhang G, Zou J, Zou X. 2022. Tumor-Promoting
ATAD2 and Its Preclinical Challenges. *Biomolecules* **12**.
Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**: 550.
Montellier E, Boussouar F, Rousseaux S, Zhang K, Buchou T, Fenaille F, Shiota H, Debernardi A,
Hery P, Curtet S et al. 2013. Chromatin-to-nucleoprotamine transition is controlled by
the histone H2B variant TH2B. *Genes Dev* **27**: 1680-1692.
Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, Puigserver P, Carlsson
E, Ridderstrale M, Laurila E et al. 2003. PGC-1alpha-responsive genes involved in
oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat
Genet* **34**: 267-273.
Moriniere J, Rousseaux S, Steuerwald U, Soler-Lopez M, Curtet S, Vitte AL, Govin J, Gaucher J,
Sadoul K, Hart DJ et al. 2009. Cooperative binding of two acetylation marks on a
histone tail by a single bromodomain. *Nature* **461**: 664-668.
Morozumi Y, Boussouar F, Tan M, Chaikuad A, Jamshidikia M, Colak G, He H, Nie L, Petosa C,
de Dieuleveult M et al. 2016. Atad2 is a generalist facilitator of chromatin dynamics in
embryonic stem cells. *J Mol Cell Biol* **8**: 349-362.
Nashun B, Hill PW, Smallwood SA, Dharmalingam G, Amouroux R, Clark SJ, Sharma V, Ndjetehe
E, Pelczar P, Festenstein RJ et al. 2015. Continuous Histone Replacement by Hira Is
Essential for Normal Transcriptional Regulation and De Novo DNA Methylation during
Mouse Oogenesis. *Mol Cell* **60**: 611-625.

Okada Y. 2022. Sperm chromatin condensation: epigenetic mechanisms to compact the
genome and spatiotemporal regulation from inside and outside the nucleus. *Genes*
Genet Syst **97**: 41-53.
Puumalainen MR, Lessel D, Ruthemann P, Kaczmarek N, Bachmann K, Ramadan K, Naegeli H.
2014. Chromatin retention of DNA damage sensors DDB2 and XPC through loss of p97
segregase causes genotoxicity. *Nat Commun* **5**: 3695.
Ramirez F, Dundar F, Diehl S, Gruning BA, Manke T. 2014. deepTools: a flexible platform for
exploring deep-sequencing data. *Nucleic Acids Res* **42**: W187-191.
Rezaei-Gazik M, Vargas A, Amiri-Yekta A, Vitte AL, Akbari A, Barral S, Esmaeili V, Chuffart F,
Sadighi-Gilani MA, Coute Y et al. 2022. Direct visualization of pre-protamine 2 detects
protamine assembly failures and predicts ICSI success. *Mol Hum Reprod* **28**.
Rousseaux S, Reynoird N, Khochbin S. 2022. NUT Is a Driver of p300-Mediated Histone
Hyperacetylation: From Spermatogenesis to Cancer. *Cancers (Basel)* **14**.
Shiota H, Barral S, Buchou T, Tan M, Coute Y, Charbonnier G, Reynoird N, Boussouar F, Gerard
M, Zhu M et al. 2018. Nut Directs p300-Dependent, Genome-Wide H4
Hyperacetylation in Male Germ Cells. *Cell Rep* **24**: 3477-3487 e3476.
Skarnes WC, Rosen B, West AP, Koutsourakis M, Bushell W, Iyer V, Mujica AO, Thomas M,
Harrow J, Cox T et al. 2011. A conditional knockout resource for the genome-wide
study of mouse gene function. *Nature* **474**: 337-342.
Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A,
Pomeroy SL, Golub TR, Lander ES et al. 2005. Gene set enrichment analysis: a
knowledge-based approach for interpreting genome-wide expression profiles. *Proc*
Natl Acad Sci U S A **102**: 15545-15550.
Tang MC, Jacobs SA, Mattiske DM, Soh YM, Graham AN, Tran A, Lim SL, Hudson DF, Kalitsis P,
O'Bryan MK et al. 2015. Contribution of the two genes encoding histone variant h3.3
to viability and fertility in mice. *PLoS Genet* **11**: e1004964.
van den Berg SJW, East S, Mitra S, Jansen LET. 2023. p97/VCP drives turnover of SUMOylated
centromeric CCAN proteins and CENP-A. *Mol Biol Cell* **34**: br6.
van den Boom J, Wolf M, Weimann L, Schulze N, Li F, Kaschani F, Riemer A, Zierhut C, Kaiser
M, Iliakis G et al. 2016. VCP/p97 Extracts Sterically Trapped Ku70/80 Rings from DNA
in Double-Strand Break Repair. *Mol Cell* **64**: 189-198.
van der Heijden GW, Derijck AA, Posfai E, Giele M, Pelczar P, Ramos L, Wansink DG, van der
Vlag J, Peters AH, de Boer P. 2007. Chromosome-wide nucleosome replacement and
H3.3 incorporation during mammalian meiotic sex chromosome inactivation. *Nat*
Genet **39**: 251-258.
Varet H, Brillet-Gueguen L, Coppee JY, Dillies MA. 2016. SARTools: A DESeq2- and EdgeR-Based
R Pipeline for Comprehensive Differential Analysis of RNA-Seq Data. *PLoS One* **11**:
e0157022.
Wang T, Perazza D, Boussouar F, Cattaneo M, Bougdour A, Chuffart F, Barral S, Vargas A,
Liakopoulou A, Puthier D et al. 2021. ATAD2 controls chromatin-bound HIRA turnover.
Life Sci Alliance **4**.
Yuen BT, Bush KM, Barrilleaux BL, Cotterman R, Knoepfler PS. 2014. Histone H3.3 regulates
dynamic chromatin states during spermatogenesis. *Development* **141**: 3483-3494.
971
972
973

 Supplementary materials

 Supplemental_Fig_S1

A   Post-meiotic H3.3-activated genes compared to differentially up-regulated
genes in D26-*Atad2* KO versus D26-*Atad2* WT testes

B   Post-meiotic H3.3-repressed genes compared to differentially down-regulated
genes in D26-KO versus D26-WT

C   Sex chromosome-linked genes compared to differentially up- and down-
regulated genes in D26-KO versus D26-WT

Figure S1: Venn diagrams showing the overlap of significantly differentially expressed genes between *Atad2* KO and *Atad2* WT samples at day 26 post-partum (PP), compared with three gene sets reported by (Fontaine et al. 2022). (A) Genes upregulated in the absence of histone H3.3 in both *Atad2* WT and *Atad2* KO samples at day 26 PP. **(B)** genes repressed under the same conditions, and **(C)** sex chromosome–linked genes. In the differential expression analysis, a fold-change greater than 1.2 between *Atad2* KO and *Atad2* WT samples was considered significant.

1015   **Supplemental_Fig_S2**

1016

Atad2 WT   Atad2 KO 1017

1042

1043   **Figure S2: *Atad2* gene inactivation does not disrupt testis morphology, acetylation of**
1044   **H4K5ac, or the cell-specific accumulation of H2A.L.2, TP2, PRM1 and PRM2.** Paraffin-
1045   embedded sections from wild-type or *Atad2* KO testes were used to visualize the
1046   accumulation of H4K5ac or expression of H2A.L.2, TP2, PRM1 and PRM2.

1047 **Supplemental_Fig_S3**

1048

1049

1050 **Figure S3: ATAD2 is required for proper pre-protamine 2 processing.** Extracts from

1051 representative samples shown in Fig. 7B, derived from wild-type or *Atad2* KO cells, were

1052 analyzed by SDS-PAGE followed by immunoblotting using our anti-prePRM2 and anti-histone

1053 H3 antibodies. In each panel, the films were aligned with the corresponding blot and

1054 photographed. M, molecular weight marker.

1071 **Supplemental_Fig_S4.**

1072

1073

Figure S4: *Atad2* depletion affects testis weight and mature sperm counts.

(**A**) Testes from six 21- to 27-week-old wild-type and *Atad2* KO mice were collected, weighted

(mg) and their weights normalized to body weight and only same-age mice were compared

pairwise. Histograms represent the mean of the measured weights and error bars the

Standard Deviation (upper panel). The *p*-value of pairwise Student t-test = 0.004. The lower

panel shows representative testes from wild-type and *Atad2* KO mice.

(B) Spermatozoa count (Cells /ml) were performed for pairs of wild type and *Atad2* KO (*n=6*).

The average numbers of spermatozoa counts are respectively 21.6 ± 5.8 and 13.7 ± 4,. and the

p-value of paired Student t-test = 0.001.

1109   **Supplemental_Fig_S5**

1110

Decompaction (minutes)

1111

1112   **Figure S5: Sperm head decompaction after DTT and heparin treatment in wild-type and**
1113   ***Atad2* KO males.** Epididymal sperm were treated with a DTT/heparin/Triton mixture, fixed,
1114   and stained with DAPI. Fluorescent images were acquired using a 63× oil-immersion objective,
1115   and sperm head size was quantified with ImageJ. Scale bars represent 5 µm.

1131 **Table S1**: Primary Antibodies

Name	Company	Reference	Host Species	Dilution
Anti-ATAD2	Homemade	-	Rabbit	1:1000
anti-H2A.L1/L2 SAB771	Homemade	-	Rabbit	1:200 for IHC
Anti-Histone H3 antibody - Nuclear Marker and ChIP Grade	Abcam	Ab1791	Rabbit	1:5000 for WB
Anti-H2B (testis specific)	Abcam	AB178426	Rabbit	1:100 for IF
anti-H3F3B	Abnova	#H00003021-M01	Mouse	1:200 for IF
Anti-H4K5Ac	PTM Biolabs	#PTM-163	Mouse	1:200 for IHC
Anti-HIRA	Cell Signaling Technology	D6O8L	Rabbit	1:1000 for WB
Pre-PRM2	Homemade	-	Rabbit	1:100 for IF
anti-Protamine 1	BRIAR Patch Biosciences	HUP1N	Mouse	1:200 for IHC/IF
anti-Protamine 2	BRIAR Patch Biosciences	HUP2B	Mouse	1:200 for IHC
Anti-TNP2 (B-2)	Santa Cruz Biotechnology	sc-393843	Mouse	1:200 for IHC
Goat anti Rabbit IgG (H + L)-HRP	Bio-rad	Cat# 1706515	-	1:5000 for WB
Goat anti-Mouse IgG (H + L)-HRP	Bio-rad	Cat# 1706516	-	1:5000 for WB
Alexa Fluor™ 568 goat Anti-mouse IgG (H+L)	Invitrogen	Cat# A11004	-	1:500 for IF
Alexa Fluor™ 488 goat Anti-rabbit IgG (H+L)	Invitrogen	Cat# A11034	-	1:500 for IF

WB: Western Blot Analysis, IF: Immunofluorescence and IHC: Immunohistochemistry.

1132   **Supplemental_Table_S2**

1133   **Table S2**: Reagents / Products used

Products	Company	Reference
Bovine Serum Albumin (BSA)	Euromedex	# 04-100-812-C
Clarity™ Western ECL Substrate	Bio-Rad	Cat # 170-5061
Clarity Max™ Western ECL Substrate	Bio-Rad	Cat # 170-5062
Complete Protease inhibitor EDTA-free	Sigma-Aldrich	# 4693159001
Dako Fluorescent Mounting Medium	Dako	S3023
DL-Dithiothreitol (DTT)	Euromedex	EU0006
DNase I recombinant	Roche	4536282001
DPBS	Gibco	14200-075
Ethylenediaminetetraacetic acid disodium salt dihydrate (EDTA)	Sigma-Aldrich	E6635
Ethylene glycol-bis (2-aminoethylether)-N,N,N',N'-tetraacetic acid (EGTA)	Sigma-Aldrich	E3889
Formalin solution, neutral buffered, 10%	Sigma-Aldrich	HT5012
Glutaraldehyde solution	Sigma-Aldrich	G5882
Heparin sodium salt from porcine intestinal mucosa	Sigma-Aldrich	H3149
Hoechst	Invitrogen	# 33342
Human Chorionic Gonadotropin (HCG)	MED-VET	Chorulon 1500 UI
Laemmli Sample Buffer 4x	BioRad	# 1610747
Magnesium Chloride (MgCl2)	Euromedex	2189
Mayer's Hematoxylin Solution	Sigma-Aldrich	MHS16
M2 medium	Sigma-Aldrich	M7167
M16 medium	Sigma-Aldrich	M7292
Nitrocellulose Blotting membrane	Amersham™ Protran®	10600015
Nonidet™ P 40 Substitute (NP 40)	Sigma-Aldrich	74385
NuPAGE™ 4-12% Bis-Tris Gel	Invitrogen	NP0335BOX

Potassium Ferricyanide	Sigma-Aldrich	P-8131
Potassium Ferrocyanide	Sigma-Aldrich	P-9387
Pregnant Mare Serum Gonadotropin (PMSG)	MED-VET	Syncro-part PMSG 600 UI
Saccharose	Sigma-Aldrich	S7903
Saponin from quillaja bark	Sigma-Aldrich	S7900
Sodium deoxycholate (NaDoc)	Sigma-Aldrich	D6750
Superfrost®slides	Thermo Scientific™	LCSF05
TMB ELISA Substrate (High Sensitivity)	Abcam	AB171523
Trans-Blot Turbo 5X Transfer Buffer	BioRad	Cat # 10026938
Triton™ X-100	Sigma-Aldrich	T9284
TRIzol™	Thermo Scientific™	# 15596026
TWEEN® 20	Sigma-Aldrich	P7949
Urea	Sigma-Aldrich	U5378
X-Gal	Invitrogen	B1690
96 Well Round (U) Bottom Plate	Thermo Scientific™	# 163320
450 nm Stop Solution for TMB Substrate	Abcam	AB171529

1134
1135
1136