An updated map of the trypanosome nuclear pore and its associated proteins

Bernardo Papini Gabiatti[1*], Johanna Odenwald[1], Silke Braune[1], Timothy Krüger[1], Martin

Zoltner[2*] and Susanne Kramer[1*]

[1] Biocenter, University of Würzburg, Würzburg, Germany

[2] Department of Parasitology, Faculty of Science, Charles University in Prague, Biocev,

Vestec, Prague, Czech Republic

* corresponding authors

Tel.: +49 931 3186785

Email: susanne.kramer@uni-wuerzburg.de

Email: martin.zoltner@natur.cuni.cz

Email: bernardo.gabiatti@uni-wuerzburg.de

short title: Trypanosome nuclear pore architecture

ABSTRACT

A nuclear pore consists of two outer rings flanking an inner ring connected to the nuclear

envelope. Asymmetric structures at both nucleoplasmic and cytoplasmic faces moderate

unidirectional transport of macromolecules. In the ancient trypanosomes, many of the pore

(associated) proteins are not conserved. In particular, asymmetric components are mostly

unknown, and the genome lacks orthologues to both, the nuclear mRNA-recruiting TREX-2

complex and the cytoplasmic mRNA remodelling complex Dbp5/Gle1. By combining

expansion microscopy with proximity labelling, we mapped most of the 75 trypanosome

nuclear pore localised proteins to specific subregions of the pore. At the nuclear site, we

identified structural homologues to TREX-2 proteins, the Ran binding protein RanBPL and

several trypanosome-unique proteins including NUP64 and NUP98. At the cytoplasmic site

we found RanGAP, RanBP1 and the heterotrimeric NUP76 complex that we show to

function in mRNA export, but no homologue to Dbp5/Gle1. Our novel, asymmetric and

comprehensive map of the trypanosomes pore and its associated proteins supports a Ran-

dependent mRNA export system with many trypanosome-unique features at both sides of

the pore.

INTRODUCTION

Nuclear pores penetrate the double-membrane of the nucleus and serve as an essential gateway for the exchange of proteins, RNAs and ribosomes between the nucleoplasm and the cytoplasm. They are among the largest macromolecular complexes in nature with more than 500 copies of ~30 different nucleoporins (NUPs) that form eight identical protomers (spokes) (Wente & Rout, 2010; Hampoelz *et al*, 2019; Schwartz, 2016; Lin & Hoelz, 2019). Each spoke is connected to the nuclear envelope (NE) as well as to the neighbouring spokes, resulting in multiple concentric rings: the inner ring (IR) at the centre of the pore is flanked by two outer rings (OR) at the cytoplasmic site (cytoplasmic ring) and nuclear site (nuclear ring). The outer rings are composed of large Y-shaped protein complexes, called the Nup84/NUP107 complex in yeast/human (Lin & Hoelz, 2019). This core structure of the pore is extended by the nuclear basket at the nuclear site and the cytoplasmic filaments at the cytoplasmic site. NUPs can be divided into three classes: (i) structured NUPs that form the scaffold of the pore, with structural features being limited to beta propellers, coiled coil and alpha-helical solenoids (ii) pore membrane proteins (POMs) that anchor the pore in the nuclear envelope via transmembrane regions and (iii) non-structured, intrinsically disordered NUPs, that contain FG (phenylalanine and glycine) repeat motifs and provide a diffusion barrier at the central channel of the pore by phase separation (Terry & Wente, 2009). While smaller molecules can pass by diffusion, the transport of larger molecules, such as most RNAs, ribonucleoprotein particles (RNPs), pre-ribosomes and most proteins, requires energy and depends on transporters. Protein transport, as well as the transport of micro-RNAs and tRNAs is mediated by importins and exportins of the karyopherin family (Wing *et al*, 2022). These transporters recognise and bind nuclear localisation signals (NLS) or nuclear export signals (NES) of their cargo and shuttle it within the phase-separated central channel of the pore by interacting with the FG-repeat NUPs (Wing *et al*, 2022). This transport is energised by the RanGTP/RanGDP gradient across the nuclear envelope maintained by the chromatin-bound guanine nucleotide exchange factor RCC1 and the cytoplasmic-localised proteins Ran-GTPase-activating protein RanGAP and RanBP1 (Wing

et al, 2022). Importins bind cargo and Ran-GTP mutually exclusively, while exportins bind

Ran-GTP and cargo cooperatively, thus allowing selective release of cargo either in the

nucleus or cytoplasm, respectively, driven by GTP hydrolysis cycles of Ran (Wing *et al*,

2022). The vast majority of messenger ribonucleoprotein particles (mRNPs) are not exported

by karyopherins but use the heterodimeric Mex67/Mtr2 complex (NXF1 or TAP/NXT1 in

human) instead (Chen *et al*, 2024; Magistris, 2021; Ashkenazy-Titelman *et al*, 2020). The

energy is provided by at least two RNA helicases, Sub2/UAP56 and Dbp5/DDX19

(yeast/human), that assemble and disassemble the Mex67/Mtr2/mRNA export complex in

the nucleus and in the cytoplasm, respectively, in events known as nuclear and cytoplasmic

mRNP remodelling (Xie & Ren, 2019). In ophistokonts, two asymmetric pore components,

the basket and the cytoplasmic filaments, ensure directionality of RNP transport. The basket

of *S. cerevisiae* consists of Nup1, Nup2, Nup60, Mlp1/2 and Pml39 and in metazoa of

NUP153 (orthologue to yeast Nup1/60), NUP50 (orthologue to yeast Nup2), TPR

(orthologue to Mlp1/2) and ZC3H1 (orthologue to Pml39). Nup60 anchors the yeast nuclear

basket to the Y-complex of the nuclear outer ring (Stankunas & Köhler, 2024). Nup1 and

NUP153 anchor the TREX-2 (three prime repair exoribonuclease 2) complex to the pore in

yeast (Jani *et al*, 2014) and humans (Umlauf *et al*, 2013), respectively. TREX-2 consists of

the large scaffolding protein Sac3/GANP bound to Thp1/PCID2, Sem1/DSS1, and

Sus1/ENY2 (yeast/human), and, in yeast only, to Cdc31. This complex functions in recruiting

the mRNP to the pore by direct interaction of the Sac3 N-terminal region with the mRNA-

loaded Mex67 (Stewart, 2020). The cytoplasmic filaments are heterotrimeric complexes of

Nup82/NUP88, Nup159/NUP214 and Nsp1/NUP62 (yeast/human); Nup159/NUP214 recruit

the mRNA remodelling helicase Dbp5/Ddx19 with its cofactor Gle1 (Fernandez-Martinez *et*

al, 2016; Bley *et al*, 2022).

Structure and composition of nuclear pore complexes has been characterised in a range of

organisms (examples (Kim *et al*, 2018; Kosinski *et al*, 2016; Lin *et al*, 2016; Allegretti *et al*,

2020; Mosalaganti *et al*, 2018)). While the general structure of the pore, in particular the

inner ring structure, is highly conserved, the more peripheral structures can differ between

organisms and even within the same organism (Fernandez-Martinez & Rout, 2021). Yeast

for example has up to three pore variants that differ in the number of nuclear outer rings and

in the presence or absence of a basket (Akey *et al*, 2022; Singh *et al*, 2024; Niepel *et al*,

2005; Galy *et al*, 2004). Trypanosomes have separated from the eukaryotic main branches

very early and their nuclear pore architecture is thus an important puzzle-stone towards a

better understanding of pore evolution. In particular, structural differences would help to

unravel which pore features constitute organism-specific adaptations and which have been

present in the LECA (last common eukaryotic ancestor) (Padilla-Mejia & Field, 2023; Obado *et*

al, 2017).

In *Trypanosoma brucei*, 22 NUPs were initially identified based mostly on structural

similarities to human and yeast NUPs, as the sequences are poorly conserved, and pore

localisation was confirmed by GFP-tagging (DeGrasse *et al*, 2009). This served as

foundation for a hallmark follow-up study, that has defined the sub-complexes, quaternary

structure and pore-associated proteins by a large set of immunoprecipitations with multiple

baits from cryomilled samples, combined with immunogold electron localisation and *in silico*

prediction tools (Obado *et al*, 2016). Similar to all other eukaryotes studied so far, the inner

ring is mostly conserved (Obado *et al*, 2016; Makarov *et al*, 2021; Fernandez-Martinez &

Rout, 2021), with the one exception of the membrane anchoring mechanism: *T. brucei* lacks

orthologues to all POMs of opisthokonts. Instead, TbNUP65 has evolved a C-terminal

transmembrane helix to connect to the nuclear envelope (Obado *et al*, 2016), replacing the

amphipathic lipid-packing sensor (ALPS) motif used by its opisthokont orthologues

ScNup53/59 and HsNUP35. Two additional POM candidates, with transmembrane helixes,

were recently identified within the interactome of lamin-like proteins (Butterfield *et al*, 2024).

The structured outer ring complex (Y-complex) was clearly defined in multiple affinity

purifications to consist of TbNUP158, TbSEC13, TbNUP41, TbNUP82, TbNUP89,

TbNUP132, TbNUP152 and, likely, TbNUP109 (Obado *et al*, 2016). This complex, named

NUP89 complex, is the equivalent to the outer ring complexes Nup84/NUP107 from

yeast/human and mostly conserved, with some lineage specific variations in the β-propeller

proteins (Obado *et al*, 2016). Three FG-NUPs, NUP64, NUP75 and NUP98, are unique to

trypanosomes and part of one complex with unknown localisation (Obado *et al*, 2016). There

were two major unexpected outcomes from this study: (i) no asymmetrically localised NUPs

were identified, with the exception of the basket proteins NUP110 and NUP92, suggested as

putative homologues to yeast Mlp1 and Mlp2. Even TbNUP76, which was co-isolated with

TbMex67 and has structural homology to the cytoplasmic-site specific yeast Nup82 that has

a function in mRNA export (Hurwitz & Blobel, 1995; Grandi *et al*, 1995) was predicted at both

outer rings by immunogold labelling; (ii) the authors could not identify any homologue to the

cytoplasmic mRNA remodelling enzyme, the DEAD-box RNA helicase Dbp5. Instead, they

found co-purification in high-stringency conditions between the conserved mRNA transporter

TbMex67 with Ran, RanBP1 and a putative RanGAP, indicating that mRNA export may be

fuelled by the Ran system. Meanwhile, many additional nuclear pore localised proteins were

identified, primarily by the genome-wide localisation database TrypTag (Billington *et al*,

2023), of which most remain functionally uncharacterised.

We were puzzled by the absence of asymmetric NUPs at the outer rings, which are viewed

as key determinants underpinning directed transport of macromolecules. We therefore

revisited the ultrastructure of the trypanosome nuclear pore using a novel, powerful

combination of expansion microscopy and proximity labelling techniques. Our approach

indeed identified a set of asymmetric components and we employed these as markers to

map all 75 nuclear pore localised proteins reported by TrypTag (Billington *et al*, 2023).

Altogether, we provide an updated, comprehensive map of the pore and its associated

proteins, including proteins of the Ran GTPase transport system. We describe many novel

proteins at the nuclear site of the pore, most of these trypanosome-unique, including three

potential TREX-2 complex proteins. We find the NUP76 complex proteins, NUP76, NUP140

and NUP149, exclusively at the cytoplasmic site and demonstrate a conserved function of

NUP76 in mRNA export, while NUP140 and NUP149 are unique to trypanosomes, and lack

any conserved binding site for Dbp5, consistent with the absence of this RNA helicase. Our

data, combined with the data of (Obado *et al*, 2016), support a model of the trypanosome

pore with a conserved core structure, but with a fundamental different mRNA remodelling

platform at the cytoplasmic site and many trypanosome-unique proteins at the basket site

that await functional characterisation.

MATERIAL AND METHODS

Bioinformatics

All sequences were retrieved from TriTrypDB (Shanmugasundram *et al*, 2023). InterPro was used for domain search based on sequence (Paysan-Lafosse *et al*, 2022). Homology search based in primary and predicted secondary alignments was done with Phyre2 (Kelley *et al*, 2015). Tertiary alignments of Trypanosomatid-optimised predicted AlphaFold2 models (Wheeler, 2021) were carried out with Foldseek (Kempen *et al*, 2024). Foldseek searches were performed on the web server (https://search.foldseek.com) covering all available databases (AlphaFold/Proteome v4, AlphaFold/Swiss-Prot v4, AlphaFold/Swiss-Prot v4, BMFD 20240623, CATH50 4.3.0, Mgnify-ESM30 v1, PDB100 20240101 and GMGCL 2204) with Mode 3Di/A. The outputs from Foldseek including the superimposed structures, the values of sequence identity, RMSD (root mean square deviation), TM (template modelling score), qTM and tTM (TM scores normalised by query and template length, respectively) values were retrieved. All structures were predicted using AlphaFold2-Multimer-v2.3.1 (Jumper *et al*, 2021; Evans *et al*, 2022) through the ColabFold version 1.5.3 with Mmseq2 (UniRef+Environmental) with 5 recycles and 5 models (doi:10.1038/s41592-022-01488-1). Predicted structures were visualized with ChimeraX (Meng *et al*, 2023). Heatmaps were generated with the ComplexHeatmap package (Gu, 2022) in R. The *t*-test difference values from the affinity purifications (detailed below) were fed in and clustered with a Pearson distance method (option cluster_rows = TRUE, cluster_columns = FALSE). The *t*-test difference values are represented as a colour scale and the colouring was made by the package. pLLDT plots of local prediction confidence over the protein length shown near the heatmaps are available and were retrieved from the Trypanosomatid-optimised AlphaFold2 database (Wheeler, 2021). Violin plots were created with the ggpubR package of R (https://rpkgs.datanovia.com/ggpubr/).

***Trypanosoma* cells**

Trypanosoma brucei Lister 427 procyclic cells in logarithmic growth were used for all

experiments. Cells were grown in SDM-79 supplemented with 5% (v/v) FCS and 75 μg/ml

hemin at 27°C, 5% CO_2, and appropriate drugs (Brun & Schönenberger, 1979). Drugs used

for transgenic cells were G418 disulfate (15 μg/ml), blasticidine S (10 μg/ml), puromycin

dihydrochloride (1 μg/ml), hygromycin B (25 μg/ml) and phleomycin (2.5 μg/ml); these

concentrations were used for maintenance and doubled during the actual selection process

after transfection. Growth was measured by sub-culturing cells daily to 10^6 cells/ml and

measuring densities 24 hours later using a Coulter Counter Z2 particle counter (Beckman

Coulter) over five days.

Transgenic trypanosomes were generated by standard procedures. Endogenous

tagging with TurboID-Ty1, 3xHA, 4xTy1 and *Os*AID-3xHA was done using a PCR-based

method and the pPOTv7 system (Dean *et al*, 2015). TurboID-Ty1, 3xHA and 4xTy1

customisations of the pPOTv7 were made in this work. 25 μl of PCR reaction (PrimeSTAR

MAX (Takara)) was used for transfections. The PCR product was precipitated with

isopropanol, washed once with 70% ethanol in a sterile hood, resuspended in 10 μl of sterile

ddH_2O and mixed with 10^7 cells in 400 μl of transfection buffer (Burkard *et al*, 2011).

Transfections were performed with Amaxa Nucleofactor IIb (Lonza Cologne AG, Germany,

program X-001) using BTX electroporation cuvettes (45-0125). Cells were recovered in 25

ml SDM-79 supplemented with 20% FCS for 18 h and diluted 1:4 in 75 ml SDM-79

supplemented with 20% FCS. Relevant drugs were added and cells plated in four 24-well

plates (1 ml/well). Drug-resistant populations were analysed after 10 days and confirmed

with Western blotting or diagnostic PCR from genomic DNA (Rotureau *et al*, 2005). For

ectopic, inducible expression of Mex67 fused to TurboID-Ty1 at the carboxy-terminus, its

open reading frame was cloned in frame with TurboID-Ty1 in a genetic cassette containing a

EP procyclin promoter controlled by 2x Tet operator flanked by sequences for integration to

the rRNA locus (Sunter *et al*, 2012).

For Auxin-inducible degron experiments, 50 µM 5-Ph-IAA (MedChem Express, HY-134653)

was added to the cultured cells from a 50 mM stock in DMSO. The auxin system was kindly

provided by the laboratory of Mark Carrington (University of Cambridge, UK) and is

described in (Gabiatti *et al*, 2024).

Plasmids and PCR products

All plasmids and PCR products used in this study are listed in Table S1. pPOTv7 variants for

TurboID-Ty1, 3xHA, 4xTy1 tagging were generated by sub-cloning the respective tag

sequence in the BamHI/HindIII sites. pPOTv7 *Os*AID-3xHA was generated in (Gabiatti *et al*,

2024).

Western blot and antibodies

Western blots were done using standard methods. Primary antibodies used for detection of

proteins were rat anti-HA (3F10, Roche) (1:1000), anti-Ty1/BB2 ((Bastin *et al*, 1996)

hybridoma supernatant 1:1000) and anti- *T.brucei* PFRA/B (L13D6) (1:10,000) (KOHL *et al*,

1999). Secondary antibodies were IRDye® 680 RD and 800 CW rat and mouse anti-goat

(LI-COR) (1:30,000). Biotinylated proteins were detected with Streptavidin-IRDye® 680 LT

(LI-COR) (1:10,000). Blots were scanned with the Odyssey Infrared Imaging System (LI-

COR Biosciences, Lincoln, NE).

Protein retention Expansion Microscopy (proExM) and ultrastructural Expansion

Microscopy (U-ExM)

The proExM and UExM methods were performed as previously described in

(Odenwald *et al*, 2024), with the following minor modifications for UExM: the primary and

secondary antibody labelling reactions were done in 6-well plates with 1 ml of antibody

diluted in PBS-T (0.1% Tween20 (v/v) in 3% BSA (w/v)). Primary antibodies were incubated

overnight at 37°C with agitation and secondary antibodies for 3 hours at 37°C with agitation.

The plates were slightly tilted to ensure that the whole gel piece was covered.

Microscopy and quantification of microscopy data

For all fluorescence microscopy experiments, images were acquired using a fully automated

iMIC microscope (TILL Photonics) equipped with a 100x, 1.4 numerical aperture objective

(Olympus, Japan) and a sensicam qe CCD camera (PCO, Germany). Z-stacks (75, 100, or

150 slices, 140 nm step size) were recorded. Exposure times ranged between 50-100 ms for

DAPI and 400-800 ms for all other fluorophores. Image stacks were deconvolved with

the Huygens™ Essential software v24.04 (SVI, Hilversum, Netherlands). Deconvolution

parameters were kept constant for all images, except for the number of iterations which were

optimised depending on the signal intensity and background. To correct aberrations due to

the refractive index mismatches occurring at different depths of the gel specimens, the

varPSF function of Huygens Essential was used, which calculates the PSFs according to

depth position. After deconvolution, images were corrected for the chromatic shift aberration

between the green and red channel in three dimensions. The Huygens™ Chromatic

Aberration Corrector was used with image stacks of TetraSpeck fluorescent microspheres

(T7279, Thermo Fisher Scientific) as template. Fiji (Schindelin *et al*, 2012) was used for

figure generation.

Streptavidin affinity purification and LC MS/MS analysis

Affinity purification of biotinylated proteins followed by tryptic digest and peptide preparation

were done as described (Moreira *et al*, 2023), except that 1 mM biotin was added to the on-

beads tryptic digests, to improve the elution. Eluted peptides were analysed by liquid

chromatography coupled to tandem mass spectrometry (LC-MS/MS) on an Ultimate3000

nano rapid separation LC system (Dionex) coupled to an Orbitrap Fusion mass spectrometer

(Thermo Fisher Scientific).

Spectra were processed using the intensity-based label-free quantification (LFQ) in

MaxQuant version 2.1.3.0 (Cox *et al*, 2014; Cox & Mann, 2008) searching the *T. brucei*

brucei 927 annotated protein database (release 64) from TriTrypDB (Aslett *et al*, 2010).

Analysis was done using Perseus (Tyanova *et al*, 2016) essentially as described in (Zoltner

et al, 2020). Briefly, known contaminants, reverse and hits only identified by site were filtered

out. LFQ intensities were $\log_2$-transformed and missing values imputed from a normal

distribution of intensities around the detection limit of the mass spectrometer. A Student's *t*-

test was used to compare the LFQ intensity values between the duplicate samples of the

bait with untagged control (WT parental cells) triplicate samples. The $-\log_{10}$ *p*-values were

plotted versus the *t*-test difference to generate multiple volcano plots (Hawaii plots). Potential

interactors were classified according to their position in the Hawaii plot, applying cut-off

curves for significant class A (SigA; FDR = 0.01, s0 = 0.1) and significant class B (SigB; FDR

= 0.05, s0 = 0.1). The cut-off is based on the false discovery rate (FDR) and the artificial

factor s0, which controls the relative importance of the *t*-test *p*-value and difference between

means (at s0 = 0 only the *p*-value matters, while at non-zero s0 the difference of means

contributes). Perseus was also used for principal component analysis (PCA), the profile plots

and to determine proteins with similar distribution in the plot profile using Pearson's

correlation. All proteomics data have been deposited at the ProteomeXchange Consortium

via the PRIDE partner repository (Perez-Riverol *et al*, 2019) with the dataset identifiers

PXD055934 (Nup75, Ran, Mex67), PXD047268 (NUP76, NUP96, NUP110) and

PXD031245 (NUP158, wt control).

RESULTS

Expansion microscopy identifies novel asymmetric components of the trypanosome

pore

We revisited the trypanosomatid nuclear pore architecture with expansion

microscopy. Therefore, we expressed target proteins fused to a small peptide epitope-tag

(3xHA or 4xTy1) to allow immunofluorescence detection via anti-HA or anti-Ty1. In some

experiments, we expressed the target protein fused to the biotin ligase TurboID (Branon *et*

al, 2018), followed by the detection of the auto-biotinylation with fluorescent streptavidin

(=streptavidin imaging). We had previously shown that labelling with streptavidin increases

the signal with no obvious loss in resolution, which is essential since expansion microscopy

causes a massive reduction in antigen density (Odenwald *et al*, 2024). Even more

importantly, streptavidin readily labels proteins within phase-separated areas, such as the

nuclear pore central channel, that we found largely inaccessible to antibodies (Odenwald *et*

al, 2024). Since TurboID will not only auto-biotinylate the bait but also adjacent proteins,

there is the possibility that the streptavidin signal may not reflect the true localisation of the

bait. Hence, throughout this work, we have confirmed all major findings derived from

streptavidin labelling with orthogonal methods, such immunofluorescence, mass

spectrometry and/or vice-versa labelling. All fusion proteins were expressed from the

endogenous loci, to avoid changes in gene expression.

First, we used Protein Retention Expansion Microscopy (proExM), a method that expands

cells after protein labelling (Tillberg *et al*, 2016). We obtained an expansion factor of 3.6 and

confirmed the isotropic expansion of the nucleus (Figure S1). We initially focused on the

proteins of the NUP76 complex (NUP76, NUP140, and NUP149) as these were previously

shown to co-precipitate with the trypanosome mRNA export factor Mex67 (Obado *et al*,

2016). These NUPs were fused to either N- or C-terminal peptide tags (NUP76-3xHA,

NUP140-4xTy1, 3xHA-NUP149). In the same cell lines, we co-expressed the nuclear basket

protein NUP110 with an amino-terminal fusion to a different epitope tag (4xTy1 or 3xHA).

Upon dual labelling with anti-Ty1 and anti-HA, we carried out expansion and imaging. All

four proteins resolved as single dots. The signals from the NUP76-complex proteins were in

all cases clearly separated from the NUP110 signal towards the cytoplasmic site of the pore

(Figure 1A and Figure S2 in supplementary material). Notably, we observed for every dot

signal originating from the NUP76 complex a corresponding NUP110 dot, indicating that

trypanosomes, unlike yeast (Galy *et al*, 2004), do not have basket-less pores. The median,

expansion-factor corrected distance to NUP110 was 129±17 nm for NUP76, 120±22 nm for

NUP140, and 137±18 nm for NUP149 (Figure 1B). The large distance of the NUP76

complex proteins to the basket protein NUP110 and the absence of "double-dots" for the

NUP76-complex strongly indicate asymmetric localization of the NUP76 complex exclusively

at the cytoplasmic site of the pore. The data disagree with previous observations derived

from immuno-electron microscopy that place NUP76, symmetrically, to both outer rings

(Obado *et al*, 2016).

Thus, we were concerned that our observed sole cytoplasmic localisation of the NUP76

complex proteins could be a technical artifact caused by (i) an insufficient resolution of the

proExM method (ii) a non-isotropic expansion across the nuclear membrane or (iii) reduced

accessibilities of antibodies to the nuclear ring in comparison to the cytoplasmic ring. To

investigate, we applied Ultrastructural Expansion Microscopy (U-ExM) (Gambarotto *et al*,

2019), which offers a higher resolution because the antibody labelling is applied after the

expansion and the linkage error (distance between the fluorophore of the secondary

antibody and the target protein) is therefore not expanded. U-ExM has been successfully

used in trypanosomes (Kalichava & Ochsenreiter, 2021; Gorilak *et al*, 2021; Hernández *et al*,

2023; Odenwald *et al*, 2024), and we achieved an expansion factor of 4.2-fold with isotropic

expansion of the nucleus (Figure S1). To prove that U-ExM provides the resolution to

resolve the nuclear outer ring, inner ring, and cytoplasmic outer ring we first imaged NUPs

that are conserved across eukaryotes. As an outer ring marker, we chose NUP89, the

trypanosome orthologue to the outer ring Y-complex component of yeast Nup84/85 and

human NUP107/75 (Obado *et al*, 2016). As an inner ring marker, we selected NUP96, the

conserved trypanosome orthologue to *S. cerevisiae* Nic96. We co-expressed NUP89-

TurboID with NUP96-3HA. U-ExM with fluorescent streptavidin and anti-HA readily resolved

NUP89 as a double dot, sandwiching the single dot signal of NUP96 (Figure 1C), proving

that the resolution of the method is sufficient to distinguish these different subregions of the

pore. However, as the NUP96 signal was weak, we searched for a better inner ring marker.

We tested NUP64, a trypanosome-unique FG-repeat NUP previously identified as a multi-

complex NUP localised mostly to the centre of the pore (Obado *et al*, 2016). To our surprise,

the resulting single NUP64 dot signal was not sandwiched by the two outer ring dots of

NUP89 but instead co-localised solely with the NUP89 dot at the nuclear site (Figure 1D).

Likewise, TbNUP98, known to form a complex with NUP64 (Obado *et al*, 2016), resolved as

a single dot that colocalised exclusively with the NUP64 dot at the nuclear site (Figure 1D).

The data indicate an asymmetric, exclusive nuclear-site localisation of NUP64 and NUP98

(Figure 1D).

Having confirmed that UExpM readily resolves the two outer rings, we reassessed the

localisation of the NUP76 complex. Using identical cell lines as for ProExM (Figure 1A) we

obtained the same result: all NUP76 complex proteins were resolved as single dots at the

cytoplasmic site, clearly separated from NUP110 (Figure S2). We further confirmed the sole

cytoplasmic localisation of the NUP76 complex by co-staining C-terminal fusions of this

complex to either 3xHA (Figure S3) or TurboID (Figure 1E) with a C-terminal 4xTy1 fusion of

our newly-identified nuclear-site marker NUP64.

In summary, we discovered five asymmetric proteins of the trypanosome nuclear pore

complex, previously assumed to be symmetrically distributed: NUP76/NUP140/NUP149 at

the cytoplasmic site and NUP64/NUP98 are at the nuclear site. With the exception of

NUP76, which is the structural orthologue to yeast NUP82 and human NUP88 (Obado *et al*,

2016), all novel asymmetric NUPs are trypanosome-specific.

A proximity map of the trypanosome nuclear pore

The novel availability of asymmetric NUPs prompted us to use mass spectrometry data from

TurboID proximity labelling experiments, to generate a proximity map of the entire

trypanosome nuclear pore. We used six available LC-MS/MS datasets from streptavidin-

affinity purifications, namely amino- and carboxy- terminal TurboID fusions of NUP110

(basket), NUP96 (inner ring) and NUP76 (cytoplasmic outer ring) (Odenwald *et al*, 2024). Of

all the proteins that were labelled by these baits, we initially focused on bona fide NUPs. For

each dataset, we colour-coded the enrichment of the NUP proteins based on *t*-test

difference to a wild-type control. Then, the NUPs were sorted by hierarchical clustering

applying a Pearson distance method (Figure 2A). We included the pLLDT plots (Wheeler,

2021) to indicate predicted structured and unstructured (mostly FG-repeat) domains for each

of the NUPs. The majority of the NUPs separated into three clearly distinct main clusters.

The first cluster contained the basket proteins (NUP110, NUP92), the lamina protein NUP2

(Maishman *et al*, 2016) and NUP64/NUP98 that we had placed asymmetrically localised to

the nuclear site by expansion microscopy (Figure 1D). The second cluster largely consisted

of NUPs previously identified as inner ring NUPs (Obado *et al*, 2016) while the third was

dominated by NUPs previously assigned to the outer ring (Obado *et al*, 2016). Eight NUPs

were not or poorly labelled by the three bait proteins, either, due to their small size (Moreira

et al, 2023) or for unknown reasons. These were manually added to the tree, using

positional information from (Obado *et al*, 2016), except for NUP48 and Gle2 which could not

be unequivocally assigned. For the vast majority of NUPs, the proximity map confirmed the

previous assignments of the NUPs from affinity capture experiments (Obado *et al*, 2016).

NUP98 and NUP64 unequivocally grouped with the basket/nuclear site (basket and inner

ring) and were labelled by NUP110 and NUP96 baits, but not with the cytoplasmic NUP76, in

line with our proExM and U-ExM data (compare Figure 1D). To our surprise, NUP75, which

shares 46% sequence identity with NUP64 and associates with both NUP98 and NUP64

(Obado *et al*, 2016), was placed to the inner ring and was not labelled by NUP110. We did a

vice versa TurboID experiment with NUP75 as bait (Figure S4, Table S2) and confirmed the

absence of NUP110 labelling, as well as the proximity of NUP75 to NUP98 and NUP64

(Figure 2B). Moreover, our previously published TurboID data with the outer ring NUP158

(carboxy-terminal fusion) strongly labelled NUP64 and slightly less NUP98, but not NUP75

(Figure 2C), further supporting the absence of NUP75 from the outer rings (Odenwald *et al*,

2024). Our data strongly suggest a model of the asymmetric NUP98/64/75 complex reaching

from the nuclear outer ring to the inner ring, with NUP98 and NUP64 being located at the

outer nuclear ring and NUP75 at the inner ring. Data from previous affinity isolation

experiments with NUP98, NUP64 and NUP75 show marked differences between the

interactomes of NUP98/64 and NUP75, including the exclusive absence of NUP110 from the

NUP75 interactome, in full agreement with our data (Obado *et al*, 2016).

Only two NUPs exhibited ambiguous placement in the proximity map. The outer ring

NUP132 is labelled strongly by all eight baits arguing against a sole localisation to the

basket/nuclear site and suggesting a symmetric distribution with the two protomers

extending to both sites of the pore. Further, the presumed inner ring NUP53a is labelled by

the outer ring NUP76. However, labelling by NUP96 is stronger and the outer ring NUP158

bait labels NUP53a very weakly, indicating that NUP53a is mostly at the inner ring,

potentially reaching out to the outer ring.

In summary, the proximity map accurately predicts the localisations for the vast majority of

NUPs. Importantly, it offers orthogonal validation of the asymmetric localisation of NUP98

and NUP64 to the nuclear site of the pore as derived from expansion microscopy. Therefore,

the NUP110/96/76 proximity map has the potential to predict localisation of proteins with

sub-pore size resolution, which prompted us to look at all 44 proteins that have nuclear pore

localisation according to TrypTag (Billington *et al*, 2023) but are not annotated as *bona fide*

NUPs.

Mapping the Ran-based mRNA export system to the pore

First, we focused on all nuclear-pore localised proteins that are involved in mRNA export:

Mex67 (Schwede *et al*, 2009) , Mtr2, Mex67b (Obado *et al*, 2022b) and, as postulated

(Obado *et al*, 2016, 2022a), Ran, RanGAP, the putative RanGDP importer NTF2 and two

Ran binding proteins, RanBP1 and RanBPL (Brasseur *et al*, 2014). The proximity map

places RanGAP, RanBP1 and Ran to the cytoplasmic site of the pore, while RanBPL is

predicted at the nucleoplasmic site (Figure 3A). For the mRNA transporters Mex67 and, in

particular for Mex67b, the labelling was weaker and less confined to a specific site. For Mtr2

and NTF2 we obtained no labelling, likely due to their small size which is problematic in

BioID (Moreira *et al*, 2023).

Next, we determined the localisation of Mex67, Ran, RanGAP, RanBP1 and RanBPL by U-

ExM, expressing TurboID fusions in a cell line that expressed NUP64-3xHA as a

nucleoplasmic-site marker (Figure 3B, C). RanGAP and RanBP1 resolved as a single dot,

well distanced from the NUP64 dot towards the cytoplasmic site, while the RanBPL signal

overlapped with the NUP64 signal at the nuclear site, in full agreement with the proximity

map. The biotinylation signal of the two proteins with suspected shuttling activity, Ran and

Mex67, resolved as a large cytoplasmic dot and a smaller nuclear dot, connected by a

string-like signal reaching through the pore. For Mex67, we observed that these bone-

shaped signals were more pronounced when the gene was 8-fold overexpressed from an

ectopic locus (Figure 3B, images on the right) which only slightly impaired growth (Figure

S5). For Ran, Mex67 and RanBPL, we observed an additional signal at the nucleolus, which

is defined by the reduction in DAPI stain (Figure S6A), and a minor signal in the

nucleoplasm. For RanBPL, the nucleolar signal was stronger than the signal at the pores,

while for Ran and Mex67 (at endogenous expression levels) the nuclear pore signal was

dominant. We attempted to confirm the nucleolar localisation by direct immunofluorescence

instead of streptavidin imaging. The nucleolus is challenging to label with antibodies due to

phase separation (Odenwald *et al*, 2024), but for Mex67 fused to 4Ty1 we could get a weak,

but distinct nucleolar antibody signal (Figure S6B). The functional implications of the

nucleolar localisation of Ran, Mex67 and RanBPL is not fully understood in trypanosomes

but not unexpected, as in ophistokonts Ran and Mex67 participate in pre-ribosome transport

(Li *et al*, 2023).

For the shuttling proteins Ran and Mex67 we confirmed the streptavidin-based imaging data

by LC-MS/MS upon streptavidin enrichment (Figure 3D, Figure S4, Table S2). Both Mex67

and Ran strongly labelled FG-NUPs lining the inner pore channel, consistently reflecting

their movement across the pore. The NUPs with the strongest labelling were the asymmetric

NUPs on both sides of the pore: NUP149 at the cytoplasmic site and NUP98, NUP64 and

the lamin-like protein NUP2 at the nuclear site (red arrows in Figure 3D). This strongly

suggests that both proteins, Mex67 and Ran, would have binding sites at both sides of the

trypanosome pore, analogous to human Ran (Wu *et al*, 1995; Mahajan *et al*, 1997; Lindsay

et al, 2002; Nakielny *et al*, 1999). There was weak labelling of structured NUPs, explaining

why the NUP76/96/110 proximity map did not resolve their shuttling (Figure 3A).

Consistently, preferential labelling of asymmetric FG NUPs over structured NUPs has also

been shown for human karyopherins tagged with the biotin ligase BirA* (Mackmull *et al*,

2017).

In conclusion, the NUP110/96/76 proximity map predicted the localisation of all non-shuttling

Ran system components confidently and in agreement with streptavidin imaging in U-ExM.

RanGAP is at the (expected) cytoplasmic site, together with RanBP1. RanBPL had not been

previously mapped, but is unequivocally placed to the nuclear site, consistent with its binding

preference for RanGTP (Brasseur *et al*, 2014). The NUP76/96/110 proximity map failed to

categorise the shuttling proteins Mex67 and Ran, likely because key interactions with non-

structured FG-NUPs are not represented in our bait repertoire. Direct BioID combined with

orthogonal assessment through expansion microscopy, was required to confidently place

these putative export factors. The derived localisations are summarised in Figure 3E and are

consistent with a mechanistically divergent Ran-dependent mRNA export in trypanosomes.

Mapping unknown proteins to the pore

To predict the localisation of the remaining 38 nuclear pore-localised proteins more

accurately, we extended the NUP76/96/110 map by the proximity labelling data of Mex67

and Ran and of the outer ring protein NUP158 (Moreira *et al*, 2023).

Fifteen of the 38 nuclear pore-localised proteins are karyopherins (Figure S7), five of which

have not been previously classified as karyopherins but have unequivocal structural

homology to importin and exportin-like folds predicted by FoldSeek (Figure S7B); these

include a putative orthologue to the importin Hikeshi (Tb927.1.1400) that is specialised on

the import of Hsp70-family proteins (Kose *et al*, 2012) (Figure S7C). Karyopherins were

mostly not, or poorly labelled by NUP76/96/110 (Figure S7A). The likely reason is their

preferred interaction with FG-NUPs rather than structured NUPs, similar to what we

observed for Mex67 and Ran (compare Figure 3A). The one exception is Xpo1 (exportin 1),

known to be involved in the transport of both mRNAs and tRNAs (Buhlmann *et al*, 2015),

which is labelled by all six bait proteins.

An additional five proteins with nuclear pore-localisation were not labelled by either of the six

proteins (Figure S8). For three of these proteins, Tb927.11.1000, Tb927.10.12200 and

Tb927.10.8160, the reason could be failed detection due to small size (Moreira *et al*, 2023).

None of these small proteins has homologues outside of trypanosomatids and their function

is unknown. Tb927.10.8160 has the strongest nuclear pore localisation ((Billington *et al*,

2023), Figure S8B) and high-throughput phenotyping indicates an essential function (Alsford

et al, 2011). The two larger proteins (Tb927.1.3230 and Tb927.9.12700) do not have very

prominent nuclear pore localisation ((Billington *et al*, 2023), Figure S8B). For Tb927.9.12700,

biochemical data indicate glycosomal localisation (Güther *et al*, 2014) and Tb927.1.3230

could be the trypanosomatid ortholog of the ribosome biogenesis factor Rix7 (Lo *et al*, 2019).

Their lack of labelling might thus be due to poor or absent nuclear pore localisation.

The remaining 18 proteins were labelled by at least one of the six bait proteins (Figure 4A).

Strikingly, none were labelled by the cytoplasmic-site marker protein NUP76, suggesting

absence of further proteins with exclusive cytoplasmic localisation, other than the NUP76

complex, RanGAP and RanBP1. Moreover, not a single protein was exclusively labelled by

the outer ring NUP158 (Moreira *et al*, 2023), with the one exception of Tb927.11.13080. The

(near) absence of combined labelling by NUP76 and NUP158 suggests that the outer-ring

proteome may be complete. Instead, these 18 proteins were either labelled exclusively by

NUP110 (7 proteins) or NUP96 (4 proteins) or by both, NUP110 and NUP96 (3 proteins).

Four proteins were not labelled by NUP110/96/76, but showed some labelling with Mex67,

Ran or NUP158.

The majority of these 18 proteins is unique to Kinetoplastida or even to Trypanosomatida

and lack functional annotation. Only two proteins have readily identifiable homologues

outside Kinetoplastida: Tb927.10.9020 has homology to the non-catalytic, tRNA substrate

binding subunit of the tRNA methyltransferase Trm6/Gcd10, responsible for adenosine(58)-

N(1) methylation, a modification present in many eukaryotic tRNAs (Anderson *et al*, 2000).

The *T. brucei* protein exhibits strong nuclear pore localisation (Billington *et al*, 2023) and

clusters with the basket in our proximity map, which is in contrast to the nuclear localisation

observed for *S. cerevisiae* and *A. thaliana* Gcd10/TmR6 (Anderson *et al*, 1998; Tang *et al*,

2020). Whilst localisation to the nuclear pore and/or envelope is not unheard of for tRNA

modifying enzymes (Simos *et al*, 1996; Clark & Abelson, 1987), this finding requires further

investigation, as, conversely, the putative *T. brucei* homolog (Tb927.11.11660) of the

corresponding catalytic subunit, TRM61/Gdc14, localises to the nuclear lumen/nucleoplasm

localisation (Billington *et al*, 2023). The second protein with homologies outside

Kinetoplastida, Tb927.9.2220, is a SUMO protease of the Ulp/SENP (ubiquitin-like protease/sentrin-specific protease) family with potential function in resolving stalled DNA replication forks (Kramarz *et al*, 2020). Both the SENP protease Ulp1 of yeast and SENP2 of human have nuclear pore localisation (Hang & Dasso, 2002; Nie & Boddy, 2015) and the latter was localised to the nuclear site of the pore, consistent with our map (Hang & Dasso, 2002).

The remaining 16 proteins contain five proteins with predicted basket or inner ring localisation that were previously identified as lamina associated proteins (LAPs), based on their interactions with the lamina-like proteins NUP1 and NUP2 (Butterfield *et al*, 2024). Two of these LAPs, LAP71 and LAP102, are basket specific in our map, as expected for lamina associated proteins. Two further LAPs, LAP73 and LAP59, have exclusive inner ring prediction. Consistently, LAP59, conserved across eukaryotes, has two N-terminal transmembrane domains and was suggested to be a POM (pore membrane protein) (Butterfield *et al*, 2024). LAP73 has a divergent NUP35/Nup53 type RNA binding domain (Butterfield *et al*, 2024) and the *T. brucei* Nup53 ortholog, TbNup65, is anchored to the nuclear envelope via a trans-membrane helix (Obado *et al*, 2016). This raises the possibility of a nuclear envelope- and thus inner ring localisation of LAP73 via binding to TbNUP65. However, immunoprecipitation assays failed to establish an inner ring association with LAP73 (Obado *et al*, 2016). It is possible that the interaction is weak and thus exclusively detectable in BioID, as our data strongly suggests inner ring proximity, consistent with LAP73 potentially bridging NUP65 and the inner ring, a hypothesis that deserves further investigation.

Of significant interest is basket/inner ring-predicted LAP173, which has a Sac3/GANP domain and was suggested to be the orthologue to Sac3 and sole representative of a potential trypanosome TREX-2 complex (Butterfield *et al*, 2024). Association with Mex67 was observed with two methods: affinity purifications (Obado *et al*, 2016) and BioID (Moreira *et al*, 2023). In fact, the Sac3/GANP region of a LAP173 model predicted by a trypanosome-

optimised AlphaFold2 (Jumper *et al*, 2021; Wheeler, 2021) displays remarkable structural

homology to the equivalent region of an experimentally resolved *S. cerevisiae* Sac3 structure

(RMSD 1.12Å, (Schneider *et al*, 2015)), despite poor sequence conservation (Figure 4B).

Motivated by the presence of putative Sac3, we used Foldseek (Kempen *et al*, 2024) to

search for structural homologues of the remaining TREX2-complex components, using

AlphaFold2 models as inputs (Jumper *et al*, 2021; Wheeler, 2021). We identified the Csn12-

like domain containing protein Tb927.11.5560 as a putative Thp1 orthologue (Figure 4C),

with structural homology to the human Tph1 homologue PCID2 (PDB entry 3T5X; TM score

0.82), while the primary sequence is, again, poorly conserved. Just like Sac3, our proximity

map places this Thp1 candidate to both, basket and inner ring. Moreover, we identified

Tb927.7.5830 as a putative Sus1 orthologue with highest structural similarity to Sus1 of the

yeast *K. phaffii* (Papai *et al*, 2020) (Figure 4D). The Sus1 candidate protein is not labelled by

the structural NUPs, presumably due to its small size. However, all three trypanosome

TREX2 complex candidates, including Sus1, are strongly labelled by Mex67, a prototypic

Sac3 interactor in ophistokonts (Ellisdon *et al*, 2012) and Ran, supportive of a potential role

in a trypanosome TREX2 complex (Figure 4A). Two further Trex-2 components, Sem1 and

Cdc31 (Stewart, 2020), were not identified by our modelling approach, that encompassed all

nuclear pore localised proteins. Sem1 appears to act as Thp1/Sac3 complex stabilising

element and also associates with other, functionally unrelated, complexes (Ellisdon *et al*,

2012). Cdc31, like Sus, binds to the Sac3 CID domain and both these factors are reported to

promote nuclear basket association (Jani *et al*, 2009). It is conceivable that Cdc31 and

Sem1 may be dispensable in a divergent trypanosome TREX2 complex, which warrants

functional investigation.

In conclusion, our extended NUP110/NUP96/NUP76/Me67/Ran proximity map granted

mapping the majority of nuclear pore localised proteins to a sub-region of the pore. We

found no evidence for the existence of any further proteins asymmetrically distributed to the

cytoplasmic-site other than the NUP76 complex, RanGAP and RanBP1, indicating

completeness of the cytoplasmic-site specific proteome of the pore. Instead, we predict a

diverse cohort of proteins with preferential localisation to the basket or nuclear site of the

pore, including three putative TREX2 complex proteins with proximity to Ran and Mex67,

indicative of a conserved function. These candidate TREX2 components, as well as the

other newly assigned basket-specific proteins now await experimental analysis, to

understand the mechanistic details of the trypanosome mRNA export platform.

The trypanosome NUP76 complex as a cytoplasmic mRNA remodelling hub

We have assigned the NUP76 complex (NUP76, NUP140, NUP149) exclusively to the

cytoplasmic site (Figure 1A, E) and a previous study has shown the interaction of this

complex with Mex67 under high stringency conditions (Obado *et al*, 2016). In combination,

these data strongly suggest that the NUP76 complex is the trypanosomes cytoplasmic

mRNP binding hub that serves as remodelling platform. In opisthokonts, the cytoplasmic

mRNP remodelling platform is based on the heterotrimeric complex composed of

Nup82/NUP88, Nup159/NUP214 and Nsp1/NUP62 (yeast/human) (Figure 5A) (Fernandez-

Martinez *et al*, 2016; Bley *et al*, 2022). The three proteins are connected via a carboxy-

terminal parallel coiled-coil structure. Both Nup82/NUP88 and Nup159/NUP214 possess N-

terminal β-propellers, that provide direct binding platforms for Nup145N/NUP98 (which

recruits Gle2/RAE1) and the RNA helicase Dbp5/DDX19 (recruiting Gle1/GLE1),

respectively, (yeast/human, Figure 5A). Nsp1/NUP62 and Nup159/NUP214 possess FG

repeat regions. *T. brucei* NUP76 has been previously suggested as Nup82/NUP88

homologue (Obado *et al*, 2016); indeed, the AlphaFold2 model of NUP76 (Jumper *et al*,

2021; Wheeler, 2021) shows an analogous structural organisation with a β-propeller at the

amino-terminus, interrupted by a long, disordered coil, and a three-helical coiled-coil at the

carboxy-terminus (Figure 5B-C and Figure S9). Moreover, *T.brucei* NUP76 may share its β-

propeller interactions with NUP82/88: orthologues to both Nup145N/NUP98 and Gle2/RAE1

can be readily identified in trypanosomes (DeGrasse *et al*, 2009; Obado *et al*, 2016).

However, the two remaining TbNUP76 complex components, TbNUP140 and TbNUP149, do not exhibit detectable structural homology to the NUP82/NUP88 partner proteins Nup159/NUP214 and Nsp1/NUP62 (Figure 5A-C) as based on AlphaFold2 predictions. TbNUP140 consists almost entirely of FG repeats of the PxFG type, apart from a small N-terminal stretch with a coiled-coil structure. predicted with low confidence. NUP149 is not FG rich, with only few FG motifs of the SxFG and of the FxFG type but contains six zinc fingers sparsed by coils and potentially a small coiled-coil region at the carboxy-terminus (Obado *et al*, 2016). Notably, zinc fingers are also present in the human cytoplasmic-filament NUP358 and the nuclear-site localised NUP153, both absent from trypanosomes, where they engage in Ran binding (Yaseen & Blobel, 1999; Higa *et al*, 2007). However, the zinc fingers of TbNUP149, confidently predicted as three β-hairpin strands with four cysteines side chains coordinating a zinc ion by AlphaFold2 and AlphaFold3 (Abramson *et al*, 2024) appear to lack obvious sequence or structural homology to the zinc fingers of human NUP358 and NUP153 (Figure S10). Whether these TbNUP149 zinc fingers promote Ran binding, analogous to human NUP358 and NUP153, remains to be investigated. Notably, NUP149 is heavily biotinylated by Ran-TurboID, indicative of a possible interaction (Figure 3B).

To investigate whether the trypanosome NUP76 complex is involved in mRNA export, we depleted the protein with an auxin-inducible degron system. Both alleles of the NUP76 gene were genetically fused to the OsAID-3HA sequence, in a cell line that expressed the necessary components for the auxin degron system (Gabiatti *et al*, 2024); the cell line was confirmed by diagnostic PCR (Figure S11). Upon induction with the auxin derivative 5-Ph-IAA, the NUP76-OsAID-3HA protein was depleted within 2 hours (Figure 6A), followed by growth arrest (Figure 6B) and accumulation of poly(A) signal in the nucleus that was saturated 4 hours post induction (Figure 6C). This phenotype is similar to the one observed upon Nup82 depletion in yeast (Hurwitz & Blobel, 1995; Grandi *et al*, 1995), strongly suggesting that NUP76 is the functional orthologue to yeast Nup82 with a crucial role in mRNA export. In order to limit the possibility that the observed blockade of mRNA export is

an indirect effect, i.e., the result of a disrupted pore architecture, we expressed a range of

NUPs as N- or C-terminal eYFP fusions in the NUP76 auxin degron cell line to test whether

their localisation to the pore is dependent on NUP76 (Figure 6 D and E and Figure S12). The

localisation of the inner ring NUP96 was not affected by NUP76 depletion, suggesting that

the overall pore structure remains intact (Figure 6D). Of the (putative) NUP76 associated

proteins, only the pore localisation of NUP140 was clearly abrogated upon NUP76 depletion,

while NUP149 and Gle2 still localised to the pore. Note that a slightly diminished pore

localisation was observed for all four proteins, likely caused by the disrupted mRNA export

and general loss in fitness rather than a specific impact on nuclear pore architecture. Thus,

NUP140 localisation to the pore is fully dependent on NUP76, while NUP149 and Gle2

appear to be anchored independent of NUP76.

In conclusion, the cytoplasmic-site localised NUP76 complex of trypanosomes, consisting of

NUP76, NUP140 and NUP149, is distinct from the cytoplasmic mRNA remodelling complex

of yeast and human. While NUP76 is the likely functional homologue to Nup82/NUP88 from

yeast/human, NUP140 and NUP149 are trypanosome-unique with no sequence or structural

homology to cytoplasmic site (filament) proteins from opisthokonts. The absence of a

Nup159/NUP214 (yeast/human) orthologue in trypanosomes correlates with the absence of

orthologues to its interaction partners Dbp5/DDX19 and Gle1/GLE1 (yeast/human),

suggestive of significant differences of mRNA remodelling mechanisms in trypanosomes.

A mechanistically divergent Ran-dependent mRNA export pathway in trypanosomes

Apart from the NUP76 complex, we mapped RANGAP and RANBP1 to the cytoplasmic site

of the nuclear pore. Their sole cytoplasmic localisation is suggestive of a conserved function

of these proteins in triggering GTP hydrolysis of RanGTP and thus disassembly of exportin-

cargo-RanGTP and importin-RanGTP complexes. Trypanosome RanGAP is

phylogenetically more closely related to a RabGAP (Gabernet-Castello *et al*, 2013) but its

proposed function as RanGAP (Obado *et al*, 2016) is further corroborated by our study.

Pore-anchoring of RanGAP and underlying mechanisms significantly vary across species,

ranging from a SUMO dependent interaction with the metazoan specific RanBP2/NUP358

(Matunis *et al*, 1998; Mahajan *et al*, 1998, 1997; Matunis *et al*, 1996), over a WPP domain

specific to plant RanGAP that interacts with a plant specific nucleoporin (Xu *et al*, 2007;

Rose & Meier, 2001), to no pore localisation at all in *S. cerevisiae* and *S. pombe* (Hopper *et*

al, 1990; Melchior *et al*, 1993). The mechanism of pore localisation of trypanosome RanGAP

is thus likely unique and may involve interactions with likewise unique trypanosome specific

NUPs such as NUP140 and/or NUP149. Trypanosome RanBP1 consists of a disordered 30-

amino acid stretch followed by a conserved RanBP domain (Figure S13A) and it remains

unclear whether it has binding sites to the pore or simply concentrates in sites of cargo

docking.

The lack of a Dbp5 homologue and the interaction between Mex67 and Ran implies that

trypanosomes employ the Ran-GTP gradient for mRNA export (Obado *et al*, 2016). In fact,

while Ran is predominantly nuclear localised in humans (Hutchins *et al*, 2009) we find

biotinylated Ran targets on both sites of the pore, possibly reflecting Ran engagement in

cargo import and export (Hutchins *et al*, 2009). This unique dual usage of the Ran pathway

for both mRNA and protein cargo presents a formidable challenge for export/import ratio

moderation. It is tempting to speculate that RanGTP is anchored at the basket site awaiting

the MEX67 bound mRNP, which is then liberated on the cytoplasmic site driven by RanGAP-

catalysed GTP hydrolysis. The lack of Dbp5 suggests that ATP-dependent mRNP

disassembly at the cytoplasmic site of the pore is dispensable in trypanosomes implying a

fundamentally different mode of interaction between MEX67 and mRNA. Indeed,

trypanosome Mex67 uniquely carries a CCCH-type zinc finger instead of the canonical RNA

recognition motif containing RNA binding domain (RRM/RBD) found in ophistokonts (Kramer

et al, 2010; Dostalova *et al*, 2013), and trypanosomes lack mRNA adaptors (SR proteins)

that would require stripping during cytoplasmic remodelling. Thus, the exported trypanosome

RNP may exhibit lower stability and complexity, making a remodelling RNA helicase

redundant.

While only five proteins are specific to the cytoplasmic site, the nuclear site of the pore appears more complex. Next to the previously described basket proteins NUP110 and NUP92 (Obado *et al*, 2016; DeGrasse *et al*, 2009) we found nuclear site localisation for the FG nucleoporins NUP64 and NUP98, as well as for RanBPL, putative TREX-complex proteins and several further proteins that are mostly unique to trypanosomes.

The FG-NUPs NUP64 and NUP98 are unique to trypanosomes and in a complex with NUP75 (Obado *et al*, 2016) which appears to extend to the inner ring. NUP64 and NUP98 were previously suggested to be the (functional) orthologues of *S. cerevisiae* Nup1 and Nup60, as they carry the same FG-type and engage in an interaction with the putative Sac3 homologue (Obado *et al*, 2016; Butterfield *et al*, 2024). Our data now show the exclusive nuclear-side position of these NUPs, in full support of this model.

RanBPL has a Ran-binding domain which is very similar to the one of cytoplasmic-site localised RanBP1 but has a longer disordered amino-terminal stretch (Figure S13) and was previously characterized as Ran binding protein with a clear preference to RanGTP over RanGDP (Brasseur *et al*, 2014). Thus, RanBPL may be the trypanosome functional counterpart to basket proteins Nup2/NUP50 (yeast/human), which also possess Ran-binding domains. Whilst the multiple roles of Nup2/NUP50 remain largely elusive, one known function is the acceleration of protein import complex disassembly through stimulation of RanGEF/RCC1 activity (Holzer & Antonin, 2022), analogous to the function of RanBP1 as enhancer of RanGAP activity at the cytoplasmic site (Seewald *et al*, 2003). In trypanosomes, a RanGEF has not yet been identified and the absence of a detectable RanGEF/RCC1 domain among the proteins biotinylated by Ran indicates that a trypanosome RanGEF/RCC1 is either absent or divergent. Theoretically, RanBPL has the potential to compensate for the absence of the canonical RanGEF/RCC1: instead of directly catalysing the GDP to GTP exchange, RanBPL1 could act by stabilizing RanGTP and preventing GTP hydrolysis, driving the equilibrium towards RanGTP. However, the observation that RanBPL

734   silencing evokes only a mild growth phenotype (Brasseur *et al*, 2014) argues against this

735   hypothesis.

736   Altogether, our study fortifies the hypothesis of a Ran-dependent mRNA export pathway in

737   trypanosomes and opens new avenues for exploration of the underlying molecular

738   mechanisms.

CONCLUSION

Our revisited map of the trypanosome nuclear pore conforms to the pattern of conservation at the core scaffold regions and diversity at the borders of the pore (Makarov *et al*, 2021). We discovered an asymmetric architecture, confidently placing the NUP76 complex exclusively to the cytoplasmic site and defining the sole localisation of the trypanosomatid-exclusive FG NUPs NUP64 and NUP98, at the basket site. Notably, this corrects the current view of a largely symmetric trypanosome nuclear pore and ultimately supports moderation of directional nucleocytoplasmic transport which is crucially dependent on asymmetric components at the nuclear pore borders in other systems. For the NUP76 complex, our data strongly indicates a crucial function as cytoplasmic mRNP remodelling hub, analogous to the Nup82/NUP88 complex in opisthokonts, while the presence of trypanosome-unique NUP140 and NUP149 implies significant mechanistic difference. Mapping of the export factors Mex67 and Ran elucidated further divergence, supporting a trypanosome-specific, Ran-dependent export system. Lastly, we present a comprehensive assignment of pore localised proteins to sub-regions of the nuclear pore that resulted in the identification of novel nuclear pore components, including three putative members of a trypanosome TREX-2 complex. Altogether, our approach delivers asymmetric and novel nuclear pore components, including positional information, which can now be interrogated for functional roles to explore trypanosome- specific adaptions of nuclear transport, export control, and mRNP remodelling. The compartmentalisation of hereditary information in the nucleus necessitated the invention of a gateway allowing mRNPs and a variety of other essential cargos to cross the nuclear envelope. Hence, the nucleus, eponymous feature of eukaryotes, must have co-evolved with the nuclear pore that constitutes this gateway. The study of nuclear pores in evolutionary divergent eukaryotes, such as the ancient trypanosomes, is fundamental to understand the evolutionary origin of the nucleus and decipher the complexity of the nuclear pore as platform with multiple cargo routes. Our study contributes a roadmap of the trypanosome nuclear pore, reporting conserved and non-conserved features and devising a plethora of new leads for further exploration.

 FIGURE LEGENDS

Figure 1. Expansion microscopy identifies novel asymmetric pore proteins

(A) Protein retention Expansion Microscopy (proExM) of cell lines co-expressing epitope-

tagged versions of NUP76, NUP140 or NUP149 in combination with NUP110. Proteins were

labelled with antibodies to the epitope tags, as indicated, before expansion. Images were

deconvolved (20 iterations for NUP149 and NUP76; 60 iterations for NUP140) and single

planes of the nuclei are shown. Further images are shown in Figure S2.

(B) The distances between NUP76/NUP140/NUP149 and NUP110 were measured from

manually selected pores that had both proteins in the same focal plane. The data were

corrected for the expansion factor are shown as a violin plot. Mean ± standard deviation is

on the right.

(C and D) Ultrastructure Expansion Microscopy (UExM) of cells expressing proteins fused to

TurboID or 3HA, as indicated. Labelling was done with fluorescent streptavidin and with anti-

HA. Images were deconvolved (20, 20, 40 iterations for NUP89/NUP96, NUP89/NUP64 and

NUP98/NUP64, respectively). A single plane image of an entire nucleus is shown on the left

and three enlarged regions from the same or another nucleus are shown on the right

(Cy=cytoplasm, Nu=nucleus). For NUP89/NUP96, yellow arrows point to pores that are in a

suitable focal plane to see the two NUP89 dots sandwiching the NUP96 dot.

(E) Ultrastructure Expansion Microscopy (UExM) of lines co-expressing TurboID-tagged

versions of NUP76 or NUP149 with NUP64-3HA. Labelling was done with fluorescent

streptavidin and anti-HA. Images were deconvolved with 60 and 20 iterations for

NUP76/NUP64 and NUP149/NUP64, respectively. A single plane image of one nucleus is

shown. Similar results were obtained for all NUP76 complex proteins (NUP76, NUP140,

NUP149) with sole antibody labelling (Figure S3).

Figure 2. A proximity map of the trypanosomatid nuclear pore.

(A) Mass spectrometry data (t-test difference values) from proximity labelling experiments

were used to create a heat map of the trypanosome nuclear pore. N- and C- terminally

tagged TurboID fusions of NUP110 (basket), NUP96 (inner ring) and NUP76 (cytoplasmic

outer ring) served as baits and the labelling (proximity) of all bona fide nuclear pore proteins

was analysed. The heat map grouped most NUPs into three main clusters, corresponding to

the basket/nuclear ring, inner ring and outer ring. Some NUPs with too little or no labelling

were manually added using data of (Obado *et al*, 2016). Only two NUPs appear potentially

misplaced (NUP132 and NUP53a) and two NUPs could not be placed (NUP48 and GILE2).

pLDDT plots from Trypanosomatid-optimized AlphaFold2 models (Wheeler, 2021) are

shown on the right.

(B and C) Mass spectrometry data (t-test difference values) from proximity labelling

experiments of NUP75-TurboID and TurboID-NUP158 are shown for all the NUPs shown in

A, using the same colour-code.

(D) The model of the trypanosome nuclear pore changes, with the discovery of five novel

asymmetrically localised proteins.

Figure 3. Mapping the trypanosome mRNA export system to the nuclear pore.

(A) Mass spectrometry data (t-test difference values) from proximity labelling experiments

with NUP110/NUP96/NUP76 were used to map the Ran-based mRNA export system to the

trypanosome pore.

(B and C) Ultra-expansion microscopy. Mex67, Ran, RanBP1, RanGAP and RanBPL were

expressed as C-terminal (Mex67, Ran RanBP1, RanGAP) or N-terminal (RanBPL) fusions to

TurboID in a cell line co-expressing the nuclear-site marker NUP64-3HA; all from the

endogenous locus. Cells were expanded and the proteins detected with streptavidin and

anti-HA. Images of single nuclei are shown (single plane of deconvolved Z-stacks with 10

iterations for Mex67, Ran, RanBP1 and RanBPL and 20 iterations for RanGAP). For Mex67,

the streptavidin signal was weak and three representative nuclei of an overexpression cell

line are shown (B, right). For the other proteins, enlarged section of the nuclear envelope of

the same or other nuclei are shown (C, right).

(D) Proximity labelling of the nuclear pore by Mex67 and Ran. Mex67 and Ran were

expressed as carboxy-terminal fusions to TurboID from the endogenous loci and biotinylated

peptides were analysed by mass spectrometry. The labelling of all bona-fide NUPs by

Mex67 (left) and Ran (right) is shown coloured based on their *t*-test difference values in

comparison to wild type cells. Asymmetric NUPs are marked with a red arrow.

(E) Schematic summary of the UExM localisation data of the Ran-based mRNA export

system.

Figure 4. Characterisation of unknown nuclear pore proteins and their localisation

over the nuclear pore complex.

(A) Mass spectrometry data (t-test difference values) from proximity labelling experiments

with NUP110/NUP96/NUP76/Mex67/Ran/NUP158 were screened for labelling of nuclear

pore localised proteins, that are not bona fide NUPs or karyopherins. All proteins that were

labelled by at least one of the baits proteins are shown. Annotations are explained in the

text. pLDDT plots from Trypanosomatid-optimized AlphaFold2 models (Wheeler, 2021) are

shown on the right.

(B-D) Structural alignments of AlphaFold2 models of the *T. brucei* TREX2 complex

candidates Sac3 (B), Thp1 (C) and Sus1 (D) with PDB structures of the respective TREX2

complex proteins from other organisms, using Foldseek (Kempen *et al*, 2024). The regions

of the proteins that were used for the structural alignments are shown in the schematics in

orange (*T. brucei*) and blue (other organisms). The root mean square deviation of atomic

positions (RMSD) and the sequence identity are shown for the superimposed regions.

Figure 5: The *T. brucei* NUP76 complex is only partially conserved

(A) Schematics of the cytoplasmic filament complex from yeast and human (modified from

(Bley *et al*, 2022), not to scale. The proteins of the trypanosome NUP76 complex are shown

for comparison (left). Note that trypanosomes do have orthologues to NUP145N and Gle2,

but it is not known whether these directly interact with NUP76.

(B) pAE and pLLDT plots of trypanosomatid-optimised AlphaFold2 models of NUP76,

NUP140 and NUP149. Each protein is also shown schematically with all predicted domains

and, for NUP140 and NUP149, with positions and types of FG repeats.

(C) Models of trypanosomatid-optimised AlphaFold2 predictions of NUP76, NUP140 and

NUP149. Structured parts are coloured, disordered regions are shown in grey.

Figure 6: Depletion of TbNUP76 causes nuclear poly(A) accumulation and loss of

NUP140 pore localisation

NUP76 protein was depleted using a degron system based on induction with the auxin

derivative 5-Ph-IAA. Both alleles of NUP76 were replaced by NUP76 fused to OsAID-3HA at

the C-terminus.

(A) The depletion of the NUP76 protein at 2, 4, and 24 hours of induction was monitored on

a western blot using anti-HA to detect NUP76 and anti-PFR as loading control. Wild type

(WT) cells served as negative controls. Data of one representative clonal cell line are shown.

(B) Growth was monitored over 4 days following NUP76 depletion. Data of three

independent clonal cell lines are shown.

(C) In situ hybridisation: cells were probed with oligo dT to monitor mRNA localisation. The

DNA is labelled with DAPI. One representative cell is shown for untreated cells and cells

after 2 and 4 hours of NUP76 depletion.

(D and E) An N-terminal eYFP fusion of NUP96 and C-terminal eYFP fusions of NUP140,

Nup149 and Gle2 were expressed from endogenous loci in the NUP76 degron cell line. The

eYFP fluorescence of seven randomly selected nuclei is shown before and after induction of

NUP76 depletion. Additional nuclei are shown in Figure S12.

SUPPLEMENTAL FIGURES

Figure S1: Establishment and validation of the expansion microscopy protocols.

Figure S2: Additional proExM images of the NUP76 complex proteins.

Figure S3. UExM images of NUP64-4Ty1 with NUP76 complex proteins tagged with 3xHA (labelled using antibodies).

Figure S4: Statistical analysis of TurboID experiments

Figure S5: Inducible overexpression of Mex67 (growth curves and Western blotting to check for expression levels).

Figure S6: Nucleolus can be identified by reduction in DAPI stain.

Figure S7: Mapping proteins with the proximity map: Karyopherins.

Figure S8: Mapping proteins with the proximity map: unlabelled proteins.

Figure S9: Structures of yeast NUP82, human NUP88 and predicted *T. brucei* NUP76.

Figure S10: The NUP149 zinc fingers in comparison to the ones from human NUP153 and NUP358.

Figure S11: Verification of NUP76 auxin degron cell lines by diagnostic PCR.

Figure S12: Effect of NUP76 depletion on NUP140, NUP149, Gle2 and NUP96 localisation: additional images.

Figure S13: AlphaFold2 models of RanBP1 and RanBPL

ACKNOWLEDGEMENTS

The project was funded by a bilateral GACR/DFG grant (project IDs.: 21-19503J and KR4017/9-1; to M. Z. and S. K., respectively) and the DFG grant KR4017/8-1 to S.K. We are grateful to the OMICS Proteomics BIOCEV core facility for excellent technical service. We like to thank Eva Kowalinski (EMBL, Grenoble, France) for expert help with Alphafold and Colabfold. We are grateful to Mark Carrington for providing the highly useful auxin degron system.

Disclosure and competing interests statement

The authors declare that they have no conflict of interest.

LITERATURE

Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L, Ballard AJ, Bambrick J, *et al* (2024) Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* 630: 493–500

Akey CW, Singh D, Ouch C, Echeverria I, Nudelman I, Varberg JM, Yu Z, Fang F, Shi Y, Wang J, *et al* (2022) Comprehensive structure and functional adaptations of the yeast nuclear pore complex. *Cell* 185: 361-378.e25

Allegretti M, Zimmerli CE, Rantos V, Wilfling F, Ronchi P, Fung HKH, Lee C-W, Hagen W, Turoňová B, Karius K, *et al* (2020) In-cell architecture of the nuclear pore and snapshots of its turnover. *Nature* 586: 796–800

Alsford S, Turner D, Obado S, Sanchez-Flores A, Glover L, Berriman M, Hertz-Fowler C & Horn D (2011) High throughput phenotyping using parallel sequencing of RNA interference targets in the African trypanosome. *Genome Research*

Anderson J, Phan L, Cuesta R, Carlson BA, Pak M, Asano K, Björk GR, Tamame M & Hinnebusch AG (1998) The essential Gcd10p–Gcd14p nuclear complex is required for 1-methyladenosine modification and maturation of initiator methionyl-tRNA. *Genes Dev* 12: 3650–3662

Anderson J, Phan L & Hinnebusch AG (2000) The Gcd10p/Gcd14p complex is the essential two-subunit tRNA(1-methyladenosine) methyltransferase of Saccharomyces cerevisiae. *Proc Natl Acad Sci* 97: 5173–5178

Ashkenazy-Titelman A, Shav-Tal Y & Kehlenbach RH (2020) Into the basket and beyond: the journey of mRNA through the nuclear pore complex. *The Biochemical journal* 477: 23–44

Aslett M, Aurrecoechea C, Berriman M, Brestelli J, Brunk BP, Carrington M, Depledge DP, Fischer S, Gajria B, Gao X, *et al* (2010) TriTrypDB: a functional genomic resource for the Trypanosomatidae. *Nucleic Acids Research* 38: D457-62

Bastin P, Bagherzadeh Z, Matthews KR & Gull K (1996) A novel epitope tag system to study protein targeting and organelle biogenesis in Trypanosoma brucei. 77: 235–239

Billington K, Halliday C, Madden R, Dyer P, Barker AR, Moreira-Leite FF, Carrington M, Vaughan S, Hertz-Fowler C, Dean S, *et al* (2023) Genome-wide subcellular protein map for the flagellate parasite Trypanosoma brucei. *Nat Microbiol* 8: 533–547

Bley CJ, Nie S, Mobbs GW, Petrovic S, Gres AT, Liu X, Mukherjee S, Harvey S, Huber FM, Lin DH, *et al* (2022) Architecture of the cytoplasmic face of the nuclear pore. *Science* 376: eabm9129

Branon TC, Bosch JA, Sanchez AD, Udeshi ND, Svinkina T, Carr SA, Feldman JL, Perrimon N & Ting AY (2018) Efficient proximity labeling in living cells and organisms with TurboID. *Nature biotechnology* 36: 880–887

Brasseur A, Bayat S, Chua XL, Zhang Y, Zhou Q, Low BC & He CY (2014) The bi-lobe-associated LRRP1 regulates Ran activity in Trypanosoma brucei. *Journal of Cell Science* 127: 4846–4856

Brun R & Schönenberger (1979) Cultivation and in vitro cloning or procyclic culture forms of Trypanosoma brucei in a semi-defined medium. Short communication. *Acta tropica* 36: 289–292

Buhlmann M, Walrad P, Rico E, Ivens A, Capewell P, Naguleswaran A, Roditi I & Matthews KR (2015) NMD3 regulates both mRNA and rRNA nuclear export in African trypanosomes via an XPOI-linked pathway. *Nucleic Acids Research*

Burkard GS, Jutzi P & Roditi I (2011) Genome-wide RNAi screens in bloodstream form trypanosomes identify drug transporters. 175: 91–94

Butterfield ER, Obado SO, Scutts SR, Zhang W, Chait BT, Rout MP & Field MC (2024) A lineage-specific protein network at the trypanosome nuclear envelope. *Nucleus* 15: 2310452

Chen S, Jiang Q, Fan J & Cheng H (2024) Nuclear mRNA export. *Acta Biochim Biophys Sin*

Clark MW & Abelson J (1987) The subnuclear localization of tRNA ligase in yeast. *J cell Biol* 105: 1515–1526

Cox J, Hein MY, Luber CA, Paron I, Nagaraj N & Mann M (2014) Accurate Proteome-wide Label-free Quantification by Delayed Normalization and Maximal Peptide Ratio Extraction, Termed MaxLFQ*. *Mol Cell Proteomics* 13: 2513–2526

Cox J & Mann M (2008) MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nature biotechnology* 26: 1367–1372

Dean S, Sunter J, Wheeler RJ, Hodkinson I, Gluenz E & Gull K (2015) A toolkit enabling efficient, scalable and reproducible gene tagging in trypanosomatids. *Open biology* 5: 140197

DeGrasse JA, DuBois KN, Devos D, Siegel TN, Sali A, Field MC, Rout MP & Chait BT (2009) Evidence for a Shared Nuclear Pore Complex Architecture That Is Conserved from the Last Common Eukaryotic Ancestor. *Molecular & Cellular Proteomics* 8: 2119–2130

Dostalova A, Käser S, Cristodero M & Schimanski B (2013) The nuclear mRNA export receptor Mex67-Mtr2 of Trypanosoma brucei contains a unique and essential zinc finger motif. *Molecular Microbiology* 88: 728–739

Ellisdon AM, Dimitrova L, Hurt E & Stewart M (2012) Structural basis for the assembly and nucleic acid binding of the TREX-2 transcription-export complex. *Nat Struct Mol Biol* 19: 328–336

Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, Žídek A, Bates R, Blackwell S, Yim J, *et al* (2022) Protein complex prediction with AlphaFold-Multimer. *bioRxiv*: 2021.10.04.463034

Fernandez-Martinez J, Kim SJ, Shi Y, Upla P, Pellarin R, Gagnon M, Chemmama IE, Wang J, Nudelman I, Zhang W, *et al* (2016) Structure and Function of the Nuclear Pore Complex Cytoplasmic mRNA Export Platform. *Cell* 167: 1215-1228.e25

Fernandez-Martinez J & Rout MP (2021) One Ring to Rule them All? Structural and Functional Diversity in the Nuclear Pore Complex. *Trends Biochem Sci* 46: 595–607

Gabernet-Castello C, O'Reilly AJ, Dacks JB & Field MC (2013) Evolution of Tre-2/Bub2/Cdc16 (TBC) Rab GTPase-activating proteins. *Molecular biology of the cell* 24: 1574–1583

Gabiatti BP, Freire ER, Odenwald J, Holetz F, Carrington M, Kramer S & Zoltner M (2024) Intron-
loss in Kinetoplastea correlates with a non-functional EJC and loss of NMD factors.

Galy V, Gadal O, Fromont-Racine M, Romano A, Jacquier A & Nehrbass U (2004) Nuclear retention
of unspliced mRNAs in yeast is mediated by perinuclear Mlp1. *Cell* 116: 63–73

Gambarotto D, Zwettler FU, Guennec ML, Schmidt-Cernohorska M, Fortun D, Borgers S, Heine J,
Schloetel JG, Reuss M, Unser M, *et al* (2019) Imaging cellular ultrastructures using expansion
microscopy (U-ExM). *Nat Methods* 16: 71–74

Gorilak P, Pružincová M, Vachova H, Olšinová M, Cernohorska MS & Varga V (2021) Expansion
microscopy facilitates quantitative super-resolution studies of cytoskeletal structures in
kinetoplastid parasites. *Open Biol* 11: 210131

Grandi P, Emig S, Weise C, Hucho F, Pohl T & Hurt EC (1995) A novel nuclear pore protein Nup82p
which specifically binds to a fraction of Nsp1p. *J cell Biol* 130: 1263–1273

Gu Z (2022) Complex heatmap visualization. *iMeta* 1: e43

Güther MLS, Urbaniak MD, Tavendale A, Prescott A & Ferguson MAJ (2014) High-confidence
glycosome proteome for procyclic form Trypanosoma brucei by epitope-tag organelle enrichment
and SILAC proteomics. *Journal of proteome research* 13: 2796–2806

Hampoelz B, Andres-Pons A, Kastritis P & Beck M (2019) Structure and Assembly of the Nuclear
Pore Complex. *Annu Rev Biophys* 48: 1–22

Hang J & Dasso M (2002) Association of the Human SUMO-1 Protease SENP2 with the Nuclear
Pore*. *J Biol Chem* 277: 19961–19966

Hernández MA de, Peralta GM, Vena R & Alonso VL (2023) Ultrastructural Expansion Microscopy
in Three In Vitro Life Cycle Stages of Trypanosoma cruzi. *J Vis Exp : JoVE*

Higa MM, Alam SL, Sundquist WI & Ullman KS (2007) Molecular Characterization of the Ran-
binding Zinc Finger Domain of Nup153*. *J Biol Chem* 282: 17090–17100

Holzer G & Antonin W (2022) Nup50 plays more than one instrument. *Cell Cycle* 21: 1785–1794

Hopper AK, Traglia HM & Dunst RW (1990) The yeast RNA1 gene product necessary for RNA
processing is located in the cytosol and apparently excluded from the nucleus. *J cell Biol* 111:
309–321

Hurwitz ME & Blobel G (1995) NUP82 is an essential yeast nucleoporin required for poly(A)+ RNA
export. *J Cell Biol* 130: 1275–1281

Hutchins JR, Moore WJ & Clarke PR (2009) Dynamic localisation of Ran GTPase during the cell
cycle. *BMC Cell Biol* 10: 66

Jani D, Lutz S, Marshall NJ, Fischer T, Köhler A, Ellisdon AM, Hurt E & Stewart M (2009) Sus1,
Cdc31, and the Sac3 CID region form a conserved interaction platform that promotes nuclear pore
association and mRNA export. *Molecular Cell* 33: 727–737

Jani D, Valkov E & Stewart M (2014) Structural basis for binding the TREX2 complex to nuclear
pores, GAL1 localisation and mRNA export. *Nucleic Acids Research* 42: 6686–6697

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, *et al* (2021) Highly accurate protein structure prediction with AlphaFold. *Nature* 596: 583–589

Kalichava A & Ochsenreiter T (2021) Ultrastructure expansion microscopy in Trypanosoma brucei. *Open Biol* 11: 210132

Kelley LA, Mezulis S, Yates CM, Wass MN & Sternberg MJE (2015) The Phyre2 web portal for protein modeling, prediction and analysis. *Nat Protoc* 10: 845–858

Kempen M van, Kim SS, Tumescheit C, Mirdita M, Lee J, Gilchrist CLM, Söding J & Steinegger M (2024) Fast and accurate protein structure search with Foldseek. *Nat Biotechnol* 42: 243–246

Kim SJ, Fernandez-Martinez J, Nudelman I, Shi Y, Zhang W, Raveh B, Herricks T, Slaughter BD, Hogan JA, Upla P, *et al* (2018) Integrative structure and functional anatomy of a nuclear pore complex. *Nature* 555: 475–482

KOHL L, SHERWIN T & GULL K (1999) Assembly of the Paraflagellar Rod and the Flagellum Attachment Zone Complex During the Trypanosoma brucei Cell Cycle. *J Eukaryot Microbiol* 46: 105–109

Kose S, Furuta M & Imamoto N (2012) Hikeshi, a Nuclear Import Carrier for Hsp70s, Protects Cells from Heat Shock-Induced Nuclear Damage. *Cell* 149: 578–589

Kosinski J, Mosalaganti S, Appen A von, Teimer R, DiGuilio AL, Wan W, Bui KH, Hagen WJH, Briggs JAG, Glavy JS, *et al* (2016) Molecular architecture of the inner ring scaffold of the human nuclear pore complex. *Science (New York, NY)* 352: 363–365

Kramarz K, Schirmeisen K, Boucherit V, Saada AA, Lovo C, Palancade B, Freudenreich C & Lambert SAE (2020) The nuclear pore primes recombination-dependent DNA synthesis at arrested forks by promoting SUMO removal. *Nat Commun* 11: 5643

Kramer S, Kimblin NC & Carrington M (2010) Genome-wide in silico screen for CCCH-type zinc finger proteins of Trypanosoma brucei, Trypanosoma cruzi and Leishmania major. *BMC genomics* 11: 283

Li Z, Chen S, Zhao L, Huang G, Xu H, Yang X, Wang P, Gao N & Sui S-F (2023) Nuclear export of pre-60S particles through the nuclear pore complex. *Nature* 618: 411–418

Lin DH & Hoelz A (2019) The Structure of the Nuclear Pore Complex (An Update). *Annu Rev Biochem* 88: 1–59

Lin DH, Stuwe T, Schilbach S, Rundlet EJ, Perriches T, Mobbs G, Fan Y, Thierbach K, Huber FM, Collins LN, *et al* (2016) Architecture of the symmetric core of the nuclear pore. *Science (New York, NY)* 352: aaf1015

Lindsay ME, Plafker K, Smith AE, Clurman BE & Macara IG (2002) Npap60/Nup50 Is a Tri-Stable Switch that Stimulates Importin-α:β-Mediated Nuclear Protein Import. *Cell* 110: 349–360

Lo Y-H, Sobhany M, Hsu AL, Ford BL, Krahn JM, Borgnia MJ & Stanley RE (2019) Cryo-EM structure of the essential ribosome assembly AAA-ATPase Rix7. *Nat Commun* 10: 513

Mackmull M, Klaus B, Heinze I, Chokkalingam M, Beyer A, Russell RB, Ori A & Beck M (2017)
Landscape of nuclear transport receptor cargo specificity. *Mol Syst Biol* 13: 962

Magistris PD (2021) The Great Escape: mRNA Export through the Nuclear Pore Complex. *Int J Mol*
Sci 22: 11767

Mahajan R, Delphin C, Guan T, Gerace L & Melchior F (1997) A Small Ubiquitin-Related
Polypeptide Involved in Targeting RanGAP1 to Nuclear Pore Complex Protein RanBP2. *Cell* 88:
97–107

Mahajan R, Gerace L & Melchior F (1998) Molecular Characterization of the SUMO-1 Modification
of RanGAP1 and Its Role in Nuclear Envelope Association. *J Cell Biol* 140: 259–270

Maishman L, Obado SO, Alsford S, Bart J-M, Chen W-M, Ratushny AV, Navarro M, Horn D,
Aitchison JD, Chait BT, *et al* (2016) Co-dependence between trypanosome nuclear lamina
components in nuclear stability and control of gene expression. *Nucleic Acids Research* 44:
10554–10570

Makarov AA, Padilla-Mejia NE & Field MC (2021) Evolution and diversification of the nuclear pore
complex. *Biochem Soc Trans* 49: 1601–1619

Matunis MJ, Coutavas E & Blobel G (1996) A novel ubiquitin-like modification modulates the
partitioning of the Ran-GTPase-activating protein RanGAP1 between the cytosol and the nuclear
pore complex. *J cell Biol* 135: 1457–1470

Matunis MJ, Wu J & Blobel G (1998) SUMO-1 Modification and Its Role in Targeting the Ran
GTPase-activating Protein, RanGAP1, to the Nuclear Pore Complex. *J Cell Biol* 140: 499–509

Melchior F, Weber K & Gerke V (1993) A functional homologue of the RNA1 gene product in
Schizosaccharomyces pombe: purification, biochemical characterization, and identification of a
leucine-rich repeat motif. *Mol Biol Cell* 4: 569–581

Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH & Ferrin TE (2023) UCSF
ChimeraX: Tools for structure building and analysis. *Protein Sci* 32: e4792

Moreira CM do N, Kelemen CD, Obado SO, Zahedifard F, Zhang N, Holetz FB, Gauglitz L,
Dallagiovanna B, Field MC, Kramer S, *et al* (2023) Impact of inherent biases built into proteomic
techniques: Proximity labeling and affinity capture compared. *J Biol Chem* 299: 102726

Mosalaganti S, Kosinski J, Albert S, Schaffer M, Strenkert D, Salomé PA, Merchant SS, Plitzko JM,
Baumeister W, Engel BD, *et al* (2018) In situ architecture of the algal nuclear pore complex. *Nat*
Commun 9: 2361

Nakielny S, Shaikh S, Burke B & Dreyfuss G (1999) Nup153 is an M9-containing mobile nucleoporin
with a novel Ran-binding domain. *EMBO J* 18: 1982–1995

Nie M & Boddy MN (2015) Pli1PIAS1 SUMO Ligase Protected by the Nuclear Pore-associated
SUMO Protease Ulp1SENP1/2 *. *J Biol Chem* 290: 22678–22685

Niepel M, Strambio-de-Castillia C, Fasolo J, Chait BT & Rout MP (2005) The nuclear pore complex–
associated protein, Mlp2p, binds to the yeast spindle pole body and promotes its efficient
assembly. *J Cell Biol* 170: 225–235

Obado SO, Brillantes M, Uryu K, Zhang W, Ketaren NE, Chait BT, Field MC & Rout MP (2016) Interactome Mapping Reveals the Evolutionary History of the Nuclear Pore Complex. *PLoS biology* 14: e1002365

Obado SO, Field MC & Rout MP (2017) Comparative interactomics provides evidence for functional specialization of the nuclear pore complex. *Nucleus (Austin, Tex)* 8: 1–13

Obado SO, Rout MP & Field MC (2022a) Sending the message: specialized RNA export mechanisms in trypanosomes. *Trends Parasitol* 38: 854–867

Obado SO, Stein M, Hegedűsová E, Zhang W, Hutchinson S, Brillantes M, Glover L, Paris Z, Chait BT, Field MC, *et al* (2022b) Mex67 paralogs mediate division of labor in trypanosome RNA processing and export. *bioRxiv*: 2022.06.27.497849

Odenwald J, Gabiatti B, Braune S, Shen S, Zoltner M & Kramer S (2024) Detection of TurboID fusion proteins by fluorescent streptavidin outcompetes antibody signals and visualises targets not accessible to antibodies. *eLife* 13: RP95028

Padilla-Mejia NE & Field MC (2023) Evolutionary, structural and functional insights in nuclear organisation and nucleocytoplasmic transport in trypanosomes. *FEBS Lett* 597: 2501–2518

Papai G, Frechard A, Kolesnikova O, Crucifix C, Schultz P & Ben-Shem A (2020) Structure of SAGA and mechanism of TBP deposition on gene promoters. *Nature* 577: 711–716

Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, Bileschi ML, Bork P, Bridge A, Colwell L, *et al* (2022) InterPro in 2022. *Nucleic Acids Res* 51: D418–D427

Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ, Inuganti A, Griss J, Mayer G, Eisenacher M, *et al* (2019) The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic Acids Research* 47: D442–D450

Rose A & Meier I (2001) A domain unique to plant RanGAP is responsible for its targeting to the plant nuclear rim. *Proc Natl Acad Sci* 98: 15377–15382

Rotureau B, Gego A & Carme B (2005) Trypanosomatid protozoa: A simplified DNA isolation procedure. *Exp Parasitol* 111: 207–209

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, *et al* (2012) Fiji: an open-source platform for biological-image analysis. *Nature methods* 9: 676–682

Schneider M, Hellerschmied D, Schubert T, Amlacher S, Vinayachandran V, Reja R, Pugh BF, Clausen T & Köhler A (2015) The Nuclear Pore-Associated TREX-2 Complex Employs Mediator to Regulate Gene Expression. *Cell* 162: 1016–1028

Schwartz TU (2016) The Structure Inventory of the Nuclear Pore Complex. *J Mol Biol* 428: 1986–2000

Schwede A, Manful T, Jha BA, Helbig C, Bercovich N, Stewart M & Clayton CE (2009) The role of deadenylation in the degradation of unstable mRNAs in trypanosomes. *Nucleic Acids Research* 37: 5511–5528

Seewald MJ, Kraemer A, Farkasovsky M, Kürner C, Wittinghofer A & Vetter IR (2003) Biochemical
Characterization of the Ran-RanBP1-RanGAP System: Are RanBP Proteins and the Acidic Tail of
RanGAP Required for the Ran-RanGAP GTPase Reaction? *Mol Cell Biol* 23: 8124–8136

Shanmugasundram A, Starns D, Böhme U, Amos B, Wilkinson PA, Harb OS, Warrenfeltz S,
Kissinger JC, McDowell MA, Roos DS, *et al* (2023) TriTrypDB: An integrated functional
genomics resource for kinetoplastida. *PLOS Neglected Trop Dis* 17: e0011058

Simos G, Tekotte H, Grosjean H, Segref A, Sharma K, Tollervey D & Hurt EC (1996) Nuclear pore
proteins are involved in the biogenesis of functional tRNA. *EMBO J* 15: 2270–84

Singh D, Soni N, Hutchings J, Echeverria I, Shaikh F, Duquette M, Suslov S, Li Z, Eeuwen T van,
Molloy K, *et al* (2024) The molecular architecture of the nuclear basket. *Cell*

Stankunas E & Köhler A (2024) Docking a flexible basket onto the core of the nuclear pore complex.
Nat Cell Biol 26: 1504–1519

Stewart M (2020) Macromolecular Protein Complexes II: Structure and Function. *Subcell Biochem*
93: 461–470

Sunter J, Wickstead B, Gull K & Carrington M (2012) A new generation of T7 RNA polymerase-
independent inducible expression plasmids for Trypanosoma brucei. *PLoS ONE* 7: e35167

Tang J, Jia P, Xin P, Chu J, Shi D-Q & Yang W-C (2020) The Arabidopsis TRM61/TRM6 complex is
a bona fide tRNA N1-methyladenosine methyltransferase. *J Exp Bot* 71: 3024–3036

Terry LJ & Wente SR (2009) Flexible Gates: Dynamic Topologies and Functions for FG
Nucleoporins in Nucleocytoplasmic Transport. *Eukaryot Cell* 8: 1814–1827

Tillberg PW, Chen F, Piatkevich KD, Zhao Y, Yu C-C (Jay), English BP, Gao L, Martorell A, Suk H-
J, Yoshida F, *et al* (2016) Protein-retention expansion microscopy of cells and tissues labeled
using standard fluorescent proteins and antibodies. *Nat Biotechnol* 34: 987–992

Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, Mann M & Cox J (2016) The
Perseus computational platform for comprehensive analysis of (prote)omics data. *Nature methods*
13: 731–740

Umlauf D, Bonnet J, Waharte F, Fournier M, Stierle M, Fischer B, Brino L, Devys D & Tora L
(2013) The human TREX-2 complex is stably associated with the nuclear pore basket. *Journal of*
Cell Science 126: 2656–2667

Wente SR & Rout MP (2010) The nuclear pore complex and nuclear transport. *Cold Spring Harbor*
Perspectives in Biology 2: a000562

Wheeler RJ (2021) A resource for improved predictions of Trypanosoma and Leishmania protein
three-dimensional structure. *Plos One* 16: e0259871

Wing CE, Fung HYJ & Chook YM (2022) Karyopherin-mediated nucleocytoplasmic transport. *Nat*
Rev Mol Cell Biol 23: 307–328

Wu J, Matunis MJ, Kraemer D, Blobel G & Coutavas E (1995) Nup358, a Cytoplasmically Exposed
Nucleoporin with Peptide Repeats, Ran-GTP Binding Sites, Zinc Fingers, a Cyclophilin A
Homologous Domain, and a Leucine-rich Region *. *J Biol Chem* 270: 14209–14213

Xie Y & Ren Y (2019) Mechanisms of nuclear mRNA export: A structural perspective. *Traffic* 20:
829–840

Xu XM, Meulia T & Meier I (2007) Anchorage of Plant RanGAP to the Nuclear Envelope Involves
Novel Nuclear-Pore-Associated Proteins. *Curr Biol* 17: 1157–1163

Yaseen NR & Blobel G (1999) Two distinct classes of Ran-binding sites on the nucleoporin Nup-358.
Proc Natl Acad Sci 96: 5516–5521

Zoltner M, Pino RCD & Field MC (2020) Sorting the Muck from the Brass: Analysis of Protein
Complexes and Cell Lysates. *Methods Mol Biology Clifton N J* 2116: 645–653

1175

Figure 1

Figure 2

Figure 3

Figure 4

A NUP76 depletion (5-Ph-IAA)

B

C mRNA (oligo dT)   DNA (DAPI)   merged

NUP76 depletion (5-Ph-IAA, hours)

D inner ring

E NUP76 complex and putative interactors