A single nucleotide polymorphism genotyping panel for efficient genetic stock identification of the Teno river Atlantic salmon (*Salmo salar*) population complex

Helena Johansson[1,2], Jaakko Erkinaro[2], Morten Falkegård[4], Anders Foldvik[4,] Panu Orell[2], Craig R Primmer[1,5], Victoria L Pritchard[1,3]

1- Organismal and Evolutionary Biology Research Program, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland
2- Natural Resources Institute, Oulu/Helsinki, Finland
3- UHI Institute for Biodiversity & Freshwater Conservation, Inverness, U.K.
4- Norwegian Institute for Nature Research (NINA), Tromsø/Trondheim, Norway
5- Institute of Biotechnology, Helsinki Institute of Life Science (HiLIFE), University of Helsinki, Finland

Corresponding author: Craig Primmer, Viikinkaari 9, 00014, Helsinki, Finland, craig.primmer@helsinki.fi

Funding: The study was supported by NINA, LUKE and the Academy of Finland

Abstract

A new single nucleotide polymorphism (SNP) panel for genetic stock identification in the Teno River Atlantic salmon fishery was developed, with a view to improving on an existing microsatellite panel. Twenty-two genetically differentiated reporting units were proposed for the Teno based on population genetic analyses of 1212 individuals collected at 37 locations in the river and genotyped for > 33,000 genome-wide SNPs. A small subset of these SNPs was selected for GSI using an iterative process that considered their diversity and differentiation across reporting units. A genotyping-by-sequencing assay was developed to simultaneously genotype 180 of these GSI SNPs plus a sexing marker. This new SNP panel showed comparative GSI power to the microsatellite panel, with anticipated improvements in terms of cost, speed, and robustness and transferability across laboratories and genotyping platforms. Mixed stock analysis of the 2018 Teno River salmon catch using the new panel inferred all 22 reporting units to contribute to the fishery. Estimated catch proportions positively scaled with an independent estimate of reporting unit productivity: target spawning female biomass. This demonstrates the usefulness and efficiency of the 180 SNP panel for Atlantic salmon GSI in the Teno river system. If conducted on a regular basis, GSI can enable fine-tuning of management strategies to promote sustainable fishing.

Keywords: fisheries, management, Atlantic salmon, genotyping by sequencing, population structure

Introduction

Fisheries often consist of a mix of stocks – that is, they harvest individuals that originate from multiple genetically distinct populations. Key for effective management of such mixed-stock fisheries is accurate knowledge of how different source populations contribute to the catch (Begg *et al.*, 1999; Palsbøll *et al.*, 2007). This enables stock-specific management strategies aimed at safeguarding vulnerable populations while allowing continued harvest of healthier stocks (Crozier *et al.*, 2004). Genetic stock identification (GSI) has been a mainstay of mixed-stock fisheries management for several decades (Anderson *et al.*, 2008; Koljonen & McKinnell, 1996; Winans *et al.*, 2004). The GSI approach involves first generating a genetic baseline of allele frequencies from populations that may contribute to the fishery. Samples from the mixed stock fishery are then genotyped with the same marker panel, and the likelihood of the baseline population allele frequencies producing the genotype in question in the fishery sample are estimated. using individual assignment-based approaches or else overall catch proportions are estimated using mixture modelling e.g. (Anderson *et al.*, 2008; Koljonen *et al.*, 2005).

In recent years, single nucleotide polymorphisms (SNPs) have replaced microsatellites as the marker of choice in a range of population and conservation genetic applications, including GSI e.g. (Larson *et al.*, 2014). Compared to microsatellites, which have commonly been typed by fragment size analysis, SNPs have advantages in terms of contemporary ease, speed and cost of genotyping, improved repeatability among laboratories and platforms, and a simpler mutation model allowing more straightforward interpretation of evolutionary and population genetic processes. One key benefit of SNPs compared to microsatellites for GSI is the potential availability of many thousands of easily genotyped loci, from which a sub-set of highly informative markers can be selected for a specific assignment task. This may enable more efficient stock assignment, improved resolution of more challenging population substructures (Larson *et al.*, 2014) and/or the inclusion of SNPs to address different research and management questions (Aykanat *et al.*, 2016; May *et al.*, 2020).

Atlantic salmon, *Salmo salar,* is an anadromous salmonid fish natively distributed across the north Atlantic coast of Europe and North America. Atlantic salmon are fished and managed across their range, and many wild populations are in long-term decline from a multitude of threats, some of which are poorly understood (Parrish *et al.*, 1998). Management of the species can be politically and logistically complicated, spanning national borders and involving conflicts about rights of use (Crozier *et al.*, 2004). One of the largest and most diverse wild Atlantic salmon (*Salmo salar*) populations in the world (Erkinaro *et al.*, 2019) breeds in the large Teno (Swedish: Tana, Sami: Deatnu) river system on the border of Finland and Norway in northernmost Europe. Previous genetic studies have revealed a highly structured population complex consisting of up to 28 genetically distinct, and temporally stable populations (Vähä *et al.*, 2011, 2017, 2007, 2008), with evidence for fine-scale local adaptation (Mobley *et al.*, 2019; Pritchard *et al.*, 2018). Until 2009, the riverine catch regularly reached 50,000 kg per year, however weak sea returns in recent years suggests that some stocks are struggling (Erkinaro *et al.*, 2019). To aid management, a microsatellite baseline (33 loci) was developed (Vähä *et al.*, 2017) that included over 3,300 individuals from 36 locations within the river system. Based on these data, a maximum of 32

reporting units for GSI was proposed, with an estimated assignment accuracy of 36-100%
depending on the reporting unit and analytical approach applied.

The aim of this study was to develop a SNP genotyping panel for Teno river salmon GSI, with a
view to improving on the microsatellite panel and producing a tool more robust to expected future
changes in genotyping technology. We first genotyped original (Vähä *et al.*, 2017) baseline
samples, augmented with a set of more recent samples, for over 38,000 genome-wide SNPs. We
used this SNP dataset to re-determine suitable reporting units and develop a 180 SNP genotyping-
by-sequencing (GT-seq) panel for GSI. We compared the assignment efficacy of the new GT-seq
panel with that of 10,000 genome-wide SNPs, and with the previous 33 microsatellite panel. The
new SNP panel was then used to perform mixed stock assignment of salmon caught in the Teno in
2018. As additional validation, GSI results were compared with an independent proxy of expected
population size, target female biomass.

Materials and Methods

Baseline samples and datasets

Archived DNA samples from juvenile salmon in the (Vähä *et al.*, 2017) baseline collection were
used as the basis of the new SNP baseline (Figure 1, Table S1). Samples were quantified using a
NanoDrop spectrophotometer (Thermo Scientific) and assessed for degradation using gel
electrophoresis. For two of the 32 baseline reporting units in (Vähä *et al.*, 2017) (Bavttájohka and
Anarjohka) there were either insufficient samples, the pre-extracted DNA was completely degraded
or the DNA did not pass the minimum quality requirements for SNP genotyping. To augment the
baseline, DNA was extracted and quantified for juvenile samples collected in 2014 from four
localities, Tsarsjoki, Lakšjohka, and the Lakšjohka tributaries Deavkkehanjohka and Gurtsjohka.
Good quality DNA samples with a minimum concentration of 20ng/ul were sent to the Center for
Integrative Genetics (CIGENE), Ås, Norway, for SNP typing on a custom 60,250 (60K) Atlantic
salmon SNP genotyping array. All subsequent merging and quality control of SNP datasets was
performed using Plink 1.90 (Chang *et al.*, 2015).

To further extend the data set, we added genetic data generated for previous studies of Teno salmon
(Barson *et al.*, 2015, Pritchard *et al.*, 2018): adult genotypes from one locality missing from the 60K
juvenile baseline (Máskejohka) and additional juvenile genotypes – also part of the original baseline
-from ten localities. These samples had been screened using a 220K SNP array, which shares
38,669 named SNPs with the 60K array. Forty-one individuals from (Pritchard *et al.*, 2018) were
typed on both arrays, enabling genotyping repeatability be examined. After merging the 60K and
220K datasets across the 38,669 markers, we removed 364 SNPs with which had $> 2/41$
mismatching genotypes across arrays and four SNPs with known off-target variants. Subsequently
we applied the following filters: SNP minor allele frequency >0.05; SNP missing genotype
frequency < 0.03; individual missing genotype frequency < 0.05. Finally, putative full-sibs were
identified and removed based on genome-wide patterns of identity-by-descent (--genome) within
and across collection locations.

Baseline population substructure

Before analyses of population structure, the baseline SNP dataset was further filtered to prune neighbouring SNPs in high linkage disequilibrium with one another (--indep 50 5 1.5). Population genetic analyses were carried out either in the R statistical environment (R Core Team, 2014), or in the Linux shell.

Overall and pairwise F_{ST}s among samples from different locations were estimated using *StAMPP* (Archer *et al.*, 2017). Significance of pairwise F_{ST} was assessed using 20,000 bootstrap replicates across loci and correcting for multiple testing using the Bonferroni procedure with alpha = 0.05. A neighbour-joining (NJ) tree based on Prevosti's distance (Prevosti *et al.*, 1975), was created and bootstrapped 100 times using *poppr* (Kamvar *et al.*, 2014) and the tree plotted using the package *ape* (Paradis & Schliep, 2019). Prevosti's distance is modelfree distance suitable for use with SNP data with missing genotypes.

Admixture (Alexander *et al.*, 2009) was used to identify the most likely number of genetically distinct ancestral populations present in the LD pruned baseline SNP dataset, independent of sampling location. Five replicates of the Admixture analysis were run, each starting from a different random number seed, allowing between 1 and 30 genetically distinct populations in the dataset. The most likely number of populations was determined using the software's cross-validation procedure. Finally, genetic variation within the dataset was also explored using principal component analysis followed by graphical visualization. PCA was performed using the package LEA in R (Frichot & Francois 2015), with number of retained PCs based on Tracy-Widom test results. The combined results from the Admixture and PCA analyses, interpreted with reference to (Vähä *et al.*, 2017) were used to select suitable new reporting units for GSI of the Teno River fishery.

Developing and testing of the GT-seq SNP panel

To identify a sub-set of SNP markers for efficient mixed stock assignment, highly variable SNPs (MAF > 0.45), were first extracted from the full baseline dataset using PLINK 1.9. Following this, genotypes from eight highly differentiated reporting units (Kevojoki, Tsarsjoki, Utsjoki, Lakšjohka, Deavkkehanjohka, Gurtsjohka, Yla-Pulmankijoki, Galddasjoki) were removed, as inclusion of these sites greatly skewed the F_{ST} distribution for SNPs, while assignment efficiency should be high for these reporting units regardless of the specific markers used. Closely linked SNPs in this new dataset were LD pruned (indep --50 5 1.5), and the remaining SNPs were ranked by calculating F_{ST} over the remaining reporting units. The 96 highest ranking SNPs were retained as the initial GSI panel. Filtering for highly variable SNPs retains a markers set that is expected to be informative across multiple reporting units, and subsequent selection of SNPs with the highest F_{ST} across units is expected to maximise accuracy for GSI (Ackerman *et al.*, 2011; Storer *et al.*, 2012). This initial SNP panel was tested for its assignment efficacy using the 'leave-one-out' individual re-assignment approach implemented in the R package *rubias* (Moran & Anderson, 2019). Markers were then added to improve discrimination among target subsets of reporting units by selecting additional SNPs with high MAF and high F_{ST} across those units alone. This process was performed iteratively, aiming for a final panel of 220-230 candidate GSI SNPs which was intended to translate into working panel of approximately 200 SNPs. As an additional check to minimize marker redundancy due to linkage, the positions of final candidate SNPs were inspected to exclude any markers within 10,000 bp of one another. Finally, SNPs linked to several genes identified as locally adaptive

candidates in the Teno River (Pritchard *et al.*, 2018) and the *sdY* sex marker (Yano *et al.*, 2013)
were added to the panel.

Primers for GT-Seq (Campbell *et al.*, 2015) were designed to amplify a 120-155bp DNA sequence
surrounding each of the target SNPs using BatchPrimer3 (Rozen & Skaletsky 1998;Untergasser *et*
al., 2012), with the following parameters: Tm 58-62°C optimum 60°C; GC% 28-68; all other
parameters default. Primers were not assessed *in silico* for dimerization potential.

All multiplex PCR was performed in 12ul reactions containing the following: 6ul Qiagen Multiplex
PCR mix, 2ul DNA at a concentration of 10-50 ng/ul, volume of multiplexed diluted primers to
yield the target concentration of each primer (see below); molecular grade water. Thermocycling
conditions were: 98°C for 2 min, 20x [98°C 10s, 62°C 30s, 72°C 15s], 7C for 10 min. PCR
products were tagged with individual-specific barcode indices, pooled, and sent to the DNA
Sequencing and Genomics Laboratory, University of Helsinki for library preparation and
sequencing. Products were single-end sequenced using Illumina150 cycle kits on an Illumina HiSeq
or MiSeq. Sequences were demultiplexed by barcode index using Generate FASTQ (Illumina) and
3' trimmed on the basis of indexing adaptor sequence using cutadapt 3.2 (Martin, 2011). Trimmed
sequences were aligned using bwa mem (BWA 0.7.17, Li & Durbin, 2010) to the *S. salar* reference
genome (ICSASG_v2) with an appended *S. salar* sdY sequence (Genbank Accession KT223110.1).
Variants at targeted sites were called using samtools mpileup piped to bcftools call (samtools 1.12,
Li & Barrett, 2011). The output VCF file was imported into Plink 1.9 with genotypes filtered for a
minimum GQ of 15 and a minimum sequencing depth of 8. For genetic sexing, sdY sequencing
depth was corrected for overall sequencing depth for each individual. Individuals were then
assigned as male (sdY present), female (sdY absent) or unknown on the basis of thresholds chosen
by examining the distribution of corrected sdY sequencing depths over all individuals.

Two rounds of multiplex PCR optimization were performed for the GT-Seq panel, each using 16
samples as the amplification target. For the first optimization round, all primers were combined in
equimolar concentrations (0.03 µM) in a single reaction. The output fastq files for the 16 test fish
were merged, and a custom bash script was used to count number of sequence reads for each
possible forward-reverse primer combination. This information was used to identify primers
producing dimer sequence, and primers were split into two multiplex pools that minimized within-
pool dimer formation. The subsequent optimization round was used to check pool composition and
to adjust primer concentrations (final range 0.03-0.06 µM) with the aim of roughly equalizing
sequencing coverage across SNPs. After these two rounds of optimisation the SNP panel was used
to genotype 74 baseline individuals to check for genotyping success and consistency of genotype
calling between GT-Seq and the 60K SNP array.

Assignment efficacy of SNP panel

The accuracy of GSI across identified Teno reporting units was tested using two methods
implemented in *rubias*: the 'leave one out' approach described above, and a '100% mixture'
approach. In the latter analysis, baseline allele frequencies are used to simulate the genotypes of fish
deriving from different populations and those fish are then assigned back to the full baseline. We
simulated 500 fish for each population and performed 100 replicates.

To compare the performance of the SNP panel to that which could be achieved by a much denser sampling of genetic variation across the genome, we randomly subsampled 10,000 SNPs from the baseline dataset and examined assignment power of this dataset via a leave-one-out analysis implemented in gsi_sim, a Linux command line version of *rubias* suitable for large datasets (https://github.com/eriqande/gsi_sim, (Anderson, 2010; Anderson *et al.*, 2008).

To assess the robustness of assignments with smaller numbers of SNPs from the SNP panel, and thus assess the effect of missing genotypes on GSI reliability, random subsets of the 180 SNPs (60, 80, 100, 120, 140 and 160 loci) were tested in 'leave-one-out' analyses against the final set of reporting units.

Comparison with the microsatellite baseline

We compared the efficacy of the new SNP GSI panel with that of the previous microsatellite panel. To do this, we used the existing Teno microsatellite baseline, which comprises 3,323 juvenile and adult salmon from 36 sites sampled within the Teno river system and typed at 33 microsatellite loci. Sampling and molecular protocols for this dataset can be found in (Vähä *et al.*, 2011, 2017, 2007, 2008). We compared the assignment performance of the SNPs and microsatellites using the *rubias* leave-one-out procedure under two reporting unit scenarios: the 22 units proposed in this paper and the 32 reporting units (two not represented in the SNP baseline) used by Vähä *et al.*, 2011.

Mixed stock assignment and validation

In 2018, scale samples from salmon caught in the Teno were collected by local fishers, coordinated by Luonnonvarakeskus (LUKE, Finland) and Norsk Institutt for Naturforskning (NINA, Norway). Fishers also collected standard data on the fish including visually determined sex and assessed whether they had escaped from fish farms. We extracted DNA from 1-2 scales for each fish (excluding two putative escapees) using a high-salt method, quantified extractions using NanoDrop, and standardized to a concentration of 10-50ng/µl. Samples from which sufficient DNA could be obtained were genotyped for the 180 GSI SNPs and sex marker as described above. Individuals missing genotypes at >60 loci were removed from the data set. Finally, we identified duplicate samples on the basis >of 90% genotype match with another sample using the close_matching_samples function in *rubias* and in each case excluded the duplicate with the fewest missing genotypes.

To estimate the proportional contribution of each of the 22 reporting units to the 2018 Teno catch, we used *rubias* to perform mixed stock analysis against the SNP baseline. We implemented the maximum likelihood approach with the default options of a uniform prior**,** 20000 sweeps and the first 1000 sweeps discarded as burn-in. We computed 95% credible intervals from the MCMC traces of mixing proportions.

To indirectly validate the results of the mixed stock analysis, we used location-specific figures for target spawning female biomass provided in (Falkegård *et al.*, 2014). This is the biomass of females required to produce the target number of eggs for each location, which is estimated by modelling habitat suitability and egg density estimates and taking stock-specific fecundity into account. It can be considered a proxy for the expected numbers of breeding adults returning to different parts of the Teno system. In (Falkegård *et al.*, 2014), tributaries were grouped by geography into somewhat

different groups than the reporting units that we use here. For the purpose of comparison, the mixed
stock analysis proportions were combined to match the (Falkegård *et al.*, 2014) geographic
groupings, and female biomass (kg) estimates were transformed into relative proportions of the total
for each group (Table S5). A Pearson correlation was performed to compare the estimated female
biomass proportions to the proportions from the mixed stock assignment using cor.test in base R.

Results

Genetic baseline

The initial genetic baseline comprised 1,309 samples, collected between 2006 and 2014 from 37
locations in the Teno (mean per location 35.3, range 11-67) and genotyped at 33,139 post-QC
SNPs. Following removal of putative full siblings and LD pruning, a total of 1,212 individuals
genotyped at 17,546 loci (Table S1) were used to investigate baseline population structure.

Global F_{ST} demonstrated substantial genetic structuring among sample sites across the Teno system
(F_{ST} =0.055, *P*=0.0099). Pairwise F_{ST}s ranged from 0.0005 (between Piltamo and Alaköngäs) to
0.1953 (between Deavkkehanjohka and Galddasjoki) and all were statistically significant, with the
exception of the comparison between Piltamo and Alaköngäs (Table S2). Nevertheless, the
generally very low pairwise F_{ST} among samples from Tana Mainstem localities (Alaköngäs,
Garnjaga, Kortsham, Luovtejohka, Outakoski, Piltamo, Sirma, Tana Bru and Yläköngäs) indicated
only weak genetic differentiation across most of the Teno mainstem.

The neighbour-joining tree reflected the pairwise F_{ST} results, showing a strong geographic signal
across sampling locations (Figure 2), with strong bootstrap support at many nodes.

Admixture inferred 21 ancestral clusters within the dataset (Figure 3), with a geographic signal in
line with the F_{ST} analysis and the neighbour-joining tree and indicative of gene flow among
neighbouring locations. The Teno mainstem and its two large headwater systems (Inari and
Iešjohka/ Kárášjohka) were separately dominated by three large ancestral clusters. The uppermost
sample from the mainstem (Outakosi) showed substantial mixing among these clusters. A fourth
genetic cluster dominated the lowermost mainstem sample, Tana Bru and its tributary Luovtejohka.
Ten smaller tributaries (Karigasjoki, Máskejohka, Galddasjoki, Ylä-Pulmankijoki, Utsjoki,
Kevojoki, Váljohka, Kietsimäjoki, Vetsijoki and Geáimmejohka) and two pairs of neighbouring
tributaries (Birkejohka/Kuoppilasjoki and Báišjohka/Nilijoki) were dominated by their own unique
ancestral clusters. Akujoki had a mixed ancestry largely from the neighbouring Karigasjoki and
Báišjohka/Nilijoki clusters. Two ancestral clusters were inferred for samples from Lakšjohka and its
two small tributaries Gurte- and Deavkkehanjohka; examination of the underlying data suggested
that this was likely caused by a segregating chromosomal rearrangement not accounted for by LD
pruning. The large sample from Tsarjoki was split into two clusters, which corresponded to year of
collection and probably reflect drift within this small tributary: these were amalgamated for
subsequent analyses.

Twenty three principal component were retained in the PCA. Graphical examination of the position
of individuals along these PCs (not shown) supported the conclusions of the Admixture analysis
with three exceptions: the Outakoski sample partly separated from the rest of the Teno mainstem
samples (Alaköngäs, Garnjarga, Kortsham, Piltamo, Sirma, Yläköngäs) along PC5, probably
reflecting the genetic mixing in this location, the Iesjohka and Karasjohka samples were separated

along PC23, and Akujoki was partly separated from other samples along PCX. Based on the
combined Admixture and PCA results, together with consideration of Vähä *et al.*, 2017, 22
reporting units were defined for Teno GSI (Table 2), with Lakš/ Gurte/ Deavkkehanjohka
considered a single reporting unit but Outakoski, Akujoki and Kárášjohka treated as separate
reporting units.

In some tributaries, a subset of juveniles were assigned very high Teno mainstem ancestry by the
Admixture analysis and clustered with mainstem fish in the PCA. Examination of available capture
locations showed that these 24 fish were sampled near tributary mouths; they were therefore
considered to be migrants from the Teno mainstem breeding population rather than part of the
genetically distinct tributary breeding population and were removed from the final GSI baseline.

SNP panel development

Primers were successfully designed for a final set of 223 candidate GSI SNPs, plus one marker each
for candidate adaptively important genes *vgll3*, *six6*, *numa1* and *btg3* and genotypic sex (sdY).
Following initial optimization and tests, SNPs were excluded due to complete non-amplification
(n=11), >10% mismatching genotypes between GT-Seq genotypes and SNP array genotypes (n=16,
mostly missing heterozygotes), and >40% missing genotypes (n= 21). One pair of primers
amplified two SNPs that were both included in the GSI panel. Primer sequences for the 180 SNPs
in the final GSI panel plus the sdY locus are provided in the Supplementary Materials.

Assignment efficacy

The leave-one-out analysis using the 180 SNP panel correctly assigned 86% of individuals to their
reporting unit of origin. (Table 1). Thirteen out of the 22 reporting units had assignment accuracies
between 90-100%, and only Outakoski, Tana Bru and Akujoki reporting units had assignment
success < 70%. Mis-assigned fish were generally assigned to geographically adjacent reporting
units (Table 1). Assessment of assignment efficacy using the 100% mixture approach gave very
similar results, with slightly better accuracy inferred for all reporting units (93% of simulated
individuals correctly assigned). (Table S3).

When testing both with the leave-one-out analysis, the 180 SNP panel performed almost as well as
10,000 genome-wide SNPs to assign samples to reporting unit, indicating that it is well captures
genome-wide patterns of population substructure (Table S4). Overall assignment accuracies
decreased gradually with each reduction in SNP number for sub-sets of the 180 GT-Seq panel
however even with 100 SNPs 80% of fish were assigned back to their correct reporting unit (Table
S4). Assignment success for the different size panels generally showed consistent patterns across
the same reporting units.

The same two analyses applied to the larger microsatellite baseline split by the new reporting units
demonstrated comparable assignment efficacy of the microsatellites (88% correct assignment using
the leave-one-out approach, 90% correct assignment for the 100% mixture approach, Table 1 &
S2). Similarly to the SNP panel the lowest assignment accuracy was for the Outakoski, Tana Bru
and Akujoki reporting units.

Mixed stock analysis

Of 2,116 scale samples received from the 2018 Teno mixed stock fishery, 187 failed genotyping
and a further 16 were identified as duplicate samples. This left 1,913 available for mixed stock
analysis against the 22 reporting unit SNP baseline (mean proportion of missing genotypes = 0.04,
range = 0 – 0.32) Results are shown in Table 2. Three of the largest reporting groups were each
assigned >10% of the fishery samples, totalling almost 55% of the total catch: Teno mainstem
(25.1%), Inari headwaters (15.9%) and Iešjohka (13.6%). Three additional reporting groups were
assigned >5% each: Kárášjohka (7.5%); Vetsijoki (7.4%) and Outakoski (5.9%) with the remaining
16 reporting groups each being assigned 3.2% or less of the catch (24.6% of the total catch). The
MSA proportions were highly and significantly correlated with the proportions of female biomass
target abundance (Figure 3, r= 0.97, p<0.000). About 9% fewer fish were assigned to Tana
Bru/Tana Mainstem/Outakoski than expected based on the female biomass target abundance
proportions, similarly, 4.5% less fish were assigned to Kárášjohka than expected. Conversely, about
5% more fish were caught from Vetsijoki, and 3% more from Utsjoki/Tsarsjoki/Kevojoki than
expected from the target abundance biomass estimates. All other localities had less than 2%
difference in proportions between MSA and target abundance female biomass estimates (Table S6).

Discussion

We have designed a new GT-seq panel of 180 SNPs for genetic stock identification within the large
Teno River Atlantic salmon fishery. The full panel performs equivalently to the previous
microsatellite panel, with advantages in terms of cost and speed of genotyping, transferability
among laboratories and flexibility for adding additional markers of interest. For example, in 2024
with an established pipeline and using a desktop short-read sequencer (Illumina MiSeq), generating
the 180 SNP genotypes from 1,000 DNA extracts is estimated to require five days of staff time and
€3,000 of laboratory consumables. It will also be more robust to anticipated future changes in
genotyping technologies – in particular, greatly reduced availability of platforms and reagents
currently used for microsatellite typing. Smaller sub-panels of the same SNPs also show good
performance indicating that the full panel will be robust to a proportion of missing genotypes, a
common outcome in GT-seq due to batch effects causing insufficient read depth over some targeted
sites.

Our re-investigation of population structure within the Teno River, for which we re-genotyped
archived samples for a large set of genome-wide SNPs, recapitulated the patterns previously found
using microsatellites. Examination of genome-wide variation can reveal population genetic
substructure that was not previously identified in studies using a smaller number of markers
(reviewed in Sunde *et al.* 2020). However, this was not the case for our study - the existing 33
microsatellites appear to have been sufficient to resolve the population genetic substructure within
the Teno Atlantic salmon population. We caution against a general assumption that examining more
variation across the genome will resolve increasingly more ecological- and management-relevant
substructure within a target species.

Vaha *et al.* (2008), genotyping archived scale samples, demonstrated that this broad-scale
population substructure in Teno Atlantic salmon had remained stable over a minimum of four
generations. Our results also support this, albeit on a smaller temporal scale: samples collected from
the same site 3-6 years apart almost invariably clustered together in the Admixture analysis and
PCA. Such temporally stable substructure is expected to arise and be perpetuated by the natal

homing of salmon to their birth river. This in turn can enable the establishment of local adapted genotypes, which further reduces gene flow among populations by reducing the fitness of migrants. There is both genetic and demographic evidence for local adaptation within the Teno salmon stock (Mobley *et al.*, 2019; Pritchard *et al.*, 2018), further emphasizing the importance of managing the fishery to avoid over-exploitation of certain subpopulations.

Despite this broad-scale stability of population structure, Vaha *et al.* (2008) did document change in allele frequencies across timepoints particularly in smaller tributary populations. We in turn find that samples collected in Tsarsjoki in 2006 and 2014 separate into two genetic clusters. Thus, any genetic baseline for Teno River GSI has the potential over time to become less representative of the contemporary allele frequencies in a subpopulation. We recommend additional juvenile collections to augment the baseline presented here, both to increase the sample size for poorly represented reporting units and to ensure that it accurately represents the contemporary genetic make-up of the Teno salmon stock. A long-term Teno monitoring programme should include plans to systematically update the baseline at pre-planned intervals and re-test the performance of the 180 SNP panel against the modified baseline.

Based on our examination of population structure, we chose to use 22 reporting units for the SNP-based GSI, fewer than the maximum of 32 that were used by (Vähä *et al.*, 2017). Specifically, we amalgamated Ylakongas with the Teno mainstem reporting unit, Luovtejohka with Tana Bru, Báišjohka with Nilojoki, Upper Iešjohka with Lower Iešjohka, and Iskurasjoki, Goššjohka, Vuomajoki and Cáščemjohka with the Inarijoki mainstem. Additionally, due to sample loss our baseline did not include two reporting units, Bavttájohka – a tributary of Kárášjohka and Anarjohka – a headwater tributary of Inarijoki. Nevertheless, investigation of the performance of the 180 SNP panel and the 33 microsatellite panel with the baseline divided into either 22 or 30 reporting units showed very similar results. Both demonstrated better overall assignment accuracy when there were fewer reporting units, while the SNPs actually performed better than the microsatellites for Báišjohka, Niljohka, Lower Iešjohka, Iskurasjoki and Goššjohka. Due to DNA degradation, our SNP baseline omitted two reporting units used by (Vähä *et al.*, 2017): Kárášjohka tributary Bavttájohka and Inarijoki tributary Anarjohka. Given the known patterns of genetic structure, we expect fish originating from these locations to be misassigned to a Kárášjohka/Iesjohka and an Inarijoki headwaters reporting unit, respectively. Bavttájohka contributes 1.8% of the target female biomass for the Teno, while the biomass of Anarjohka is included in the Inarijoki and not calculated separately; thus, while we recommend addition of these locations into the SNP baseline we do not anticipate that their current absence is substantially biasing our results.

Average accuracy rates for other Atlantic salmon GSI systems range from 87% (33 microsatellites, 182 rivers in the northernmost Norway and Finland and in the northwest of Russia; (Ozerov *et al.*, 2017), to 53.1 % at the level of the individual river and 72.1% at the regional level (31 microsatellite markers, 14 Norwegian rivers (Harvey *et al.*, 2019). Considering the spatial scale of the Teno basin, the resolution using this GSI system is very high. Further increases in accuracy would be possible by merging rivers into broader, regional, reporting units, however, for management purposes tributary-based reporting units are more informative.

We caution that results from the microsatellites and SNP panel are not entirely comparable. In addition to differences in baseline sample sizes, the panel of GSI SNPs were selected and tested

using the same individuals, an ascertainment bias that is expected to somewhat over-estimate their true assignment efficacy. We used this approach instead of dividing the baseline into training and test datasets to more accurately capture the true population allele frequency where we had low numbers of samples. Nevertheless, the proportion of the Teno River 2018 salmon catch assigned to the 22 baseline reporting units in the Mixed Stock Analysis closely scaled with the target female spawning biomass of those reporting units, suggesting that the SNP panel is indeed performing well: only three comparisons deviated more than 2% from the expected. Salmon are not fished at random from the rivers in the Tana river system; fishing permits are required for the different fish stocks, and the quota that can be fished in any given year is determined based on the vulnerability of overexploitation of the different stocks (Anon., 2020). Based on the comparison in this study alone, the outtake of fish in 2018 reflected quite accurately the potential size in female biomass of the populations (Falkegård *et al.*, 2014).

In conclusion, we present a new genetic tool for adaptive management of the Teno River Atlantic salmon fishery, which targets one of the largest wild salmon stocks remaining in Europe. These genotyping-by-sequencing SNP panel performs equivalently to a previous microsatellite panel and could be easily modified with the addition of new markers. The new panel could also form the foundation of a more broadly applicable panel for Genetic Stock Identification of Atlantic salmon throughout northern Europe.

Acknowledgements

This project was funded by LUKE, NINA and the Academy of Finland. We thank
Teno/Tana/Deatnu fishers for providing scale samples and Reeta Partanen for laboratory assistance.

Contributions
Project conceptualization: CRP, VLP, HJ, JE, MF, PO
Project management/ Sample coordination: - PO, MF, AF, JE, CRP, VLP
Methods development: VLP
Laboratory analysis: VLP,
Bioinformatics: VLP
Data analysis: HJ, VLP
Manuscript drafting: HJ, CRP, VLP
Visualization: HJ, VLP
Manuscript finalization: all authors.

Figure 1 (high res version to be added) Map of the Teno river system baseline sampling localities
and Neighbour-joining tree using Prevosti's distance on 17,546 SNPs. Geographic regions indicated
by colour in map and in tree.

468   Figure 2: Results of Admixture analysis.

473   Figure 3. The proportion of female biomass per river/tributary plotted against the estimated catch
474   proportion in the mixed fishery for the same reporting units.
475

476
477

478   **Table 1**. Percent of correct assignment of the 180 SNPs into the 22 reporting units, mis-assignments locations and rates.

Reporting unit	SNPs			Microsatellites	
	Baseline sample size	% correct assignment	Largest misassignment to:	Baseline sample size	% correct assignment
Teno Mainstem					
TanaBru/Luovtejohka	69	61%	Teno Mainstem (37%)	127	66%
Middle Mainstem*	220	83%	Tana Bru (12%)	446	83%
Outakoski	48	52%	Teno MS (18%); Inari HW (15%)	91	54%
Lower Tributaries					
Máskejohka	37	86%	Teno Mainstem (8%)	121	97%
Vetsijoki	36	92%	Iešjohka (5%)	212	96%
Ylä-Pulmankijoki	17	100%	-	279	99%
Galddasjoki	19	100%	-	85	99%
Utsjoki	38	97%	Vetsijoki (3%)	129	99%
Tsarsjoki	65	97%	Kevojoki (3%)	196	97%
Kevojoki	33	100%	-	165	99%
Lakš/Deavkke/Gurtsjohka	98	97%	Karigas; Máske; Vetsi (1% each)	71	100%
Upper Tributaries					
Levajohka	31	74%	Outakoski (6%); Báiš/Nilijoki (6%)	51	69%
Báišjohka/Nilijoki	73	85%	Akujoki (6%)	106	77%
Akujoki	33	70%	Báišjohka/Nilijoki (12%)	53	40%
Karigasjoki	40	83%	Akujoki (10%)	75	79%
Kuoppilasjoki/Birkejohka	33	91%	Báišjohka/Nilijoki (10%)	112	91%
Váljohka	30	93%	Inari HW (7%)	270	94%
Iešjohka Headwaters					
Iešjohka	54	93%	Kárášjohka (5%)	201	89%
Kárášjohka	40	83%	Iešjohka (3%); Vetsijoki (3%)	270	94%
Geáimmejohka	18	94%	Akujoki (5%)	52	92%
Inari Headwaters					
Inari Headwaters**	132	91%	Karigasjoki (4%)	177	85%
Kietsimäjoki	26	73%	Inari HW (15%)	61	74%

479
480   *Teno Mainstem reporting group: Alaköngäs, Piltamo, Sirma, Garnjaga, Kortsham and Yläköngäs
481   **Inari Headwaters reporting group: Inari Mainstem, Cášcemjohka,Vuomajoki, Goššjohka, Iškurasjohka
482
483

Table 2. Mixed Stock Analysis of Tana 2018 Mainstem Catch.

Reporting Unit	Estimated mixture proportion	Lower 95% CI	Upper 95% CI
Tana Mainstem			
TanaBru/Luovtejohka	0.029	0.0214	0.0378
Teno Mainstem*	0.251	0.2335	0.2689
Outakoski	0.059	0.0480	0.0700
Lower Tributaries			
Máskejohka	0.023	0.0171	0.0297
Vetsijoki	0.074	0.0641	0.0851
Váljohka	0.021	0.0155	0.0265
Ylä-Pulmankijoki	0.007	0.0044	0.0109
Galddasjoki	0.001	0.0003	0.0030
Utsjoki	0.019	0.0144	0.0250
Tsarsjoki	0.019	0.0140	0.0243
Kevojoki	0.032	0.0253	0.0386
Lakš/Deavkkehan/Gurtsjohka	0.007	0.0044	0.0107
Geáimmejohka	0.013	0.0086	0.0171
Upper Tributaries			
Levajohka	0.018	0.0133	0.0237
Báišjohka/Nilijoki	0.023	0.0171	0.0285
Akujoki	0.009	0.0048	0.0129
Karigasjoki	0.017	0.0120	0.0232
Kuoppilasjoki/Birkejohka	0.006	0.0031	0.0090
Iešjohka Headwaters			
Iešjohka Upper/Lower	0.136	0.1227	0.1507
Kárášjohka	0.075	0.0639	0.0854
Inari Headwaters			
Inari Headwaters**	0.159	0.1445	0.1733
Kietsimäjoki	0.003	0.0009	0.0050

*Teno Mainstem reporting group: Alaköngäs, Piltamo, Sirma, Garnjaga, Kortsham and Yläköngäs

**Inari Headwaters reporting group: Inari Mainstem, Cášcemjohka,Vuomajoki, Goššjohka, Iškurasjohka

References

Ackerman, M. W., Habicht, C., & Seeb, L. W. (2011). Single-Nucleotide Polymorphisms (SNPs) under Diversifying Selection Provide Increased Accuracy and Precision in Mixed-Stock Analyses of Sockeye Salmon from the Copper River, Alaska. *Transactions of the American Fisheries Society*, *140*, 865–881.

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, *19*, 1655–1664.

Anderson, E. C. (2010). Assessing the power of informative subsets of loci for population assignment: standard methods are upwardly biased. *Molecular Ecology Resources*, *10*, 701–710.

Anderson, E. C., Waples, R. S., & Kalinowski, S. T. (2008). An improved method for predicting the accuracy of genetic stock identification. *Canadian Journal of Fisheries and Aquatic Sciences*, *65*, 1475–1486.

Anon. (2020). *Status of the Tana / Teno River salmon populations in 2020*. Vol. 1/2020.

Archer, F. I., Adams, P. E., & Schneiders, B. B. (2017). STRATAG: An R package for manipulating, summarizing and analysing population genetic data. *Molecular Ecology Resources*, *17*, 5–11.

Aykanat, T., Lindqvist, M., Pritchard, V. L., Primmer, C. R., Lindqvist, M., & Primmer, C. R. (2016). From population genomics to conservation and management: a workflow for targeted analysis of markers identified using genome-wide approaches in Atlantic salmon Salmo salar. *Journal of Fish Biology*, *89*, 2658–2679.

Barson, N. J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G. H., Fiske, P., … Primmer, C. R. (2015). Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature*, *528*, 405–408.

Begg, G. A., Friedland, K. D., & Pearce, J. B. (1999). Stock identification and its role in stock assessment and fisheries management: an overview. *Fisheries Research*, *43*, 1–8.

Campbell, N. R., Harmon, S. A., & Narum, S. R. (2015). Genotyping-in-Thousands by sequencing (GT-seq): A cost effective SNP genotyping method based on custom amplicon sequencing. *Molecular Ecology Resources*, *15*, 855–867.

Chang, C. C., Chow, C. C., Tellier, L. C. A. M., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-generation PLINK: Rising to the challenge of larger and richer datasets. *GigaScience*, *4*, 1–16.

Crozier, W. W., Schön, P.-J., Chaput, G., Potter, E. C. E., Maoileidigh, N. O., & MacLean, J. C. (2004). Managing Atlantic salmon (Salmo salar L.) in the mixed stock environment: challenges and considerations. *ICES Journal of Marine Science*, *61*, 1344–1358.

Erkinaro, J., Czorlich, Y., Orell, P., Kuusela, J., Falkegård, M., Länsman, M., … Niemelä, E. (2019). Life history variation across four decades in a diverse population complex of Atlantic salmon in a large subarctic river. *Canadian Journal of Fisheries and Aquatic Sciences*, *76*, 42–55.

Falkegård, M., Foldvik, A., Fiske, P., Erkinaro, J., Orell, P., Niemelä, E., … Hindar, K. (2014). *Revised first-generation spawning targets for the Tana/Teno river system*. 1–68 pp. Vol. 1087.

Fischer, M. C., Rellstab, C., Leuzinger, M., Roumet, M., Gugerli, F., Shimizu, K. K., … Widmer, A. (2017). Estimating genomic diversity and population differentiation - an empirical comparison of microsatellite and SNP variation in Arabidopsis halleri. *BMC Genomics*, *18*, 1–

15.

Frichot, E., & Francois, O. (2015). LEA: An R package for landscape and ecological association
studies. *Methods in Ecology and Evolution*, *6*, 925–929.

Harvey, A. C., Quintela, M., Glover, K. A., Karlsen, Ø., Nilsen, R., Skaala, Ø., … Wennevik, V.
(2019). Inferring Atlantic salmon post-smolt migration patterns using genetic assignment.
Royal Society Open Science, *6*, 190426.

Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: an R package for genetic analysis
of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*, *2*, e281.

Koljonen, M.-L., & McKinnell, S. (1996). *Assessing seasonal changes in stock composition of*
Atlantic salmon catches in the Baltic Sea with genetic stock identification. 998–1018 pp. Vol.
49.

Koljonen, M., Pella, J. J., & Masuda, M. (2005). Classical individual assignments versus mixture
modeling to estimate stock proportions in Atlantic salmon (Salmo salar) catches from DNA
microsatellite data. *2158*, 2143–2158.

Larson, W. A., Seeb, J. E., Pascal, C. E., Templin, W. D., & Seeb, L. W. (2014). Single-nucleotide
polymorphisms (snps) identified through genotyping-by-sequencing improve genetic stock
identification of chinook salmon (oncorhynchus tshawytscha) from western alaska. *Canadian*
Journal of Fisheries and Aquatic Sciences, *71*, 698–708.

Li, H., & Barrett, J. (2011). A statistical framework for SNP calling, mutation discovery,
association mapping and population genetical parameter estimation from sequencing data. *27*,
2987–2993.

Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows–Wheeler
transform. *Bioinformatics*, *26*, 589–595.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet Journal, *17*, 10–12.

May, S. A., McKinney, G. J., Hilborn, R., Hauser, L., & Naish, K. A. (2020). Power of a dual-use
SNP panel for pedigree reconstruction and population assignment. *Ecology and Evolution*, *10*,
9522–9531.

Mobley, K. B., Granroth-Wilding, H., Ellmen, M., Vähä, J.-P., Aykanat, T., Johnston, S. E., …
Primmer, C. R. (2019). Home ground advantage: Local Atlantic salmon have higher
reproductive fitness than dispersers in the wild. *Science Advances*, *5*, eaav1112.

Moran, B. M., & Anderson, E. C. (2019). Bayesian inference from the conditional genetic stock
identification model. *Canadian Journal of Fisheries and Aquatic Sciences*, *76*, 551–560.

Ozerov, M., Vähä, J.-P., Wennevik, V., Niemelä, E., Svenning, M.-A., Prusov, S., … Christiansen,
B. (2017). Comprehensive microsatellite baseline for genetic stock identification of Atlantic
salmon (Salmo salar L.) in northernmost Europe. *ICES Journal of Marine Science*, *74*, 2159–
2169.

Palsbøll, P. J., Bérubé, M., & Allendorf, F. W. (2007). Identification of management units using
population genetic data. *Trends in Ecology and Evolution*, *22*, 11–16.

Paradis, E., & Schliep, K. (2019). Phylogenetics ape 5.0: an environment for modern phylogenetics
and evolutionary analyses in R. *Bioinformatics*, *35*, 526–528.

Parrish, D. L., Behnke, R. J., Gephard, S. R., McCormick, S. D., & Reeves, G. H. (1998). Why
aren't there more Atlantic salmon (*Salmo salar*)? *Canadian Journal of Fisheries and Aquatic*
Sciences, *55*, 281–287.

Prevosti, A., Ocana, J., & Alonso, G. (1975). Distances between Populations of Drosophila
subobscura, Based on Chromosome Arrangement Frequencies. *Theoretical and Applied*
Geentics, *45*, 231–241.
Pritchard, V. L., Mäkinen, H., Vähä, J. P., Erkinaro, J., Orell, P., & Primmer, C. R. (2018).
Genomic signatures of fine-scale local selection in Atlantic salmon suggest involvement of
sexual maturation, energy homeostasis and immune defence-related genes. *Molecular Ecology*,
27, 2560–2575.
R Core Team. (2014). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. *URL http://www.R-project.org/.*
Storer, C. G., Pascal, C. E., Roberts, S. B., Templin, W. D., Seeb, L. W., & Seeb, J. E. (2012). Rank
and Order: Evaluating the Performance of SNPs for Individual Assignment in a Non-Model
Organism. *PLoS ONE*, *7*, 49018.
Sunde J, Yıldırım Y, Tibblin P, Forsman A. Comparing the Performance of Microsatellites and
RADseq in Population Genetic Studies: Analysis of Data for Pike (*Esox lucius*) and a
Synthesis of Previous Studies. Front Genet. 2020 Mar 13;11:218. doi:
10.3389/fgene.2020.00218. PMID: 32231687; PMCID: PMC7082332.
Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B. C., Remm, M., & Rozen, S. G.
(2012). Primer3-new capabilities and interfaces. *Nucleic Acids Research*, *40*, e115.
Vähä, J.-P., Erkinaro, J., Niemelä, E., Primmer, C. R., Saloniemi, I., Johansen, M., … Brørs, S.
(2011). Temporally stable population-specific differences in run timing of one-sea-winter
Atlantic salmon returning to a large river system. *Evolutionary Applications*, *4*.
Vähä, J.-P., Erkinaro, J., Falkegård, M., Orell, P., & Niemelä, E. (2017). Genetic stock
identification of Atlantic salmon and its evaluation in a large population complex. *Canadian*
Journal of Fisheries and Aquatic Science, *74*, 327–338.
Vähä, J.-P., Erkinaro, J., Niemelä, E., & Primmer, C. R. (2007). Life-history and habitat features
influence the within-river genetic structure of Atlantic salmon. *Molecular Ecology*, *16*, 2638–
2654.
Vähä, J.-P., Erkinaro, J., Niemelä, E., & Primmer, C. R. (2008). Temporally stable genetic structure
and low migration in an Atlantic salmon population complex: implications for conservation
and management. *Evolutionary Applications*, *1*, 137–154.
Winans, G. A., Paquin, M. M., Van Doornik, D. M., Baker, B. M., Thornton, P., Rawding, D., …
Kalinowski, S. (2004). *Genetic Stock Identification of Steelhead in the Columbia River Basin:*
An Evaluation of Different Molecular Markers; Genetic Stock Identification of Steelhead in the
Columbia River Basin: An Evaluation of Different Molecular Markers. 672–685 pp. Vol. 24.
Yano, A., Nicol, B., Jouanno, E., Quillet, E., Fostier, A., Guyomard, R., & Guiguen, Y. (2013). The
sexually dimorphic on the Y-chromosome gene (sdY) is a conserved male-specific Y-
chromosome sequence in many salmonids. *Evolutionary Applications*, *6*, 486–496.
617