TITLE

Melanocyte differentiation and mechanosensation are differentially modulated by distinct

extracellular matrix proteins

AUTHORS

Carole Luthold[1,2,3*], Marie Didion[1,2,3], Vanessa Samira Rácz[1,2,3], Emilio Benedum[1,2,3], Ann-

Kathrin Burkhart[1,2,3], Nina Demmerle[1,2,3], Evelyn Wirth[1,2,3], Gubesh Gunaratnam[4], Sudharshini

Thangamurugan[5], Volkhard Helms[5], Markus Bischoff[4], Annika Ridzal[1,3], and Sandra Iden[1,2,3*]

AFFILIATIONS

[1] Cell & Developmental Biology, Saarland University, Faculty of Medicine, Homburg/Saar,

Germany

[2] Center for Gender-specific Biology & Medicine (CGBM), Saarland University, Faculty of

Medicine, Homburg/Saar, Germany

[3] Center for Human & Molecular Biology (ZHMB), Saarland University, Germany

[4] Institute for Medical Microbiology and Hygiene, Homburg/Saar, Germany

[5] Center for Bioinformatics, Saarland University, Saarbrücken, Germany

CONTACT DETAILS

CL, SI: Cell & Developmental Biology, Saarland University, Faculty of Medicine, Kirrberger

Strasse, Building 61.4, 66421 Homburg/Saar, Germany

CORRESPONDENCE

carole.luthold@uni-saarland.de, sandra.iden@uks.eu

ORCIDs

CL: 0009-0009-9297-7516

MD: 0009-0008-4114-8841

VSR: 0009-0007-4884-1051

EB: 0009-0006-0751-6087

AKB: 0000-0003-4128-6677

ND: 0009-0007-4436-3048

GG: 0000-0002-6533-8681

VH: 0000-0002-2180-9154

MB: 0000-0001-6734-2732

AR: 0000-0002-9038-0734

SI: 0000-0003-2333-9827

RUNNING TITLE ECM molecules and substrate stiffness steer melanocyte functions

WORD COUNT 6253 (abstract, introduction, results, discussion, conclusion)

DISPLAY ITEMS 7 Figures, 6 Supplementary Figures, 2 Supplementary Tables

ABSTRACT

Dysfunctions in melanocytes can lead to pigmentation disorders, such as albinism, or contribute to the development of melanoma, the most aggressive form of skin cancer. Epidermal melanocytes typically interact with the collagen IV-rich basement membrane, but upon injury or in pathological conditions, they can encounter environments rich in collagen I or fibronectin. While alterations in extracellular matrix (ECM) composition and stiffness are known to impact cell behavior, the specific roles of each of these cues for melanocyte functions remain unclear. To explore the impact of these extrinsic cues, we here exposed murine melanocytes to different ECM proteins as well as varying substrate stiffnesses. This study identified MITF, a key regulator of melanocyte differentiation and function, as an ECM- and mechanosensitive transcription factor. We further revealed that distinct ECM proteins and substrate stiffness engage a FAK/MEK/ERK/MITF signaling axis to control melanocyte functions. Exposure of melanocytes to collagen I restricted FAK and ERK activation, promoting high nuclear MITF levels associated with melanocyte proliferation and differentiation. Conversely, fibronectin elicited elevated FAK and ERK activation, leading to low nuclear MITF, correlating with a dedifferentiated and motile phenotype. Consistent with these observations, RNA sequencing revealed that collagen I supports a differentiated gene expression program, whereas fibronectin induces a dedifferentiated transcriptomic signature. Importantly, inhibiting MEK or ERK activity in melanocytes cultured on fibronectin led to increased MITF nuclear localization and enhanced melanogenesis. Additionally, on fibronectin FAK inhibition reduced ERK activation and enhanced melanogenesis, supporting a role for FAK upstream of ERK. Finally, we uncovered that melanocyte mechanoresponses differ depending on the specific ECM environment. Together, these findings reveal a substantial effect of extrinsic cues on melanocyte function, with a context-dependent MITF regulation downstream of ERK, offering new perspectives for our understanding of melanocyte-related pathologies.

KEYWORDS

Melanocytes; MITF; Differentiation; Extracellular matrix; Type IV Collagen; Type I Collagen; Fibronectin; Stiffness; ERK signaling; Melanogenesis; Mechanosensation.

INTRODUCTION

Melanocytes (MCs) are pigment-producing cells found in the basal layer of the skin epidermis, and in various organs such as the brain, heart, and eyes (Centeno et al., 2023). The pronounced dendritic morphology of cutaneous MCs enables an efficient melanosome transfer to surrounding keratinocytes, providing photoprotection (Benito-Martínez et al., 2021; Cui & Man, 2023; Domingues et al., 2020; Hirobe, 2014; Prospéri et al., 2024). Dysfunctions of MCs can cause pigmentation disorders such as albinism and vitiligo (Coutant et al., 2024), and can lead to malignant transformation into melanoma, an aggressive form of skin cancer responsible for most skin cancer-related deaths (Matthews et al., 2017).

The specification and differentiation of MCs from the neural lineage are governed by the melanocyte-inducing transcription factor (MITF). MITF regulates MC differentiation by controlling the expression of pigmentation genes (e.g., *Tyr*, *Trp1*, *Trp2*) and maintains cellular homeostasis by modulating genes involved, among others, in cell-cycle progression (e.g., *Cdk2*) and apoptosis (e.g., *Bcl2*) (Goding & Arnheiter, 2019; Kawakami & Fisher, 2017). MITF is also a major player of melanoma progression, influencing both, the melanoma cell differentiation state and plasticity, contributing to high tumor heterogeneity. Within a single tumor, different cell states co-exist: highly differentiated, proliferative melanocytic cells, which are associated with high MITF levels, whereas slow-cycling, dedifferentiated, invasive stem-like cells correlate with low MITF levels (Arozarena & Wellbrock, 2019; Carreira et al., 2006; Cheli et al., 2011; Hoek et al., 2008; Müller et al., 2014; Popovic & Tartare-Deckert, 2022; Rambow et al., 2019; Tirosh et al., 2016; Tsoi et al., 2018; Wouters et al., 2020). Interestingly, differentiated MCs can give rise to melanoma through a process that includes their reprograming and dedifferentiation (Köhler et al., 2017). However, the specific factors and mechanisms driving this dedifferentiation of mature MCs, and the role of MITF in these processes, remain poorly understood. Therefore, understanding how MITF affects MC plasticity and identifying novel factors that regulate MITF expression and activity could provide new insights into the mechanisms underlying MC transformation.

In addition to the heterotypic cell-cell interaction with keratinocytes, MCs are in direct contact with the basement membrane, a specialized extracellular matrix (ECM) rich in type IV collagen (COL IV), which forms the junction with the underlying dermis, comprising type I collagen (COL I) and fibronectin (FN) (Pfisterer et al., 2021). This implies that under physiological conditions, COL IV is the predominant ECM type epidermal MCs are exposed to, whereas interactions with dermal COL I and FN can occur due to altered basement membrane integrity, for instance, following injury or chronic UV exposure (Amano, 2009, 2016; Fisher & Rittié, 2018; Iriyama et al., 2011, 2020; Yoshihisa et al., 2014). Within the skin, a wide range of stiffness has been reported for different compartments, ranging from lower kPa (dermis-like)

to lower MPa (epidermis-like) values (Biggs et al., 2020; Feng et al., 2022; Graham et al., 2019). Importantly, the cellular microenvironment, including ECM components and tissue stiffness, significantly impacts cell fate and function (Bonnans et al., 2014; Guilak et al., 2009; Walma & Yamada, 2020). Cells adapt to these molecular and mechanical parameters by transmitting environmental signals to intracellular signal transduction pathways, including mitogen-activated protein kinase (MAPK)/ extracellular signal-regulated kinase (ERK) pathways (Tan et al., 2023). Crucial for this process are focal adhesions (FAs), dynamic multi-protein complexes at the plasma membrane that link the ECM to the actin cytoskeleton via transmembrane receptors such as integrins. Notably, focal adhesion kinase (FAK), a key signaling protein within FAs, is activated through integrin engagement with the ECM, resulting in autophosphorylation and activation of downstream signaling pathways like MAPK/ERK (Paszek et al., 2005). In various cell systems, such signal activation downstream of ECM cues has been reported to affect the localization and activation levels of various transcription factors and coregulatory factors, such as Yes-associated protein (YAP), leading to gene expression changes that impact processes like proliferation, survival, or differentiation (Ishihara & Haga, 2022).

In melanoma cells, MAPK signaling components have recently been reported to negatively regulate MITF nuclear localization and activity in melanoma cells: rapidly accelerated fibrosarcoma (RAF) proteins, acting upstream of ERK, interact directly with MITF, causing its cytoplasmic retention and reduced transcriptional activity (Estrada et al., 2022), while MAPK/ERK signaling, in collaboration with glycogen synthase kinase 3 (GSK3), controls MITF nuclear export (Ngeow et al., 2018). The transcriptional co-activator YAP, known to translocate into the nucleus in response to ECM stiffening in various cell types (Cai et al., 2021; Dupont et al., 2011; Y. Huang et al., 2022; Miskolczi et al., 2018), promotes MITF expression in uveal and cutaneous melanoma cells (Barbosa et al., 2023; Miskolczi et al., 2018) and serves as MITF cofactor in uveal melanoma (Barbosa et al., 2023). This indicates a dual role of YAP in the control of MITF in melanoma. However, whether and how ECM cues modulate MITF remains to be elucidated.

So far, studies have separately explored the roles of ECM type and substrate stiffness for MC biology. For example, Hara et al. reported that COL IV stimulates dendricity in human MCs on glass substrates (considered as ultra-hard stiffness) (Hara et al., 1994), while melanin production on stiff PDMS substrates (5.5 MPa) coated with laminin is higher compared to softer counterparts (from 50 kPa to 1.8 MPa) (Choi et al., 2014). However, the potential synergistic effects of substrate stiffness and ECM components on MC functions remain to be defined. Understanding such effects is particularly important since the ECM undergoes continuous remodeling that includes de novo synthesis and degradation (Pfisterer et al., 2021). Various

factors, such as skin aging, inflammation, sunlight exposure, wound healing, and fibrosis, can promote ECM remodeling, potentially resulting in aberrant ECM modifications (Pfisterer et al., 2021), which may modify the presentation of ECM ligands and alter substrate stiffness. For instance, during wound healing, the ECM transitions from a FN-rich provisional matrix to a COL I–rich structure as the tissue repairs. Nevertheless, it is still insufficiently understood how substrate stiffness influences the cellular responses to specific ECM components, and, conversely, whether cells need to engage in particular ECM interactions in order to sense environmental stiffness features.

This study investigated whether and how substrate stiffness and ECM components act together to modulate MC functions. We compared the combined effects of various ECM proteins and substrate stiffness on MC behavior. Given that MITF plays a central role in integrating environmental signals to regulate key aspects of MC identity, function, and plasticity—and considering its known regulation by the MAPK/ERK pathway, which mediates ECM signal transduction—we hypothesized that MITF may serve as a key effector of ECM-dependent cues in MCs. By investigating how ECM composition and mechanical properties affect MITF activity, we aimed to uncover mechanisms through which the microenvironment shapes MC behavior and plasticity. Our results revealed that ECM components control MC differentiation and function via a FAK-MEK-ERK-MITF axis, with different ECM types determining the ability of MCs to perceive and respond to mechanical stimuli in their environment.

RESULTS

ECM components differentially affect MC morphology, adhesion and migration

To assess how extracellular mechanical cues affect MC behavior, we employed PDMS substrates of varying stiffnesses: ≈45 kPa defined here as soft (fig. S1A), ≈140 kPa as intermediate (fig. S1B) and ≈900 kPa as stiff (fig. S1C). This range was chosen to reflect the wide range of stiffness found within the skin and because increased environmental stiffness is associated with multiple pathologies (Diazzi et al., 2020; Pfisterer et al., 2021; K. Wang et al., 2023). To mimic both normal and altered ECM conditions, COL IV-, COL I-, and FN-coated substrates of varying stiffness were used for cultures of either primary (pMCs) or immortalized murine MCs (stably expressing membrane-targeted tandem dimer Tomato; iMCs, generated for this study), followed by molecular, cellular and functional assays.

Interestingly, visual inspection in phase-contrast and fluorescence microscopy revealed distinct ECM-specific phenotypes, with COL I and FN eliciting opposing cellular responses. iMCs cultured on FN-coated substrates exhibited a more dendritic shape compared to the mostly bi- or tripolar MCs grown on COL I (fig. 1A). To examine this phenotype further, we

performed a Sholl analysis (fig. 1A-C; fig. S2A; Binley et al., 2014; Sholl, 1953). Quantification

of the dendritic complexity of MCs (fig. 1B) and the corresponding Sholl profile (fig. 1C; fig.

S2A) confirmed that most iMCs exposed to COL IV and FN displayed between 4 to 7 dendrites,

whereas MCs grown on COL I exhibited mostly 2 to 3 dendrites (fig. 1B). Noteworthy, over the

range of stiffnesses tested, MC dendricity was not impacted by substrate stiffness, irrespective

of the ECM protein type used (fig. S2A).

Since cell spreading is regulated by coordinated changes in adhesions to ECM and

cytoskeletal reorganization, we next examined the cell area as well as key cell adhesion

parameters. Among these, FAK activity by means of its autophosphorylation (p-FAK) and

quantification of the number of focal adhesions (FAs) per cell served as indicators of responses

resulting from MC-ECM interactions. In both pMCs and iMCs, the smallest cell area was

detected on COL I (fig. 1D,E; fig. S3A) and correlated with the lowest p-FAK levels (fig. 1D,F;

fig. S3B), whereas the largest cell areas (fig. 1D,E; fig. S3A) were found on FN, correlating

with high p-FAK signals (fig. 1D,F; fig. S3B). Substrate stiffness, however, seemed to have

less influence on the cell area: apart from a slight increase from soft to intermediate stiffness

in iMCs cultured on COL I, we did not note significant changes of the cell area across the

different substrate stiffnesses, regardless of the ECM protein used (fig. 1E; fig. S3A). In

contrast, both p-FAK and the number of FAs per μm^2 exhibited a significant stiffness-mediated

increase in MCs grown on FN-coated substrates (fig. 1F,G; fig. S3B,C). In contrast, the overall

low p-FAK levels in COL I-exposed iMCs even decreased with stiffness (fig. 1F), highlighting

that stiffness sensing in MCs depends on the ECM type.

Finally, since cell morphology and FAs are closely linked to cell movement, we investigated

whether the distinct ECM molecules influence MC motility. Live cell imaging of iMCs revealed

that both the distance traveled (fig. 2A-C) and velocity (fig. 2D) of cell migration decreased on

COL I compared to COL IV and FN, whereas FN-grown iMCs showed increased motility (fig.

2A-C) and faster migration (fig. 2D) relative to their COL IV-grown counterparts.

Together, these data show that COL I and FN trigger opposite morphological, adhesive and

migratory phenotypes in MCs.

COL I and FN have opposite effects on melanin production and MC proliferation

Given the observed role of ECM components for MC morphology and migration, we next

investigated whether ECM cues influence other MC functions. As a hallmark task of MCs, we

examined their melanin production by measuring both intracellular melanin contents and

melanin released into the culture medium. iMC grown on COL I-coated substrates exhibited

the highest intra- and extracellular melanin content, while melanin levels on FN were lowest

compared to COL I and COL IV (fig. 3A,B). Notably, the expression of *Tyr*, an essential gene

for melanogenesis, correlated with melanin production, exhibiting a higher expression in iMCs

exposed to COL I than in those exposed to FN (fig. 3C). Like iMCs, pMCs cultured on FN
produced least melanin on soft and intermediate stiffness (fig. S3D). For all ECM types tested,
highest values for melanin production were observed at high stiffness (fig. S3D, fig. 3A,B).
Collectively, these data demonstrated that compared to COL I, FN restricts melanin
production.

Using a BrdU assay, we next evaluated MC proliferation. While stiffness had no effect on MC
proliferation in the presence of COL IV and FN, we observed a higher proportion of BrdU-
positive iMCs when exposed to stiff COL I-coated substrates, indicating a synergistic effect of
COL I and stiffness on MC proliferation (fig. 3D). Among the three ECM types tested, FN
exposure resulted in the lowest proliferation rates (fig. 3D), a finding that could be further
confirmed by Ki67 immunostaining (fig. 3E), as well as live cell imaging followed by
quantification of the percentage of dividing cells (fig. 3F). Overall, these experiments revealed
a stiffness-dependent proliferative phenotype triggered by COL I, while FN appears to restrict
cell division of MCs.

Together, our findings show that melanin production and *Tyr* expression are highest in MCs
exposed to COL I and lowest in FN-grown MCs. Moreover, cell proliferation was lowest on FN,
and stiffness-mediated increase in proliferation was only observed on COL I-coated
substrates, suggesting that MC mechanoresponses depend on the ECM type.

MITF nuclear localization is modulated by ECM molecules and substrate stiffness but
does not correlate with nuclear YAP
Our data indicate an ECM-dependent regulation of MC proliferation and melanin production,
with COL I and FN driving opposite effects on these cellular functions. Considering the central
role of the transcription factor MITF in the control of both melanogenesis and cell cycle
progression (Goding & Arnheiter, 2019), and taking into account the observed increase of the
MITF target gene *Tyr* on COL I (fig. 3C), we next examined the impact of ECM cues on the
protein levels and subcellular localization of MITF (fig. 4A-C; fig. S3E). Although overall MITF
levels were lowest on COL I, nuclear MITF levels were highest on this ECM type (fig. 4A-C),
consistent with pronounced melanin production. In contrast, MCs cultured on FN displayed low
nuclear MITF levels compared to COL I-coated substrates (fig. 4A-C; fig. S3E).

Furthermore, a stiffness-mediated increase of MITF nuclear localization was noted in MCs on
COL I (fig. 4B,C; fig. S3E), thus identifying MITF as an ECM- and mechanosensitive
transcription factor. The contrasting phenotypes of MCs grown on COL I vs. FN were also
evident when correlating nuclear MITF levels to the cell area (fig. S4A): while MCs grown on
COL I were characterized by a low ratio of cell area to nuclear MITF compared to COL IV, FN-
exposed MCs showed high ratios, i.e. a large cell area was associated with low MITF.
Interestingly, this ratio decreased with substrate stiffness on COL I and FN but not on COL IV

(fig. S4A), suggesting that changes in nuclear MITF are not linearly related to morphological

changes in response to ECM. siRNA-mediated knock-down of MITF (fig. S4B) confirmed the

principal role of MITF for melanin production on both COL I and FN (fig. 4D).

Given that YAP is a transcriptional co-activator known to translocate into the nucleus in

response to ECM stiffening in various cell types, including melanoma cells (Miskolczi et al.,

2018), and that YAP may serve as potential MITF partner in uveal melanoma cells (Barbosa

et al., 2023), we wondered whether YAP could be associated with the observed ECM-

dependent MC phenotypes. We examined YAP subcellular localization and, consistent with

previous reports, observed a stiffness-mediated increase in nuclear YAP, with the strongest

response at intermediate stiffness levels on COL IV and COL I, and on stiff FN-coated

substrates (fig. 4E,F). Importantly, however, unlike nuclear MITF, we hardly observed

significant differences of nuclear YAP when comparing the three ECM types, making it unlikely

that MCs utilize YAP for ECM-specific melanin production.

Collectively, our findings suggest that ECM subtypes regulate MC behavior and function likely

through modulation of MITF localization and activity, independently of YAP.

Bulk RNA sequencing confirms that ECM types elicit significant changes in expression

of genes associated with MC pigmentation and differentiation

Our data so far indicate that ECM composition modulates MITF localization and activity in MCs,

suggesting that ECM components may influence MITF-regulated transcriptional programs. To

investigate this further and gain a broader understanding of how ECM components shape the

MC phenotype, we performed bulk RNA sequencing (RNAseq) on MCs cultured on COL IV,

COL I, and FN (fig. 5; fig. S5). To capture global transcriptomic changes associated with

distinct ECM environments and to identify gene signatures linked to MC differentiation,

plasticity, and other relevant pathways, we analyzed the expression of MITF target genes, MC-

specific genes, and broader signaling networks that are associated with the phenotypic shifts

observed across ECM conditions.

Comparison of global transcriptomic profiles across the ECM types used revealed a range of

differentially expressed genes (DEGs) (fig. 5; fig. S5B,C). Consistent with the reduced melanin

production observed for MCs on FN (fig. 3B), several MITF-dependent target genes, which are

also pigmentation-related (e.g., *Dct*, *Pmel*, *Tyr*, *Gpr143*), were significantly downregulated on

FN compared to COL I (fig. 5A,B). In addition, further components involved in melanin

biosynthesis and trafficking—such as *Oca2*, *Adcy2*, and *Rab27a*—also showed reduced

expression on FN (fig. 5A,B). Of note, some of the MITF-dependent DEGs have previously

been linked to apoptosis *(Trpm1)* and cell cycle progression (*Ccnd1*, *Cdkn1a*) (fig. 5A). Global

gene expression analysis indeed revealed further DEGs related to cell cycle control and

apoptosis (fig. 5C,D), in line with the observed decrease in proliferation of MCs cultured on FN

when compared to COL I (fig. 3D,E). In addition, FN elicited differential expression of genes associated with cellular senescence (fig. 5E). Taken together, these findings suggest that adhesion to FN promotes a transcriptional program in MCs that may counteract cell cycle progression.

Considering that MITF-dependent gene expression overlapped with transcriptional programs involved in melanocyte development, and that these programs were downregulated in MCs cultured on FN (fig. 5A), we extended our analysis to gene expression signatures associated with melanoma plasticity and differentiation states (Arozarena & Wellbrock, 2019; Durand et al., 2024; F. Huang et al., 2021; Konieczkowski et al., 2014; Pessoa et al., 2021; Rambow et al., 2015, 2019; Tsoi et al., 2018). Comparing FN to COL I revealed downregulation of genes associated with melanocytic differentiation on FN, such as *Apoe,* next to the previously mentioned MITF target genes (e.g., *Dct*, *Pmel*, *Tyr*, *Trpm1*, *Gpr143*, *Irf4*) and melanin biosynthesis-related genes (e.g., *Oca2*, *Adcy2)* (fig. 5A,B,F). However, core melanocytic factors such as *Sox10*, *Lef1*, *Creb1*, *Pax3*, and *Mlana* (fig. 5F) were not significantly altered between COL I and FN, suggesting maintenance of lineage identity. In contrast, several genes associated with cell plasticity *(Tead1, Fzd4, Tagln2)* and neural crest-like features (*Ngfr, Erbb3*) were significantly upregulated on FN relative to COL I (fig. 5G). Additionally, genes previously associated with a dedifferentiated state, such as *Axl, Itga8*, *Timp1*, *Hmga1*, and *Nkd2*, were upregulated on FN compared to COL I (fig. 5H), further supporting the emergence of a phenotype characterized by increased plasticity and reduced differentiation on FN.

Gene set enrichment analysis (GSEA) identified "ECM organization" as the top enriched Reactome pathway in iMCs cultured on FN compared to COL I (fig. S5E), indicating transcriptional remodeling of ECM-associated genes in response to ECM molecules. Among these, collagen genes (*Col1a2* and *Col9a3;* fig. 5I) were downregulated on FN, while non-collagenous ECM components, including *Fn1*, *Tnc*, *Hspg2* and *Thbs1,* showed increased expression (fig. 5I). In addition, ECM-modifying genes (*Sparc*, *Adam12*, *Serpine1;* Fig. 5I) were also upregulated, supporting a potential active ECM remodeling on FN. Together, these gene expression patterns align with the emergence of a cell state characterized by increased migratory potential and plasticity in MCs grown on FN.

Interestingly, several integrin subunits were also differentially expressed across ECM conditions (fig. 5I). A broad repertoire of integrins was found to be expressed in iMCs grown on COL IV, COL I, and FN, including subunits commonly reported in MCs (Arias-Mejias et al., 2020; Hara et al., 1994; Morelli et al., 1993; Pinon & Wehrle-Haller, 2011; Scott et al., 1992), such as *Itga2, Itga3, Itga5, ItgaV, Itgb1,* and *Itgb3*, which mediate adhesion to collagens and fibronectin (fig. 5J, K). While most of these were expressed independently of the ECM type used, we also noted that *Itga6*, *Itga7*, *Itga8*, and *Itgb8* were upregulated on FN, while *Itga9*

315 was upregulated on COL I (fig. 5I). Taken together, our data suggests that while ECM

316 composition influences integrin gene expression to some extent, changes in integrin profiles

317 alone may not fully account for the phenotypic differences observed between the distinct ECM

318 components.

ECM-dependent FAK/MEK/ERK activation controls MITF localization and activity, as well as melanogenesis

Given the ECM-dependent phenotypic shifts in MCs, we next analyzed expression changes in

key signaling pathways— PI3K/Akt (Larribere et al., 2004; Phung et al., 2011; Shi et al., 2016;

C. Wang et al., 2016), Wnt (Colombo et al., 2022; Katkat et al., 2023; Sinnberg et al., 2018),

and MAPK (Buscà & Ballotti, 2000; Estrada et al., 2022; Ngeow et al., 2018; Ostojić et al.,

2021; Wellbrock & Arozarena, 2015)— known to regulate MITF activity, melanocyte

differentiation, and plasticity. While only a few PI3K/Akt-associated genes were differentially

expressed, with *Brca1* and *Sgk2* downregulated and *Sgk1* upregulated on FN (fig. 6A), Wnt

signaling showed a broader modulation (fig. 6B). Among the DEGs, multiple components—

including the canonical target *Ccnd1*, the positive regulators *Frat2*, *Fzd4*, and *Cd88c*, and the

feedback inhibitor *Nkd2*—were upregulated on FN, suggesting an activation of both canonical

and non-canonical Wnt signaling pathways, potentially accompanied by feedback regulation

(fig. 6B). The most pronounced transcriptional changes, however, were observed in MAPK (fig.

6C) and ERK signaling (fig. 6D), reflected by a high number of DEGs associated with these

pathways. Some upstream activators of ERK (*Cnksr1*, *Rasgrp3*, *Pak3*) were downregulated

on FN compared to COL I (fig. 6C). In contrast, genes enhancing ERK signaling (*Igf2bp1*,

Irak2, *Il1rap*, *Ngfr*) and transcriptional ERK targets (*Etv4*, *Il6*, *Tnc*, *Lif*, *Col1a2*) were

upregulated on FN-coated substrates (fig. 6C, D). In addition, upstream modulators of ERK,

such as *Erbb3*, *Axl*, and *Itga6*, were also upregulated (fig. 6C, D).

These data pointed to ECM-driven remodeling of MAPK/ERK signaling, prompting us to

directly assess ERK activation in iMCs grown on COL IV, COL I, and FN. Immunoblot analyses

of ERK1/2 phosphorylation (p-ERK) revealed an ECM-dependent modulation of ERK activity,

with FN-grown iMCs featuring highest p-ERK levels, and COL I-exposed MCs displaying

lowest p-ERK levels (fig. 6E). Considering these differential p-ERK signals observed with the

various ECM types and given that in melanoma cells, MAPK/ERK has recently been shown to

negatively control MITF nuclear localization and activity (Estrada et al., 2022; Ngeow et al.,

2018), we next asked whether MEK/ERK activation was causal for the observed ECM-

dependent MC responses. For this, we used Trametinib, a pharmacological inhibitor targeting

MEK, the upstream kinase mediating ERK phosphorylation and activation (Gilmartin et al.,

2011). Next to the expected loss of ERK phosphorylation (fig. S6A), Trametinib treatment

(indicated as MEKi) increased *Tyr* mRNA expression (fig. 6F) and resulted in increased

nuclear MITF in iMCs exposed to COL IV and FN, reaching the nuclear MITF levels observed on COL I (fig. 6G,H). Congruent with this, melanin production by iMCs cultured on both COL IV- and FN-coated substrates and treated with Trametinib were significantly elevated towards those levels observed in COL I conditions (fig. 6I). Similar results were obtained using an ERK inhibitor Ravoxertinib (Blake et al., 2016; Varga et al., 2020) (fig. 6J-L). This indicates that MEK/ERK activity in MCs negatively regulates nuclear MITF levels. We next aimed to delineate signals that might transduce ECM sensing to activation of ERK. Given the correlation between elevated FAK and ERK activity in FN-exposed MCs and considering that FAK has been reported to transduce ECM signals through MAPK/ERK (Paszek et al., 2005), we inhibited FAK in MCs. As expected, treatment of MCs with the FAK inhibitor Ifebemtinib (Li et al., 2021) resulted in reduced p-FAK levels as well as a reduced number of FAs in MCs (fig. S6B, C). More importantly, FAK inhibition resulted in a reduction of p-ERK levels as assessed by immunocytochemistry (fig. 6M,N) and caused a significant increase in melanin production (fig. 6O). Together, these data demonstrate that FN-mediated MEK/ERK activation counteracts MITF nuclear accumulation and melanin synthesis, suggesting that MEK/ERK — potentially downstream of FAK— act as negative regulators of melanogenesis in response to ECM cues.

In summary, we deciphered that distinct ECM proteins in the MC environment steer melanogenesis through differential activation of the MEK/ERK pathway and subsequent regulation of MITF localization and activity. We further showed that COL I and FN exert opposite effects on MC behavior and functions: COL I elicits a highly pigmented and proliferative, but non-motile MC state, while FN triggers a less pigmented, low proliferative and highly motile MC state. Finally, by combining varying substrate stiffness and distinct ECM types we delineated that stiffness-mediated MC functions rely on specific ECM components.

DISCUSSION

ECM components fine-tune MC phenotypes and functions

Our study identifies ECM components as critical environmental triggers that instruct MC behavior. Through dynamic interactions with the ECM, MCs engage adhesion-dependent signaling, such as FAK activation, enabling them to decode contextual ECM inputs and adapt their phenotype accordingly. Specifically, we observed that compared to the abundant physiological matrix protein that epidermal MCs face (COL IV), COL I and FN – representative of a dermal matrix and thus an altered environment – elicit notable phenotypic shifts in MCs. Relative to COL IV, COL I reduced MC migration and increased melanin production, while FN promoted migration and decreased both proliferation and melanin production. These contrasting effects suggest that ECM composition can selectively modulate distinct aspects of MC behavior. Notably, the intermediate state observed on COL IV supports a model in which

this basement membrane component enables MCs to maintain phenotypic flexibility — for example, allowing them to increase melanin production in response to external stimuli such as UV or inflammation. The opposing responses of MCs to FN and COL I underscore the importance of ECM composition in regulating MC function, suggesting that context-dependent ECM remodeling can actively shape MC behavior, with relevance not only for MC homeostasis but potentially also for pathophysiological states and stress responses.

Our findings open the possibility that ECM alterations can disrupt MC homeostasis, with potential beneficial or detrimental consequences for skin health depending on the context. For instance, wound healing reflects a complex physiological process in which MC phenotypic shifts could have a significant impact. Upon skin injury, the initial stage of tissue repair entails the formation of a provisional FN-rich environment (Potekaev et al., 2021), which may facilitate the repopulation of MCs within the regenerated tissue (Snell, 1963). Taking into account our observation that FN enhances MC motility in vitro (fig. 2), it is tempting to speculate that this FN-enriched tissue enables MCs to efficiently migrate into wound sites and re-establish their protective function, such as melanin-based protection from UV-induced damage. Conversely, in fibrotic conditions such as scleroderma, marked by stiffening of the skin due to excessive deposition of COL I, cases of localized hyperpigmentation have been reported in patients, possibly reflecting COL I-driven MC reprogramming. Vitiligo, which is characterized by the loss of epidermal MCs and non-pigmented patches, is associated with a decrease in COL IV and FN and a concomitant increase in COL I content (Rani et al., 2023). The repigmentation process requires MCs to migrate into depigmented areas and synthesize melanin effectively (Norris et al., 1994). Considering our observation of strongly reduced MC migration on COL I, it seems plausible that COL I enrichment in vitiligo lesions interferes with the efficient MC redistribution into depigmented areas in the skin. Furthermore, melasma is a multifactorial skin condition that is characterized by focal hypermelanosis. Sex hormones and UV radiation have been implicated in the development of melasma (Espósito et al., 2022). Interestingly, disruption of the basement membrane has also been reported in 96% of melasma lesions, with MCs protruding into the dermal layer in 66% of cases (Torres-Álvarez et al., 2011). While the mechanism of this dermal invasion by MCs remains open, it has been proposed that the migration of "hyperactive" MCs into the dermis leads to the constant hyperpigmentation in melasma (Phansuk et al., 2022; Torres-Álvarez et al., 2011). This concept would be in line with our findings of COL I-triggered melanogenesis, opening the possibility that MC hyperactivity in melasma could, at least in part, result from the exposure to dermal COL I.

Together, these examples illustrate how context-dependent ECM remodeling could shape MC behavior, with implications not only for physiological repair but also for pathological skin remodeling. Though further investigation is warranted, these observations open new avenues to explore how shifting ECM landscapes influence MC phenotypes in vivo.

FN-induced phenotypic reprogramming rewires MCs toward a dedifferentiated state

Our study also highlight for the first that ECM components are pivotal in regulating the phenotypic plasticity of MCs. We demonstrate that a FN-rich environment rewires MCs toward a dedifferentiated state, marked by reduced melanin production, slow cycling, and increased motility, while COL I elicits features of a differentiated phenotype, promoting melanin synthesis but limiting migration. In addition to these phenotypic observations, our data indicates that FN exposure induces a distinct transcriptional program associated with MC dedifferentiation. Transcriptomic profiling revealed that FN-cultured MCs downregulate melanocytic differentiation markers and upregulate genes linked to plasticity, stemness, and neural crest-like features. This phenotype is indicative of an adaptive, dedifferentiated state, distinct from the more stable melanogenic profile observed on both COL I and COL IV, and suggests that FN may act as a cue for reprogramming MC identity.

Notably, this dedifferentiated signature observed in FN-exposed MCs is reminiscent of the early phenotypic changes reported during malignant transformation of MCs. Indeed, in a mouse model recapitulating features of human melanomagenesis, it has been shown that mature MCs expressing a B-Raf oncogene can undergo transcriptional reprogramming, associated with a loss of their differentiated characteristics and eventual invasion into the dermis (Köhler et al., 2017). This suggests that MC dedifferentiation precedes MC transformation and melanoma development. Hence, the dedifferentiated phenotype observed in FN-exposed MCs could reflect one of several steps that are required during early stages of melanoma initiation. Taken together, this raises the possibility that a FN-rich environment, combined with oncogenic mutations, could render MCs susceptible to transformation and ultimately melanomagenesis.

In conclusion, our findings underscore the remarkable plasticity of MCs and their ability to shift differentiation states in response to environmental cues, highlighting the importance of ECM molecules in regulating MC behavior. By modulating the balance between melanogenic identity and motile potential, the ECM — particularly FN — may influence not only physiological processes like tissue repair but also pathological trajectories including pigmentation disorders and oncogenesis. Understanding how different conditions elicit these shifts could provide valuable insights into normal skin physiology, the mechanisms underlying melanocyte-related conditions like melanoma, and potential therapeutic targets for restoring normal melanocyte function in pigmentation disorders. In particular, elucidating how ECM remodeling contributes to MC dedifferentiation could open new therapeutic avenues in both regenerative medicine and cancer biology.

ECM-dependent ERK activation controls MITF localization and activity, and melanogenesis

Given the well-established role of MITF as a master regulator of MC differentiation,
pigmentation, and survival, and its known regulation by ERK signaling, we reasoned that it
could function as a key integrator of ECM-derived cues. We therefore centered our analysis
on MITF, aiming to understand how its localization and activity might be shaped by the
extracellular microenvironment. This hypothesis was supported by our finding that ECM-
dependent phenotypic shifts of MCs are tightly linked to differential activation of the MEK/ERK
pathway, which in turn governs the localization and output of MITF. Specifically, COL I limits
ERK activity, triggering high nuclear MITF, while FN stimulates high ERK activity resulting in
reduced nuclear MITF levels. This is further associated with adaptation of MC functions under
the control of MITF, whereby proliferation and melanin production are enhanced on COL I but
reduced on FN. In line with previous studies that linked RAF and MEK/ERK activation to
cytoplasmic retention and nuclear export of MITF in melanoma cells, respectively (Estrada et
al., 2022; Ngeow et al., 2018), we here demonstrate in MCs that MEK and ERK inhibition
reverted the low nuclear MITF and melanin production in FN-exposed MCs. This, together with
our findings on ECM-dependent differential ERK activation in MCs, identifies a hitherto
unrecognized role of ECM cues in MITF nuclear localization through the regulation of ERK
activity. Our data further suggest that FAK may act upstream of ERK, as FAK is more activated
on FN and its inhibition on FN reduced ERK activation and melanin production.

Our transcriptomic analysis also pointed to ECM-induced expression changes in Wnt pathway
components, e.g. an upregulation of *Frat2* and *Ccnd1* on FN. FRAT2 is known to stabilize β-
catenin by inhibiting its GSK3β-mediated degradation (van Amerongen & Berns, 2005),
thereby supporting β-catenin–dependent transcription. Since β-catenin can enhance MITF
transcription through TCF/LEF-mediated activity (Widlund et al., 2002; Yasumoto et al., 2002),
and given that MITF itself can interact with β-catenin to regulate gene expression (Schepsky
et al., 2006), these findings open the possibility that Wnt signaling -next to FAK/MEK/ERK
signalling- could contribute to ECM-dependent modulation of MITF levels and function.

Overall, further studies are needed to dissect the specific molecular signals downstream of FN
that drive the shift of mature MCs toward less differentiated states and to delineate the
respective contributions of individual signaling pathways in ECM-mediated responses.

Interplay between ECM components and substrate stiffness in the mechanosensation
of MCs
Our findings underpin a crucial role for ECM components in regulating mechanosensation in
MCs and show that ECM components and substrate stiffness govern MC functions. So far,
studies have predominantly explored either the roles of different ECM subtypes using ultra-
hard substrates like glass or plastic (Hara et al., 1994), or focused on the impact of substrate
stiffness with a single ECM type (Choi et al., 2014). In contrast, our approach, combining

varying substrate stiffness with different ECM proteins, demonstrated that 1) over the range of stiffness and ECM molecules tested, matrix protein subtypes can dominate over mechanical substrate surface properties, as altering ECM proteins produced more pronounced effects on MC behavior compared to changes in substrate stiffness; and 2) the ability of MCs to respond to mechanical cues is highly dependent on the specific ECM proteins present. Notably, COL I supports stiffness-induced MITF nuclear localization and MC proliferation, while FN enhances mechanosensitive responses at the level of FAK activation and number of FAs. Seong et al. previously reported that in a human fibrosarcoma cell line stiffness-mediated FAK activation can be observed on FN- but not on COL I-coated substrates (Seong et al., 2013). While our observations in MCs are consistent with such effects on FN, iMCs cultured on COL I even showed reduced FAK activity with substrate stiffness, and pMCs had lowest FAK activity at intermediate stiffness. These observations suggest cell type-specific roles of ECM molecules in mechanoresponses.

Strikingly, our data identified MITF as a novel ECM- and mechanosensitive transcription factor, with a stiffness-dependent increase as well as the highest nuclear localization observed on COL I compared to COL IV and FN. Such fine-tuning of MITF subcellular localization may ensure that MC responses are appropriately matched to the environmental matrix protein composition and mechanical features of the ECM. Overall, this highlights the context-dependent nature of MITF regulation downstream of ERK, revealing a hitherto unknown aspect of MITF modulation by ECM cues. Furthermore, our findings suggest that ECM components and substrate stiffness do not merely exert additive effects; instead, individual ECM types demonstrate synergistic interactions with stiffness in regulating MC behavior. This underscores the importance of considering both ECM protein composition and mechanical properties in future studies, as the interplay between these factors significantly influences how cells interpret and respond to mechanical signals, thereby affecting cellular function and differentiation pathways.

Conclusion

In summary, we report a novel ECM-dependent FAK/MEK/ERK/MITF axis that controls the plasticity of MCs in response to interactions with their environment. Our findings underscore the critical role of the ECM in orchestrating MC function and differentiation, highlighting the need for further studies to fully understand the intricate crosstalk between mechanical parameters and ECM ligands in the cellular microenvironment (fig. 7). Moreover, our data point to a complex link between ECM protein types and substrate stiffness, which can influence MC mechanosensation (fig. 7). Elucidating the specific molecular mechanisms driving this interplay, particularly the roles of MITF and other key regulators, will be crucial for a comprehensive understanding of MC function and pathology.

METHODS

Preparation of PDMS substrates with tunable stiffness and ECM protein coating

Polydimethylsiloxane (PDMS) substrates were prepared using a SYLGARD 184 Elastomer kit (Dow Silicones, USA) and varying the crosslinker:silicone base agent ratio to 1:60, 1:35, and 1:10, respectively (fig. S1). Once mixed, the PDMS elastomers were cured at 60°C for 16 h. To enhance the hydrophilicity of the substrates and facilitate binding of ECM proteins, all substrates underwent argon plasma treatment using a PDC-002-CE plasma cleaner (Harrick Plasma, USA) at 300 mTorr for 2 min. The substrates were then coated for 2 h at 37°C with 30 µg/ml of mouse collagen IV (Cultrex™, #3410-010-02, Bio-Techne, Germany), 30 µg/ml of type I collagen from calf skin (Collagen G, L7213, Sigma-Aldrich, USA), or 10 µg/ml of bovine plasma fibronectin (F1141, Sigma-Aldrich, USA) diluted in sterile D-PBS (Gibco, USA). After incubation, the excess ECM solution was aspirated, and substrates were washed with sterile D-PBS prior to cell seeding.

Determination of the elastic modulus of uncoated and coated PDMS surfaces

Atomic Force Microscopy (AFM) served to determine the elastic modulus (Young's Modulus) of uncoated and protein-coated PDMS surfaces used in this study. For this, the PDMS elastomers (ratio 1:10, 1:35 or 1:60) were prepared in an AFM-suitable imaging dish (ibiTreat µ-Dish, Ibidi, Germany) and transferred to the sample stage of the microscope (Nanowizard 4, Bruker Nano GmbH, Germany). An AFM cantilever with a spherical tip (Biosphere B2000, Nanotools, Germany) with a nominal spring constant of 0.2 N/m and a sphere radius of 2 µm was used to perform nanoindentation measurements. AFM force-distance measurements were conducted in a grid of 4 x 4 points with a lateral distance of 4 µm (= one tip diameter) between two points to probe native spots. The AFM cantilever was extended to the PDMS surface with a speed of 1 µm/s and pressed on it with a loading force of 20 nN. Afterwards, the cantilever was retracted with 1 µm/s to complete the force-distance measurements. For each force-distance curve, the Hertz fit (Horvath et al., 2019) was applied to the contact region of the extend curve to evaluate the Young's Modulus with the JPK SPM Data Processing Software v8.0.1 (Bruker Nano GmbH, Germany).

Mice

All mice were housed and maintained according to federal guidelines within the specific pathogen free (SPF) animal facility in Building 61.4 of the Saarland University in Homburg under the breeding license of Sandra Iden's lab (§11 Az. 2.4.1.1- Iden). All experiments were performed in line with German animal welfare laws as well as institutional guidelines and were reported to the responsible authorities (Vorabmeldung VM2023-16 and previous notifications). Wildtype (wt) mice were either purchased from the Jackson Laboratory (Charles River Germany) or were taken from in-house breedings of transgenic mouse lines. In those cases,

F1 and F2 generation were genotyped to verify the wt alleles of the genes of interest of the
respective mouse line.

Isolation and culture of primary mouse melanocytes
Primary mouse MCs were isolated from the epidermis of newborn wt mice and maintained up
to passage 3 as previously described (Mescher et al., 2017). Briefly, to separate epidermis
from dermis, whole skin from mice at postnatal day (P)0-P3 were incubated in a solution of 5
mg/ml Dispase II (#D4693, Sigma-Aldrich, USA) diluted in RPMI medium (Gibco, USA)
supplemented with 10% fetal calf serum (FCS, #S0615, Sigma-Aldrich, USA), penicillin (100
U/ml), streptomycin (100 µg/ml), 100 nM sodium pyruvate, and 10 mM non-essential amino
acids (all Gibco, USA) at 4°C overnight. The next day, the epidermis was incubated for 20 min
in TrypLE select (#12563-011, Gibco, USA) at room temperature. Dissociated cells were
collected and cultured in RPMI medium containing 200 nM TPA (12-O-tetradecanoylphorbol-
13-acetate, #P8139, Sigma-Aldrich, USA) and 200 pM cholera toxin (#C8052, Sigma-Aldrich,
USA), hereafter referred to as RPMI+. At passage 0, the cell cultures contained non pigmented
melanoblasts, MCs, and keratinocytes. After 7 days, cells were passaged, and due to terminal
differentiation, keratinocytes were largely removed, resulting in MC monocultures suitable for
further experiments.

Generation of stable murine melanocyte cell lines (iMCs)
Immortalized MCs (iMCs) were generated by immortalizing mouse primary MCs (mTmG-
positive) with SV40 large T antigen, followed by subcloning. The cells express membrane-
targeted tandem dimer Tomato (dt; Muzumdar et al., 2007), yielding red fluorescence. iMCs
exhibit morphological and molecular characteristics common to normal murine skin MCs. iMCs
were maintained in culture in RPMI+ at 37°C with 5% CO_2.

Immunofluorescence
To assess cell area and morphology, and for quantification of p-FAK, Ki67, MITF, YAP and p-
ERK signal intensities, MCs were seeded on the different substrate combinations at a density
of 2,500 cells/cm², to avoid overcrowding and allow single cell measurements. 24 h later, MCs
were fixed in 4% PFA/PBS for 10 min. For MITF, YAP and p-FAK stainings, cells were blocked
for 1 h at room temperature with 5% bovine serum albumin (BSA) in TBS containing 0.2%
Triton X-100 (named blocking buffer hereafter). Primary antibody incubation was performed in
blocking buffer diluted at a ratio of 1:5 in TBS at 4°C overnight. For Ki67 and p-ERK stainings,
blocking and primary antibody dilution were performed using PB buffer composed of 0.05%
milk powder (#T145.2, Carl Roth, Germany), 0.25% fish gelatin (#G7765, Sigma-Aldrich,
USA), 0.5% Triton X-100 (#BP151-100, Fisher Bioreagents, Germany), 20 mM HEPES (pH
7.2, #9105.4, Carl Roth, Germany), and 0.9% NaCl (#3957.1, Carl Roth, Germany). For all,
after 3x washing in 0.2% Tween/TBS, MCs were stained with AlexaFluor-conjugated

secondary antibodies, DAPI (Carl Roth, Germany) and phalloidin to stain actin for pMCs for 1 h at room temperature. After washing, PDMS scaffolds were mounted onto glass coverslips using Fluoromount-G™ (#00-4958-02, ThermoFisher Scientific, USA). Antibodies used for immunostainings are listed in the Antibodies and reagents section.

BrdU Assay

48 h post-plating, MCs were incubated with 160 µg/ml of BrdU (5-Brom-2′-desoxyuridin; #10280879001, Roche, Germany) for 2 h at 37°C and then washed with PBS and fixed with 4% PFA for 10 min at room temperature. Cells were treated with 2 N HCl for 10 min at room temperature to achieve DNA denaturation, followed by a PBS wash and subsequent immunofluorescence staining as previously described. BrdU-positive MCs were visualized using fluorescence microscopy (see below) and quantified from at least 20 randomly positioned micrographs. The background signal was subtracted from the measured signal of each nucleus to identify BrdU-positive cells. The proliferation rate was then determined by counting BrdU-positive cells and dividing this number by the total number of cells, identified via DAPI staining.

Microscopy

Epifluorescence micrographs were acquired with an AxioObserver Z1 inverted microscope equipped with a Colibri 7 LED light source (Zeiss, Germany) using a Plan-Apochromat 20x/0.8 air objective an EC Plan Neofluar 40x/1.3 oil objective, a Plan-Apochromat 63x/1.4 oil objective and an Axiocam 305 camera (Zeiss, Germany) or Rolera ECM2 camera coupled to the ZEN Blue imaging software V2.6 (Zeiss, Germany).

Image quantification

Image analysis was performed using Fiji/ImageJ software (Schindelin et al., 2012; Schneider et al., 2012) to quantify cell area, morphology, FAK phosphorylation and focal adhesion numbers, ERK phosphorylation, as well as YAP and MITF nuclear translocation in MCs 24 h post-plating.

MC dendricity and cell area: Epifluorescence micrographs of actin (phalloidin staining) or td-Tomato, for pMCs and iMCs respectively, were preprocessed using "mean" filter to smoothen images, by reducing noise and averaging pixel values, followed by the "median" filter, which further reduced noise while preserving edges and details. Images were then converted into binary images by using the "Threshold" function to obtain a mask of the cell shape. The binary mask was used either for cell area quantification or for the quantitative assessment of morphological characteristics of MCs. Using the "Sholl Analysis" plugin, concentric circles (named radius), were drawn on the binary image at regular intervals (5 µm) from the center of the cell body. The number of intersections between dendrites and each radius was counted

and then plotted to create a Sholl profile, which shows the number of dendrite intersections at
increasing distances from the cell body (Binley et al., 2014; Sholl, 1953).

p-FAK immunoreactivity: To quantify p-FAK immunoreactivity in MCs, actin (phalloidin
staining) or td-Tomato signals were used to detect the cell contour of pMCs and iMCs,
respectively. This contour was then superimposed on p-FAK signal, followed by measurement
of the integrated density of the signal of individual MCs. For iMCs, data were pooled after
normalization for each experiment as two different microscope cameras have been used for
these experimental series.

Focal adhesion number: To quantify the number of focal adhesions based on p-FAK signals,
first, a background subtraction was performed using a rolling ball algorithm. Then, a suitable
threshold was applied to isolate the focal adhesion signal from the background. Individual focal
adhesions were detected using the "analyze particles" function (0.05-5 μm). Finally, the
number of focal adhesions per cell and per unit area was determined.

YAP and MITF subcellullar localization: Immunodetection of MITF and YAP were performed in
separate experiments and analyzed independently. For both stainings, appropriate negative
controls were included (secondary antibody alone without primary antibody), which showed no
detectable signal.
To evaluate YAP nuclear localization in response to ECM cues, DAPI was used to delineate
nuclear contours, which were overlaid onto the YAP signal to quantify nuclear integrated
intensity in individual MC. To account for signal variation across experiments and due to the
use of two different cameras along this experimental series, values were normalized within
each independent experiment by dividing each individual measurement by the mean nuclear
YAP intensity across all conditions.
For MITF quantification, phalloidin-based F-actin staining (in case of pMCs) or td-Tomato
signal (for iMCs) was employed to delineate whole-cell contours and DAPI served as nuclear
counterstain. These masks were overlaid onto the MITF signal to extract integrated intensity
values from the nuclear and whole-cell compartments of individual MCs. Cytoplasmic MITF
intensity was calculated by subtracting the nuclear signal from the whole-cell signal. Then,
nuclear and cytoplasmic intensity values were normalized within each independent experiment
by dividing the values of each individual measurement by the mean total intensity across all
conditions, thereby normalizing for inter-experimental variability while preserving relative
differences between ECM conditions. Finally, MITF nuclear localization was expressed as the
percentage of nuclear signal relative to total cellular signal (value for immunoreactivity in the
nucleus/ value for immunoreactivity in the whole cell x 100), allowing comparison of nuclear
enrichment across ECM conditions.

p-ERK immunoreactivity: To quantify p-ERK immunoreactivity in iMCs, the td-Tomato signal was used to detect the cell contour, which was then superimposed on the p-ERK immunostaining signal, followed by measurement of the integrated density of the signal of individual iMCs. Values were normalized within each independent experiment by dividing the values of each individual measurement by the mean nuclear intensity across all conditions. This approach allowed us to account for variability between experiments while preserving relative differences between ECM conditions.

Live-cell imaging for analysis of MC motility

Live-cell imaging was conducted to investigate dynamic cellular behaviors in response to ECM. Taking advantage of the stable expression of the fluorescent membrane-targeted Tomato, MCs were seeded at a density of 1,000 cells/cm² in plastic chamber slide coated with the distinct ECM proteins (μ-Slide 8 Well, Ibidi, Germany), and placed in a live cell imaging system (Zeiss CellDiscoverer 7) equipped with a temperature- and CO_2-controlled chamber. Time-lapse images were acquired every 20 min for 20 h using the epifluorescence microscopy mode. Two methodologies were employed to enable real-time dynamic visualization and quantitative analysis of cell migration. To quantify migration parameters, i.e., total distance migrated by cells, and their speed, cells were tracked using the "Manual tracking" plugin in Fiji software. Additionally, the "Temporal Color Code" in Fiji software was applied to track cell shape and movement over time, allowing to visualize changes of MC dynamics over time. Briefly, different colors represent the shape and position of cells at each time point, with the color gradient indicating the progression of time. The overlay of different colors provides information on the dynamic behavior of cells throughout the observation period.

RT-qPCR

For quantitative RT-PCRs, RNA was isolated from MCs using TRIzol reagent according to the manufacturer's protocol (Invitrogen#15596026, ThermoFisher Scientific, USA). RNA was reversely transcribed using a QuantiTect Reverse Transcription kit (Qiagen, Germany) and amplified with iTaq Universal Probes Supermix (Biorad, USA). Target gene expression was detected using TaqMan probes (GAPDH mouse Mm99999915_g1; Tyr mouse Mm00495817_m1; ThermoFisher Scientific, USA). Gene expression changes were calculated using the comparative CT ($\Delta\Delta$CT) method, normalized to GAPDH expression, and compared to cell lysates from soft substrates coated with COL IV.

Melanin content assay

The melanin assay was performed to quantify the average melanin amount produced by a cell. Both intracellular melanin and extracellular melanin content (i.e., melanin released into the medium) were measured from the same batch of cells. For melanin assays, each condition

was performed in duplicate: one well was used for melanin extraction, and the other for cell counting. MCs were cultured in 12-well plates for 72 h in 1 mL phenol red-free RPMI medium (Gibco, USA) to avoid interference with extracellular melanin measurements. The medium was then collected and analyzed to measure extracellular melanin content. Intracellular melanin was quantified following an adapted protocol from Ito and Wakamatsu (Ito & Wakamatsu, 2003). MCs were lysed in 400 µL of 10% DMSO in 1 N NaOH (extraction buffer) and incubated with agitation (24 h at 100°C for pMCs and 18 h at 80°C for iMCs). Extracellular extracts were quickly centrifuged for 10 sec to sediment cellular debris, and intracellular extracts were centrifuged for 5 min at 13.000 rpm. Extra- and intracellular extracts and their respective blanks, RPMI media or lysis buffer, were loaded onto a 96-well plate and absorbance was measured at 405 nm using a Multiskan™ FC Microplate Photometer (ThermoFisher Scientific, USA). For analysis, the OD values were subtracted by the blank to remove background signals caused by the medium or lysis buffer. Next, extracellular values were multiplied by 5 and intracellular by 2 to account for the stock volume, and the values were divided by the cell number to obtain the relative melanin content per cell.

For intracellular melanin quantification following MITF depletion, iMCs were transfected with 50 nM siMITF (ON-TARGETplus Mouse Mitf siRNA SMARTPool, L-047441-00-0005, Dharmacon, UK) or control siRNA (siGENOME Control Pool, D-001206-14-20, Dharmacon, UK) using the Viromer® Blue kit (Lipocalyx) 4 h after plating. Cells were harvested and lysed for melanin extraction 72 h post-transfection using the same procedure as above.

Transcriptomic analysis

RNA extraction, quality control, library preparation, and sequencing

Total RNA was extracted using TRIzol reagent (Invitrogen, ThermoFisher Scientific, USA) or the RNeasy Mini Kit (Qiagen, Germany), following the manufacturer's instructions. For each condition, RNA was prepared from three biological replicates. RNA quantity and purity were assessed using a NanoDrop One© spectrophotometer (ThermoFisher Scientific, USA), and RNA integrity was verified using the RNA Integrity Number (RIN). Only samples with RIN values ≥4 were processed further. As additional quality controls, melanin content assays and RT-qPCR were performed in parallel, targeting melanocyte-specific markers *Mitf* and *Tyr*.

RNA sequencing was performed by Novogene (Munich, Germany) using standard Illumina protocols. Polyadenylated mRNAs were enriched, fragmented, and reverse-transcribed into cDNA. Strand-specific libraries were prepared and quality-checked using Qubit, real-time PCR, and a Bioanalyzer, then sequenced on Illumina platforms (2×150 bp), generating approximately 25 million paired-end reads per sample. Raw reads were processed with fastp to remove adapters and low-quality reads. Clean reads were aligned to the mouse genome (GRCm39/mm39) using HISAT2, and gene expression was quantified with featureCounts and

normalized as FPKM values. RNAseq data have been deposited at GEO (accession GSE297747).

<u>Bioinformatics analysis</u>

Gene expression data from cells cultured on COL1, COL4, and FN were processed using Pandas (McKinney, 2010) and NumPy (Harris et al., 2020) for sorting and initial exploration. Differential gene expression analysis was performed using the Bioconductor package DESeq2 (Love et al., 2014) to identify genes significantly regulated between the conditions. Genes with an adjusted p-value < 0.05 and |log2 fold change| ≥ 1 were considered differentially expressed. Gene expression values were Euclidean-normalized, and heatmaps were created using Seaborn (Waskom, 2021) to visualize expression patterns across conditions. Results were visualized using volcano plots generated with Matplotlib (Hunter, 2007). To explore biological relevance, association of significantly regulated genes with biochemical pathways was analyzed using Reactome Pathway (Milacic et al., 2024), KEGG Pathway (Kanehisa et al., 2025), and WikiPathways (Agrawal et al., 2024). In addition, Gene Ontology (GO) (Thomas et al., 2022) enrichment analysis was performed to identify overrepresented biological processes, providing insight into the functional impact of ECM proteins on gene regulation.

Preparation of cell extracts, SDS-PAGE, and immunoblotting

MCs cultures were lysed using boiled SDS/EDTA solution (1% SDS, 10 mM EDTA) and genomic DNA was sheared by passing the lysates through a 27 G x ¾ canula. Protein concentrations were quantified using the BCA assay according to the manufacturer's protocol (Pierce, ThermoFisher Scientific, USA). SDS-PAGE was carried out using 8-10% polyacrylamide gels. After completing gel electrophoresis, the proteins were transferred onto a PVDF membrane at 20 V for 60 min. Following the transfer, the membrane was blocked for 1 h at room temperature using 5% BSA/TBST (TBS containing 0.2% Tween 20). The primary antibody was incubated in 5% BSA/TBST overnight at 4°C on a roller. Membranes were then washed 3x for 10 min with TBST and incubated with an HRP-conjugated secondary antibody in 5% BSA/TBST for 1 h at room temperature on a roller. Membranes were washed 3x for 10 min with TBST, excess TBST was removed, and membranes were incubated 1 min with in a 1:1 mixture of ECL solutions A and B (PerkinElmer, USA). Excess ECL solution was removed, and the membrane was imaged using the Amersham ImageQuant™ 800 detector (Cytiva, USA). Following a 30-min wash at room temperature with a buffer containing 0.2 M glycine and 0.1% SDS (pH 2.5) to strip the membranes, they were blocked and immunoblotted for total ERK using the same membranes previously used for phospho-ERK detection.

Quantification of phospho-ERK signal in immunoblot analyses

The protein band intensity of non-saturated Western blot signals was quantified using Fiji software. Each band was selected on the original .tiff image obtained with the Amersham ImageQuant™ 800 Detector, and the "Plot Lanes" tool generated intensity profiles. Peaks corresponding to individual bands (related to total or phosphorylated protein) were identified, and the area under them was measured. The p-ERK/total ERK ratio was calculated to assess relative ERK activity under the different conditions. The values were normalized to the sum of each membrane's values to determine the mean and standard deviation across experiments.

Inhibitor treatment

To assess the role of ERK activity for ECM-dependent MC functions, MCs were seeded 4 h prior a 24h- or 72h-treatment with 100 nM Trametinib (MEK inhibitor, #10999, Biozol, Germany), 10 µM of Ravoxertinib (ERK inhibitor, #S7554, Selleckchem, Germany) or 10 µM of Ifebemtinib (FAK inhibitor, #E1114, Selleckchem, Germany) diluted in RPMI+.

Antibodies

Details on antibodies and dyes used in this study are listed in the following Table S1.

Statistical analyses

Statistical analysis was performed using GraphPad Prism software (GraphPad, version 10.2.1). Measures of pooled data, from at least three independent experiments, are represented by mean and SD or SEM, as indicated in the figure legends. All data sets were subjected to normality tests (Anderson-Darling test, D'Agostino & Pearson test, Kolmogorov-Smirnov test, or Shapiro-Wilk test) when applicable. Then, statistical significance was determined using One-way ANOVA with Welch's correction (unequal variances) with Dunnett's T3 post-hoc test; RM one-way ANOVA with Geisser-Greenhouse's correction (unequal variances) with Tukey's post-hoc test; Ordinary two-way ANOVA Tukey's post-hoc test; Multiple t tests with the Holm-Sidak method's correction with Dunn's post-hoc test; and for non-Gaussian datasets using Kruskal-Wallis test with Dunn's post-hoc test, as indicated in the figure legends. P-values are ranged as follows: *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; and ****, $P < 0.0001$, as detailed in the Table S2. The number of independent biological replicates and number of single cells analyzed is specified in each figure legend and in the Table S2.

Software

Data collection used the following software: Microscopy: ZEN Blue 2.6 (Zeiss, Germany); Immunoblot: Amersham ImageQuant™ 800 control software (Cytiva, USA); qRT-PCR: CFX Opus 96 (BioRad, USA). For data analysis, the following softwares were used: GraphPad PRISM, version 10.2.1, ImageJ/Fiji, JPK SPM Data Processing Software v8.0.1 (Bruker, Nano, Germany).

ACKNOWLEDGEMENTS

We thank the animal facility of the Faculty of Medicine, Saarland University, for important

services, the Robert Ernst lab (Medical Biochemistry and Molecular Biology) for access to the

AxioObserver Z1 microscope, the Department of Cellular Neurophysiology (CIPMM, Saarland

University) for kindly providing access to the plasma cleaner, and Dr. Nicole Ludwig for access

to a TapeStation for RIN measurements. We further thank all members of the Iden laboratory

for stimulating discussions.

CONFLICT OF INTEREST

The authors declare no competing interests.

FUNDING

This project was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation) Projektnummer 200049484; (SFB 1027 projects A12 to S.I., B2 to M.B., and ZX

to V.H.), Projektnummer 519828155 (DFG INST 256/583-1 FUGG) and Projektnummer

466742891 (DFG INST 256/555-1 FUGG).

DATA AVAILABILITY

Correspondence and requests for materials related to this study should be sent to

carole.luthold@uni-saarland.de and sandra.iden@uni-saarland.de. All data supporting the

findings of this study are available within the paper and its supplemental information, or from

the corresponding author on reasonable request.

AUTHOR CONTRIBUTIONS

Conceptualization and Supervision: C.L., S.I.; **Methodology**: C.L., M.D., E.B., V.S.R., G.G.,

S.T., A.K.B., A.R., S.I.; **Validation**: C.L., M.D., E.B., V.S.R., G.G., V.H., S.I.; **Investigation**:

C.L., M.D., V.S.R., E.B., N.D., E.W., G.G., S.T.; **Formal analysis**: C.L., M.D., V.S.R., E.B.,

G.G., S.T.; **Visualization**: C.L., S.T., S.I.; **Resources**: A.K.B., A.R., V.H., M.B., S.I.; **Writing -**

first concept: C.L.; **Writing - original draft**: C.L., S.I.; **Review and editing of manuscript**:

C.L., S.I., A.K.B., V.R:, M.B, V.H.; and **Funding acquisition**: S.I.

REFERENCES

Agrawal, A., Balcı, H., Hanspers, K., Coort, S. L., Martens, M., Slenter, D. N., Ehrhart, F., Digles, D., Waagmeester,
A., Wassink, I., Abbassi-Daloii, T., Lopes, E. N., Iyer, A., Acosta, J. M., Willighagen, L. G., Nishida, K., Riutta,
A., Basaric, H., Evelo, C. T., … Pico, A. R. (2024). WikiPathways 2024: Next generation pathway database.
Nucleic Acids Research, *52*(D1), D679–D689. https://doi.org/10.1093/nar/gkad960
Amano, S. (2009). Possible Involvement of Basement Membrane Damage in Skin Photoaging. *Journal of*
Investigative Dermatology Symposium Proceedings, *14*(1), 2–7. https://doi.org/10.1038/jidsymp.2009.5
Amano, S. (2016). Characterization and mechanisms of photoageing-related changes in skin. Damages of
basement membrane and dermal structures. *Experimental Dermatology*, *25*(S3), 14–19.
https://doi.org/10.1111/exd.13085

Arias-Mejias, S. M., Warda, K. Y., Quattrocchi, E., Alonso-Quinones, H., Sominidi-Damodaran, S., & Meves, A. (2020). The Role of Integrins in Melanoma: A Review. *International Journal of Dermatology*, *59*(5), 525–534. https://doi.org/10.1111/ijd.14850

Arozarena, I., & Wellbrock, C. (2019). Phenotype plasticity as enabler of melanoma progression and therapy resistance. *Nature Reviews Cancer*, *19*(7), 377–391. https://doi.org/10.1038/s41568-019-0154-4

Barbosa, I. A. M., Gopalakrishnan, R., Mercan, S., Mourikis, T. P., Martin, T., Wengert, S., Sheng, C., Ji, F., Lopes, R., Knehr, J., Altorfer, M., Lindeman, A., Russ, C., Naumann, U., Golji, J., Sprouffske, K., Barys, L., Tordella, L., Schübeler, D., … Galli, G. G. (2023). Cancer lineage-specific regulation of YAP responsive elements revealed through large-scale functional epigenomic screens. *Nature Communications*, *14*(1), 3907. https://doi.org/10.1038/s41467-023-39527-w

Benito-Martínez, S., Salavessa, L., Raposo, G., Marks, M. S., & Delevoye, C. (2021). Melanin Transfer and Fate within Keratinocytes in Human Skin Pigmentation. *Integrative and Comparative Biology*, *61*(4), 1546–1555. https://doi.org/10.1093/icb/icab094

Biggs, L. C., Kim, C. S., Miroshnikova, Y. A., & Wickström, S. A. (2020). Mechanical Forces in the Skin: Roles in Tissue Architecture, Stability, and Function. *Journal of Investigative Dermatology*, *140*(2), 284–290. https://doi.org/10.1016/j.jid.2019.06.137

Binley, K. E., Ng, W. S., Tribble, J. R., Song, B., & Morgan, J. E. (2014). Sholl analysis: A quantitative comparison of semi-automated methods. *Journal of Neuroscience Methods*, *225*, 65–70. https://doi.org/10.1016/j.jneumeth.2014.01.017

Blake, J. F., Burkard, M., Chan, J., Chen, H., Chou, K.-J., Diaz, D., Dudley, D. A., Gaudino, J. J., Gould, S. E., Grina, J., Hunsaker, T., Liu, L., Martinson, M., Moreno, D., Mueller, L., Orr, C., Pacheco, P., Qin, A., Rasor, K., … Schwarz, J. B. (2016). Discovery of (S)-1-(1-(4-Chloro-3-fluorophenyl)-2-hydroxyethyl)-4-(2-((1-methyl-1H-pyrazol-5-yl)amino)pyrimidin-4-yl)pyridin-2(1H)-one (GDC-0994), an Extracellular Signal-Regulated Kinase 1/2 (ERK1/2) Inhibitor in Early Clinical Development. *Journal of Medicinal Chemistry*, *59*(12), 5650–5660. https://doi.org/10.1021/acs.jmedchem.6b00389

Bonnans, C., Chou, J., & Werb, Z. (2014). Remodelling the extracellular matrix in development and disease. *Nature Reviews. Molecular Cell Biology*, *15*(12), 786–801. https://doi.org/10.1038/nrm3904

Buscà, R., & Ballotti, R. (2000). Cyclic AMP a key messenger in the regulation of skin pigmentation. *Pigment Cell Research*, *13*(2), 60–69. https://doi.org/10.1034/j.1600-0749.2000.130203.x

Cai, X., Wang, K.-C., & Meng, Z. (2021). Mechanoregulation of YAP and TAZ in Cellular Homeostasis and Disease Progression. *Frontiers in Cell and Developmental Biology*, *9*. https://doi.org/10.3389/fcell.2021.673599

Carreira, S., Goodall, J., Denat, L., Rodriguez, M., Nuciforo, P., Hoek, K. S., Testori, A., Larue, L., & Goding, C. R. (2006). Mitf regulation of Dia1 controls melanoma proliferation and invasiveness. *Genes & Development*, *20*(24), 3426–3439. https://doi.org/10.1101/gad.406406

Centeno, P. P., Pavet, V., & Marais, R. (2023). The journey from melanocytes to melanoma. *Nature Reviews Cancer*, *23*(6), 372–390. https://doi.org/10.1038/s41568-023-00565-7

Cheli, Y., Giuliano, S., Botton, T., Rocchi, S., Hofman, V., Hofman, P., Bahadoran, P., Bertolotto, C., & Ballotti, R. (2011). Mitf is the key molecular switch between mouse or human melanoma initiating cells and their differentiated progeny. *Oncogene*, *30*(20), 2307–2318. https://doi.org/10.1038/onc.2010.598

Choi, H., Kim, M., Ahn, S. I., Cho, E.-G., Lee, T. R., & Shin, J. H. (2014). Regulation of pigmentation by substrate elasticity in normal human melanocytes and melanotic MNT1 human melanoma cells. *Experimental Dermatology*, *23*(3), 172–177. https://doi.org/10.1111/exd.12343

Colombo, S., Petit, V., Wagner, R. Y., Champeval, D., Yajima, I., Gesbert, F., Aktary, Z., Davidson, I., Delmas, V., & Larue, L. (2022). Stabilization of β-catenin promotes melanocyte specification at the expense of the Schwann cell lineage. *Development*, *149*(2), dev194407. https://doi.org/10.1242/dev.194407

Coutant, K., Magne, B., Ferland, K., Fuentes-Rodriguez, A., Chancy, O., Mitchell, A., Germain, L., & Landreville, S. (2024). Melanocytes in regenerative medicine applications and disease modeling. *Journal of Translational Medicine*, *22*(1), 336. https://doi.org/10.1186/s12967-024-05113-x

Cui, Y.-Z., & Man, X.-Y. (2023). Biology of melanocytes in mammals. *Frontiers in Cell and Developmental Biology*, *11*, 1309557. https://doi.org/10.3389/fcell.2023.1309557

Diazzi, S., Tartare-Deckert, S., & Deckert, M. (2020). Bad Neighborhood: Fibrotic Stroma as a New Player in Melanoma Resistance to Targeted Therapies. *Cancers*, *12*(6), 1364. https://doi.org/10.3390/cancers12061364

Domingues, L., Hurbain, I., Gilles-Marsens, F., Sirés-Campos, J., André, N., Dewulf, M., Romao, M., Viaris de Lesegno, C., Macé, A.-S., Blouin, C., Guéré, C., Vié, K., Raposo, G., Lamaze, C., & Delevoye, C. (2020). Coupling of melanocyte signaling and mechanics by caveolae is required for human skin pigmentation. *Nature Communications*, *11*(1), 2988. https://doi.org/10.1038/s41467-020-16738-z

Dupont, S., Morsut, L., Aragona, M., Enzo, E., Giulitti, S., Cordenonsi, M., Zanconato, F., Digabel, J. L., Forcato, M., Bicciato, S., Elvassore, N., & Piccolo, S. (2011). Role of YAP/TAZ in mechanotransduction. *Nature*, *474*(7350), 179–183. https://doi.org/10.1038/nature10137

Durand, S., Tang, Y., Pommier, R. M., Benboubker, V., Grimont, M., Boivin, F., Barbollat-Boutrand, L., Cumunel, E., Dupeuble, F., Eberhardt, A., Plaschka, M., Dalle, S., & Caramel, J. (2024). ZEB1 controls a lineage-specific transcriptional program essential for melanoma cell state transitions. *Oncogene*, *43*(20), 1489–1505. https://doi.org/10.1038/s41388-024-03010-7

Ebrahimighaei, R., Tarassova, N., Bond, S. C., McNeill, M. C., Hathway, T., Vohra, H., Newby, A. C., & Bond, M. (2024). Extracellular matrix stiffness controls cardiac fibroblast proliferation via the nuclear factor-Y (NF-Y) transcription factor. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, *1871*(2), 119640. https://doi.org/10.1016/j.bbamcr.2023.119640

Espósito, A. C. C., Cassiano, D. P., da Silva, C. N., Lima, P. B., Dias, J. A. F., Hassun, K., Bagatin, E., Miot, L. D. B., & Miot, H. A. (2022). Update on Melasma-Part I: Pathogenesis. *Dermatology and Therapy*, *12*(9), 1967–1988. https://doi.org/10.1007/s13555-022-00779-x

Estrada, C., Mirabal-Ortega, L., Méry, L., Dingli, F., Besse, L., Messaoudi, C., Loew, D., Pouponnot, C., Bertolotto, C., Eychène, A., & Druillennec, S. (2022). MITF activity is regulated by a direct interaction with RAF proteins in melanoma cells. *Communications Biology*, *5*(1), Article 1. https://doi.org/10.1038/s42003-022-03049-w

Feng, X., Li, G.-Y., Ramier, A., Eltony, A. M., & Yun, S.-H. (2022). In vivo stiffness measurement of epidermis, dermis, and hypodermis using broadband Rayleigh-wave optical coherence elastography. *Acta Biomaterialia*. https://doi.org/10.1016/j.actbio.2022.04.030

Fisher, G., & Rittié, L. (2018). Restoration of the basement membrane after wounding: A hallmark of young human skin altered with aging. *Journal of Cell Communication and Signaling*, *12*(1), 401–411. https://doi.org/10.1007/s12079-017-0417-3

Gilmartin, A. G., Bleam, M. R., Groy, A., Moss, K. G., Minthorn, E. A., Kulkarni, S. G., Rominger, C. M., Erskine, S., Fisher, K. E., Yang, J., Zappacosta, F., Annan, R., Sutton, D., & Laquerre, S. G. (2011). GSK1120212 (JTP-74057) Is an Inhibitor of MEK Activity and Activation with Favorable Pharmacokinetic Properties for Sustained In Vivo Pathway Inhibition. *Clinical Cancer Research*, *17*(5), 989–1000. https://doi.org/10.1158/1078-0432.CCR-10-2200

Goding, C. R., & Arnheiter, H. (2019). MITF—the first 25 years. *Genes & Development*, *33*(15–16), 983–1007. https://doi.org/10.1101/gad.324657.119

Graham, H. K., McConnell, J. C., Limbert, G., & Sherratt, M. J. (2019). How stiff is skin? *Experimental Dermatology*, *28 Suppl 1*, 4–9. https://doi.org/10.1111/exd.13826

Guilak, F., Cohen, D. M., Estes, B. T., Gimble, J. M., Liedtke, W., & Chen, C. S. (2009). Control of stem cell fate by physical interactions with the extracellular matrix. *Cell Stem Cell*, *5*(1), 17–26. https://doi.org/10.1016/j.stem.2009.06.016

Hara, M., Yaar, M., Tang, A., Eller, M. S., Reenstra, W., & Gilchrest, B. A. (1994). Role of integrins in melanocyte attachment and dendricity. *Journal of Cell Science*, *107*(10), 2739–2748. https://doi.org/10.1242/jcs.107.10.2739

Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van Kerkwijk, M. H., Brett, M., Haldane, A., del Río, J. F., Wiebe, M., Peterson, P., … Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, *585*(7825), 357–362. https://doi.org/10.1038/s41586-020-2649-2

Hirobe, T. (2014). Keratinocytes regulate the function of melanocytes. *Dermatologica Sinica*, *32*(4), 200–204. https://doi.org/10.1016/j.dsi.2014.05.002

Hoek, K. S., Eichhoff, O. M., Schlegel, N. C., Döbbeling, U., Kobert, N., Schaerer, L., Hemmi, S., & Dummer, R. (2008). In vivo switching of human melanoma cells between proliferative and invasive states. *Cancer Research*, *68*(3), 650–656. https://doi.org/10.1158/0008-5472.CAN-07-2491

Horvath, A. N., Holenstein, C. N., Silvan, U., & Snedeker, J. G. (2019). The Protein Mat(ters)-Revealing the Biologically Relevant Mechanical Contribution of Collagen- and Fibronectin-Coated Micropatterns. *ACS Applied Materials & Interfaces*, *11*(44), 41791–41798. https://doi.org/10.1021/acsami.9b12430

Huang, F., Santinon, F., Flores González, R. E., & del Rincón, S. V. (2021). Melanoma Plasticity: Promoter of Metastasis and Resistance to Therapy. *Frontiers in Oncology*, *11*. https://doi.org/10.3389/fonc.2021.756001

Huang, Y., Su, J., Liu, J., Yi, X., Zhou, F., Zhang, J., Wang, J., Meng, X., Si, L., & Wu, C. (2022). YAP Activation in Promoting Negative Durotaxis and Acral Melanoma Progression. *Cells*, *11*(22), 3543. https://doi.org/10.3390/cells11223543

Hunter, J. D. (2007). Matplotlib: A 2D Graphics Environment. *Computing in Science & Engineering*, *9*(3), 90–95. https://doi.org/10.1109/MCSE.2007.55

Iriyama, S., Matsunaga, Y., Takahashi, K., Matsuzaki, K., Kumagai, N., & Amano, S. (2011). Activation of heparanase by ultraviolet B irradiation leads to functional loss of basement membrane at the dermal–epidermal junction in human skin. *Archives of Dermatological Research*, *303*(4), 253–261. https://doi.org/10.1007/s00403-010-1117-5

Iriyama, S., Yasuda, M., Nishikawa, S., Takai, E., Hosoi, J., & Amano, S. (2020). Decrease of laminin-511 in the basement membrane due to photoaging reduces epidermal stem/progenitor cells. *Scientific Reports*, *10*(1), 12592. https://doi.org/10.1038/s41598-020-69558-y

Ishihara, S., & Haga, H. (2022). Matrix Stiffness Contributes to Cancer Progression by Regulating Transcription Factors. *Cancers*, *14*(4), 1049. https://doi.org/10.3390/cancers14041049

Ito, S., & Wakamatsu, K. (2003). Quantitative Analysis of Eumelanin and Pheomelanin in Humans, Mice, and Other Animals: A Comparative Review. *Pigment Cell Research*, *16*(5), 523–531. https://doi.org/10.1034/j.1600-0749.2003.00072.x

Kanehisa, M., Furumichi, M., Sato, Y., Matsuura, Y., & Ishiguro-Watanabe, M. (2025). KEGG: Biological systems database as a model of the real world. *Nucleic Acids Research*, *53*(D1), D672–D677. https://doi.org/10.1093/nar/gkae909

Katkat, E., Demirci, Y., Heger, G., Karagulle, D., Papatheodorou, I., Brazma, A., & Ozhan, G. (2023). Canonical Wnt and TGF-β/BMP signaling enhance melanocyte regeneration but suppress invasiveness, migration, and proliferation of melanoma cells. *Frontiers in Cell and Developmental Biology*, *11*. https://doi.org/10.3389/fcell.2023.1297910

Kawakami, A., & Fisher, D. E. (2017). The master role of microphthalmia-associated transcription factor in melanocyte and melanoma biology. *Laboratory Investigation*, *97*(6), Article 6. https://doi.org/10.1038/labinvest.2017.9

Köhler, C., Nittner, D., Rambow, F., Radaelli, E., Stanchi, F., Vandamme, N., Baggiolini, A., Sommer, L., Berx, G., van den Oord, J. J., Gerhardt, H., Blanpain, C., & Marine, J.-C. (2017). Mouse Cutaneous Melanoma Induced by Mutant BRaf Arises from Expansion and Dedifferentiation of Mature Pigmented Melanocytes. *Cell Stem Cell*, *21*(5), 679-693.e6. https://doi.org/10.1016/j.stem.2017.08.003

Konieczkowski, D. J., Johannessen, C. M., Abudayyeh, O., Kim, J. W., Cooper, Z. A., Piris, A., Frederick, D. T., Barzily-Rokni, M., Straussman, R., Haq, R., Fisher, D. E., Mesirov, J. P., Hahn, W. C., Flaherty, K. T., Wargo, J. A., Tamayo, P., & Garraway, L. A. (2014). A melanoma cell state distinction influences sensitivity to MAPK pathway inhibitors. *Cancer Discovery*, *4*(7), 816–827. https://doi.org/10.1158/2159-8290.CD-13-0424

Larribere, L., Khaled, M., Tartare-Deckert, S., Busca, R., Luciano, F., Bille, K., Valony, G., Eychene, A., Auberger, P., Ortonne, J. P., Ballotti, R., & Bertolotto, C. (2004). PI3K mediates protection against TRAIL-induced apoptosis in primary human melanocytes. *Cell Death & Differentiation*, *11*(10), 1084–1091. https://doi.org/10.1038/sj.cdd.4401475

Li, H., Gao, Y., & Ren, C. (2021). Focal adhesion kinase inhibitor BI 853520 inhibits cell proliferation, migration and EMT process through PI3K/AKT/mTOR signaling pathway in ovarian cancer. *Discover. Oncology*, *12*, 29. https://doi.org/10.1007/s12672-021-00425-6

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550. https://doi.org/10.1186/s13059-014-0550-8

Matthews, N. H., Li, W.-Q., Qureshi, A. A., Weinstock, M. A., & Cho, E. (2017). Epidemiology of Melanoma. In W. H. Ward & J. M. Farma (Eds.), *Cutaneous Melanoma: Etiology and Therapy*. Codon Publications. http://www.ncbi.nlm.nih.gov/books/NBK481862/

McKinney, W. (2010). Data Structures for Statistical Computing in Python. *Scipy*. https://doi.org/10.25080/Majora-92bf1922-00a

Mescher, M., Jeong, P., Knapp, S. K., Rübsam, M., Saynisch, M., Kranen, M., Landsberg, J., Schlaak, M., Mauch, C., Tüting, T., Niessen, C. M., & Iden, S. (2017). The epidermal polarity protein Par3 is a non–cell autonomous suppressor of malignant melanoma. *Journal of Experimental Medicine*, *214*(2), 339–358. https://doi.org/10.1084/jem.20160596

Milacic, M., Beavers, D., Conley, P., Gong, C., Gillespie, M., Griss, J., Haw, R., Jassal, B., Matthews, L., May, B., Petryszak, R., Ragueneau, E., Rothfels, K., Sevilla, C., Shamovsky, V., Stephan, R., Tiwari, K., Varusai, T., Weiser, J., … D'Eustachio, P. (2024). The Reactome Pathway Knowledgebase 2024. *Nucleic Acids Research*, *52*(D1), D672–D678. https://doi.org/10.1093/nar/gkad1025

Miskolczi, Z., Smith, M. P., Rowling, E. J., Ferguson, J., Barriuso, J., & Wellbrock, C. (2018). Collagen abundance controls melanoma phenotypes through lineage-specific microenvironment sensing. *Oncogene*, *37*(23), 3166–3182. https://doi.org/10.1038/s41388-018-0209-0

Morelli, J. G., Yohn, J. J., Zekman, T., & Norris, D. A. (1993). Melanocyte Movement In Vitro: Role of Matrix Proteins and Integrin Receptors. *Journal of Investigative Dermatology*, *101*(4), 605–608. https://doi.org/10.1111/1523-1747.ep12366064

Müller, J., Krijgsman, O., Tsoi, J., Robert, L., Hugo, W., Song, C., Kong, X., Possik, P. A., Cornelissen-Steijger, P. D. M., Geukes Foppen, M. H., Kemper, K., Goding, C. R., McDermott, U., Blank, C., Haanen, J., Graeber, T. G., Ribas, A., Lo, R. S., & Peeper, D. S. (2014). Low MITF/AXL ratio predicts early resistance to multiple targeted drugs in melanoma. *Nature Communications*, *5*, 5712. https://doi.org/10.1038/ncomms6712

Muzumdar, M. D., Tasic, B., Miyamichi, K., Li, L., & Luo, L. (2007). A global double-fluorescent Cre reporter mouse. *Genesis (New York, N.Y.: 2000)*, *45*(9), 593–605. https://doi.org/10.1002/dvg.20335

Ngeow, K. C., Friedrichsen, H. J., Li, L., Zeng, Z., Andrews, S., Volpon, L., Brunsdon, H., Berridge, G., Picaud, S., Fischer, R., Lisle, R., Knapp, S., Filippakopoulos, P., Knowles, H., Steingrímsson, E., Borden, K. L. B., Patton, E. E., & Goding, C. R. (2018). BRAF/MAPK and GSK3 signaling converges to control MITF nuclear export. *Proceedings of the National Academy of Sciences*, *115*(37), E8668–E8677. https://doi.org/10.1073/pnas.1810498115

Norris, D. A., Horikawa, T., & Morelli, J. G. (1994). Melanocyte Destruction and Repopulation in Vitiligo. *Pigment Cell Research*, *7*(4), 193–203. https://doi.org/10.1111/j.1600-0749.1994.tb00049.x

Ostojić, J., Yoon, Y.-S., Sonntag, T., Nguyen, B., Vaughan, J. M., Shokhirev, M., & Montminy, M. (2021). Transcriptional co-activator regulates melanocyte differentiation and oncogenesis by integrating cAMP and MAPK/ERK pathways. *Cell Reports*, *35*(7), 109136. https://doi.org/10.1016/j.celrep.2021.109136

Paszek, M. J., Zahir, N., Johnson, K. R., Lakins, J. N., Rozenberg, G. I., Gefen, A., Reinhart-King, C. A., Margulies, S. S., Dembo, M., Boettiger, D., Hammer, D. A., & Weaver, V. M. (2005). Tensional homeostasis and the malignant phenotype. *Cancer Cell*, *8*(3), 241–254. https://doi.org/10.1016/j.ccr.2005.08.010

Pessoa, D. de O., Rius, F. E., Papaiz, D. D., Ayub, A. L. P., Morais, A. S., de Souza, C. F., da Paixão, V. F., Setubal, J. C., Newton-Bishop, J., Nsengimana, J., Azevedo, H., Reis, E. M., & Jasiulionis, M. G. (2021). Transcriptional signatures underlying dynamic phenotypic switching and novel disease biomarkers in a linear cellular model of melanoma progression. *Neoplasia*, *23*(4), 439–455. https://doi.org/10.1016/j.neo.2021.03.007

Pfisterer, K., Shaw, L. E., Symmank, D., & Weninger, W. (2021). The Extracellular Matrix in Skin Inflammation and Infection. *Frontiers in Cell and Developmental Biology*, *9*. https://www.frontiersin.org/article/10.3389/fcell.2021.682414

Phansuk, K., Vachiramon, V., Jurairattanaporn, N., Chanprapaph, K., & Rattananukrom, T. (2022). Dermal Pathology in Melasma: An Update Review. *Clinical, Cosmetic and Investigational Dermatology*, *15*, 11–19. https://doi.org/10.2147/CCID.S343332

Phung, B., Sun, J., Schepsky, A., Steingrimsson, E., & Rönnstrand, L. (2011). C-KIT Signaling Depends on Microphthalmia-Associated Transcription Factor for Effects on Cell Proliferation. *PLoS ONE*, *6*(8), e24064. https://doi.org/10.1371/journal.pone.0024064

Pinon, P., & Wehrle-Haller, B. (2011). Integrins: Versatile receptors controlling melanocyte adhesion, migration and proliferation. *Pigment Cell & Melanoma Research*, *24*(2), 282–294. https://doi.org/10.1111/j.1755-148X.2010.00806.x

Popovic, A., & Tartare-Deckert, S. (2022). Role of extracellular matrix architecture and signaling in melanoma therapeutic resistance. *Frontiers in Oncology*, *12*. https://www.frontiersin.org/articles/10.3389/fonc.2022.924553

Potekaev, N. N., Borzykh, O. B., Medvedev, G. V., Pushkin, D. V., Petrova, M. M., Petrov, A. V., Dmitrenko, D. V., Karpova, E. I., Demina, O. M., & Shnayder, N. A. (2021). The Role of Extracellular Matrix in Skin Wound Healing. *Journal of Clinical Medicine*, *10*(24), 5947. https://doi.org/10.3390/jcm10245947

Prospéri, M.-T., Giordano, C., Gomez-Duro, M., Hurbain, I., Macé, A.-S., Raposo, G., & D'Angelo, G. (2024). Extracellular vesicles released by keratinocytes regulate melanosome maturation, melanocyte dendricity, and pigment transfer. *Proceedings of the National Academy of Sciences of the United States of America*, *121*(16), e2321323121. https://doi.org/10.1073/pnas.2321323121

Rambow, F., Job, B., Petit, V., Gesbert, F., Delmas, V., Seberg, H., Meurice, G., Van Otterloo, E., Dessen, P., Robert, C., Gautheret, D., Cornell, R. A., Sarasin, A., & Larue, L. (2015). New Functional Signatures for Understanding Melanoma Biology from Tumor Cell Lineage-Specific Analysis. *Cell Reports*, *13*(4), 840–853. https://doi.org/10.1016/j.celrep.2015.09.037

Rambow, F., Marine, J.-C., & Goding, C. R. (2019). Melanoma plasticity and phenotypic diversity: Therapeutic barriers and opportunities. *Genes & Development*, *33*(19–20), 1295–1318. https://doi.org/10.1101/gad.329771.119

Rani, S., Pervaiz, N., Parsad, D., & Kumar, R. (2023). Differential expression of extracellular matrix proteins in the lesional skin of vitiligo patients. *Archives of Dermatological Research*. https://doi.org/10.1007/s00403-023-02628-z

Schepsky, A., Bruser, K., Gunnarsson, G. J., Goodall, J., Hallsson, J. H., Goding, C. R., Steingrimsson, E., & Hecht, A. (2006). The microphthalmia-associated transcription factor Mitf interacts with beta-catenin to determine target gene expression. *Molecular and Cellular Biology*, *26*(23), 8914–8927. https://doi.org/10.1128/MCB.02299-05

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, *9*(7), 671–675. https://doi.org/10.1038/nmeth.2089

Scott, G., Ryan, D. H., & McCarthy, J. B. (1992). Molecular Mechanisms of Human Melanocyte Attachment to Fibronectin. *Journal of Investigative Dermatology*, *99*(6), 787–794. https://doi.org/10.1111/1523-1747.ep12614749

Seong, J., Tajik, A., Sun, J., Guan, J.-L., Humphries, M. J., Craig, S. E., Shekaran, A., García, A. J., Lu, S., Lin, M. Z., Wang, N., & Wang, Y. (2013). Distinct biophysical mechanisms of focal adhesion kinase mechanoactivation by different extracellular matrix proteins. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(48), 19372–19377. https://doi.org/10.1073/pnas.1307405110

Shi, H., Lin, B., Huang, Y., Wu, J., Zhang, H., Lin, C., Wang, Z., Zhu, J., Zhao, Y., Fu, X., Lou, Z., Li, X., & Xiao, J. (2016). Basic fibroblast growth factor promotes melanocyte migration via activating PI3K/Akt-Rac1-FAK-JNK and ERK signaling pathways. *IUBMB Life*, *68*(9), 735–747. https://doi.org/10.1002/iub.1531

Sholl, D. A. (1953). Dendritic organization in the neurons of the visual and motor cortices of the cat. *Journal of Anatomy*, *87*(4), 387–406.

Sinnberg, T., Levesque, M. P., Krochmann, J., Cheng, P. F., Ikenberg, K., Meraz-Torres, F., Niessner, H., Garbe, C., & Busch, C. (2018). Wnt-signaling enhances neural crest migration of melanoma cells and induces an invasive phenotype. *Molecular Cancer*, *17*(1), 59. https://doi.org/10.1186/s12943-018-0773-5

Snell, R. S. (1963). A study of the melanocytes and melanin in a healing deep wound. *Journal of Anatomy*, *97*(Pt 2), 243–253.

Tan, X., Yan, Y., Song, B., Zhu, S., Mei, Q., & Wu, K. (2023). Focal adhesion kinase: From biological functions to therapeutic strategies. *Experimental Hematology & Oncology*, *12*(1), 83. https://doi.org/10.1186/s40164-023-00446-7

Thomas, P. D., Ebert, D., Muruganujan, A., Mushayahama, T., Albou, L.-P., & Mi, H. (2022). PANTHER: Making genome-scale phylogenetics accessible to all. *Protein Science*, *31*(1), 8–22. https://doi.org/10.1002/pro.4218

Tirosh, I., Izar, B., Prakadan, S. M., Wadsworth, M. H., Treacy, D., Trombetta, J. J., Rotem, A., Rodman, C., Lian, C., Murphy, G., Fallahi-Sichani, M., Dutton-Regester, K., Lin, J.-R., Cohen, O., Shah, P., Lu, D., Genshaft, A. S., Hughes, T. K., Ziegler, C. G. K., … Garraway, L. A. (2016). Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science (New York, N.Y.)*, *352*(6282), 189–196. https://doi.org/10.1126/science.aad0501

Torres-Álvarez, B., Mesa-Garza, I. G., Castanedo-Cázares, J. P., Fuentes-Ahumada, C., Oros-Ovalle, C., Navarrete-Solis, J., & Moncada, B. (2011). Histochemical and immunohistochemical study in melasma: Evidence of damage in the basal membrane. *The American Journal of Dermatopathology*, *33*(3), 291–295. https://doi.org/10.1097/DAD.0b013e3181ef2d45

Tsoi, J., Robert, L., Paraiso, K., Galvan, C., Sheu, K. M., Lay, J., Wong, D. J. L., Atefi, M., Shirazi, R., Wang, X., Braas, D., Grasso, C. S., Palaskas, N., Ribas, A., & Graeber, T. G. (2018). Multi-stage differentiation defines melanoma subtypes with differential vulnerability to drug-induced iron-dependent oxidative stress. *Cancer Cell*, *33*(5), 890-904.e5. https://doi.org/10.1016/j.ccell.2018.03.017

van Amerongen, R., & Berns, A. (2005). Re-evaluating the role of Frat in Wnt-signal transduction. *Cell Cycle (Georgetown, Tex.)*, *4*(8), 1065–1072.

Varga, A., Soria, J.-C., Hollebecque, A., LoRusso, P., Bendell, J., Huang, S.-M. A., Wagle, M.-C., Okrah, K., Liu, L., Murray, E., Sanabria-Bohorquez, S. M., Tagen, M., Dokainish, H., Mueller, L., & Burris, H. (2020). A First-in-Human Phase I Study to Evaluate the ERK1/2 Inhibitor GDC-0994 in Patients with Advanced Solid Tumors. *Clinical Cancer Research*, *26*(6), 1229–1236. https://doi.org/10.1158/1078-0432.CCR-19-2574

Walma, D. A. C., & Yamada, K. M. (2020). The extracellular matrix in development. *Development*, *147*(10), dev175596. https://doi.org/10.1242/dev.175596

Wang, C., Zhao, L., Su, Q., Fan, X., Wang, Y., Gao, S., Wang, H., Chen, H., Chan, C. B., & Liu, Z. (2016).
Phosphorylation of MITF by AKT affects its downstream targets and causes TP53-dependent cell
senescence. *The International Journal of Biochemistry & Cell Biology*, *80*, 132–142.
https://doi.org/10.1016/j.biocel.2016.09.029
Wang, K., Wen, D., Xu, X., Zhao, R., Jiang, F., Yuan, S., Zhang, Y., Gao, Y., & Li, Q. (2023). Extracellular matrix
stiffness-The central cue for skin fibrosis. *Frontiers in Molecular Biosciences*, *10*, 1132353.
https://doi.org/10.3389/fmolb.2023.1132353
Waskom, M. L. (2021). seaborn: Statistical data visualization. *Journal of Open Source Software*, *6*(60), 3021.
https://doi.org/10.21105/joss.03021
Wellbrock, C., & Arozarena, I. (2015). Microphthalmia-associated transcription factor in melanoma development
and MAP-kinase pathway targeted therapy. *Pigment Cell & Melanoma Research*, *28*(4), 390–406.
https://doi.org/10.1111/pcmr.12370
Widlund, H. R., Horstmann, M. A., Price, E. R., Cui, J., Lessnick, S. L., Wu, M., He, X., & Fisher, D. E. (2002). Beta-
catenin-induced melanoma growth requires the downstream target Microphthalmia-associated
transcription factor. *The Journal of Cell Biology*, *158*(6), 1079–1087.
https://doi.org/10.1083/jcb.200202049
Wouters, J., Kalender-Atak, Z., Minnoye, L., Spanier, K. I., De Waegeneer, M., Bravo González-Blas, C., Mauduit,
D., Davie, K., Hulselmans, G., Najem, A., Dewaele, M., Pedri, D., Rambow, F., Makhzami, S., Christiaens,
V., Ceyssens, F., Ghanem, G., Marine, J.-C., Poovathingal, S., & Aerts, S. (2020). Robust gene expression
programs underlie recurrent cell states and phenotype switching in melanoma. *Nature Cell Biology*,
22(8), 986–998. https://doi.org/10.1038/s41556-020-0547-3
Yasumoto, K., Takeda, K., Saito, H., Watanabe, K., Takahashi, K., & Shibahara, S. (2002). Microphthalmia-
associated transcription factor interacts with LEF-1, a mediator of Wnt signaling. *The EMBO Journal*,
21(11), 2703–2714. https://doi.org/10.1093/emboj/21.11.2703
Yoshihisa, Y., Norisugi, O., Matsunaga, K., Nishihira, J., & Shimizu, T. (2014). Involvement of MIF in basement
membrane damage in chronically UVB-exposed skin in mice. *PloS One*, *9*(2), e89569.
https://doi.org/10.1371/journal.pone.0089569

FIGURE LEGENDS

Figure 1: ECM components determine the morphology and adhesion of immortalized
melanocytes. iMCs were cultured overnight on substrates of varying stiffness defined here
as soft (PDMS ratio 1:60 ≈40kPa), intermediate (PDMS ratio 1:35 ≈130kPa) and stiff (PDMS
ratio 1:10 ≈1MPa) coated with COL IV, COL I or FN. **A. Top:** Representative micrographs of
iMCs, expressing td-Tomato, cultured overnight on stiff substrates coated with COL IV, COL I
or FN; scale bar: 20µm. **Bottom**: Representative binary image of cells used for Sholl analysis,
with radius drawn every 5µm. **B.** Quantification of the number of dendrites of iMCs; graph
indicating the percentage of cells exhibiting 2 or 3 dendrites, between 4 and 7, or more than
7 dendrites; mean±SEM, N=3, n(cells)≥85. **C.** Graphs showing the Sholl profile analysis,
plotting the number of dendrite intersections against the distance from the cell body. The
SEM are represented by the connecting curve (dotted line); N=3, n(cells)≥85. Each curve
represents an ECM type, and each graph represents a substrate stiffness condition; Welch
ANOVA test: **: $p<0.005$; *: $p<0.05$. **D.** Representative micrographs of iMCs (expressing
membrane-targeted tandem dimer (td) Tomato) cultured on stiff substrates coated with
COL IV, COL I or FN and immunostained for p-Y397-FAK and with DAPI; scale bar: 20µm.
E. Quantification of the cell area of iMCs; violin plots display the medians and distributions of
cell area in each condition; N=3, n(cells)≥69; Kruskal–Wallis test: ****: $p<0.0001$; **: $p<0.005$;

*: p<0.05. **F.** Quantification of p-FAK levels; violin plots show the medians and distributions of the integrated density of p-FAK per cell; N≥4, n(cells)≥82; Kruskal–Wallis test: ****: p<0.0001; ***: p<0.001; *: p<0.05. **G.** Quantification of the number of focal adhesions per µm^2 per cell with violin plots displaying medians and distributions; N≥3, n(cells)≥65; Kruskal–Wallis test: ****: p<0.0001; *: p<0.05. Abbreviations: A.U., arbitrary units.

Figure 2: Differential effects of ECM proteins on iMC motility, with COL I decreasing and FN increasing motility. A. Image showing the temporal color code analysis from live-cell imaging of iMCs cultured overnight in plastic chamber coated with COL IV, COL I or FN to visualize cell dynamics over time. The different colors represent morphology and position of the iMCs at a given time (20min intervals), with the color gradient indicating time progression (from 0 to 240min); scale bar: 50µm. **B.** Tracks showing the path and distance of cells (one cell = one track) relative to the point of origin (time point zero) in the x and y plane. **C.** Quantification of total distance traveled by iMCs; violin plots show the medians and distributions; N=3, n(cells)≥113; Kruskal–Wallis test: ****: p<0.0001; *: p<0.05. **D.** Quantification of mean speed of iMCs; violin plots show the medians and distributions N=3, n(cells)≥113; Kruskal–Wallis test: ****: p<0.0001.

Figure 3: COL I stimulates melanin production, and FN counteracts iMC proliferation. A. Representative example of total melanin produced by iMCs cultured 72 h on substrates of varying stiffness (soft, intermediate or stiff) coated with COL IV, COL I or FN. **B.** Quantification of intra- and extracellular melanin content by spectrophotometry at 405 nm from iMCs cultured as in (A); means±SD, N≥5; Welch ANOVA test: **: p<0.005; *: p<0.05. **C.** Quantification of *Tyr* mRNA expression of iMCs cultured overnight on substrates of varying stiffness (soft, intermediate or stiff) coated with COL IV, COL I or FN; means±SD, N=5; Kruskal–Wallis test: **: p<0.005; *: p<0.05. **D.** Percentage of BrdU-positive cells from iMCs cultured 48 h on substrates of varying stiffness (soft, intermediate or stiff) coated with COL IV, COL I or FN; means±SD, N≥4; ordinary two-way ANOVA: ****: p<0.0001; ***: p<0.001; **: p<0.005; *: p<0.05. **E.** Percentage of Ki67-positive iMCs cultured 48 h on stiff substrates; means±SD, N=4; Welch ANOVA test: *: $p < 0.05$. **F.** Percentage of dividing cells from live-cell imaging of iMCs cultured overnight on COL IV, COL I and FN; means±SD, N=3; RM one-way ANOVA test: *: p<0.05. Abbreviations: A.U., arbitrary units.

Figure 4: MITF nuclear localization is modulated by both ECM molecules and substrate stiffness. A. Representative micrographs of iMCs expressing td-Tomato cultured overnight on stiff substrates coated with COL IV, COL I or FN and stained for MITF and DAPI; scale bar: 20µm. A pseudo-color intensity scale ('fire' color map) was applied to enhance visualization of MITF nuclear localization. **B.** Quantification of cytoplasmic and nuclear MITF

levels; graphs show the mean±SEM of nuclear and cytoplasmic MITF intensities per cell 24

h post-plating on soft, intermediate, or stiff substrates coated with COL IV, COL I, or FN; N≥3,

n(cells)≥89; Kruskal–Wallis test for nuclear MITF: ****: p < 0.0001; *: p<0.05; Kruskal–Wallis

test for total MITF: $$$$: p < 0.0001; $$$: p < 0.001; $$: p<0.005; $: p<0.05.**C**. Quantification

of nuclear MITF levels; violin plots show the medians and distributions of the percentage of

nuclear MITF per cell; N≥3, n(cells)≥67; Kruskal–Wallis test: ****: p<0.0001; *: p<0.05. **D**.

Quantification of intracellular melanin levels following siRNA-mediated MITF knockdown.

iMCs were transfected with control siRNA (siCtrl) or MITF-targeting siRNA (siMITF) 4 h after

plating on COL I or FN. Seventy-two hours post-transfection, cells were harvested and

melanin content was measured by spectrophotometry at 405 nm; means±SD, N=3; Welch

ANOVA test: **: p<0.005; *: p<0.05. **E**. Representative micrographs of iMCs expressing td-

Tomato cultured overnight on stiff substrates coated with COL IV, COL I or FN and stained

for YAP and DAPI; scale bar: 20µm. A pseudo-color intensity scale ('fire' color map) was

applied to enhance visualization of YAP nuclear localization. **F**. Quantification of nuclear YAP

levels; violin plots show the medians and distributions of the nuclear integrated density of

YAP per cell; N=3, n(cells)≥71; Kruskal–Wallis test: ***: p<0.001; *: p<0.05. Abbreviations:

A.U., arbitrary units; td-Tomato, membrane-targeted tandem dimer Tomato.

Figure 5: ECM-dependent transcriptional changes in iMCs. iMCs were cultured overnight

on stiff substrates coated with COL IV, COL I or FN. Gene expression heatmaps showing

transcript abundance across ECM conditions. Darker blue indicates higher expression; lighter

blue indicates lower expression; bold labels indicate statistically differentially expressed

genes. Displayed values represent normalized expression levels per gene across conditions

(e.g., standard scaling), enabling comparison of the same gene between ECM conditions

despite differences in expression range. Panels show: **A.** differentially expressed MITF-

dependent genes; **B.** genes involved in melanin biosynthesis; **C–E.** genes related to

apoptosis, cell cycle, and senescence, respectively; **F–H.** signature genes of the

differentiated, intermediate, and dedifferentiated states; genes were grouped by clustering of

expression profiles; **I.** genes involved in ECM dynamics; **J.** schematic overview of integrins

commonly expressed in MCs; and **K.** integrin gene expression across COL IV, COL I, and

FN conditions.

Figure 6: ECM-mediated ERK activation regulates MITF localization and

melanogenesis. A-D. iMCs were cultured overnight on stiff substrates coated with COL IV,

COL I or FN. Gene expression heatmaps of signaling pathway components across ECM

conditions. Darker blue indicates higher transcript abundance; lighter blue indicates lower

abundance; bold labels denote statistically differentially expressed genes. Heatmaps

illustrate: **A.** genes involved in the PI3K/Akt signaling; **B.** components of the Wnt signaling;

C. genes associated with MAPK signaling; and **D.** genes associated with MAPK signaling. **E.** iMCs were cultured overnight on substrates of varying stiffness (soft, intermediate or stiff) coated with COL IV, COL I or FN. Representative Western blot and associated graph depicting the quantification of p-ERK1/2 Thr202/Tyr204 to total ERK2 ratio; means±SD; COL IV, N=4; COL I and FN, N=5; Welch ANOVA test: **: $p<0.005$ and ‡ : $p<0.05$; and Kruskal–Wallis test: ¤ : $p<0.05$. **F-O.** iMCs were cultured on stiff substrates coated with COL IV, COL I or FN. 4 h post-plating, a 16-hrs (F-H) or 72-hrs (I) treatment with DMSO or 100 nM of MEK inhibitor (MEKi = Trametinib), a 16-hrs (J, K) or 72-hrs (L) treatment with DMSO or 10 µM of ERK inhibitor (ERKi = Ravoxertinib) or a 16-hrs (M, N) or 72-hrs (O) treatment with DMSO or 10 µM of FAK inhibitor (FAKi = Ifebemtinib) was commenced. **F.** Quantification of *Tyr* mRNA expression after MEK inhibitor treatment; means±SD, N=3; multiple t tests: **: $p<0.005$; *: $p<0.05$. **G**. Representative immunostainings for MITF following MEK inhibition; scale bar: 10µm. A pseudo-color intensity scale ("orange hot" color map) was applied to enhance visualization of MITF nuclear localization. **H**. Quantification of (G); violin plots showing the medians and distributions of the integrated density of nuclear MITF per cell; N=3, n(cells)≥63; Kruskal–Wallis test: ****: $p<0.0001$; ***: $p<0.001$. **I**. Quantification of intra- and extracellular melanin content following MEK inhibition by spectrophotometry at 405 nm; means±SD, N=3; multiple t tests: ***: $p<0.001$; *: $p<0.05$. **J**. Representative immunostainings for MITF following ERK inhibition; scale bar: 20µm. A pseudo-color intensity scale ("orange hot" color map) was applied to enhance visualization of MITF nuclear localization. **K**. Quantification of (J); violin plots showing the medians and distributions of the integrated density of nuclear MITF per cell; N=3, n(cells)≥89; Kruskal–Wallis test: ****: $p<0.0001$; *: $p<0.05$. **L**. Quantification of intra- and extracellular melanin content following ERK inhibition by spectrophotometry at 405 nm; means±SD, N=4; Welch ANOVA tests: ***: $p<0.001$; **: $p<0.005$; *: $p<0.05$. **M**. Representative immunostainings for p-ERK following FAK inhibition; scale bar: 20µm. A pseudo-color intensity scale ("fire" color map) was applied to enhance visualization of p-ERK intensity. **N**. Quantification of (M); violin plots showing the medians and distributions of the integrated density of p-ERK per cell; N=3, n(cells)≥83; Kruskal–Wallis test: ****: $p<0.0001$; ***: $p<0.001$; **: $p<0.005$; *: $p<0.05$. **O**. Quantification of intra- and extracellular melanin content following FAK inhibition by spectrophotometry at 405 nm; means±SD, N=5; multiple t tests: ***: $p<0.001$; **: $p<0.005$; *: $p<0.05$. Abbreviations: A.U., arbitrary units.

Figure 7 (graphical abstract). Upper panel: MC differentiation is differentially modulated by ECM components. COL I promotes a differentiated phenotype by limiting ERK activation, leading to high nuclear MITF levels, associated with increased pigmentation and proliferation but reduced motility. In contrast, FN rewires MCs towards dedifferentiation, characterized by enhanced ERK activity resulting in reduced nuclear MITF and decreased pigmentation, and

associated with a slow-cycling phenotype as well as increased motility. Reducing ERK activity, through FAK, MEK or ERK inhibition, restores the differentiation of MCs exposed to FN. **Lower panel:** ECM components influence how cells interpret and respond to mechanical signals: COL I enables stiffness-mediated MITF nuclear localization and MC proliferation, while FN supports a stiffness-dependent increase in focal adhesion number and FAK activation.

Figure S1: Validation of PDMS elasticity using Atomic Force Microscopy. Young's modulus of PDMS substrates with varying crosslinker ratios. **A**. PDMS substrate with a crosslinker:silicone base agent ratio of 1:60; means±SD, N=3; Welch ANOVA tests: ns, $p > 0.05$. **B**. PDMS substrate with a 1:35 ratio; means±SD, N≥3; Welch ANOVA tests: n.s.: $p > 0.05$. **C**. PDMS substrate with a 1:10 ratio; means±SD, N=3; Welch ANOVA tests: n.s.: $p > 0.05$.

Figure S2: Sholl profiles for iMCs exposed to different ECM molecules. A. Graphs showing the Sholl profile analysis, plotting the number of dendrite intersections against the distance from the cell body. The SEM are represented by the connecting curve (dotted line); N=3, n(cells)≥85. Each curve represents a substrate stiffness condition, and each graph represents an ECM type (same cells as analyzed in fig. 1C, different plotting of the data); Welch ANOVA test: n.s.: $p > 0.05$.

Figure S3: Response of primary MCs to distinct ECM cues. pMCs were cultured overnight on substrates of varying stiffness (soft, intermediate or stiff) coated with COL IV, COL I or FN. **A.** Violin plots showing the medians and distributions of the cell area of pMCs; N≥5, n(cells)≥96; Kruskal–Wallis test: ****: $p < 0.0001$. **B.** Quantification of p-FAK levels; violin plots showing the medians and distributions of the integrated density of total p-FAK per cell; N=3, n(cells)≥52; Kruskal–Wallis test: ****: $p < 0.0001$; ***: $p < 0.001$; **: $p < 0.005$. **C.** Quantification of FAs; violin plots showing the medians and distributions of the number of focal adhesions per µm² per cell; N=3, n(cells)≥65; Kruskal–Wallis test: ****: $p < 0.0001$; **: $p < 0.005$; *: $p < 0.05$. **D.** Quantification of intra- and extracellular melanin content by spectrophotometry at 405 nm from pMCs cultured 72 h on substrates of varying stiffness (soft, intermediate or stiff) coated with COL IV, COL I or FN; means±SD, N=3; multiple t tests: *: $p < 0.05$. **E.** Quantification of nuclear MITF; violin plots showing the medians and distributions of the percentage of nuclear MITF per cell; N=3, n(cells)≥67; Kruskal–Wallis test: ****: $p < 0.0001$; ***: $p < 0.001$; **: $p < 0.005$; *: $p < 0.05$. Abbreviations: A.U., arbitrary units.

Figure S4: Negative correlation between nuclear MITF levels and MC area and siMITF validation. A. The percentage of nuclear MITF was plotted against the cell area of the corresponding MC for all stiffnesses and ECM types tested. The bar diagram depicts the

means±SEM of the ratio of cell area to percentage of nuclear MITF per cell; N=3, n(cells)≥47; Kruskal–Wallis test: ****: $p<0.0001$; ***: $p<0.001$; **: $p<0.005$; *: $p<0.05$. **B.** Western blot analysis of MITF expression in iMCs plated on COL I or FN and transfected for 72 hours with siCtrl or siMITF, performed in parallel with melanin quantification shown in Fig. 4D.

Figure S5: Transcriptomic profiling across ECM conditions. A. Principal Component Analysis (PCA) of gene expression across substrates. Each dot represents one sample: COL IV (green), COL I (blue), and FN (purple). Axes indicate principal components capturing the highest variance. **B.** Venn diagram showing the number of genes expressed in the samples grown on COL IV, COL I and FN substrates. **C.** Volcano plots depicting differentially expressed genes between substrates: (i) COL I vs. COL IV, (ii) FN vs. COL I, and (iii) FN vs. COL IV. Significantly downregulated genes are shown in blue; upregulated genes in yellow. **D.** Gene Ontology (GO) Biological Process enrichment analysis of significantly regulated genes across substrates. The x-axis indicates the adjusted p-value of Fisher's Exact test; the y-axis lists the most significantly enriched biological processes. **E.** Reactome pathway enrichment analysis performed on differentially expressed genes across ECM conditions. The x-axis represents the adjusted p-value of Fisher's Exact test; the y-axis lists the top significantly enriched Reactome pathways.

Figure S6: Functional validation of MEK and FAK inhibitors in iMCs cultured on ECM-coated substrates. iMCs were cultured on stiff substrates coated with COL IV, COL I or FN. 4 h post-plating, a 16-hrs treatment with DMSO, 100 nM of MEK inhibitor (MEKi = Trametinib) or FAK inhibitor (FAKi = Ifebemtinib) was commenced. **A.** Representative Western blot showing the inhibition of ERK1/2 phosphorylation with Trametinib treatment. **B**. Quantification of p-FAK (Y397) intensity per cell following treatment with DMSO or FAK inhibitor (FAKi); violin plots display median and distribution; N=3, n(cells)≥88; Kruskal–Wallis test: ****: $p<0.0001$. **C**. Quantification of focal adhesion (FA) number per cell following FAKi treatment; violin plots show median and distribution; N=3, n(cells)≥70; Kruskal–Wallis test: ****: $p<0.0001$.

A COL IV COL I FN

time color scale 0 / 240 min

B COL IV COL I FN

migration (in µm) vs migration (in µm)

C Cell migration

Total distance migrated by a cell (µm)

COL IV COL I FN

D Cell velocity

Ccell velocity in µm/h

COL IV COL I FN

A   **Total melanin production**

B   **Melanin production**

D   **Cell proliferation**

C   *Tyr* expression

E   **Cell proliferation**

F   **Cell division**

A   **Correlation between nuclear MITF and cell area**

B

	siCTL	siMITF	
MITF			
Calnexin			
COL I	+	-	+	-
FN	-	+	-	+

Luthold et al **Supplementary tables**

Supplementary Table S1

Primary Antibodies		Company	Catalog Nr.
IF	Anti-MITF antibody [C5]	Abcam	Ab12039
	Phospho-FAK (Tyr397) antibody	Cell Signaling Technology	3283S
	YAP (D8H1X) XP®	Cell Signaling Technology	14074S
	BrdU	Biolegend	339802
	Ki67	Biolegend	652401
	pERK1/2	Cell Signaling Technology	4370
WB	pERK1/2	Cell Signaling Technology	4370
	ERK2	BD Pharmingen	610104
	MITF	Cell Signaling	97800
	Calnexin	Enzo Life Sciences	ADI-SPA-860-D
Conjugated Secondary Antibodies			
IF	Alexa Fluor™ 488 donkey anti-rabbit	Invitrogen	A-21206
	Alexa Fluor™ 647 goat anti-mouse	Invitrogen	A-21235
WB	HRP donkey anti-rabbit	Invitrogen	A16023
	HRP sheep anti-mouse	Amersham	NA931V
Dyes			
IF	Alexa Fluor™ 568 Phalloidin	Molecular Probes	A12380
	DAPI	Carl Roth	6335.1

Supplementary Table S2

		COL IV			COL I			FN		
		1:60	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**
	N exp	3	3	3	3	3	3	3	3	3
	n cells	85	89	107	138	136	153	126	140	146
mean	**2 to 3**	15,5555556	20,5	24,6187364	60,0461425	59,359738	53,2537914	7,15242122	3,12178388	9,71516972
	4 to 7	68,8888889	61,2083333	61,4379085	38,6231082	38,2012376	39,3942329	67,1086278	68,2443539	61,9410319
	>7	15,5555556	18,2916667	13,9433551	1,33074935	2,43902439	7,35197563	25,738951	28,6338622	28,3437983
SEM	**2 to 3**	1,41219758	1,0261342	1,8258221	1,66918658	0,57102161	1,4985177	1,1611834	1,76150372	2,02460515
	4 to 7	1,32577898	0,82816685	0,75070439	1,90380086	0,22108701	1,18338293	0,29768777	0,64682354	1,4000745
	>7	1,41219758	2,19464868	3,77926573	0,84755901	2,43902439	1,69692285	1,00441296	1,61600055	1,42235041

fig 1B - MC dendricity complexity

fig 1C - sholl profiles

descriptive statistics

	COL IV			COL I			FN		
	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10
N exp	3	3	3	3	3	3	3	3	3
n cells	85	89	107	138	136	153	126	140	146
Area under the curve	**258,2**	**259,2**	**249,3**	**148,9**	**132,1**	**149,6**	**303**	**318,5**	**340,1**
Std. Error	28,97	27,3	24,41	14,97	15,55	20,01	30,29	29,88	29,15

	Significant?	P Value
1:60 COL I 1:60 vs. FN 1:60	*	0,0101
COL I 1:60 vs. COL IV 1:60	*	0,024
FN 1:60 vs. COL IV 1:60	ns	0,3119
1:35 COL I 1:35 vs. FN 1:35	**	0,0057
COL I 1:35 vs. COL IV 1:35	*	0,0142
FN 1:35 vs. COL IV 1:35	ns	0,1536
1:10 COL I 1:10 vs. FN 1:10	**	0,0019
COL I 1:10 vs. COL IV 1:10	*	0,0138
FN 1:10 vs. COL IV 1:10	*	0,0362

	COL IV			COL I			FN		
	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10
N exp	3	3	3	3	3	3	3	3	3
n cells	85	89	107	138	136	153	126	140	146
Area under the curve	**258,2**	**259,2**	**249,3**	**148,9**	**132,1**	**149,6**	**303**	**318,5**	**340,1**
Std. Error	28,97	27,3	24,41	14,97	15,55	20,01	30,29	29,88	29,15

fig 1E - MC area

descriptive statistics

	COL IV			COL I			FN		
	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10
N exp	3								
n cells	69	90	97	93	88	93	96	103	101
Mean	1727	1775	1950	804,2	1178	1027	2386	2048	2414
SD	993,5	1018	1187	507,7	654,7	736,1	1310	997,7	1356
SEM	119,6	107,3	120,5	52,64	69,79	76,33	133,7	98,3	134,9
Minimum	408,1	235	311,1	162,1	320,4	217,1	431,8	416,6	640,8
25% Percentile	993,3	1019	992,5	446,9	652,3	458	1447	1290	1412
Median	1386	1523	1625	609,8	933,1	707,9	1940	1758	2044
75% Percentile	2396	2336	2607	1017	1659	1448	3061	2586	3136
Maximum	4418	4696	5158	2086	3042	2765	6009	4929	6133
Range	4010	4461	4847	1924	2722	2548	5577	4512	5492

stastical test

		Significant?	P Value
Pass normality test? NO			
Kruskal-Wallis			
Dunn's multiple comparisons test			
COL IV	COL IV 1:60 vs. COL IV 1:35	ns	>0,9999
	COL IV 1:60 vs. COL IV 1:10	ns	0,8765
	COL IV 1:35 vs. COL IV 1:10	ns	>0,9999
COL I	COL I 1:60 vs. COL I 1:35	****	<0,0001
	COL I 1:60 vs. COL I 1:10	ns	0,2148
	COL I 1:35 vs. COL I 1:10	*	0,0234
FN	FN 1:60 vs. FN 1:35	ns	0,3094
	FN 1:60 vs. FN 1:10	ns	>0,9999
	FN 1:35 vs. FN 1:10	ns	0,3022
1:60	COL I 1:60 vs. FN 1:60	****	<0,0001
	COL I 1:60 vs. COL IV 1:60	****	<0,0001
	FN 1:60 vs. COL IV 1:60	*	0,0482
1:35	COL I 1:35 vs. FN 1:35	****	<0,0001
	COL I 1:35 vs. COL IV 1:35	**	0,0033
	FN 1:35 vs. COL IV 1:35	ns	>0,9999
1:10	COL I 1:10 vs. FN 1:10	****	<0,0001
	COL I 1:10 vs. COL IV 1:10	****	<0,0001
	FN 1:10 vs. COL IV 1:10	ns	0,4259

fig 1F - pFAK levels

Pass normality test? NO		
Kruskal-Wallis		
Dunn's multiple comparisons test		
	Significant?	**P Value**
COL IV COL IV 1:60 vs. COL IV 1:35	ns	0,4894
COL IV 1:60 vs. COL IV 1:10	ns	0,0928
COL IV 1:35 vs. COL IV 1:10	ns	>0,9999
COL I COL I 1:60 vs. COL I 1:35	ns	0,3868
COL I 1:60 vs. COL I 1:10	****	<0,0001
COL I 1:35 vs. COL I 1:10	****	<0,0001
FN FN 1:60 vs. FN 1:35	ns	0,282
FN 1:60 vs. FN 1:10	*	0,0359
FN 1:35 vs. FN 1:10	ns	0,8607
1:60 COL I 1:60 vs. FN 1:60	*	0,0256
COL I 1:60 vs. COL IV 1:60	ns	0,0604
FN 1:60 vs. COL IV 1:60	ns	>0,9999
1:35 COL I 1:35 vs. FN 1:35	****	<0,0001
COL I 1:35 vs. COL IV 1:35	*	0,0427
FN 1:35 vs. COL IV 1:35	*	0,0311
1:10 COL I 1:10 vs. FN 1:10	****	<0,0001
COL I 1:10 vs. COL IV 1:10	****	<0,0001
FN 1:10 vs. COL IV 1:10	***	0,0005

	COL IV			COL I			FN		
	1:60	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**
N exp	4	4	4	4	4	5	4	5	3
n cells	92	85	109	100	97	133	82	121	83
Mean	**1,148**	**1,086**	**1,001**	**0,8877**	**0,767**	**0,5429**	**1,208**	**1,348**	**1,553**
SD	0,5978	0,7068	0,5609	0,4956	0,4253	0,2994	0,6544	0,6813	0,9443
SEM	0,06233	0,07666	0,05372	0,04956	0,04319	0,02596	0,07226	0,06193	0,1037
Minimum	**0,3374**	**0,1596**	**0,2573**	**0,1754**	**0,2115**	**0,1354**	**0,3054**	**0,3427**	**0,319**
25% Percentile	0,7303	0,5859	0,5975	0,4996	0,4524	0,32	0,7271	0,8487	0,8731
Median	**0,945**	**0,9032**	**0,802**	**0,7677**	**0,6746**	**0,453**	**1,016**	**1,188**	**1,35**
75% Percentile	1,38	1,315	1,228	1,165	0,9045	0,7177	1,614	1,756	2,12
Maximum	**3,02**	**3,13**	**2,58**	**2,444**	**2,02**	**1,45**	**3,141**	**3,153**	**4,34**
Range	2,683	2,97	2,323	2,268	1,808	1,314	2,836	2,81	4,021

fig 1G - number of FAs

<table>
<tr><td colspan="3">stastical test</td></tr>
<tr><td colspan="3">Pass normality test? NO</td></tr>
<tr><td colspan="3">Kruskal-Wallis</td></tr>
<tr><td colspan="3">Dunn's multiple comparisons test</td></tr>
<tr><td></td><td>Significant?</td><td>P Value</td></tr>
<tr><td rowspan="3">COL IV</td><td>COL IV 1:60 vs. COL IV 1:35</td><td>ns</td><td>>0,9999</td></tr>
</table>

		Significant?	P Value
COL IV	COL IV 1:60 vs. COL IV 1:35	ns	>0,9999
	COL IV 1:60 vs. COL IV 1:10	ns	0,9173
	COL IV 1:35 vs. COL IV 1:10	ns	>0,9999
COL I	COL I 1:60 vs. COL I 1:35	ns	>0,9999
	COL I 1:60 vs. COL I 1:10	ns	>0,9999
	COL I 1:35 vs. COL I 1:10	ns	0,375
FN	FN 1:60 vs. FN 1:35	*	0,0363
	FN 1:60 vs. FN 1:10	****	<0,0001
	FN 1:35 vs. FN 1:10	****	<0,0001
1:60	COL I 1:60 vs. FN 1:60	ns	>0,9999
	COL I 1:60 vs. COL IV 1:60	ns	>0,9999
	FN 1:60 vs. COL IV 1:60	ns	0,0733
1:35	COL I 1:35 vs. FN 1:35	ns	>0,9999
	COL I 1:35 vs. COL IV 1:35	ns	>0,9999
	FN 1:35 vs. COL IV 1:35	ns	>0,9999
1:10	COL I 1:10 vs. FN 1:10	****	<0,0001
	COL I 1:10 vs. COL IV 1:10	*	0,017
	FN 1:10 vs. COL IV 1:10	ns	>0,9999

descriptive statistics

	COL IV			COL I			FN		
	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10
N exp	3	3	3	3	3	5	3	4	3
n cells	69	65	89	84	79	133	69	99	83
Mean	**0,08505**	**0,08922**	**0,09498**	**0,06871**	**0,07679**	**0,06747**	**0,05413**	**0,0731**	**0,1162**
SD	0,05818	0,05545	0,05905	0,04149	0,04721	0,04487	0,03012	0,04481	0,07241
SEM	0,007005	0,006878	0,006259	0,004527	0,005312	0,003891	0,003626	0,004503	0,007948
Minimum	**0,009269**	**0,01108**	**0,01926**	**0,01692**	**0,0106**	**0,005748**	**0,01374**	**0,01228**	**0,006993**
25% Percentile	0,04006	0,03893	0,03926	0,03586	0,03606	0,02871	0,03361	0,03966	0,05707
Median	**0,06392**	**0,07813**	**0,09005**	**0,0572**	**0,0712**	**0,05866**	**0,0471**	**0,05687**	**0,1068**
75% Percentile	0,1148	0,1273	0,1365	0,09822	0,1145	0,09245	0,06381	0,1039	0,1591
Maximum	**0,2405**	**0,2367**	**0,2921**	**0,1659**	**0,1936**	**0,1977**	**0,1478**	**0,1929**	**0,3405**
Range	0,2313	0,2256	0,2728	0,149	0,183	0,1919	0,1341	0,1806	0,3335

fig 2C and 2D - cell migration and velocity

stastical test - total distance migrated

Pass normality test? **NO**		
Kruskal-Wallis		
Dunn's multiple comparisons test		
	Significant?	**P Value**
COL I vs. FN	****	<0,0001
COL I vs. COL IV	****	<0,0001
FN vs. COL IV	*	0,0239

descriptive statistics - total distance migrated

	COL IV	COL I	FN
N exp	3	3	3
n cells	179	158	113
Mean	**214,2**	**91,46**	**267,2**
SD	142,9	57,47	159,5
SEM	10,68	4,572	15,01
Minimum	**3,088**	**3,431**	**21,14**
25% Percentile	104,1	50,29	136,5
Median	**171,5**	**78,16**	**239,5**
75% Percentile	309,7	120,8	375,7
Maximum	**627,6**	**253,2**	**700,8**
Range	624,6	249,7	679,7

stastical test - cell velocity

Pass normality test? **NO**		
Kruskal-Wallis		
Dunn's multiple comparisons test		
	Significant?	**P Value**
COL I vs. FN	****	<0,0001
COL I vs. COL IV	****	<0,0001
FN vs. COL IV	ns	0,0792

descriptive statistics - cell velocity

	COL IV	COL I	FN
N exp	3	3	3
n cells	181	152	114
Mean	**20,64**	**7,06**	**24,52**
SD	11,12	3,804	12,06
SEM	0,8262	0,3085	1,129
Minimum	**0,6176**	**0,4901**	**2,846**
25% Percentile	12,33	4,165	15,23
Median	**19,56**	**6,367**	**23,41**
75% Percentile	27,55	9,048	32,49
Maximum	**57,23**	**18,43**	**61,04**
Range	56,61	17,94	58,19

	fig 3B - melanin production										

Left table — stastical test

		Significant?	P Value
Pass normality test? YES (Shapiro-Wilk test)			
Welch ANOVA tests			
Dunnett's T3 multiple comparisons test			
COL IV	COL IV 1:60 vs. COL IV 1:35	ns	0,9634
	COL IV 1:60 vs. COL IV 1:10	ns	0,6945
	COL IV 1:35 vs. COL IV 1:10	ns	0,9057
COL I	COL I 1:60 vs. COL I 1:35	ns	0,9214
	COL I 1:60 vs. COL I 1:10	ns	0,7689
	COL I 1:35 vs. COL I 1:10	ns	0,9936
FN	FN 1:60 vs. FN 1:35	ns	0,994
	FN 1:60 vs. FN 1:10	ns	0,3219
	FN 1:35 vs. FN 1:10	ns	0,4965
1:60	COL I 1:60 vs. FN 1:60	**	0,0033
	COL I 1:60 vs. COL IV 1:60	*	0,0206
	FN 1:60 vs. COL IV 1:60	ns	0,7957
1:35	COL I 1:35 vs. FN 1:35	*	0,0154
	COL I 1:35 vs. COL IV 1:35	ns	0,0517
	FN 1:35 vs. COL IV 1:35	ns	0,6287
1:10	COL I 1:10 vs. FN 1:10	*	0,0193
	COL I 1:10 vs. COL IV 1:10	ns	0,0515
	FN 1:10 vs. COL IV 1:10	ns	0,8636

Right table — descriptive statistics

	COL IV			COL I			FN		
	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10
N exp	5	5	5	7	7	7	7	7	7
Mean	0,1785	0,1959	0,2216	0,3294	0,3637	0,3796	0,1502	0,1563	0,1952
SD	0,06573	0,06492	0,07062	0,08978	0,1346	0,1231	0,04679	0,05516	0,05592
SEM	0,0294	0,02903	0,03158	0,03393	0,05089	0,04651	0,01768	0,02085	0,02114

stastical test			
Pass normality test? **YES (Shapiro-Wilk test)**			
Welch ANOVA tests			
Dunnett's T3 multiple comparisons test			
		Significant?	P Value
1:60	COL 4 vs. COL 1	*	0,0144
	COL 4 vs. FN	ns	0,0722
	COL 1 vs. FN	**	0,004
1:35	COL 4 vs. COL 1	***	0,0002
	COL 4 vs. FN	**	0,0089
	COL 1 vs. FN	****	<0,0001
1:10	COL 4 vs. COL 1	*	0,0241
	COL 4 vs. FN	ns	0,4891
	COL 1 vs. FN	**	0,0083
COL IV	1:60 COL 4 vs. 1:35 COL 4	ns	0,221
	1:60 COL 4 vs. 1:10 COL 4	ns	0,4313
	1:35 COL 4 vs. 1:10 COL 4	ns	0,9644
COL I	1:60 COL 1 vs. 1:35 COL 1	ns	0,7861
	1:60 COL1 vs. 1:10 COL1	ns	0,9965
	1:35 COL 1 vs. 1:10 COL 1	ns	0,9095
FN	1:60 FN vs. 1:35 FN	ns	0,9733
	1:60 FN vs. 1:10 FN	ns	0,0926
	1:35 FN vs. 1:10 FN	ns	0,0636

fig 3C - *Tyr* expression level

descriptive statistics

	COL IV			COL I			FN		
	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10
N exp	5	5	5	5	4	5	5	5	5
Mean	**0,09176**	**0,1017**	**0,1059**	**0,1579**	**0,1691**	**0,1611**	**0,07816**	**0,07589**	**0,09226**
SD	0,005307	0,009627	0,02039	0,02684	0,01087	0,02762	0,009056	0,009989	0,008451

	fig 3D - cell proliferation BrdU assay								

stastical test

Pass normality test? **YES (Shapiro-Wilk test)**			
Ordinary two-way ANOVA			
Tukey's multiple comparisons test			
		Significant?	**P Value**
COL IV	1:60 vs. 1:35	ns	0,9984
	1:60 vs. 1:10	ns	>0,9999
	1:35 vs. 1:10	ns	>0,9999
COL I	1:60 vs. 1:35	ns	>0,9999
	1:60 vs. 1:10	*	0,0296
	1:35 vs. 1:10	**	0,0063
FN	1:60 vs. 1:35	ns	0,6252
	1:60 vs. 1:10	ns	>0,9999
	1:35 vs. 1:10	ns	0,7993
1:60	COL I 1:60 vs. FN 1:60	ns	0,0965
	COL I 1:60 vs. COL IV 1:60	ns	0,9351
	FN 1:60 vs. COL IV 1:60	ns	0,6423
1:35	COL I 1:35 vs. FN 1:35	**	0,0014
	COL I 1:35 vs. COL IV 1:35	ns	>0,9999
	FN 1:35 vs. COL IV 1:35	**	0,0025
1:10	COL I 1:10 vs. FN 1:10	****	<0,0001
	COL I 1:10 vs. COL IV 1:10	**	0,0019
	FN 1:10 vs. COL IV 1:10	ns	0,2238

descriptive statistics

	COL IV			COL I			FN		
	1:60	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**
N exp	5	5	5	4	5	4	5	4	5
Mean	32,07364	33,42575	32,93229	34,65256	33,83667	42,24035	28,48825	24,85156	27,92726
SD	2,49960	2,85814	1,85663	3,17603	1,43913	6,63614	3,04371	2,65949	1,36758

fig 3E - cell proliferation Ki67 staining			

		COL IV	COL I	FN
Pass normality test? **YES (Shapiro-Wilk test)**	N exp	4	4	4
Welch ANOVA tests	n cells	1461	1444	1026
Dunnett's T3 multiple comparisons test				
	Mean	**44,31**	**46,72**	**31,82**

	Significant?	P Value
COL I vs. FN	ns	0,9366
COL I vs. COL IV	ns	0,1935
FN vs. COL IV	*	0,0128

	COL IV	COL I	FN
SD	6,306	4,091	8,261
SEM	3,641	2,362	4,769

fig 4D - cell division

	Significant?	P Value			COL IV	COL I	FN
Pass normality test? **YES (Shapiro-Wilk test)**				**N exp**	3	3	3
RM one-way ANOVA				**n cells**	258	230	243
Tukey's multiple comparisons test							
	Significant?	**P Value**		**Mean**	**44,31**	**46,72**	**31,82**
COL I vs. FN	ns	0,0913		SD	6,306	4,091	8,261
COL I vs. COL IV	ns	0,7956		SEM	3,641	2,362	4,769
FN vs. COL IV	*	0,0471					

Pass normality test? **NO**		
Kruskal-Wallis **Dunn's multiple comparisons test**		
Nuclear intensity	**Significant?**	**P Value**
COL IV COL IV 1:60 vs. COL IV 1:35	ns	0,5977
COL IV 1:60 vs. COL IV 1:10	ns	>0,9999
COL IV 1:35 vs. COL IV 1:10	ns	0,1299
COL I COL I 1:60 vs. COL I 1:35	ns	0,5303
COL I 1:60 vs. COL I 1:10	*	0,0156
COL I 1:35 vs. COL I 1:10	****	<0,0001
FN FN 1:60 vs. FN 1:35	*	0,0357
FN 1:60 vs. FN 1:10	ns	>0,9999
FN 1:35 vs. FN 1:10	ns	0,453
Total MITF intensity	**Significant?**	**P Value**
1:60 COL I 1:60 vs. FN 1:60	***	0,0002
COL I 1:60 vs. COL IV 1:60	**	0,0057
FN 1:60 vs. COL IV 1:60	ns	>0,9999
1:35 COL I 1:35 vs. FN 1:35	****	<0,0001
COL I 1:35 vs. COL IV 1:35	****	<0,0001
FN 1:35 vs. COL IV 1:35	ns	0,1296
1:10 COL I 1:10 vs. FN 1:10	****	<0,0001
COL I 1:10 vs. COL IV 1:10	ns	>0,9999
FN 1:10 vs. COL IV 1:10	*	0,0232

fig 4B - MITF levels

descriptive statistics

		COL IV			COL I			FN		
		1:60	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**
	N exp	4	3	4	4	4	5	5	5	4
	n cells	126	89	124	101	114	137	136	146	113
total	**Mean**	**0,8737**	**0,8317**	**0,835**	**0,6952**	**0,5717**	**0,7648**	**0,896**	**0,9951**	**1,021**
	SD	0,3796	0,4033	0,3009	0,3156	0,2286	0,3255	0,352	0,4769	0,4014
	SEM	0,03382	0,04275	0,02702	0,0314	0,02141	0,02781	0,03018	0,03947	0,03776
cytoplasm	**Mean**	**0,7016**	**0,6462**	**0,6757**	**0,4499**	**0,3563**	**0,4591**	**0,7373**	**0,8089**	**0,8506**
	SD	0,3389	0,3765	0,2687	0,2411	0,1618	0,2509	0,3263	0,4519	0,3746
	SEM	0,03019	0,03991	0,02413	0,02399	0,01515	0,02159	0,02798	0,0374	0,03524
nucleus	**Mean**	**0,1721**	**0,1854**	**0,1593**	**0,2453**	**0,2154**	**0,317**	**0,1588**	**0,1862**	**0,1701**
	SD	0,08989	0,08335	0,07319	0,1432	0,1094	0,1943	0,07582	0,0916	0,08375
	SEM	0,008008	0,008835	0,006573	0,01425	0,01025	0,01672	0,006501	0,007581	0,007879

Pass normality test? NO		
Kruskal-Wallis		
Dunn's multiple comparisons test		
	Significant?	P Value
COL IV COL IV 1:60 vs. COL IV 1:35	ns	0,5743
COL IV 1:60 vs. COL IV 1:10	ns	>0,9999
COL IV 1:35 vs. COL IV 1:10	ns	0,8231
COL I COL I 1:60 vs. COL I 1:35	ns	>0,9999
COL I 1:60 vs. COL I 1:10	*	0,0305
COL I 1:35 vs. COL I 1:10	**	0,004
FN FN 1:60 vs. FN 1:35	ns	0,0601
FN 1:60 vs. FN 1:10	ns	>0,9999
FN 1:35 vs. FN 1:10	ns	0,5688
1:60 COL I 1:60 vs. FN 1:60	****	<0,0001
COL I 1:60 vs. COL IV 1:60	****	<0,0001
FN 1:60 vs. COL IV 1:60	ns	>0,9999
1:35 COL I 1:35 vs. FN 1:35	****	<0,0001
COL I 1:35 vs. COL IV 1:35	****	<0,0001
FN 1:35 vs. COL IV 1:35	ns	0,8645
1:10 COL I 1:10 vs. FN 1:10	****	<0,0001
COL I 1:10 vs. COL IV 1:10	****	<0,0001
FN 1:10 vs. COL IV 1:10	ns	0,4252

descriptive statistics

	COL IV			COL I			FN		
	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10
N exp	4	3	4	4	4	5	5	5	4
n cells	137	81	135	130	137	161	155	143	124
Mean	**20,11**	**18,36**	**19,79**	**40,39**	**39,28**	**44,73**	**17,17**	**14,9**	**16,17**
SD	9,17	8,121	9,24	11,68	12,89	13,9	7,471	5,907	7,058
SEM	0,7834	0,9024	0,7952	1,024	1,101	1,095	0,6001	0,494	0,6338
Minimum	**6,155**	**6,953**	**4,964**	**17,41**	**16,81**	**15,37**	**5,292**	**5,924**	**4,966**
25% Percentile	11,98	11,53	12,79	31,71	29,73	33,99	11,25	10,8	11,13
Median	**18,39**	**15,84**	**18,64**	**37,92**	**38,23**	**41,94**	**14,89**	**13,7**	**14,69**
75% Percentile	26,67	23,43	24,68	47,31	46,77	55,23	22,54	17,49	20,49
Maximum	**48,45**	**37,98**	**54,73**	**68,05**	**73,36**	**81,54**	**38,99**	**31,72**	**37,69**
Range	42,3	31,03	49,77	50,64	56,55	66,17	33,7	25,8	32,73

fig 4D - melanin production following MITF depletion

Pass normality test? YES (Shapiro-Wilk test)			descriptive statistics					

Welch ANOVA tests Dunnett's T3 multiple comparisons test		
	Significant?	P Value
COL1 - siCTL vs. FN - siCTL	**	0,0062
COL1 - siCTL vs. COL1 - siMITF	*	0,0108
COL1 - siCTL vs. FN - siMITF	*	0,0368
FN - siCTL vs. COL1 - siMITF	ns	0,9827
FN - siCTL vs. FN - siMITF	ns	0,8274
COL1 - siMITF vs. FN - siMITF	ns	0,671

	COL I		FN	
	siCTL	siMITF	siCTL	siMITF
N exp	3	3	3	3
Mean	0,3333	0,2281	0,2339	0,2046
SD	0,01749	0,01535	0,007638	0,0338

fig 4F - nuclear YAP

stastical test

Pass normality test? NO		
Kruskal-Wallis		
Dunn's multiple comparisons test		
	Significant?	P Value
COL IV COL IV 1:60 vs. COL IV 1:35	*	0,0147
COL IV 1:60 vs. COL IV 1:10	ns	0,3972
COL IV 1:35 vs. COL IV 1:10	ns	0,4846
COL I COL I 1:60 vs. COL I 1:35	ns	0,1505
COL I 1:60 vs. COL I 1:10	ns	>0,9999
COL I 1:35 vs. COL I 1:10	*	0,0485
FN FN 1:60 vs. FN 1:35	*	0,0112
FN 1:60 vs. FN 1:10	***	0,0004
FN 1:35 vs. FN 1:10	ns	0,8248
1:60 COL I 1:60 vs. FN 1:60	*	0,0198
COL I 1:60 vs. COL IV 1:60	ns	>0,9999
FN 1:60 vs. COL IV 1:60	ns	>0,9999
1:35 COL I 1:35 vs. FN 1:35	ns	0,6123
COL I 1:35 vs. COL IV 1:35	ns	>0,9999
FN 1:35 vs. COL IV 1:35	ns	>0,9999
1:10 COL I 1:10 vs. FN 1:10	ns	>0,9999
COL I 1:10 vs. COL IV 1:10	ns	>0,9999
FN 1:10 vs. COL IV 1:10	ns	>0,9999

descriptive statistics

	COL IV			COL I			FN		
	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10
N exp	3	3	3	3	3	3	3	3	3
n cells	82	78	91	91	83	92	71	86	73
Mean	0,5423	0,6992	0,6433	0,6432	0,7406	0,6014	0,4559	0,5928	0,6631
SD	0,2521	0,3472	0,3528	0,3424	0,3623	0,2587	0,2884	0,2818	0,3257
SEM	0,02784	0,03932	0,03698	0,03589	0,03977	0,02698	0,03423	0,03038	0,03812
Minimum	0,1667	0,1983	0,1992	0,1532	0,1454	0,2045	0,05028	0,1372	0,1402
25% Percentile	0,3803	0,426	0,3843	0,4171	0,4401	0,4173	0,2454	0,4014	0,4313
Median	0,4783	0,6247	0,5114	0,6154	0,6862	0,5318	0,4376	0,5621	0,6251
75% Percentile	0,6758	0,9298	0,8007	0,8057	0,991	0,7518	0,6727	0,7665	0,8444
Maximum	1,338	1,727	1,561	1,669	1,65	1,273	1,151	1,555	1,699
Range	1,171	1,529	1,362	1,516	1,505	1,068	1,101	1,418	1,559

Statistical test

		Significant?	P Value
Pass normality test? YES (Shapiro-Wilk test)			
Welch ANOVA tests			
Dunnett's T3 multiple comparisons test			
COL IV	1:60 vs. 1:35	ns	0,9977
	1:60 vs. 1:10	ns	0,758
	1:35 vs. 1:10	ns	0,654
COL I	1:60 vs. 1:35	ns	>0,9999
	1:60 vs. 1:10	ns	0,8712
	1:35 vs. 1:10	ns	0,88
1:60	COL I 1:60 vs. FN 1:60	**	0,0066
	COL I 1:60 vs. COL IV 1:60	ns	0,8299
	FN 1:60 vs. COL IV 1:60	ns	0,1007
1:10	COL I 1:10 vs. FN 1:10	*	0,0393
	COL I 1:10 vs. COL IV 1:10	ns	0,2418
	FN 1:10 vs. COL IV 1:10	ns	0,1137

		Significant?	P Value
Pass normality test? NO			
Kruskal-Wallis			
Dunn's multiple comparisons test			
FN	1:60 vs. 1:35	ns	0,3116
	1:60 vs. 1:10	ns	0,0851
	1:35 vs. 1:10	ns	>0,9999
1:35	COL I 1:35 vs. FN 1:35	*	0,0195
	COL I 1:35 vs. COL IV 1:35	ns	>0,9999
	FN 1:35 vs. COL IV 1:35	*	0,0485

descriptive statistics

	COL IV			COL I			FN		
	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10
N exp	4	4	4	5	5	5	5	5	5
Mean	0,0697	0,06543	0,1008	0,05055	0,05032	0,03889	0,1379	0,1863	0,2496
SD	0,04109	0,03388	0,05607	0,03239	0,03279	0,02078	0,03064	0,0443	0,1149

fig 6F - *Tyr* expression following MEK inhibition								

stastical test

stastical test				descriptive statistics						
Pass normality test? **YES (Shapiro-Wilk test)**					**COL IV**		**COL I**		**FN**	
Multiple t tests **Dunn's multiple comparisons test**					DMSO	MEKi	DMSO	MEKi	DMSO	MEKi
		Significant?	Adjusted P Value	**N exp**	3	3	3	3	3	3
COL IV vs. COL I	**DMSO**	*	0,02948	**Mean**	**0,1162**	**0,186**	**0,1964**	**0,2354**	**0,1037**	**0,1622**
	MEKi	*	0,02948	SD	0,01089	0,01735	0,03204	0,01816	0,003171	0,0245
COL IV vs. FN	**DMSO**	ns	0,239076							
	MEKi	ns	0,24224							
COL I vs. FN	**DMSO**	*	0,015087							
	MEKi	*	0,015087							
DMSO vs MEKi	**COL IV**	*	0,012357							
	COL I	ns	0,139977							
	FN	*	0,029362							

fig 6H - MITF nuclear localization following MEK inhibition								
stastical test			descriptive statistics					
Pass normality test? NO				COL IV		COL I		FN	
Kruskal-Wallis				DMSO	MEKi	DMSO	MEKi	DMSO	MEKi
Dunn's multiple comparisons test			N exp	3	3	3	3	3	3
	Significant?	P Value	n cells	75	87	83	86	63	80
COL IV DMSO vs. COL I DMSO	****	<0,0001	Mean	2,99404	4,73727	4,80857	4,12824	2,347	5,04157
COL IV DMSO vs. FN DMSO	ns	0,6551	SD	1,656	2,68814	2,53877	2,27062	1,43225	2,7861
COL I DMSO vs. FN DMSO	****	<0,0001	SEM	0,19122	0,2882	0,27867	0,24485	0,18045	0,3115
COL IV DMSO vs. COL IV MEKi	***	0,0001	Minimum	0,74372	1,24148	1,55216	1,092	0,46891	1,15777
COL I DMSO vs. COL I MEKi	ns	>0,9999	25% Percentile	1,54947	2,39813	3,05737	2,41256	1,28225	2,56296
FN DMSO vs. FN MEKi	****	<0,0001	Median	2,57809	4,20123	3,87857	3,6125	1,90062	4,1922
			75% Percentile	3,93021	6,3675	6,63351	4,99515	2,696	6,89667
			Maximum	6,41366	12,48933	13,12165	10,28927	6,35344	12,25691
			Range	5,66994	11,24785	11,56949	9,19728	5,88453	11,09914

fig 6I - melanin production following MEK inhibition							

stastical test

stastical test		
Pass normality test? **YES (Shapiro-Wilk test)**		
Multiple t tests		
Dunn's multiple comparisons test		
	Significant?	**P Value**
COL IV DMSO vs. COL I DMSO	ns	0,092307
COL IV DMSO vs. FN DMSO	*	0,019629
COL I DMSO vs. FN DMSO	*	0,019109
COL IV DMSO vs. COL IV MEKi	***	0,000252
COL I DMSO vs. COL I MEKi	ns	0,532253
FN DMSO vs. FN MEKi	*	0,027828

	descriptive statistics					
	COL IV		**COL I**		**FN**	
	DMSO	MEKi	DMSO	MEKi	DMSO	MEKi
N exp	3	3	3	3	3	3
Mean	**0,1582**	**0,3522**	**0,2711**	**0,3278**	**0,04884**	**0,3277**
SD	0,008314	0,02605	0,08836	0,1135	0,04957	0,1341

fig 6K - MITF nuclear localization following ERK inhibition

descriptive statistics

	COL IV		COL I		FN	
	DMSO	ERKi	DMSO	ERKi	DMSO	ERKi
N exp	3	3	3	3	3	3
n cells	94	89	97	97	97	92
Mean	3,1148	5,3699	4,4835	6,2824	3,01119	4,85466
SD	1,73341	3,02697	2,1617	3,60689	1,5851	2,23513
SEM	0,17879	0,32086	0,21949	0,36622	0,16094	0,23303
Minimum	0,75315	1,31929	0,88298	1,55145	0,93948	1,24764
25% Percentile	1,63225	2,81743	2,85909	3,6587	1,78728	3,16784
Median	2,82252	4,56534	4,30881	5,09271	2,58744	4,4764
75% Percentile	4,26161	7,52305	5,66406	7,9799	4,04162	5,91185
Maximum	9,09394	13,51866	11,29825	15,82687	7,84094	11,48497
Range	8,34079	12,19937	10,41527	14,27541	6,90146	10,23732

stastical test		
Pass normality test? NO		
Kruskal-Wallis		
Dunn's multiple comparisons test		
	Significant?	P Value
COL IV DMSO vs. COL I DMSO	****	<0,0001
COL IV DMSO vs. FN DMSO	ns	>0,9999
COL I DMSO vs. FN DMSO	****	<0,0001
COL IV DMSO vs. COL IV ERKi	****	<0,0001
COL I DMSO vs. COL I ERKi	*	0,0334
FN DMSO vs. FN ERKi	****	<0,0001
COL1 DMSO vs. COL4 ERKi	ns	>0,9999
COL1 DMSO vs. FN ERKi	ns	>0,9999

fig 6I - melanin production following ERK inhibition

stastical test				descriptive statistics						
Pass normality test? **YES (Shapiro-Wilk test)**					**COL IV**		**COL I**		**FN**	
Welch ANOVA tests					DMSO	ERKi	DMSO	ERKi	DMSO	ERKi
Dunnett's T3 multiple comparisons test				N exp	4	4	4	4	4	4
	Significant?	**P Value**		Mean	**0,1118**	**0,2185**	**0,1959**	**0,2336**	**0,06181**	**0,2115**
COL4 DMSO vs. COL1 DMSO	ns	0,0693		SD	0,0434	0,03022	0,03761	0,02197	0,02676	0,0268
COL4 DMSO vs. FN DMSO	ns	0,2556								
COL1 DMSO vs. FN DMSO	**	0,0057								
COL4 ERKi vs. COL4 DMSO	*	0,0186								
COL4 ERKi vs. COL1 DMSO	ns	0,6035								
FN ERKi vs. COL1 DMSO	ns	0,7619								
FN ERKi vs. FN DMSO	***	**0,0004**								

				fig 6N - pERK levels following FAK inhibition			
stastical test				descriptive statistics			
Pass normality test? **NO**					COL IV		COL I		FN	
Kruskal-Wallis					DMSO	FAKi	DMSO	FAKi	DMSO	FAKi
Dunn's multiple comparisons test										
	Significant?	**P Value**		**N exp**	3	3	3	3	3	3
				n cells	88	93	83	95	91	96
COL IV DMSO vs. COL I DMSO	*	0,0483		**Mean**	**0,9769**	**0,6028**	**0,6439**	**0,2983**	**1,495**	**0,8919**
COL IV DMSO vs. FN DMSO	***	0,0008		SD	0,6168	0,3233	0,3451	0,1479	0,8128	0,4077
COL I DMSO vs. FN DMSO	****	<0,0001		SEM	0,06575	0,03352	0,03788	0,01517	0,08521	0,04161
COL IV DMSO vs. COL IV FAKi	**	0,0021		**Minimum**	**0,181**	**0,1852**	**0,2348**	**0,09372**	**0,2913**	**0,2552**
COL I DMSO vs. COL I FAKi	****	<0,0001		25% Percentile	0,5427	0,3677	0,3974	0,1878	0,8962	0,5872
FN DMSO vs. FN FAKi	***	0,0004		**Median**	**0,8007**	**0,549**	**0,5299**	**0,2638**	**1,272**	**0,7708**
				75% Percentile	1,362	0,7253	0,7643	0,3867	1,991	1,079
				Maximum	**2,791**	**1,526**	**1,607**	**0,6314**	**4,009**	**2,061**
				Range	2,61	1,341	1,372	0,5377	3,718	1,806

fig 6O - melanin production following FAK inhibition								
stastical test				descriptive statistics				
Pass normality test? **YES (Shapiro-Wilk test)**				**COL IV**		**COL I**		**FN**	
Multiple t tests				DMSO	FAKi	DMSO	FAKi	DMSO	FAKi
Dunn's multiple comparisons test			**N exp**	5	5	5	5	5	5
	Significant?	**P Value**	**Mean**	**0,1601**	**0,2118**	**0,1925**	**0,2608**	**0,105**	**0,203**
COL IV DMSO vs. COL I DMSO		0,258394	SD	0,04429	0,009109	0,0398	0,04417	0,03397	0,04146
COL IV DMSO vs. FN DMSO		0,058323							
COL I DMSO vs. FN DMSO	*	0,005712							
COL IV DMSO vs. COL IV FAKi	*	0,033822							
COL I DMSO vs. COL I FAKi	*	0,033196							
FN DMSO vs. FN FAKi	**	0,003493							

fig S1 - PDMS's Young's modulus

descriptive statistics

	uncoated			COL IV			COL I			FN		
	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10
N exp	3	5	3	3	3	3	3	4	3	3	3	3
n cells	47,91	151	964,2	52,78	161,2	748	44,14	114,8	815	39,3	132	896,6
SD	3,725	28,24	86,42	14,89	11,75	144,8	12,37	22,4	188,6	6,823	25,75	33,65

fig S3A - pMC area

stastical test

		Significant?	P Value
Pass normality test? **NO**			
Kruskal-Wallis			
Dunn's multiple comparisons test			
COL IV	COL IV 1:60 vs. COL IV 1:35	ns	0,5372
	COL IV 1:60 vs. COL IV 1:10	ns	>0,9999
	COL IV 1:35 vs. COL IV 1:10	ns	0,1807
COL I	COL I 1:60 vs. COL I 1:35	ns	0,1552
	COL I 1:60 vs. COL I 1:10	ns	0,7629
	COL I 1:35 vs. COL I 1:10	ns	>0,9999
FN	FN 1:60 vs. FN 1:35	ns	0,4268
	FN 1:60 vs. FN 1:10	ns	>0,9999
	FN 1:35 vs. FN 1:10	ns	0,0972
1:60	COL I 1:60 vs. FN 1:60	****	<0,0001
	COL I 1:60 vs. COL IV 1:60	****	<0,0001
	FN 1:60 vs. COL IV 1:60	****	<0,0001
1:35	COL I 1:35 vs. FN 1:35	****	<0,0001
	COL I 1:35 vs. COL IV 1:35	****	<0,0001
	FN 1:35 vs. COL IV 1:35	****	<0,0001
1:10	COL I 1:10 vs. FN 1:10	****	<0,0001
	COL I 1:10 vs. COL IV 1:10	****	<0,0001
	FN 1:10 vs. COL IV 1:10	****	<0,0001

descriptive statistics

	COL IV			COL I			FN		
	1:60	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**
N exp	5	6	6	5	6	6	6	6	6
n cells	152	126	166	96	142	161	169	142	188
Mean	905,5	783,5	935,4	422	364,9	389,6	1449	1363	1429
SD	466,5	382,8	499,4	189,2	135	158,5	574,2	578,8	474,1
SEM	37,84	34,11	38,76	19,31	11,33	12,49	44,17	48,57	34,57
Minimum	169,6	231,4	221	139,9	170,8	152,8	444,8	71,08	407,9
25% Percentile	522,6	430,3	529,7	297,6	262,3	263,3	1061	921,6	1042
Median	808,8	702,9	849	366,9	329,5	366,5	1349	1324	1364
75% Percentile	1180	1056	1210	513,9	443,3	492,6	1802	1641	1713
Maximum	2222	2043	2564	960,7	783,2	893,3	3364	3145	2892
Range	2053	1811	2343	820,8	612,4	740,5	2919	3074	2484

fig S3B - pMCs pFAK

stastical test

		Significant?	P Value
Pass normality test? NO			
Kruskal-Wallis			
Dunn's multiple comparisons test			
COL IV	COL IV 1:60 vs. COL IV 1:35	***	0,0004
	COL IV 1:60 vs. COL IV 1:10	ns	>0,9999
	COL IV 1:35 vs. COL IV 1:10	***	0,0005
COL I	COL I 1:60 vs. COL I 1:35	**	0,002
	COL I 1:60 vs. COL I 1:10	ns	>0,9999
	COL I 1:35 vs. COL I 1:10	**	0,0024
FN	FN 1:60 vs. FN 1:35	ns	0,3084
	FN 1:60 vs. FN 1:10	**	0,0035
	FN 1:35 vs. FN 1:10	ns	0,7979
1:60	COL I 1:60 vs. FN 1:60	****	<0,0001
	COL I 1:60 vs. COL IV 1:60	****	<0,0001
	FN 1:60 vs. COL IV 1:60	ns	0,1
1:35	COL I 1:35 vs. FN 1:35	****	<0,0001
	COL I 1:35 vs. COL IV 1:35	***	0,0003
	FN 1:35 vs. COL IV 1:35	****	<0,0001
1:10	COL I 1:10 vs. FN 1:10	****	<0,0001
	COL I 1:10 vs. COL IV 1:10	****	<0,0001
	FN 1:10 vs. COL IV 1:10	****	<0,0001

descriptive statistics

	COL IV			COL I			FN		
	1:60	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**
N exp	3	3	3	3	3	3	3	3	3
n cells	83	56	94	61	60	94	88	52	99
Mean	**53960627**	**37850823**	**53121944**	**26372614**	**20865783**	**25719541**	**68881323**	**75897376**	**83136623**
SD	25959332	18026407	25987786	9547453	7649541	9749189	27956374	24175255	29375298
SEM	2849407	2408880	2680435	1222426	987552	1005552	2980159	3352505	2952329
Minimum	**16159373**	**14867328**	**15530967**	**12048802**	**10645512**	**11290650**	**24167920**	**28310931**	**28420452**
25% Percentile	31375362	23863710	35167508	18606713	15836489	17918260	43958642	56234925	62293546
Median	**48248282**	**32846970**	**49118005**	**24166665**	**19697711**	**24944454**	**64671823**	**74186191**	**78139052**
75% Percentile	72133456	52504500	68597073	30522200	24534243	31539422	87383057	98534832	97487149
Maximum	**142920545**	**84526895**	**132503709**	**50515757**	**42573146**	**54848768**	**154650751**	**133684495**	**167690367**
Range	126761172	69659567	116972742	38466955	31927634	43558118	130482831	105373564	139269915

fig S3C - pMCs number of focal adhesions

stastical test

Pass normality test? NO		
Kruskal-Wallis		
Dunn's multiple comparisons test		
	Significant?	**P Value**
COL IV		
COL IV 1:60 vs. COL IV 1:35	**	0,0081
COL IV 1:60 vs. COL IV 1:10	ns	0,507
COL IV 1:35 vs. COL IV 1:10	****	<0,0001
COL I		
COL I 1:60 vs. COL I 1:35	ns	0,7166
COL I 1:60 vs. COL I 1:10	ns	>0,9999
COL I 1:35 vs. COL I 1:10	ns	0,1991
FN		
FN 1:60 vs. FN 1:35	*	0,0117
FN 1:60 vs. FN 1:10	**	0,0071
FN 1:35 vs. FN 1:10	ns	>0,9999
1:60		
COL I 1:60 vs. FN 1:60	ns	>0,9999
COL I 1:60 vs. COL IV 1:60	ns	>0,9999
FN 1:60 vs. COL IV 1:60	ns	>0,9999
1:35		
COL I 1:35 vs. FN 1:35	*	0,0167
COL I 1:35 vs. COL IV 1:35	ns	0,5377
FN 1:35 vs. COL IV 1:35	****	<0,0001
1:10		
COL I 1:10 vs. FN 1:10	ns	>0,9999
COL I 1:10 vs. COL IV 1:10	ns	>0,9999
FN 1:10 vs. COL IV 1:10	ns	>0,9999

descriptive statistics

	COL IV			COL I			FN		
	1:60	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**	**1:60**	**1:35**	**1:10**
N exp	3	3	3	3	3	3	3	3	3
n cells	82	70	94	65	69	90	88	70	97
Mean	0,03502	0,02651	0,04527	0,03746	0,03459	0,04262	0,03773	0,04632	0,0479
SD	0,01668	0,01922	0,03103	0,02026	0,02325	0,02915	0,01517	0,01932	0,02315
SEM	0,001842	0,002298	0,0032	0,002514	0,002799	0,003073	0,001617	0,002309	0,002351
Minimum	0,004765	0,002161	0,002673	0,007635	0	0	0,01128	0,008768	0,01145
25% Percentile	0,02149	0,009788	0,02045	0,02308	0,01966	0,02116	0,0265	0,03449	0,03123
Median	0,03417	0,0236	0,03601	0,03389	0,02637	0,03505	0,03531	0,04149	0,041
75% Percentile	0,04701	0,03708	0,06262	0,04974	0,04412	0,05678	0,04717	0,05816	0,06374
Maximum	0,08189	0,07847	0,1341	0,09445	0,09355	0,1294	0,09034	0,1014	0,1084
Range	0,07712	0,07631	0,1314	0,08681	0,09355	0,1294	0,07907	0,0926	0,09695

fig S3E - pMCs MITF nuclear localization

stastical test			
Pass normality test? NO			
Kruskal-Wallis			
Dunn's multiple comparisons test			
		Significant?	P Value
COL IV	COL IV 1:60 vs. COL IV 1:35	ns	0,2081
	COL IV 1:60 vs. COL IV 1:10	ns	>0,9999
	COL IV 1:35 vs. COL IV 1:10	ns	0,3965
COL I	COL I 1:60 vs. COL I 1:35	*	0,0186
	COL I 1:60 vs. COL I 1:10	**	0,0034
	COL I 1:35 vs. COL I 1:10	ns	>0,9999
FN	FN 1:60 vs. FN 1:35	ns	0,7181
	FN 1:60 vs. FN 1:10	ns	>0,9999
	FN 1:35 vs. FN 1:10	ns	0,2589
1:60	COL I 1:60 vs. FN 1:60	****	<0,0001
	COL I 1:60 vs. COL IV 1:60	**	0,0056
	FN 1:60 vs. COL IV 1:60	**	0,0013
1:35	COL I 1:35 vs. FN 1:35	****	<0,0001
	COL I 1:35 vs. COL IV 1:35	**	0,0046
	FN 1:35 vs. COL IV 1:35	****	<0,0001
1:10	COL I 1:10 vs. FN 1:10	****	<0,0001
	COL I 1:10 vs. COL IV 1:10	****	<0,0001
	FN 1:10 vs. COL IV 1:10	***	0,0009

descriptive statistics

	COL IV			COL I			FN		
	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10
N exp	3	3	3	3	3	3	3	3	3
n cells	86	70	96	67	74	91	88	71	99
Mean	29,68	34,32	30,3	38,36	43,09	43,84	21,27	19,69	22,29
SD	12,05	15,48	11,44	11,58	11,34	10,83	7,844	7,228	9,502
SEM	1,299	1,85	1,168	1,415	1,318	1,135	0,8361	0,8578	0,955
Minimum	8,542	8,892	9,374	11,74	10,65	8,77	7,304	8,562	6,159
25% Percentile	19,53	19,53	21,04	31,47	38,51	38,46	14,77	14,37	14,59
Median	29,08	34,24	28,87	39,91	45,31	45,85	20,37	18,57	20,52
75% Percentile	38,82	47,69	38,33	45,05	50,73	51,04	27,42	24,11	30,15
Maximum	61,13	61,1	56,35	65,56	65,54	65,54	41,3	40,22	44,81
Range	52,58	52,21	46,98	53,81	54,89	56,77	34	31,66	38,65

fig S4 - ratio cell area to % nuclear MITF

descriptive statistics

	COL IV			COL I			FN		
	1:60	1:35	1:10	1:60	1:35	1:10	1:60	1:35	1:10
N exp	3	3	3	3	3	3	3	3	3
n cells	67	47	93	81	67	83	78	82	75
Mean	61,63	62,14	62,59	21,29	16,54	12,92	136,2	132,9	75,83
SD	46,88	51,37	61,61	14,85	9,843	8,021	100,1	91,7	49,73
SEM	5,727	7,493	6,389	1,65	1,203	0,8804	11,34	10,13	5,742

fig S6B - pFAK levels following FAK inhibition

	Significant?	P Value
Pass normality test? **NO**		
Kruskal-Wallis		
Dunn's multiple comparisons test		
	Significant?	**P Value**
COL4 DMSO vs. COL1 DMSO	****	<0,0001
COL4 DMSO vs. FN DMSO	ns	>0,9999
COL1 DMSO vs. FN DMSO	****	<0,0001
COL4 DMSO vs. COL4 FAKi 24h	****	<0,0001
COL1 DMSO vs. COL1 FAKi 24h	****	<0,0001
FN DMSO vs. FN FAKi 24h	****	<0,0001

descriptive statistics

	COL IV		COL I		FN	
	DMSO	FAKi	DMSO	FAKi	DMSO	FAKi
N exp	3	3	3	3	3	3
n cells	91	89	89	92	89	88
Mean	**1,422**	**0,4279**	**0,8189**	**0,3294**	**1,894**	**0,5285**
SD	0,6885	0,2765	0,3942	0,1955	1,046	0,3096
SEM	0,07217	0,0293	0,04179	0,02038	0,1109	0,033
Minimum	**0,4204**	**0,0598**	**0,21**	**0,07309**	**0,4163**	**0,123**
25% Percentile	0,9394	0,2165	0,5411	0,162	1,041	0,2775
Median	**1,239**	**0,3447**	**0,6963**	**0,322**	**1,705**	**0,4613**
75% Percentile	1,797	0,5666	1,066	0,4396	2,607	0,712
Maximum	**3,494**	**1,087**	**1,902**	**0,9362**	**4,908**	**1,327**
Range	3,074	1,027	1,692	0,8631	4,491	1,204

fig S6C - FA number following FAK inhibition

Pass normality test? NO		
Kruskal-Wallis		
Dunn's multiple comparisons test		
	Significant?	**P Value**
COL4 DMSO vs. COL1 DMSO	****	<0,0001
COL4 DMSO vs. FN DMSO	ns	0,6525
COL1 DMSO vs. FN DMSO	****	<0,0001
COL4 DMSO vs. COL4 FAKi	****	<0,0001
COL1 DMSO vs. COL1 FAKi	****	<0,0001
FN DMSO vs. FN FAKi	****	<0,0001

descriptive statistics

	COL IV		COL I		FN	
	DMSO	FAKi	DMSO	FAKi	DMSO	FAKi
N exp	3	3	3	3	3	3
n cells	76	77	76	70	72	75
Mean	**119,2**	**38,78**	**39,96**	**12,67**	**163,1**	**51,13**
SD	66,71	25,91	25,97	8,937	88,47	29,88
SEM	7,652	2,953	2,978	1,068	10,43	3,45
Minimum	**21**	**2**	**5**	**1**	**64**	**9**
25% Percentile	67	16	20	6	88,5	28
Median	**108**	**32**	**33,5**	**11**	**136**	**43**
75% Percentile	157,5	57	54,25	18	204,5	70
Maximum	**314**	**97**	**119**	**38**	**373**	**140**
Range	293	95	114	37	309	131