A genotype-first approach identifies variants for orofacial clefts and other phenotypes in

dogs

Reuben M Buckley[1], Nüket Bilgen[2], Alexander C Harris[1], Peter Savolainen[3], Cafer Tepeli[4], Metin

Erdoğan[5], Aitor Serres Armero[1], Dayna L Dreger[1], Frank G van Steenbeek[6], Marjo K

Hytönen[7,8,9], Jessica Hale[1], Hannes Lohi[7,8,9], Bengi Çınar Kul[2], Adam R Boyko[10,11], Elaine A

Ostrander[1]*

[1]National Human Genome Research Institute, National Institutes of Health, Bethesda, MD

20892, USA.

[2]Department of Animal Genetics, Faculty of Veterinary Medicine, University of Ankara, Ankara

06110, Türkiye

[3]KTH Royal Institute of Technology, School of Chemistry, Biotechnology and Health, Science for

Life Laboratory, Stockholm, Sweden

[4]Department of Animal Science, University of Selcuk, Faculty of Veterinary Medicine, Konya,

Türkiye

[5]Department of Veterinary Biology and Genetics, Faculty of Veterinary Medicine, Afyon

Kocatepe University, Afyonkarahisar, Türkiye

[6]Utrecht University, Faculty of Veterinary Medicine, Dept. of Clinical Sciences, The Netherlands

[7]Department of Medical and Clinical Genetics, University of Helsinki, 00014 Helsinki, Finland

[8]Department of Veterinary Biosciences, University of Helsinki, 00014 Helsinki, Finland

[9]Folkhälsan Research Center, 00290 Helsinki, Finland

[10]Department of Biomedical Sciences, College of Veterinary Medicine, Cornell University, Ithaca,

NY 14853, USA

[11]Embark Veterinary, Inc., Boston, MA 02210, USA

*Corresponding author

Author info:

Reuben M Buckley (RMB) (0000-0001-6334-3906) - reuben.buckley@nih.gov

Nüket Bilgen (NB) (0000-0003-2324-7965) - nbilgen@ankara.edu.tr

Alexander C Harris (ACH) (0000-0002-6868-5363)- alex.harris@nih.gov

Peter Savolainen (PS) (0000-0002-1495-8338) - savo@kth.se

Cafer Tepeli (CT) (0000-0003-2966-8330) - ctepeli@selcuk.edu.tr

Metin Erdoğan (ME) (0000-0003-0975-1989) - erdogan@aku.edu.tr

Aitor Serres Armero (ASA) (0000-0003-1679-811X) - aitor.serresarmero@nih.gov

Dayna L Dreger (DLD) (0000-0002-3218-6431) - dayna.dreger@nih.gov

Frank G van Steenbeek (FGVS) (0000-0001-5460-6540) - F.G.vanSteenbeek@uu.nl

Marjo K Hytönen (MKH) (0000-0003-1976-5874) - marjo.hytonen@helsinki.fi

Jessica Hale (JH) (0009-0008-4953-9556) - jessica.hale@nih.gov

Hannes Lohi (HL) (0000-0003-1087-5532) - hannes.lohi@helsinki.fi

Bengi Çınar Kul (BCK) (0000-0002-8955-0097) - bengikul@gmail.com

Adam R Boyko (ARB) (0000-0002-9125-6874) - boyko@cornell.edu

Elaine A Ostrander (EAO) (0000-0001-6075-9738) - eostrand@mail.nih.gov

Abstract

Dog breeding promotes within-group homogeneity through conformation to strict breed

standards, and also drives between-group heterogeneity in pursuit of characteristic breed traits.

There are over 350 recognized dog breeds that provide the foundation for investigating the

genetic basis of phenotypic diversity. Typically, breed standard phenotypes such as stature, fur

length, and craniofacial structure are analyzed in genetic association studies. However, such

analyses are limited to the assayed phenotypes, leaving difficult to measure phenotypic

subtleties potentially overlooked. In this study, the genotype-first approach was adapted to the

dog genome to investigate coding variation from over 2000 dogs, leading to discoveries of new

mutations related to craniofacial morphology and stature. Breed-enriched variants were

prioritized according to gene constraint, which was calculated using a mutation model derived

from trinucleotide substitution probabilities in the dog. Among the discovered variants was a

splice-acceptor mutation in *PDGFRA* associated with bifid nose, a characteristic trait of

Çatalburun dogs, implicating the gene's role in midline closure, and a frameshift mutation in

LCORL associated with large canine body size, thus highlighting the importance of allelic

heterogeneity in selection for breed traits. Most priority variants were not associated with

genomic signatures for breed differentiation, as these regions were enriched for constrained

genes intolerant to nonsynonymous variation, suggesting a model of breed phenotype

diversification based on regulatory changes to essential genes. Identification of trait-associated

variants in dogs informs new biological roles for genes. Improved collection of breed disease

risk data, along with increased breed representation, will drive further discoveries.

Introduction

The genomes of domestic species hold immense promise for the discovery of novel

genotype-phenotype associations (Andersson and Purugganan 2022; Buckley and Ostrander

2024). The domestic dog comprises over 350 breeds (Rogers and Brace 1995; Fogle 2000),

many of which exhibit extreme levels of phenotypic diversity, including those associated with

morphology, behavior, and disease susceptibility (Ostrander et al. 2017; Leeb et al. 2023).

Identifying genetic variants that drive canine phenotypic variation will further define gene

functions. Genetic association studies typically require an initial and robust characterization of a

phenotype, a practice that ultimately limits the scope of potential findings to only those that can

be recognized and sufficiently defined.

Alternatively, genetic associations can be identified using a "genotype-first" approach,

where the first step is to identify candidate genotypes with potential phenotypic impacts

(Stessman et al. 2014; Wilczewski et al. 2023). Based on the impacted genes, including

predicted functions and characteristics of individuals with candidate variants, such as breed

disease prevalence, researchers can generate new hypotheses for potential genotype-

phenotype associations. In turn, this spurs additional collection of comprehensive phenotypes

relevant to the trait of interest and allows for the hypothesized associations to be tested.

The genotype-first approach was initially formulated to investigate the genetic basis of

phenotypic variation associated with complex human disorders which are frequently polygenic,

non-coding, incompletely penetrant, and influenced by environmental factors (Stessman et al.

2014; Wilczewski et al. 2023). Complex traits often vary widely in presentation, and the

presence of disease subtypes can require information regarding family history, disease

presentation, treatment response, and histology. Genotype-first approaches, however, focus on

classifying disease subtypes according to profiles of causative/pathogenic genetic variants. This

alternate structure of disease classification can then guide comprehensive clinical phenotyping,

making it possible to determine how specific alleles contribute to disease variability (Stessman

et al. 2014; Leitsalu et al. 2021; Wenger et al. 2021; Park et al. 2022; Garcia-Pelaez et al. 2023;

Wilczewski et al. 2023).

The adoption of the genotype-first approach in complex disease research has been

largely driven by extensive genomic data collection in case cohorts, availability of electronic

health records, and ability to recontact patients, supporting the identification of

causative/pathogenic variants among cases and facilitating the collection of comprehensive

phenotypes (Lyon et al. 2023; Safonov et al. 2023; Wilczewski et al. 2023; Wright et al. 2024). In

canine genomics, there is far less availability of genomic and phenotypic resources. However, a

rigid population structure, paired with extreme phenotypic diversity across breeds means that

genotype-first approaches, when applied to dogs, can be used to ask different kinds of

questions, specifically addressing the roles of specific genetic variants in shaping not only

disease risk, but the striking differences in body morphology and behavior that characterize

different breeds. By definition, dog breeds are closed populations, where members of a breed

are only crossed with those of the same breed. Thus, every registered dog is descended from a

small number of founders and shares extensive genetic homogeneity (Mellanby et al. 2013;

Lampi et al. 2020). Breed members have multigeneration pedigrees and are judged against

morphologic standards that further enforce genetic homogeneity, resulting in a reduction in the

locus heterogeneity that typifies human traits, where dozens of genes may be foundational for a

given complex phenotype. The ease of association studies in dogs is further enhanced by the

fact that most breeds were developed within the last 200 years, such that there are minimal

numbers of generations for meiotic recombination to occur (Worboys 2018). This unique

population structure means that breed membership can be used as a surrogate when mapping

breed-associated morphologic traits, thus avoiding the time and expense of collecting individual

metrics (Gough et al. 2018; Plassais et al. 2019; Dutrow et al. 2022; Morrill et al. 2022).

In humans, candidate trait and disease variants are generally identified using a

combination of strategies including multiple species alignment, protein structure predictions,

gene mutation tolerance, and machine learning (Ng and Henikoff 2001; Adzhubei et al. 2010; Havrilla et al. 2019; Rentzsch et al. 2019; Karczewski et al. 2020; Cheng et al. 2023; Gao et al. 2023; Sullivan et al. 2023). Most such tools benefit from extensive catalogues of human variation featuring large amounts of clinical and personal data (Landrum et al. 2014; Lek et al. 2016; Bycroft et al. 2018; All of Us Research Program Genomics 2024). Although efforts to catalogue trait-associated genetic variation exist for domestic mammals (Nicholas et al. 1995; Plassais et al. 2019; Nicholas 2021; Meadows et al. 2023), the resources for extensive databases are limiting. While mapping functional gene/genomic annotations over from humans can theoretically be used for improving variant annotation, it is estimated that ~13% of human pathogenic variants are likely benign in dogs (Gao et al. 2023). Systems such as the dog may, therefore, be rife with lost opportunities for identifying genes and variants that play fundamental roles in mammalian biology. Nevertheless, the recent Dog10K release of whole genome sequence (WGS) data from over 2000 dogs (Ostrander et al. 2019; Meadows et al. 2023), inclusive of 321 breeds and populations, means the time is ripe to use genotype-first methods, in combination with measures of functional and evolutionary constraint, to identify sequence level variants with a high probability of impacting gene function un the dog

This study focuses, specifically, on identifying genotypes that are rare, overall, in dogs but common within a single breed or groups of related breeds that share a particular trait. If the genotype highlights a variant of potential impact, it is selected for additional study. We characterize coding variation across 2,291 canine genomes from over 200 phylogenetically informed breeds/subpopulations. We use a mutation model to investigate the accuracy of three different gene/site constraint metrics for predicting variant phenotypic impacts. We then combine constraint metrics with variant data to apply the genotype-first approaches to canines, identifying several candidate trait-associated variants. Finally, we investigate the contribution of coding variation to breed differentiation, identifying the importance of constrained genes for shaping phenotypic differences between breeds.

Results

Canine genetic population structure and coding sequence constraint

To represent the breadth of canine coding variation, we combined the Dog10K cohort with a

panel of dogs compiled from WGS data from the sequence read archive (SRA). After extensive

sample and variant filtering, the combined dataset retained 821,202 coding sequence (CDS)

sites with derived alleles from across 2,291 samples, including both splice donor and acceptor

sites (**Fig. 1A**) (**Supplemental Table S1**) (**Methods**). Using the final dataset of 2,018 breed

dogs, 219 village dogs, 53 wolves, and one coyote we further refined the dataset to 611,644

single nucleotide variants (SNVs), 28,970 deletions, and 28,123 insertions. Approximately 40%

of all CDS SNVs occur only once in the dataset, indicating a vast landscape of uncharacterized

rare variation within canids (**Fig. 1B**). We also identified 16,974 multiallelic sites, which

consisted of 12,677 SNVs, representing 2.1% of all SNVs within the entire dataset, thus

highlighting the extent of data loss from filtering for biallelic variants only, which can be expected

to increase as datasets continue to grow (**Fig. 1C**). Dog breeds were also characterized as

monophyletic clusters using a phylogenetic clustering approach (**Methods**) (**Supplemental**

Data S1). A total of 200 breeds/groups with five or more WGS samples were identified

(**Supplemental Table S2**).

1
Figure 1: Coding variation and population genetics of 2,291 canine samples. A) Variant
data from SRA and Dog10K were filtered and combined to create a single dataset used
throughout the analysis. B) Allele count distribution of mutation types. C) Multiallelic variants
occupying the same sites. D) The allele frequency distributions of SNVs according to mutation
type and constraint level for three different constraint metrics. Constraint is expressed as a
decile, where the highest deciles represent the most constrained genes/sites.
8

Variants were annotated according to three different constraint metrics: constrained

CDS, human ortholog LOEUF (loss of function observed over expected upper fraction), and

variant phyloP score. Constrained CDS and variant phyloP score both represent evolutionary

conservation derived from the Zoonomia consortium's alignment of 240 mammals. Variant

phyloP score is the level of conservation of an individual nucleotide position. Constrained CDS

is the fraction of nucleotides within a gene's CDS that have a phyloP score >2.27 (genome-wide

FDR > 0.05) (Christmas et al. 2023; Sullivan et al. 2023). Human ortholog LOEUF indicates

whether a human-dog ortholog is depleted of predicted loss of function (LoF) variation in

humans (Karczewski et al. 2020), where LoF here specifically represents nonsense, splice

disruptor, and frameshift variant annotations (MacArthur et al. 2012). The values used are from

GNOMAD v4.1.0 and are based on expected rates of LoF variation from over 800,000 humans

(Chen et al. 2024). The key differences between the constraint metrics are 1) whether they

reflect evolutionary constraint (constrained CDS and variant phyloP score) or functional

variation constraint (human ortholog LOEUF), and 2) whether the metric represents an entire

gene (constrained CDS and human ortholog LOEUF) or a specific base pair (variant phyloP

score). These differences are important, as evolutionary-based metrics, which depended on

across species sequence conservation, are blind to DNA sequences with essential lineage-

specific biological roles, and gene-based metrics have too low resolution for evaluating selection

effects on gene regions or protein domains.

To characterize each constraint metric in dogs, we analyzed the distribution of SNV

mutation types and allele counts according to constraint level (**Fig. 1D**). As expected, genes

under the greatest constraint, according to constrained CDS and human ortholog LOEUF, had

the lowest fraction of LoF variants. Conversely, LoF variants were most common among sites

with high variant phyloP scores, the opposite of what was observed for gene-based metrics.

This outcome is caused by codon position effects on phyloP scores. Positions susceptible to

LoF substitutions are going to be the most conserved across species, indicating that LoF

variants are going to occur more frequently at these positions (Pollard et al. 2010; Sullivan et al.

2023). The inverse of these effects is observed with synonymous variants, which are most

abundant in highly constrained genes and almost completely absent from highly constrained

sites, due to their often-negligible impact on fitness.

In dogs, allele count distributions, a proxy for allele frequencies, also varied according to

constraint level. LoF SNVs in constrained genes/sites were usually rare, while common LoF

SNVs (AC >100) were more frequently found within genes/sites not under strong constraint. Although low allele frequency and density of functional variants is an indicator of negative selection acting on deleterious mutations, the observed difference in SNV density across constraint metrics suggests that the utility of each metric for variant prioritization may vary significantly.

A framework for evaluating gene sequence constraint metrics

Genomic sequence constraint metrics are often used to predict whether specific mutations are likely to have deleterious impacts. Characteristic breed traits in dogs are often phenotypic extremes and can be linked to deleterious health outcomes, such as in brachycephalic breeds (Marchant et al. 2019; O'Neill et al. 2020). Sequence constraint, therefore, has potential for identifying trait-causing variants in dogs. However, the constraint metrics used here do not measure canine coding constraint directly. Instead, evolutionary-based metrics identify sites and genes conserved across many species and are blind to lineage-specific effects. Similarly, LOEUF values were calculated in human genes, and are not only blind to canid lineage-specific selection but also capture human lineage-specific effects that may not be relevant to canines. We therefore evaluated the capacity of each constraint measure to predict gene tolerance to mutations within dogs by measuring their concordance with the ratio of observed to expected mutation rates in canine genes.

Calculation of the observed mutation rates for this analysis must only capture negative selection effects and no other population genetic effects. One approach is to use recent spontaneous mutations, whose presence in the genome is unaffected by population dynamics. Similarly, expected mutation rates must also be unbiased. The observed genetic variation within dogs is a result of their complex demographic history, including recurrent population bottlenecks and strong selective sweeps (Marsden et al. 2016). It is difficult to determine whether a LoF mutation has persisted because it has a low deleterious impact or because it has a high

deleterious impact but is in linkage disequilibrium (LD) for a desired trait. Canine trinucleotide

substitution rates and coding sequence composition was therefore used to build a canine gene

mutation model, an approach similar to calculating LOEUF expected mutation rates (Samocha

et al. 2014) (**Methods**).

Singleton variants represent the best candidates for recent spontaneous mutations. To

further enrich for recent mutations, dogs with highly represented ancestry were identified using

individual singleton counts, as excess singletons per individual is an indicator of

underrepresented ancestry. Most dogs, ~70%, carried <40 singletons each within their coding

sequences, allowing this value to serve as a cutoff (**Supplemental Fig. S1A**). These dogs

belonged to breeds that had significantly more members in the dataset (greater Wilcoxon rank

sum test; P-value < 10^{-15}), and larger fractions of their genomes were within runs of

homozygosity (F_{ROH}) (greater Wilcoxon rank sum test; P-value < 10^{-15}) than was observed for

individuals with >40 singletons (**Supplemental Fig. S1B**). Only 11.2% of all singletons were

inferred to be recent spontaneous mutations by this method, and most low-frequency variants

were contributed by a minority of individuals with poorly represented ancestry (**Supplemental**

Fig. S1C).

Next, the canine mutation model was evaluated by testing whether it could predict the

number of synonymous mutations per gene. Since synonymous mutations typically have little

biological impact on coding sequence, their observed frequency should be highly correlated with

gene mutation probabilities. The Pearson correlation coefficient was used to evaluate model fit.

A total of 8,660 autosomal synonymous singletons were extracted from dogs with highly

represented ancestry and used as observations. A modest correlation of 0.53 was achieved, yet

due to the low number of observations, there was likely an insufficient number of events for

expected trends in mutation accumulation to emerge.

Sparsity in observations was addressed first by relaxing criteria for defining recent

spontaneous mutations, which involved increasing maximum allele count thresholds and

including variants from all dogs, and second by grouping genes based on mutation probabilities.

The 95% confidence intervals for expected fit were also calculated, allowing the mutation model

to be evaluated as a departure from expected fit values, which is less sensitive to the number of

observed events than the Pearson correlation coefficient. Although relaxing criteria for defining

recent mutations led to increased correlations, the departure from expected model fit also grew

(**Fig. 2A**), indicating that inclusion of additional variants to the analysis diluted the fraction of

true recent spontaneous mutations. The alternative strategy of placing genes into groups based

on mutation probability was more successful for testing model fit. For 1000 groups or less,

correlations greater than 0.9 were reached and the model fit remained within expected ranges,

unlike a competing model based on CDS length (**Fig. 2B**).

Together, these analyses show that accumulation of recent spontaneous synonymous

mutations across genes closely follows expected mutation probabilities, demonstrating that it is

possible to identify mutations in dogs where presence in the genome is largely independent of

canine population history, and that the proposed canine mutation model can accurately

calculate expected mutation frequencies for a given number of observations. This framework

can therefore be used to determine if genes have significantly fewer nonsynonymous mutations

than expected, thus detecting negative selection effects in the dog.

1
Figure 2: Recent spontaneous mutations in dogs and selection effects on constrained
regions. A) Pearson correlation coefficient between observed rare variants and gene mutation
probabilities. Variants are organized by the group of samples they were observed in and the
maximum allele count threshold for defining rare variants, which is displayed near each
datapoint. The grey band represents the 95% expected interval of the model fit based on
simulated accumulation of synonymous mutations. B) Pearson correlation coefficient of rare
variant frequency and gene mutation probability across gene groups. Genes were grouped

according to their mutation probabilities. The x-axis shows the number of groups used to
calculate correlation coefficients. The number of groups decreases as group sizes get larger.
Two models are tested, one is the mutation probabilities calculated using the mutation model
and the other is mutation probabilities according to gene CDS length. The 95% confidence
interval for the expected model fit are also displayed. C) Mutation frequency in genes ranked by
different constraint metrics. The expected number of mutations is calculated following the
mutation model. The observed number of mutations are from singletons in dogs with highly
represented ancestry. Error bars indicate whether the observed mutation frequency is outside
the 95% expected bounds. D) Enrichment of OMIA variants after variant/gene constraint
filtering. Vertical bars show the fraction of OMIA and all other variants that remain after
implementing each filtering threshold. Significance was determined using a Fisher test. "*"
indicates P-value < 0.05, "**" indicates P-value < 0.01, and "***" indicates P-value < 0.001.
13

The functional constraint of dog genes

Using 907 LoF and 14,947 missense singleton mutations from dogs with highly

represented ancestry, we tested whether the constraint metrics were associated with negative

selection effects. For gene-based metrics, dog genes were ranked and grouped into deciles by

constraint level, and mutation probabilities were summed accordingly. For variant phyloP

scores, a site-based metric, the individual CDS positions, were grouped into constraint deciles,

and base-pair mutation probabilities were summed according to each decile. Also, since phyloP

scores are heavily stratified by mutation type (**Fig. 1D**), breaks between phyloP score deciles

were made mutation type specific, thus balancing the expected frequencies of mutations across

the full spectrum of phyloP constraint.

Results show that gene-based constraint metrics in dogs are significantly associated

with intolerance to LoF mutations (**Fig. 2C**). The top three deciles for constrained CDS and top

two deciles for human ortholog LOEUF were significantly depleted of LoF mutations, indicating

that for this group of genes a significant portion of the expected LoF mutations likely had

deleterious impacts on fitness. The strongest selection effects were observed in the top human

ortholog LOEUF decile, where only half the expected number of LoF mutations were observed

(**Fig. 2C**). Conversely, no significant negative selection effects were observed for LoF mutations

among the top phyloP constraint deciles, indicating that evolutionary conservation across

species does not always correlate with mutation impacts on fitness.

In dogs, gene level constraint was also associated with intolerance to missense

mutations (**Fig. 2C**). These effects were less pronounced than LoF selection effects, as

functional consequences from missense mutations are often less severe. For variant phyloP

scores, intolerance to missense mutations was observed across the top four constraint deciles.

For synonymous mutations, no significant negative selection effects were observed amongst the

top gene-based constraint deciles, consistent with synonymous mutations having mostly

negligible impacts on gene function. Unexpectedly, top phyloP score deciles appeared

significantly intolerant to synonymous substitutions. However, since phyloP scores are site-

based, the observed intolerance may be in response to synonymous mutations that can cause

cryptic-splice sites, impact mRNA stability, have reduced tRNA availability, or induce change in

functionally relevant GC content (Sarkar et al. 2022; Gudkov et al. 2024).

We next tested whether constraint metrics were associated with known canine trait

variants. We increased the minimum threshold for defining constraint by decile increments and

measured enrichment of trait-associations within the remaining fraction variants. Canine trait-

associated variants were extracted from the online mendelian inheritance in animals (OMIA)

database (**Supplemental Table S3**), a catalogue of inherited disorders and other traits, and

associated genes in non-model animal species (Nicholas 2021; Nicholas 2024)

(https://www.omia.org/home/). Although OMIA is the largest collection of animal trait-associated

variants, supporting evidence for trait-associations vary widely, ranging from candidate alleles

found in single individuals, to mutations shown to be causative in functional studies. We focused

only on OMIA variants with an allele count > 2 within the combined cohort. This cutoff was used

to ensure that we would only analyze variants that could be identified using genotype-first

approaches. Altogether, 31 out of 305 LoF and 50 out of 154 missense OMIA variants met these

criteria (**Supplemental Table S3**).

For evolutionary-based constraint metrics, we detected enrichment of both LoF and

missense OMIA variants (**Fig. 2D**). However, in most cases increases to the minimum constraint

threshold beyond a certain point led to a loss of OMIA enrichment (**Supplemental Table S4)**.

Similarly, no enrichment of LoF OMIA variants were detected for human ortholog LOEUF

constraint, which was associated with the strongest negative selection effects. This outcome

indicates that OMIA LoF variants likely do not have severe deleterious impacts on fitness. Such

variants would be unlikely to persist in canine populations, unlike the variants analyzed here,

which were all present in multiple dogs within the cohort.

Genotype-first approaches applied to canine variation

For most dogs in this dataset, the only individual information available was sex and

breed. We therefore focused on genotypes that were enriched in a particular breed. We also

investigated the breed specificity of OMIA variants across a range of minimum breed frequency

cutoffs. We found that increases to the minimum breed allele frequency cutoff result in fewer

discoverable OMIA variants. At a 20% minimum breed allele frequency, the fraction of the

remaining variants found in only one breed is at its highest (**Supplemental Fig. S2**). For the

remainder of the analyses, we therefore only considered variants with a minimum breed allele

frequency >20% in at least one breed.

Lists of potential LoF and missense candidates for canine traits were defined using each

constraint metric. Where possible, cutoffs were selected at the points where negative selection

effects could be detected (**Fig 2C**). When there was no such effect, such as for variant phyloP

scores and LoF mutations, the highest percentile cutoffs where OMIA enrichment was observed

were used instead (**Fig 2D**). Variant lists were filtered using a minimum breed allele frequency

of 20%, a minimum allele count of three, a minimum breed allele number of six, and a maximum

non-breed allele frequency of 5%. Variant lists were then ranked according to the number of

breeds the variant was found in and the proportion of individuals in those breeds carrying the

variant (**Fig 3.**) (**Supplemental Table S5-S10**).

Table 1: Breed genotypes and allele frequencies of select candidate variants.

Variant	Breed	AF	Ref	Het	Hom
chr13:47440160, A/C, *PDGFRA*, splice acceptor	Catalburun	0.5	0	3	0
chr1:66494619, C/T, *RSPO3*, stop gained	Irish Water Spaniel	0.33	2	4	0
	Unassigned	0	324	1	0
chr6:42492671, C/A, *CSF1*, stop gained	Chow Chow	1	0	0	6
	Elo	0.1	4	1	0
chr3:91871030, AACAG/A, *LCORL*, frameshift variant[1]	Spanish Mastiff	1	0	0	6
	Berger Picard	0.77	0	5	6
	Briard	0.67	0	4	2
	Petit Bleu de Gascogne	0.58	1	3	2
	Bluetick Coonhound	0.5	0	2	0
	Bracco Italiano	0.5	3	0	3
	Spinone Italiano	0.43	2	4	1
	Pyrenean Mastiff	0.36	3	3	1
	Leonberger	0.32	9	8	2
	Beauceron	0.3	2	3	0
	Mastiff	0.28	5	3	1
	German Hound	0.25	3	3	0
	Griffon Nivernais	0.25	2	2	0
	Ariege Pointer	0.2	3	2	0
	Karakachan	0.2	3	2	0
	Picardy Spaniel	0.2	3	2	0
chr30:32785671, ACT/A, *ITGA11*, frameshift variant[1]	Pug	1	0	0	3
	Japanese Chin	1	0	0	4
	Pekingese	1	0	0	4
	Shih Tzu	1	0	0	4
	Tibetan Spaniel	1	0	0	4
	Boston Terrier	1	0	0	6
	French Bulldog	1	0	0	6
	Lhasa Apso	1	0	0	6
	Chihuahua	0.36	4	1	2
	Biewer Terrier	0.3	2	3	0
	Portuguese Podengo Pequeno	0.25	1	1	0
	Russian Tsvetnaya Bolonka	0.25	3	3	0
	Chesapeake Bay Retriever	0.21	4	3	0

[1]Only breeds with AF ≥0.2 are displayed

1
Fig 3: List of candidate LoF/missense variants for trait associations. Variants are ordered
along the y-axis according to number of breeds they are found in and the proportion of
individuals in those breeds carrying the variant. Breeds carrying the variant are ordered along
the x-axis according to the proportion of the alt alleles they are carrying. Gene names printed in
black text contain OMIA variants. Gene names in red are non-OMIA variants that were identified
after manual curation of the variant list. Variant type and constraint cutoff are displayed for each
panel.
9

For LoF constraint, the number of variants returned after filtering by breed allele

frequency ranged from 191 for human ortholog LOEUF to 584 for variant phyloP score. For

missense mutations, the number of returned variants ranged from 1,637 for constrained CDS to

10,351 for phyloP score (**Fig. 3**). Since LoF constraint filtering returned fewer variants, and is

more highly associated with negative selection effects, the associated variant lists were

manually investigated to identify new candidate genotype-phenotype relationships. Based on

their potential for phenotypic impact, a total of six variants were selected for comment (**Table 1**).

The most interesting candidate was a splice-acceptor within the *PDGFRA* gene that was

found in a heterozygous state in all three Çatalburun dogs within the cohort. Members of this

breed, also called the Turkish pointer, are notably characterized by a split/bifid nose (**Fig 4A**)

(Yilmaz et al. 2012). Indeed, the English translation of "Çatalburun" is "forked nose" (Özarslan et

al. 2021). Mouse knockout models demonstrate that *PDGFRA* plays an important role in medial

nasal process development and palatogenesis (He and Soriano 2013; Qian et al. 2017), thus

aligning with the hypothesis that the *PDGFRA* splice-acceptor variant causes the Çatalburun's

bifid nose phenotype. The variant was genotyped in fourteen additional Çatalburun dogs,

although phenotypes were not available, the variant was heterozygous in all but one dog which

was homozygous for the reference allele (**Supplemental Table S11**).

To further investigate the genetics of this phenotype, we recruited and performed WGS

on eight additional bifid nose dogs, all lacking any apparent Çatalburun ancestry. Six of the dogs

were unrelated to one another, while two were litter mates. An unaffected parent of the two

affected siblings was also in included (**Supplemental Fig. S3**). Variants were prioritized

according to evolutionary constraint and potential functional impact on orofacial cleft associated

genes. Twelve heterozygous candidate variants were identified across all eight non-Çatalburun

bifid nose dogs (**Supplemental Table S12**). One sample, 40931 (**Fig. 4B**), carried a

heterozygous insertion that also disrupted a *PDGFRA* splice-acceptor but at a different position

(**Fig. 4C**). When considering all 2,291 dogs in the entire cohort, only those with the bifid nose

phenotype carried LoF mutations in *PDGFRA*, thus defining the existence of a naturally

occurring model for study of *PDGFRA's* role in craniofacial development and midline closure. An

additional breed, the Pachón Navarro, which originates in Spain, is also known for its bifid nose

phenotype. Unfortunately, no samples from this breed were available for study. Interestingly,

however, masonry on the outside of the 18th century Palace of Bertemati (Andalusia, Spain)

shows a gargoyle-like dog with a bifid nose (Cobo 2017), attesting to the continuous selection

for this trait in the Pachón Navarro breed (**Fig. 4D**).

2 **Figure 4: Phenotypic variant discovery.** A) Çatalburun dog with distinctive bifid nose
3 phenotype. B) Golden retriever mix (Dog ID: 40931) displaying a bifid nose phenotype and is
4 heterozygous for an additional *PDGFRA* splice-acceptor variant. C) Canine *PDGFRA* transcript
5 showing the locations of each splice acceptor variant found within dogs with orofacial clefts. The

first variant is the çatalburun variant with the second variant belonging to dog 40931. D)
Photograph of dog statue with split-nose phenotype. The statue is located at the Palace of
Bertemati in Andalusia, Spain. The palace was constructed in the 18[th] Century. E) Mutations
found in the long isoform of *LCORL*. Early termination of translation and loss of the DUF4553
domain caused by frameshift mutations appears to be associated with larger dog size (>
25kg/>55cm). F) *LCORL* genotypes and breed standard weight. G) *LCORL* genotypes and
breed standard height.
8

Another variant with potential implications on craniofacial morphology is a frameshift

within *ITGA11* (**Table 1**). It is fixed in small brachycephalic breeds, and mouse knockouts of the

gene demonstrate its association with small size and craniofacial abnormalities, specifically the

eruption of incisors (Popova et al. 2007). In dogs, an association between the locus and

brachycephaly was described previously but was sensitive to corrections for canine size

(Schoenebeck et al. 2012; Marchant et al. 2017). The variant could cause a range of complex

phenotypes, contributing to both size and craniofacial structure. However, additional functional

analyses are needed.

A new mutation within *LCORL*, a previously described canine size-associated gene

(Hayward et al. 2016; Plassais et al. 2019), was also identified. The previously described

variant, c.3768_2769insA, causes a frameshift and premature stop codon in the long isoform of

the gene. The variant is one of the largest contributors to canine size, explaining ~15% of the

variance in canine standard breed weight (Plassais et al. 2019). Here, a newly identified *LCORL*

variant, c.1978_1981delACAG, a deletion upstream of the previously described insertion, also

causes a frameshift and premature stop (**Fig. 4E**). The c.1978_1981delACAG variant was fixed

in Spanish Mastiffs, an extremely large breed whose standard has no upper limit on size, and

segregates in several other large breeds, i.e., those weighing > 25kg (**Table 1**). At an allele

frequency of 2.9%, the newly identified c.1978_1981delACAG variant is less common across

the entire cohort than the c.3768_2769insA, which has an allele frequency of 31.2%. Among

large dogs, the frequency of the c.1978_1981delACAG is 6.4% and 37.9% for

c.3768_2769insA. Both alleles were also significantly associated with both breed standard

weight (**Fig. 4F**) and height (**Fig. 4G**) (P-value < 10^{-10}). There were 30 instances where dogs

had one copy of each allele and no instances of a dog having two copies of one allele and one

or two copies of the other, indicating that the c.3768_2769insA and c.1978_1981delACAG

alleles are most likely on different haplotypes.

Another variant that may correspond to a breed trait is a stop-gain at position p.Glu391*

in *colony stimulating factor 1* (*CSF1*). The variant was homozygous in all six unrelated Chow

Chows (Chow), heterozygous in one of five Elos, and absent from two Eurasiers, the latter two

breeds being the product of Chow admixture (**Table 1**), thus indicating a high degree of breed

specificity. The Chow is a medium-sized spitz-type breed that traces its origins to China (Yang et

al. 2017). It is characterized by its thick double coat, black pigmented tongue, straight hind legs,

broad square skull and muzzle, and supernumerary third incisors (Yang et al. 2017). The region

containing *CSF1* is under strong positive selection in Chows (Yang et al. 2017), suggesting

genes in this region contribute to the breed's characteristic phenotypes. The p.Glu391* mutation

is located in exon six of the secreted isoform of the CSF1 protein (Cosman et al. 1988). Rodent

knockout models implicate *CSF1* in skeletal and craniofacial development, specifically a domed

skull, missing incisors, short thickened limb bones, fertility, and macrophage differentiation

(Marks and Lane 1976; Yoshida et al. 1990; Dai et al. 2002; Van Wesenbeeck et al. 2002;

Sehgal et al. 2021). While the precise function of the p.Glu391* mutation in dogs is unknown,

the rodent phenotypes of a domed skull and altered number of incisors, suggest p.Glu391* may

contribute to the Chow's broad square skull and two additional incisors, which contrast sharply

with the Eurasier and Elo breed requirements.

One variant with potential medical consequences was a stop gain within *RSPO3* and

was found in a heterozygous state in four out of six Irish Water Spaniels and one Xoloitzcuintle,

the latter being a hairless Mexican breed whose members were largely unassigned to a

breed/group during phylogenetic clustering (**Table 1**). This mutation is of particular interest as a

167 bp insertion in the 3'UTR of a gene from the same family, *RSPO2*, is responsible for

"furnishings", defined as the long moustache and increased eyebrow hair observed in breeds

like the schnauzer (Cadieu et al. 2009). Depletion of Rspo3 within the mouse dermal papillae

causes delayed hair regrowth, indicating it is required to activate proliferation of dermal stem

cells within the hair follicle, although *RSPO2* may provide some functional redundancy (Hagner

et al. 2020). Since the *RSPO3* mutation in Irish water spaniels is not found within every breed

member, it is unlikely to be connected to a breed-characteristic trait. Rather, the variant may

explain why some Irish Water Spaniels suffer alopecia from the trunk or thighs as they age,

which occurs with high frequency (Cerundolo et al. 1999).

Coding variation within genomic regions exhibiting differentiation across breeds

Artificial selection for characteristic breed traits has left its mark on the dog genome.

Within breeds, there are extended regions of homozygosity that are also associated with strong

signatures of population differentiation between breeds (Akey et al. 2010; Vaysse et al. 2011;

Parker et al. 2017a; Morrill et al. 2022). Moreover, trait-associated variants for breed

characteristic traits, such as ear shape, body size, and sport-hunting ability are often within

breed-differentiated genomic regions, where genetic variation is extremely stratified by breed

(Boyko et al. 2010; Kim et al. 2018; Plassais et al. 2019). Like genotype-first approaches, breed-

differentiated regions provide a genomic signature potentially associated with canine breed

traits. However, unlike genotype-first approaches, breed-differentiated loci often encapsulate

multiple genes, making it difficult to identify specific variants that contribute to breed

characteristic traits. To enhance detection of trait-associated variants, potentially functional

variants in constrained genes were analyzed for their association with signatures of breed

differentiation.

Pairwise F_{ST} between 200 breeds/groups was calculated for 50kb genomic windows.

Windows with $F_{ST} > 0.8$ signified extreme differentiation between a breed/group pair. Loci with >

100 extremely differentiated breed/groups pairs were classified as breed differentiated regions.

These regions were further refined using peak detection software (Locher 2024). Genomic

segments of extended breed homozygosity were also identified and were also used to further

refine breed-differentiated regions by providing an additional marker of artificial selection

(**Methods**) (**Fig. 5A**). In total, there were 1,743 breed-differentiated regions, containing 2,085

genes (**Supplemental Table S13**). However, 92.3% and 55.2% of breed-differentiated regions

were within 1 Mb and 100 kb of each other, respectively, indicating that most breed-

differentiated regions are likely part of larger selective sweeps. Seventeen of 25 autosomal trait-

associated loci described in Plassais et al. (2019) were located within 100kb of a breed-

differentiated region, indicating that the regions identified here are associated with breed traits

under selection (**Fig. 5B**). Population branch statistics from an analysis of 101 different breeds

inclusive of >12 dogs were also used to identify breed-differentiated regions, capturing a total of

3,016 genes (Morrill et al. 2022). Of these, 726 were within breed-differentiated regions

identified in this analysis.

Fig 5: Non-coding variation at constrained genes drives breed differentiation. A)
Identification of breed differentiated regions at chromosome 15 IGF1 locus. Genomic windows
with $F_{ST} > 0.8$ between breed pairs were plotted across the genome, and peaks were then
identified as regions of breed differentiation. High F_{ST} peaks are shown in the top panel and
highlighted in red. The second panel shows genes found in the region. The third panel shows
the density of regions of breed homozygosity that were identified as 1 Mb genomic regions with
breed nucleotide diversity < 0.02. The final panel shows black bars as individual breed
homozygosity stretches. B) Genome-wide display of breed-differentiated regions that overlap
previously characterized trait loci from Plassais et al (2019), which are displayed as lower-case
characters. Loci in red are within 100kb of a breed-differentiated region. C) Number of genes in
highly differentiated regions carrying potentially functional variants that pass minimum constraint
thresholds. The High F_{ST} column is the number of genes that carry variants of interest. The Low
pi column is the count of genes that carry variants that are in breeds with low nucleotide
diversity across the differentiated region. The final column is the number of genes with variants

enriched in breeds that share the same haplotype within the differentiated region. D) Observed over expected frequency of genes at different constraint levels within highly differentiated regions.

There were between 10 and 34 genes across all breed-differentiated regions that

contained LoF mutations with potential functional consequences (**Fig. 5C**). Of these, only two

were found in breeds that exhibited signatures of differentiation at these loci, specifically the

CSF1 premature stop codon found in Chows (Table 1) and the *SGK3* mutation causing

hairlessness in American Hairless Terriers (Parker et al. 2017b). Although not significantly

associated with breed differentiation, a subset of mutations was within breed-differentiated

regions and segregating in the dogs whose breeds contained long stretches of homozygosity.

One example is a frameshift mutation caused by a single base pair insertion in *CYP4A11*, a

monooxygenase involved in drug metabolism and synthesis of cholesterol, steroids and other

lipids, associated with cardiovascular and cerebrovascular disease and hypertension in humans

(Sugimoto et al. 2008; Yu et al. 2018) (**Supplemental Table S14**). The canine variant may also

be associated with health outcomes in dogs and may persist because it is in LD with a trait

under positive selection.

Altogether, 70 genes contained missense mutations with potential functional impacts that

were significantly associated with breed differentiation (**Fig. 5C**). The strongest result was for a

missense variant in *RYR1* (**Supplemental Table S14**), a calcium ion channel primarily

expressed in the sarcoplasmic reticulum of skeletal muscle and involved with muscle

contraction (Hernandez-Ochoa et al. 2015). The variant was almost exclusively found within Bull

Terriers and American Staffordshire Terriers, two breeds with shared ancestry that are known for

their musculature and agility. Another member of the gene family, *RYR3*, which is also involved

with muscle contraction, is under positive selection in sport-hunting breeds (Kim et al. 2018).

The vast majority of breed-differentiated regions were not associated with

nonsynonymous variants, indicating that breed selection may instead be acting upon non-

coding variants that likely impact gene expression, which is consistent with most mapping

studies for breed-defining traits and breed/lineage differentiation (Dutrow et al. 2022; Morrill et

al. 2022). The genotype-first approach described here focuses on identifying variants with

potential phenotypic impacts, prioritizing variants in constrained genes. Since breed-

differentiated regions often represent loci under selection for breed characteristic phenotypes

(Plassais et al. 2019; Morrill et al. 2022), these regions should also be comprised of constrained

genes. We tested for enrichment across constraint deciles for gene-based metrics and found

that human ortholog LOEUF constraint was significantly overrepresented (**Fig 5D**). This

supports a model of phenotypic breed differentiation driven by regulatory variation of highly

constrained genes rather than alterations to gene function from nonsynonymous mutations.

Discussion

Artificial selection for desired traits has made the domestic dog one of the most

phenotypically diverse mammals in the world (Wayne and vonHoldt 2012). Rigid population

structure, enforced by breed standards, has promoted high levels of homozygosity within breeds

while also maintaining isolation between breeds (Parker et al. 2004; Karlsson and Lindblad-Toh

2008). Together, these factors make the domestic dog an invaluable system for identifying

genes related to breed characteristics, which are often morphological extremes (Ostrander et al.

2017; Buckley and Ostrander 2024). We therefore investigated the genetic basis of phenotypic

breed differences using a genotype-first approach. Specifically, we used metrics for genomic

constraint to identify variants with potential phenotypic impacts. We then ranked variants

according to breed enrichment. Variants common to one or a few breeds but rare throughout the

population are potential candidates for breed traits. This work provides significant advances to

the field of canine genetics by: 1) defining a variant set whose presence in the genome is

unaffected by canine population dynamics, thus making it possible to determine genes under

negative selection; 2) evaluating three constraint metrics for prioritizing variants for potential

phenotypic impacts; 3) identifying new CDS variants with strong impacts on craniofacial

morphology and body size; and 4) applying genomic constraint to support a model of breed

differentiation based on non-coding variation.

Using the genotype-first approach, we identified strong candidates for phenotypic

impacts as previously unreported LoF variants in *PDGFRA* and *LCORL,* genes both under

strong human ortholog LOEUF constraint. Two *PDGFRA* variants were identified, one was in

Çatalburun dogs, a breed largely characterized by a bifid nose phenotype (Fig. 4A), and the

other was in a Golden Retriever mix. This gene may therefore be one of a few for which

functional mutations affecting midline closure occur, but are tolerated, as long as they are in the

heterozygous state. In humans, a missense variant in this gene within a single family is

associated with left non-syndromic cleft-lip and palate but is not fully penetrant (Yu et al. 2022).

PDGFRA is a type III receptor tyrosine kinase that consists of an extracellular region with five

immunoglobulin-like domains for ligand binding, a transmembrane helix, and an intracellular

tyrosine kinase domain (Williams 1989; Chen et al. 2013). The Çatalburun variant occurs in the

seventh intron of the gene within the fourth immunoglobulin-like domain, while the other variant

in the Golden Retriever mix is further downstream within the protein kinase domain (**Fig 4C**).

Both mutations are splice-acceptors, making it difficult to determine their impact on transcription.

The affected exons may be skipped entirely, or an alternative splice site may be used, resulting

in either partial exon retention or retention of nearby intronic sequence. Dogs carrying the

PDGFRA mutations had no reported health concerns, which is surprising as *PDGFRA* is a

broadly expressed developmental gene and strongly associated with gastrointestinal stromal

tumor-plus syndrome in humans (Manley et al. 2018; Huang et al. 2022).

Two other breeds share a similar phenotype as the Çatalburun, the Andean tiger hound,

about which little is known, and the Spanish Pachón Navarro. The Pachón Navarro, like the

Çatalburun, is a pointing breed, and not all breed members have a bifid nose. Evidence for bifid

nose dogs in Spain can be observed not only in 18[th] century architecture (**Fig. 4D**), but also in a

1623 painting of a hunting party titled "Diane et ses nymphes s'apprêtant à partir pour la

chasse" by Pierre Paul Rubens and Jan Brueghel l'Ancien. Other works from around the turn of

the 20th century describe bifid nose pointing dogs throughout the south of France (Arkwright

1906). Although it is unknown whether the bifid nose phenotype observed in the Çatalburun and

Pachón Navarro breeds has a common origin, Turkey and Spain were both under the rule of the

Umayyad Caliphate in the 8th century CE, providing a possible route for cultural exchange

between the two regions. Given our observation of multiple variant alleles among bifid dogs

recruited for this study and the recurrent artistic depictions, the phenotype could have arisen

multiple times in pointers.

The newly discovered *LCORL* c.1978_1981delACAG allele was identified because of its

high frequency in Spanish Mastiffs, Berger Picards, and Briards (**Table 1**). Like the earlier

discovered *LCORL* c.3768_2769insA variant, the c.1978_1981delACAG allele causes a

frameshift and premature stop, resulting in loss of the DUF4553 DNA-binding domain from the

long isoform of LCORL (Plassais et al. 2019). Both alleles are associated with larger size and

stature in dogs (**Fig 4F,G**). *LCORL* is also a primary determinant of stature across many

domesticated mammals including goats (Saif et al. 2020; Graber et al. 2022), cattle (Bouwman

et al. 2018; Chen et al. 2020), pigs (Rubin et al. 2012), and horses (Metzger et al. 2013; Tetens

et al. 2013). Given that the gene is under selection in multiple species, it is unsurprising that

more than one *LCORL* allele is under selection. The presence of both alleles within the same

breeds also suggests that both variants predate breed development. Further investigation into

the mechanisms by which trait variants affect phenotypes may reveal other instances where

selection for different alleles achieves redundancy, thus assuring the same phenotypic outcome.

The association between breed-differentiated regions and potential trait variants in

constrained genes returned few associated CDS variants. Importantly, none of these

associations included OMIA variants prioritized using genotype-first approaches, indicating that

signatures for breed enrichment are distinct from signatures for selection/breed differentiation.

This is because many OMIA variants are disease alleles and, while overrepresented in a small

subset of breeds, are not under positive selection (Boyko et al. 2010; Kim et al. 2018; Plassais

et al. 2019). With additional disease risk phenotyping for dog breeds, genotype-first approaches

may be useful for expanding lists for disease risk alleles. Most genes in breed-differentiated

regions had no association with potentially functional CDS variants, consistent with non-coding

variation being the primary driver of phenotypic variation across breeds, a repeated finding from

other large-scale multi-breed canine genetic analyses in dogs (Sahlen et al. 2021; Dutrow et al.

2022; Morrill et al. 2022) and humans (Halldorsson et al. 2022). However, breed-differentiated

regions were enriched for constrained genes, emphasizing the importance of constraint in

predicting phenotypic impacts.

One of the defining features of the canine system is that often few loci contribute to large

phenotypic variation, particularly for morphological traits like breed standard weight and height

or ear shape and position (Rimbault et al. 2013; Plassais et al. 2019), while in the human

genome there are often large numbers of loci associated with the same phenotypes (Yengo et

al. 2022). Loci contributing to extreme phenotypic differences between breeds overwhelmingly

exhibit strong signatures of positive selection because of artificial selection for desired traits

(Plassais et al. 2019; Morrill et al. 2022; Plassais et al. 2022). Our analysis shows that highly

constrained genes are overrepresented in such regions and are therefore the likely targets of

trait selection, driving phenotypic variation. This suggests an explanation for why CDS variation

is rarely associated with morphological differences between breeds and more often associated

with disease. Constrained genes are often essential for development and survival and are

frequently implicated in highly penetrant genetic diseases and syndromes (Lek et al. 2016;

Karczewski et al. 2020; Sun et al. 2024; Zeng et al. 2024). Trait selection in dogs is therefore a

balancing act between preserving essential gene functions and maximizing regulatory variation

of essential genes to drive phenotypic extremes. Since breed variation is driven by the

expression of genes with essential biological roles, future work should focus on assigning non-

coding variants under positive selection to their target genes.

Central to the analysis is the identification of variants with phenotypic impacts, which

was assessed according to evolutionary/functional constraint acting on genes/sites containing

the variant in question. However, specific factors impact the interpretation of these results. For

instance, the measured selection effects only capture negative selection acting on heterozygous

variants with highly deleterious impacts on fitness (Fuller et al. 2019). Our analysis is therefore

blind to recessive deleterious alleles and genes associated with late age of onset conditions

such as breed-prevalent diseases like cancer. Since singletons are the input to the mutation

model, depletion of variants with little to no deleterious impact in a heterozygous state remain

unobservable, as do variants that produce no phenotype until after a dog's reproductive years

have elapsed. This is relevant to canines where a high degree of inbreeding makes dogs

especially susceptible to genetic load and recessive disease allele burden (Marsden et al. 2016;

Mooney et al. 2021; Donner et al. 2023). Moreover, OMIA variants segregating in canine

populations likely persist because they do not have strong effects on fitness in a heterozygous

state. In fact, many of the OMIA variants underlie traits, such as coat color or recessive and late-

onset diseases, such as the many forms of progressive retinal atrophy (Miyadera et al. 2012;

Downs et al. 2014; Bunel et al. 2019). Therefore, strong human ortholog LOEUF constraint

lacks an association with OMIA variants because OMIA LoF variants have a low impact on

fitness, and human ortholog LOEUF constraint indicates genes where heterozygous LoF

mutations have a strong deleterious impact on fitness. Another factor they may impact

interpretation of results is whether the unit for measuring constraint is gene or base-pair. The

increased resolution provided by site-specific phyloP scores is only beneficial for analyzing

missense mutations as mutations annotated as LoF, defined throughout as premature stop

codons, frameshifts, and splice-site disruptions, often impact the entire transcript downstream of

the mutation. Missense mutations, alternatively, have more localized effects and usually cause

negligible impacts on biology unless interrupting important functional domains (Gerasimavicius et al. 2022). Therefore, in dogs, human ortholog LOEUF is the best of the three constraint metrics for prioritizing LoF mutations for phenotypes with severe impacts on fitness. Conversely, variant phyloP scores are best for prioritizing missense mutations.

While the genotype first approach is ideal for a species with a complicated population structure like dogs, it is facilitated by sufficient breed representation and availability of accurate phenotype information. While mapping breed standard traits has proven fruitful, more refined phenotyping and mapping of more nuanced traits will require individualized measures of each dog. For instance, CT scans are now fundamental to genetic studies of canine skull morphology, as reliance on owner measurement and breed descriptors lacks precision (Marchant et al. 2017).

The dog genome project began with two goals, and both remain a high priority today: improve the health of canine companions through informed genetics and translate those findings to advance human health. While many breeds feature largely healthy, long-lived animals, others are characterized by an excess of disease alleles and associated health problems (Axelsson et al. 2021). These breeds offer a unique opportunity to make use of the approaches and data provided here to identify underlying genetic flaws, particularly for diseases of relevance to human health. Of equal priority are studies of morphologic and behavioral genetics, as identification of alleles that control extreme phenotypes associated with these traits can improve our understanding of human growth and development.

Methods

Defining a canine mutation model and callable gene set

We built a mutation model in dogs following the methods described in (Samocha et al. 2014). Briefly, trinucleotide substitution rates, normalized according to the generation mutation rate and trinucleotide frequencies, were applied to each possible substitution for each site within gene

models. For each gene model, substitution probabilities were summed according to their

consequences on coding sequence, such as synonymous, missense, stop gain, stop loss,

etcetera. The result was per gene probabilities for each mutation type per generation

(**Supplemental Fig. S4**). Dog10K variants at least 20 bp from an exon and within callable loci

were used to calculate trinucleotide substation rates (Meadows et al. 2023). A canine generation

mutation rate of 9.02×10^{-9} was used for normalization (Bergeron et al. 2023) (**Supplemental**

Fig. S5). We calculated transcript-specific mutation probabilities using NCBI gene annotations

from Dog10K, only considering sites that were callable loci, obviating the need to normalize

probabilities by mean exon coverage (**Supplemental Table S15**). There was insufficient variant

data to adjust mutation probabilities by evolutionary conservation as was performed in

(Samocha et al. 2014). We classified genes according to intact start and stop codons, low-

quality annotation, and partial annotation. Also, for each gene, a representative transcript was

chosen based on reciprocal best hit blast results with protein sequences of human

representative transcripts. Where no blast hit was found, the first annotated transcript was used

as a representative. Blast hits were also defined as dog-human orthologs and used to assign

human LOEUF constraint metrics to dog genes. For phyloP constraint, we used the Zoonomia

values mapped to UU_Cfam_GSD_1.0 as part of Dog10K, which were further summarized

according to gene models (**Supplemental Table S16**). Expected mutations per gene were

calculated by sampling *n* gene names with replacement weighted by gene/gene group mutation

probabilities, where *n* is the number of observations the expected rate is being compared to.

Sampled gene names are tabulated to get the expected per gene mutation count. This process

is replicated 10,000 times to generate 95% expected ranges.

Preparing genotypes for analysis

Genotyping and filtering procedures are described in full in the supplemental methods. Briefly, a

VCF comprised of SRA genomes was constructed using GATK to complement the publicly

available Dog10K VCF (Poplin et al. 2017). All WGS data was mapped to the

UU_Cfam_GSD_1.0 reference assembly. For each VCF, samples were filtered according to a

variety of genotype metrics (**Supplemental Table S17**), which included coverage statistics

(**Supplemental Fig. S6**), heterozygous allele balance (**Supplemental Fig. S7**), as well as

several others (**Supplemental Fig. S8**). A total of 1,146 SRA and 1,778 Dog10K samples

remained (**Supplemental Table S18**). Duplicates and first-degree relatives across both

datasets were removed from the analysis (**Supplemental Fig. S9**). A total of 2,291 samples

remained (**Supplemental Table S19**). Sites were filtered according to variant quality score

recalibration (VQRS) using a common training set (**Supplemental Fig. S10**). Autosome and

chromosome X PAR sites (**Supplemental Table S20**) were analyzed separately from

chromosome X non-PAR sites (**Supplemental Table S21**). Sites that passed VQSR filtering in

both datasets were used to create a merged VCF for population-scale analyses. Sites that

passed filtering in either dataset and were within the CDS underwent further filtering

(**Supplemental Fig. S11**) (**Supplemental Table S22**). Criteria for filtering CDS genotypes

depended on chromosome, sex, and genotype (**Supplemental Table S23**). CDS indels and

SNVs were reclassified and atomized in both datasets to improve harmonization

(**Supplemental Fig. S12**) (**Supplemental Table S24**). Over 95% of SNVs and over 80% of

indels passed filtering (**Supplemental Table S25**). Genotype quality was the criteria that failed

most often (**Supplemental Table S26**). Sample genotype pass rates were higher for SNVs than

for indels (**Supplemental Table S27**). CDS filtering removed genotyping biases between each

dataset for SNVs (**Supplemental Fig. S13**) and indels (**Supplemental Fig. S14**). Genotypes

were merged and annotated with variant effect predictor (VEP) (McLaren et al. 2016). All VCF-

based operations were performed with bcftools (Danecek et al. 2021), plink (Purcell et al. 2007),

vcftools (Danecek et al. 2011), and vcflib (Garrison et al. 2022).

Defining breeds and groups

Illumina CanineHD array genotypes from 2,156 samples were lifted over to the

UU_Cfam_GSD_1.0 reference genome using LiftoverVCF (**Supplemental Table S28**). WGS

sites were downsampled to match array positions and datasets were merged. After filtering at

MAF > 0.01, genotyping rate > 0.9, and LD < 0.5, a total of 76,570 sites remained (Purcell et al.

2007). Using Phylip, a consensus neighbor joining tree was built with Golden Jackal set as the

outgroup (Felsenstein 1993). Branch support was calculated from 10% jackknifing 10,000 times.

Branches needed at least 50% support. Breed/group clusters were determined by optimizing for

shared breed membership on the same branch. Samples that did not match the majority breed

on a branch were assigned new breed membership, matching the majority breed. Branches that

consisted of overlapping breeds were merged. Breeds that could not form a branch of at least

three members were placed in the unassigned category.

Canine population statistics and breed differentiated loci

Population analyses were only performed on the 200 breeds/groups that had five or more

members. Genome-wide population statistics were calculated across 200,000 randomly

sampled sites. Nucleotide diversity within breeds was measured as pi, the average distance

between two randomly sampled sequences in a breed (Nei and Li 1979). Breed differentiation

was measured as F_{ST}, using the Weir and Cockerham estimator on a per-site basis (Weir and

Cockerham 1984). F_{ROH} was calculated using the same procedures as specified in (Meadows et

al. 2023). We detected breed-differentiated regions by calculating F_{ST} between all 200 pairs of

breeds with five or more members across 50kb windows using a 10kb step size. Windows with

F_{ST} values > 0.8, which corresponded to ~0.1% of all pairwise comparisons, were extracted for

further analysis. The density of high F_{ST} windows was then measured across the genome and

the "peaks" function of IDPmisc, a peak-calling software R package, was used to define breed

differentiated regions (Locher 2024). Next, for each region, we extracted breeds with pairwise

F_{ST} values > 0.8 within the region's peak window. Extracted breed pairs with F_{ST} < 0.05 were

defined as sharing the same haplotype and were assigned the same cluster ID. We also defined

genomic segments of extended breed homozygosity if half of the 50kb windows along a

megabase segment had pi values < 0.02. Breed-differentiated regions were defined as selection

loci in individuals where there was overlap with extended homozygosity regions.

Identification of additional orofacial cleft genes

Blood or saliva samples were collected from nine additional dogs, eight with a bifid nose, of

which two were littermates. The non-bifid nose dog is a parent to the littermates. Samples were

whole genome sequenced at mean 20x coverage and genotyped following the above methods

used for SRA samples. Variant sites shared with the Dog10K cohort were filtered out from the

analysis. Next, SVNs were discarded if QD < 2, SOR > 3, FS > 60, MQ < 40, MQRankSum < -

12.5, or ReadPosRankSum < -8.0. Indels were discarded if QD < 2, FS > 200,

ReadPosRankSum < -20.0, or variant length was > 10 bp. Since orofacial clefts within this

cohort were likely sporadic, except for the siblings and parent, we focused specifically on high

quality heterozygous genotypes that met the following criteria: GT = 0/1, GQ >= 20, DP >= 10,

AD ratio > 0.5, and AD ratio < 2.0. Next, we selected variants as candidates if they were found

in any orofacial cleft candidate genes, which were determined according to membership within

the following gene lists; DisGeNET associations for craniofacial abnormalities (C0376634), cleft

Palate (C0008925), or cleft upper lip (C0008924) (Pinero et al. 2017), experimentally identified

orofacial cleft genes (Xu et al. 2021), and genes with restricted craniofacial expression during

human development (Yankee et al. 2023).

Acknowledgements

We thank Tatiana Feuerborn and Heidi Parker for their helpful advice and insight throughout the

project and Gabby Spatola for assistance in recruiting samples. We also thank Zachary

Lounsberry and Embark Veterinary for assistance in recruiting samples and thank Sruthi Hundi

for bioinformatic assistance. For careful reading of the manuscript and providing helpful

feedback, we also thank Christopher B. Kaelin and Wesley C. Warren. Most importantly, we

thank all citizen scientists who donated samples for analysis, without whom this work would be

impossible. E.A.O., R.M.B., A.C.H., D.L.D., A.S.A are funded by the Intramural Program of the

National Human Genome Research Institute at the National Institutes of Health (HG200377).

H.L. is funded by the Jane and Aatos Erkko Foundation.

Competing interest statement

A.R.B. is a co-founder and board member of Embark Veterinary, Inc., a private canine DNA

testing company. All other authors declare no competing interests.

CRediT author statement

Conceptualization: R.M.B. and E.A.O.

Methodology: R.M.B. and A.S.A.

Software: R.M.B. and A.S.A.

Validation: N.B., F.G.V.S., and M.K.H.

Formal analysis: R.M.B.

Investigation: R.M.B., N.B., M.K.H., and D.L.D.

Resources: N.B., P.S., C.T., M.E., F.G.V.S., M.K.H., J.H., H.L., B.C.K., A.R.B., D.L.D., and

R.M.B.

Data Curation: R.M.B. and A.C.H.

Writing - Original Draft: R.M.B. and E.A.O.

Writing - Review & Editing: R.M.B., E.A.O., and D.L.D.

Visualization: R.M.B.

Supervision: E.A.O.

Project administration: R.M.B. and E.A.O.

Funding acquisition: E.A.O.

References

Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, Kondrashov AS, Sunyaev SR. 2010. A method and server for predicting damaging missense mutations. *Nat Methods* **7**: 248-249.

Akey JM, Ruhe AL, Akey DT, Wong AK, Connelly CF, Madeoy J, Nicholas TJ, Neff MW. 2010. Tracking footprints of artificial selection in the dog genome. *Proc Natl Acad Sci U S A* **107**: 1160-1165.

All of Us Research Program Genomics I. 2024. Genomic data in the All of Us Research Program. *Nature* **627**: 340-346.

Andersson L, Purugganan M. 2022. Molecular genetic variation of animals and plants under domestication. *Proc Natl Acad Sci U S A* **119**: e2122150119.

Arkwright W. 1906. *The Pointer and His Predecessors: An Illustrated History of the Pointing Dog from the Earliest Times*. AL Humphreys.

Axelsson E, Ljungvall I, Bhoumik P, Conn LB, Muren E, Ohlsson A, Olsen LH, Engdahl K, Hagman R, Hanson J et al. 2021. The genetic consequences of dog breed formation-Accumulation of deleterious genetic variation and fixation of mutations associated with myxomatous mitral valve disease in cavalier King Charles spaniels. *PLoS Genet* **17**: e1009726.

Bergeron LA, Besenbacher S, Zheng J, Li P, Bertelsen MF, Quintard B, Hoffman JI, Li Z, St Leger J, Shao C et al. 2023. Evolution of the germline mutation rate across vertebrates. *Nature* **615**: 285-291.

Bouwman AC, Daetwyler HD, Chamberlain AJ, Ponce CH, Sargolzaei M, Schenkel FS, Sahana G, Govignon-Gion A, Boitard S, Dolezal M et al. 2018. Meta-analysis of genome-wide

association studies for cattle stature identifies common genes that regulate body size in mammals. *Nat Genet* **50**: 362-367.

Boyko AR, Quignon P, Li L, Schoenebeck JJ, Degenhardt JD, Lohmueller KE, Zhao K, Brisbin A, Parker HG, vonHoldt BM et al. 2010. A simple genetic architecture underlies morphological variation in dogs. *PLoS Biol* **8**: e1000451.

Buckley RM, Ostrander EA. 2024. Large-scale genomic analysis of the domestic dog informs biological discovery. *Genome Res* doi:10.1101/gr.278569.123.

Bunel M, Chaudieu G, Hamel C, Lagoutte L, Manes G, Botherel N, Brabet P, Pilorge P, Andre C, Quignon P. 2019. Natural models for retinitis pigmentosa: progressive retinal atrophy in dog breeds. *Hum Genet* **138**: 441-453.

Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, Motyer A, Vukcevic D, Delaneau O, O'Connell J et al. 2018. The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**: 203-209.

Cadieu E, Neff MW, Quignon P, Walsh K, Chase K, Parker HG, Vonholdt BM, Rhue A, Boyko A, Byers A et al. 2009. Coat variation in the domestic dog is governed by variants in three genes. *Science* **326**: 150-153.

Cerundolo R, Lloyd DH, Pidduck HG. 1999. Studies on the inheritance of hair loss in the Irish water spaniel. *Vet Rec* **145**: 542-544.

Chen PH, Chen X, He X. 2013. Platelet-derived growth factors and their receptors: structural and functional perspectives. *Biochim Biophys Acta* **1834**: 2176-2186.

Chen Q, Huang B, Zhan J, Wang J, Qu K, Zhang F, Shen J, Jia P, Ning Q, Zhang J et al. 2020. Whole-genome analyses identify loci and selective signals associated with body size in cattle. *J Anim Sci* **98**.

Chen S, Francioli LC, Goodrich JK, Collins RL, Kanai M, Wang Q, Alfoldi J, Watts NA, Vittal C, Gauthier LD et al. 2024. A genomic mutational constraint map using variation in 76,156 human genomes. *Nature* **625**: 92-100.

Cheng J, Novati G, Pan J, Bycroft C, Zemgulyte A, Applebaum T, Pritzel A, Wong LH, Zielinski

M, Sargeant T et al. 2023. Accurate proteome-wide missense variant effect prediction

with AlphaMissense. *Science* **381**: eadg7492.

Christmas MJ, Kaplow IM, Genereux DP, Dong MX, Hughes GM, Li X, Sullivan PF, Hindle AG,

Andrews G, Armstrong JC et al. 2023. Evolutionary constraint and innovation across

hundreds of placental mammals. *Science* **380**: eabn3943.

Cobo AA. 2017. El palacio de Bertemati. Iconografía y posible influencia en otros palacios de la

zona. In *El Barroco: Universo de Experiencias*, pp. 227-248. Asociación Hurtado

Izquierdo.

Cosman D, Wignall J, Anderson D, Tushinski J, Gallis B, Urdal D, Cerretti DP. 1988. Human

macrophage colony stimulating factor (M-CSF): alternate RNA splicing generates three

different proteins that are expressed on the cell surface and secreted. *Behring Inst Mitt*:

15-26.

Dai XM, Ryan GR, Hapel AJ, Dominguez MG, Russell RG, Kapp S, Sylvestre V, Stanley ER.

2002. Targeted disruption of the mouse colony-stimulating factor 1 receptor gene results

in osteopetrosis, mononuclear phagocyte deficiency, increased primitive progenitor cell

frequencies, and reproductive defects. *Blood* **99**: 111-120.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,

Marth GT, Sherry ST et al. 2011. The variant call format and VCFtools. *Bioinformatics*

27: 2156-2158.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,

McCarthy SA, Davies RM, Li H. 2021. Twelve years of SAMtools and BCFtools.

Gigascience **10**.

Donner J, Freyer J, Davison S, Anderson H, Blades M, Honkanen L, Inman L, Brookhart-Knox

CA, Louviere A, Forman OP, Chodroff Foran R. 2023. Genetic prevalence and clinical

relevance of canine Mendelian disease variants in over one million dogs. *PLoS Genet* **19**: e1010651.

Downs LM, Hitti R, Pregnolato S, Mellersh CS. 2014. Genetic screening for PRA-associated mutations in multiple dog breeds shows that PRA is heterogeneous within and between breeds. *Vet Ophthalmol* **17**: 126-130.

Dutrow EV, Serpell JA, Ostrander EA. 2022. Domestic dog lineages reveal genetic drivers of behavioral diversification. *Cell* **185**: 4737-4755 e4718.

Felsenstein J. 1993. PHYLIP (phylogeny inference package), version 3.5 c. *Produced and distributed by author/Department of Genetics, University of Washington*.

Fogle B. 2000. *The new encyclopedia of the dog*. Dorling Kindersley Publishing, Inc., New York.

Fuller ZL, Berg JJ, Mostafavi H, Sella G, Przeworski M. 2019. Measuring intolerance to mutation in human genetics. *Nat Genet* **51**: 772-776.

Gao H, Hamp T, Ede J, Schraiber JG, McRae J, Singer-Berk M, Yang Y, Dietrich ASD, Fiziev PP, Kuderna LFK et al. 2023. The landscape of tolerated genetic variation in humans and primates. *Science* **380**: eabn8153.

Garcia-Pelaez J, Barbosa-Matos R, Lobo S, Dias A, Garrido L, Castedo S, Sousa S, Pinheiro H, Sousa L, Monteiro R et al. 2023. Genotype-first approach to identify associations between CDH1 germline variants and cancer phenotypes: a multicentre study by the European Reference Network on Genetic Tumour Risk Syndromes. *Lancet Oncol* **24**: 91-106.

Garrison E, Kronenberg ZN, Dawson ET, Pedersen BS, Prins P. 2022. A spectrum of free software tools for processing the VCF variant call format: vcflib, bio-vcf, cyvcf2, hts-nim and slivar. *PLoS Comput Biol* **18**: e1009123.

Gerasimavicius L, Livesey BJ, Marsh JA. 2022. Loss-of-function, gain-of-function and dominant-negative mutations have profoundly different effects on protein structure. *Nat Commun* **13**: 3895.

Gough A, Thomas A, O'Neill D. 2018. *Breed predispositions to disease in dogs and cats*. John

Wiley & Sons.

Graber JK, Signer-Hasler H, Burren A, Drogemuller C. 2022. Evaluation of truncating variants in

the LCORL gene in relation to body size of goats from Switzerland. *Anim Genet* **53**: 237-

239.

Gudkov M, Thibaut L, Giannoulatou E. 2024. Quantifying negative selection on synonymous

variants. *HGG Adv* **5**: 100262.

Hagner A, Shin W, Sinha S, Alpaugh W, Workentine M, Abbasi S, Rahmani W, Agabalyan N,

Sharma N, Sparks H et al. 2020. Transcriptional Profiling of the Adult Hair Follicle

Mesenchyme Reveals R-spondin as a Novel Regulator of Dermal Progenitor Function.

iScience **23**: 101019.

Halldorsson BV, Eggertsson HP, Moore KHS, Hauswedell H, Eiriksson O, Ulfarsson MO,

Palsson G, Hardarson MT, Oddsson A, Jensson BO et al. 2022. The sequences of

150,119 genomes in the UK Biobank. *Nature* **607**: 732-740.

Havrilla JM, Pedersen BS, Layer RM, Quinlan AR. 2019. A map of constrained coding regions in

the human genome. *Nat Genet* **51**: 88-95.

Hayward JJ, Castelhano MG, Oliveira KC, Corey E, Balkman C, Baxter TL, Casal ML, Center

SA, Fang M, Garrison SJ et al. 2016. Complex disease and phenotype mapping in the

domestic dog. *Nat Commun* **7**: 10460.

He F, Soriano P. 2013. A critical role for PDGFRalpha signaling in medial nasal process

development. *PLoS Genet* **9**: e1003851.

Hernandez-Ochoa EO, Pratt SJP, Lovering RM, Schneider MF. 2015. Critical Role of

Intracellular RyR1 Calcium Release Channels in Skeletal Muscle Function and Disease.

Front Physiol **6**: 420.

Huang W, Yuan W, Ren L, Liang H, Du X, Sun X, Fang Y, Gao X, Fu M, Sun Y et al. 2022.

Clinicopathological and therapeutic analysis of PDGFRA mutated gastrointestinal

stromal tumor. *Pathol Res Pract* **239**: 154138.

Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alfoldi J, Wang Q, Collins RL, Laricchia

KM, Ganna A, Birnbaum DP et al. 2020. The mutational constraint spectrum quantified

from variation in 141,456 humans. *Nature* **581**: 434-443.

Karlsson EK, Lindblad-Toh K. 2008. Leader of the pack: gene mapping in dogs and other model

organisms. *Nat Rev Genet* **9**: 713-725.

Kim J, Williams FJ, Dreger DL, Plassais J, Davis BW, Parker HG, Ostrander EA. 2018. Genetic

selection of athletic success in sport-hunting dogs. *Proc Natl Acad Sci U S A* **115**:

E7212-E7221.

Lampi S, Donner J, Anderson H, Pohjoismaki J. 2020. Variation in breeding practices and

geographic isolation drive subpopulation differentiation, contributing to the loss of

genetic diversity within dog breed lineages. *Canine Med Genet* **7**: 5.

Landrum MJ, Lee JM, Riley GR, Jang W, Rubinstein WS, Church DM, Maglott DR. 2014.

ClinVar: public archive of relationships among sequence variation and human

phenotype. *Nucleic Acids Res* **42**: D980-985.

Leeb T, Bannasch D, Schoenebeck JJ. 2023. Identification of Genetic Risk Factors for

Monogenic and Complex Canine Diseases. *Annu Rev Anim Biosci* **11**: 183-205.

Leitsalu L, Palover M, Sikka TT, Reigo A, Kals M, Parn K, Nikopensius T, Esko T, Metspalu A,

Padrik P, Tonisson N. 2021. Genotype-first approach to the detection of hereditary breast

and ovarian cancer risk, and effects of risk disclosure to biobank participants. *Eur J Hum*

Genet **29**: 471-481.

Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, O'Donnell-Luria AH, Ware

JS, Hill AJ, Cummings BB et al. 2016. Analysis of protein-coding genetic variation in

60,706 humans. *Nature* **536**: 285-291.

Locher R. 2024. IDPmisc: Utilities of Institute of Data Analyses and

Process Design (www.zhaw.ch/idp).

Lyon GJ, Vedaie M, Beisheim T, Park A, Marchi E, Gottlieb L, Hsieh TC, Klinkhammer H,

Sandomirsky K, Cheng H et al. 2023. Expanding the phenotypic spectrum of NAA10-

related neurodevelopmental syndrome and NAA15-related neurodevelopmental

syndrome. *Eur J Hum Genet* **31**: 824-833.

MacArthur DG, Balasubramanian S, Frankish A, Huang N, Morris J, Walter K, Jostins L,

Habegger L, Pickrell JK, Montgomery SB et al. 2012. A systematic survey of loss-of-

function variants in human protein-coding genes. *Science* **335**: 823-828.

Manley PN, Abu-Abed S, Kirsch R, Hawrysh A, Perrier N, Feilotter H, Pollett A, Riddell RH,

Hookey L, Walia JS. 2018. Familial PDGFRA-mutation syndrome: somatic and

gastrointestinal phenotype. *Hum Pathol* **76**: 52-57.

Marchant TW, Dietschi E, Rytz U, Schawalder P, Jagannathan V, Hadji Rasouliha S, Gurtner C,

Waldvogel AS, Harrington RS, Drogemuller M et al. 2019. An ADAMTS3 missense

variant is associated with Norwich Terrier upper airway syndrome. *PLoS Genet* **15**:

e1008102.

Marchant TW, Johnson EJ, McTeir L, Johnson CI, Gow A, Liuti T, Kuehn D, Svenson K,

Bermingham ML, Drogemuller M et al. 2017. Canine Brachycephaly Is Associated with a

Retrotransposon-Mediated Missplicing of SMOC2. *Curr Biol* **27**: 1573-1584 e1576.

Marks SC, Jr., Lane PW. 1976. Osteopetrosis, a new recessive skeletal mutation on

chromosome 12 of the mouse. *J Hered* **67**: 11-18.

Marsden CD, Ortega-Del Vecchyo D, O'Brien DP, Taylor JF, Ramirez O, Vila C, Marques-Bonet

T, Schnabel RD, Wayne RK, Lohmueller KE. 2016. Bottlenecks and selective sweeps

during domestication have increased deleterious genetic variation in dogs. *Proc Natl*

Acad Sci U S A **113**: 152-157.

McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GR, Thormann A, Flicek P, Cunningham F. 2016. The Ensembl Variant Effect Predictor. *Genome Biol* **17**: 122.

Meadows JRS, Kidd JM, Wang GD, Parker HG, Schall PZ, Bianchi M, Christmas MJ, Bougiouri K, Buckley RM, Hitte C et al. 2023. Genome sequencing of 2000 canids by the Dog10K consortium advances the understanding of demography, genome function and architecture. *Genome Biol* **24**: 187.

Mellanby RJ, Ogden R, Clements DN, French AT, Gow AG, Powell R, Corcoran B, Schoeman JP, Summers KM. 2013. Population structure and genetic heterogeneity in popular dog breeds in the UK. *Vet J* **196**: 92-97.

Metzger J, Schrimpf R, Philipp U, Distl O. 2013. Expression levels of LCORL are associated with body size in horses. *PLoS One* **8**: e56497.

Miyadera K, Acland GM, Aguirre GD. 2012. Genetic and phenotypic variations of inherited retinal diseases in dogs: the power of within- and across-breed studies. *Mamm Genome* **23**: 40-61.

Mooney JA, Yohannes A, Lohmueller KE. 2021. The impact of identity by descent on fitness and disease in dogs. *Proc Natl Acad Sci U S A* **118**.

Morrill K, Hekman J, Li X, McClure J, Logan B, Goodman L, Gao M, Dong Y, Alonso M, Carmichael E et al. 2022. Ancestry-inclusive dog genomics challenges popular breed stereotypes. *Science* **376**: eabk0639.

Nei M, Li WH. 1979. Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc Natl Acad Sci U S A* **76**: 5269-5273.

Ng PC, Henikoff S. 2001. Predicting deleterious amino acid substitutions. *Genome Res* **11**: 863-874.

Nicholas F, Tammen I, Sydney Informatics H. 1995. Online Mendelian Inheritance in Animals (OMIA). doi:10.25910/2amr-pv70. The University of Sydney.

Nicholas FW. 2021. Online Mendelian Inheritance in Animals (OMIA): a record of advances in
    animal genetics, freely available on the Internet for 25 years. *Anim Genet* **52**: 3-9.

Nicholas FW, Tammen, I., Sydney Informatics Hub. 2024. Online Mendelian Inheritance in
    Animals (OMIA). doi:https://doi.org/10.25910/2AMR-PV70.

O'Neill DG, Pegram C, Crocker P, Brodbelt DC, Church DB, Packer RMA. 2020. Unravelling the
    health status of brachycephalic dogs in the UK using multivariable analysis. *Sci Rep* **10**:
    17251.

Ostrander EA, Wang GD, Larson G, vonHoldt BM, Davis BW, Jagannathan V, Hitte C, Wayne
    RK, Zhang YP, Dog KC. 2019. Dog10K: an international sequencing effort to advance
    studies of canine domestication, phenotypes and health. *Natl Sci Rev* **6**: 810-824.

Ostrander EA, Wayne RK, Freedman AH, Davis BW. 2017. Demographic history, selection and
    functional diversity of the canine genome. *Nat Rev Genet* **18**: 705-720.

Özarslan B, Atasoy F, Erdogan M, Özkul BY, Özbaser FT. 2021. Evaluation of some
    morphological and genetic characteristics of Catalburun dog breed. *Turk J Vet Anim Sci*
    45: 649-656.

Park J, Packard EA, Levin MG, Judy RL, Regeneron Genetics C, Damrauer SM, Day SM,
    Ritchie MD, Rader DJ. 2022. A genome-first approach to rare variants in hypertrophic
    cardiomyopathy genes MYBPC3 and MYH7 in a medical biobank. *Hum Mol Genet* **31**:
    827-837.

Parker HG, Dreger DL, Rimbault M, Davis BW, Mullen AB, Carpintero-Ramirez G, Ostrander
    EA. 2017a. Genomic Analyses Reveal the Influence of Geographic Origin, Migration,
    and Hybridization on Modern Dog Breed Development. *Cell Rep* **19**: 697-708.

Parker HG, Harris A, Dreger DL, Davis BW, Ostrander EA. 2017b. The bald and the beautiful:
    hairlessness in domestic dog breeds. *Philos Trans R Soc Lond B Biol Sci* **372**.

Parker HG, Kim LV, Sutter NB, Carlson S, Lorentzen TD, Malek TB, Johnson GS, DeFrance HB,

Ostrander EA, Kruglyak L. 2004. Genetic structure of the purebred domestic dog.

Science **304**: 1160-1164.

Pinero J, Bravo A, Queralt-Rosinach N, Gutierrez-Sacristan A, Deu-Pons J, Centeno E, Garcia-

Garcia J, Sanz F, Furlong LI. 2017. DisGeNET: a comprehensive platform integrating

information on human disease-associated genes and variants. *Nucleic Acids Res* **45**:

D833-D839.

Plassais J, Kim J, Davis BW, Karyadi DM, Hogan AN, Harris AC, Decker B, Parker HG,

Ostrander EA. 2019. Whole genome sequencing of canids reveals genomic regions

under selection and variants influencing morphology. *Nat Commun* **10**: 1489.

Plassais J, vonHoldt BM, Parker HG, Carmagnini A, Dubos N, Papa I, Bevant K, Derrien T,

Hennelly LM, Whitaker DT et al. 2022. Natural and human-driven selection of a single

non-coding body size variant in ancient and modern canids. *Curr Biol* **32**: 889-897 e889.

Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. 2010. Detection of nonneutral substitution

rates on mammalian phylogenies. *Genome Res* **20**: 110-121.

Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Van der Auwera GA, Kling

DE, Gauthier LD, Levy-Moonshine A, Roazen D. 2017. Scaling accurate genetic variant

discovery to tens of thousands of samples. *BioRxiv*: 201178.

Popova SN, Barczyk M, Tiger CF, Beertsen W, Zigrino P, Aszodi A, Miosge N, Forsberg E,

Gullberg D. 2007. Alpha11 beta1 integrin-dependent regulation of periodontal ligament

function in the erupting mouse incisor. *Mol Cell Biol* **27**: 4306-4316.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de

Bakker PI, Daly MJ, Sham PC. 2007. PLINK: a tool set for whole-genome association

and population-based linkage analyses. *Am J Hum Genet* **81**: 559-575.

Qian C, Wong CWY, Wu Z, He Q, Xia H, Tam PKH, Wong KKY, Lui VCH. 2017. Stage specific requirement of platelet-derived growth factor receptor-alpha in embryonic development. *PLoS One* **12**: e0184473.

Rentzsch P, Witten D, Cooper GM, Shendure J, Kircher M. 2019. CADD: predicting the deleteriousness of variants throughout the human genome. *Nucleic Acids Res* **47**: D886-D894.

Rimbault M, Beale HC, Schoenebeck JJ, Hoopes BC, Allen JJ, Kilroy-Glynn P, Wayne RK, Sutter NB, Ostrander EA. 2013. Derived variants at six genes explain nearly half of size reduction in dog breeds. *Genome Res* **23**: 1985-1995.

Rogers CA, Brace AH. 1995. *The international encyclopedia of dogs*. Howell Book House, New York.

Rubin CJ, Megens HJ, Martinez Barrio A, Maqbool K, Sayyab S, Schwochow D, Wang C, Carlborg O, Jern P, Jorgensen CB et al. 2012. Strong signatures of selection in the domestic pig genome. *Proc Natl Acad Sci U S A* **109**: 19529-19536.

Safonov A, Nomakuchi TT, Chao E, Horton C, Dolinsky JS, Yussuf A, Richardson M, Speare V, Li S, Bogus ZC et al. 2023. A genotype-first approach identifies high incidence of NF1 pathogenic variants with distinct disease associations. *medRxiv* doi:10.1101/2023.08.08.23293676.

Sahlen P, Yanhu L, Xu J, Kubinyi E, Wang GD, Savolainen P. 2021. Variants That Differentiate Wolf and Dog Populations Are Enriched in Regulatory Elements. *Genome Biol Evol* **13**.

Saif R, Henkel J, Jagannathan V, Drogemuller C, Flury C, Leeb T. 2020. The LCORL Locus is under Selection in Large-Sized Pakistani Goat Breeds. *Genes (Basel)* **11**.

Samocha KE, Robinson EB, Sanders SJ, Stevens C, Sabo A, McGrath LM, Kosmicki JA, Rehnstrom K, Mallick S, Kirby A et al. 2014. A framework for the interpretation of de novo mutation in human disease. *Nat Genet* **46**: 944-950.

Sarkar A, Panati K, Narala VR. 2022. Code inside the codon: The role of synonymous mutations in regulating splicing machinery and its impact on disease. *Mutat Res Rev Mutat Res* **790**: 108444.

Schoenebeck JJ, Hutchinson SA, Byers A, Beale HC, Carrington B, Faden DL, Rimbault M, Decker B, Kidd JM, Sood R et al. 2012. Variation of BMP3 contributes to dog breed skull diversity. *PLoS Genet* **8**: e1002849.

Sehgal A, Irvine KM, Hume DA. 2021. Functions of macrophage colony-stimulating factor (CSF1) in development, homeostasis, and tissue repair. *Semin Immunol* **54**: 101509.

Stessman HA, Bernier R, Eichler EE. 2014. A genotype-first approach to defining the subtypes of a complex disease. *Cell* **156**: 872-877.

Sugimoto K, Akasaka H, Katsuya T, Node K, Fujisawa T, Shimaoka I, Yasuda O, Ohishi M, Ogihara T, Shimamoto K, Rakugi H. 2008. A polymorphism regulates CYP4A11 transcriptional activity and is associated with hypertension in a Japanese population. *Hypertension* **52**: 1142-1148.

Sullivan PF, Meadows JRS, Gazal S, Phan BN, Li X, Genereux DP, Dong MX, Bianchi M, Andrews G, Sakthikumar S et al. 2023. Leveraging base-pair mammalian constraint to understand genetic variation and human disease. *Science* **380**: eabn2937.

Sun KY, Bai X, Chen S, Bao S, Zhang C, Kapoor M, Backman J, Joseph T, Maxwell E, Mitra G et al. 2024. A deep catalogue of protein-coding variation in 983,578 individuals. *Nature* **631**: 583-592.

Tetens J, Widmann P, Kuhn C, Thaller G. 2013. A genome-wide association study indicates LCORL/NCAPG as a candidate locus for withers height in German Warmblood horses. *Anim Genet* **44**: 467-471.

Van Wesenbeeck L, Odgren PR, MacKay CA, D'Angelo M, Safadi FF, Popoff SN, Van Hul W, Marks SC, Jr. 2002. The osteopetrotic mutation toothless (tl) is a loss-of-function frameshift mutation in the rat Csf1 gene: Evidence of a crucial role for CSF-1 in

osteoclastogenesis and endochondral ossification. *Proc Natl Acad Sci U S A* **99**: 14303-14308.

Vaysse A, Ratnakumar A, Derrien T, Axelsson E, Rosengren Pielberg G, Sigurdsson S, Fall T, Seppala EH, Hansen MS, Lawley CT et al. 2011. Identification of genomic regions associated with phenotypic variation between dog breeds using selection mapping. *PLoS Genet* **7**: e1002316.

Wayne RK, vonHoldt BM. 2012. Evolutionary genomics of dog domestication. *Mamm Genome* **23**: 3-18.

Weir BS, Cockerham CC. 1984. Estimating F-Statistics for the Analysis of Population Structure. *Evolution* **38**: 1358-1370.

Wenger BM, Patel N, Lui M, Moscati A, Do R, Stewart DR, Tartaglia M, Muino-Mosquera L, De Backer J, Kontorovich AR, Gelb BD. 2021. A genotype-first approach to exploring Mendelian cardiovascular traits with clear external manifestations. *Genet Med* **23**: 94-102.

Wilczewski CM, Obasohan J, Paschall JE, Zhang S, Singh S, Maxwell GL, Similuk M, Wolfsberg TG, Turner C, Biesecker LG, Katz AE. 2023. Genotype first: Clinical genomics research through a reverse phenotyping approach. *Am J Hum Genet* **110**: 3-12.

Williams LT. 1989. Signal transduction by the platelet-derived growth factor receptor. *Science* **243**: 1564-1570.

Worboys M, Strange, J.M., Pemberton, N. . 2018. *The invention of the modern dog: breed and blood in Victorian Britain*. Johns Hopkins University Press, Baltimore.

Wright CF, Sharp LN, Jackson L, Murray A, Ware JS, MacArthur DG, Rehm HL, Patel KA, Weedon MN. 2024. Guidance for estimating penetrance of monogenic disease-causing variants in population cohorts. *Nat Genet* doi:10.1038/s41588-024-01842-3.

Xu H, Yan F, Hu R, Suzuki A, Iwaya C, Jia P, Iwata J, Zhao Z. 2021. CleftGeneDB: a resource for annotating genes associated with cleft lip and cleft palate. *Sci Bull (Beijing)* **66**: 2340-2342.

Yang H, Wang G, Wang M, Ma Y, Yin T, Fan R, Wu H, Zhong L, Irwin DM, Zhai W, Zhang Y. 2017. The origin of chow chows in the light of the East Asian breeds. *BMC Genomics* **18**: 174.

Yankee TN, Oh S, Winchester EW, Wilderman A, Robinson K, Gordon T, Rosenfeld JA, VanOudenhove J, Scott DA, Leslie EJ, Cotney J. 2023. Integrative analysis of transcriptome dynamics during human craniofacial development identifies candidate disease genes. *Nat Commun* **14**: 4623.

Yengo L Vedantam S Marouli E Sidorenko J Bartell E Sakaue S Graff M Eliasen AU Jiang Y Raghavan S et al. 2022. A saturated map of common genetic variants associated with human height. *Nature* **610**: 704-712.

Yilmaz O, Ertugrul M, Wilson RT. 2012. The domestic livestock resources of Turkey: breed descriptions and status of guard and hunting dogs. In *63rd Annual Meeting of the EAAP*, pp. 27-31.

Yoshida H, Hayashi S, Kunisada T, Ogawa M, Nishikawa S, Okamura H, Sudo T, Shultz LD, Nishikawa S. 1990. The murine mutation osteopetrosis is in the coding region of the macrophage colony stimulating factor gene. *Nature* **345**: 442-444.

Yu K, Zhang T, Li X. 2018. Genetic role of CYP4A11 polymorphisms in the risk of developing cardiovascular and cerebrovascular diseases. *Ann Hum Genet* **82**: 370-381.

Yu Y, Alvarado R, Petty LE, Bohlender RJ, Shaw DM, Below JE, Bejar N, Ruiz OE, Tandon B, Eisenhoffer GT et al. 2022. Polygenic risk impacts PDGFRA mutation penetrance in non-syndromic cleft lip and palate. *Hum Mol Genet* **31**: 2348-2357.

Zeng T, Spence JP, Mostafavi H, Pritchard JK. 2024. Bayesian estimation of gene constraint from an evolutionary model with gene features. *Nat Genet* **56**: 1632-1643.