Early auditory and adult mating experiences interact with singer identity to shape neural responses to song in female zebra finches

Isabella Catalano[1,2] and Sarah C. Woolley[1,2,3]

[1] Integrated Program in Neuroscience, McGill University, Montreal, QC, Canada

[2] Center for Research on Brain, Language, and Music, Montreal, QC, Canada

[3] Department of Biology, McGill University, Montreal, QC, Canada

ABSTRACT

Social and sensory experiences across the lifespan can shape social interactions, however, experience-dependent plasticity is widely studied within discrete life stages. In the socially monogamous zebra finch, in which females use learned vocal signals to identify individuals and form long-lasting pair bonds, developmental exposure to song is key for females to show species-typical song perception and preferences. While adult mating experience can still lead to pair-bonding and song preference learning even in birds with limited previous song exposure ("song-naïve"), whether similarities in adult behavioral plasticity between normally-reared and song-naïve females reflect convergent patterns of neural activity is unknown. We investigated this using expression of a marker of neural activity and plasticity (phosphorylated S6) in mated normally-reared and song-naïve females in response to song from either their mate, a neighbor, or an unfamiliar male. We found that, in portions of a secondary auditory region (the caudomedial nidopallium NCM) and in dopaminergic neurons of the caudal ventral tegmental area, hearing the mate's song significantly increased pS6 expression in females from both rearing conditions. In contrast, within other NCM subregions, song-identity drove different patterns of pS6 expression depending on the rearing condition. These data suggest that developmental experiences can have long-lasting impacts on the neural signatures of behaviors acquired in adulthood and that socially-driven behavioral plasticity in adults may arise through both shared and divergent neural circuits depending on an individual's developmental experiences.

INTRODUCTION

Early sensory experiences adaptively shape perception as well as the structure and function of neural systems. Developmental sensory experience can have especially long-lasting effects on auditory perception and discrimination. For example, language exposure in infants leads to phonemic tuning whereby infants lose the ability to discriminate between a broad variety of sounds and become tuned to their own language(s) (see Moore & Linthicum, 2007). In oscine songbirds, juvenile exposure to song is similarly important. Classic studies in white-crowned sparrows found that juvenile birds will learn the 'dialect' of song they are exposed to, rather than the one of their genetic parents (Marler & Tamura, 1964). Early exposure to communication signals also tunes neural activity. For example, male zebra finches, who learn song from a tutor, have cells in a secondary auditory cortical region, the caudomedial nidopallium (NCM), that become selective for their tutor's song over the course of development while males raised without tutor exposure do not show similar tuning (Yanagihara & Yazaki-Sugiyama, 2016), and produce species-atypical song in adulthood (Eales, 1985; Jones et al., 1996).

Song perception is also shaped by early exposure to song in female zebra finches. Even though female zebra finches do not learn to sing, juvenile females still form a lasting memory of the tutor's song (Clayton, 1988; Riebel, 2000) and become tuned to the songs or dialect from their own colony (Le Maguer et al., 2021). Additionally, hearing song during development is critical for both auditory discrimination and species-typical song preferences in females. For example, compared to normally-reared females, female zebra finches who are raised without exposure to song ("song-naïve") show impaired frequency discrimination abilities (Sturdy et al., 2001) and do not show species-typical preferences for courtship song over non-courtship song (Chen et al., 2017). Further, developmental experience also impacts the evoked neural responses to preferred and less-preferred songs. Unlike normally-reared females, song-naïve females do not show greater responses to courtship compared to non-courtship songs in either a secondary auditory pallial region (caudomedial nidopallium [NCM], Chen et al., 2017) or in dopaminergic neurons in the caudal ventral tegmental area (cVTA; Barr & Woolley, 2018). Thus, developmental sensory experience can dramatically impact sensory perception and neural responses throughout life.

While experience with conspecific song early in development shapes discrimination and species-typical preferences, adult social experiences can also lead to lasting changes to perception and preference in both song-naïve and normally-reared birds. Among adult songbirds, song is important to determine the species and identity of the singer and, in many species, is used in mate selection (see Nowicki & Searcy, 2005 for review). In zebra finches, males and females form lifelong pair-bonds (Riebel, 2009) and pair-bonded females strongly prefer to hear the song of their partner over the song of an unfamiliar bird (Miller, 1979; Wall & Woolley, 2024; Woolley & Doupe, 2008). Intriguingly, adult pair-bonding experience can drive song preference learning in song-naïve females and even promote species-typical preferences for unfamiliar courtship songs like those seen in normally-reared birds (Wall & Woolley, 2024). However, while song-naïve females form pair-bonds with a partner and preferences for his song, there were important differences. For example, the relationships between pair bonding behaviors and the strength of preference differed between normally-reared and song-naïve females, indicating that there could be differences in the way that birds learn to associate the song with particular social interactions. These data hint that while early exposure to song is not

47 necessary for adult social experiences to drive preference learning, differences in
48 developmental experience could lead to differences in the manner in which these preferences
49 are formed and maintained, perhaps through differential effects on the neural substrates of song
50 preferences.
51
Here, we investigated whether the ability of song-naïve birds to learn preferences for a partner's
songs was associated with similar patterns of activity throughout the brain, especially in the
auditory forebrain and catecholamine producing neurons in the mid- and hindbrain, compared to
normally-reared birds. We quantified expression of an activity-dependent neural marker,
phosphorylated S6 (pS6), in pair-bonded females from both rearing conditions in response to
songs with differing degrees of familiarity. Specifically, within a cohort, all females heard the
same song, but each female had a different relationship to the singer: he was either their mate,
a familiar neighbor, or an unfamiliar male. We found that the impact of rearing condition on the
response to stimulus identity varied across regions, highlighting differential effects of mating
experience on evoked activity depending on an individual's developmental experience.
62
63 RESULTS
64
For female zebra finches, developmental exposure to song is key for tuning auditory responses
and the display of species-typical song preferences. Here, we investigated whether neural
circuits important for learned song preferences in adult females are affected by exposure to
song when birds are juveniles. We created experimental cohorts of normally-reared and song-
naïve females (see Methods for details on rearing). For each cohort in each rearing condition,
we housed a male-female pair in a cage within a sound-attenuating chamber ("soundbox").
Within the same soundbox, a second male-female pair was housed in a different, adjacent
cage. Cages were separated by a visual barrier which allowed birds in each cage to hear but
not see or physically interact with birds in the neighboring cage. We repeated this configuration
across two soundboxes for each rearing condition. After two weeks of cohabitation, females
were housed in isolation and all females within a cohort heard playback of the song of the same
male or silence. Importantly, while all females heard the same song, because of the
experimental design, each bird had a different relationship with the singer: he was either their
partner, an acoustically familiar male (same soundbox), or an unfamiliar male (different
soundbox; Fig. 1). We then quantified expression of pS6 catecholemine-producing cells in the
mid- and hindbrain (caudal and rostral ventral tegmental area [cVTA, rVTA], substantia nigra
pars compacta [SNc], periaqueductal grey [PAG], locus coeruleus [LC]; Fig1e, red in figures), in
ascending (lateral mesencephalic nucleus [MLd], nucleus ovoidalis [Ov]; Fig. 1f, teal), primary
(caudolateral mesopallium [CLM], Fig. 1f, green), and secondary auditory regions (caudomedial
mesopallium [CMM], Fig. 1f, green; caudomedial nidopallium [NCM], Fig. 1f, blue), and two
regions in the social behavior network, the lateral septum (LS) and ventromedial hypothalamus
(VMH; Fig. 1f, orange). We found that the dopamine-producing cVTA as well as the auditory
regions MLd, CMM, and NCM all showed significant increases in pS6 in birds that heard song
compared to silence (cVTA: $X^2_{(1, 39)}$ = 5.3865, p = 0.0203; MLd: $X^2_{(1, 39)}$ = 9.9837, p = 0.0016;

FIGURE 1

Fig.1 Song-evoked pS6 expression. (A) Females were reared either with both parents (normally-reared) or with just their mothers (song-naïve). (B) Then, as adults, females from both rearing conditions were housed for two weeks with a male partner. Two pairs of birds were housed within the same soundbox. (C) After two weeks, all birds were played the song of the same male, but birds differed in their relationship to that male: he was either their mate (red), a familiar neighbor (green) or unfamiliar (blue). A fourth group heard no songs (yellow). (D) Illustrations of sagittal sections of the avian brain (green) indicating the locations of pS6+ cell counts (white boxes) in catecholaminergic, auditory, and social behavior network regions. (E) Box and whisker plots of the normalized percent of tyrosine hydroxylase cells that co-localize with pS6 in catecholamine-positive regions and the (F) normalized density of pS6 expressing cells in auditory and SBN regions. Points are mean counts per individual for normally-reared (closed circles) and song-naïve females (open circles). Each box spans the interquartile range, horizontal white lines indicate the median and whiskers show the minima and maxima. * indicates a difference at p < 0.05.

CMM: $X^2_{(1, 39)}$ = 21.4878, p <0.0001; NCM: $X^2_{(1, 39)}$ = 47.5448, p < 0.0001; all other regions p >

0.13; Fig. 1e-f). We focused on these song-responsive regions for further analysis.

Mate's song increases activation of dopaminergic neurons in females from both rearing

conditions

Within the VTA, we quantified the number of active catecholaminergic neurons by counting both

the number of pS6-immunoreactive cells (IR) and tyrosine hydroxylase immunoreactive cells

(TH-IR), then calculating the percentage of active TH-IR cells (colocalization of TH-IR and pS6-

IR). Based on previous work (Bambico et al., 2019; Barr & Woolley, 2018; Goodson et al., 2009;

Lammel et al., 2008; Matheson & Sakata, 2015), we separated the VTA into rostral (rVTA) and

caudal (cVTA) regions. In the cVTA there was a significant main effect of stimulus, ($X^2_{(3, 39)}$ =

10.4291, p = 0.0152; Fig 2). Specifically, pS6 expression in dopaminergic neurons in the cVTA

was significantly greater in birds that heard their mate's song compared to those that heard an

unfamiliar song (p = 0.0243) or silence (p = 0.0021). Females that heard familiar song had pS6

expression that was intermediate between those that heard mate's song and those that heard

unfamiliar song or silence (p > 0.09 for all comparisons). This effect was consistent across

rearing conditions and parallels the similarity in song preference previously reported between

normally-reared and song-naïve females (Wall & Woolley, 2024).

FIGURE 2

Fig. 2. cVTA results.(A) Images of staining for pS6 in red (top), tyrosine hydroxylase in cyan (middle) and the overlay (bottom) in the cVTA. Arrows highlight cells expressing both pS6 and TH. (B) The percent of TH neurons expressing pS6 was significantly greater in females that heard their mate's song compared to females that heard an unfamiliar song or silence. Points are normalized mean counts per individual for normally-reared (closed circles) and song-naïve females (open circles). * indicates a difference at p < 0.05.

Mate's song drives high levels of pS6 expression across the NCM in normally-reared

females

The NCM is a broad region that encompasses much of the pallium caudal to the primary

auditory region Field L. Previous work has found somewhat uniform induction of immediate

early genes, such as EGR1, throughout the NCM. In contrast, pS6 expression in the NCM has

been described as more irregular, with clusters of pS6+ cells within particular subregions

(Ahmadiantehrani et al., 2018). We found similarly clustered expression of pS6 as has been

previously described. Notably, we saw dense pS6 expression in a dorsocaudal region, an
ventroanterior region, and in a central diagonal band that runs parallel to the lamina
mesopallialis separating the meso and nidopallium and, more laterally, with the border of Field
L3 (see Fig. 3a). We quantified pS6 expression in four locations within these regions that could
be consistently identified using local landmarks in DAPI: one in the dorsocaudal population
(NCMd), one in the ventroanterior region (NCMv), one centrally within the diagonal band of pS6
expression (NCMc), and one at the junction of the hippocampus and posterior NCM (NCMp; see
Fig. 3b). Across all regions, the greatest increase in pS6 expression was generally in response
to the mate's song, while the lowest expression was in silent controls. However, there was also
variation between regions in the level of pS6 induction in the response to stimulus identity ($X^2_{(9,}$
$_{39)}$ = 22.2655, p = 0.0081) as well as significant interactions between stimulus identity and
developmental song exposure ($X^2_{(3, 39)}$ = 21.2266, p <0.0001).
132
In the most dorsal population, NCMd, pS6 expression was modulated by stimulus identity ($X^2_{(3,}$
$_{39)}$ = 51.8815, p <0.0001), and there was a trend towards an interaction of rearing and stimulus
($X^2_{(3, 39)}$ = 7.6967, p = 0.0527; see Fig. 3c). Specifically, while all birds had higher pS6
expression for song playback compared to silence ($X^2_{(1, 39)}$ = 28.2628, p <0.0001), females
varied in the degree to which pS6 expression was modulated by the bird's relationship to the
singer. In normally-reared females, pS6 expression was greatest when the song belonged to the
mate (versus familiar: p < 0.0001; versus unfamiliar: p = 0.0406; versus silence: p < 0.0001).
Expression was also high for the song of an unfamiliar male compared to a familiar song or
silence (p = 0.0109 and p < 0.0001, respectively) while the familiar song did not induce
significant pS6 expression (p = 0.0741 compared to silence). In striking contrast, among song-
naïve birds, all song playbacks induced significant expression of pS6 compared to silence (p <
0.0001 for all); however, there was no significant variation based on the identity of the song (p >
0.5 for all comparisons). A similar pattern of expression was apparent in the NCMp, where there
was an interaction between rearing and stimulus identity ($X^2_{(3, 39)}$ = 9.4253, p = 0.0241; Fig. 3d).
Among normally-reared females, those that heard their mate's song had significantly higher pS6
expression than those that heard familiar song (p = 0.0030), unfamiliar song (p = 0.0252), or
silence (p < 0.0001). Females that heard an unfamiliar song also had greater pS6 expression
than those that heard silence (p = 0.0107), but there was no difference between those that
heard familiar song versus silence (p = 0.0609) or unfamiliar versus familiar song (p = 0.4632).
In contrast, among song-naïve females, there was greater response to song compared to
silence for all stimuli (p < 0.01 for all). Intriguingly, unfamiliar song elicited the greatest increase
in pS6 expression in song-naïve females, which was significantly higher than expression in
response to the mate's song (p = 0.0488) and trending towards significantly higher than
expression in response to a familiar song (p = 0.0550).
157
158   In the central region, NCMc, there was a main effect of stimulus but no differences between the
159   rearing conditions (stimulus: $X^2_{(3, 39)}$ = 10.9736, p = 0.0119; rearing: $X^2_{(1, 39)}$ = 2.5006, p = 0.1138;
160   Fig. 3e). In particular, the response to the mate's song was significantly greater than either
161   unfamiliar song (p = 0.0187) or silence (p = 0.0025). Hearing the familiar song also led to
162

FIGURE 3

Fig. 3. NCM results differ by subregion. (A) Representative photomicrographs of pS6 expression (white) throughout the caudal meso- and nidopallium for all 8 groups. (B) Photomicrograph of the caudal auditory forebrain. White boxes indicate the four regions that were counted within the NCM. (C-E) Box and whisker plots of pS6 expression varied between stimuli and rearing conditions in the (C) dorsal (NCMd), (D) posterior (NCMp), and (E) caudal (NCMc) NCM. (F) There were no significant effects of the stimulus or rearing condition on pS6 expression in the ventral NCM (NCMv). CMM Caudomedial mesopallium; Hp hippocampus; NCMc central caudomedial nidopallium; NCMd dorsal caudomedial nidopallium; NCMp posterior caudomedial nidopallium; NCMv ventral caudomedial nidopallium. Points are mean counts per individual for normally-reared (filled bars in C-D, closed circles in E-F) and song-naïve females (open bars in C-D, open circles in E-F). Each box spans the interquartile range, horizontal white lines indicate the median and whiskers show the minima and maxima. * indicates a difference at p < 0.05. # indicates a difference at p<0.06.

increased pS6 expression compared to silence (p = 0.0483) while hearing unfamiliar song did
not increase pS6 expression over silence (p = 0.4473). Finally, in the NCMv, there were no
significantt effects of rearing, stimulus identity, or their interaction (p > 0.10 for all; Fig. 3f).

Stimulus identity also modulates activity in the ascending auditory system

In the caudomedial mesopallium (CMM) there were, on average, significantly more pS6-positive
cells in normally-reared females compared to song-naïve females ($X^2_{(3, 39)}$ = 5.6584, p = 0.0174;
Fig. 4f). However, pS6 expression in females from both rearing conditions was similarly
modulated by stimulus identity. In particular, pS6 expression was greater in females that heard a
song compared to those that heard silence (all pairwise comparisons p < 0.0005; Fig. 4g). The
increase in pS6 expression was significantly greater in females that heard unfamiliar songs
compared to those that heard a familiar song (p = 0.0481), but not compared to those that heard
the mate's song (p = 0.6863). There was also no significant difference in pS6 expression
between females who heard the mate's song versus familiar song (p = 0.1225). These results
are similar to previously published results where EGR1 expression was greater, though not
significantly, in females that heard unfamiliar songs compared to those that heard the song of a
mate (Woolley & Doupe, 2008).

In the MLd there was no significant effect of rearing ($X^2_{(1, 39)}$ = 1.9019, p = 0.1679); however,
there was a significant effect of stimulus ($X^2_{(3, 39)}$ = 17.5096, p = 0.0006; Fig. 5f). In particular,
pS6 was significantly greater in birds that heard unfamiliar song and the mate's song compared
to silence (both p < 0.005), while, unexpectedly, pS6 was not significantly increased in females
that heard familiar song compared to those exposed to silence (p = 0.1832). In females that
heard familiar song, pS6 expression was significantly lower than in females that heard the
mate's song (p = 0.0090), but not significantly different compared to females that heard an
unfamiliar song (p = 0.3203).

FIGURE 4

Fig. 4. Song-evoked activity in the CMM (A) Widefield view of the caudal auditory forebrain. The white box indicates the location of pS6 counts in the caudomedial mesopallium (CMM). Representative photomicrographs of pS6+ cells (red) in the CMM in (B) silence controls, and birds that heard (C) an unfamiliar song, (D) a familiar neighbor's song, or (E) their mate's song. White lines are 25 μm. (F) Box-and-whisker plots showing a higher density of pS6 expressing cells in the CMM of normally-reared females compared to song-naïve females. (G) Box-and-whisker plots showing song evoked activity in females from both rearing conditions. Points are mean counts per individual for normally-reared (closed circles) and song-naïve females (open circles). Each box spans the interquartile range, horizontal white lines indicate the median and whiskers show the minima and maxima.* indicates a difference at p < 0.05.

FIGURE 5

Fig. 5. Song-evoked activity in the MLd . (A) Widefield view of the mesencephalic nucleus (MLd). Dashed lines indicate the location of pS6 counts. Representative photomicrographs of pS6+ cells (red) in (B) silence controls, and birds that heard (C) an unfamiliar song, (D) a familiar neighbor's song, or (E) their mate's song. White lines are 25 μms. (F) Box-and-whisker plots showing that birds that heard the mate's song and unfamiliar song had greater pS6 expression than those that heard silence. Points are mean counts per individual for normally-reared (closed circles) and song-naïve females (open circles). Each box spans the interquartile range, horizontal white lines indicate the median and whiskers show the minima and maxima.* indicates a difference at p < 0.05.

196
197
198

DISCUSSION

Developmental experiences can shape sensory tuning and have sustained, often interactive effects on adult behaviors (Kundakovic & Champagne, 2015; Prounis et al., 2015). While social interactions in adulthood may ameliorate behavioral deficits that arise from aberrant developmental experience, whether these behavioral similarities arise from similarity in the underlying neural circuits is unclear. In zebra finches, a lack of exposure to conspecific song during development has lasting effects on song perception and neural responses and may define or constrain the way these are shaped by adult social experiences. Here, we tested how developmental experience affects adult neural responses to songs that vary in their social valence or familiarity in areas of the avian brain known to be involved in song identification, learning, and preference. While most catecholaminergic regions examined did not show any significant differences, the caudal VTA showed greater activity in the mate song condition, regardless of rearing, an effect that correlates with previous findings of similarities in pair-bonding behaviors and song preferences. In auditory processing areas such as the MLd and CMM, responses also varied by stimulus but not by rearing condition. In contrast, in a region important for song recognition, learning, and memory, the NCM, there were differences in response across subregions. In a central region of the NCM, both normally-reared and song-naïve females had similar patterns of pS6 expression, with the greatest induction of pS6 in response to the mate's song. In contrast, in two more dorsal subregions there was an interaction of rearing and stimulus: normally-reared females showed greater pS6 expression in response to the mate's song, while the peak pS6 expression in song-naïve females was in response to the unfamiliar song. Taken together, these data highlight that socially-driven behavioral plasticity in adults may arise through both shared and divergent neural circuits depending on an individual's developmental experiences.

The VTA is significant in sexual and partner preferences in a diversity of species (Beny-Shefer et al., 2017; Frye & Paris, 2011; Gossman et al., 2024; Nowicki et al., 2020). In voles, the activation of VTA neurons can lead to the formation of pair bonds even in the absence of mating (Curtis & Wang, 2005). Similarly, in zebra finches, peripheral injections of D2 receptor agonists or D2 antagonists can hasten or inhibit the formation of pair-bonds, respectively (Day et al., 2019). More broadly, activity in the caudal VTA is greater for preferred songs (Barr et al., 2021; Barr & Woolley, 2018). In particular, in normally-reared females, cFOS expression in dopaminergic neurons of the VTA is greater for courtship song compared to non-courtship song as well as for normally-tutored versus experimentally tutored song. Early experience modulates courtship song preference, with less consistent preferences for courtship song among song-naïve females. In addition, song-naïve females also do not show an increase in cFOS expression in the cVTA for courtship song compared to non-courtship song. Together, these data hint at a link between song preferences and the level of activity of dopaminergic neurons in the cVTA. Here, we find that, regardless of rearing condition, activation of DA neurons in the cVTA is greater for the mate's song. Additionally, previous work has also shown that mating ameliorates the preference for courtship song in song-naïve birds (Wall & Woolley, 2024). Whether this means that the activity of the cVTA in response to courtship song in mated, song-naïve females also parallels the activity for normally-reared females is unknown. Future work investigating the degree to which activity in cVTA DA neurons drives or is driven by the formation of the pair bond and the mate's song preference will lend needed insight into the specific role of the cVTA in song preference learning.

246
The NCM is a secondary or associative auditory region that has been implicated in song
recognition and memory. In adult birds, song from a familiar individual, such as a mate or tutor,
or song to which birds were passively exposed through repeated playback induces lower
expression of the immediate early gene EGR1 and lower electrophysiological activity within
subregions of the NCM compared to a novel song (Chew et al., 1995, 1996; Mello et al., 1995;
Mello et al., 1992; Woolley & Doupe, 2008; Woolley & Woolley, 2020). Moreover, NCM lesions
and pharmacological manipulations can impair song recognition (Gobes & Bolhuis, 2007;
Macedo-Lima & Remage-Healey, 2020; Tomaszycki & Blaine, 2014; Yu et al., 2023). During
development, the NCM is implicated in the memorization of the tutor song in juvenile
males. Formation of a tutor song memory is associated with changes to neural tuning and the
emergence of tutor-song selectivity in the NCM (Phan et al., 2006; Yanagihara & Yazaki-
Sugiyama, 2016) and the strength of song learning is correlated with the degree of induction of
immediate early genes in response to the tutor's song (Bolhuis et al., 2000; Bolhuis & Gahr,
2006). Hearing song early in development switches EGR1 expression from constitutively active
to inducible (Kudo et al., 2020; Stripling et al., 2001) and disruption of ERK pathways upstream
from EGR1 affects tutor song memorization in juvenile birds (London & Clayton, 2008). In
female zebra finches, IEG responses in the NCM are also modulated by the attractiveness of
song. Across species, females exhibit greater IEG expression to preferred songs compared to
less preferred songs (Chen et al., 2017; Gentner et al., 2001; Monbureau et al., 2015). For
example, there is greater induction of EGR1 in response to preferred long-bout songs compared
to short-bout songs in females starlings (Gentner et al., 2001) and in response to courtship song
relative to non-courtship song in female zebra finches (Chen et al., 2017). In zebra finches,
silencing or pharmacological manipulation of the NCM alters female song preferences (Barr et
al., 2021; Tomaszycki & Blaine, 2014). We hypothesize that the NCM is important in individual
recognition and preference and could be a locus encoding the mate's song memory. Our data
support a potential role for the NCM in mate's song recognition as we find a number of
subregions within the NCM, especially in the dorsal and rostral regions, that have higher pS6
expression for the mate's song over an unfamiliar song or silence.
275
While our data showing modulation of pS6 expression by singer identity generally support a role
for the NCM in mate's song recognition, we did find region-dependent variation between the
rearing conditions in the strength of response to the mate's song. Developmental song exposure
affects both song preferences and the induction of immediate-early genes in response to
species-typical song variation in females (Chen et al., 2017). However, song-naïve females
learn to prefer the song of a mate in a manner similar to normally-reared females (Wall &
Woolley, 2024). Given the ability of both normally-reared and song-naïve females to recognize
and prefer the mate's song, we expected to find regions of the NCM in which responses to the
mate's song was high for both rearing conditions. In the NCMc, pS6 expression was greatest for
the mate's song in females from both rearing conditions. In contrast, pS6 expression in regions
dorsal to the NCMc, including the NCMp and NCMd, differed between the rearing conditions.
Mate's song induced the greatest pS6 expression in normally-reared females while in song-
naïve females unfamiliar song elicited the greatest pS6 expression. The degree to which this
reflects recognition of the mate's song or a preference for the mate's song is unclear. However,
these data highlight potential topographic differences in the shaping of auditory responses by
developmental song exposure as well as in the processing of the mate's song.
292

In normally-reared females, across a number of NCM subregions, there was significantly greater
pS6 expression for the mate's song compared to the song of a familiar male. Importantly, our
experimental design ensured that, within a cohort of females, one female's mate's song was
simultaneously the familiar song for a female housed in the same soundbox, and an unfamiliar
song for a female in a different soundbox. Thus, the same song was used for all females in a
cohort, only the relationship of each individual bird with the singer of the song differed. This
design accounts for any male-specific variation in song rate during the two weeks of co-
habitation or spectral properties. Females could physically interact with their mate, but could
only hear the familiar male. The difference in pS6 expression suggests that the mate's song is
encoded differently than songs that have been merely heard many times. In particular, it seems
that the social aspect of the pair-bond, and not a mechanism related to habituation to a familiar
stimulus, drives a change in the neural encoding of the mate's song in normally-reared females.
Further investigation of the mechanisms driving memory-formation for a mate's song versus the
habituation to a repeated stimulus will lend needed insight into the mechanisms of auditory
recognition and preference.
308
The NCM is highly dopamine sensitive, with extensive innervation by tyrosine hydroxylase
terminals (Van Ruijssevelt et al., 2018; Von Eugen et al., 2020) coming from the ventral
tegmental area, including the cVTA and substantia nigra pars compacta (Barr et al., 2021) and
high expression of dopamine receptors (Kubikova et al., 2010; Macedo-Lima & Remage-Healey,
2021). Moreover, infusion of dopamine agonists or antagonists during song playback can switch
a female's song preference (Barr et al., 2021). Combined with the results of the current
experiment, dopaminergic inputs from the VTA appear to be important in forming song
preferences, regardless of how the NCM itself processes song. Further research is needed to
confirm whether the VTA neurons that are active when hearing the mate's song are the same
neurons that send projections to NCM, and if so, whether these projections are of equal weight
in song-naïve and normally-reared females.
320
In the MLd there was greater pS6 expression for the mate's song or an unfamiliar song
compared to a familiar song. This result was surprising given previous work that has found that
responses in the MLd vary based on the quantity of sound but not necessarily the
spectrotemporal features or identity of the song (Dai et al., 2018; Schneider & Woolley, 2010;
Woolley & Doupe, 2008). However, unlike the current experiment, in most instances in which
the MLd has been studied, birds were presented with different songs. Here, we used the same
song but varied the categorial relationship between the female and the singer (i.e. the song
could be from a mate or an unfamiliar male). In bats, activity in the inferior colliculus (the
mammalian homologue of the MLd) has been found to vary in response to sounds that differ in
their categorical representation (Lawlor et al., 2023; Salles et al., 2020, 2024). It is also possible
that pS6 highlights a different population of neurons than those that have been previously
investigated with either EGR1 or electrophysiology.
333
The patterns of pS6 induction in both the MLd and the NCM were similar, but not identical to
those seen previously with EGR1 expression in response to mate's song or an unfamiliar song
in normally-reared birds (Chen et al., 2017; Woolley & Doupe, 2008). While both pS6 and
immediate early genes such as EGR1 lie downstream to similar pathways, there may be both
coordination and independence to their activation, as seen with how cFOS, EGR1, and Arc
show both overlapping and distinct patters of expression (Leitner et al., 2005; Sockman et al.,

2005; Velho et al., 2005; Wada et al., 2006). In NCM specifically, EGR1 expression in is greater
in response to a novel song compared to the mate's song (Woolley & Doupe, 2008), which is
contrary to the results we have found in this paper. However, previous work describes pS6
expression in the NCM as clustered and sparse compared to EGR1 expression
(Ahmadiantehrani et al., 2018), an effect that our data replicates. Moreover, we saw stimulus
specific expression patterns that highlight a diagonal band of activity, parallel to Field L3 and the
lamina mesopallialis, especially in females that heard the mate's song. These results provide a
foundation for uncovering the degree to which there is topography or subregion level
organization within the NCM.
349
In the wild, zebra finches live in large, gregarious flocks, and thus are exposed to many
individual songs over the course of their lives. Within those flocks, finches form long-term,
monogamous male-female pairs, with high partner fidelity, and females form strong, lasting
preferences for the song of their mate (Griffith, 2019; Zann, 1996). Our data, finding tuning to
the mate's song in regions of the NCM and VTA fits with the ascendency of pair bonding and
mate preference in the species' life history and ecology (Griffith, 2019). The widespread,
preferential response to the mate's song in the NCM in normally-reared females highlights the
significance of this social relationship and raises interesting questions about the mechanisms by
which the neural signature for the mate's song arises over the course of pair-bonding.
Moreover, while the similarity in neural activation in the NCMc and VTA between normally-
reared and song naïve birds highlights potential subregions that may be important in learning of
a preference for a mate's song, we see lasting evidence of the effects of developmental
differences in sensory experience in other NCM subregions, which suggest differences in the
neuronal encoding mechanisms of learned song in adulthood that is dependent upon the song
exposure or lack thereof received during development. Taken together, these highlight the
importance of early life experience in shaping neural functioning throughout adulthood.
366
367   METHODS
368
369   *Animals*
370
371   Zebra finches (N = 67, n = 39 females, n = 28 males, all >90 days post-hatch) had *ad libitum*
372   access to seed, water, and grit and were kept on a 14:10 light:dark cycle. Egg and egg
373   supplements were provided twice a week. Bird care and experimental procedures were
374   approved by the McGill University Animal Care Committee and were performed in accordance
375   with the Canadian Council on Animal Care guidelines.
376
377   Birds were reared in one of two conditions. Normally-reared birds were reared with both parents
378   and were exposed to the songs of other males in a colony setting, while song-naïve birds had
379   their father removed 5-7 dph, and were reared by their mother only in sound-attenuating boxes
380   (TRA Acoustics, Cornwell, Ontario). Song-naïve birds remained in the nest until 60 dph, at which
381   point they were moved to a female-only colony setting.
382
383   *Experimental Design*
384
385   Female birds were housed with a male in a cage (10 in. x 10 in.) within a sound-attenuating
386   chamber ("soundbox"). Within the same soundbox, a second male-female pair was housed in a

different, adjacent cage. Cages were separated by a visual barrier which allowed birds in each cage to hear but not see or physically interact with birds in the neighboring cage. All pairs were given access to nesting material and an empty dish in which to nest. This setup was repeated across two soundboxes for each rearing condition Thus, for each experimental "batch" there were a total of four soundboxes containing a total of eight pairs of mated birds (four with song-naïve females, four with normally-reared females). We created 5 experimental batches (n = 20 total normally-reared females and n = 19 total song-naïve females). In one exceptional instance, a silent control female was not mated. Birds were housed together for two weeks, which previous research indicates is sufficient time to form a pair bond (Wall & Woolley, 2024).

Stimulus creation

Prior to mating, males were recorded singing both directed (female-oriented) and undirected song using Sound Analysis Pro (SAP; Tchernichovski et al., 2000) as previously described (Chen et al., 2017; Schubloom & Woolley, 2016; Wall & Woolley, 2024; Woolley & Doupe, 2008). Briefly, directed song was elicited using stimulus females (not used in the study), who were presented to the male for 30 seconds and then removed. Recorded song was then bandpass filtered (300-10 kHz), normalized by the maximum amplitude, and saved as a .wav file (44.1 kHz) using custom Matlab code (Mathworks, Natick, Massachusetts, USA). On average 9 song clips (range 8-10) were selected per male to be used as stimuli. Because within a batch, song-naïve and normally-reared females heard different males, males with similar song durations and numbers of motifs were selected to ensure similar quantities of sound were heard across rearing conditions.

Stimulus playback

Female birds were separated from their mates overnight. The following morning, lights went on for 30 minutes, beginning one hour and 45 minutes prior to perfusion. Lights were then turned off, and 30 minutes later stimuli were played for the following 30 minutes, also in the dark. Birds were perfused fifteen minutes after playback of the last stimulus.

Birds were organized into 5 immunocytochemical batches, with each batch containing females from both rearing conditions (normally-reared and song-naïve) and all stimulus conditions (mate, familiar, unfamiliar, silence). Specifically, stimuli were such that for each set of four normally-reared and song-naïve females, the song from one male was chosen as the stimulus. This male's song was played back to three females who each had a different relationship with the male. For one female, the song was from her mate ('Mate's song'), for one female the song was from the bird in the adjacent cage within the same sound-attenuating box ('Familiar song'), and for one bird the song was completely unfamiliar, from a bird in a different sound-attenuating box ('Unfamiliar song'). The fourth bird in each batch was a silent control, and received silence in place of sound playback.

Immunocytochemistry and Imaging

After stimulus playback, females were acutely anesthetized via isoflurane inhalation and then transcardially perfused with 12mg heparin in 0.9% saline, followed by 150mL of 4%

434 paraformaldehyde. Brains were removed and stored overnight in 4% paraformaldehyde, after
435 which they were transferred to 30% sucrose for cryoprotection. Brains were sliced in 40 μm
436 sagittal sections and kept in 0.025M phosphate-buffered saline (PBS) with sodium azide before
437 further procedures.
438
We performed double-label immunocytochemistry to antibodies for phosphorylated S6 (pS6) as
a marker of neural activity, and tyrosine hydroxylase (TH) as a marker of catecholaminergic
neurons as described previously (Barr & Woolley, 2018; Chen et al., 2017; Fan et al.,
2022). Reactions were performed in five immunocytochemical batches with all rearing
conditions and stimulus groups represented in each batch, using every third section for each
bird. In brief, sections were washed 3x10 minutes with 0.025M PBS, then transferred to blocking
for one hour in 5% donkey serum and 0.3% Triton X-100. After blocking, sections were
incubated for 48 hours in primary antibody: rabbit-anti-pS6 (1:500 dilution, Cell Signalling
Technologies, Danvers, Massachusetts, USA) and sheep-anti-TH (1:1000 dilution, Novus
Biologicals, Toronto, Ontario, Canada) at 4° C. After 48 hours, slices were once again washed
2x 30 minutes in 0.025M PBS, and then incubated at room temperature in the secondary
antibody: donkey anti-rabbit conjugated to Alexa Fluor 594 (Life Technologies, Eugene, Oregon,
USA) and donkey anti-sheep conjugated to Alexa Fluor 488 (Life Technologies). Finally, the
sections were washed 3x10 minutes in 0.025 PBS and stored in 0.025 PBS. Sections were
mounted on chromium-aluminum subbed glass slides and cover slipped using ProLong Gold
Antifade Reagent (Life Sciences, Burlington, Ontario, Canada).
455
Stained tissue was imaged using Zen microscopy software (Carl Zeiss, Germany) with a 40x
objective on a Zeiss Axio Imager upright microscope and an AxioCam MRm Zeiss camera (Carl
Zeiss, Germany), using 594 and 488 channels. Exposure times were calculated by auto-
determining the exposure of individual areas through Zen, and then averaging the exposure
number of each brain region per batch. Exposures > 1000s were removed from the average to
avoid overexposing sections that contained large amounts of fluorescence.
462
Each section was imaged using TH as an anatomical reference (see Fig. 1d), and each brain
region was imaged in both left and right hemispheres. We imaged five auditory regions, the
caudomedial nidopallium (NCM), caudomedial mesopallium (CMM), caudolateral mesopallium
(CLM), the nucleus ovoidalis (Ov), and the nucleus mesencephalicus lateralis pars dorsalis
(MLd), and five catecholamine producing regions, the ventral tegmental area (VTA), substantia
nigra pars compacta (SN), periaqueductal gray (PAG), and locus coeruleus (LC), as well as the
lateral septum (LS) and ventromedial hypothalamus (VMH). The NCM was imaged four times
per slice (see Fig. 3b; avg 4.6, range 2-11 sections per region per bird), starting from the most
medial slice on which tissue in that region was visible. The most dorsal region was imaged at
the very top of NCM (NCMd), the next at the very bottom (NCMv), and the third in the center
(NCMc). All of these had one image at 20x zoom per slice, two images per bird. NCMp was
imaged as two parts at 40x zoom, the first from where the hippocampus met the NCM. The
viewfield was then moved rostrally (towards Field L) until there was no overlap with the previous
part and a second part was imaged. Both of these images were counted separately, but
analyzed as belonging to NCMp. All subsequent regions were imaged once per slice at 40x
zoom. The CMM was imaged where Field L met a ventricle under the hippocampus and formed
a triangular shape (avg. 8.6, range 4-11 sections per bird). The CLM was imaged approximately
1.5mm lateral from the midpoint, under the hippocampus and above field L (avg. 7.5, range 5-10

per bird). Ovoidalis was located using autofluorescence to identify the nucleus, approximately 1-
1.2mm lateral (avg. 3.9, range 2-4 sections per bird). The MLd was imaged approximately
2.5mm laterally (avg. 7.1, range 4-10 sections per bird). The VTA was imaged twice, in distinct
subregions. The first, the rostral VTA, was imaged when the VTA met the edge of the tissue
(avg. 4.3, range 2-6 sections per bird). The caudal VTA was measured at the opposite end of
the VTA but on the same slices, where it narrowed into the rest of the midbrain (avg. 4.3, range
2-6 sections per bird). The SNc was imaged more laterally than the VTA, when it starts to look
rounded on one end and smeared on the other, central in the midbrain (avg. 6.0, range 4-8
sections per bird). The PAG was imaged in roughly the same slices in which the VTA was
imaged, along the caudal edge of the midbrain where a ventricle separates the hippocampus
(avg. 4.2, range 3-5 sections per bird). The LC was imaged slightly rostral to where the
hippocampus joined the midbrain, and identified by a bright cluster of cells (avg. 4.2, range 2-8
sections per bird). The LS was imaged medially, under Field L when it formed a curved or bump
shape (avg. 2.8, range 1-4 sections per bird). Finally, the VMH was imaged medially, behind the
optic chiasm (avg. 3.1, range 1-4 sections per bird).
496
Individual cells were counted from raw images using FIJI/ImageJ imaging software (Schindelin
et al., 2012; Schneider et al., 2012) by individuals blinded to the conditions of the study. For
double-labelled images, cells were counted first on one stain, then separately on the other,
before being overlaid to count how many cells were marked on both layers. Counts were then
normalized by batch for each area. Each immunocytochemical batch, described in more detail
above, consisted of eight birds including females from both rearing conditions (normally-reared
and song-naïve) and all stimulus conditions (mate, familiar, unfamiliar, silence). For pS6, the
number of cells per image was divided by the size (area) of the image to obtain a count
normalized by area. Then, these densities were averaged for each batch, and then the density
for each individual image was divided by the batch average. We then took the mean normalized
density for each individual. For TH-IR areas, the percentage of active TH cells on each
individual image was divided by the batch average of percent active cells. We then took the
mean normalized percentage for each individual.
510
511   Images used in paper figures were edited for easier visual contrast, and composite images were
512   formed by taking multiple 5x-zoom images and digitally stitching them together to form a
513   complete larger map. However, as mentioned, all data counting used raw images directly with
514   no photomanipulation.
515
516   *Statistical analysis*
517
Analysis was conducted using JMP Statistical Processing Software (SAS Institute, Cary, North
Carolina, USA). We calculated the density of neurons for each TH- region and batch-normalized
the density values as described above to account for variations in immunohistochemical
analysis. For TH+ regions, the percentage of TH+ cells colocalizing pS6 was calculated. We
performed a Poisson-distributed generalized linear model (GLM) with Log link function. Rearing,
stimulus, and their interaction served as independent variables. We used least-squared Tukey's
HSD for all post-hoc tests and set $\alpha < 0.05$ for all tests unless otherwise noted.
525
526
527

BIBLIOGRAPHY

Ahmadiantehrani, S., Gores, E. O., & London, S. E. (2018). A complex mTOR response in

   habituation paradigms for a social signal in adult songbirds. *Learning & Memory*, *25*(6),

   273–282.

Bambico, F. R., Li, Z., Oliveira, C., Mcneill, S., Diwan, M., Raymond, R., & Nobrega, J. N.

   (2019). Rostrocaudal subregions of the ventral tegmental area are differentially impacted

   by chronic stress. *Psychopharmacology*, *236*(6), 1917–1929.

   https://doi.org/10.1007/s00213-019-5177-8

Barr, H. J., Wall, E. M., & Woolley, S. C. (2021). Dopamine in the songbird auditory cortex

   shapes auditory preference. *Current Biology*, *31*(20), 4547-4559.e5.

   https://doi.org/10.1016/j.cub.2021.08.005

Barr, H. J., & Woolley, S. C. (2018). Developmental auditory exposure shapes responses of

   catecholaminergic neurons to socially-modulated song. *Scientific Reports*, *8*(1), 11717.

   https://doi.org/10.1038/s41598-018-30039-y

Beny-Shefer, Y., Zilkha, N., Lavi-Avnon, Y., Bezalel, N., Rogachev, I., Brandis, A., Dayan, M., &

   Kimchi, T. (2017). Nucleus Accumbens Dopamine Signaling Regulates Sexual

   Preference for Females in Male Mice. *Cell Reports*, *21*(11), 3079–3088.

   https://doi.org/10.1016/j.celrep.2017.11.062

Bolhuis, J. J., & Gahr, M. (2006). Neural mechanisms of birdsong memory. *Nature Reviews

   Neuroscience*, *7*(5), 347–357. https://doi.org/10.1038/nrn1904

Bolhuis, J. J., Zijlstra, G. G. O., Den Boer-Visser, A. M., & Van Der Zee, E. A. (2000). Localized

   neuronal activation in the zebra finch brain is related to the strength of song learning.

   Proceedings of the National Academy of Sciences, *97*(5), 2282–2285.

   https://doi.org/10.1073/pnas.030539097

Chen, Y., Clark, O., & Woolley, S. C. (2017). Courtship song preferences in female zebra

   finches are shaped by developmental auditory experience. *Proceedings of the Royal*

Society B: Biological Sciences, *284*(1855), 20170054.

https://doi.org/10.1098/rspb.2017.0054

Chew, S. J., Mello, C., Nottebohm, F., Jarvis, E., & Vicario, D. S. (1995). Decrements in auditory

responses to a repeated conspecific song are long-lasting and require two periods of

protein synthesis in the songbird forebrain. *Proceedings of the National Academy of*

Sciences, *92*(8), 3406–3410. https://doi.org/10.1073/pnas.92.8.3406

Chew, S. J., Vicario, D. S., & Nottebohm, F. (1996). A large-capacity memory system that

recognizes the calls and songs of individual birds. *Proceedings of the National Academy*

of Sciences, *93*(5), 1950–1955. https://doi.org/10.1073/pnas.93.5.1950

Clayton, N. S. (1988). Song discrimination learning in zebra finches. *Animal Behaviour*, *36*(4),

1016–1024. https://doi.org/10.1016/S0003-3472(88)80061-7

Curtis, J., & Wang, Z. (2005). Ventral tegmental area involvement in pair bonding in male prairie

voles. *Physiology & Behavior*, *86*(3), 338–346.

https://doi.org/10.1016/j.physbeh.2005.08.022

Dai, J. B., Chen, Y., & Sakata, J. T. (2018). EGR-1 Expression in Catecholamine-synthesizing

Neurons Reflects Auditory Learning and Correlates with Responses in Auditory

Processing Areas. *Neuroscience*, *379*, 415–427.

https://doi.org/10.1016/j.neuroscience.2018.03.032

Day, N. F., Saxon, D., Robbins, A., Harris, L., Nee, E., Shroff-Mehta, N., Stout, K., Sun, J., Lillie,

N., Burns, M., Korn, C., & Coleman, M. J. (2019). D2 dopamine receptor activation

induces female preference for male song in the monogamous zebra finch. *Journal of*

Experimental Biology, *222*(5). https://doi.org/10.1242/jeb.191510

Eales, L. A. (1985). Song learning in zebra finches: Some effects of song model availability on

what is learnt and when. *Animal Behaviour*, *33*(4), 1293–1300.

https://doi.org/10.1016/S0003-3472(85)80189-5

Fan, R., Reader, S. M., & Sakata, J. T. (2022). Alarm cues and alarmed conspecifics: Neural

    activity during social learning from different cues in Trinidadian guppies. *Proceedings of*

    the Royal Society B: Biological Sciences, *289*(1981), 20220829.

    https://doi.org/10.1098/rspb.2022.0829

Frye, C. A., & Paris, J. J. (2011). Effects of neurosteroid actions at N-methyl-d-aspartate and

    GABAA receptors in the midbrain ventral tegmental area for anxiety-like and mating

    behavior of female rats. *Psychopharmacology*, *213*(1), 93–103.

    https://doi.org/10.1007/s00213-010-2016-3

Gentner, T. Q., Hulse, S. H., Duffy, D., & Ball, G. F. (2001). Response biases in auditory

    forebrain regions of female songbirds following exposure to sexually relevant variation in

    male song. *Journal of Neurobiology*, *46*(1), 48–58. https://doi.org/10.1002/1097-

    4695(200101)46:1<48::AID-NEU5>3.0.CO;2-3

Gobes, S. M. H., & Bolhuis, J. J. (2007). Birdsong Memory: A Neural Dissociation between Song

    Recognition and Production. *Current Biology*, *17*(9), 789–793.

    https://doi.org/10.1016/j.cub.2007.03.059

Goodson, J. L., Kabelik, D., Kelly, A. M., Rinaldi, J., & Klatt, J. D. (2009). Midbrain dopamine

    neurons reflect affiliation phenotypes in finches and are tightly coupled to courtship.

    Proceedings of the National Academy of Sciences, *106*(21), 8737–8742.

    https://doi.org/10.1073/pnas.0811821106

Gossman, K. R., Lowe, C. S., Kirckof, A., Vanmeerhaeghe, S., & Smith, A. S. (2024).

    Corticotropin-releasing factor and GABA in the ventral tegmental area modulate partner

    preference formation in male and female prairie voles (Microtus ochrogaster). *Frontiers*

    in Neuroscience, *18*. https://doi.org/10.3389/fnins.2024.1430447

Griffith, S. C. (2019). Cooperation and Coordination in Socially Monogamous Birds: Moving

    Away From a Focus on Sexual Conflict. *Frontiers in Ecology and Evolution*, *7*.

    https://doi.org/10.3389/fevo.2019.00455

Jones, A. E., ten Cate, C., & Slater, P. J. B. (1996). Early experience and plasticity of song in adult male zebra finches (Taeniopygia guttata). *Journal of Comparative Psychology*, *110*(4), 354–369. https://doi.org/10.1037/0735-7036.110.4.354

Kubikova, L., Wada, K., & Jarvis, E. D. (2010). Dopamine receptors in a songbird brain. *Journal of Comparative Neurology*, *518*(6), 741–769. https://doi.org/10.1002/cne.22255

Kudo, T., Morohashi, Y., & Yazaki-Sugiyama, Y. (2020). Early Auditory Experience Modifies Neuronal Firing Properties in the Zebra Finch Auditory Cortex. *Frontiers in Neural Circuits*, *14*. https://doi.org/10.3389/fncir.2020.570174

Kundakovic, M., & Champagne, F. A. (2015). Early-Life Experience, Epigenetics, and the Developing Brain. *Neuropsychopharmacology*, *40*(1), 141–153. https://doi.org/10.1038/npp.2014.140

Lammel, S., Hetzel, A., Häckel, O., Jones, I., Liss, B., & Roeper, J. (2008). Unique properties of mesoprefrontal neurons within a dual mesocorticolimbic dopamine system. *Neuron*, *57*(5), 760–773.

Lawlor, J., Wohlgemuth, M. J., Moss, C. F., & Kuchibhotla, K. V. (2023). Spatially clustered neurons encode vocalization categories in the bat midbrain. *bioRxiv*.

Le Maguer, L., Derégnaucourt, S., & Geberzahn, N. (2021). Female preference for artificial song dialects in the zebra finch (Taeniopygia guttata). *Ethology*, *127*(7), 537–549. https://doi.org/10.1111/eth.13159

Leitner, S., Voigt, C., Metzdorf, R., & Catchpole, C. K. (2005). Immediate early gene (ZENK, Arc) expression in the auditory forebrain of female canaries varies in response to male song quality. *Journal of Neurobiology*, *64*(3), 275–284. https://doi.org/10.1002/neu.20135

London, S. E., & Clayton, D. F. (2008). Functional identification of sensory mechanisms required for developmental song learning. *Nature Neuroscience*, *11*(5), 579–586. https://doi.org/10.1038/nn.2103

Macedo-Lima, M., & Remage-Healey, L. (2020). Auditory learning in an operant task with social

reinforcement is dependent on neuroestrogen synthesis in the male songbird auditory

cortex. *Hormones and Behavior*, *121*, 104713.

https://doi.org/10.1016/j.yhbeh.2020.104713

Macedo-Lima, M., & Remage-Healey, L. (2021). Dopamine Modulation of Motor and Sensory

Cortical Plasticity among Vertebrates. *Integrative and Comparative Biology*, *icab019*.

https://doi.org/10.1093/icb/icab019

Marler, P., & Tamura, M. (1964). Culturally Transmitted Patterns of Vocal Behavior in Sparrows.

Science, *146*(3650), 1483–1486. https://doi.org/10.1126/science.146.3650.1483

Matheson, L. E., & Sakata, J. T. (2015). Catecholaminergic contributions to vocal

communication signals. *European Journal of Neuroscience*, *41*(9), 1180–1194.

https://doi.org/10.1111/ejn.12885

Mello, C., Nottebohm, F., & Clayton, D. (1995). Repeated exposure to one song leads to a rapid

and persistent decline in an immediate early gene's response to that song in zebra finch

telencephalon. *The Journal of Neuroscience*, *15*(10), 6919–6925.

https://doi.org/10.1523/jneurosci.15-10-06919.1995

Mello, C. V., Vicario, D. S., & Clayton, D. F. (1992). Song presentation induces gene expression

in the songbird forebrain. *Proceedings of the National Academy of Sciences*, *89*(15),

6818–6822.

Miller, D. B. (1979). The acoustic basis of mate recognition by female Zebra finches

(Taeniopygia guttata). *Animal Behaviour*, *27*, 376–380. https://doi.org/10.1016/0003-

3472(79)90172-6

Monbureau, M., Barker, J. M., Leboucher, G., & Balthazart, J. (2015). Male song quality

modulates c-Fos expression in the auditory forebrain of the female canary. *Physiology &*

Behavior, *147*, 7–15. https://doi.org/10.1016/j.physbeh.2015.04.005

Moore, J. K., & Linthicum, F. H. (2007). The human auditory system: A timeline of development.

International Journal of Audiology, *46*(9), 460–478.

https://doi.org/10.1080/14992020701383019

Nowicki, J. P., Pratchett, M. S., Walker, S. P. W., Coker, D. J., & O'Connell, L. A. (2020). Gene

expression correlates of social evolution in coral reef butterflyfishes. *Proceedings of the*

Royal Society B: Biological Sciences, *287*(1929), 20200239.

https://doi.org/10.1098/rspb.2020.0239

Nowicki, S., & Searcy, W. A. (2005). Song and Mate Choice in Birds: How The Development of

Behavior Helps us Understand Function. *The Auk*, *122*(1), 1–14.

https://doi.org/10.1093/auk/122.1.1

Phan, M. L., Pytte, C. L., & Vicario, D. S. (2006). Early auditory experience generates long-

lasting memories that may subserve vocal learning in songbirds. *Proceedings of the*

National Academy of Sciences, *103*(4), 1088–1093.

https://doi.org/10.1073/pnas.0510136103

Prounis, G. S., Foley, L., Rehman, A., & Ophir, A. G. (2015). Perinatal and juvenile social

environments interact to shape cognitive behaviour and neural phenotype in prairie

voles. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1819), 20152236.

https://doi.org/10.1098/rspb.2015.2236

Riebel, K. (2000). Early exposure leads to repeatable preferences for male song in female zebra

finches. *Proceedings of the Royal Society of London. Series B: Biological Sciences*,

267(1461), 2553–2558. https://doi.org/10.1098/rspb.2000.1320

Riebel, K. (2009). Chapter 6 Song and Female Mate Choice in Zebra Finches: A Review. In

Advances in the Study of Behavior (Vol. 40, pp. 197–238). Academic Press.

https://doi.org/10.1016/S0065-3454(09)40006-8

Salles, A., Loscalzo, E., Montoya, J., Mendoza, R., Boergens, K. M., & Moss, C. F. (2024).

Auditory processing of communication calls in interacting bats. *iScience*, *27*(6).

https://doi.org/10.1016/j.isci.2024.109872

Salles, A., Park, S., Sundar, H., Macías, S., Elhilali, M., & Moss, C. F. (2020). Neural Response

Selectivity to Natural Sounds in the Bat Midbrain. *Neuroscience*, *434*, 200–211.

https://doi.org/10.1016/j.neuroscience.2019.11.047

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,

Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri,

K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source platform for biological-

image analysis. *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of

image analysis. *Nature Methods*, *9*(7), 671–675. https://doi.org/10.1038/nmeth.2089

Schneider, D. M., & Woolley, S. M. N. (2010). Discrimination of Communication Vocalizations by

Single Neurons and Groups of Neurons in the Auditory Midbrain. *Journal of*

Neurophysiology, *103*(6), 3248–3265. https://doi.org/10.1152/jn.01131.2009

Schubloom, H. E., & Woolley, S. C. (2016). Variation in social relationships relates to song

preferences and EGR1 expression in a female songbird. *Developmental Neurobiology*,

76(9), 1029–1040. https://doi.org/10.1002/dneu.22373

Sockman, K. W., Gentner, T. Q., & Ball, G. F. (2005). Complementary neural systems for the

experience-dependent integration of mate-choice cues in European starlings. *Journal of*

Neurobiology, *62*(1), 72–81. https://doi.org/10.1002/neu.20068

Stripling, R., Kruse, A. A., & Clayton, D. F. (2001). Development of song responses in the zebra

finch caudomedial neostriatum: Role of genomic and electrophysiological activities.

Journal of Neurobiology, *48*(3), 163–180. https://doi.org/10.1002/neu.1049

Sturdy, C. B., Phillmore, L. S., Sartor, J. J., & Weisman, R. G. (2001). Reduced social contact

causes auditory perceptual deficits in zebra finches, Taeniopygia guttata. *Animal*

Behaviour, *62*(6), 1207–1218. https://doi.org/10.1006/anbe.2001.1864

Tchernichovski, O., Nottebohm, F., Ho, C. E., Pesaran, B., & Mitra, P. P. (2000). A procedure for

an automated measurement of song similarity. *Animal Behaviour*, *59*(6), 1167–1176.

https://doi.org/10.1006/anbe.1999.1416

Tomaszycki, M. L., & Blaine, S. K. (2014). Temporary inactivation of NCM, an auditory region,

increases social interaction and decreases song perception in female zebra finches.

Behavioural Processes, *108*, 65–70. https://doi.org/10.1016/j.beproc.2014.09.031

Van Ruijssevelt, L., Chen, Y., Von Eugen, K., Hamaide, J., De Groof, G., Verhoye, M.,

Güntürkün, O., Woolley, S. C., & Van Der Linden, A. (2018). fMRI Reveals a Novel

Region for Evaluating Acoustic Information for Mate Choice in a Female Songbird.

Current Biology, *28*(5), 711-721.e6. https://doi.org/10.1016/j.cub.2018.01.048

Velho, T. A. F., Pinaud, R., Rodrigues, P. V., & Mello, C. V. (2005). Co-induction of activity-

dependent genes in songbirds. *European Journal of Neuroscience*, *22*(7), 1667–1678.

https://doi.org/10.1111/j.1460-9568.2005.04369.x

Von Eugen, K., Tabrik, S., Güntürkün, O., & Ströckens, F. (2020). A comparative analysis of the

dopaminergic innervation of the executive caudal nidopallium in pigeon, chicken, zebra

finch, and carrion crow. *Journal of Comparative Neurology*, *528*(17), 2929–2955.

https://doi.org/10.1002/cne.24878

Wada, K., Howard, J. T., McConnell, P., Whitney, O., Lints, T., Rivas, M. V., Horita, H., Patterson,

M. A., White, S. A., Scharff, C., Haesler, S., Zhao, S., Sakaguchi, H., Hagiwara, M.,

Shiraki, T., Hirozane-Kishikawa, T., Skene, P., Hayashizaki, Y., Carninci, P., & Jarvis, E.

D. (2006). A molecular neuroethological approach for identifying and characterizing a

cascade of behaviorally regulated genes. *Proceedings of the National Academy of*

Sciences, *103*(41), 15212–15217. https://doi.org/10.1073/pnas.0607098103

Wall, E. M., & Woolley, S. C. (2024). Social experiences shape song preference learning

independently of developmental exposure to song. *Proceedings of the Royal Society B:*

Biological Sciences, *291*(2024). https://doi.org/10.1098/rspb.2024.0358

Woolley, S. C., & Doupe, A. J. (2008). Social Context–Induced Song Variation Affects Female

Behavior and Gene Expression. *PLOS Biology*, *6*(3), e62.

https://doi.org/10.1371/journal.pbio.0060062

Woolley, S. C., & Woolley, S. M. (2020). Integrating Form and Function in the Songbird Auditory

Forebrain. In *J. T. Sakata, S. C. Woolley, R. R. Fay, & A. N. Popper (Eds.), The*

Neuroethology of Birdsong (p. (pp. 127-155)). Springer International Publishing.

Yanagihara, S., & Yazaki-Sugiyama, Y. (2016). Auditory experience-dependent cortical circuit

shaping for memory formation in bird song learning. *Nature Communications*, *7*(1),

11946. https://doi.org/10.1038/ncomms11946

Yu, K., Wood, W. E., Johnston, L. G., & Theunissen, F. E. (2023). Lesions to Caudomedial

Nidopallium Impair Individual Vocal Recognition in the Zebra Finch. *The Journal of*

Neuroscience, *43*(14), 2579–2596. https://doi.org/10.1523/jneurosci.0643-22.2023

Zann, R. A. (1996). *The Zebra Finch: A Synthesis of Field and Laboratory Studies*. Oxford

University Press.