Transcriptomic Insights into Hypothalamic Aging During Menopause: A Comparative

Analysis of Human and Mouse Models.

Authors: Jordana C. B. Bloom[1,2], Encarnacion Torres[2,3*], Sidney A. Pereira[2,3*], Liliana Arvizu-

Sanchez[2], Audrey N. Fontes[2], Hadine Joffe[3,4,5], David C. Page[1,6,7], Victor M. Navarro[2,3,8]

Affiliations:

1. Whitehead Institute for Biomedical Research, Cambridge, MA 02142, USA

2. Division of Endocrinology, Diabetes and Hypertension, Brigham and Women's Hospital,

9   Boston, MA 02115, USA

10   3. Harvard Medical School, Boston, MA 02115, USA

11   4. Division of Sleep Medicine, Brigham and Women's Hospital, Boston, MA 02115, USA

12   5. Mary Horrigan Connors Center for Women's Health and Gender Biology, Brigham and

13   Women's Hospital, Boston MA 02115, USA

14   6. Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139,

15   USA

16   7. Howard Hughes Medical Institute, Cambridge, MA 02142, USA

17   8. Harvard Graduate Program in Neuroscience, Boston, MA 02115, USA

18   *Equal contribution.

Abstract

Menopause significantly impacts women's health, yet hypothalamic changes during this transition remain poorly understood. We analyzed gene expression in human hypothalamic samples across menopausal stages, revealing significant changes in inflammatory pathways, KNDy neuron signaling, and thermoregulatory genes. This analysis also identified novel central markers of menopause, including AKAP5 and CDKN1A (p21). To further investigate these changes, we developed a novel mouse model based on long-term (4 months) ovariectomy (OVX) that mimics the gonadotropin and temperature fluctuations of menopausal transition. Remarkably, the posterior hypothalamus of 4-month OVX mice closely mirrored the transcriptional changes observed in perimenopausal women. Both species exhibited increased inflammatory signaling and glial activation, alongside altered KNDy neuron activity. Our findings provide new insights into the central mechanisms of menopausal symptoms and offer a valuable preclinical model for testing interventions.

Introduction

Menopause marks a significant transition in women's lives, characterized by profound physiological changes that extend beyond reproductive function[1]. Understanding the neurobiological underpinnings of these changes is crucial for advancing women's health and developing targeted interventions for menopausal symptoms[2]. While much research has focused on ovarian function, our understanding of the central nervous system's role in menopausal transition, particularly the hypothalamus, remains limited[3].

The hypothalamus houses neuroendocrine centers that both control and are regulated by estrogens[4,5]. This bidirectional relationship positions the hypothalamus as a key player in the menopausal transition. As ovarian function declines and estrogen levels fluctuate, the hypothalamus undergoes significant adaptations, which in turn influence various physiological processes throughout the body[1,6-10]. Notably, these hypothalamic changes are implicated in the disruption of thermoregulation, sleep patterns, and the increased prevalence of mood disorders observed during menopause[7,10-16].

A subset of hypothalamic neurons known as KNDy neurons (named for their co-expression of kisspeptin, neurokinin B (NKB), and dynorphin A) have emerged as central players in the onset of vasomotor symptoms (VMS)[17]. VMS, which include hot flashes and night sweats, are among the most common and disruptive symptoms experienced by women during menopause[18]. The identification of KNDy neurons as triggers for VMS has opened new avenues for understanding and treating these symptoms[17,19]. These findings built on early works from Naomi Rance's group that described hypertrophy of NKB and substance P (SP) expressing neurons in the hypothalamus of post-menopausal women[4,20-23]. Despite these advances, our knowledge of the broader changes occurring in the hypothalamus during and after menopause, particularly regarding the onset, duration, and eventual resolution of menopausal symptoms, remains incomplete.

This gap in our understanding is further compounded by the limitations of current animal

models of menopause[24]. While animal studies have provided valuable insights into some aspects

of menopause, they often fail to fully recapitulate the complex changes observed in women during

this transition. This limitation highlights a critical need for the development of new mouse models

that more closely mimic the human menopausal experience, particularly with respect to

hypothalamic changes.

The lack of comprehensive transcriptomic data from human hypothalamic tissue across

different stages of menopause represents a significant barrier to progress in this field. Such data

would not only enhance our understanding of the molecular mechanisms underlying menopausal

symptoms, but also inform the development of more accurate animal models and potentially lead

to novel therapeutic targets.

In this study, we address these knowledge gaps through a dual approach. First, we present

a complete, unbiased transcriptomic analysis of hypothalamic samples from women across all

menopausal stages. This unique analysis provides a detailed map of gene expression changes

in the human hypothalamus throughout the menopausal transition, offering new insights into the

molecular pathways involved in this process. Second, leveraging the insights gained from our

human transcriptomic analysis, we have developed and characterized a new mouse model that

closely resembles the changes observed in the human hypothalamus during menopause. This

model represents a significant advancement in the field, providing a more accurate platform for

studying menopausal processes and testing potential interventions.

By combining human and mouse transcriptomic data, our study aims to significantly

enhance our understanding of the hypothalamic changes associated with menopause, while

offering a preclinical platform to develop new approaches to manage menopausal symptoms.

Results

Identification and characterization of gene expression changes in the human

hypothalamus across the menopausal transition

Utilizing publicly available RNA-sequencing data from the Genotype-Tissue Expression

(GTEx) project[25], we investigated gene expression changes across the menopausal transition in

the human hypothalamus. Female donors with hypothalamic tissue were included in our analyses

if they passed GTEx donor eligibility criteria and their sequenced tissue had an RNA integrity

number of 6 or greater (Supplementary Table 1). 31 samples from donors between the ages of

28 and 65 years old met our inclusion criteria (Fig. 1a) and samples were divided into five groups

based upon menopause status[26,27], including pre-menopausal (< 45 years old), early peri-

menopausal (46-50 years old), late peri-menopausal (51-55 years old), post-menopausal (56-60

years old) and late post-menopausal (61-65 years old) groups (Fig. 1b). Cell type deconvolution

analyses of the bulk RNA-sequencing data revealed the presence of previously reported

hypothalamic cell types[28] across all samples (Supplementary Fig. 1, Supplementary Table 2). To

assess gene expression changes across the menopausal transition, we began by examining

expression patterns across age in the cohort using locally estimated scatterplot smoothing

(LOESS). Hierarchical clustering of LOESS expression profiles revealed distinct patterns across

age, including genes with nearly constant (Fig. 1c), decreased (Fig. 1d), and increased (Fig. 1e)

expression across age. Of particular interest was the identification of many genes with specific

decreases (Fig. 1f) or increases (Fig. 1g) coinciding with menopausal transition timing

(Supplementary Table 3).

The menopausal transition induces alterations in thermoregulatory signaling and

hypothalamic inflammation

Next, we performed differential gene expression analyses[29] between the pre-menopausal

group and the peri-menopausal and post-menopausal groups. Notably, gene set enrichment

analysis (GSEA)[30] revealed the highest and most significant enrichments in inflammatory pathway

signaling during late peri-menopause (Fig. 1h and Supplementary Table 4). Furthermore,

STRING analysis[31] of TNF-alpha hallmark pathway leading edge genes between late peri-

menopausal and pre-menopausal groups highlighted genes previously identified[32,33] to be

involved in thermoregulation, including fever responses (Fig. 1i).

Given the mounting scientific evidence implicating KNDy neurons in the onset of

vasomotor symptoms (VMS), we examined the expression of key ligand-receptor systems that

constitute the KNDy model, as well as critical regulatory genes, including *TAC1/TACR1* and

estrogen receptor α (*ESR1*). We plotted LOESS trajectories of these genes across age in the

human hypothalamus (Fig 2a). Our analysis revealed that while *KISS1R, TAC3, OPRK1*, and

TACR1 did not display major changes throughout menopausal transition, they tended to increase

with age after menopause. However, *KISS1* and *ESR1* showed a marked increase throughout

menopause, as expected due to the decline in circulating estrogen, followed by a decline in

expression approximately 10 years after their menopausal peak. The expression of aromatase

(*CYP19A1*) showed a progressive increase with age but was unaffected by menopausal

transition. Interestingly, the changes in *KISS1* expression were inversely correlated with that of

the inhibitory dynorphin (*PDYN*) during this same transition, decreasing during menopause and

rising shortly after. *TACR3* expression, which is at the center of the mechanistic pathway that

triggers VMS, showed a marked increase in the years leading up to menopause, reaching

maximum expression at approximately 45 years of age. Thereafter, *TACR3* displayed a constant

decrease in expression, reaching minimal levels around 10 years post-menopause (Fig 2a).

We also examined the prostaglandin systems, which are involved in thermoregulation,

including fever[32,33]. Prostaglandin synthases 1 and 2 (*PTGS1/2*) increased their hypothalamic

expression with age during postmenopausal years. However, *PTGS2* also showed a marked

increase right at the beginning of menopausal transition (~50 years of age). Prostaglandin E

receptor 3 (*PTGER3*) increased right after menopause, followed by a sharp decrease around age

60 (Fig 2a). We also observed an increase with age in the expression of translocator protein

(*TSPO*), a marker of activated microglia, in late post-menopausal years, as previously reported[34].

Additionally, we noted an increase in the astrocyte activation markers *GFAP* and *VIM* during early

post-menopause. These trajectories in pro-inflammatory markers align with the pathways

identified in our GSEA (Fig 1h, i).

Based on our identification of thermoregulatory and inflammatory pathway changes during

the menopausal transition period, we performed an unbiased approach to identify genes of

interest. We conducted differential gene expression analysis between the late peri-menopausal

and pre-menopausal groups (Fig. 2b, Supplementary Table 5). We examined expression of the

identified differentially expressed genes across the defined age groups (Fig. 2c, Supplementary

Table 6) and plotted LOESS trajectories across age for the two identified genes with the most

statistically significant adjusted p-values, *AKAP5* and *CDKN1A* (Fig. 2d, e). We also assessed

expression of canonical aging-associated genes[35] across our groups (Supplementary Fig. 2a) and

plotted trajectories for genes with previously reported functions in brain aging and cognitive

decline[36-40] (Supplementary Fig. 2b). Many of the trajectories exhibited notable inflection points

during the late peri-menopausal years.

Characterization of a novel mouse model of reproductive aging

To experimentally model reproductive aging, we assessed a series of time points post-

OVX in mice (Fig. 3a). Serum luteinizing hormone (LH) concentrations increased progressively

compared to the intact model, with levels peaking at 2 months post-OVX before declining at 4

months post-OVX (Fig. 3b). Importantly, this observation mirrors what is observed over the course

of reproductive aging in women[26]. To characterize changes in thermoregulation, core temperature

was measured across time points in the model with a peak in core temperature observed at 14

days post-OVX (Fig. 3c). The increase in LH and temperature observed at 14 days post-OVX

resembles the changes observed in peri- and early menopause in women, while the decline in LH

and normalization of core temperature seen at 4 months post-OVX resemble the changes

observed in late post-menopausal women.

To further characterize this mouse model and determine whether these physiological

correlations between the rodent and the human extend to hypothalamic transcriptomic changes,

we performed bulk RNA-sequencing of the intact, 14 days (short-term) post-OVX and 4 months

(long-term) post-OVX models. We selected two regions of the hypothalamus to sequence based

on the localization of neurons controlling temperature regulation (preoptic area, POA) and the

localization of neurons controlling the metabolic and reproductive centers (posterior

hypothalamus, PH). Cell type deconvolution analyses of the bulk RNA-sequencing data revealed

the presence of previously reported hypothalamic cell types[28] across all samples just as in the

human dataset (Supplementary Fig. 3).

We then conducted differential gene expression[29] analyses between intact, short-term and

long-term post-OVX models in both the PH and POA. Among the differentially expressed genes,

Gm28040, which shares overlapping CDS with *Kiss1*, shows the largest expression change,

which was increased at 14 days and 4 month OVX in the PH and decreased in both groups in the

POA, resembling the known regulation of *Kiss1* in both hypothalamic areas[41]. This suggests that

in the mouse, *Gm28040* encodes a kisspeptin isoform (Fig 3 e-h, Supplementary Table 7).

Global gene set enrichment analysis revealed the greatest enrichment in inflammatory

pathways similar to what we identified in the human hypothalamus (Fig. 3d, Supplementary Table

8). STRING analysis[31] of TNF-alpha hallmark pathway leading edge genes in the POA also

identified genes related to thermoregulation (Supplemental Fig. 4). Volcano plots highlight

differentially expressed genes between labeled conditions and regions (Fig. 3e-h and

Supplemental Fig. 5a and b), with heatmaps showing how the identified genes are expressed

across conditions in the PH (Fig. 3i) and POA (Fig. 3j) (Supplementary Table 9).

Next, we examined the expression of previously identified marker genes for reproduction,

thermoregulation, and inflammation (particularly related to astrocyte and microglia activity) in both

the PH (Fig. 4a) and the POA (Fig. 4b), as we performed previously in human transcriptomic

samples (Fig 2). The expression of inflammation and astrocytic markers from gonad intact to 4

months post-OVX showed similar increasing trends to that observed in the human hypothalamus

across menopausal transition, with significant increases in *Vim*, *Iba1*, and *Tspo* in the PH, and in

the prostaglandin pathway (*Ptgs2*) in the POA. The "KNDy" genes, involved in reproductive and

thermoregulatory (VMS) control, are highly regulated by sex steroids[42], and thus they showed the

expected increase in expression in the PH (where the arcuate nucleus is located) in proportion to

time after OVX. The specific markers of temperature regulation assessed in this initial analysis

did not show significant changes after OVX.

Comparative analyses of human and mouse hypothalamic reproductive aging

A large proportion of expressed genes are shared between the human and mouse

hypothalamus (~77%) (Supplementary Fig. 6). Comparison of the GSEA results from the two

species revealed statistically significant pathway enrichment correlations between the two

datasets (Fig. 5a), further suggesting the suitability of this mouse model to assess hypothalamic

changes during menopause. To assess whether this correlation can be observed at the single

gene level, we selected *TACR3*, which has a documented role in the onset and severity of VMS[43].

Co-expression analyses identified genes whose expression positively correlated with *TACR3* (in

human) or *Tacr3* (in mouse) LOESS trajectories. These correlations revealed statistically

significant overlap in the positively regulated genes with *TACR3* in the two species (Fig. 5b,

Supplementary Table 10). We then hypothesized that if the short and long term OVX mouse

model is a good model to study hypothalamic transcriptomic changes throughout menopause, the

differentially expressed genes identified in the human samples should show similar trends in our

mouse model and vice versa. We depicted gene expression across the groups as heatmaps,

which show differentially expressed genes from the human dataset across mouse conditions (Fig.

5c, d) and mouse PH and POA datasets across human age groups (Fig. 5e, f). Remarkably, 2/3

of the differentially expressed genes identified in the human between 51-55 years showed a clear

increase at 4 months post-OVX, while the remaining 1/3 were captured at 14d post OVX in both

PH and POA. The opposite analysis revealed a majority of the genes that presented the highest

changes at 4 months post-OVX were upregulated in the 51-55 year old, late peri-menopausal,

group.

In sum, these findings indicate that the short-term and long-term post-OVX mouse models

described can be used as an experimental system to model hypothalamic changes in menopausal

women with a high degree of fidelity. Furthermore, our data reveal that while LH changes occur

relatively fast after OVX, the majority of hypothalamic transcriptomic changes that correlate with

perimenopause are not evident until 4 months post OVX (Fig 3), as also evidenced by pathway

correlation (Fig 5a). These data also reveal that the mouse PH resembles the whole human

hypothalamus better than the mouse POA.

Neurokinin B distribution and neuroinflammatory markers in the reproductive aging

mouse models

Further characterization of this mouse model was carried out at the protein level. To this

end, we continued our focus on the hypothalamic NKB/NK3R system. Unlike *Tac3* expression,

which increases with time after OVX (Fig 4), the expression of NKB decreased after OVX (Fig. 6).

This effect is expected for highly released neuropeptides, suggesting that this ligand is actively

being released rather than stored. To assess the activity of the arcuate nucleus (site of KNDy

neurons) after OVX, we stained for the early activation marker cFos. Importantly, this assay

revealed that cFos expression in the arcuate nucleus is significantly lower at 4 months compared

to 14 days post-OVX, and this was correlated with an increase in astrocytic activation, as shown

by Gfap staining, which correlates with the transcriptomic findings observed in human and mouse

transcriptomes. Nonetheless, these data could not answer whether the decrease in activity was

also at the level of the KNDy neuron or due to changes in other cells of the arcuate. Thus, we

quantified the changes in cFos within NKB neurons, which revealed a 50% decrease in activation.

Overall, we present a novel mouse model to study hypothalamic changes in menopausal

women and we used genes in the KNDy system to demonstrate the validity of this model.

Discussion

Our study utilizes transcriptomics from human hypothalamic samples across adult ages

to identify specific gene expression trends associated with aging and the menopausal transition

in women. Overall our findings offer new insights into the complex molecular landscape of the

aging hypothalamus and its response to declining ovarian function.

A key finding from our analysis is the increased inflammatory signaling in the

hypothalamus during late perimenopause, consistent with the concept of "inflammaging" that

develops with advanced age and contributes to various age-related pathologies[44]. Gene set

enrichment analysis revealed significant upregulation of inflammatory pathways, particularly TNF-

alpha signaling via NF-κB, during this transition period (Fig. 1h). Our STRING analysis of

differentially expressed genes highlighted several that are involved in thermoregulation and fever

responses (Fig. 1i), providing a potential mechanistic link between hypothalamic inflammation

and vasomotor symptoms commonly experienced during menopause. This neuroinflammatory

signature was further evidenced by increased expression of astrocyte activation markers *GFAP*

and VIM during early post-menopause, as well as the microglial marker *TSPO* in late

postmenopausal years (Fig. 2a). These findings align with previous studies showing astrocyte

and microglial reactivity in the aging brain[45]. Additionally, we observed age-associated increases

in prostaglandin synthesis enzymes *PTGS1/2*, which may contribute to neuroinflammation and

altered pain sensitivity in older women. Importantly, our mouse ovariectomy model recapitulated

these inflammatory changes, with gene set enrichment analysis showing significant enrichment

of inflammatory pathways at 4 months post-OVX (Fig. 3d), closely matching our findings in

perimenopausal women. We observed upregulation of glial activation markers and prostaglandin

synthesis enzymes in both the posterior hypothalamus (PH) and preoptic area (POA) (Fig. 4a,b),

mirroring the neuroinflammatory signature seen in aging human hypothalamus. This conserved

inflammatory response across species underscores its potential importance in driving age-related

changes in hypothalamic function.

We also assessed the changes in the expression of genes involved in reproductive

function and thermoregulation across the menopausal transition. As expected, *KISS1* expression

increased markedly during menopause[3,46], likely due to loss of estrogen negative feedback,

followed by a decline approximately 10 years post-menopause (Fig. 2a). This late-life reduction

in kisspeptin signaling may contribute to the eventual decline in gonadotropin levels seen years

after menopause onset[47,48]. Intriguingly, *TACR3* expression, which GWAS studies have

associated with the severity of VMS[43], showed a distinct pattern - increasing in the years leading

up to menopause, peaking around age 45, and then progressively declining (Fig. 2a). Given the

proposed role of neurokinin B signaling in triggering vasomotor symptoms[19,49], this expression

pattern aligns well with the typical time course of hot flashes, which often begin in perimenopause

and resolve within 5-10 years [50].

Unbiased differential expression analysis identified two genes, *AKAP5* and *CDKN1A*, as

particularly notable markers of the menopausal transition (Fig. 2b-e). *AKAP5* encodes a

scaffolding protein involved in synaptic plasticity[51-54] and showed increased expression in early

perimenopause. *CDKN1A*, encoding the cell cycle regulator p21,[55] was upregulated in late

perimenopause. Interestingly, p21 has been implicated in promoting senescence-associated

secretory phenotype (SASP)[55-58], in which senescent cells increase the secretion of inflammatory

factors, thus potentially contributing to the inflammatory phenotype we observed.

To determine whether these human transcriptomic changes could be modeled

experimentally, we developed a mouse ovariectomy (OVX) model with both short-term (14 day)

and long-term (4 month) timepoints. Remarkably, we found that the 4-month OVX timepoint

recapitulated many key features of human perimenopause. LH levels showed an initial increase

followed by decline (Fig. 3b), mirroring the human menopausal transition[26]. Core body

temperature was elevated at 14 days post-OVX but normalized by 4 months (Fig. 3c), paralleling

the typical time course of vasomotor symptoms[50]. Transcriptomic analysis of the mouse PH and

POA revealed striking similarities to the human dataset. Gene set enrichment analysis showed

significant enrichment of inflammatory pathways at 4 months post-OVX (Fig. 3d), closely matching

our findings in perimenopausal women. We observed upregulation of glial activation markers and

prostaglandin synthesis enzymes in both the PH and POA (Fig. 4a,b), recapitulating the

neuroinflammatory signature seen in aging human hypothalamus. Importantly, cross-species

comparison demonstrated significant correlation between the transcriptomic changes in our

human and mouse datasets (Fig. 5a). Nearly two-thirds of genes differentially expressed in late

peri-menopausal women showed concordant changes at 4 months post-OVX in mice, particularly

at the level of the PH where the early marker of menopause in humans *AKAP5* peaks 14d post

OVX and the maker of late peri-menopause, *CDKN1A*, showed the highest expression 4 months

post OVX (Fig 5c). This high degree of conservation suggests that our long term OVX model

effectively captures key molecular features of the human menopausal transition.

To further validate our model, we examined protein changes in neurokinin B (NKB)

expression and neuronal activation. Immunofluorescence staining revealed decreased NKB

levels in the arcuate nucleus at both 14 days and 4 months post-OVX (Fig. 6a-c,j). This contrasts

with the increased *Tac2* mRNA expression we observed and suggests enhanced NKB release

and depletion of peptide stores. Importantly, we found reduced c-Fos expression in NKB neurons

at 4 months post-OVX (Fig. 6m,n), indicating decreased neuronal activity. This aligns with the

declining *TACR3* expression seen in postmenopausal women and may reflect adaptive changes

to prolonged estrogen deficiency.

The age-related transcriptomic changes we identified likely have significant functional

consequences for hypothalamic physiology. Increased inflammatory signaling may disrupt the

precise neuroendocrine control of thermoregulation, metabolism, sleep, mood or cognition, all

critical symptoms of menopause. Astrocyte and microglial activation may impair normal neuronal

function through mechanisms that include altered glutamate uptake and synaptic pruning.

In conclusion, our findings offer significant potential for clinical applications. The

comprehensive transcriptomic data we've generated serves as a valuable resource for future

investigations into hypothalamic aging and the menopausal transition. Our validated mouse model

complements these findings by providing a platform for experimental manipulations, and

potentially allowing to determine the functional significance of specific molecular changes

observed in human samples. Together, these tools and insights pave the way for a deeper

understanding of menopause and the development of interventions that could significantly

improve women's health during this life stage.

Methods

Mouse studies

Study approval and animals

The animal studies were approved by the Brigham and Women's Hospital Institutional

Animal Care and Use Committee (IACUC) in the Center for Comparative Medicine. Adult wild-

type (WT) C57/BL6 female mice were group housed under constant conditions of temperature

(22–24°C) and light (12:12-hour light:dark cycle), fed with standard mouse chow and ad libitum

access to tap water.

Experimental design

To investigate the effects of sex steroids withdrawal on LH levels, core temperature (Tc), and hypothalamic transcriptome profile, intact WT female mice were bilaterally ovariectomized (OVX) to generate mouse models that would resemble pre-menopause, menopause transition and early post-menopause in women. Blood LH levels and Tc were measured at baseline, when females were still intact, at 7-, 14- and 21-days post OVX and at 1-, 2- and 4-months post OVX. Additional females were euthanized before OVX in diestrus (pre-menopause group), after 14 days of OVX (menopause transition group) or after 4 months of OVX (early post-menopause group) and their posterior (PH) and preoptic area (POA) of the hypothalamus were collected for transcriptome profile analysis by bulk RNA-seq.

Ovariectomy (OVX)

Adult (3-4 months-old) female mice were bilaterally OVX under isoflurane anesthesia. Mice were treated with Buprenex SR (0.1 mg/kg, SC) and meloxicam (5 mg/kg, SC) on the day of surgery. Mice were allowed to recover for at least 1 week before the start of experiments.

Whole blood samples and LH Elisa

For LH measurements, a 4μl blood sample was collected with a pipette. Whole blood was immediately diluted in 116μl of 0.05% PBST [phosphate buffer saline (Boston Bio Products, catalog #BM220) containing Tween-20 (Sigma, catalog #P2287)], vortexed, and frozen on dry ice. Samples were stored at 80˚C until analyzed using an in-house sandwich enzyme-linked immunosorbent assay as previously described Kreisman, et al. [59].

Core temperature (Tc)

For Tc measurements, intact female mice were implanted (intraperitoneally) with an IPTT-

300 temperature transponder (BioMedic Data Systems, Inc.). Tc from the transponder were

collected with a DAS-8007 Reader and DASHost 8000 software (BioMedic Data Systems, Inc.).

Immunohistochemistry

Animals were anesthetized with a ketamine/xylazine/saline cocktail and transcardially

perfused with saline (0.9% NaCl) followed by 4% paraformaldehyde (PFA) diluted in phosphate

buffer (PB) (Boston BioProducts, Inc). Brains were removed, stored in 4% PFA overnight and then

transferred into sucrose solution (30% sucrose in 0.1 M PB, Thermo Fisher Scientific) at 4°C. After

sucrose infiltration, the tissue was frozen in Tissue-Tek® O.C.T. Compound (Sakura® Finetek-

4583) and cut on a freezing stage microtome (Microm HM 450, Thermo Fisher Scientific) into 30

μm coronal sections. Free-floating sections were washed in tris buffered saline (TBS, Boston

BioProducts, Inc) and blocked for 1 hour at room temperature in incubation solution [TBS +

0.3%Triton™ X-100 (Millipore Sigma) + 2% normal goat serum (NGS)]. Sections were

subsequently incubated for 48 hours at 4°C with primary antibodies: rabbit-anti neurokinin B

(1:1000, Novus Biologicals, catalog #NB300-201), guinea pig anti-cFos (1:1000, Synaptic

System, catalog #226308) and mouse anti-GFAP (1:300, Cell Signaling, catalog #3670) diluted

in incubation solution. Next, sections were incubated for 1 hour at room temperature with

secondary antibodies: anti-rabbit Alexa Fluor 594 (1:500, Thermo Fisher Scientific, catalog

#A11012), anti-guinea pig Alexa Fluor 488 (1:500, Jackson ImmunoResearch, catalog

#706545148) and anti-mouse donkey Alexa 594 (1:500, Invitrogen, catalog #A32744). Sections

were mounted onto Super-frost plus glass slides (Fisher Laboratories), air dried and cover slipped

with Vectashield mounting medium with DAPI (Vector laboratories).

Microscopy and image analysis

The images were captured by using a fluorescent microscope (Leica DM 2500) in JPG

format and processed using the open-source software ImageJ, Fiji (National Institutes of Health,

Bethesda, Maryland). For image quantification, representative images of the areas of interest (i.e.,

ARC) were delimited using as reference the mouse section from Allen Brain Atlas [60]. Both sides

of bilateral ARC were counted on a similar number of sections and animal, and replicate values

were averaged.

RNA extraction, library preparation, and sequencing

Mouse posterior hypothalamus and preoptic area regions were dissected and flash frozen

prior to extracting RNA using a Zymo Quick RNA microprep kit according to the manufacturer's

instructions. RNA concentration was measured using a Qubit 4.0 fluorometer and the Quibit RNA

HS Assay Kit (ThermoFisher). All samples had RNA integrity numbers (RIN) greater than 7 as

measured on a Bioanalyzer 2100 instrument (Aligent). RNA sequencing libraries were prepared

using the KAPA mRNA HyperPrep–Stranded Kit (Roche) and sequenced on a NovaSeq SP

(Illumina) with 50bp paired-end reads.

RNA-seq data processing and analysis

We performed all analyses using mouse genome build mm39. To pseudoalign reads to

the transcriptome annotation and estimate expression levels of each transcript, we used kallisto[61]

in the tximport package[62] (v1.24.0). We imported the resulting count data (abundance.h5 file) into

R for analysis in DESeq2[29] (v1.36.0). To adjust for unknown, unmodeled or latent sources of

noise in the data we performed surrogate variable analysis (SVA) using the smartSVA package[63]

(v0.1.3) and used SV1 as a covariate in the differential gene expression analysis. Both

pseudoalignment and DESeq2 were performed on the entire transcriptome annotation, but, in

some cases, we restricted our downstream analysis to protein-coding genes, as annotated in

GENCODE version M30[64]. For gene set enrichment analysis (GSEA), we performed analyses

on pre-ranked ordered gene lists using the fgsea package[30] (v1.22.0).

Prediction of protein interactions

The first 30 leading edge genes from the comparison of the POA between female gonadal

intact vs. 4-month post-OVX mice were selected as the target gene list for TNF-α signaling

through NF-κB. The gene list was imported into STRING[65] for network reconstruction,

visualization and enrichment analysis [65]. Medium confidence, as the minimum required interaction

scored, was selected (0.400). Enrichment biological processed defined by gene ontology (GO)

were assessed. Of the top 10 variables identified by the GO terms, highlighted genes related to

thermoregulation and inflammation response to menopause transition are shown.

Human studies

Human hypothalamus inclusion criteria

Bulk RNA-sequencing read count data from human hypothalamus tissue was obtained

from publicly available Genotype-Tissue Expression (GTEx) Project[25] data. We included donor

tissue as part of our dataset if a sample met the following criteria: 1) donor eligibility criteria

(INCEXC) == TRUE, 2) tissue RNA quality score (RIN) >= 6, 3) donors were female (SEX == 2).

Human data analysis

After the sub-setting the GTEx RNA expression file for our samples of interest, we adjusted

for unknown, latent sources of noise in the data using SVA with the smartSVA package[63] (v0.1.3).

Differential expression analyses across age bins were performed using DESeq2[29] (v1.36.0) and

GSEA was performed on pre-ranked ordered gene lists using the fgsea package[30] (v1.22.0).

Gene expression trajectories across age (and parallel analyses across OVX conditions in mouse)

were performed on protein-coding genes using locally estimated scatterplot smoothing (LOESS)

similarly to Piehl et al. 2022[66]. Gene level output from LOESS across age was subjected to

expression trajectory correlation analysis. Genes were ranked based on their correlation with the

TACR3/Tacr3 expression pattern. STRING interaction analyses were performed as described

above.

Cell type decomposition analysis

We performed cell type decomposition analyses to estimate the cell type composition of

our bulk RNA-seq datasets from human and mouse using the BisqueRNA package[67] (v1.0.5).

Table S5 lists the marker genes used to annotation cell types[28].

Quantification and statistical analyses

Various statistical tests were used to calculate p values as indicated in the methods

section, figure legends, or text, where appropriate. All statistics were calculated using R software,

version 4.2.1 unless stated otherwise. Results were considered statistically significant when p <

0.05 or FDR < 0.05 when multiple hypothesis correction was applied.

Data and code availability

• Raw RNA-seq data from mouse posterior hypothalamus and preoptic area regions have

been deposited to dbGAP and are publicly available as of the date of publication.

• This paper analyzes existing, publicly available GTEx consortium data available on the

GTEx Portal (https://www.gtexportal.org/home/).

• Original code to perform analyses will be publicly available as of the date of publication.

• Any additional information required to reanalyze the data reported in this study is available

from the lead contact upon request.

Acknowledgements: The Genotype-Tissue Expression (GTEx) Project was supported by the Common Fund of the Office of the Director of the National Institutes of Health (NIH), and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. The data used for the analyses described in this manuscript were obtained from: the GTEx Portal on 08/10/22.

We also thank Whitehead Institute Genome Technology Core facility for library preparation and sequencing of mouse tissue samples.

Funding: This research was funded by NIH NICHD grants HD090151, HD099084, and NIA/ORWH grant U54 AG062322 to VMN and by The Brit Jepson d'Arbeloff Center on Women's Health to DCP.

Author contributions: J.C.B.B. performed all analyses of the human and mouse RNA-sequencing datasets. E.T.J. and S.A.P. performed all additional mouse experiments. J.C.B.B and V.M.N. wrote the manuscript. D.C.P., E.T.J. and S.A.P. revised the manuscript. D.C.P. and V.M.N. supervised all aspects of the work.

References

Davis, S. R. *et al.* Menopause. *Nature Reviews Disease Primers* **1**, 15004 (2015). https://doi.org/10.1038/nrdp.2015.4

Brinton, R. D., Yao, J., Yin, F., Mack, W. J. & Cadenas, E. Perimenopause as a neurological transition state. *Nature reviews. Endocrinology* **11**, 393-405 (2015). https://doi.org/10.1038/nrendo.2015.82

Rance, N. E. Menopause and the human hypothalamus: evidence for the role of kisspeptin/neurokinin B neurons in the regulation of estrogen negative feedback. *Peptides* **30**, 111-122 (2009). https://doi.org/10.1016/j.peptides.2008.05.016

Rance, N. E. & Uswandi, S. V. Gonadotropin-releasing hormone gene expression is increased in the medial basal hypothalamus of postmenopausal women. *J Clin Endocrinol Metab* **81**, 3540-3546 (1996). https://doi.org/10.1210/jcem.81.10.8855798

Herbison, A. E. in *Physiology of Reproduction* (ed J. Neil and E. Knobil) 1415–1482 (Academic Press, 2006).

Morselli, E. *et al.* The effects of oestrogens and their receptors on cardiometabolic health. *Nature reviews. Endocrinology* **13**, 352-364 (2017). https://doi.org/10.1038/nrendo.2017.12

Cohn, A. Y. *et al.* Effects of Sleep Fragmentation and Estradiol Decline on Cortisol in a Human Experimental Model of Menopause. *J Clin Endocrinol Metab* **108**, e1347-e1357 (2023). https://doi.org/10.1210/clinem/dgad285

Deecher, D. C. & Dorries, K. Understanding the pathophysiology of vasomotor symptoms (hot flushes and night sweats) that occur in perimenopause, menopause, and postmenopause life stages. *Arch Womens Ment Health* **10**, 247-257 (2007). https://doi.org/10.1007/s00737-007-0209-5

Maki, P. M. & Thurston, R. C. Menopause and Brain Health: Hormonal Changes Are Only Part of the Story. *Front Neurol* **11**, 562275 (2020). https://doi.org/10.3389/fneur.2020.562275

Reed, S. D., Newton, K. M., LaCroix, A. Z., Grothaus, L. C. & Ehrlich, K. Night sweats, sleep disturbance, and depression associated with diminished libido in late menopausal transition and early postmenopause: baseline data from the Herbal Alternatives for Menopause Trial (HALT). *Am J Obstet Gynecol* **196**, 593 e591-597; discussion 593 e597 (2007). https://doi.org/10.1016/j.ajog.2007.03.008

Ameratunga, D., Goldin, J. & Hickey, M. Sleep disturbance in menopause. *Intern Med J* **42**, 742-747 (2012). https://doi.org/10.1111/j.1445-5994.2012.02723.x

Augoulea, A. *et al.* Psychosomatic and vasomotor symptom changes during transition to menopause. *Prz Menopauzalny* **18**, 110-115 (2019). https://doi.org/10.5114/pm.2019.86835

Coborn, J. *et al.* Disruption of Sleep Continuity During the Perimenopause: Associations with Female Reproductive Hormone Profiles. *J Clin Endocrinol Metab* **107**, e4144-e4153 (2022). https://doi.org/10.1210/clinem/dgac447

Greendale, G. A., Karlamangla, A. S. & Maki, P. M. The Menopause Transition and Cognition. *JAMA* **323**, 1495-1496 (2020). https://doi.org/10.1001/jama.2020.1757

Kravitz, H. M. & Joffe, H. Sleep during the perimenopause: a SWAN story. *Obstet Gynecol Clin North Am* **38**, 567-586 (2011). https://doi.org/10.1016/j.ogc.2011.06.002

Landis, C. A. & Moe, K. E. Sleep and menopause. *Nurs Clin North Am* **39**, 97-115 (2004). https://doi.org/10.1016/j.cnur.2003.11.006

Koysombat, K., McGown, P., Nyunt, S., Abbara, A. & Dhillo, W. S. New advances in menopause symptom management. *Best Pract Res Clin Endocrinol Metab* **38**, 101774 (2024). https://doi.org/10.1016/j.beem.2023.101774

Freeman, E. W., Sammel, M. D., Lin, H., Liu, Z. & Gracia, C. R. Duration of menopausal hot flushes and associated risk factors. *Obstet Gynecol* **117**, 1095-1104 (2011). https://doi.org/10.1097/AOG.0b013e318214f0de

Prague, J. K. *et al.* Neurokinin 3 receptor antagonism as a novel treatment for menopausal hot flushes: a phase 2, randomised, double-blind, placebo-controlled trial. *Lancet* **389**, 1809-1820 (2017). https://doi.org/10.1016/s0140-6736(17)30823-1

Rance, N. E., McMullen, N. T., Smialek, J. E., Price, D. L. & Young, W. S., 3rd. Postmenopausal hypertrophy of neurons expressing the estrogen receptor gene in the human hypothalamus. *J Clin Endocrinol Metab* **71**, 79-85 (1990). https://doi.org/10.1210/jcem-71-1-79

Rance, N. E. & Young, W. S., 3rd. Hypertrophy and increased gene expression of neurons containing neurokinin-B and substance-P messenger ribonucleic acids in the hypothalami of postmenopausal women. *Endocrinology* **128**, 2239-2247 (1991).

Rance, N. E. Hormonal influences on morphology and neuropeptide gene expression in the infundibular nucleus of postmenopausal women. *Prog Brain Res* **93**, 221-235; discussion 235-226 (1992).

Chawla, M. K., Gutierrez, G. M., Young, W. S., 3rd, McMullen, N. T. & Rance, N. E. Localization of neurons expressing substance P and neurokinin B gene transcripts in the human hypothalamus and basal forebrain. *The Journal of comparative neurology* **384**, 429-442 (1997).

Koebele, S. V. & Bimonte-Nelson, H. A. Modeling menopause: The utility of rodents in translational behavioral endocrinology research. *Maturitas* **87**, 5-17 (2016). https://doi.org/10.1016/j.maturitas.2016.01.015

Lonsdale, J. *et al.* The Genotype-Tissue Expression (GTEx) project. *Nature Genetics 2013 45:6* **45** (2013-05-29). https://doi.org/10.1038/ng.2653

Soares, A. G. *et al.* Longitudinal changes in reproductive hormones through the menopause transition in the Avon Longitudinal Study of Parents and Children (ALSPAC). *Scientific Reports 2020 10:1* **10** (2020-12-04). https://doi.org/10.1038/s41598-020-77871-9

Khoudary, S. R. E. *et al.* The menopause transition and women's health at midlife: a progress report from the Study of Women's Health Across the Nation (SWAN). *Menopause (New York, N.y.)* **26** (2019/10). https://doi.org/10.1097/GME.0000000000001424

Hajdarovic, K. H. *et al.* Single-cell analysis of the aging female mouse hypothalamus. *Nature Aging 2022 2:7* **2** (2022-07-04). https://doi.org/10.1038/s43587-022-00246-4

29   Love, M. I. *et al.* Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. *Genome Biology 2014 15:12* **15** (2014-12-05).
https://doi.org/10.1186/s13059-014-0550-8

30   Korotkevich, G. *et al.* Fast gene set enrichment analysis. *BioRxiv* (2021-02-01).
https://doi.org/10.1101/060012

31   The STRING database in 2021: customizable protein-protein networks, and functional
characterization of user-uploaded gene/measurement sets - PubMed. *Nucleic Acids Res*
49 (01/08/2021). https://doi.org/10.1093/nar/gkaa1074

32   Blomqvist, A. & Engblom, D. Neural Mechanisms of Inflammation-Induced Fever. *The*
Neuroscientist **24** (2018/08). https://doi.org/10.1177/1073858418760481

33   Prostaglandin production selectively in brain endothelial cells is both necessary and
sufficient for eliciting fever - PubMed. *Proc Natl Acad Sci USA* **119** (10/25/2022).
https://doi.org/10.1073/pnas.2122562119

34   Butler, T. *et al.* Positron Emission Tomography reveals age-associated hypothalamic
microglial activation in women. *Sci Rep* **12**, 13351 (2022).
https://doi.org/10.1038/s41598-022-17315-8

35   Tacutu, R. *et al.* Human Ageing Genomic Resources: new and updated databases. *Nucleic*
Acids Res **46** (2018/01/01). https://doi.org/10.1093/nar/gkx1042

36   O'Day, D. H. Calmodulin Binding Proteins and Alzheimer's Disease: Biomarkers,
Regulatory Enzymes and Receptors That Are Regulated by Calmodulin. *Int J Mol Sci* **21**
(2020/10). https://doi.org/10.3390/ijms21197344

37   Kim, J., Basak, J. M. & Holtzman, D. M. The Role of Apolipoprotein E in Alzheimer's
Disease. *Neuron* **63** (2009/08/13). https://doi.org/10.1016/j.neuron.2009.06.026

38   Bateman, R. J. *et al.* Clinical and Biomarker Changes in Dominantly Inherited Alzheimer's
Disease. *New England Journal of Medicine* **367** (2012-08-30).
https://doi.org/10.1056/NEJMoa1202753

39   SIRT1 in Neurodevelopment and Brain Senescence. *Neuron* **81** (2014/02/05).
https://doi.org/10.1016/j.neuron.2014.01.028

40   Catechol-O-methyltransferase polymorphisms and some implications for cognitive
therapeutics - PubMed. *NeuroRx : the journal of the American Society for Experimental*
NeuroTherapeutics **3** (2006 Jan). https://doi.org/10.1016/j.nurx.2005.12.010

41   Smith, J. T., Cunningham, M. J., Rissman, E. F., Clifton, D. K. & Steiner, R. A. Regulation of
Kiss1 gene expression in the brain of the female mouse. *Endocrinology* **146**, 3686-3692
(2005). https://doi.org/10.1210/en.2005-0488

42   Goodman, R. L., Herbison, A. E., Lehman, M. N. & Navarro, V. M. Neuroendocrine control
of gonadotropin-releasing hormone: Pulsatile and surge modes of secretion. *Journal of*
neuroendocrinology **34**, e13094 (2022). https://doi.org/10.1111/jne.13094

43   Crandall, C. J. *et al.* Association of genetic variation in the tachykinin receptor 3 locus
with hot flashes and night sweats in the Women's Health Initiative Study. *Menopause* **24**,
252-261 (2017). https://doi.org/10.1097/gme.0000000000000763

44   Franceschi, C., Garagnani, P., Vitale, G., Capri, M. & Salvioli, S. Inflammaging and 'Garb-
aging'. *Trends Endocrinol Metab* **28**, 199-212 (2017).
https://doi.org/10.1016/j.tem.2016.09.005

45 Clarke, L. E. *et al.* Normal aging induces A1-like astrocyte reactivity. *Proceedings of the*
National Academy of Sciences of the United States of America **115**, E1896-e1905 (2018).
https://doi.org/10.1073/pnas.1800165115

46 Rometo, A. M., Krajewski, S. J., Voytko, M. L. & Rance, N. E. Hypertrophy and increased
kisspeptin gene expression in the hypothalamic infundibular nucleus of postmenopausal
women and ovariectomized monkeys. *J Clin Endocrinol Metab* **92**, 2744-2750 (2007).
https://doi.org/10.1210/jc.2007-0553

47 Hall, J. E. Neuroendocrine physiology of the early and late menopause. *Endocrinology*
and Metabolism Clinics **33** (2004/12/01). https://doi.org/10.1016/j.ecl.2004.08.002

48 Hall, J. E., Lavoie, H. B., Marsh, E. E. & Martin, K. A. Decrease in Gonadotropin-Releasing
Hormone (GnRH) Pulse Frequency with Aging in Postmenopausal Women*. *The Journal*
of Clinical Endocrinology & Metabolism **85** (2000/05/01).
https://doi.org/10.1210/jcem.85.5.6612

49 Christ, J. P., Navarro, V. M. & Reed, S. D. Nonhormonal Therapies for Menopausal
Vasomotor Symptoms. *Jama* **330**, 1278-1279 (2023).
https://doi.org/10.1001/jama.2023.15965

50 Avis, N. E. *et al.* Duration of Menopausal Vasomotor Symptoms. *JAMA Internal Medicine*
175 (2015/04/01). https://doi.org/10.1001/jamainternmed.2014.8063

51 He, Z. *et al.* Novel insight into the role of A-kinase anchoring proteins (AKAPs) in
ischemic stroke and therapeutic potentials. *Biomedicine & Pharmacotherapy* **175**,
116715 (2024). https://doi.org/https://doi.org/10.1016/j.biopha.2024.116715

52 Koster, K. P. *et al.* Akap5 links synaptic dysfunction to neuroinflammatory signaling in a
mouse model of infantile neuronal ceroid lipofuscinosis. *Front Synaptic Neurosci* **16**,
1384625 (2024). https://doi.org/10.3389/fnsyn.2024.1384625

53 Sanderson, J. L. & Dell'Acqua, M. L. AKAP signaling complexes in regulation of excitatory
synaptic plasticity. *Neuroscientist* **17**, 321-336 (2011).
https://doi.org/10.1177/1073858410384740

54 Weisenhaus, M. *et al.* Mutations in AKAP5 disrupt dendritic signaling complexes and
lead to electrophysiological and behavioral phenotypes in mice. *PloS one* **5**, e10325
(2010). https://doi.org/10.1371/journal.pone.0010325

55 Engeland, K. Cell cycle regulation: p53-p21-RB signaling. *Cell Death & Differentiation* **29**,
946-960 (2022). https://doi.org/10.1038/s41418-022-00988-z

56 Wagner, K. D. & Wagner, N. The Senescence Markers p16INK4A, p14ARF/p19ARF, and
p21 in Organ Development and Homeostasis. *Cells* **11** (2022).
https://doi.org/10.3390/cells11121966

57 Wang, L., Lankhorst, L. & Bernards, R. Exploiting senescence for the treatment of cancer.
Nature Reviews Cancer **22**, 340-355 (2022). https://doi.org/10.1038/s41568-022-00450-
9

58 Chinta, S. J. *et al.* Cellular senescence and the aging brain. *Exp Gerontol* **68**, 3-7 (2015).
https://doi.org/10.1016/j.exger.2014.09.018

59 Kreisman, M. J., McCosh, R. B. & Breen, K. M. A Modified Ultra-Sensitive ELISA for
Measurement of LH in Mice. *Endocrinology* **163** (2022/09).
https://doi.org/10.1210/endocr/bqac109

60   Wang, Q. *et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D Reference
684   Atlas. *Cell* **181**, 936-953 e920 (2020). https://doi.org/10.1016/j.cell.2020.04.007
685   61   Bray, N. L. *et al.* Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology*
686   *2016 34:5* **34** (2016-04-04). https://doi.org/10.1038/nbt.3519
687   62   Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq: transcript-
688   level estimates improve gene-level inferences - PubMed. *F1000Res* **4** (12/30/2015).
689   https://doi.org/10.12688/f1000research.7563.2
690   63   Chen, J. *et al.* Fast and robust adjustment of cell mixtures in epigenome-wide association
691   studies with SmartSVA - PubMed. *BMC genomics* **18** (05/26/2017).
692   https://doi.org/10.1186/s12864-017-3808-1
693   64   Frankish, A. *et al.* GENCODE reference annotation for the human and mouse genomes.
694   *Nucleic Acids Res* **47** (2019/01/01). https://doi.org/10.1093/nar/gky955
695   65   Szklarczyk, D. *et al.* STRING v11: protein-protein association networks with increased
696   coverage, supporting functional discovery in genome-wide experimental datasets.
697   *Nucleic Acids Res* **47**, D607-D613 (2019). https://doi.org/10.1093/nar/gky1131
698   66   Piehl, N. *et al.* Cerebrospinal fluid immune dysregulation during healthy brain aging and
699   cognitive impairment. *Cell* **185** (2022/12/12). https://doi.org/10.1016/j.cell.2022.11.019
700   67   Jew, B. *et al.* Accurate estimation of cell composition in bulk expression through robust
701   integration of single-cell information. *Nature Communications 2020 11:1* **11** (2020-04-
702   24). https://doi.org/10.1038/s41467-020-15816-6
703

Figure 1. Gene expression changes across the menopausal transition in the human hypothalamus reveal alterations in thermoregulatory and inflammatory signaling.

a) Demographic overview of the GTEx donor subset used in this study.

b) Distribution of ages within the donor cohort and the defined age groupings for subsequent analyses.

c-g) Sets of genes grouped by hierarchical clustering and displayed using LOESS trajectories, **c)** constant expression across age, **d)** decreased expression across age, **e)** increased expression across age, **f)** decreased expression during the peri-menopausal transition, **g)** increased expression during the peri-menopausal transition.

h) Heatmap of normalized enrichment scores (NES) for Hallmark gene sets in which there was a statistically significant pathway enrichment in at least one of the three comparisons; significant enrichments indicated by asterisks (* *p*-adj <0.05; ** *p*-adj < 0.01; *** *p*-adj < 0.001)

i) STRING analysis of TNF-alpha hallmark pathway leading edge genes between the late peri-menopausal vs. pre-menopausal groups.

Figure 2. Expression of reproductive- and aging-related genes during the menopausal transition in the human hypothalamus.

a) Expression of KNDy genes and their receptors, thermoregulatory genes, astrocyte-, and microglia-marker genes across age in human hypothalamus tissue.

b) Volcano plot highlighting significantly differentially expressed genes between late peri- and pre-menopausal groups. Genes with a FDR < 0.05 are highlighted in pink.

c) Heatmap of genes identified in **b** and their expression across age groups.

d) Expression of *AKAP5* across age in human hypothalamus tissue.

e) Expression of *CDKN1A* across age in human hypothalamus tissue.

Figure 3

Figure 3. Physiological and transcriptomic changes in post-ovariectomy mouse models recapitulate key features of reproductive aging.

    a) Schematic representation of the experimental design.

    b) Luteinizing Hormone (LH) concentration in serum isolated from whole blood 7-, 14-, 21-days post-OVX and 1-, 2- and 4-months post-OVX in wild-type female mice (n=20, per condition).

    c) Core temperature (ºC) measurements at 7-, 14-, 21-days post-OVX and 1-, 2- and 4-months post-OVX in wild-type female mice (n=20, per condition). b-c) Data values are shown as mean ± SEM. Groups with different letters are significantly from each other, P < 0.05 by One-way ANOVA/Tukey.

    d) Heatmap of normalized enrichment scores (NES) for Hallmark gene sets in the Posterior Hypothalamus (PH) and the Preoptic Area (POA); significant enrichments indicated by asterisks (* p-adj <0.05; ** p-adj < 0.01; *** p-adj < 0.001)

e-h) Volcano plots highlighting significantly differentially expressed genes between labeled conditions and regions. Genes with a FDR < 0.05 are labeled in pink.

i) Heatmap of PH DE genes identified in **b** and their expression across conditions.

j) Heatmap of POA DE genes identified in **d** and their expression across conditions.

Figure 4. Expression of reproduction-, thermoregulation- and inflammation-related genes in short- and long-term post-ovariectomy mouse models.

    a) Expression of KNDy genes and their receptors, thermoregulatory genes, astrocyte, and microglia marker genes between groups in the posterior hypothalamus (PH).

    b) Expression of KNDy genes and their receptors, thermoregulatory genes, astrocyte, and microglia marker genes between groups in the preoptic area (POA).

762
Figure 5. Comparative analysis of human and mouse hypothalamic reproductive aging datasets.

a) Heatmap of GSEA pathway correlations between human and mouse datasets; *p*-value from Pearson's correlation.

b) Overlap of *TACR3/Tacr3* positively correlated co-expressed genes between human and mouse; *p*-value from hypergeometric test.

Heatmap of identified DE genes from the human dataset and their expression across
conditions in the mouse **c)** PH and **d)** POA datasets.
e) Heatmap of identified DE genes from the mouse PH and their expression across age
groups in the human dataset.
f) Heatmap of identified DE genes from the mouse POA and their expression across age
groups in the human dataset.
775
776
777
778

781 **Figure 6. Differential neurokinin B distribution, neuronal activity and astrocyte reactivity**
782 **within the arcuate nucleus from the short- and long-term post-ovariectomy mouse models.**

Representative immunofluorescence staining of NKB, cFos and Gfap in the arcuate nucleus (ARC) from intact **(a-c)**, 14 days post-OVX **(d-f)**, and 4 months post-OVX female mice **(g-i)**. Quantification of NKB **(j)**, cFos **(k)**, and Gfap **(l)** in the ARC from intact, 14 days post- and 4 months post-OVX female mice.

m) Co-localization of NKB and cFos in the ARC from 14 days post- and 4 months post-OVX female mice. Data from **j-l** and **n** shown as mean ± SEM. Statistical significance was determinate by t-student Mann-Whitney test (intact vs. OVX).