Long-term post-ovariectomy mouse model recapitulates hypothalamic transcriptomic changes observed during the menopausal transition in women

Authors: Jordana C. B. Bloom[1,2], Encarnacion Torres[2,3]†, Sidney A. Pereira[2,3]†, Liliana Arvizu-Sanchez[2], Audrey N. Fontes[2], Hadine Joffe[3,4], David C. Page[1,5,6], Victor M. Navarro[2,3,7]*

Affiliations:

1. Whitehead Institute for Biomedical Research, Cambridge, MA 02142, USA
2. Division of Endocrinology, Diabetes and Hypertension, Brigham and Women's Hospital, Boston, MA 02115, USA
3. Harvard Medical School, Boston, MA 02115, USA
4. Department of Psychiatry, Beth Israel Deaconess Medical Center, Boston, MA 02115, USA
5. Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
6. Howard Hughes Medical Institute, Cambridge, MA 02142, USA
7. Harvard Graduate Program in Neuroscience, Boston, MA 02115, USA

*Email: vnavarro@bwh.harvard.edu

†These authors contributed equally to this work

Abstract

Menopause significantly impacts women's health, yet hypothalamic changes during this transition

remain poorly understood. We analyzed gene expression in human hypothalamic samples across

menopausal transition, revealing significant changes in inflammatory pathways, KNDy neuron

signaling, and thermoregulatory genes. This analysis also identified novel central markers of

menopause, including *AKAP5* and *CDKN1A* (*p21*). To further investigate these changes, we

developed a novel mouse model based on long-term (4 months) post-ovariectomy (OVX) that

mimics the gonadotropin and temperature fluctuations of menopausal transition. Remarkably, the

posterior hypothalamus of 4-month post-OVX mice closely mirrored the transcriptional changes

observed in women at the late peri-/early post-menopausal age range (ages 51-55). Both species

exhibited increased inflammatory signaling and glial activation, alongside altered KNDy neuron

activity. Our findings provide new insights into the central mechanisms of menopausal symptoms

and offer a valuable preclinical model for testing interventions.

Introduction

Menopause represents a significant life transition for women, characterized by physiological changes extending beyond reproductive function[1]. Understanding the neurobiological basis of these changes is crucial for advancing women's health and developing targeted interventions for menopausal symptoms[2]. While much attention has been focused on ovarian function, research focused on characterizing the changes that happen at the level of the brain—particularly the hypothalamus—has lagged. The hypothalamus contains neuroendocrine centers that both control and respond to estrogens[3,4], and hypothalamic changes are implicated in disrupted thermoregulation, sleep patterns, and increased mood disorders during menopause[5-12].

Hypothalamic KNDy neurons (expressing kisspeptin, neurokinin B, and dynorphin A) have emerged as central players in vasomotor symptoms (VMS), including hot flashes and night sweats, which are among the most common and disruptive menopausal symptoms[13,14]. The identification of KNDy neurons as VMS triggers has opened new avenues for understanding and treating these symptoms[13,15]. However, our knowledge of broader hypothalamic changes during and after menopause remains incomplete, compounded by limitations in current animal models[16]. By combining human and mouse transcriptomic data, this study enhances our understanding while offering a preclinical platform for developing new approaches to manage menopausal symptoms.

Results

Utilizing publicly available bulk RNA-sequencing data from the Genotype-Tissue Expression (GTEx) project[17], we investigated gene expression changes across the menopausal transition in the human hypothalamus. Female donors with hypothalamic tissue were included in our analyses if they passed GTEx donor eligibility criteria and their sequenced tissue had an RNA integrity number of 6 or greater (Supplementary Table 1). 31 samples from donors including

those in the menopausal transition (between the ages of 28 and 65 years old) met our inclusion

criteria (Fig. 1a). Surrogate variable analysis (SVA)[18] was performed on the RNA expression

matrix prior to gene expression analyses to control for latent, confounding sources of variation in

the data, such as postmortem interval, donor cause of death, and hormone therapy status.

Additionally, cell type deconvolution of the bulk RNA-sequencing data revealed the presence of

previously reported hypothalamic cell types[19] (Supplementary Fig. 1a, Supplementary Table 2)

and the presence of all eight major transcription factors (*FEZF1, FOXB1, LHX6, MEIS2, OTP,*

SIM1, SIX3, TBX3) identified in the human hypothalamus from the HYPOMAP single-nucleus

RNA-seq reference atlas[20] (Supplementary Fig. 1b) indicating that there are no major differences

in cell type composition between samples in the cohort.

Having established that there is comparable cell-type representation across the

hypothalamus samples, we next assessed patterns of gene expression changes across the

menopausal transition by examining gene expression across age using locally estimated

scatterplot smoothing (LOESS). LOESS expression profiles were grouped together based on

Euclidean distance hierarchical clustering which revealed distinct patterns across age, including

genes with nearly constant (Fig. 1b), decreased (Fig. 1c), and increased (Fig. 1d) expression

across age. Of particular interest was the identification of many genes with specific decreases

(Fig. 1e) or increases (Fig. 1f) coinciding with expected menopausal transition timing

(Supplementary Table 3).

Next, we divided the cohort into groups in age bins that correspond to the average age

range for each menopausal stage based on large longitudinal studies using the age of final

menstrual period as stage-defining factor[21-23] (Fig. 2a). We then performed differential gene

expression analyses[24] between groups of donors younger than 45 and those between the ages

of 46-50, 51-55, and 56-60. Notably, gene set enrichment analysis (GSEA)[25] revealed the highest

and most significant enrichments in inflammatory pathway signaling in the group encompassing

donors between the ages of 51-55 (which often includes late peri-menopausal/early post-

menopausal individuals) compared to individuals younger than 45 (often pre-menopausal) (Fig.

2b and Supplementary Table 4). Furthermore, STRING analysis[26] of TNF-alpha hallmark

pathway leading edge genes between the late peri-menopausal/early post-menopausal and pre-

menopausal groups highlighted genes previously identified[27,28] to be involved in thermoregulation,

including fever responses (Fig. 2c).

Given the mounting scientific evidence implicating KNDy neurons in the onset of VMS, we

examined the expression of key ligand-receptor systems that constitute the KNDy model, as well

as critical regulatory genes, including *TAC1/TACR1* and estrogen receptor α (*ESR1*). We plotted

LOESS trajectories of these genes in the human hypothalamus (Fig 2d) to visualize their

expression patterns across chronological age. Our analysis revealed that while *KISS1R, TAC3,*

OPRK1, and *TACR1* did not appear to display major changes with age throughout menopausal

transition, they tended to increase after age 50, coinciding with the typical age of menopause.

However, *KISS1* and *ESR1* showed a marked increase throughout menopause, as expected due

to the decline in circulating estrogen, followed by a decline in expression approximately 10 years

after their menopausal peak. The expression of aromatase (*CYP19A1*) showed a progressive

increase with age but was unaffected by menopausal transition. Interestingly, the changes in

KISS1 expression were inversely correlated with that of the inhibitory dynorphin (*PDYN*) during

this same transition, decreasing during typical menopause years (50-55) and rising shortly after.

TACR3 expression, which is at the center of the mechanistic pathway that triggers VMS, showed

a marked increase in the years leading up to menopause, reaching maximum expression at

approximately 45 years of age. Thereafter, *TACR3* displayed a constant decrease in expression,

reaching minimal levels around 10 years post-menopause (Fig 2d). Additionally, we assessed the

expression of *PGR*, *GPER1*, and *ESR2* (Supplementary Fig. 2) based on a recent publication that

examined expression of these genes as potential postmortem biomarkers in the hypothalamus[29].

We also examined the prostaglandin systems, which are involved in thermoregulation,

including fever[27,28]. Prostaglandin synthases 1 and 2 (*PTGS1/2*) increased their hypothalamic

expression with age during post-menopausal years. However, Prostaglandin E receptor 3

(*PTGER3*) increased right after menopause, followed by a sharp decrease around age 60 (Fig

2d). We also observed an increase with age in the expression of translocator protein (*TSPO*), a

marker of activated microglia, in late post-menopausal years, as previously reported[30].

Additionally, the pro-inflammatory astrocyte activation markers *GFAP* and *VIM* increase during

the years corresponding to the average ages of late peri-/early post-menopause, which aligns

with the increase in inflammatory signaling pathways identified in our GSEA (Fig 2b, c).

Based on our identification of thermoregulatory and inflammatory pathway changes during

the menopausal transition period, we performed an unbiased approach to identify novel genes of

interest. We conducted differential gene expression analysis between age bins corresponding to

normal late peri-/early post-menopausal and pre-menopausal periods (Fig. 3a, Supplementary

Table 5). We examined expression of the identified significantly differentially expressed genes

across the defined age groups (Fig. 3b, Supplementary Table 6) and plotted LOESS trajectories

across age for the two identified genes with the most statistically significant adjusted p-values,

AKAP5 and *CDKN1A* (Fig. 2c, d). We used the Allen Brain Cell Atlas Brain Knowledge Platform

(https://knowledge.brain-map.org/) to visualize transcript localization in the mouse brain

(Supplementary Fig. 3a, b). Additionally, we assessed expression of canonical aging-associated

genes[31] across our groups (Supplementary Fig. 4a) and plotted trajectories for genes with

previously reported functions in brain aging and cognitive decline[32-36] (Supplementary Fig. 4b, c).

Many of the trajectories exhibited notable inflection points during the age range corresponding

with late peri-/early post-menopausal years.

To experimentally model the time without estrogen exposure that occurs during

reproductive aging, we assessed a series of time points post-OVX in mice (Fig. 4a). Serum

luteinizing hormone (LH) concentrations increased progressively compared to the intact model,

with levels peaking at 2 months post-OVX before declining at 4 months post-OVX (Fig. 4b).

Importantly, this observation mirrors what is observed over the course of reproductive aging in

women[21]. To characterize changes in thermoregulation, core temperature was measured across

time points in the model with a peak in core temperature observed at 14 days post-OVX (Fig. 4c).

The increase in LH and temperature observed at 14 days post-OVX resembles the changes

observed in peri- and early menopause in women, while the decline in LH and normalization of

core temperature seen at 4 months post-OVX resemble the changes observed in late post-

menopausal women.

To further characterize this mouse model and determine whether these physiological

correlations between the rodent and the human extend to hypothalamic transcriptomic changes,

we performed bulk RNA-sequencing of the intact, 14 days (short-term) post-OVX and 4 months

(long-term) post-OVX models. We selected two regions of the hypothalamus to sequence based

on the localization of neurons controlling temperature regulation (preoptic area, POA) and the

localization of neurons controlling the metabolic and reproductive centers (posterior

hypothalamus, PH). Cell type deconvolution analyses of the bulk RNA-sequencing data revealed

the presence of previously reported hypothalamic cell types[19] across all samples just as in the

human dataset (Supplementary Fig. 5).

Global gene set enrichment analysis revealed the greatest enrichment in inflammatory

pathways similar to what we identified in the human hypothalamus (Fig. 4d, Supplementary Table

7). STRING analysis[26] of TNF-alpha hallmark pathway leading edge genes in the POA also

identified genes related to thermoregulation (Supplemental Fig. 6). We then conducted

differential gene expression[24] analyses between intact, short-term and long-term post-OVX

models in both the PH and POA. Volcano plots highlight significantly differentially expressed

genes between labeled conditions and regions (Fig. 4e-h, Supplemental Fig. 7a and b,

Supplementary Table 8), with heatmaps showing how the identified genes are expressed across

conditions in the PH (Fig. 4i) and POA (Fig. 4j) (Supplementary Table 9). Among the differentially

expressed genes, *Gm28040*, which shares overlapping CDS with *Kiss1*, shows the largest

expression change, which was increased at 14 days and 4 month OVX in the PH and decreased

in both groups in the POA, resembling the known regulation of *Kiss1* in both hypothalamic areas[37].

This suggests that in the mouse, *Gm28040* encodes a kisspeptin isoform (Fig 4 e-h,

Supplementary Table 8).

Next, we examined the expression of previously identified marker genes for reproduction,

thermoregulation, and inflammation (particularly related to astrocyte and microglia activity) in both

the PH (Supplemental Fig. 8a) and the POA (Supplemental Fig. 8b), as we performed previously

in human transcriptomic samples (Fig 2). While none of the genes reached statistical significance

after adjusting for multiple hypothesis testing, the expression of inflammation and astrocytic

markers from gonad intact to 4 months post-OVX showed similar increasing trends to that

observed in the human hypothalamus across menopausal transition, with notable increases in

Vim, *Iba1*, and *Tspo* in the PH, and in the prostaglandin pathway (*Ptgs2*) in the POA. The KNDy

genes, involved in reproductive and thermoregulatory (VMS) control, are highly regulated by sex

steroids[38], and thus they showed the expected increase in expression in the PH (where the

arcuate nucleus is located) in proportion to time after OVX.

A substantial proportion of expressed genes are shared between the human and mouse

hypothalamus (~77%; Supplementary Fig. 9). Comparison of the GSEA results between the

human GTEx dataset and OVX mouse model revealed statistically significant pathway enrichment

correlations between the two datasets (Fig. 5a), further supporting the suitability of this mouse

model to study hypothalamic changes during the menopausal transition. We also compared

GSEA results between our OVX mouse model and a published single-cell RNA-seq dataset of

the aging female mouse hypothalamus[19] to assess whether the mouse models capture similar

hypothalamic gene expression changes. Of the 26 Hallmark pathways significantly impacted in

the human dataset (Fig. 2b, Supplementary Table 4) only eight were similarly altered in the aged

mouse hypothalamus[19], compared with 14 in the OVX model. Notably, the aged mouse study[19]

did not report any changes in the estrogen response or TNFα signaling pathways, which are

consistently affected in both the human dataset and the OVX mouse model and highly relevant

to the menopausal transition[39]. At the cell-type level, the observed pathway changes in

macrophages and microglia from the aged mouse hypothalamus dataset exhibited the greatest

similarity to those changes observed in the human and OVX mouse model (Supplementary Table

10). Despite this partial overlap between mouse models, the two only share a single significantly

differentially expressed gene in common (*Thsd4)*, suggesting that they capture largely distinct

hypothalamic transcriptional programs.

To determine whether the cross-species similarities extend beyond pathway-level

analyses to individual genes, we selected *TACR3*, which has a documented role in the onset and

severity of VMS[40] and performed co-expression analyses to identify genes whose expression

positively correlates with *TACR3* (in human) or *Tacr3* (in the OVX mouse model). These

correlations revealed statistically significant overlap in the positively regulated genes with *TACR3*

in the two species (Fig. 5b, Supplementary Table 11). We then hypothesized that if the short and

long term OVX mouse model is a suitable model to study hypothalamic transcriptomic changes

throughout menopause, the differentially expressed genes identified in the human samples should

show similar trends in our mouse model and vice versa. We depicted gene expression across the

groups as heatmaps, which show differentially expressed genes from the human dataset across

mouse conditions (Fig. 5c, d) and mouse PH and POA datasets across human age groups (Fig.

5e, f). Remarkably, two-thirds of the differentially expressed genes identified in the human

between 51-55 years showed a clear increase at 4 months post-OVX, while the remaining one-

third were captured at 14 days post-OVX in both PH and POA. The opposite analysis revealed a

majority of the genes that presented the highest changes at 4 months post-OVX were upregulated

in the 51-55 year old age group, usually corresponding to individuals in the late peri-/early post-

menopausal stage.

In sum, these findings indicate that the short-term and long-term post-OVX mouse models

described can be used as an experimental system to model hypothalamic changes in menopausal

women with a high degree of fidelity. Furthermore, our data reveal that while LH changes occur

relatively fast after OVX, the majority of hypothalamic transcriptomic changes that correlate with

perimenopause are not evident until 4 months post OVX (Figs 4, 5a). These data also reveal that

the mouse PH resembles the whole human hypothalamus better than the mouse POA.

Further characterization of this mouse model was carried out at the protein level. To this

end, we continued our focus on the hypothalamic NKB/NK3R system. Unlike *Tac2* expression,

which increases with time after OVX (Supplemental Fig. 8), the expression of NKB decreased

after OVX (Supplemental Fig. 10). This effect is expected for highly released neuropeptides,

suggesting that this ligand is actively being released rather than stored. To assess the activity of

the arcuate nucleus (site of KNDy neurons) after OVX, we stained for the early activation marker

cFos. This assay revealed that cFos expression in the arcuate nucleus is significantly lower at 4

months compared to 14 days post-OVX, and this was correlated with an increase in inflammation,

as shown by Gfap staining, which correlates with the findings observed in human and mouse

transcriptomes. Nonetheless, these data could not answer whether the decrease in activity was

also at the level of the KNDy neuron or due to changes in other cells of the arcuate. Thus, we

quantified the changes in cFos within NKB neurons, which revealed a 50% decrease in NKB

neuron activation.

Discussion

Our study utilizes transcriptomics from human hypothalamic samples across adult ages

to identify specific gene expression trends associated with aging and the menopausal transition

in women. Overall, our findings offer new genome-wide insights into the complex molecular

landscape of the aging hypothalamus and its response to declining ovarian function.

A key finding from our analysis is the increased inflammatory signaling in the

hypothalamus during the age range usually corresponding to individuals in late peri-/early post-

menopause, consistent with the concept of "inflammaging" that develops with advanced age and

contributes to various age-related pathologies[41]. Gene set enrichment analysis revealed

significant upregulation of inflammatory pathways, particularly TNF-alpha signaling via NF-κB, during this transition period (Fig. 2b). Our STRING analysis of differentially expressed genes highlighted several that are involved in thermoregulation and fever responses (Fig. 2c), providing a potential mechanistic link between hypothalamic inflammation and vasomotor symptoms commonly experienced during menopause. This neuroinflammatory signature was further evidenced by increased expression of astrocyte activation markers *GFAP* and VIM during normal early post-menopause years, as well as the microglial marker *TSPO* in late post-menopausal years (Fig. 2d). Additionally, we observed age-associated increases in prostaglandin synthesis enzymes *PTGS1/2*, which may contribute to neuroinflammation and altered pain sensitivity in older women. Importantly, our mouse ovariectomy model recapitulated these inflammatory changes, with gene set enrichment analysis showing significant enrichment of inflammatory pathways at 4 months post-OVX (Fig. 4d), closely matching our findings in the late peri-/early post-menopausal women. These hypothalamic inflammatory and glial changes may have functional consequences extending beyond the central nervous system. Activated astrocytes and microglia in thermoregulatory and metabolic hypothalamic nuclei can modulate local neuronal excitability through altered neurotransmitter reuptake, gap junction coupling or cytokine release [42]. The prostaglandin pathway changes we observed (*PTGS1/2, PTGER3*) directly link to autonomic output controlling cutaneous vasodilation, which is the effector mechanism of VMS [43,44]. Furthermore, hypothalamic inflammation can influence systemic metabolic regulation through altered brown adipose tissue thermogenesis and energy expenditure, potentially contributing to the weight gain and metabolic dysregulation commonly observed during the menopausal transition [44,45]. The upregulation of NF-κB-mediated inflammatory signaling in the hypothalamus may also propagate to peripheral tissues through neuroendocrine and autonomic pathways, contributing to the systemic low-grade inflammation ("inflammaging") associated with increased cardiometabolic risk in postmenopausal women [46]. While our study identifies these

pathways as correlatively altered, future mechanistic studies using the OVX model that we describe could directly test these hypothesized central-peripheral links.

We also assessed the changes in the expression of genes involved in reproductive function and thermoregulation across the menopausal transition. As expected, *KISS1* expression increased markedly during menopause[47,48], likely due to loss of estrogen negative feedback, followed by a decline approximately 10 years post-menopause (Fig. 2d). This late-life reduction in kisspeptin signaling may contribute to the eventual decline in gonadotropin levels seen years after menopause onset[49,50]. Intriguingly, *TACR3* expression, which GWAS studies have associated with the severity of VMS[40], showed a distinct pattern, increasing in the years leading up to menopause, peaking around age 45, and then progressively declining (Fig. 2d). Given the proposed role of neurokinin B signaling in triggering vasomotor symptoms[15,51], this expression pattern aligns well with the typical time course of hot flashes, which often begin in perimenopause and resolve within 5-10 years [52]. Thus, our data offer novel insights into the mechanisms that could mediate the decrease in LH levels and termination of hot flashes years after menopause, likely mediated by the significant drop in hypothalamic *KISS1* and *TACR3* expression.

Unbiased differential expression analysis identified two genes, *AKAP5* and *CDKN1A*, as particularly notable markers of the menopausal transition (Fig. 3a-d). *AKAP5* (A-kinase anchoring protein 5) encodes a scaffolding protein involved in synaptic plasticity[53-56] and showed decreased expression in late peri-/early post-menopause ages. *CDKN1A*, encoding the cell cycle regulator p21,[57] was upregulated in late peri-/early post-menopause ages. Interestingly, p21 has been implicated in promoting senescence-associated secretory phenotype (SASP)[57-60], in which senescent cells increase the secretion of inflammatory factors, thus potentially contributing to the inflammatory phenotype we observed. The identification of *CDKN1A* (p21) upregulation in the 51-55 age range suggests that cellular senescence may contribute to hypothalamic aging. Whether this effect is mediated by p21 action at the glial level to promote senescence remains to be investigated.

To determine whether these human transcriptomic changes could be modeled experimentally, we developed a mouse ovariectomy (OVX) model with both short-term (14 day) and long-term (4 month) timepoints. Remarkably, we found that the 4-month post-OVX timepoint recapitulated many key features of the human menopausal transition. LH levels showed an initial increase followed by decline (Fig. 4b), mirroring the menopausal transition[21] in humans. Core body temperature was elevated at 14 days post-OVX but normalized by 4 months (Fig. 4c), paralleling the typical time course of vasomotor symptoms[52]. Transcriptomic analysis of the mouse PH and POA revealed striking similarities to the human dataset. Gene set enrichment analysis showed significant enrichment of inflammatory pathways at 4 months post-OVX (Fig. 4d), closely matching our findings in women undergoing the menopausal transition. Importantly, cross-species comparison demonstrated significant correlation between the transcriptomic changes in our human and mouse datasets (Fig. 5a). Nearly two-thirds of genes differentially expressed in women at age ranges between normal late peri-/early post-menopause and pre-menopause showed concordant changes at 4 months post-OVX in mice, particularly at the level of the PH. This high degree of conservation suggests that our long term OVX model effectively captures key molecular features of the human menopausal transition.

To further validate our model, we examined protein changes in neurokinin B (NKB) expression and neuronal activation. Immunofluorescence staining revealed decreased NKB levels in the arcuate nucleus at both 14 days and 4 months post-OVX (Supplemental Fig. 10a-c,j). This contrasts with the increased *Tac2* mRNA expression we observed and suggests enhanced NKB release and depletion of peptide stores. Importantly, we found reduced c-Fos expression in NKB neurons at 4 months post-OVX (Supplemental Fig. 10m,n), indicating decreased neuronal activity. This aligns with the declining *TACR3* expression seen in postmenopausal women and may reflect adaptive changes to prolonged estrogen deficiency.

Of note, while multiple rodent models exist to study menopausal biology, we selected long-term OVX for this study because it provides precise control over the timing of estrogen withdrawal,

enabling direct alignment of endocrine, thermoregulatory, and transcriptomic changes with human age-associated windows. The VCD (4-vinylcyclohexene diepoxide) model offers an alternative approach that more closely mimics gradual follicular depletion characteristic of spontaneous menopause. However, VCD induces variable timing of ovarian failure (typically 15-160 days depending on dose and strain)[61], residual ovarian interstitial tissue can produce androgens for months[61-63] and repeated dosing and potential off-target effects add variables that are absent in OVX[64,65]. Our OVX model, while representing abrupt surgical menopause, provides a robust and widely validated framework for studying chronic hypoestrogenism and serves as a complementary approach to gradual-depletion models. This model also provides a platform for future studies examining whether estrogen replacement reverses specific transcriptomic changes associated with menopausal-like transitions in the mouse that could be translated into humans.

In conclusion, this study serves as a valuable resource for future investigations into hypothalamic aging and the menopausal transition. We demonstrate that our OVX mice model can provide a physiologically relevant platform for experimental manipulations to determine the functional significance of specific molecular changes observed in the human samples. Together, these insights and tools pave the way for a deeper understanding of menopause and the development of interventions that could significantly improve women's health during this life stage. Nonetheless, several limitations should be considered when interpreting our findings. First, both human and mouse analyses relied on bulk RNA-sequencing, which captures population-averaged gene expression but cannot resolve cell-type-specific transcriptomic changes. While we employed marker-based deconvolution approaches using the Hajdarovic et al. single-cell reference dataset [66], these methods provide estimates rather than direct measurements of cellular composition changes. In this context, the recent publication of comprehensive human hypothalamic single-cell atlases [67,68] provides an important resource for future studies aimed at characterizing cell-type-specific responses to menopause. Second, protein-level validation in our study focused on NKB, c-Fos, and GFAP immunostaining in the mouse model. While

transcriptomic changes provide insight into cellular responses, post-transcriptional regulation may result in discordance between mRNA and protein levels, particularly for secreted neuropeptides where release dynamics affect detectable protein stores, as observed for NKB. Third, the human GTEx dataset presents inherent confounders including variation in postmortem interval, unknown hormone therapy history, and heterogeneous causes of death. We addressed latent sources of variation computationally using Surrogate Variable Analysis (SVA), but cannot fully account for unmeasured biological and technical variables.

Materials and Methods

Mouse studies

Study approval and animals

The animal studies were approved by the Brigham and Women's Hospital Institutional Animal Care and Use Committee (IACUC) in the Center for Comparative Medicine. Adult wild-type (WT) C57/BL6 female mice were group housed under constant conditions of temperature (22–24°C) and light (12:12-hour light:dark cycle), fed with standard mouse chow and ad libitum access to tap water.

Experimental design

To investigate the effects of sex steroids withdrawal on LH levels, core temperature (Tc), and hypothalamic transcriptome profile, intact WT female mice were bilaterally ovariectomized (OVX) to generate mouse models that would resemble pre-menopause, menopause transition and early post-menopause in women. Blood LH levels and Tc were measured at baseline, when females were still intact, at 7-, 14- and 21-days post OVX and at 1-, 2- and 4-months post OVX. Additional females were euthanized before OVX in diestrus (pre-menopause group), after 14 days of OVX (menopause transition group) or after 4 months of OVX (early post-menopause group)

and their posterior (PH) and preoptic area (POA) of the hypothalamus were collected for

transcriptome profile analysis by bulk RNA-seq.

Ovariectomy (OVX)

Adult (3-4 months-old) female mice were bilaterally OVX under isoflurane anesthesia.

Mice were treated with Buprenex SR (0.1 mg/kg, SC) and meloxicam (5 mg/kg, SC) on the day

of surgery. Mice were allowed to recover for at least 1 week before the start of experiments.

Whole blood samples and LH ELISA

For LH measurements, a 4µl blood sample was collected with a pipette. Whole blood was

immediately diluted in 116µl of 0.05% PBST [phosphate buffer saline (Boston Bio Products,

catalog #BM220) containing Tween-20 (Sigma, catalog #P2287)], vortexed, and frozen on dry ice.

Samples were stored at 80˚C until analyzed using an in-house sandwich enzyme-linked

immunosorbent assay as previously described Kreisman, et al. [69].

Core temperature (Tc)

For Tc measurements, intact female mice were implanted (intraperitoneally) with an IPTT-

300 temperature transponder (BioMedic Data Systems, Inc.). Tc from the transponder were

collected with a DAS-8007 Reader and DASHost 8000 software (BioMedic Data Systems, Inc.).

Immunohistochemistry

Animals were anesthetized with a ketamine/xylazine/saline cocktail and transcardially

perfused with saline (0.9% NaCl) followed by 4% paraformaldehyde (PFA) diluted in phosphate

buffer (PB) (Boston BioProducts, Inc). Brains were removed, stored in 4% PFA overnight and then

transferred into sucrose solution (30% sucrose in 0.1 M PB, Thermo Fisher Scientific) at 4˚C. After

sucrose infiltration, the tissue was frozen in Tissue-Tek® O.C.T. Compound (Sakura® Finetek-

4583) and cut on a freezing stage microtome (Microm HM 450, Thermo Fisher Scientific) into 30

μm coronal sections. Free-floating sections were washed in tris buffered saline (TBS, Boston

BioProducts, Inc) and blocked for 1 hour at room temperature in incubation solution [TBS +

0.3%Triton™ X-100 (Millipore Sigma) + 2% normal goat serum (NGS)]. Sections were

subsequently incubated for 48 hours at 4°C with primary antibodies: rabbit-anti neurokinin B

(1:1000, Novus Biologicals, catalog #NB300-201), guinea pig anti-cFos (1:1000, Synaptic

System, catalog #226308) and mouse anti-GFAP (1:300, Cell Signaling, catalog #3670) diluted

in incubation solution. Next, sections were incubated for 1 hour at room temperature with

secondary antibodies: anti-rabbit Alexa Fluor 594 (1:500, Thermo Fisher Scientific, catalog

#A11012), anti-guinea pig Alexa Fluor 488 (1:500, Jackson ImmunoResearch, catalog

#706545148) and anti-mouse donkey Alexa 594 (1:500, Invitrogen, catalog #A32744). Sections

were mounted onto Super-frost plus glass slides (Fisher Laboratories), air dried and cover slipped

with Vectashield mounting medium with DAPI (Vector laboratories).

Microscopy and image analysis

The images were captured by using a fluorescent microscope (Leica DM 2500) in JPG

format and processed using the open-source software ImageJ, Fiji (National Institutes of Health,

Bethesda, Maryland). For image quantification, representative images of the areas of interest (i.e.,

ARC) were delimited using as reference the mouse section from Allen Brain Atlas [70]. Both sides

of bilateral ARC were counted on a similar number of sections and animal, and replicate values

were averaged.

RNA extraction, library preparation, and sequencing

Mouse posterior hypothalamus and preoptic area regions were dissected and flash frozen

prior to extracting RNA using a Zymo Quick RNA microprep kit according to the manufacturer's

instructions. RNA concentration was measured using a Qubit 4.0 fluorometer and the Quibit RNA

HS Assay Kit (ThermoFisher). All samples had RNA integrity numbers (RIN) greater than 7 as

measured on a Bioanalyzer 2100 instrument (Aligent). RNA sequencing libraries were prepared

using the KAPA mRNA HyperPrep–Stranded Kit (Roche) and sequenced on a NovaSeq SP

(Illumina) with 50bp paired-end reads.

RNA-seq data processing and analysis

We performed all analyses using mouse genome build mm39. To pseudoalign reads to

the transcriptome annotation and estimate expression levels of each transcript, we used kallisto[71]

in the tximport package[72] (v1.24.0). We imported the resulting count data (abundance.h5 file) into

R for analysis in DESeq2[24] (v1.36.0). To adjust for unknown, unmodeled or latent sources of

noise in the data we performed surrogate variable analysis (SVA) using the smartSVA package[18]

(v0.1.3) and used SV1 as a covariate in the differential gene expression analysis. Both

pseudoalignment and DESeq2 were performed on the entire transcriptome annotation, but, in

some cases, we restricted our downstream analysis to protein-coding genes, as annotated in

GENCODE version M30[73]. For gene set enrichment analysis (GSEA), we performed analyses

on pre-ranked ordered gene lists using the fgsea package[25] (v1.22.0).

Prediction of protein interactions

The first 30 leading edge genes from the comparison of the POA between female gonadal

intact vs. 4-month post-OVX mice were selected as the target gene list for TNF-α signaling

through NF-κB. The gene list was imported into STRING[74] for network reconstruction,

visualization and enrichment analysis [74]. Medium confidence, as the minimum required interaction

scored, was selected (0.400). Enrichment biological processed defined by gene ontology (GO)

were assessed. Of the top 10 variables identified by the GO terms, highlighted genes related to

thermoregulation and inflammation response to menopause transition are shown.

Human studies

Human hypothalamus inclusion criteria

Bulk RNA-sequencing read count data from human hypothalamus tissue was obtained from publicly available Genotype-Tissue Expression (GTEx) Project[17] data. We included donor tissue as part of our dataset if a sample met the following criteria: 1) donor eligibility criteria (INCEXC) == TRUE, 2) tissue RNA quality score (RIN) >= 6, 3) donors were female (SEX == 2).

Sex and Gender Reporting

All human hypothalamic samples from the GTEx database were from donors recorded as female (biological sex). This study specifically focuses on women's health, examining transcriptomic changes across the female reproductive lifespan from premenopausal through postmenopausal years. Sex was determined by GTEx based on donor records. We use the term "women" throughout to reflect the clinical population most directly affected by natural menopause, while acknowledging that individuals of diverse gender identities with intact ovarian function may also experience menopausal transition.

Human data analysis

After the sub-setting the GTEx RNA expression file for our samples of interest, we adjusted for unknown, latent sources of noise in the data using SVA with the smartSVA package[18] (v0.1.3). Differential expression analyses across age bins were performed using DESeq2[24] (v1.36.0) and GSEA was performed on pre-ranked ordered gene lists using the fgsea package[25] (v1.22.0). Gene expression trajectories across age (and parallel analyses across OVX conditions in mouse) were performed on protein-coding genes using locally estimated scatterplot smoothing (LOESS) similarly to Piehl et al. 2022[75]. Gene level output from LOESS across age was subjected to expression trajectory correlation analysis. Genes were ranked based on their correlation with the

TACR3/Tacr3 expression pattern. STRING interaction analyses were performed as described

above. Given the unavailability of hormone measurements in GTEx, we used chronological age

bins as proxies for reproductive stage, informed by epidemiological data from the Study of

Women's Health Across the Nation (SWAN) where median age at final menstrual period is 51-52

years[76]. Samples were grouped into five age ranges: <45 years (corresponding to typical

premenopausal years), 46-50 years (early perimenopausal age range), 51-55 years (late peri-

/early postmenopausal age range), 56-60 years (postmenopausal years), and 61-65 years (late

postmenopausal years). We acknowledge that individual variation in menopausal timing means

our age-based classification of human samples represents chronological age bins that correspond

to typical pre-, peri-, and postmenopausal years in population studies, but we cannot definitively

assign clinical menopausal staging or vasomotor symptom status to individual donors. The GTEx

consortium did not collect serum samples, precluding direct hormone measurements or

application of recently developed gene-based menopausal biomarkers.

Cell type decomposition analysis

We performed cell type decomposition analyses to estimate the cell type composition of

our bulk RNA-seq datasets from human and mouse using the BisqueRNA package[77] (v1.0.5).

Table S2 lists the marker genes used to annotation cell types[19].

Quantification and statistical analyses

Various statistical tests were used to calculate p values as indicated in the methods

section, figure legends, or text, where appropriate. All statistics were calculated using R software,

version 4.2.1 unless stated otherwise. Differential gene expression was performed using DESeq2

with Benjamini-Hochberg false discovery rate (FDR) correction; genes with adjusted p-values <

0.05 were considered significant. Prior to DE analysis, we performed Surrogate Variable Analysis

(SVA) using the smartSVA package to adjust for unknown latent sources of variation in both

human and mouse datasets. Gene set enrichment analysis (GSEA) was performed using the fgsea package on pre-ranked gene lists, with FDR-corrected normalized enrichment scores (NES) reported. LOESS trajectory analysis was adapted from Piehl et al. 2022 to visualize continuous expression patterns across age; genes with similar trajectories were identified by hierarchical clustering using Euclidean distance. Statistical comparisons of physiological measurements (LH, core temperature) and immunofluorescence quantification used one-way ANOVA with Tukey's post-hoc test or Mann-Whitney U test as appropriate. $P < 0.05$ or FDR < 0.05 was considered statistically significant.

Funding

This work has been supported by the National Institutes of Health (NIH) grant HD090151, DK133760, HD099084 and NIH Research Grant U54 AG062322 (ROSA Center) funded by The National Institute on Aging (NIA) and Office of Research on Women's Health (ORWH) to (ETJ, SAP, LAS, ANF, VMN), The Brit Jepson d'Arbeloff Center on Women's Health (JCBB, DCP)

Author contributions

Conceptualization: JCBB, ETJ, SAP, VMN

Methodology: JCBB, ETJ, SAP, LAS, ANF

Investigation: JCBB, ETJ, SAP

Visualization: JCBB, ETJ, SAP

Supervision: DCP, VMN

Writing—original draft: JCBB, VMN

Writing—review & editing: JCBB, ETJ, SAP, HJ, DCP, VMN

Competing interests Authors declare that they have no competing interests.

Data and materials availability:

• Raw RNA-seq data from mouse posterior hypothalamus and preoptic area regions have

been deposited to dbGAP and are publicly available as of the date of publication.

• This paper analyzes existing, publicly available GTEx consortium data available on the

GTEx Portal (https://www.gtexportal.org/home/).

• Original code to perform analyses has been deposited at Github and is publicly available

as of the date of publication.

• Any additional information required to reanalyze the data reported in this study is available

from the lead contact upon request.

References

1   Davis, S. R. *et al.* Menopause. *Nature Reviews Disease Primers* **1**, 15004 (2015).
https://doi.org/10.1038/nrdp.2015.4
2   Brinton, R. D., Yao, J., Yin, F., Mack, W. J. & Cadenas, E. Perimenopause as a neurological
transition state. *Nature reviews. Endocrinology* **11**, 393-405 (2015).
https://doi.org/10.1038/nrendo.2015.82
3   Rance, N. E. & Uswandi, S. V. Gonadotropin-releasing hormone gene expression is
increased in the medial basal hypothalamus of postmenopausal women. *J Clin*
Endocrinol Metab **81**, 3540-3546 (1996). https://doi.org/10.1210/jcem.81.10.8855798
4   Herbison, A. E. in *Physiology of Reproduction*   (ed J. Neil and E. Knobil) 1415–1482
(Academic Press, 2006).
5   Ameratunga, D., Goldin, J. & Hickey, M. Sleep disturbance in menopause. *Intern Med J*
42, 742-747 (2012). https://doi.org/10.1111/j.1445-5994.2012.02723.x
6   Augoulea, A. *et al.* Psychosomatic and vasomotor symptom changes during transition to
menopause. *Prz Menopauzalny* **18**, 110-115 (2019).
https://doi.org/10.5114/pm.2019.86835
7   Coborn, J. *et al.* Disruption of Sleep Continuity During the Perimenopause: Associations
with Female Reproductive Hormone Profiles. *J Clin Endocrinol Metab* **107**, e4144-e4153
(2022). https://doi.org/10.1210/clinem/dgac447
8   Cohn, A. Y. *et al.* Effects of Sleep Fragmentation and Estradiol Decline on Cortisol in a
Human Experimental Model of Menopause. *J Clin Endocrinol Metab* **108**, e1347-e1357
(2023). https://doi.org/10.1210/clinem/dgad285

9   Greendale, G. A., Karlamangla, A. S. & Maki, P. M. The Menopause Transition and
Cognition. *JAMA* **323**, 1495-1496 (2020). https://doi.org/10.1001/jama.2020.1757

10   Kravitz, H. M. J., H. . Sleep during the perimenopause: a SWAN story. *Obstet Gynecol Clin*
North Am **38**, 567-586 (2011).

11   Landis, C. A. & Moe, K. E. Sleep and menopause. *Nurs Clin North Am* **39**, 97-115 (2004).
https://doi.org/10.1016/j.cnur.2003.11.006

12   Reed, S. D., Newton, K. M., LaCroix, A. Z., Grothaus, L. C. & Ehrlich, K. Night sweats, sleep
disturbance, and depression associated with diminished libido in late menopausal
transition and early postmenopause: baseline data from the Herbal Alternatives for
Menopause Trial (HALT). *Am J Obstet Gynecol* **196**, 591-597 (2007).

13   Koysombat, K., McGown, P., Nyunt, S., Abbara, A. & Dhillo, W. S. New advances in
menopause symptom management. *Best Pract Res Clin Endocrinol Metab* **38**, 101774
(2024). https://doi.org/10.1016/j.beem.2023.101774

14   Freeman, E. W., Sammel, M. D., Lin, H., Liu, Z. & Gracia, C. R. Duration of menopausal hot
flushes and associated risk factors. *Obstet Gynecol* **117**, 1095-1104 (2011).
https://doi.org/10.1097/AOG.0b013e318214f0de

15   Prague, J. K. *et al.* Neurokinin 3 receptor antagonism as a novel treatment for
menopausal hot flushes: a phase 2, randomised, double-blind, placebo-controlled trial.
Lancet **389**, 1809-1820 (2017). https://doi.org/10.1016/S0140-6736(17)30823-1

16   Koebele, S. V. & Bimonte-Nelson, H. A. Modeling menopause: The utility of rodents in
translational behavioral endocrinology research. *Maturitas* **87**, 5-17 (2016).
https://doi.org/10.1016/j.maturitas.2016.01.015

17   Lonsdale, J. *et al.* The Genotype-Tissue Expression (GTEx) project. *Nature Genetics 2013*
45:6 **45** (2013-05-29). https://doi.org/10.1038/ng.2653

18   Chen, J. *et al.* Fast and robust adjustment of cell mixtures in epigenome-wide association
studies with SmartSVA - PubMed. *BMC genomics* **18** (05/26/2017).
https://doi.org/10.1186/s12864-017-3808-1

19   Hajdarovic, K. H. *et al.* Single-cell analysis of the aging female mouse hypothalamus.
Nature Aging 2022 2:7 **2** (2022-07-04). https://doi.org/10.1038/s43587-022-00246-4

20   Tadross, J. A. *et al.* A comprehensive spatio-cellular map of the human hypothalamus.
Nature **639** (2025 Feb 5). https://doi.org/10.1038/s41586-024-08504-8

21   Soares, A. G. *et al.* Longitudinal changes in reproductive hormones through the
menopause transition in the Avon Longitudinal Study of Parents and Children (ALSPAC).
Scientific Reports 2020 10:1 **10** (2020-12-04). https://doi.org/10.1038/s41598-020-
77871-9

22   Khoudary, S. R. E. *et al.* The menopause transition and women's health at midlife: a
progress report from the Study of Women's Health Across the Nation (SWAN).
Menopause (New York, N.y.) **26** (2019/10).
https://doi.org/10.1097/GME.0000000000001424

23   Gold, E. B. *et al.* Factors related to age at natural menopause: longitudinal analyses from
SWAN. *Am J Epidemiol* **178**, 70-83 (2013). https://doi.org/10.1093/aje/kws421

24   Love, M. I. *et al.* Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. *Genome Biology 2014 15:12* **15** (2014-12-05).
https://doi.org/10.1186/s13059-014-0550-8

25 Korotkevich, G. *et al.* Fast gene set enrichment analysis. *BioRxiv* (2021-02-01).
https://doi.org/10.1101/060012
26 The STRING database in 2021: customizable protein-protein networks, and functional
characterization of user-uploaded gene/measurement sets - PubMed. *Nucleic Acids Res*
49 (01/08/2021). https://doi.org/10.1093/nar/gkaa1074
27 Blomqvist, A. & Engblom, D. Neural Mechanisms of Inflammation-Induced Fever. *The*
Neuroscientist **24** (2018/08). https://doi.org/10.1177/1073858418760481
28 Prostaglandin production selectively in brain endothelial cells is both necessary and
sufficient for eliciting fever - PubMed. *Proc Natl Acad Sci USA* **119** (10/25/2022).
https://doi.org/10.1073/pnas.2122562119
29 Tickerhoof, M. *et al.* Postmortem tissue biomarkers of menopausal transition. *Molecular*
psychiatry (2025 Aug 27). https://doi.org/10.1038/s41380-025-03177-9
30 Butler, T. *et al.* Positron Emission Tomography reveals age-associated hypothalamic
microglial activation in women. *Sci Rep* **12**, 13351 (2022).
https://doi.org/10.1038/s41598-022-17315-8
31 Tacutu, R. *et al.* Human Ageing Genomic Resources: new and updated databases. *Nucleic*
Acids Res **46** (2018/01/01). https://doi.org/10.1093/nar/gkx1042
32 O'Day, D. H. Calmodulin Binding Proteins and Alzheimer's Disease: Biomarkers,
Regulatory Enzymes and Receptors That Are Regulated by Calmodulin. *Int J Mol Sci* **21**
(2020/10). https://doi.org/10.3390/ijms21197344
33 Kim, J., Basak, J. M. & Holtzman, D. M. The Role of Apolipoprotein E in Alzheimer's
Disease. *Neuron* **63** (2009/08/13). https://doi.org/10.1016/j.neuron.2009.06.026
34 Bateman, R. J. *et al.* Clinical and Biomarker Changes in Dominantly Inherited Alzheimer's
Disease. *New England Journal of Medicine* **367** (2012-08-30).
https://doi.org/10.1056/NEJMoa1202753
35 SIRT1 in Neurodevelopment and Brain Senescence. *Neuron* **81** (2014/02/05).
https://doi.org/10.1016/j.neuron.2014.01.028
36 Catechol-O-methyltransferase polymorphisms and some implications for cognitive
therapeutics - PubMed. *NeuroRx : the journal of the American Society for Experimental*
NeuroTherapeutics **3** (2006 Jan). https://doi.org/10.1016/j.nurx.2005.12.010
37 Smith, J. T., Cunningham, M. J., Rissman, E. F., Clifton, D. K. & Steiner, R. A. Regulation of
Kiss1 gene expression in the brain of the female mouse. *Endocrinology* **146**, 3686-3692
(2005). https://doi.org/10.1210/en.2005-0488
38 Goodman, R. L., Herbison, A. E., Lehman, M. N. & Navarro, V. M. Neuroendocrine control
of gonadotropin-releasing hormone: Pulsatile and surge modes of secretion. *J*
Neuroendocrinol **34**, e13094 (2022). https://doi.org/10.1111/jne.13094
39 J, P., R, K., M, P. & H, S. Changes in proinflammatory cytokine activity after menopause -
PubMed. *Endocr Rev* **23** (2002 Feb). https://doi.org/10.1210/edrv.23.1.0456
40 Crandall, C. J. *et al.* Association of genetic variation in the tachykinin receptor 3 locus
with hot flashes and night sweats in the Women's Health Initiative Study. *Menopause* **24**,
252-261 (2017). https://doi.org/10.1097/gme.0000000000000763
41 Franceschi, C., Garagnani, P., Vitale, G., Capri, M. & Salvioli, S. Inflammaging and 'Garb-
aging'. *Trends Endocrinol Metab* **28**, 199-212 (2017).
https://doi.org/10.1016/j.tem.2016.09.005

42 Pascual, O., Ben Achour, S., Rostaing, P., Triller, A. & Bessis, A. Microglia activation
triggers astrocyte-mediated modulation of excitatory neurotransmission. *Proc Natl Acad*
Sci U S A **109**, E197-205 (2012). https://doi.org/10.1073/pnas.1111098109
43 Nakamura, Y. *et al.* Prostaglandin EP3 receptor-expressing preoptic neurons
bidirectionally control body temperature via tonic GABAergic signaling. *Sci Adv* **8**,
eadd5463 (2022). https://doi.org/10.1126/sciadv.add5463
44 Aten, S., Lynch, N., Saper, C. B. & Machado, N. L. S. A brain-body perspective on
thermoregulatory adaptation. *Curr Biol* **35**, R1016-r1028 (2025).
https://doi.org/10.1016/j.cub.2025.09.023
45 Arruda, A. P., Milanski, M. & Velloso, L. A. Hypothalamic inflammation and
thermogenesis: the brown adipose tissue connection. *J Bioenerg Biomembr* **43**, 53-58
(2011). https://doi.org/10.1007/s10863-011-9325-z
46 Zhang, G. *et al.* Hypothalamic programming of systemic ageing involving IKK-β, NF-κB
and GnRH. *Nature* **497**, 211-216 (2013). https://doi.org/10.1038/nature12143
47 Rance, N. E. Menopause and the human hypothalamus: evidence for the role of
kisspeptin/neurokinin B neurons in the regulation of estrogen negative feedback.
Peptides **30**, 111-122 (2009). https://doi.org/10.1016/j.peptides.2008.05.016
48 Rometo, A. M., Krajewski, S. J., Voytko, M. L. & Rance, N. E. Hypertrophy and increased
kisspeptin gene expression in the hypothalamic infundibular nucleus of postmenopausal
women and ovariectomized monkeys. *J Clin Endocrinol Metab* **92**, 2744-2750 (2007).
https://doi.org/10.1210/jc.2007-0553
49 Hall, J. E. Neuroendocrine physiology of the early and late menopause. *Endocrinology*
and Metabolism Clinics **33** (2004/12/01). https://doi.org/10.1016/j.ecl.2004.08.002
50 Hall, J. E., Lavoie, H. B., Marsh, E. E. & Martin, K. A. Decrease in Gonadotropin-Releasing
Hormone (GnRH) Pulse Frequency with Aging in Postmenopausal Women*. *The Journal*
of Clinical Endocrinology & Metabolism **85** (2000/05/01).
https://doi.org/10.1210/jcem.85.5.6612
51 Christ, J. P., Navarro, V. M. & Reed, S. D. Nonhormonal Therapies for Menopausal
Vasomotor Symptoms. *Jama* **330**, 1278-1279 (2023).
https://doi.org/10.1001/jama.2023.15965
52 Avis, N. E. *et al.* Duration of Menopausal Vasomotor Symptoms. *JAMA Internal Medicine*
175 (2015/04/01). https://doi.org/10.1001/jamainternmed.2014.8063
53 He, Z. *et al.* Novel insight into the role of A-kinase anchoring proteins (AKAPs) in
ischemic stroke and therapeutic potentials. *Biomedicine & Pharmacotherapy* **175**,
116715 (2024). https://doi.org/https://doi.org/10.1016/j.biopha.2024.116715
54 Koster, K. P. *et al.* Akap5 links synaptic dysfunction to neuroinflammatory signaling in a
mouse model of infantile neuronal ceroid lipofuscinosis. *Front Synaptic Neurosci* **16**,
1384625 (2024). https://doi.org/10.3389/fnsyn.2024.1384625
55 Sanderson, J. L. & Dell'Acqua, M. L. AKAP signaling complexes in regulation of excitatory
synaptic plasticity. *Neuroscientist* **17**, 321-336 (2011).
https://doi.org/10.1177/1073858410384740
56 Weisenhaus, M. *et al.* Mutations in AKAP5 disrupt dendritic signaling complexes and
lead to electrophysiological and behavioral phenotypes in mice. *PloS one* **5**, e10325
(2010). https://doi.org/10.1371/journal.pone.0010325

57 Engeland, K. Cell cycle regulation: p53-p21-RB signaling. *Cell Death & Differentiation* **29**,
946-960 (2022). https://doi.org/10.1038/s41418-022-00988-z
58 Wagner, K. D. & Wagner, N. The Senescence Markers p16INK4A, p14ARF/p19ARF, and
p21 in Organ Development and Homeostasis. *Cells* **11** (2022).
https://doi.org/10.3390/cells11121966
59 Wang, L., Lankhorst, L. & Bernards, R. Exploiting senescence for the treatment of cancer.
Nature Reviews Cancer **22**, 340-355 (2022). https://doi.org/10.1038/s41568-022-00450-
9
60 Chinta, S. J. *et al.* Cellular senescence and the aging brain. *Exp Gerontol* **68**, 3-7 (2015).
https://doi.org/10.1016/j.exger.2014.09.018
61 Lohff, J. C., Christian, P. J., Marion, S. L. & Hoyer, P. B. Effect of duration of dosing on
onset of ovarian failure in a chemical-induced mouse model of perimenopause.
Menopause **13**, 482-488 (2006). https://doi.org/10.1097/01.gme.0000191883.59799.2e
62 Van Kempen, T. A., Milner, T. A. & Waters, E. M. Accelerated ovarian failure: a novel,
chemically induced animal model of menopause. *Brain Res* **1379**, 176-187 (2011).
https://doi.org/10.1016/j.brainres.2010.12.064
63 Mayer, L. P., Devine, P. J., Dyer, C. A. & Hoyer, P. B. The follicle-deplete mouse ovary
produces androgen. *Biol Reprod* **71**, 130-138 (2004).
https://doi.org/10.1095/biolreprod.103.016113
64 Cao, L. B. *et al.* Systemic changes in a mouse model of VCD-induced premature ovarian
failure. *Life Sci* **262**, 118543 (2020). https://doi.org/10.1016/j.lfs.2020.118543
65 Cao, L. B. *et al.* Hormone-Like Effects of 4-Vinylcyclohexene Diepoxide on Follicular
Development. *Front Cell Dev Biol* **8**, 587 (2020).
https://doi.org/10.3389/fcell.2020.00587
66 Hajdarovic, K. H. *et al.* Single-cell analysis of the aging female mouse hypothalamus. *Nat*
Aging **2**, 662-678 (2022). https://doi.org/10.1038/s43587-022-00246-4
67 Siletti, K. *et al.* Transcriptomic diversity of cell types across the adult human brain.
Science **382**, eadd7046 (2023). https://doi.org/doi:10.1126/science.add7046
68 Tadross, J. A. *et al.* A comprehensive spatio-cellular map of the human hypothalamus.
Nature **639**, 708-716 (2025). https://doi.org/10.1038/s41586-024-08504-8
69 Kreisman, M. J., McCosh, R. B. & Breen, K. M. A Modified Ultra-Sensitive ELISA for
Measurement of LH in Mice. *Endocrinology* **163** (2022/09).
https://doi.org/10.1210/endocr/bqac109
70 Wang, Q. *et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D Reference
Atlas. *Cell* **181**, 936-953 e920 (2020). https://doi.org/10.1016/j.cell.2020.04.007
71 Bray, N. L. *et al.* Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology*
2016 34:5 **34** (2016-04-04). https://doi.org/10.1038/nbt.3519
72 Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences - PubMed. *F1000Res* **4** (12/30/2015).
https://doi.org/10.12688/f1000research.7563.2
73 Frankish, A. *et al.* GENCODE reference annotation for the human and mouse genomes.
Nucleic Acids Res **47** (2019/01/01). https://doi.org/10.1093/nar/gky955

74 Szklarczyk, D. *et al.* STRING v11: protein-protein association networks with increased
coverage, supporting functional discovery in genome-wide experimental datasets.
Nucleic Acids Res **47**, D607-D613 (2019). https://doi.org/10.1093/nar/gky1131
75 Piehl, N. *et al.* Cerebrospinal fluid immune dysregulation during healthy brain aging and
cognitive impairment. *Cell* **185** (2022/12/12). https://doi.org/10.1016/j.cell.2022.11.019
76 El Khoudary, S. R. *et al.* The menopause transition and women's health at midlife: a
progress report from the Study of Women's Health Across the Nation (SWAN).
Menopause **26**, 1213-1227 (2019). https://doi.org/10.1097/gme.0000000000001424
77 Jew, B. *et al.* Accurate estimation of cell composition in bulk expression through robust
integration of single-cell information. *Nature Communications 2020 11:1* **11** (2020-04-
24). https://doi.org/10.1038/s41467-020-15816-6
762

Figure 1. Gene expression trajectories in the human female hypothalamus across chronological age.
a) Demographic overview of the GTEx donor subset used in this study.
b-f) Sets of genes grouped by hierarchical clustering and displayed using LOESS trajectories exhibit distinct patterns, including, **b)** constant expression across age, **c)** decreased expression across age, **d)** increased expression across age, **e)** decreased expression during the menopausal transition, **f)** increased expression during the menopausal transition.

Figure 2. Expression of reproductive- and aging-related genes during the menopausal transition in the human hypothalamus.

a) Distribution of ages within the donor cohort and the defined age groupings for subsequent analyses.

b) Heatmap of normalized enrichment scores (NES) for Hallmark gene sets in which there was a statistically significant pathway enrichment in at least one of the three comparisons; significant enrichments indicated by asterisks (* p-adj <0.05; ** p-adj < 0.01; *** p-adj < 0.001)

c) STRING analysis of TNF-alpha hallmark pathway leading edge genes between the late peri-/early post-menopausal vs. pre-menopausal groups.

d) LOESS trajectories of KNDy genes and their receptors, thermoregulatory genes, astrocyte-, and microglia-marker genes across age in human hypothalamus tissue.

Figure 3. Identification of differentially expressed genes between the late peri-/early post-menopausal donors and the pre-menopausal donors.

a) Volcano plot highlighting significantly differentially expressed genes between late peri-/early post-menopausal and pre-menopausal groups. Genes with a FDR < 0.05 are highlighted in pink.
b) Expression of genes across age groups with a FDR < 0.05.
c) LOESS trajectory of *AKAP5* across age.
d) LOESS trajectory of *CDKN1A* across age.

Figure 4

Figure 4. Physiological and transcriptomic changes in post-ovariectomy mouse models recapitulate key features of reproductive aging.

a) Schematic representation of the experimental design.

b) Luteinizing Hormone (LH) concentration in serum isolated from whole blood 7-, 14-, 21-days post-OVX and 1-, 2- and 4-months post-OVX in wild-type female mice (n=20, per condition).

c) Core temperature (ºC) measurements at 7-, 14-, 21-days post-OVX and 1-, 2- and 4-months post-OVX in wild-type female mice (n=20, per condition). b-c) Data values are shown as mean ± SEM. Groups with different letters are significantly from each other, $P < 0.05$ by One-way ANOVA/Tukey.

d) Heatmap of normalized enrichment scores (NES) for Hallmark gene sets in the Posterior Hypothalamus (PH) and the Preoptic Area (POA); significant enrichments indicated by asterisks (* *p*-adj <0.05; ** *p*-adj < 0.01; *** *p*-adj < 0.001)

e-h) Volcano plots highlighting significantly differentially expressed genes between labeled conditions and regions. Genes with a FDR < 0.05 are labeled in pink.

i) Expression of PH DE genes identified in **b** across conditions.

j) Expression of POA DE genes identified in **d** across conditions.

Figure 5. Comparative analysis of human and mouse hypothalamic reproductive aging datasets.

a) Heatmap of GSEA pathway correlations between human and mouse datasets; *p*-value from Pearson's correlation.

b) Overlap of *TACR3/Tacr3* positively correlated co-expressed genes between human and mouse; *p*-value from hypergeometric test.

Expression of identified DE genes from the human dataset across conditions in the mouse **c)** PH and **d)** POA datasets.

e) Expression of identified DE genes from the mouse PH across age groups in the human dataset.

f) Expression of identified DE genes from the mouse POA across age groups in the human dataset.