Volatiles from low R:FR-treated maize plants increase the emission of herbivore-induced plant volatiles in their neighbors

Rocío Escobar-Bravo[1,2] *, Bernardus C.J. Schimmel[1,3], Matthias Erb[1]

[1]Institute of Plant Sciences, University of Bern, 3013, Bern, Switzerland.

[2]Current address: Enza Zaden, Haling 1-E,1602 DB, Enkhuizen, The Netherlands

[3]Current address: Institute of Environmental Biology, Utrecht University, Padualaan 8, 3584 CH, Utrecht, The Netherlands

*Author for correspondence: Rocío Escobar-Bravo (rocioebe@hotmail.com)

ABSTRACT

Low Red (R) to Far Red (FR) light ratios, a light signal associated with vegetation shade, can prompt intact maize (*Zea mays*) plants to constitutively emit more volatiles when exposed to herbivory-induced plant volatiles (HIPVs). Here we investigated how simulated shading affects priming responses in the context of volatile-mediated plant-plant interactions. Receiver maize plants were exposed to either constitutive volatile organic compounds (cVOCs) or HIPVs from emitter maize plants, while we manipulated R:FR light conditions of receivers or emitters. Priming responses in the receivers were then assessed by measuring real-time volatile emissions following simulated herbivory. We show that low R:FR light enhances HIPVs emissions in plants previously exposed to HIPVs from neighbours independently of the light conditions of emitters. We also demonstrate that both cVOCs and HIPVs emitted by maize grown under low R:FR amplify HIPVs emissions in their neighbours. This amplified response could not be explained by FR-mediated changes in the release of green leaf volatiles or terpenoids by emitters, thus suggesting the involvement of other VOCs. We conclude that volatile-mediated plant-plant interactions can be expected to become more intense in denser canopies due to light-mediated amplification of volatile emission and responsiveness.

Key Words

Far-red light, herbivores, maize, plant defenses, priming, volatiles.

INTRODUCTION

Plants can sense the presence of neighbors through light, chemical, and mechanical cues (Huber et al., 2020; Ninkovic et al., 2021; Pantazopoulou et al., 2023). This ability to detect nearby plants helps them to assess their competitive environment, including the availability of light and soil nutrients, as well as the health and potential threats posed by pests and pathogens. By integrating these signals, plants can strategically allocate their resources towards growth or defense, thus enhancing their chances of survival.

Neighbor detection through light cues occurs via the photoreceptor-mediated perception of light spectral changes in the canopy. Blue (λ =400-500 nm) and red (R) (λ = 600-700 nm) light wavelengths are preferentially absorbed by photosynthetic pigments of plant leaves while far red (FR) light is mainly reflected or transmitted. In dense canopies, this results in a reduction in R:FR ratios, a light cue that can promote upward bending of the leaves (hyponasty) and stem elongation (Pierik and Ballaré 2021). These photomorphogenic changes help the plant to reach more illuminated areas in the canopy and outcompete their neighbors (Pierik and Ballaré 2021) In addition, FR photons can support growth by promoting photosynthesis in dicotyledonous and C3 and C4 monocotyledonous species, including maize (*Zea mays*), when added to a background of white light (Zhen and Bugbee 2020; Escobar￿Bravo et al. 2024; Huber et al. 2024).

Beyond light, plants also perceive their environment through volatile organic compounds (VOCs) (Wang and Erb 2022). These compounds include volatile hormones (e.g., ethylene, methyl salicylate, and methyl jasmonate), terpenoids, benzenoids (e.g., indole), phenylpropanoids, and fatty acid-derived molecules such as green leaf volatiles (GLVs). Plant VOCs can be constitutively emitted and modulated by the plant interactions with the environment (Piesik et al. 2015; Escobar-Bravo et al. 2023). Changes in their emission therefore reflect the plant status and the abiotic and/or biotic stresses it encounters. Upon herbivory, for instance, plants can emit herbivory-induced plant volatiles (HIPVs) that repel the attacker, attract natural predators of the herbivore, and/or alter the defense status in non-attacked neighbors (Turlings and Erb 2018; Hu et al. 2021).

Some HIPVs, such as the green leaf volatiles (*Z*)-3-hexenal [(*Z*)-3-HAL], (*Z*)-3-hexen-1-ol [(*Z*)-3-HOL], or (*Z*)-3-hexenyl acetate [(*Z*)-3-HAC], can directly induce the emission of volatiles in intact plants and prime their volatile emissions (Farag and Pare 2002; Engelberth et al. 2004; Hu et al. 2019; Escobar￿Bravo et al. 2024). Similarly, other HIPVs like indole, the monoterpene β-ocimene, and the homoterpene (*E*)-4,8-dimethyl-1,3,7-nonatriene (DMNT) can induce or prime plant defenses (Erb et al. 2015; Jing et al. 2021; Onosato et al. 2022). Defense priming can confer ecological benefits to plants, as the exposure to the priming stimulus prepares them to display a faster, stronger, and more lasting defense response when the actual herbivore or pathogen attack takes place (Conrath et al. 2015; Martinez-Medina et al. 2016). The mechanisms of volatile-mediated defense priming downstream of volatile perception have been amply characterized. Once HIPVs get into plant tissues, they induce the expression of biotic and abiotic stress-responsive genes in a Ca^{2+} dependent manner, including the activation of the jasmonic acid (JA) defense signaling pathway that controls the emission of HIPVs (Zebelo et al. 2012; Christensen et al. 2013; Engelberth et al. 2004; Hu et al. 2019; Ye et al. 2019; Jing et al. 2021; Aratani et al. 2023; Aguirre et al. 2023; Wang et al. 2023).

How VOC-mediated transfer of information between plants occurs in dense canopies is not
well known. The FR-enriched light environment within vegetation canopies might play a crucial role in
these interactions by altering (1) the emitter's volatile emissions, (2) the ability of the receiver to
perceive and respond to the volatile cue/s, or (3) both. For instance, Kegge et al. (2015) showed that
barley (*Hordeum vulgare*) emits different volatiles under low R:FR ratios, influencing the carbon
allocation in neighboring plants. Aguirre et al. (2023) reported that maize exposure to (*Z*)-3-HAC in the
dark does not prime plant defenses, suggesting that light-mediated induction of stomatal opening is
required for signal integration of (*Z*)-3-HAC in receiver plants. More recently, we have demonstrated
that low R:FR light conditions can change the perception of HIPVs in maize plants (Escobar￭Bravo et
al. 2024). Our study showed that FR-light enrichment prompts intact maize plants to emit more
volatiles when exposed to HIPVs from *Spodoptera littoralis*- infested neighbors or to (*Z*)-3-HAC alone.
We also showed that low R:FR-treated plants pre-exposed to (*Z*)-3-HAC displayed stronger HIPVs
emissions after simulated herbivory and, therefore, a stronger priming response. Yet, whether low
R:FR-induced priming response also occurs in plants exposed to the whole bouquet of HIPVs from
herbivore-infested neighbors was not determined. Furthermore, whether FR light enrichment can alter
volatile-mediated interactions between maize plants by inducing changes in the VOCs emissions of
emitter plants is unknown.

Here, we have investigated whether low R:FR affects HIPVs-induced priming responses in
maize by altering (1) the volatile perception of the receivers, and/or (2) the volatile emissions of the
emitters. For this, we first exposed plants to low R:FR ratios as a light cue for vegetation shade or
high R:FR ratios as control light conditions, and we determined volatile emissions in simulated
herbivory-induced plants that were previously exposed to constitutive volatile organic compounds
(cVOCs) or HIPVs emissions from intact or *S. littoralis*-infested plants, respectively. Next, we
conducted a multifactorial experiment where we tested whether HIPVs emitted by intact or simulated
herbivory-induced maize plants growing under low or high R:FR conditions trigger different defense
priming responses in the receivers, and whether these differences could be explained by changes in
the GLVs emissions of the emitters. Our findings show that maize seedlings exposed to enriched FR
conditions not only display a higher emission of HIPVs after exposure to volatile cues from neighbors,
but they also emit volatiles that increase the emission of HIPVs in neighboring plants.

METHODS & MATERIALS

Plants

Experiments were conducted with the maize (*Zea mays*) inbred line B73. Seeds were sown in
transparent cylindric plastic pots (11x4 cm) filled with commercial soil (Selmaterra, BiglerSamen,
Switzerland) and wrapped with aluminium foil to prevent root exposure to light. Plants were grown in a
ventilated greenhouse supplemented with artificial lighting under 50-70% relative humidity, 14/10 h
light/dark photoperiod, and 14-22/10-14°C day/night temperatures.

Insects

Eggs of *Spodoptera littoralis* were provided by Oliver Kindler (Syngenta, Stein, CHE) and the larvae were reared on artificial diet as described in (Maag et al. 2014). *S. littoralis* oral secretions were collected from third- to fourth-instar larvae as described in Hu *et al.*, 2019 and stored at – 80°C until use.

Light Treatments

Light treatments were conducted in a custom-made high-throughput phenotyping platform designed for volatile analyses previously described in Escobar☐Bravo et al. (2024). Eleven-day old maize seedlings were exposed to low (~0.5) or high (~2.65) R:FR ratios (R, λ = 600-700 nm; FR, λ = 700-800 nm) by providing supplemental FR to the same white light background (120 ± 15 µmol m^{-2} s^{-1}) (Supplemental Fig. S1) with a photoperiod of 16/8 h light/dark. This low R:FR corresponds to the ratios observed in high density maize canopies (Maddonni et al. 2002), where low PAR levels as the ones used in our experimental set-up are commonly observed (Xue et al. 2016). Light treatments were separated by white opaque curtains and started between 9:00 and 10:00 AM two days before the plant-plant interaction experiments.

Plant-Plant Interaction Experiments

In the first experiment (Supplemental Fig. S2), we exposed receiver plants to low or high R:FR light conditions and the emitters to low R:FR light conditions. After two days, low R:FR-treated emitter plants were infested with ten second-instar *S. littoralis* larvae or left uninfested. Right after infestation, low and high R:FR-treated receiver plants were exposed to volatiles emitted by *S. littoralis*-infested or non-infested low R:FR treated plants. The chambers of emitter and receiver plants were connected using Teflon tubing. Clean air flow moved from the emitter's chamber towards the receiver's chamber. After 20 h, the Teflon tubing connecting emitters and receivers' chambers were removed, and receiver plants were induced with simulated herbivory. Receiver's volatile emissions were then recorded in time-series analysis using a PTR-TOF-MS system. In the second experiment (Supplemental Fig. S3), we employed the same set-up, but emitters were exposed to high R:FR light conditions.

In the third experiment (Supplemental Fig. S4), we conducted a multifactorial experiment where we exposed both emitter and receiver plants to high or low R:FR conditions for 2 days. Next, emitter and receivers' chambers were connected via Teflon tubing as in previous experiments and the emitter plants were immediately induced with simulated herbivory or left intact. By using simulated herbivory, we standardized the leaf damage and minimized differences in herbivore-associated damage patterns between low and high R:FR-treated emitter plants. After 6 h, the Teflon tubing connecting emitters and receivers' chambers was removed, and all the receiver plants were induced with simulated herbivory. Although the duration of the exposure to volatiles was shorter than in Experiments 1 and 2, our previous work has shown that HIPVs emissions in simulated herbivory induced maize plants steadily increases and reaches their maximum over the first six hours after induction, and that exposure to GLVs for 6 h can prime volatile emissions (Escobar☐Bravo et al.

2024). Receiver's volatile emissions were recorded in time-series analysis using a PTR-TOF-MS
system. This experiment was repeated three times using three replicates per combined treatment
each time. Data from the three repetitions were pooled prior to the statistical analysis.

PTR-TOF-MS Volatile Emission Analysis

Plant volatile emission was determined using the automated high-throughput real-time phenotyping
platform and the methods described in Escobar□Bravo et al. (2024)

Simulated Herbivory

Two leaves (leaf 2 and 3 from the bottom) per plant were wounded over an area of ca. 0.5 cm^2 parallel
to the central vein using a haemostat followed by the application of 10 µL of *S. littoralis* larval oral
secretions (diluted 1:1 in autoclaved distilled water) in each wounding site. This treatment induces
plant defence responses comparable to real herbivory (Erb et al. 2009).

Statistical Analysis

Data analysis was conducted in R version 4.0.4 (R core Team, 2016). Effects of (1) sampling time, (2)
light receiver, (3) light emitter, (4) volatiles from the emitter, and (5) their interactions on volatile
emissions in the time course experiments (i.e., Figures 1, 2, 3, and 5) were tested in linear
mixed-effects models (LME) using the nlme package. Individual plants were included in the model as
a random factor and a correlation structure when autocorrelation among residuals was found to be
significant ($p < 0.05$). Fitted models were subjected to type III analyses of variance (ANOVAs) to
produce a summary of the F- and p statistics (car package) (Fox et al. 2012). When needed, data
from volatiles signatures were log10 or squared root transformed prior analysis to correct for
heteroscedasticity. Cumulative emission of the sum of all volatile's signatures depicted in Fig. 4 was
analysed by using a mixed-effects model where we the replication of the experiment was included as
random factor. Cumulative emission of individual volatiles over time depicted in Fig. S5 and S6 was
analysed by using linear models. Differences among treatments were tested by the calculation of
estimated marginal means (EMMs) followed by pairwise comparisons using Tukey's Honest
Significant Difference (HSD) test (emmeans package; Lenth, 2021).

RESULTS

Low R:FR Light Enhances HIPVs-Mediated Priming of Volatile Emissions in Maize

Low R:FR light ratios can increase the emission of volatiles in intact maize plants when these are
exposed to HIPVs emitted by low R:FR-treated neighbors (Escobar□Bravo et al. 2024). To determine
whether low R:FR and pre-exposure to HIPVs emitted by herbivore-infested neighbors affects
subsequent maize responses to herbivory, we conducted a series of plant-plant interactions
experiments.
In the first experiment, we exposed 11-day old B73 maize seedlings to low or high R:FR light
conditions for two days and measured their volatile emissions after exposure to volatiles emitted by *S.*
littoralis-infested or non-infested low R:FR treated plants followed by induction with wounding and the
application of *S. littoralis* oral secretions (Fig. S2 and Fig. 1ab). Upon simulated herbivory, emission of

all the volatile signatures increased over time (Fig. 1b). This induction was stronger under low R:FR
irrespective of the pre-exposure to emitters' volatiles. Emissions of monoterpenes, sesquiterpenes,
DMNT, and TMTT [(E, E)-4,8,12-trimethyltrideca-1,3,7,11-tetraene] were higher in low R:FR treated
plants compared to high R:FR treated plants, except for indole. Light also influenced maize responses
to volatiles. The emissions of sesquiterpenes, monoterpenes, TMTT, and DMNT were higher in low
R:FR-treated plants exposed to HIPVs from herbivore-infested plants, whereas no induction, or even
the opposite pattern, was observed in high R:FR-treated plants. Cumulative emissions of
monoterpenes and sesquiterpenes were higher in low R:FR treated plants pre-exposed to HIPVs in
comparison to their controls pre-exposed to cVOCs (Fig. S5)
In the second experiment, we used the same set-up, but plants were instead exposed to
volatiles emitted by high R:FR treated plants that were either infested with *S. littoralis* larvae or non-
infested (Fig S3, and Fig. 2). Emission of all the volatiles increased over time upon simulated
herbivory in low and high R:FR treated plants. Unlike the first experiment, plants exposed to low R:FR
and volatiles from high R:FR treated plants emitted less indole, monoterpenes, DMNT and TMTT than
plants exposed to high R:FR and the same volatiles (Fig. 2 and Fig. S6). Prior exposure to volatiles
from high R:FR treated plants infested with *S. littoralis* larvae slightly enhanced the emission of
monoterpenes and sesquiterpenes in low R:FR-treated plants upon simulated herbivory. In plants
irradiated with high R:FR ratios, prior exposure to HIPVs slightly decreased the emission of
monoterpenes, sesquiterpenes, and DMNT over time, and the opposite was observed for indole.
Similar patterns were found when the cumulative emissions of individual volatiles were analyzed (Fig.
S6). Low R:FR treated plants emitted less indole, monoterpenes and TMTT than high R:FR-treated
plants upon exposure to emitter's volatiles and simulated herbivory.
These results suggest that low R:FR can enhance plant responses to HIPVs, but this
induction seems to be stronger when plants were exposed to volatiles emitted by neighbors growing
under low R:FR ratios

.

Figure 1. Low R:FR enhances priming of monoterpenes, sesquiterpenes and TMTT emissions in maize plants pre-exposed to HIPVs from low R:FR treated plants. (a) Schematic overview of the emitters and receivers' treatments. (b) Time-series analysis of volatile emissions (mean ± SEM, $n = 6$) collected from receiver plants pre-exposed to cVOCs or HIPVs from low R:FR-treated plants during 20 h and followed by simulated herbivory (wounding + oral secretions from *S. littoralis*). The effects of sampling time (T), light treatment of receivers (Light), volatiles from emitters (Vol), and their interactions on receivers' volatile emissions were tested using linear mixed-effects models. Statistically significant effects are shown in each graph (*$p < 0.05$, ** $p < 0.001$, *** $p < 0.001$). DMNT and TMTT stand for the homoterpenes (*E*)-4,8-dimethyl-1,3,7-nonatriene and (*E, E*)-4,8,12-trimethyltrideca-1,3,7,11-tetraene, respectively. AU refers to arbitrary units.

Figure 2. Low R:FR enhances priming of monoterpenes and sesquiterpenes emissions in maize plants pre-exposed to HIPVs from high R:FR treated plants. (a) Schematic overview of the emitters and receivers' treatments. (b) Time-series analysis of volatile emissions (mean ± SEM, *n* = 6) collected from receiver plants pre-exposed to cVOCs or HIPVs from low R:FR-treated plants during 20 h and followed by simulated herbivory (wounding + oral secretions from *S. littoralis*). The effects of sampling time (T), light treatment of receivers (Light), volatiles from emitters (Vol), and their interactions on receivers' volatile emissions were tested using linear mixed-effects models. Statistically significant effects are shown in each graph (**p* < 0.05, ** *p* < 0.001, *** *p* < 0.001). DMNT and TMTT stand for the homoterpenes (*E*)-4,8-dimethyl-1,3,7-nonatriene and (*E, E*)-4,8,12-trimethyltrideca-1,3,7,11-tetraene, respectively. AU refers to arbitrary units.

Volatiles Emitted by Low R:FR Treated Maize Plants Increase the Emission Of Herbivore-Induced Plant Volatiles in Receiver Plants

To further test whether maize plants respond differently to volatiles emitted by plants under low R:FR ratios (i.e., simulated shade) and control for potential HIPVs differences arising from herbivore feeding patterns, we conducted a factorial experiment where high and low R:FR-treated plants were exposed to volatiles emitted by low or high R:FR treated plants that were either induced with simulated herbivory or left intact (Supplemental Fig. S4). This set-up allowed us to compare in the same experiment the magnitude of the priming induction when the emitters differ in their light environment conditions but similar herbivory induction. After 6 h of exposure to constitutive or herbivory-induced volatiles from emitters, receiver plants were induced with simulated herbivory and their volatile emissions were determined by PTR-TOF-MS in time series analyses.

For the sake of clarity, we present the results of this experiment in two panels grouped by the light treatment of the receiver (Fig. 3). Maize plants exposed to low R:FR responded differently to volatiles emitted by low and high R:FR-treated plants (Fig. 3a). Upon simulated herbivory, monoterpenes, sesquiterpenes and linalool emissions were higher when low R:FR-treated plants were exposed to volatiles emitted by low R:FR-treated plants than by emitter plants irradiated with high R:FR (Fig. 3a). This induction was statistically significant for sesquiterpenes, and it was independent of the herbivory treatment of the emitters. In addition, indole emissions were increased in low R:FR-treated plants when they were pre-exposed to volatiles from not-wounded low R:FR-treated plants.

Receiver plants exposed to high R:FR ratios and volatiles emitted by low R:FR-treated plants showed a slight increase in the emission of monoterpenes and sesquiterpenes, but this was not statistically significant for any of the volatile signatures (Fig. 3b).

Further analysis on the sum of all the volatiles emitted over the sampling period in both data sets showed that (1) low R:FR-treated plants overall emitted more volatiles after exposure to volatiles from emitters and subsequent simulated herbivory and (2) volatiles emitted by intact (cVOCs) and herbivory-induced (HIPVs) low R:FR-treated plants enhance the emission of HIPVs in neighboring plants (Fig. 4).

Figure 3. Volatiles emitted by low R:FR-treated plants enhance emissions of HIPVs in low R:FR-treated plants. Time-series analysis of volatile emissions (mean ± SEM, $n = 9$) collected from **(a)** low R:FR-treated or **(b)** high R:FR-treated receiver plants pre-exposed to cVOCs or HIPVs from low or high R:FR-treated plants during 6 h and followed by simulated herbivory (wounding + oral secretions from *S. littoralis*). Pooled data from three independent experiments are shown. The effects of sampling time (T), light treatment of emitters (Light Emitter), volatiles from emitters (Vol Emitter), and their interactions on receivers' volatile emissions were tested using linear mixed-effects models. Statistically significant effects are shown in each graph (*$p < 0.05$, ** $p < 0.001$, *** $p < 0.001$). DMNT and TMTT stand for the homoterpenes (*E*)-4,8-dimethyl-1,3,7-nonatriene and (*E*, *E*)-4,8,12-trimethyltrideca-1,3,7,11-tetraene, respectively. AU refers to arbitrary units.

Volatile treatment

■ cVOCs from high R:FR treated plants
■ HIPVs from high R:FR treated plants
■ cVOCs from low R:FR treated plants
■ HIPVs from low R:FR treated plants

Light Receiver***
Light Emitter**
Volatiles Emitter*

281

Figure 4. Sum (mean ± SEM, *n* = 9) of the emission of all the volatiles signatures (indole, monoterpenes, sesquiterpenes, DMNT, and TMTT) measured over a period of 5 h upon simulated herbivory induction in receiver plants. Low or high R:FR-treated receiver plants were pre-exposed to cVOCs or HIPVs from low or high R:FR-treated plants during 6 h followed by simulated herbivory (wounding + oral secretions from *S. littoralis*).The effects of light treatment of the receiver (Light Receiver), light treatment of the emitter (Light Emitter), the volatiles from the emitters (Volatiles Emitter), and their interactions on receivers' volatile emissions were tested using mixed-effects models followed by the calculation of estimated marginal means (EMMs) and pairwise comparisons using Tukey's Honest Significant Difference (HSD) test. Statistically significant effects are shown in the graph (*p < 0.05, **p < 0.01, ***p < 0.001). Different letters denote significant differences among groups at p < 0.05.

Low R:FR Slightly Increases (Z)-3-HAL Emission but not (Z)-3-HOL, and (Z)-3-HAC

Green leaf volatiles such as (*Z*)-3-HAL, (*Z*)-3-HOL, and (*Z*)-3-HAC, and indole are HIPVs emitted by maize that can primed defenses in neighboring plants (Farag and Pare 2002; Engelberth et al. 2004; Erb et al. 2015). Yet, our previous study has shown that indole emissions in low R:FR treated plants do not differ from high R:FR-treated plants upon simulated herbivory (Escobar□Bravo et al. 2024). To further determine whether FR light affect the emission of GLVs, we compared (*Z*)-3-HAL, (*Z*)-3-HOL, and (*Z*)-3-HAC emissions in simulated herbivory-induced and unwounded plants that were previously exposed to low or high R:FR conditions for 2 days (Fig. 5).

Simulated herbivory significantly induced the emissions of (*Z*)-3-HAL, (*Z*)-3-HOL, and (*Z*)-3-HAC in both low and high R:FR treated plants (Fig. 5). Low R:FR did not significantly affect (*Z*)-3-HOL, and (*Z*)-3-HAC emissions in unwounded and simulated herbivory-induced plants (Fig. 5bc). (*Z*)-3-HAL emission levels were higher in low R:FR-treated plants compared to high R:FR treated plants (Fig. 5a).

Figure 5. Time series analysis of green leaf volatile emissions (mean ± SEM, *n* = 8) determined by PTR-TOF-MS at intervals of 37 min in high and low R:FR-treated B73 maize (*Zea mays*) plants after simulated herbivory. Volatile measurements started at 12:30 pm, 10 min after simulated herbivory treatment (wounding and application of *Spodoptera littoralis* oral secretions; W+OS). The effects of sampling time (Time), light treatment (Light), simulated herbivory (Herbivory) and their interactions on volatile emission were tested using linear mixed-effects models. Plant unit was included as the random intercept as well as a

correlation structure when autocorrelation among residuals was found significant ($p < 0.05$). Statistically
significant effects are indicated in each graph. *** $p < 0.001$.

DISCUSSION

Our study shows that maize plants growing under FR-enriched light conditions (low R:FR ratios) not
only display a higher emission of HIPVs after exposure to volatile cues from neighbors, but they also
emit volatiles that increase the emission of HIPVs in neighboring plants. These findings contribute to
unraveling the complexity of volatile-mediated interplay between plants in vegetation canopies where
shade-associated light cues might play a key role.

Our previous study demonstrated that low R:FR enhances HIPVs emissions in plants
exposed to (Z)-3-HAC alone followed by simulated herbivory (Escobar-Bravo et al., 2024). Here we
show that maize plants exposed to low R:FR also display a stronger priming response upon pre-
exposure to the HIPVs emitted by herbivore-infested plants. We hypothesize that this response might
be partially explained by FR-mediated positive effects on stomata opening and photosynthesis (Zhen
and Bugbee 2020; Escobar⬜Bravo et al. 2024; Huber et al. 2024). As stomata are the entry ports of
VOCs into intact plants (Aratani et al. 2023) and modulate indole and (Z)-3-HAL-mediated priming
responses in maize (Aguirre et al. 2023), changes in stomatal aperture might have affected maize
perception and responses to neighbor's volatiles. In other words, FR-treated plants might take up
more volatiles than high R:FR-treated plants, resulting in a stronger priming stimulus. Alternatively, FR
positive effects on photosynthesis and stomatal opening might have increased the availability of
precursors for volatile biosynthesis (Arimura et al. 2008) and/or the release rate of these volatiles.
Further studies are needed to determine the mechanisms involved in FR-mediated modulation of
plant perception and responses to HIPVs.

Next, we showed that maize plants emit more HIPVs after pre-exposure to constitutive or
herbivory-induced volatiles from low R:FR-treated plants. A possible explanation for this phenomenon
is that a higher emission of volatiles by low R:FR-treated plants might trigger a stronger priming
response in the receiver. For instance, exposure to increasing concentrations of GLVs has been
shown to parallel increases in plasma membrane potential depolarization and cytosolic calcium
concentrations in plant leaves (Zebelo et al. 2012; Aratani et al. 2023). These cellular responses are
associated with the activation of JA defense-related genes (Aratani et al. 2023) and might be part of
the priming stimulus in the receiver plant. Our analysis of GLV emissions in intact plants and plants
subjected to simulated herbivory showed slightly higher levels of hexenal in low R:FR-treated plants
compared to the high R:FR treatment, but no differences in (Z)-3-HAC and (Z)-3-HOL emissions.
Thus, it is unlikely that these differences are responsible for the increase in HIPVs observed in
receiver plants.

Additionally, our previous study revealed that an enriched FR light environment enhances the
emission of monoterpenes and the homoterpene DMMT in herbivory-induced B73 maize plants
(Escobar⬜Bravo et al. 2024). Yet, in the same experiments, we did not observe significant differences
in constitutive volatile emissions in unwounded FR-treated plants compared to high R:FR light
conditions (Escobar⬜Bravo et al. 2024). As both unwounded and herbivory-induced low R:FR treated

plants enhanced the volatile emissions in receiver plants, they might emit a common volatile signal
that was not detected in our PTR-TOF-MS system. This would be the case of volatile compounds that
cannot be ionized by the primary ion (H_3O^+) used in our system. A candidate signal could be the
volatile hormone ethylene. Ethylene emission is induced by low R:FR light conditions in several plant
species, including *Arabidopsis thaliana* and *Nicotiana tabacum*, playing an important role in the
development of shade avoidance traits (Pierik et al. 2003, 2004; Kegge et al. 2013). Furthermore,
ethylene exposure has been shown to increase the emission of HIPVs in maize plants when these
were challenged with simulated herbivory (Schmelz et al. 2003) or (*Z*)-3-HOL exposure (Ruther and
Kleier 2005). Further analysis determining ethylene emissions in low R:FR-treated maize plants
together with the artificial manipulation of ethylene signaling could shed light on the role of this volatile
hormone in the stronger volatile-mediated priming response under enriched FR light conditions.

Finally, our data showed that while 20 h of exposure to HIPVs from plants irradiated with low
R:FR ratios and infested with *S. littoralis* larvae induced a clear priming response in low R:FR treated
plants, shorter exposure (6 h) to HIPVs from simulated herbivory-induced plants did not. We
hypothesize that exposure time and differences in GLVs emission patterns between real and
simulated herbivory might explain these differences and this would need further investigation.

In summary, our findings suggest an enhanced transmission of volatile signals within maize
canopies via FR-mediated effects on the emitter's volatile emissions and the perception of these
volatiles by the receiver. Additional experiments are needed to determine (1) whether this translates
into a larger spatial distribution of warning volatile signals in maize fields, (2) if this information
transfer is species-specific, and (3) what are the benefits and tradeoffs of such a phenomenon. Under
field conditions, for instance, the propagation of volatile signals involving several plants located at
different distances from the emitter might be affected by the diffusion properties and breakdown of
volatile compounds after release (Niinemets et al. 2014; Schuman 2023), as well as by their
absorption and reabsorption by the plant/s closer to the volatile source (Himanen et al. 2010;
Sugimoto et al. 2014, 2023). In addition, maize plants might be exposed to other plant-competition
cues such as limited water and nutrient resources, and root-derived exudates from neighboring plants
(Pierik et al. 2013; Wang et al. 2021) that might affect volatile emissions and perception. For instance,
drought can modulate herbivory-induced (Catola et al. 2018; Lin et al. 2022; Vázquez☐González et al.
2022) and constitutive volatile emissions (Jin et al. 2021), the latter affecting the drought tolerance in
neighboring plants (Jin et al. 2021). Finally, we hypothesize that an enhanced transmission of volatile
priming signals in maize fields might increase HIPVs emission at the plant community level which can
boost the attraction of the natural enemies of herbivores (Aartsma et al. 2017). This might be
especially relevant in dense monocultures, where the limited air movement within a dense canopy can
reduce the dispersion of plant volatile compounds, potentially leading to volatile entrapment. Thus, our
study not only provides a better understanding of the environmental modulation of the transmission of
volatile priming signals in an agronomic relevant species but also sets the basis to further determine
their ecological functions.

ACKNOWLEDGEMENTS

The work of BCJS was supported by the Marie Sklodowska-Curie Action Individual Fellowship (European Union Horizon 2020, Grant Nr. 794947). This work was supported by the Velux Foundation (Grant Nr. 1231), the Swiss National Science Foundation (Grant. Nr. 200355), the State Secretariat for Education, Research, and Innovation SERI (Project CANWAS) and the University of Bern.

COMPETING INTERESTS

The authors have no relevant financial or non-financial interests to disclose

AUTHORS CONTRIBUTION

ME conceived the study. ME, RE-B, BCJS designed experiments. RE-B and BCJS conducted the experiments. RE-B and BCJS performed the PTR-TOF-MS measurements and analyses. RE-B analyzed the data. RE-B, BCJS, and ME interpreted the data. RE-B wrote the manuscript, and all the authors contributed to revisions.

DATA AVAILABILITY

The data generated for this manuscript and the code to reproduce the statistical analysis can be downloaded from GitHub (https://github.com/RocioEscobarBravo/Escobar-Bravo-et-al.-JCE).

REFERENCES

Aartsma Y, Bianchi FJJA, Van Der Werf W, et al (2017) Herbivore-induced plant volatiles and tritrophic interactions across spatial scales. New Phytol 216:1054–1063. https://doi.org/10.1111/nph.14475

Aguirre NM, Grunseich JM, Lima AF, et al (2023) Plant communication across different environmental contexts suggests a role for stomata in volatile perception. Plant Cell Environ 46:2017–2030. https://doi.org/10.1111/pce.14601

Aratani Y, Uemura T, Hagihara T, et al (2023) Green leaf volatile sensory calcium transduction in Arabidopsis. Nat Commun 14:6236

Arimura G, Köpke S, Kunert M, et al (2008) Effects of feeding Spodoptera littoralis on lima bean leaves: IV. Diurnal and nocturnal damage differentially initiate plant volatile emission. Plant Physiol 146:965–973

Catola S, Centritto M, Cascone P, et al (2018) Effects of single or combined water deficit and aphid attack on tomato volatile organic compound (VOC) emission and plant-plant communication. Environ Exp Bot 153:54–62

Christensen SA, Nemchenko A, Borrego E, et al (2013) The maize lipoxygenase, *Zm LOX 10*, mediates green leaf volatile, jasmonate and herbivore-induced plant volatile production for defense against insect attack. Plant J 74:59–73. https://doi.org/10.1111/tpj.12101

Conrath U, Beckers GJM, Langenbach CJG, Jaskiewicz MR (2015) Priming for Enhanced Defense. Annu Rev Phytopathol 53:97–119. https://doi.org/10.1146/annurev-phyto-080614-120132

Engelberth J, Alborn HT, Schmelz EA, Tumlinson JH (2004) Airborne signals prime plants against insect herbivore attack. Proc Natl Acad Sci 101:1781–1785. https://doi.org/10.1073/pnas.0308037100

Erb M, Flors V, Karlen D, et al (2009) Signal signature of aboveground-induced resistance upon belowground herbivory in maize. Plant J 59:292–302. https://doi.org/10.1111/j.1365-313X.2009.03868.x

Erb M, Veyrat N, Robert CA, et al (2015) Indole is an essential herbivore-induced volatile priming signal in
maize. Nat Commun 6:6273

Escobar-Bravo R, Lin P-A, Waterman JM, Erb M (2023) Dynamic environmental interactions shaped by
vegetative plant volatiles. Nat Prod Rep 40:840–865

Escobar-Bravo R, Schimmel BCJ, Zhang Y, et al (2024) Far-red light increases maize volatile emissions in
response to volatile cues from neighbouring plants. Plant Cell Environ 47:3979–3998.
https://doi.org/10.1111/pce.14995

Farag MA, Pare PW (2002) C6-Green leaf volatiles trigger local and systemic VOC emissions in tomato.
Phytochemistry 61:545–554

Fox J, Weisberg S, Adler D, et al (2012) Package 'car.' Vienna R Found Stat Comput 16:333

Himanen SJ, Blande JD, Klemola T, et al (2010) Birch (Betula spp.) leaves adsorb and re-release volatiles specific
to neighbouring plants – a mechanism for associational herbivore resistance? New Phytol 186:722–
732. https://doi.org/10.1111/j.1469-8137.2010.03220.x

Hu L, Ye M, Erb M (2019) Integration of two herbivore-induced plant volatiles results in synergistic effects on
plant defence and resistance. Plant Cell Environ 42:959–971. https://doi.org/10.1111/pce.13443

Hu L, Zhang K, Wu Z, et al (2021) Plant volatiles as regulators of plant defense and herbivore immunity:
molecular mechanisms and unanswered questions. Curr Opin Insect Sci 44:82–88

Huber M, De Boer HJ, Romanowski A, et al (2024) Far-red light enrichment affects gene expression and
architecture as well as growth and photosynthesis in rice. Plant Cell Environ 47:2936–2953.
https://doi.org/10.1111/pce.14909

Jin J, Zhao M, Gao T, et al (2021) Amplification of early drought responses caused by volatile cues emitted from
neighboring tea plants. Hortic Res 8:

Jing T, Qian X, Du W, et al (2021) Herbivore-induced volatiles influence moth preference by increasing the
β-Ocimene emission of neighbouring tea plants. Plant Cell Environ 44:3667–3680.
https://doi.org/10.1111/pce.14174

Kegge W, Ninkovic V, Glinwood R, et al (2015) Red: far-red light conditions affect the emission of volatile
organic compounds from barley (Hordeum vulgare), leading to altered biomass allocation in
neighbouring plants. Ann Bot 115:961–970

Kegge W, Weldegergis BT, Soler R, et al (2013) Canopy light cues affect emission of constitutive and methyl
jasmonate-induced volatile organic compounds in A rabidopsis thaliana . New Phytol 200:861–874.
https://doi.org/10.1111/nph.12407

Lin P, Paudel S, Bin Zainuddin N, et al (2022) Low water availability enhances volatile-mediated direct defences
but disturbs indirect defences against herbivores. J Ecol 110:2759–2771.
https://doi.org/10.1111/1365-2745.13987

Maag D, Dalvit C, Thevenet D, et al (2014) 3-β-D-Glucopyranosyl-6-methoxy-2-benzoxazolinone (MBOA-N-Glc)
is an insect detoxification product of maize 1, 4-benzoxazin-3-ones. Phytochemistry 102:97–105

Maddonni GA, Otegui ME, Andrieu B, et al (2002) Maize leaves turn away from neighbors. Plant Physiol
130:1181–1189

Martinez-Medina A, Flors V, Heil M, et al (2016) Recognizing plant defense priming. Trends Plant Sci 21:818–
822

Niinemets Ü, Fares S, Harley P, Jardine KJ (2014) Bidirectional exchange of biogenic volatiles with vegetation:
emission sources, reactions, breakdown and deposition. Plant Cell Environ 37:1790–1809.
https://doi.org/10.1111/pce.12322

Onosato H, Fujimoto G, Higami T, et al (2022) Sustained defense response via volatile signaling and its
epigenetic transcriptional regulation. Plant Physiol 189:922–933

Pierik R, Ballaré CL (2021) Control of plant growth and defense by photoreceptors: from mechanisms to
opportunities in agriculture. Mol Plant 14:61–76

Pierik R, Mommer L, Voesenek LA (2013) Molecular mechanisms of plant competition: neighbour detection
and response strategies. Funct Ecol 27:841–853. https://doi.org/10.1111/1365-2435.12010

Pierik R, Visser EJW, De Kroon H, Voesenek LACJ (2003) Ethylene is required in tobacco to successfully compete
with proximate neighbours. Plant Cell Environ 26:1229–1234. https://doi.org/10.1046/j.1365-
3040.2003.01045.x

Pierik R, Whitelam GC, Voesenek LACJ, et al (2004) Canopy studies on ethylene-insensitive tobacco identify
ethylene as a novel element in blue light and plant–plant signalling. Plant J 38:310–319.
https://doi.org/10.1111/j.1365-313X.2004.02044.x

Piesik D, Miler N, Lemańczyk G, et al (2015) *Botrytis cinerea* infection in three cultivars of chrysanthemum in
'Alchimist' and its mutants: Volatile induction of pathogen-infected plants. Sci Hortic 193:127–135.
https://doi.org/10.1016/j.scienta.2015.06.040

Ruther J, Kleier S (2005) Plant–Plant Signaling: Ethylene Synergizes Volatile Emission In Zea mays Induced by
Exposure to (Z)-3-Hexen-1-ol. J Chem Ecol 31:2217–2222. https://doi.org/10.1007/s10886-005-6413-8

Schmelz EA, Alborn HT, Tumlinson JH (2003) Synergistic interactions between volicitin, jasmonic acid and
ethylene mediate insect-induced volatile emission in *Zea mays*. Physiol Plant 117:403–412.
https://doi.org/10.1034/j.1399-3054.2003.00054.x

Schuman MC (2023) Where, When, and Why Do Plant Volatiles Mediate Ecological Signaling? The Answer Is
Blowing in the Wind. Annu Rev Plant Biol 74:609–633. https://doi.org/10.1146/annurev-arplant-
040121-114908

Sugimoto K, Matsui K, Iijima Y, et al (2014) Intake and transformation to a glycoside of (Z)-3-hexenol from
infested neighbors reveals a mode of plant odor reception and defense. Proc Natl Acad Sci 111:7144–
7149. https://doi.org/10.1073/pnas.1320660111

Sugimoto K, Ono E, Inaba T, et al (2023) Identification of a tomato UDP-arabinosyltransferase for airborne
volatile reception. Nat Commun 14:677

Turlings TCJ, Erb M (2018) Tritrophic Interactions Mediated by Herbivore-Induced Plant Volatiles: Mechanisms,
Ecological Relevance, and Application Potential. Annu Rev Entomol 63:433–452.
https://doi.org/10.1146/annurev-ento-020117-043507

Vázquez-González C, Pombo-Salinas L, Martín-Cacheda L, et al (2022) Effect of water availability on
volatile-mediated communication between potato plants in response to insect herbivory. Funct Ecol
36:2763–2773. https://doi.org/10.1111/1365-2435.14159

Wang L, Erb M (2022) Volatile uptake, transport, perception, and signaling shape a plant's nose. Essays
Biochem 66:695–702. https://doi.org/10.1042/EBC20210092

Wang L, Jäggi S, Cofer TM, et al (2023) Immature leaves are the dominant volatile-sensing organs of maize. Curr
Biol 33:3679–3689

Wang N, Kong C, Wang P, Meiners SJ (2021) Root exudate signals in plant–plant interactions. Plant Cell Environ
44:1044–1058. https://doi.org/10.1111/pce.13892

Xue J, Gou L, Zhao Y, et al (2016) Effects of light intensity within the canopy on maize lodging. Field Crops Res
188:133–141

Ye M, Glauser G, Lou Y, et al (2019) Molecular dissection of early defense signaling underlying volatile-mediated
defense regulation and herbivore resistance in rice. Plant Cell 31:687–698

Zebelo SA, Matsui K, Ozawa R, Maffei ME (2012) Plasma membrane potential depolarization and cytosolic
calcium flux are early events involved in tomato (Solanum lycopersicon) plant-to-plant
communication. Plant Sci 196:93–100

Zhen S, Bugbee B (2020) Far-red photons have equivalent efficiency to traditional photosynthetic photons:
Implications for redefining photosynthetically active radiation. Plant Cell Environ 43:1259–1272.
https://doi.org/10.1111/pce.13730

SUPPLEMENTARY MATERIAL

Figure S1. Light spectral composition determined in far-red supplementation experiments.

Figure S2. Detailed schematic overview of the experimental set-up in Fig. 1

Figure S3. Detailed schematic overview of the experimental set-up in Fig. 2.

Figure S4. Detailed schematic overview of the experimental set-up in Fig. 3 and Fig. 4.

Figure S5. Cumulative emission of individual volatiles in Fig. 1.

Figure S6. Cumulative emission of individual volatiles in Fig. 2.