Neural dynamics for working memory and evidence integration during olfactory navigation in *Drosophila*

Nicholas D. Kathman, Aaron J. Lanz, Jacob D. Freed, Katherine I. Nagel

Department of Neuroscience, NYU School of Medicine

Abstract

Working memory and evidence integration are fundamental components of cognition thought to arise from distributed circuits throughout the brain[1-2]. Theoretical[3,4] and behavioral studies[5,6] argue that both processes are required for plume navigation, an innate task in which animals use stochastic sensory cues to navigate towards the unknown location of an odor source[7-10]. Here we identify a small population of local neurons in the navigation center of *Drosophila*[11-13] that exhibits both evidence integration and working memory dynamics during goal-directed olfactory navigation. Developing a closed-loop virtual plume navigation paradigm, we show that a bump of activity in this population ramps up with successive odor encounters, and can persist for variable intervals after odor loss. While bump activity persists, the fly maintains the goal heading it adopted during odor. Silencing these neurons impairs the persistence of upwind heading after odor loss. Simulations show that the time constant of persistence observed in these neurons optimizes navigation in a turbulent boundary layer plume. Our work localizes working memory and evidence integration to a specific group of genetically-identified neurons, which will facilitate the mechanistic dissection of these building blocks of cognition.

Introduction

In many real-world navigation tasks, sensory information about the direction and location of a goal is only intermittently available. For example, an owl hunting a mouse by sound might get only sporadic and uncertain information about the location of its prey, or a leopard chasing an antelope by sight might have its view interrupted by brush. In such tasks, the ability to accumulate sensory information about the direction of a target over time, and to hold that information in working memory during loss of sensory input are critical to success. While the neural mechanisms underlying the processes of evidence integration and working memory have been studied extensively in non-human primates and rodents[1,2], the distributed nature of these circuits has made it challenging to causally link neural activity patterns to behavior[14,15].

Many animals use odor cues to navigate towards the unknown location of a food source[16,17]. Because odor plumes are turbulent, they generate stochastic cues that must be integrated and stored in memory in order to navigate effectively[17]. Behavioral studies in plumes argue that insects integrate odor encounters over time to drive navigational decisions[5,6], while theoretical studies argue that an optimal searching agent should possess a working memory that allows it to continue moving in the goal direction during short blanks in the odor stimulus[3,4]. However, the potential neural substrate of these computations is unknown.

Recent evidence suggests that the fan-shaped body (FB), a part of the insect navigation center called the Central Complex (CX), is a key location for computing a goal direction during navigation. Output neurons of the FB can compare the fly's current heading to a goal signal expressed in allocentric (world-centered) coordinates to steer the fly in a target direction[11,13,18]. Patterned activation of certain FB local

neurons can drive goal-directed walking relative to a visual landmark[13], or to the wind[19], consistent with the idea that FB local neurons can encode allocentric goals.

The FB is ideally positioned to play a role in odor-guided navigation. It receives odor information through a set of wide-field tangential neurons that are downstream of the associative learning center in the mushroom body[19-21]. Activation of some of these tangential inputs, and also of some FB local neurons, can drive orientation upwind[19,22], while broad developmental disruption of the CX completely abolishes odor-gated wind orientation behavior[23]. However, the *in vivo* dynamics of FB goal signals and their precise role in natural navigation tasks are not well understood.

In a recent study, we identified FB local neurons labeled by the line VT062617-Gal4 whose sparse activation can drive walking in a goal direction relative to the wind[19]. Here we developed a closed-loop virtual paradigm to examine the activity of these neurons during odor-guided navigation. We find that these neurons exhibit a bump of activity that is activated by odor but persists after odor offset for variable intervals. While the bump remains active, we find that flies continue running in the direction adopted during odor, arguing that these neurons represent a short-term directional goal memory. By simulating a turbulent plume in closed-loop, we show that these neurons integrate olfactory evidence over a period of several seconds, a property not shared by a second parallel group of local neurons. Optogenetic silencing of these neurons impairs persistent directional running after odor loss. Simulating these dynamics in a model shows that a short-term memory for goal direction with a time constant similar to the one we measure in these neurons improves navigation in a turbulent boundary layer plume. Our findings reveal internal dynamics that shape the transformation of stochastic sensory input into goal-directed navigation behavior.

Results

A persistent bump of activity correlates with maintaining a goal direction after odor loss

To investigate the activity of VT062617 neurons during on-going navigation, we developed a closed-loop virtual odor navigation paradigm. In this paradigm (Fig. 1a), the fly controls the direction of the wind in closed-loop, while a pulse of attractive odor (apple cider vinegar, 0.5% dilution in water) is delivered in open loop (see Methods). Similar to previous observations in both freely-walking[5,6] and tethered flies[24,25], we observed that flies walked more strongly upwind during odor compared to the pre-wind baseline period (Fig, 1b, Ext. Data Fig.1a). Odor-evoked trajectories were straighter than baseline walking (Ext. Data Fig. 1a), and biased towards, but not uniformly, upwind (Ext Data Fig. 1b). In some trials, flies showed a looping search-like behavior at odor offset, similar to search behavior evoked by odor loss in freely-walking flies[5,24,26] (Fig. 1b, left-most trace). On many trials, however, flies maintained their trajectory for a variable number of seconds following the loss of odor (Fig 1b, right-most trace, Ext Data Fig. 1b,c). Straight trajectories after odor lasted significantly longer than those observed during the baseline period (Ext Data Fig. 1b,c).

Imaging from FB local neurons labeled by VT062617 (Ext Data Fig. 1d), revealed a localized bump of activity with dynamics related to both sensory input and ongoing navigation behavior (Fig. 1c). Bump activity was most strongly and consistently activated by odor (Fig. 1d,e, Ext. Data Fig.1e), weakly activated by wind onset in some trials (Fig. 1d), and was absent in interleaved trials without airflow (Ext. Data Fig. 1f). Localized to a few columns (Ext. Data Fig.1g), bump activity persisted for variable durations after measured odor offset (Fig. 1c,f,g, Ext. Data Fig. 1h,i). Unlike activity in the compass system[27], we observed little movement of the bump during turns, and poor correlation with heading (Ext. Data Fig.1j,k). We observed no systematic change in bump position when the wind was rotated during odor, leading to a compensatory change in heading (Ext. Data Fig.1l,m). We conclude that activity in these neurons does not track heading,

but is more consistent with representing a goal direction[13,28.]

In several example trajectories, we observed that persistent bump activity in VT062617 neurons correlated with maintained straight movement in the direction adopted during odor, ending when the fly turned (Fig. 1c). To test whether this observation held across trials and flies, we computed a straightness index (see Methods), and examined the mean straightness for each fly in the presence and absence of a bump. During the odor period, flies tended to walk straight regardless of whether a bump was present. However, during the post-odor period, the presence of a bump was associated with greater straightness (Fig. 1h). To examine the relationship between bump activity and navigational trajectories in greater detail, we detected bump offsets during the post-odor period and compared trajectories after bump offset to equivalent trajectories when the bump stayed on (see Methods, Fig. 1i,j). We observed that when the bump remained on, trajectories remained close to the "goal trajectory" adopted during the odor period. In contrast, when the bump turned off, trajectories deviated from this goal direction. We conclude that the presence of an activity bump in VT062617 correlates with maintaining a goal direction following the loss of odor.

Integration of olfactory evidence during turbulent plume navigation

In natural environments, odors form turbulent plumes in which odor fluctuates stochastically as a function of distance and direction to the source[29-32]. Therefore, we next sought to examine the activity of VT062617 neurons as flies navigated a turbulent plume. To implement a virtual plume environment, we leveraged a previously recorded movie of odor concentrations in a boundary layer plume at approximately fly height and windspeed comparable to our experiment[32] (Methods, Fig. 2a). As the fly navigated in 2D, we used time and the virtual position of the fly to index into this movie and determine the odor concentration at the virtual location of the fly. We then used an adaptive compression algorithm[5] that mimics aspects of peripheral olfactory encoding[33] to binarize this concentration relative to an adaptive threshold, and delivered the resulting odor pulses to the fly. As a result of this algorithm, a fly navigating our virtual odor plume encounters a stochastic series of odor pulses that increase in frequency as the fly approaches the virtual source.

Plume navigation revealed additional dynamics in VT062617 neurons. As the fly navigated the plume, we observed initially weak responses to odor that grew with repeated odor encounters (Fig. 2b). We also observed instances where a series of odor encounters ignited a long-lasting bump of activity, that was again associated with maintaining a straight trajectory (Fig. 2c). To ask whether bump activity correlates with a maintained trajectory in this more complex environment, we computed angular speed and deviation from goal in periods following odor offset (Fig. 2d). Similar to our observations with isolated odor pulses, we found that persistent bump activity was associated with lower goal deviation, and that bump offset was associated with significant increases in both angular speed and deviation from the goal direction adopted during odor. To quantify how bump dynamics depend on odor in this more variable sensory environment, we computed instantaneous odor encounter frequency (see Methods), then plotted the probability of observing a bump as a function of odor encounter frequency (Fig. 2e). We observed that bump probability grew with odor encounter frequency. These observation suggest that VT062617 activity does not simply follow the odor stimulus, but reflects an integration of odor information over time.

During our closed-loop navigation tasks, odor, locomotor parameters, and navigation behavior are all correlated with one another, making it challenging to disentangle the contribution of each parameter to the signal observed in VT062617 neurons. To address this issue, we took advantage of the stochastic nature of odor encounters in our plume paradigm to compute decorrelated linear filters between bump amplitude and

odor, forward velocity, and angular speed. Adopting techniques from the sensory coding literature[34], we computed the correlation matrix between these variables, and applied a regularized pseudo-inverse of this matrix to the cross-correlogram of each parameter and bump amplitude (see Methods) to compute decorrelated filters relating odor and behavior to neural activity. This analysis reveals distinct encoding dynamics for both sensory input and ongoing behavior. The filter describing the relationship between odor and neural activity (Fig. 2f, Ext. Data Fig. 2a) had an "integrating" shape with a single peak that extends ~10 seconds into the past, supporting the idea that these neurons integrate odor information over several seconds. In contrast, the filter relating angular speed (turning) to activity was entirely negative (Fig. 2f, Ext. Data Fig. 2b), consistent with the observation that activity is strongest during straight walking. We also observed a positive integrating filter for forward velocity (Fig. 2f, Ext. Data Fig. 2c), although this filter was narrower than the filter for odor. This forward velocity filter supports observations during odor pulse trials, where bump onset was associated with an increase in forward velocity, and forward velocity was higher when a bump was present (Ext. Data Fig. 2d,e). Our plume experiments thus reveal a temporal integration of odor input, and support our previous finding that persistent activity in VT062617 neurons is associated with faster, straighter trajectories.

Persistent activity and odor integration are not widespread in the FB

The dynamics we observe in VT062617 neurons might arise within these neurons or might be more broadly shared across the fan-shaped body. To address the specificity of these dynamics, we imaged from a second line, 52G12-Gal4, that labels a large set of ventral FB local neurons (Ext. Data Fig. 3a,b). Imaging from 52G12 neurons in the plume paradigm revealed transient responses to repeated odor encounters without the gradual ramping and persistence we observed in VT062617 neurons (Fig. 3a). The filter relating odor to activity for 52G12 was much narrower than the filter for VT062617, indicating a shorter integration time (Fig. 3b). The filter relating angular speed to activity had a positive peak for 52G12, in contrast to the negative peak for VT062617, indicating a correlation with turning rather than straight walking (Fig. 3c). Consistent with these observations, we found that 52G12 responses to single odor pulses were transient rather than persistent (Ext. Data Fig. 3c,d), and correlated with turning rather than straight walking (Ext. Data Fig. 3e). Thus, the coding features of VT062617 neurons— gradual integration of olfactory evidence and persistent activity associated with straight trajectories— are not shared across the FB, but are specific to a subset of local neurons.

While neurons labeled by 52G12 and VT062617 showed distinct local encoding dynamics, both showed similar modulation on the slower timescale of trials (Fig. 3d). Many flies walked upwind only on a subset of trials in an experiment. Runs of trials with high upwind displacement could occur both early and late in the experiment for both genotypes, arguing that lack of engagement does not reflect simple sensory habituation. We observed greater activity in both VT062617 and 52G12 neurons during periods when the fly showed strong upwind displacement. Upwind displacement and total bump activity were positively correlated across both genotypes (Fig. 3e), significantly greater than in shuffled trials. These data indicate that while different FB local neurons track distinct local features of odor dynamics and self-motion, activity across the FB may be co-regulated by behavioral state on slower timescales.

VT062617 neurons play a causal role in persistent goal-directed navigation

Our plume experiments argue that VT062617 neurons integrate odor information over time and maintain this information for seconds after odor offset. We therefore wondered if these neurons contribute causally to persistent goal-directed navigation behavior. We first asked whether freely-walking flies exhibit

similar persistent trajectories to those we observed in our virtual paradigm. We presented freely walking flies in a laminar wind tunnel with variable numbers of odor pulses at 1Hz, and examined the dynamics of their orientation behavior (Fig. 4a-c). We found that increasing numbers of odor pulses led to a greater probability of upwind walking, that saturated around 4 pulses (Fig. 4b,c). Upwind walking persisted for 5-10 seconds after the end of the last odor pulse, and decayed with a uniform time course across pulse numbers (Fig. 4b). These results indicate that persistent upwind walking that outlasts the odor stimulus also occurs in freely-walking flies.

To ask whether VT062617 neurons play a role in these dynamics, we presented flies expressing the optogenetic silencing agent UAS-GtACR1[35] in VT062617 neurons with variable numbers of odor pulses. We measured upwind walking in interleaved trials with blue light that silences the neurons, and in the absence of light. We found that silencing VT062617 neurons led to reduced persistence in upwind walking across pulse numbers (Fig. 4d). The probability of walking upwind in the 10s period after odor OFF was significantly reduced for all stimulus configurations (Fig. 4f). In genetic control flies, light had no effect on the persistence of upwind walking (Fig. 4e,f). To ask if this effect is specific to fan-shaped body neurons labeled in this line, we repeated our experiment using a split-Gal4 line specific for hΔK neurons[36], the dominant FB cell type in VT062617[23,] to drive GtACR1. We also observed a decrease in persistent upwind running using this split-Gal4 line, although a higher light level was required to obtain a behavioral effect and we only observed a significant reduction with the longest pulse train (Fig. 4g, Ext Data Fig. 4a,b). We conclude that FB local neurons labeled by VT062617 neurons contribute causally to maintaining a trajectory after odor loss.

A persistent goal representation improves navigation in turbulent environments

Our data argue that VT062617 neurons are not required for a fly to select an initial upwind trajectory in response to odor, but that their activity allows the fly to maintain its trajectory for some time after odor loss. Intriguingly, this type of working memory was recently predicted by two theoretical studies of plume navigation using purely computational methods[3,4]. Both studies found that an optimal agent should integrate odor information over a window related to the average duration of blanks in an odor plume, allowing the agent to determine whether a gap in odor is expected (meaning the agent is within the plume envelope) or unexpected (because the agent has lost the plume). To test whether the dynamics we observed in VT062617 neurons could perform such a function, we integrated previous models of fly olfactory navigation[5,26] with the finite state controller framework of Verano et al.[3] In this model, the fly can adopt one of three navigational states: baseline walking, goal-directed upwind walking, or search. Critically, the transition from goal-directed behavior to search is probabilistic with a time constant τ_p, allowing the system to remain in the goal-directed state for variable intervals after odor offset (Fig. 5a,b, Methods). We fit coefficients of the model to reproduce the distribution of forward and angular velocities measured in our walking ball paradigm (Fig. 5c)

We ran this model in the same previously recorded odor plume we used for our plume experiments and varied the time constant τ_p of memory for the goal state (Fig. 5d,e). When this time constant was small (Fig 5d, bottom row), we found that flies began searching during brief pauses in odor within the plume. Consequently, they often remained trapped near the base of the plume and did not make strong progress upwind. As the time constant increased (Fig 5d, middle row), flies made more effective upwind progress, and search behavior became concentrated at the edges of the plume envelope. At longer time constants, flies ran past the source, and search success decreased (Fig 5d, top row). Thus, consistent with previous studies, we find that an optimal memory time constant allows flies to integrate stochastic odor encounters within the plume, and correctly localize search behavior at the edges of the plume envelope. Strikingly, the optimal memory time constant for our model was very similar to time constant of persistence we observed in

VT062617 neurons (Fig. 5e, red arrow). We conclude that the dynamics of persistent activity observed in these neurons provide an optimal integration of olfactory information for a fly navigating a turbulent plume.

Discussion

A persistent goal representation supports navigation with sparse sensory input

During natural navigation tasks, sensory cues indicating the direction or location of a target can be sparse. Turbulent plumes epitomize this type of task, as odor encounters in a plume are intermittent, with statistics that depend in complex ways on medium, windspeed, lateral and longitudinal distance from the source, and distance from the ground, among other factors[29-32]. Ideally, an agent trying to find the source of a plume should maintain its goal heading if it is within the plume envelope, and initiate search or recovery behaviors only when this envelope has been lost[3,4,17]. However, instantaneous measurements of odor concentration are insufficient to discriminate between expected gaps due to turbulence and gaps due to plume loss[37].

One solution for an animal navigating a stochastic environment is to integrate sensory information over time. By integrating odor input as it moves, a fly could generate an internal estimate of whether it is in the plume or not, and use this to more effectively determine whether it should continue pursuing its goal, or initial search behaviors[5,6,17]. However, to be effective, such a signal should be tuned to the expected duration of gaps within a plume.

Here we identify a neural goal signal that exhibits integration and persistence tuned to the properties of odor plumes that a walking fly is likely to encounter. In a previous study, we showed that sparse activation of this population of neurons can drive movement in a reproducible direction relative to the wind, suggesting that it can encode a navigational goal[19]. Here we show that this population exhibits a localized bump of activity that turns on in response to odor, and remains on while the fly maintains a goal heading relative to wind. Rotation of the wind does not cause a rotation of the bump, consistent with the idea that it represents an allocentric goal direction, rather than a heading estimate[13,28]. Silencing of these neurons curtailed persistent directional running after odor offset, arguing that they play a causal role in maintaining a goal direction. Interestingly, integration and persistence in these neurons appears to be probabilistic, rather than fixed. Simulations with a similar probabilistic persistence time showed effective integration in a boundary layer plume, arguing that the dynamics we observed *in vivo* are well-suited to navigating the types of environment a walking fly is likely to encounter. In our imaging data, different flies showed different mean persistence durations (Ext Data Fig. 1i), suggesting that this integration time might be plastic. Future experiments might explore whether the duration of persistence in these neurons can change depending on the statistics of the odor environment or the fly's mode of locomotion.

Recent studies of the FB suggest that it translates variables into allocentric coordinates[13,18,38,39], and can represent goal directions for navigation in this world-centric system[13,19,28]. An allocentric representation of estimated source direction could allow the fly to weigh and accumulate evidence for or against a particular direction in world coordinates, rather than having to update these representations with each turn. Behavioral experiments argue that insects integrate many different cues to navigate towards an odor source, including the temporal pattern of odor encounters[40], differences in odor concentration across the antennae[41,42], the motion of the odor itself[43], and the direction of the wind[44]. A centralized goal signal could allow these diverse cues to be integrated into a single estimate of goal direction. The direction signal we identified could also be inverted, perhaps through the system of hΔ neurons that translates signals by 180° across the array of the

FB, in order to allow a fly to relocate the edge of an odor after leaving it[45,46]. Future experiments could investigate the integration of convergent and divergent cues to source location at the level of the signal we describe.

Working memory and evidence integration in a genetically tractable central brain circuit

The persistent goal signal we characterize here shares properties with working memory signals described in vertebrate systems[2]. Classical experiments identified persistent stimulus-tuned activity in the prefrontal cortex of monkeys as a potential neural correlate of working memory[47-48], while ramping activity in several cortical and subcortical regions has been linked to integration of noisy sensory evidence[1,14,49]. While optogenetic perturbation experiments have provided indirect evidence for recurrent attractor networks[50,51], direct evidence for these models has been challenging to obtain, given the distributed nature of vertebrate working memory circuitry[14,15,50]. Dominant models of working memory rely on attractors generated by recurrent excitatory connectivity[2,15,52,53]. The major local neuron type labeled by VT062617 is hΔK[23], which shows strong recurrent connections with another CX cell type known as PFG[12]. In contrast, 52G12, which shows much more rapid and transient dynamics, appears to label a large number of FC1 types (Ext. Data Fig. 3a), which represent one of the most direct feedforward pathways from input to output within the FB[12]. Future recording and perturbation experiments using more specific genetic labels should allow us to directly test whether the distinct dynamics of these two populations reflect their distinct circuit connectivity, or some other form of intrinsic or synaptic dynamics.

Understanding how different species represent goals for navigation remains an area of active research[54-56]. *Drosophila* larvae, which navigate in much more stable odor gradients[57], lack a fully developed CX[58], and may require only egocentric computations to navigate. While allocentric representations of goal direction are now established in adult flies[13], it remains unclear if the FB also encodes goal locations. A recent theoretical model has suggested that insects could use a pair of FB vectors, one pointed towards home, and one estimating a goal direction, to triangulate the location of a goal[59]. In contrast to flies, rodents foraging for odorous food rapidly switch from using odor cues to remembered spatial positions[60]. In rodents performing odor-guided localization tasks, mixed odor and spatial representations develop at many places along the pathway from odor input to hippocampus[61,62]. How the distinct navigation systems of vertebrates and invertebrates might enable different behaviors remains an open question[63]. Using ethological tasks to tease apart the nature and causal role of goal representations can provide insight into the structure and organization of navigation circuits across species.

References

1. Brody CD, Hanks TD. 2016. Neural underpinnings of the evidence accumulator. *Curr Opin Neurobiol* **37**: 149–157. doi: 10.1016/j.conb.2016.01.003
2. Wang X-J. 2021. 50 years of mnemonic persistent activity: quo vadis? *Trends Neurosci* **44**(11): 888–902. doi: 10.1016/j.tins.2021.09.001
3. Verano KVB, Panizon E, Celani A. 2023. Olfactory search with finite-state controllers. *Proc Natl Acad Sci* **120**(34): e2304230120. doi: 10.1073/pnas.2304230120
4. Rando M, James, M, Verri A, Rosasco L, Seminara A. 2024. Q-learning to navigate turbulence without a map. *arXiv*:2404.17495 [physics.bio-ph]
5. Álvarez-Salvado E, Licata AM, Connor EG, McHugh MK, King BM, Stavropoulos N, Victor JD, Crimaldi JP, Nagel KI. 2018. Elementary sensory-motor transformations underlying olfactory navigation in walking fruit-flies. *Elife* **7**: e37815. doi: 10.7554/elife.37815

6. Demir M, Kadakia N, Anderson HD, Clark DA, Emonet T. 2020. Walking Drosophila navigate complex plumes using stochastic decisions biased by the timing of odor encounters. *eLife* **9**: e57524. doi: 10.7554/elife.57524

7. David CT, Kennedy JS, Ludlow AR. 1983. Finding of a sex pheromone source by gypsy moths released in the field. *Nature* **303**(5920): 804–806. doi: 10.1038/303804a0

8. Dekker T, Cardé RT. 2011. Moment-to-moment flight manoeuvres of the female yellow fever mosquito (Aedes aegypti L.) in response to plumes of carbon dioxide and human skin odour. *J Exp Biol* **214**(20): 3480–3494. doi: 10.1242/jeb.055186

9. Riffell JA, Shlizerman E, Sanders E, Abrell L, Medina B, Hinterwirth AJ, Kutz JN. 2014. Flower discrimination by pollinators in a dynamic chemical environment. *Science* **344**(6191): 1515–1518. doi: 10.1126/science.1251041

10. van Breugel F, Dickinson MH. 2014. Plume-Tracking Behavior of Flying Drosophila Emerges from a Set of Distinct Sensory-Motor Reflexes. *Curr Biol* **24**(3): 274–286. doi: 10.1016/j.cub.2013.12.023

11. Stone T, Webb B, Adden A, Weddig NB, Honkanen A, Templin R, Wcislo W, Scimeca L, Warrant E, Heinze S. 2017. An Anatomically Constrained Model for Path Integration in the Bee Brain. *Curr Biol* **27**(20): 3069-3085.e11. doi: 10.1016/j.cub.2017.08.052

12. Hulse BK, Haberkern H, Franconville R, Turner-Evans D, Takemura S, Wolff T, Noorman M, Dreher M, Dan C, Parekh R, et al. 2021. A connectome of the Drosophila central complex reveals network motifs suitable for flexible navigation and context dependent action selection. *eLife* **10**: e66039. doi: 10.7554/elife.66039

13. Mussells-Pires P, Zhang L, Parache V, Abbott LF, Maimon G. 2024. Converting an allocentric goal into an egocentric steering signal. *Nature* **626**(8000): 808–818. doi: 10.1038/s41586-023-07006-3

14. Gupta D, Kopec CD, Bondy AG, Luo TZ, Elliott VA, Brody CD. 2024. A multi-region recurrent circuit for evidence accumulation in rats. *bioRxiv*: 2024.07.08.602544. doi: 10.1101/2024.07.08.602544

15. Khona M, Fiete IR. 2022. Attractor and integrator networks in the brain. *Nat Rev Neurosci* **23**(12): 744–766. doi: 10.1038/s41583-022-00642-0

16. Baker KL, Dickinson M, Findley TM, Gire DH, Louis M, Suver MP, Verhagen JV, Nagel KI, Smear MC. 2018. Algorithms for Olfactory Search across Species. *J Neurosci* **38**(44): 9383–9389. doi: 10.1523/jneurosci.1668-18.2018

17. Reddy, G., Murthy, V.N. and Vergassola, M., 2022. Olfactory sensing and navigation in turbulent environments. *Annual Review of Condensed Matter Physics*, 13(1), pp.191-213.

18. Westeinde EA, Kellogg E, Dawson PM, Lu J, Hamburg L, Midler B, Druckmann S, Wilson RI. 2024. Transforming a head direction signal into a goal-oriented steering command. *Nature* **626**(8000): 819–826. doi: 10.1038/s41586-024-07039-2

19. Matheson AMM, Lanz AJ, Medina AM, Licata AM, Currier TA, Syed MH, Nagel KI. 2022. A neural circuit for wind-guided olfactory navigation. *Nat Commun* **13**(1): 4613. doi: 10.1038/s41467-022-32247-7

20. Li F, Lindsey JW, Marin EC, Otto N, Dreher M, Dempsey G, Stark I, Bates AS, Pleijzier MW, Schlegel P, et al. 2020. The connectome of the adult Drosophila mushroom body provides insights into function. *eLife* **9**: e62576. doi: 10.7554/elife.62576

21. Scaplen KM, Talay M, Fisher JD, Cohn R, Sorkaç A, Aso Y, Barnea G, Kaun KR. 2021. Transsynaptic mapping of Drosophila mushroom body output neurons. *eLife* **10**: e63379. doi: 10.7554/elife.63379

22. Aso Y, Yamada D, Bushey D, Hibbard KL, Sammons M, Otsuna H, Shuai Y, Hige T. 2023. Neural circuit mechanisms for transforming learned olfactory valences into wind oriented movement. *eLife* **12**: e85756. doi: 10.7554/elife.85756

23. Hamid A, Gattuso H, Caglar AN, Pillai M, Steele T, Gonzalez A, Nagel K, Syed MH. 2024. The conserved RNA-binding protein Imp is required for the specification and function of olfactory navigation circuitry in Drosophila. *Curr Biol* **34**(3): 473-488.e6. doi: 10.1016/j.cub.2023.12.020

24. Handler A, Graham TGW, Cohn R, Morantte I, Siliciano AF, Zeng J, Li Y, Ruta V. 2019. Distinct Dopamine Receptor Pathways Underlie the Temporal Sensitivity of Associative Learning. *Cell* **178**(1): 60-75.e19. doi: 10.1016/j.cell.2019.05.040

25. Zolin A, Cohn R, Pang R, Siliciano AF, Fairhall AL, Ruta V. 2021. Context-dependent representations of movement in Drosophila dopaminergic reinforcement pathways. *Nat Neurosci* **24**(11): 1555–1566. doi: 10.1038/s41593-021-00929-y

26. Gattuso H, Nuñez K, Rea B de la, Ermentrout B, Victor J, Nagel K. 2024. Inhibitory control of locomotor statistics in walking Drosophila. *bioRxiv*: 2024.04.15.589655. doi: 10.1101/2024.04.15.589655

27. Seelig JD, Jayaraman V. 2015. Neural dynamics for landmark orientation and angular path integration. *Nature* **521**(7551): 186–191. doi: 10.1038/nature14446

28. Beetz, M.J., Kraus, C. and El Jundi, B., 2023. Neural representation of goal direction in the monarch butterfly brain. *Nature communications*, *14*(1), p.5859.

29. Celani A, Villermaux E, Vergassola M. 2014. Odor landscapes in turbulent environments. *Physical Review X*. **4**(4):041015.

30. Ackels T, Erskine A, Dasgupta D, Marin AC, Warner TPA, Tootoonian S, Fukunaga I, Harris JJ, Schaefer AT. 2021. Fast odour dynamics are encoded in the olfactory system and guide behaviour. *Nature* **593**(7860): 558–563. doi: 10.1038/s41586-021-03514-2

31. Nag A, van Breugel F. 2024. Odour source distance is predictable from a time history of odour statistics for large scale outdoor plumes. *Journal of the Royal Society Interface* **21**(216):20240169.

32. Connor EG, McHugh MK, Crimaldi JP. 2018. Quantification of airborne odor plumes using planar laser-induced fluorescence. *Exp Fluids* **59**(9): 137. doi: 10.1007/s00348 018-2591-3

33. Victor JD, Boie SD, Connor EG, Crimaldi JP, Ermentrout GB, Nagel KI. 2019. Olfactory Navigation and the Receptor Nonlinearity. *J Neurosci* **39**(19): 3713–3727. doi: 10.1523/jneurosci.2512-18.2019

34. Nagel KI, Doupe AJ. 2008. Organizing Principles of Spectro-Temporal Encoding in the Avian Primary Auditory Area Field L. *Neuron* **58**(6): 938–955. doi: 10.1016/j.neuron.2008.04.028

35. Mohammad F, Stewart JC, Ott S, Chlebikova K, Chua JY, Koh TW, Ho J, Claridge-Chang A. 2017. Optogenetic inhibition of behavior with anion channelrhodopsins. *Nature methods*. **14**(3):271-4.

36. Wolff, T., Eddison, M., Chen, N., Nern, A., Sundaramurthi, P., Sitaraman, D. and Rubin, G.M., 2025. Cell type-specific driver lines targeting the Drosophila central complex and their use to investigate neuropeptide expression and sleep regulation. *ELife*, *14*, p.RP104764.

37. Murlis, J., Elkinton, J.S. and Carde, R.T., 1992. Odor plumes and how insects use them. *Annual review of entomology*, *37*(1), pp.505-532.

38. Lu J, Behbahani AH, Hamburg L, Westeinde EA, Dawson PM, Lyu C, Maimon G, Dickinson MH, Druckmann S, Wilson RI. 2022. Transforming representations of movement from body- to world-centric space. *Nature* **601**(7891): 98–104. doi: 10.1038/s41586-021-04191-x

39. Lyu C, Abbott LF, Maimon G. 2022. Building an allocentric travelling direction signal via vector computation. *Nature* **601**(7891): 92–97. doi: 10.1038/s41586-021-04067-0

40. Mafra-Neto A, Cardé RT. 1994. Fine-scale structure of pheromone plumes modulates upwind orientation of flying moths. *Nature* **369**(6476): 142–144. doi: 10.1038/369142a0

41. Borst A, Heisenberg M. 1982. Osmotropotaxis in Drosophila melanogaster. *J Comp Physiol* **147**(4): 479–484. doi: 10.1007/bf00612013

42. Gaudry, Q., Hong, E.J., Kain, J., de Bivort, B.L. and Wilson, R.I., 2013. Asymmetric neurotransmitter release enables rapid odour lateralization in Drosophila. *Nature*, *493*(7432), pp.424-428.

43. Kadakia N, Demir M, Michaelis BT, DeAngelis BD, Reidenbach MA, Clark DA, Emonet T. 2022. Odour motion sensing enhances navigation of complex plumes. *Nature* **611**(7937): 754–761. doi: 10.1038/s41586-022-05423-4

44. van Breugel F, Jewell R, Houle J. 2022. Active anemosensing hypothesis: how flying insects could estimate ambient wind direction through sensory integration and active movement. *J R Soc Interface* **19**(193): 20220258. doi: 10.1098/rsif.2022.0258

45. Tao, L., Ozarkar, S. and Bhandawat, V., 2020. Mechanisms underlying attraction to odors in walking Drosophila. *PLoS computational biology*, *16*(3), p.e1007718.

46. Siliciano, A.F., Minni, S., Morton, C., Dowell, C.K., Eghbali, N.B., Rhee, J.Y., Abbott, L.F. and Ruta, V., 2025. A vector-based strategy for olfactory navigation in Drosophila. *bioRxiv*.

47. Fuster JM. 1973. Unit activity in prefrontal cortex during delayed-response performance: neuronal correlates of transient memory. *J Neurophysiol* **36**(1): 61–78. doi: 10.1152/jn.1973.36.1.61

48. Funahashi S, Bruce CJ, Goldman-Rakic PS. 1989. Mnemonic coding of visual space in the monkey's dorsolateral prefrontal cortex. *J Neurophysiol* **61**(2): 331–349. doi: 10.1152/jn.1989.61.2.331

49. Shadlen MN, Newsome WT. 2001. Neural Basis of a Perceptual Decision in the Parietal Cortex (Area LIP) of the Rhesus Monkey. *J Neurophysiol* **86**(4): 1916–1936. doi: 10.1152/jn.2001.86.4.1916

50. Kopec CD, Erlich JC, Brunton BW, Deisseroth K, Brody CD. 2015. Cortical and Subcortical Contributions to Short-Term Memory for Orienting Movements. *Neuron* **88**(2): 367–377. doi: 10.1016/j.neuron.2015.08.033

51. Inagaki HK, Fontolan L, Romani S, Svoboda K. 2019. Discrete attractor dynamics underlies persistent activity in the frontal cortex. *Nature* **566**(7743): 212–217. doi: 10.1038/s41586-019-0919-7

52. Goldman-Rakic PS. 1995. Cellular basis of working memory. *Neuron* **14**(3): 477–485. doi: 10.1016/0896-6273(95)90304-6

53. Compte, A., Brunel, N., Goldman-Rakic, P.S. and Wang, X.J., 2000. Synaptic mechanisms and network dynamics underlying spatial working memory in a cortical network model. *Cerebral cortex*, *10*(9), pp.910-923.

54. Sarel A, Finkelstein A, Las L, Ulanovsky N. 2017. Vectorial representation of spatial goals in the hippocampus of bats. *Science* **355**(6321): 176–180. doi: 10.1126/science.aak9589

55. Basu R, Gebauer R, Herfurth T, Kolb S, Golipour Z, Tchumatchenko T, Ito HT. 2021. The orbitofrontal cortex maps future navigational goals. *Nature* **599**(7885): 449–452. doi: 10.1038/s41586-021-04042-9

56. Ormond J, O'Keefe J. 2022. Hippocampal place cells have goal-oriented vector fields during navigation. *Nature* **607**(7920): 741–746. doi: 10.1038/s41586-022-04913-9

57. Gomez-Marin A, Stephens GJ, Louis M. 2011. Active sampling and decision making in Drosophila chemotaxis. *Nat Commun* **2**(1): 441. doi: 10.1038/ncomms1455

58. Andrade IV, Riebli N, Nguyen B-CM, Omoto JJ, Cardona A, Hartenstein V. 2019. Developmentally Arrested Precursors of Pontine Neurons Establish an Embryonic Blueprint of the Drosophila Central Complex. *Curr Biol* **29**(3): 412-425.e3. doi: 10.1016/j.cub.2018.12.012

59. Goulard R, Heinze S, Webb B. 2023. Emergent spatial goals in an integrative model of the insect central complex. *PLOS Comput Biol* **19**(12): e1011480. doi: 10.1371/journal.pcbi.1011480

60. Gire DH, Kapoor V, Arrighi-Allisan A, Seminara A, Murthy VN. 2016. Mice Develop Efficient Strategies for Foraging and Navigation Using Complex Natural Stimuli. *Curr Biol* **26**(10): 1261–1273. doi: 10.1016/j.cub.2016.03.040

61. Fischler-Ruiz, W., Clark, D.G., Joshi, N.R., Devi-Chou, V., Kitch, L., Schnitzer, M., Abbott, L.F. and Axel, R., 2021. Olfactory landmarks and path integration converge to form a cognitive spatial map. *Neuron*, **109**(24), pp.4036-4049.

62. Poo C, Agarwal G, Bonacchi N, Mainen ZF. 2022. Spatial maps in piriform cortex during olfactory navigation. *Nature* **601**(7894): 595–599. doi: 10.1038/s41586-021-04242-3

63. Basu J, Nagel KI. 2024. Neural circuits for goal-directed navigation across species. *TINS* doi: 10.1016/j.tins.2024.09.005

Methods

Animals

<u>Fly strains and husbandry</u>
All flies were raised at room temp (approximately 24°-26°C) on a 12 hour light/dark cycle on standard cornmeal-agar food. For imaging experiments we used 7-12 day old female flies that were starved for ~24 hours. Experiments were run from 0-9 hours ZT. For freely walking wind tunnel experiments, we used 3-7 day old female flies. Flies were light-shifted for at least three days and then starved for 18-24 hours before experimentation. For GtACR1 flies, fly food was supplemented with hydrated potato flakes containing all-trans-retinal (35mM stock: Sigma, R2500, dissolved in ethanol, stored at −20°C).

Genotypes used in each figure were as follows:

Figure 1-3, Ext. Data 1,2	VT062617-Gal4 > UAS-GCaMP7f,UAS-tdTomato
Figure 3, Ext. Data 3	52G12-Gal4 > UAS-GCaMP7f,UAS-tdTomato
	VT062617-Gal4 > UAS-GCaMP7f,UAS-tdTomato
Figure 4, Ext. Data 4	w^{1118} 5905 (4a-c)
	VT062617-Gal4 > UAS-GtACR1 (4d,f)
	Empty-Gal4 > UAS-GtACR1 (4e,f)
	VT062617AD;VT037223DB > UAS-GtACR1 (4g, Ext. Data 4b)

Experimental Procedures

<u>Closed-loop olfactory navigation paradigm:</u>

We designed a wind and odor delivery system in which the angular position of a wind tube and the status of a binary odor valve could be controlled in either open- or closed-loop by parameters of the fly's locomotion, as measured on an air-supported walking ball. Wind and odor delivery was accomplished via a rotary union (Dynamic Sealing Technologies, # LT-2141-OF-ES12-F1-F2-C2) with three isolated channels that allowed continuous rotation of wind and odor outputs while maintaining isolation and pressure. The three channels carried wind, odor, and compensatory air for the odor OFF condition so that equal airspeed was maintained with and without odor. Air to all lines first passed through a pressure regulator (Cole-Parmer MIR2NA), and charcoal filter (Drierite 26800 Drying column, with desiccant replaced by activated charcoal). The wind line was then humidified and passed through a mass flow controller (Aalborg Model GFC17), and set to one of three flow rates: 0.2, 0.4, or 0.6 L/min. Air to the odor (0.5% apple cider vinegar) and comp (humidified air) lines was set to a flow rate of 0.3 L/min using a flowmeter (Cole-Parmer PMK1-010608). Air from the wind and odor/comp lines were mixed in a customized 3D-printed chamber that led to a 3.5cm Teflon laminarizing tube (5mm ID) extending to 2 cm from the fly's antennae. Wind, comp, and odor entered the mixing chamber sequentially to prevent back flow of odor into clean air lines.

The wind ON condition was achieved by opening a solenoid valve (Cole-Parmer Masterflex P/N 01540-11) on the wind line and a high-speed three-way solenoid valve (Lee company, LHDA1233115HA) to divert air through the comp line rather than the odor line. In the ODOR ON condition air was instead diverted through the odor solution. All three valves were positioned immediately before the rotary union. Wind and odor lines were of Tygon tubing of either 1/16" ID or 1/8" ID.

Fly behavior was monitored using a lightweight walking ball (9 mm diameter General plastics FR-7110), supported by air from a custom 3D-printed holder. Airflow supporting the ball was controlled by a flow meter (Cole-Parmer PMK1-010608) at ~0.4 L/sec. The ball was painted with spots of black ink for tracking by the FicTrac sphere tracking software[1] using a Point Grey Camera (100 fps, Grasshopper, 94mm/0.5x InfiniStix Proximity Series lens, Edmund Optics Focal Length Extenders, working distance=15cm) and illuminated using two IR LED fiber optic cables (Thorlabs M118L03 flat cleave patch cables, M850F2 fiber coupled LEDs, and LEDD1B drivers). Two secondary cameras (30fps, FLIR Blackfly, Computar 8.5mm 1:1.3 adjustable lens) viewed the front and side of the fly for positioning.

The fly was held in place using a custom 3D-printed holder made of opaque resin (Grey Resin V5, Formlabs), with a tapered saline well that opens to a .9 x .9 mm imaging window at its nadir. Resin thickness around this window is necessarily thin (~25-50 microns) to accommodate the objective's working distance to the exposed brain below the window. The holder was held in place onto a fixed steel platform using a magnetic mount (Siskiyou MGB/20).

<u>Closed-loop wind position from a walking ball</u>

During closed-loop wind delivery, we used a microcontroller (TeensyDuino +USB) and stepper motor (Oriental PKP566FMN24A, with controller CVD524-K) to rotate the rotary union on which the wind tube was mounted, as shown in Figure 1A. The amount of rotation per 10ms interval was set to be equal to the rotation around the z axis of the walking ball. Wind tube position was updated based on a running average of measured z-rotations over 10 samples at 100 samples/s. Z-rotations above 1000°/s were clipped. In

addition, the motor controller filtered out some rapid changes, leading to small amounts of error accumulation (~0-90° per trial, depending on the amount of rapid turning). Therefore, an IR beam-break (Adafruit #2168) was used to reposition the wind at 0° from the front of the fly before each trial.

Functional imaging

All two-photon calcium imaging experiments were performed under a 20x water dipping objective (Olympus Plan Fluorite XLUMP 20x/1.0). The GCaMP7f calcium and tdTomato indicators were excited using an infrared laser (SpectraPhysics MaiTai HP) at 920nm at ~20-40mW of power at the sample. Emitted fluorescence was detected using GaAsP photomultiplier tubes, and separated using green (GCaMP7f, 525/50) and red (tdTomato, 607/70) bandpass filters. Images were collected using ThorLabs LS 3.0 software, where 3-4 optical sections (typically 122 x 74 μm imaging window) were taken 8-12μm apart in the z plane, all within the fan-shaped body (determined using the tdTom signal). Volumes were taken at 4-8 vol/s with a 1.2μs dwell time.

Software control

Control of wind speed, wind position, and wind/odor state were managed by custom Python software and integrated with Fictrac ball-tracking software using an interface adapted from Python Ball Motion Tracking software from the lab of Mala Murthy. A NiDAQ board (NI PCIe-6321) and breakout board (NI BNC-2110) were used for data acquisition and communication between controllers as well as to send synchronization signals to the 2-photon microscope. Behavior data from the walking ball was collected and saved using the Fictrac image processing software at 100fps. The Fictrac timestamps (also at 100 fps) were used to trigger wind and odor changes (see experimental paradigms for timing) as well triggering 2-photon imaging (at 3s from the beginning of ball tracking).

Fly tethering and preparation

Flies were first anesthetized over ice in a glass scintillation vial for 5-10s, then held in a chilled aluminum "sarcophagus" (4-10°C) while being glued (McNett Aquaseal UV Fast Fix) into a custom 3D printed holder. Flies were only attached to the holder across the eyes and in front of the forelegs (episternum), so that the antennae and legs were clear and free to move. The head was tilted down to expose the back surface of the head in the saline well above. A minimal portion of this surface cuticle was removed directly above the FB to minimize tissue damage, prior to positioning the fly on the walking ball. Extracellular saline (103 mM NaCl, 3 mM KCl, 5 mM TES, 8 mM trehalose dihydrate, 10 mM glucose, 26 mM NaHCO$_3$, 1 mM NaH$_2$PO$_4$H$_2$0, 1.5 mM CaCl$_2$2H$_2$O, and 4 mM MgCl$_2$6H$_2$O, pH 7.1–7.4, osmolarity 270–274 mOsm), bubbled with carbogen (5% CO$_2$, 95% O$_2$) and warmed to 33°C, was continuously perfused at ~1mL/m.

Experimental paradigms

Three types of experimental paradigms were used during imaging trials: 1) Closed-loop wind, open-loop odor trials (Fig. 1, Ext Data Fig. 3c,d), where fly orientation was in constant control of the wind position and odor presentation was experimentally controlled , 2) wind-shift trials (Ext. Data Fig. 1l,m) where the wind position was transiently shifted during the odor period (+/- 90° at 45°/s) and 3) plume trials (Fig. 2,3), where the wind direction was in closed loop with the animal orientation and odor was in closed-loop with fly position in a fictive 2D space (see Figure 3A).

For plume experiments, the state of the odor valve was determined by an adaptive compression algorithm acting on odor concentrations based on virtual movement through a turbulent plume.

At each frame, fly position in virtual x,y space was first determined from FicTrac monitoring of ball motion. Fly coordinates in x,y, and time were used to index into a previously recorded movie of a turbulent boundary layer acetone plume at 6mm above the bed in 10 cm/s wind recorded with a UV laser light sheet[2]. Plume concentrations were normalized such that the peak concentration was 1.

Instantaneous odor concentration values were fed into an adaptive compression algorithm based on previous studies of fly behavior[3] and of temporal encoding at the olfactory periphery[4]. Odor concentration at the ith frame C_i was first passed through a Hill function of the form

$$x_i = \frac{C_i}{C_i + A_i + \beta}$$

where $\beta = 0.025$ is the baseline threshold and A is an adaptation term the shifts the threshold to the right based on the recent history of odor exposures:

$$A_{i+1} = A_i + (C_i - A_i)/\tau$$

where $\tau = 500$ samples (equal to 5s at 100Hz). The odor valve was turned on when $x > 0.5$ and was off otherwise.

In all experiments, we first performed 5-20 50s trials of closed-loop wind with no 2-photon imaging to acclimatize the fly to closed-loop navigation paradigm. During these trials, where wind was on for the duration of the trial, odor was presented from 20-30s, and wind speed was randomly set to one of 3 speed settings (16, 25, or 33 cm/s at the fly). After several trials, we examined behavior to see which speed was optimal for straight up-wind walking during odor and increased turning after odor offset. This speed was used for the remainder of experimental trials, although we sometimes adjusted wind speed in later trials if the fly walked less. We excluded flies that did not walk during at least 5 of these preliminary trials or only had bump activity in 2 or fewer trials.

For imaging trials, the closed-loop paradigm consisted of trial onset where wind position was randomly set to +/- 45, 135, or 0°. First the wind tube moved in closed-loop with the fly position without wind for 10s, then wind was presented at the determined speed for 15s. Odor was then presented with the wind for 15s, followed by 15s more of wind-only and 7s of no wind (total of 65s including the 3s pre-trigger time).

Experiments with plume trials also began with closed-loop speed-setting trials, followed by 10-30 plume trials that were 100s long. The fly was positioned 140 fictive mm downwind from the midline of the plume source and oriented randomly at +/- 45, 135, or 0° at the beginning of each trail. All flies were allowed to walk freely in the plume space, with no intervention if they left the plume. In this case, the fly would not receive odor until it walked back into the plume. In one fly, plume trials were run after a set of interleaved closed-loop/open-loop trials.

Behavioral experiments in wind tunnels

Freely walking wind tunnel experiments (Fig. 4, Ext. Data 4) were performed as described previously[2,5]. Briefly, the position and orientation of walking flies were tracked in shallow acrylic arenas using IR LEDS (850 nm, Environmental Lights) and a camera (Basler acA1920-155um). Constant laminar airflow was maintained at ~12 cm/s. Odor pulses of 1% apple cider vinegar (Pastorelli) were delivered at 1

Hz (500 ms duration) using solenoid valves (LHDA1233115HA, Lee Company) mounted directly below the wind tunnel. For GtACR1 inactivation experiments, blue light was delivered using a panel of LEDs (470nm, Environmental Lights) at an intensity of 61 µW/mm^2 (measured at 460 nm). For high light conditions, the intensity was set to 76 µW/mm^2 (measured at 460 nm). Flies were briefly anesthetized with ice for 30-60 seconds to load them into the wind tunnels.

Data Analysis

All analysis was performed in MATLAB (Mathworks, Natick, MA)

Processing of walking ball data

For walking ball data, angular velocity and forward velocity were calculated as the change in rotation of the ball around the z and x axes, respectively. For close-loop wind trials, heading was reported as the position of the wind tube, which was controlled by the change in z rotation, but would also account for any error in the control. For plotting and bump detection, heading, angular velocity, and forward velocity signals were smoothed using a sliding averaging window of 100 frames (0.5s). For statistical tests (e.g. Fig 1i,j Ext. Data Fig. 1a,m) and linear filter calculations (Fig. 2f, 3b,c, Ext. Data Fig. 2a-c), we used unfiltered values as described above.

Speed was calculated as the velocity in direction of heading based on the fictive fly path derived from integrating all three rotational velocities to map position on to a 2D plane[1]. X and Y position of the fly were calculated as the cumulative sum of the speed multiplied by the sine or cosine, respectively, of heading. For plume trials, X and Y position values were calculated online to enable closed-loop odor control. Upwind velocity (Fig. 3, Ext. Data Fig. 1a,3b) was calculated as the cosine of heading multiplied by speed.

Straightness (Fig. 1h, Ext. Data Fig.1a, 3e) was computed as the ratio of displacement (the distance of a straight line from the beginning and end points of a window) to path length (the sum of all displacements within the window) over a sliding 1.5s window. This value was then squared. Persistent walking duration (Ext. Data Fig. 1b,c) was determined as the amount of time walking in the goal direction before a heading deflection of more than ±45° in trials with bump activity (see below bump detection methods) and pooled across flies. Goal direction (Ext. Data Fig. 1b) was determined as the heading at odor OFF for odor walks, and heading at trial start for baseline. The distribution of durations was compared to the distribution of straight walking from the trial onset (Ext. Data. Fig. 1c), using identical criteria as post odor runs.

Exponential curves were fit to these distributions using a non-linear least-squares process. The curve was defined as:

$$A \times e^{-x/tau} + C$$

where A is the amplitude of the starting value, tau is the decay rate, and C is the offset. Optimal fit was determined by minimizing mean squared error. A 95% confidence interval (CI) for tau was given by:

$$tau \pm t\sqrt{S}$$

where *t* is computed using the inverse of Student's *t* cumulative distribution function, and *S* is a vector of the diagonal elements from the estimated covariance matrix of the coefficient estimates, $(X^TX)^{-1}s^2$, where *X* is the design matrix, X^T is the transpose of *X*, and s^2 is the mean squared error (using the Curve Fitting Toolbox from Mathworks).

Processing of imaging data

For fluorescence extraction, all image volumes were first reduced to single planes as maximum intensity projections across each z plane. Maximum intensity projections were highly correlated with mean projections, but had slightly higher signal-to-noise. We then concatenated projections for all trials of an experiment, and used the NoRMCorre algorithm[6] to correct for rigid motion across both red and green channels. We drew ROIs by hand in ImageJ, using a maximum intensity projection across all frames of a single trial (typically the first trial) on the red tdTOM channel. We then imported these ROIs into Matlab, and aligned them to subsequent images using register_rois2[7]. For both VT062617 and 52G12 imaging, we drew ROIs for each of 8 putative columns in the dorsal layer of the FB. A baseline value for each ROI was calculated as the mean of the lowest 3% of fluorescent values across all trials. $\Delta F/F$ was then calculated by subtracting this baseline value from the green fluorescence values and dividing them by the baseline value. Fluorescence timeseries data was synchronized to the timestamps of behavior data using the 3s trigger time.

Analysis of neural encoding properties

To detect activity bumps, we used a threshold-based algorithm, where the threshold for $\Delta F/F$ was set to 0.5 standard deviations above the mean $\Delta F/F$ of the entire experiment. A minimum bump duration was set at 3s and bumps with gaps <3s from a consecutive bump were merged. Using this detector, we classified samples of behavior data as either bump or no-bump samples, and detected bump onset and offset times. Only flies with more than 2 trials showing bump activity were included.

To generate distributions of post-odor bump durations (Fig. 1f,g, and Ext Data. Fig 1i), we analyzed all bumps that began during the odor period and ended after odor OFF.

To analyze behavior associated with bump offset or bump onset (Fig. 1i,j, Fig. 2d, Ext. Data Fig. 2d), we again only included bumps that turned ON during odor presentation and turned OFF after odor OFF. We computed locomotor parameters (absolute value of angular velocity and forward velocity) as described above. For bump offset analysis, we also computed goal deviation as the absolute value of the difference between the absolute value of heading at each time point after odor OFF and a goal heading (θ_{goal}), defined as the mean heading during the odor period. We then aligned each behavioral parameter by the time of bump OFF or ON and took the mean across all trials from all flies. For comparison (control), we aligned behavioral data to the midpoint of each bump. Statistical comparisons were made over a window from 0-1s after bump off (vs bump midpoint) for angular speed, and for a window from 0-5s after bump off (vs bump midpoint) for goal deviation. For bump onset (Ext. Data Fig. 2d), we compared mean forward velocity over a window -7 to -4 s before bump onset to a window -1.5 to 1.5 s from onset of each bump. A paired two-sample t-test was used for all comparisons.

To compute bump width in Ext Data Fig. 1g, we aligned each bump sample to the ROI with the peak $\Delta F/F$, and then wrapped the other ROI values to the 8 column space using the modulus function in Matlab. Aligned bumps were averaged across trials and flies.

To compute bump position in Ext. Data Fig. 1j-m, we first smoothed fluorescence traces using a 1s sliding window in the time domain and a two-column window in the spatial domain. We then computed bump position as the maximum ROI (or FB column) for each sample where a bump was detected using our bump finder algorithm above. Positions were then unwrapped to show absolute column shifts. We computed correlations between bump position and heading for all bump instances (Ext. Data Fig. 1k top) and averaged across each fly with bump activity in at least 5 trials (Ext. Data Fig. 1k bottom) using Pearson's correlation coefficient.

To compute bump position shifts associated with a rotation in wind direction, in Ext. Data Fig. 1l,m),

we analyzed only trials in which a bump was present before and during the shift. Wind driven heading shifts were detected when mean angular velocity during or up to 2s after the shift was greater than a time window of equal duration before the shift. Heading and position for right shifts were multiplied by -1 so all shifts could be directly compared on the same axes. Heading and bump position were both converted to degrees to facilitate comparison. Displacements were calculated over time windows 2s before and after the wind shift onset.

To compute the dependence of bump probability on odor frequency in Fig 2e, we analyzed all trials in which flies experienced repeated odor pulses. Odor frequency was estimated by convolving the binary odor-on signal with an exponential decay kernel ($\tau = 2$ s), creating a continuous signal that reflected the recent history of odor encounters. For each odor offset, we extracted the odor frequency at the moment of offset and quantified bump probability during the subsequent 0–2.5 second window. Bump probability was defined as the proportion of time points in this window where the bump was active. Each data point represents one odor offset event from an individual trial. Trials were grouped into 3 1Hz odor frequency bins based on the offset value, and bump probabilities were averaged across events within each bin. Final values reflect mean bump probabilities per bin for each fly.

To compute linear filters relating odor experience and locomotor parameters to neural activity in Fig. 2f, Fig. 3b,c, Ext. Data Fig. 2a-c, we first computed bump amplitude, forward velocity, and angular velocity as described above. The first 400 samples of each trace were discarded as the bump amplitude typically contained an artifact due to shutter opening. The remaining traces were concatenated to form a single vector for each parameter and all traces were mean subtracted. Forward velocity and absolute angular velocity were normalized to have unit standard deviation. We then computed the cross-correlogram between bump amplitude and odor, forward velocity, and absolute angular velocity in the frequency domain.

To account for correlations between odor and locomotor parameters, we adapted Methods from[9] to account for correlations between odor, forward velocity, and angular velocity. Briefly, we computed the complex autocorrelation matrix A at each of 256 frequencies from 0.2 to 5 Hz of the odor o(t), forward velocity f(t), and absolute angular velocity a(t) traces:

$$A_\omega = \tilde{S}'_\omega \tilde{S}_\omega$$

where

$$S_\omega = \begin{bmatrix} \overrightarrow{o_\omega} & \overrightarrow{f_\omega} & \overrightarrow{a_\omega} \end{bmatrix}$$

We then computed the pseudo-inverse of A for each frequency ω by taking the singular value decomposition:

$$U\Sigma V' = A_\omega$$

and inverting the diagonal terms of Σ with a regularization parameter λ that was set to 0.0001.

$$\Sigma_{inv}(s, s) = \frac{1}{\Sigma(s, s)} + \lambda$$

The decorrelated filter was equal to:

$$\tilde{D} = V\Sigma_{inv}U'\tilde{C}$$

Where $\tilde{C}$ is the Fourier transform of the raw cross-correlogram for each parameter:

$$C_{\omega} = \begin{bmatrix} \overrightarrow{C_{o\omega}} & \overrightarrow{C_{f\omega}} & \overrightarrow{C_{a\omega}} \end{bmatrix}$$

To compare filters from different fly lines in Fig. 3b,c, we normalized all filters by dividing by the standard deviation of the filter and subtracting minimum standard deviation values. To compare half-widths of filters, we first smoothed each filter using a sliding window of 1s. We then found the half-prominence width of peaks determined as local maxima where the prominence was a minimum of 1.5 times the mean for each fly (Matlab findpeak function, with defined prominence properties). To compare angular velocity filters, we found the area under the curve for each fly using a trapezoidal estimation for each data point (Matlab trapz function).

To compute the correlation between upwind velocity and bump activity on a trial-by-trial basis in Fig. 3d,e, we measured upwind velocity during the odor period for each trial. We then computed Pearson's correlation coefficient between these metrics and the mean ΔF/F for the same periods and trials. To assess significance, we compare these to correlation coefficients between metrics and ΔF/F for shuffled trials (null condition).

Analysis of freely walking wind tunnel behavior data:

Freely walking behavioral data were preprocessed as described previously[2,5]: trials with less than 25 mm of movement were excluded, and position and orientation raw data were low pass filtered at 2.5 Hz. As upwind orientation could persist if a fly tracks the arena walls or stops walking, we further filtered our data by excluding data points where the fly's position was within 3 mm of a wall or the fly's velocity was below 0.5 mm/s.

Upwind walking probability was determined by binarizing the orientation data depending on whether the orientations were inside or outside of a 90° cone centered on upwind. This binarized data was further processed in three ways to evaluate persistence and evidence accumulation. First, the binarized data were averaged at each timepoint to compute the upwind walking probability over time (Fig. 4b,d,e). Second, the upwind walking probabilities over time were averaged between 2 s and 6 s following odor onset to compute peak responses (Fig.4c). Third, the binarized data between 0 and 10 s following odor offset were averaged across trials for each fly to compute post-odor upwind orientation probabilities (Fig. 4f,g).

Modeling persistent goal-directed navigation in an odor plume

We implemented a three-state behavioral model to simulate fly navigation in a natural odor plume environment. At each time step, the agent occupied one of three discrete states: baseline walking, goal-directed upwind walking, or local search. Transitions between states were governed by odor inputs and probabilistic rules inspired by our observed neural dynamics. The agent received odor concentration values sampled from the same measured plume used in imaging experiments described above, and the raw odor signal was passed through the same adaptive filter and compression function with $\beta = 0.025$ and $\tau_A = 5s$.

When the compressed odor signal C_t exceeded a threshold (θ=0.5), the agent transitioned from baseline or search into the goal-directed state. The agent transitioned from the goal state to the search state with a probability defined by the persistence time constant τ:

$$P_{leave} = \frac{1}{\tau_p \cdot f_s}$$

where f_s is the simulation frame rate (15 Hz). From the search state, it could return to the goal state if odor reappeared, or stochastically transition back to baseline with a fixed time constant (6.7s).

Behavior in each state was described by a modified version of the model in[26]. Briefly, locomotion was controlled by a network of 5 descending neuron-like units that receive independent Gaussian noise $\vec{\omega}(t)$ and interact through a connectivity matrix M. U_3 drives stopping while the remaining units drive forward and angular velocity through their sums and differences:

$$v_t = \alpha_1 \cdot \max(0, U_1 + U_5) + \alpha_2 \cdot \max(0, U_2 + U_4)$$
$$a_t = \gamma_1 \cdot (U_1 + U_5) + \gamma_2 \cdot (U_2 + U_4)$$

We adjusted the parameters α_1, α_2, γ_1, γ_2 to approximately match the distribution of forward and angular velocities measured on our walking ball during boundary layer plume navigation (Fig. 5c).

	Walking	Flight
α_1	0.7°/s	40°/s
α_2	0.3°/s	30°/s
γ_1	0.1mm/s	0.9mm/s
γ_2	0.2mm/s	0.9mm/s

The matrix M has the structure

$$M = \begin{bmatrix} A & 0 & I_s & I_c & I_c \\ 0 & A & I_s & I_c & I_c \\ I_s & I_s & A & I_s & I_s \\ I_c & I_c & I_s & A & 0 \\ I_c & I_c & I_s & 0 & A \end{bmatrix}$$

During the baseline and goal states, parameters were $A = 0.8$, $I_s = -0.03$, $I_c = 0$. During the search state, $I_c = -0.035$. The inhibition between the two sides present during the search state reduces forward velocity and increases both angular velocity and the duration of runs of turns in the same direction, as described in[26]. During the goal state, a sinusoidal function of wind direction was added to the locomotor units to promote an upwind heading and the stop unit was inhibited to promote greater locomotion as in[26]:

$$\vec{u}(t + \Delta t) = a(M\vec{u}(t) + [1\ 1\ 0 -1 -1]'sin(\theta(t) - \psi) + [0\ 0 -0.1\ 0\ 0]' + \vec{\omega}(t))$$

Activity in each unit was bounded by a sigmoidal activation function that prevents runaway excitation:

$$a(x) = \frac{20}{1 + e^{-x/4}} - 10$$

To assess how the persistence parameter τ_p influences navigation success, we ran the model across a range of τ_p values in the near-surface (boundary layer) air plume described above[3] (Fig. 5d,e). The plume environment measured 300.44mm x 159.84 mm. For each τ_p, 500 trials were simulated with randomized start positions downwind from the source by 222±12mm and centered along the midline of the plume's long axis, ±100mm. Success was quantified by calculating the proportion of time the agent's position fell within a 15 mm radius of the odor source.

To visualize where the agent engaged in each behavioral state (Fig. 5d), we computed spatial heatmaps showing the probability of occupying the baseline, goal-directed, or search state across the plume environment. The agent's position and corresponding state label were divided by 15cm x 15cm bins, and the relative frequency of each state within each bin was calculated. These probability maps were smoothed using a Gaussian filter (σ = 15mm) to highlight spatial patterns.

Table 1: Reagents and Resources

Resource	Source	Identifier
FLY LINES		
VT062617-Gal4	VDRC	VDRC: 206875
52G12-Gal4	BDSC	RRID: BDSC_49581
Empty-Gal4	BDSC	RRID: BDSC_68384
w^{1118} 5905	Alvarez-Salvado et al. 2018	
UAS-GCaMP7f,UAS-tdTomato	BDSC/Nagel lab	RRID: BDSC_80906, 36327
w^{1118},s/CyO;UAS-GtACR1-EYFP/TM6b	Claude Desplan	
VT062617AD/[CyO,tb];VT037223DB/TM6b	Tanya Wolff	Janelia: SS63158
p65ADZp, ZpGAL4DBD (Empty Split-Gal4)	BDSC	RRID: BDSC_79603
SOFTWARE		
Python		Version 3.8
MATLAB	Mathworks	Version R2023a
Image J		Version 1.53t
FicTrac		Version 2.0 (build date: Nov 2 2018)
Pybmt	Mala Murthy lab	
CalmAn		
Register_roi2	Jason Moore	

Data Availability

Data will be made publicly available upon publication at Zenodo.

Code Availability

Code will be made publicly available upon publication at Github.

Methods References

1. Moore RJD, Taylor GJ, Paulk AC, Pearson T, Swinderen B van, Srinivasan MV. 2014. FicTrac: A visual method for tracking spherical motion and generating fictive animal paths. *J Neurosci Methods* **225**: 106–119. doi: 10.1016/j.jneumeth.2014.01.010

2. Connor EG, McHugh MK, Crimaldi JP. 2018. Quantification of airborne odor plumes using planar laser-induced fluorescence. *Exp Fluids* **59**(9): 137. doi: 10.1007/s00348 018-2591-3

3. Álvarez-Salvado E, Licata AM, Connor EG, McHugh MK, King BM, Stavropoulos N, Victor JD, Crimaldi JP, Nagel KI. 2018. Elementary sensory-motor transformations underlying olfactory navigation in walking fruit-flies. *Elife* **7**: e37815. doi: 10.7554/elife.37815

4. Nagel KI, Wilson RI. 2011. Biophysical mechanisms underlying olfactory receptor neuron dynamics. *Nat Neurosci* **14**(2): 208–216. doi: 10.1038/nn.2725

5. Matheson AMM, Lanz AJ, Medina AM, Licata AM, Currier TA, Syed MH, Nagel KI. 2022. A neural circuit for wind-guided olfactory navigation. *Nat Commun* **13**(1): 4613. doi: 10.1038/s41467-022-32247-7

6. Pnevmatikakis EA, Soudry D, Gao Y, Machado TA, Merel J, Pfau D, Reardon T, Mu Y, Lacefield C, Yang W, et al. 2016. Simultaneous Denoising, Deconvolution, and Demixing of Calcium Imaging Data. *Neuron* **89**(2): 285–299. doi: 10.1016/j.neuron.2015.11.037

7. Giovannucci A, Friedrich J, Gunn P, Kalfon J, Brown BL, Koay SA, Taxidis J, Najafi F, Gauthier JL, Zhou P, et al. 2019. CaImAn an open source tool for scalable calcium imaging data analysis. *eLife* **8**: e38173. doi: 10.7554/elife.38173

8. Demir M, Kadakia N, Anderson HD, Clark DA, Emonet T. 2020. Walking Drosophila navigate complex plumes using stochastic decisions biased by the timing of odor encounters. *eLife* **9**: e57524. doi: 10.7554/elife.57524

9. Nagel KI, Doupe AJ. 2008. Organizing Principles of Spectro-Temporal Encoding in the Avian Primary Auditory Area Field L. *Neuron* **58**(6): 938–955. doi: 10.1016/j.neuron.2008.04.028

Acknowledgements

The authors would like to thank Mala Murthy and her lab for providing modified FicTrac code (Pybmt) and advice on closed-loop behavior, John Crimaldi and his lab for plume data, and Margot Elmaleh in Michael Long's lab for advice on 3D printing. Claude Desplan's lab provided GtACR1 flies. David Schoppik provided assistance with 2-photon microscopy. Christina May generated imaging stocks and provided support for 2-photon microscopy. Emily Hao provided assistance with behavioral experiments. Jason Moore provided assistance with imaging analysis. Jonathan Victor provided advice on regularizing the filter decorrelation. Michael Long, Jonathan Victor, Bard Ermentrout, Floris van Breugel, and members of the Nagel and Schoppik labs provided input on the manuscript. This work was supported by NINDS Brain Initiative NS127129, NIDCD DC017979, and NSF 2014217 Odor2Action to K.I.N.

Author Contributions

N.D.K. and K.I.N designed the study. N.D.K. developed the closed-loop wind and odor apparatus, developed all all hardware and software related to the apparatus, and collected and analyzed all behavior and imaging data using this apparatus. A.J.L designed, ran, and analyzed freely walking behavioral data and contributed analysis methods and code. K.I.N. wrote code for the filter analysis. J.D.F. obtained confocal data. N.D.K. and K.I.N. wrote the paper with input from A.J.L.

Competing Interest Declaration

The authors declare no competing interests.

Correspondence

Correspondence and requests for materials should be addressed to Katherine Nagel at
katherine.nagel@nyumc.org.

Figures and Legends

Figure 1: Persistent odor-evoked activity in VT062617 FB local neurons is associated with persistent goal-directed walking. a: Schematic of the preparation. A head-fixed fly walking on an air-supported ball can control wind direction (through a motorized rotary union) and odor (apple cider vinegar 0.5%) in open- or closed-loop. An IR camera tracks ball rotations. A 2-photon microscope images concurrent neural activity. **b**: Example trajectories from one fly performing closed-loop wind navigation. Odor evokes straight trajectories that are biased towards upwind directions. Yellow star: starting position, purple bars: odor ON, odor OFF. **c**: Examples of VT062617 activity during closed-loop wind with an odor pulse. Top: virtual trajectories, color-coded by bump amplitude (max ΔF/F across columns). Black bars indicate wind ON and wind OFF, purple

bars indicate odor ON and odor OFF and red arrows indicates bump OFF. Note strongest activity during and after odor. Middle heatmap: activity across FB columns as a function of time. Wind and odor periods indicated above heatmaps, red arrow indicates time of bump OFF (left: 5.1s, right: 5.8s after odor OFF). Bottom: heading trace, 0° is upwind, black arrow indicates start of turn after straight walk **d**: Bump amplitude as a function of time for one fly across trials. Note robust responses to odor with variable offset timing, and more variable and transient responses to wind ON. **e**: Mean time course of bump activity relative to odor (purple box). Black: mean for each fly (n=13-39 trials), red: mean across flies (N=14 flies). Note activity following odor OFF (black arrow). **f:** Normalized bump amplitude for all trials with bump activity during odor (n = 135), aligned at odor OFF and sorted by bump persistence duration. **g**: Probability distribution (black) of bump persistence durations (n=135 trials from N=14 flies). Exponential fit (dashed line) with confidence intervals (shaded in grey, tau = 7.8±1.5s). **h**: Trajectory straightness in the presence vs absence of a bump for odor and post-odor periods for each fly (N=14 flies). Mean straightness was higher in the post-odor period for samples with a bump (paired t-test, p=0.02), but not during odor (p=0.26). **i**: Trajectory features associated with bump offset in the post-odor period (red), compared to periods when the bump remained on (black). Lines show mean bump amplitude (top), angular speed (middle), and deviation from goal (bottom, see Methods) for n = 135 bumps from N=14 flies. Shaded regions represent S.E. across bumps. **j**: Deviation from goal direction (bottom) increases significantly after bump OFF, but angular speed does not (comparison periods in blue in **i**, paired t-test, avel: p=0.21, goal dev.: p=0.02).

Figure 2: Evidence integration dynamics during closed-loop plume navigation. a: Closed-loop plume navigation paradigm. Time and fictive fly position are used to index into a previously recorded movie of plume dynamics. Measured odor concentrations are converted into binary odor signals based on an adaptive compression algorithm (see Methods). The frequency of the resulting odor pulses depends on location and distance from the source. **b**: Example trial from a fly navigating the virtual plume. Left: virtual 2D trajectory color-coded by bump amplitude, superimposed on a still of the plume movie. Purple dots indicate fictive locations at which odor was presented. Right: activity across FB columns (top), bump amplitude (middle), and heading as a function of time (bottom). Purple bars indicate odor periods. Note minimal responses to early odor presentations and gradual ramping of activity over time. **c**: Trajectory and activity for another example trial. Note prolonged period of activity associated with straight walking. **d**: Left:

Analysis of locomotor features around the bump OFF period (red) compared to the bump ON period (of equal length, black) during plume navigation (n = 127 bumps from N= 8 flies). Shaded regions represent S.E. across bumps. Right: Both angular speed and goal deviation were significantly higher over the time window shaded in blue (left) (paired two-sample t-test, avel: 1s time window, p = 0.03, goal dev: 5s time window, p = 0.002, N=8 flies) **e**: Probability of bump activity as a function of odor encounter frequency (see Methods) from 0 to 2.5s after last odor OFF for each fly (N=6 flies with at least 5 trials with bump activity). Bins centered at 0.5 and 2.5 were significantly different across fly means (black dots show means for each fly, red line is mean across flies, two-sample t-test, p = 0.04). **f.** Linear filters relating neural activity to odor, angular speed, and forward velocity, each mean +/- S.E. for N=6 flies. Odor and forward velocities have an integrating shape with a longer integration time for odor than for forward velocity. Angular speed filter is negative indicating a correlation with decreased angular speed (straight walking).

Figure 3: Distinct local dynamics and shared slow dynamics in a population of ventral local neurons.
a: Example trial from a fly expressing GCaMP in 52G12 neurons navigating the virtual plume. Left: trajectory color-coded by bump amplitude. Right: activity across FB columns (top), bump amplitude (middle), and heading (bottom) as a function of time. Purple bars indicate odor periods. **b**: Linear filters relating odor to activity in 52G12 neurons (green, mean +/- S.E. from N=5 flies) and VT062617 neurons (orange, replotted from Fig. 2). 52G12 filters are significantly narrower than VT062617 filters (two-sample t-test, p=0.03). **c**: Linear filters relating angular speed to activity in 52G12 versus VT062617 neurons. 52G12 filters have a positive area under the curve, indicating a correlation with turning, while VT062617 filters have a negative area (two-sample t-test, p=0.015). **d**: Example data from 4 flies (top: VT062617, bottom: 52G12) showing upwind velocity during odor (color bar, left) and bump amplitude (right) across trials. **e**: Trial-by-trial correlation coefficients between upwind velocity and total bump amplitude for each fly are significantly higher than for trial-shuffled control data (paired t-test, p-values: VT062617, p=0.001; 52G12, p=0.013).

Figure 4. VT062617 neurons are causally required for persistent upwind navigation. **a**: Schematic of behavioral apparatus used to track freely moving flies, showing odor and wind inputs. **b**: Probability of upwind walking as a function of time, for different numbers of odor pulses (500 ms each) in wild type flies (1 pulse: 246 trials from N=30 flies; 4 pulses: n=226 trials from N=28 flies; 7 pulses: 213 trials from N=29 flies; 10 pulses: 246 trials from N=30 flies). **c**: Mean probability of upwind walking during peak response period (2-6 s, relative to odor start) for each pulse number. Data are mean +/- S.E. across flies. **d**: Probability of upwind walking in VT062617>GtACR flies responding to 1, 5, or 10 pulses of odor with light OFF (black, 1 pulse: n=381 trials from N=30 flies; 5 pulses: n=418 trials from N=30 flies; 10 pulses: n=463 trials from N=30 flies) or light ON (blue, 1 pulse: n=405 trials from N=30 flies; 5 pulses: n=422 trials from N=30 flies; 10 pulses: n=450 trials from N=30 flies). Note decrease in upwind walking probability after odor OFF on light ON trials across stimuli. **e**: Probability of upwind walking in empty-gal4>GtACR flies responding to the same stimuli with light OFF (black, 1 pulse: n=261 trials from N=26 flies; 5 pulses: n=246 trials from N=26 flies; 10 pulses: n=318 trials from N=28 flies) and light ON (green, 1 pulse: n=294 trials from N=26 flies; 5 pulses: n=310 trials from N=28 flies; 10 pulses: n=292 trials from N=27 flies). **f**: Left: Mean probability of upwind walking during the 10 s post-odor period in VT062617 > GtACR1 flies during light ON (blue) is significantly lower than in light OFF (grey) trials (Bonferroni-corrected (α=0.0167) ranksum test: 1 pulse (N=26): p<0.0001, 5 pulses (N=26): p=0.0046, 10 pulses (N=26): p=0.0004). Right: same comparison, but with empty-gal4 > GtACR1 flies during light ON (green) and light OFF (grey) trials: 1 pulse (N=25): p=0.474, 5 pulses (N=27): p=0.046, 10 pulses (N=28): p=0.399 **g**: Left: Mean probability of upwind walking during the

10 s post-odor period in hΔK split > GtACR1 flies during light ON (red) is significantly lower than in light OFF (grey) trials after 10 pulses (Bonferroni-corrected (α=0.017) ranksum test: 1 pulse (*N*=23): *p*=0.276, 5 pulses (*N*=24): *p*=0.065, 10 pulses (*N*=24): *p*=0.0079). Right: same comparison, but with empty-gal4 > GtACR1 flies at the same light level during light ON (green) and light OFF (grey) trials: 1 pulse (*N*=26): *p*=0.764, 5 pulses (*N*=28): *p*=0.991, 10 pulses (*N*=29): *p*=0.417

Figure 5. Measured persistence times closely match optimal memory durations for a walking fly navigating a boundary-layer plume.

a. Schematic of the state transition model. The agent switches between three behavioral states, baseline (gray), goal-directed upwind walking (blue), and local search (green) based on odor inputs and internal probabilistic dynamics. Transitions between the goal state and the search state are governed by a tunable time constant τ_p. **b.** Still frame of a boundary layer plume as the simulation environment. Dashed black rectangle is starting zone for each trial. Symbols define graphical elements used in subsequent panels: yellow start is trial start location, red square and triangle is the plume sampled area and odor envelope estimate, red circle is the "success" area for finding the source location; colored lines are state annotations for example trajectories. **c**. Distributions of locomotor parameters (forward and angular velocity) for real flies in our walking ball data (red) and simulated agents (black) during baseline walking. Mean forward velocity was 2.4 mm/s and mean angular velocity was 39.1 deg/s. **d.** Example trajectories from the model (left) and state-specific occupancy maps (right) showing positions during baseline (top), goal (middle), and search

(bottom) states for three different values of τ_p (rows). At the optimal τ_p, the agent adopts that goal state at most locations within the plume envelope and the search state is concentrated at the edges of the plume envelope. **e.** Success rate (see Methods) as a function of τ_p for 500 runs of the model. Peak performance occurs at an intermediate τ_p value 6.4s. Red line is the mean, blue box shows middle quartiles. Red arrow shows measured τ_{real} = 7.8s from VT062617 imaging.

Ext. Data Fig. 1: Additional data on VT062617 FB local neuron responses.
a: Characteristics of virtual trajectories during baseline versus odor periods for all flies (N=15). Each line represents the mean for one fly. Mean upwind velocity and straightness are significantly higher during odor than baseline (paired t-test, p =0.001, p=0.035). Forward velocity is not significantly different (p=0.43). **b.** Heading distribution of persistent walking during baseline (left) and odor OFF (right) periods for all trials with bump activity (n = 135 trials from N=14 flies). Mean resultant vector during post-odor (red) is biased upwind

(mean: -1.5°). The length of each line represents the duration the animal walked straight after the respective time (odor: odor OFF, baseline: trial start). c. Distribution of persistent straight walking durations after odor (purple) and during baseline (grey) for each fly (N=14). Exponential curves (dashed) with functional confidence intervals (shaded) were fit to the data, providing exponential decay values (tau_{odor} = 8.8±6.7s, tau_{base} = 2.5±0.3s). **d**: Top: Confocal image of VT062617>GCaMP brain, stained for GCaMP (pink) and nc82 (blue). Fluorescence signals were imaged from the dorsal (output) projections of FB local neurons. Scale bar is 50µm. Bottom: table of cell body counts for left and right hemispheres in two brains. **e**: Distribution of ΔF/F values across all recordings in the presence of odor (purple), during wind only (green) and in the absence of stimuli (black). Total of 328 trials from 15 flies for each condition. **f**: Average bump amplitude in trials with no wind or odor stimuli, but motor still in closed-loop with heading (n=3-7 trials, N=3 flies). **g**: Mean bump profile across columns. Thin lines represent mean across trials (9 to 52 trials for each fly) and thick lines represent mean across flies (N=14 flies with both odor and no-odor imaging data). **h**: Photoionization detector (PID) recording of odor concentration during typical wind and odor pulse trials (n=6 trials, black: individual trials, red: mean across trials) at the location of the fly. Wind and odor control signals above. **i**: Bump persistence distributions of individual flies are significantly different from one another (one-way ANOVA, p = 0.004, n = 135, N = 14). Each row is from a separate fly, colored individually, and ordered along the y axis by mean duration. **j**: Example trial where bump position does not change with changes in heading. Top: fictive 2-D trajectory color-coded by bump amplitude; symbols as in Fig. 1c. Bottom: Heatmap shows activity across columns with bump position (green) and heading (red) overlaid. Heading for whole trial shown below. **k**: Distribution of correlation coefficients between bump position and heading for all trials from all flies (top, n=196 trials) and averaged for each fly (N=14 flies). Pooled data were not significantly different from zero (one-sample t-test, mean=0.043, p=0.220). Fly averages showed a very small positive correlation (mean=0.095, p=0.03). **l**: Response of a single fly to a 90° wind shift during odor. Top: fictive 2-D trajectory color-coded by bump amplitude; symbols as in Fig. 1c. Bottom: allocentric wind direction (top), VT062617 activity across columns (middle), and heading (bottom). Note stability of bump position during heading deviation following wind shift. **m**: Left: Heading response to open-loop wind shift in 8 trials from 4 flies that responded significantly to the wind shift (see Methods). Dashed lines represent start and stop of shift. Middle: Bump position for the same trials shown to the left, offset to an initial position of 0. Note lack of consistent movement. Right: Displacement of heading and bump position for the 8 trials shown. Heading displacement is significantly different from zero (mean: 25.41°, one-way t-test: p=0.006) while bump displacement is not (mean: 9.87°; one-way t-test: p=0.072).

Ext. Data Fig. 2: Additional data on plume related activity

a: Linear filters for odor (**a**), angular speed (**b**), and forward velocity (**c**) for each fly. **d**: Analysis of locomotor activity relative to bump onset. Left: lower trace shows forward velocity aligned by bump onset (upper), which peaks at bump onset. Red traces represent means across flies (N=14 flies). Right: Mean forward velocity increases during bump onset (pre-onset mean=2.13, post-onset mean=3.60, paired t-test p=0.046, blue boxes indicate analyzed periods.) **e**: Mean forward velocity during odor is higher in samples with a bump than without (paired t-test, p=0.032, N=14).

Cell Count

Line/brain	Upper Left	Lower Left	Upper Right	Lower Right	Total
52G12-1	59	30	66	24	179
52G12-2	75	34	67	36	212
Mean	67	32	66.5	30	195.5

Ext. Data Fig. 3: Additional data on responses of ventral local neurons labeled by 52G12. a: Confocal image of 52G12>GCaMP brains, stained for GCaMP (pink) and nc82 (blue). Scale bar is 50µm. Bottom: table of cell body counts for left/right/upper/lower quadrants of two brains imaged. **b**: Mean upwind velocities of expressing GCaMP in 52G12 neurons during odor versus baseline (N=11 flies, p=0.018). **c**: Mean activity in 52G12 neurons over time for all trials for one fly. Note transient responses to wind and odor onset. **d**: Time course of activity relative to measured odor time course (purple box) for all recorded flies. Black: mean for each fly (n=6 to 30 trials each), red: mean across flies (N=11 flies). **e**. Left: Mean straightness for 52G12 flies (N=11) in the odor and post-odor periods is not significantly different in the presence versus absence of a bump (paired t-test, during odor p=0.966, post-odor p=0.893). Right: Angular speed during odor is significantly higher in the presence of a bump (paired t-test, p=0.029)

Ext. Data Fig. 4: Additional data on silencing of hΔK neurons

a: Probability of upwind walking in hΔK split > GtACR flies responding to 1, 5, or 10 pulses of odor (500 ms at 1Hz) with light OFF (black, 1 pulse: n=287 trials from N=23 flies; 5 pulses: n=271 trials from N=24 flies; 10 pulses: n=297 trials from N=24 flies) or light ON (red, 1 pulse: n=362 trials from N=23 flies; 5 pulses: n=367 trials from N=24 flies; 10 pulses: n=322 trials from N=24 flies). Light level for these experiments was 76 μW/mm^2. We observed a decrease in upwind walking probability after odor OFF on light ON trials for the 10 pulse condition. **b**: Probability of upwind walking in empty-split > GtACR flies responding to the same stimuli with light OFF (black, 1 pulse: n=150 trials from N=26 flies; 5 pulses: n=234 trials from N=28 flies; 10 pulses: n=213 trials from N=29 flies) and light ON (green, 1 pulse: n=357 trials from N=26 flies; 5 pulses: n=324 trials from N=28 flies; 10 pulses: n=358 trials from N=29 flies).