Title: Gradual consolidation of skilled sequential movements in primary motor cortex of non-human primates

Short Title: Consolidation of sequential movements in motor cortex

Authors: Machiko Ohbayashi[1,2]*, Nathalie Picard[1,2]

Affiliations:

1. Department of Neurobiology, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania 15261, USA.
2. Systems Neuroscience Center, University of Pittsburgh, Pittsburgh, Pennsylvania 15261, USA.

* Corresponding author: Machiko Ohbayashi (machiko@pitt.edu)

Abstract

Many of our daily actions rely on skilled sequential movements. Sequence performance improves with practice and can reach expert levels with years of repeated practice, truly exemplifying the saying "Practice makes perfect". Human and non-human primate studies have shown structural and functional changes in the primary motor cortex (M1) following extended practice of sequential movements, suggesting M1's involvement in acquisition and retention of skilled sequences. However, it has been challenging to causally examine M1's role in sequence learning, as inactivation or lesion of M1 impairs movement production. Here, we causally examined M1's contribution to sequence learning by locally inhibiting protein synthesis in M1. Our results show that protein synthesis inhibition in M1 disrupted memory-guided sequential movements at all stages of learning without affecting movement production, though the effects decreased with continued practice. These findings suggest that neural traces for sequential movements are repeatedly consolidated in M1 through protein synthesis, with the rate of consolidation slowing as learning progresses.

Teaser

Repetitive practice strengthens neural representations of sequential movements in primary motor cortex of monkeys.

Introduction

Many daily activities, such as playing a musical instrument and typing, rely on attaining a high level of proficiency in performing sequential movements. The sequential movements can be acquired and improved to an expert level through extensive practice, eventually becoming a stable motor skill. Once the structure of a sequence is learned during the early phase of learning, continued daily practice over an extended period leads to gradual improvement in speed and accuracy, key indicators of motor skill learning. This slow incremental learning is considered a fundamental aspect of motor skill development, capturing the idea that "practice makes perfect."

Emerging evidence indicates that multiple cortical motor areas alter their activity during the learning of sequential movements (*1-11*). The primary motor cortex (M1), in particular, exhibits functional and structural changes associated with the learning of the sequential movements, not only during the early stages of learning but also after extensive practice. In humans, short-term training induces measurable changes in fMRI activity within M1 (*1-9, 12, 13*). Similarly, studies in monkeys demonstrate that neural activity in M1 is modulated by sequence components during the early stage of learning (*14, 15*).

With extensive practice, additional changes in M1 have been reported in both human and non-human primate. In humans, years of extensive practice are associated with structural modifications and altered functional activation of M1 (*9, 11, 16-20*). For instance, professional musicians who have spent years practicing complex sequential movements exhibit reduced or more focused M1 activation during the performance of sequential tasks compared with non-musicians or amateurs (*16-20*). This reduced activation is considered as evidence of increased efficiency of neural circuits, or the need for a smaller number of active neurons to perform a highly trained set of sequential movements. Furthermore, the volume of M1 has been reported to be larger in professional musicians compared to amateurs or non-musicians (*21-28*). These structural changes have been proposed to arise form synaptic processes, including intracortical remodeling of dendritic spines and axonal terminals, glial hypertrophy, and synaptogenesis as a result of extensive practice (*24-26, 29*). Similarly, plasticity of the white matter structure of M1 has been correlated with the amount of time devoted to practice, suggesting that myelination may be increased by neural activity in fiber tracts during training (*30, 31*).

In non-human primates, after years of extensive training, M1 neurons were differentially active during the performance of visually guided and internally generated sequential reaching (*32, 33*). Furthermore, the uptake of 2DG, a measure associated with presynaptic activity, was lower in M1 of monkeys that performed highly practiced, internally generated sequences of movements compared with M1 of monkeys that performed visually guided reaching (*33*). Changes in M1 observed at the levels of neuronal activity, functional activation, and gray matter structure in humans and non-human primates may all reflect plasticity underlying the learning of sequential movements through extended, repetitive practice.

Together, these diverse findings converge to the concept that M1 contributes to the learning and maintenance of motor skills such as sequential movements through extensive practice. While these studies demonstrate that activity patterns in M1 change both during the

early phase of learning and after extensive practice, they do not conclusively indicate whether
M1 is critical for sequence learning nor when and how plasticity occurs within M1 during
learning. The change after extensive training could be the outcome of plasticity during the early
phase of the training. Thus, it remains unclear whether M1 undergoes changes throughout all
phases of learning or only during specific stages, and whether M1 plays a critical role in learning
of sequential movements. Moreover, the mechanisms by which M1 contributes to the slow,
gradual improvements in sequence performance are not yet fully understood.

A traditional approach to these questions would be to assess the effect of lesion or
inactivation of M1 on learning performance. However, M1 poses a special challenge to causally
examine when and how plastic changes occur during learning, because M1 is critical for
implementing motor output. A lesion or inactivation of M1 will abolish the motor commands to
the spinal cord that generate muscle activity, so that it would be impossible to evaluate M1's
involvement in learning. Therefore, we targeted information storage in M1 instead of motor
output. We injected an inhibitor for protein synthesis in M1 to interfere with information storage
and, consequently, learning.

Inhibitors for protein synthesis have been widely used to dissect and analyze memory
systems in rodents, such as fear conditioning. Local injection of an inhibitor of protein synthesis
(e.g., anisomycin) into amygdala, hippocampus or motor cortex of rodents disrupted the learning,
maintenance or reconsolidation of memory (*34-44*). These studies indicated that protein
synthesis inhibition can interfere with formation and storage of memory. Building on this
approach, we previously injected a protein synthesis inhibitor into M1 of monkeys after ~100
days of training and found that it successfully disrupted performance of well-trained sequential
movements without affecting the performance of visually guided reaching (*45*). Here, we applied
this approach to examine when and how M1 contributes to learning of sequential movements
during repetitive practice over extensive time in non-human primates.

Results
Learning of sequential reaching task

We trained three monkeys (*Cebus apella*) on two types of reaching tasks (Random and
Repeating tasks, Fig. 1, for details see Materials and methods). In the Repeating task, we
presented visual instruction cues according to a repeating sequence of 3 elements (e.g., 5-3-1-5-
3-1- ..., Fig. 1A, B) 400 msec after monkey's contact of the preceding target. If the monkey made
a correct response during the 400 msec delay, the instruction cue was not presented, and the task
proceeded to the next target in the sequence. With practice, monkeys memorized the sequence
and performed the correct sequence in the Repeating task using predictive responses instead of
using visual instructions to guide their reaching (Fig 1C, D). Thus, the monkeys performed
memory-guided sequential movements during the Repeating task. As a control task, we trained
the monkeys to perform the Radom task in which the visual instruction cues were presented in
pseudo random order. During the Random task, cues were presented 100 ms after the monkey
contacted the target, thereby preventing predictive responses. Thus, the monkeys performed

reaching movements guided by visual cues (Random task). The two tasks were performed
continuously in alternating blocks of 200-500 trials in each daily session. The monkeys predicted
the targets in a Repeating sequence in more than 80% of trials after 50 days of training (Fig. 1C).

We trained monkeys to perform two to three sequences of the Repeating task. We added
new sequences one at a time on a staggered schedule. A new sequence was introduced only after
the previous one had been practiced for more than 50 days (Fig. 1D; see Methods). During each
training session, the monkey performed each sequence and the Random task in alternating blocks
of 200-500 trials. As a result, the training duration for each sequence differed at the time of
experiments. For example, in Fig. 1D, on day 393, training duration for the first sequence was
357 days, 151 days for the second sequence, and 42 days for the third sequence. As shown in
Figure 1D, Response Time (RT) for each sequence decreased independently as training
proceeded, which provided time points when the sequences were in different learning phases. RT
for the Random task (gray dots in Fig. 1D) reached a plateau after approximately 20 days of
training. This staggered design enabled us to examine whether the effect of pharmacological
injection depended on training duration and learning phase, both by comparing the effect on
sequences within a single injection session in addition to a comparison across different injection
sessions.

Protein synthesis inhibition disrupted performance of memory guided sequences at all
stages of learning, while sparing visually-guided reaching.

When the monkey was able to perform the Random task and more than two sequences of
the Repeating task, we made micro injections of the protein synthesis inhibitor anisomycin (100
mg/ml) at sites in M1 where intracortical microstimulation evoked shoulder or elbow movements
(Fig. 2, see Methods). For each session, we analyzed the monkeys' behavior before and after the
injection, separately evaluating each movement in both the Random and Repeating task.

In the injection session shown in Figure 3, monkey N performed three sequences with
different training durations: a short duration (42 days) for sequence '2-3-4' (Fig. 3 A-C), a long
duration (151 days) for sequence '1-2-4' (Fig. 3 D-F) and an extensive duration (357 days) for
sequence '5-3-1' (Fig. 3 G-I). The injection of anisomycin into M1 significantly disrupted
performance of all sequences of the Repeating task. The injections resulted in a significant
increase in the number of incorrect responses (Fig. 3B, E, H; $\chi2$ test, $p<0.0001$) and a significant
decrease in the number of predictive responses, an indicator of sequence learning, (Figure 3C, F,
I; $\chi2$ test, $p < 0.01$) for all moves across all three sequences of the Repeating task. The effect of
anisomycin injection was more pronounced for certain movements within the learned sequences
(increase of error rate: sequence '2-3-4' - move '4-2': 19.25%, '2-3': -5.78%, '3-4': 37.05%;
sequence '1-2-4' - move '4-1': 34.16%, '1-2': 6.89%, '2-4': 9.68%; sequence '5-3-1' - move '3-
1': 0.94%, '5-3': 14.11%, '1-5': 30.48%)(*45*). In contrast, performance of visually guided
movements during the Random task was not significantly disrupted or slightly improved (Fig. 3
A, D, G, $\chi2$ test, p>0.05).

Analysis of reaching endpoints on the screen revealed that the monkey made two types of
error responses during the Repeating task: errors of accuracy and errors in direction. An accuracy
error is a reach performed in the correct direction, but to an endpoint outside of the correct target.
As shown in Figure 3D (sequence 1-2-4), after the anisomycin injection, during the Repeating
task, the animal moved his arm in the correct direction to the left from target "4", but the reach
fell short in 50% of trials (Fig. 3D, bottom row). This type of error response was more frequent
for movements starting from the targets at the edges of the touch monitor, target 1 and 5. A
direction error is a reach performed in the direction opposite to the correct target. For example,
movement "3–4" requires a rightward movement from target "3" to "4" (Fig. 3A). After the
anisomycin injection, during the Repeating task, the animal instead moved his arm to the left
from target "3" and made direction errors on 57% of the trials (Fig 3A, bottom row). Direction
errors were not possible for movements starting from target 1 and 5, as movements in the
opposite direction from these edge targets would fall outside the touch monitor. The direction
errors suggest deficits in selecting the appropriate movement component within a sequence and
in transitioning from one movement to the next. Given the increase of direction errors and
unaffected performance in the Random task, we attribute the observed deficits in the Repeating
task to the injection's effect on memory for sequential movements.

We used the percentage of predictive trials and the Response Time (RT) as additional
indicators to assess the effect of injections on task performance (Fig. 3C, F, I). After the
anisomycin injection, RT for the affected movements in the Repeating task increased. Trials with
longer RT (RT > 150 msec) were categorized as non-predictive responses (see Methods). The
percentage of predictive trials of affected movements significantly decreased for all three
sequences of the Repeating task (Figure 3C, F, I; $\chi2$ *test, p* < 0.01). During the Repeating task,
there were times when the monkey paused in mid-reach and redirected the arm after the visual
cue was presented to make a correct response. This resulted in longer RTs and non-predictive
responses. This behavior suggests that the monkey relied on the visual cue information to make a
correct response, which was possible because the injection did not affect the performance of
visually-guided movements (i.e., Random task). Therefore, the increase of RT after the
anisomycin injection further supports the conclusion that the injection disrupted the memory-
guided sequential movements.

Notably, the effect of anisomycin injection on error rate was more pronounced for the
'Short-term' trained sequence compared to the 'Long-term trained' and 'Extensively' trained
sequences (Fig. 4A; GLM analysis, $p<0.001$; see Methods). Similarly, the decrease in the
percentage of predictive trials and the increase of the Response Time (RT) for the 'Short-term'
trained sequence were significantly larger than those for the 'Extensively trained' sequence (Fig.
4B, C: GLM analysis for predictive trials, two-way ANOVA for RT, $p<0.001$; see Methods for
the definition of predictive trials, RT and statistical analysis). These results suggest that the
neural representation of 'Long-term' and 'Extensively' trained sequence is less susceptible to
interference.

Performance of Repeating sequences recovered after a few days of training.

After the injection, we continued daily training of the monkeys and monitored their performance. Figures 4D-L show performance data of the Repeating task from 4 days before to 7 days after the anisomycin injection session shown in Fig 3. Following the anisomycin injection into M1, the number of correct responses (Fig. 4D, G and J) and the percentage of predictive trials (Fig. 4 E, H and K) decreased across all three sequences of the Repeating task (χ^2 test, $p < 0.05$). Additionally, the average Response Time (RT) increased for all three sequences (Fig. 4F, I and L) (*t-test*, $p<0.05$). However, after 2-3 days of training, the monkey's task performance recovered to the baseline levels for all sequences of the Repeating task (Fig. 4) ($\chi2$ test for performance and predictive, *t-test* for RT, $p<0.05$). No significant differences were observed in recovery periods between sequences with varying training durations. The performance recovery period following the protein synthesis inhibition was consistently short, 2-3 days (48-72 hours), regardless of training duration. This recovery timeline aligns with rodent studies showing that anisomycin injection inhibits protein synthesis in the injected brain area for 48 hours (*40*). These findings suggest that while protein synthesis inhibition disrupted memory traces around the injection site the effect on task performance is reversible with additional practice.

The effect of protein synthesis inhibition is a specific consequence of the induced translational blockade during learning.

Although anisomycin has been widely used in rodents to study memory systems, our studies are the first to apply it in non-human primates to investigate memory systems (*45*). We verified that the observed behavioral effects (Fig. 3) were specifically due to protein synthesis inhibition affecting memory, using three different approaches. First, we inactivated M1 by injecting muscimol, a GABA$_A$ agonist, into the shoulder representation area of M1 and tested its effect on performance of the Random and Repeating tasks (Fig. 5A-C). The inactivation resulted in a significant increase in the number of incorrect responses both in the Random and Repeating tasks (Fig. 5B, C: $\chi2$ test, $p<0.01$). The effect of injection was more pronounced for certain movements, and the same movements were affected in both the Random and Repeating tasks (increase of error rate: Random – move '3-1': 44.6%, '5-3': 1.8%, '1-5': 17.6%; Repeating – '3-1': 19.2%, '5-3': 4.3%, '1-5': -2.8%). In both tasks, the monkey's reach fell short of the correct target after the muscimol injection (accuracy errors, 41% for Random; 32% for Repeating) (Fig. 5A, B). The increase in the number of direction errors after the injection was small in both tasks (Random, pre: 0%; post: 4%; Repeating, pre: 4%; post: 6%). Contrary to the effects of anisomycin, the effects of muscimol injections on task performance were primarily on the accuracy of movements in both the Random and the Repeating tasks (n=2 injection sessions in monkey S). The difference suggests that the deficit in task performance after the anisomycin injection is likely to be a specific consequence of the induced translational blockade, rather than any nonspecific inhibition or dysfunction of neural activity in M1.

Second, to validate the notion that the anisomycin effect is due to its inhibition of protein synthesis, we injected another type of protein synthesis inhibitor, cycloheximide (*39*).

Anisomycin and cycloheximide have distinct mechanisms of action and different side effects
(*39*). The injection of cycloheximide in M1 resulted in a significant increase in the number of
incorrect responses during the Repeating task, but did not have an effect on performance of the
Random task. The effect of injection was more pronounced for certain movements of Repeating
sequences. During the Repeating task, the error rate increased from 3.6 % to 42.5% for the
movement "1-2" for the sequence '1-2-4' ($\chi2$ *test, p<0.05*). On the other hand, the error rate of
the movement '1-2' performed during the Random task did not increase at all (error rate, pre: 0%,
post: 0%). The consistency of the results with the two pharmacological agents suggests that the
effect on behavior is due to the common effect of the agents, inhibition of protein synthesis.

Third, we recorded the activity of neurons before and after the anisomycin injections and
evaluated the effect of injection on neural activity (Fig. 5D-F). Figure 5D shows activity of two
M1 neurons during the Random task before and after injections. The neurons were recorded 1
mm away from the injection site. We analyzed neural data collected during the Radom task
because the anisomycin injection caused a large increase in number of errors during the
Repeating task, as we showed previously (Fig. 3), which prevented the fair comparison of neuron
activity during similar movements. Both neurons had movement selectivity and similar
maximum firing rates during the preferred movements. We recorded the activity of 107 M1
neurons before and after the injections (pre: n=62; post: n=45, recorded within 1-2 mm from the
injection sites, pre: 7 sessions, post: 4 sessions) and obtained the maximum mean firing rate and
a selectivity index (SI) for each neuron (Fig. 5E and F) (see Methods). There were no significant
differences in these measures between pre and post injections (*t-test*, *p>0.05*) (Fig. 5E and F).
The results suggest that the deficit in task performance after the anisomycin injection was likely
to be a specific consequence of its translational blockade, rather than any nonspecific inhibition
or dysfunction of neural activity.

These observations suggest that we successfully manipulated the sequence memory by
injecting a protein synthesis inhibitor. The inhibition of protein synthesis in M1 disrupted the
performance of memory guided sequential movements at all stages of learning without disrupting
motor production. Observations from M1 inactivation using muscimol, recording of neural
activity after the anisomycin injection and the injection of cycloheximide, another type of protein
synthesis inhibitor, all support the conclusion that the deficit in the performance of the Repeating
task after the anisomycin injection was a specific consequence of the induced translational
blockade (*43, 46*), but not of nonspecific neural dysfunction.

The effect of protein synthesis inhibition in M1 on memory guided sequential movements
decreased as learning progressed.

The injection results in Figures 3 and 4 demonstrated that, within an injection session, the
effect of protein synthesis inhibition in M1 varied with the training duration of each Repeating
sequence. To determine whether this effect of protein synthesis inhibition in M1 is consistently
correlated with the training duration, we repeated anisomycin injections after the monkey's
performance had recovered to the pre-injection level and tested its effect on performance of the

tasks (Fig. 6, Monkey N, *n=5*; Monkey R, *n=4*). Injection sessions were separated by more than two weeks of training (see Methods), as performance recovered within 2-3 days of training regardless of training duration (Fig. 4). All injections were placed within the shoulder or elbow area of M1, but at slightly different sites to minimize the tissue damage. After each injection session, the monkeys were trained daily, and recovery of task performance was confirmed. The duration of recovery was consistently 2-3 days across all injection sessions and training durations: 'Short-term' (<100 days), 'Long-term' (100–200 days), and 'Extended' (>200 days) (Fig. 6A, B; see Methods). Overall, the effect of injections on the performance of the Repeating task was consistently observed and significant (χ2 test for error rate and predictive responses, *t-test* for RT, $p < 0.05$). In contrast, no significant decrease of performance was observed in the Random task ($p > 0.05$).

To examine whether the strength of the injection's effect on performance depended on training duration, the most affected movement of each Repeating sequence from the two monkeys was classified into one of three groups based on training duration: 'Short-term' (< 100 days), "Long-term" (100-200 days), and 'Extended' (> 200 days). Magnitude of changes in RT after the injection decreased as training proceeded (Fig. 6C). RTs were significantly different across all groups (see Methods for details; p<0.001 for all pairs) (Fig. 6C). These results indicate that the effect of injections on RTs depended strongly on training duration. On the other hand, changes in performance measured by success/error rates across training duration were not as consistent between two monkeys. Pairwise comparisons of error rates showed significant differences among all three groups for both animals (Monkey N, Short vs Long: *p*=1.678e-21; Short vs Extended: *p*=0.006036; Long vs Extended: *p*=1.756e-31; Monkey R, Short vs Long: *p*=1.942e-13; Short vs Extended: *p*=6.211e-26; Long vs Extended: *p*=1.348e-10; see Methods), but not always in the same direction (Fig. 6A). The inconsistent effects of training duration on error rates suggest that other regions involved in sequence learning, such as premotor areas, may compensate for M1's disfunction differently across learning stages (*5, 53, 67*). For example, inactivation studies show that the pre-SMA contributes only to the very early stage of learning a new sequence structure (*5, 53, 55*). Other possible factors include uncontrolled elements of performance (e.g., motivation, the number of trials in a session, fatigue), the monkey's option to avoid errors by waiting for the visual cue in the Repeating task, differences in reaching distances (e.g., long move '1-5' vs short move '2-3'), variability in the likelihood of direction errors, or the non-linear improvement of success rates as movement chunking evolves across days. Consequently, changes in error rates are less robust, and persist for longer during training. Nevertheless, because protein synthesis inhibition disrupted success rate at all stages of sequence learning, our findings indicate that M1 contributes to learning of sequence structure throughout all stages of learning, potentially in collaboration with premotor areas in a stage-dependent manner.

We further examined whether the effect of injection on performance of the Repeating task is correlated with training duration, eliminating the use of arbitrary grouping based on training duration (See Methods). We found that the magnitude of the injection's effect on the RT for each movement was significantly correlated with the training duration of a sequence (Fig.

6D; *Pearson's correlation coefficient, r = -* 0.49644, *p* = 5.4791e-05). On the other hand, the
magnitude of the injection's effect on the error rate for each movement was not correlated with
the training duration of a sequence (*Pearson's correlation coefficient*, *r* = - 0.089275, *p* = 0.497).
Furthermore, anisomycin injections had no significant effect on MT and performance of the
Random task (i.e., visually guided reaching) (*45*). These findings strongly support M1's role in
speed improvement during extended repetitive practice and suggest that the transition time from
one movement to the next becomes shorter, leading to shorter RT in our task.

The results show that the neural representations of 'Long-term' trained sequence and
'Extended' trained sequence are less susceptible to protein synthesis inhibition, and thus less
vulnerable to interference in terms of speed. The decreased injection effects indicate that the rate
of memory consolidation declines as learning proceeds. Our results suggest that the neural traces
of sequential movements in M1 are gradually stabilized through repetitive practice to support a
high level of performance.

Consolidation during repetitive practice

Analysis of Response Time (RT) provided further insight into this consolidation process
during repetitive practice (Fig. 7; see Methods for RT definition). At the beginning of the post-
injection session, RTs in both the Random and Repeating tasks were comparable to RTs
observed in the pre-injection session (Random: ~360 msec; Repeating: ~ < 0 sec). However, as
the session progressed, RTs increased only in the Repeating task. For example, in the Repeating
sequence '5-3-1' (blue), RTs increased incrementally across blocks, rising from negative values
early in the session (mean of the first 20 trails of movement '3-1': -118.50 ± 62.93 ms) to
approximately 100-300 msec by the end (mean of last 20 trails: 135.00 ± 230.02 ms: *t-test* of
first vs last, *p*<0.001; Fig. 7B). In contrast, RTs in the Random task showed no significant
change within a session either before or after the injection (*t-test*, pre: *p*=0.55; post: *p*=0.37; Fig
7B). Likewise, RTs for the movement '3-1'in the Repeating task did not change significantly
within a training session before the injection (*t-test*, *p*=0.30, Fig. 7B bottom). A significant
increase in RT after the injection was observed for the affected movements of the Repeating task
in eight of nine injections (*t-test*, *p*<0.05). These findings suggest that performing the task
following the anisomycin injection disrupted the neural representation of the Repeating
sequences in M1. This is consistent with findings of rodent studies proposing that synapses
within the memory networks would be initially destabilized by protein degradation upon memory
reactivation during task performance, and then restabilized through protein synthesis to update
the memory network (*35, 36, 41, 42, 44, 47-49*).

We next asked whether exposure to anisomycin alone disrupts the performance of the
Repeating task or whether performing the Repeating task within the drug effect window is
necessary to disrupt performance of the Repeating task (Fig. 7C) (*35, 36*). We injected
anisomycin into M1 of a highly trained monkey and assessed task performance after three days
without training, at which time the injected anisomycin was no longer effective (40). We did not
observe a significant decrease in task performance after three days of inactivity in either the

Random or Repeating tasks (Fig. 8C; χ^2 *test* for error rate and *t-test* for RT, $p > 0.05$, n = 2
injections). The results suggest that exposure of M1 to anisomycin alone did not disrupt the task
performance but performing the Repeating task within the drug exposure window was necessary
to disrupt the neural representations of sequential movements stored in M1. Our data suggest that
anisomycin prevented the synthesis of the proteins needed to consolidate the neural
representations of sequential movements (i.e., synaptic connections) in M1 during sequence
practice as proposed in rodent studies (*35, 36, 41, 42, 44, 47-49*).

Discussion
In this study, we trained monkeys to learn sequential movements and interfered with
information storage in the primary motor cortex (M1) at multiple time points during learning by
injecting a protein synthesis inhibitor, anisomycin, into the arm area of M1. The anisomycin
injection disrupted the performance of memory guided sequential reaching at all stages of
learning without affecting movement production. Notably, the effect of anisomycin injection on
sequence performance was more pronounced for the 'Short-term' trained sequence compared to
the 'Long-term trained' and 'Extended' trained sequences. Further analysis demonstrated that the
magnitude of the injection effect on the RT of each movement was significantly correlated with
the training duration of a sequence. These results suggest that the neural representation of 'Long-
term' and 'Extended' trained sequence is less susceptible to protein synthesis inhibition and thus
less vulnerable to interference. The decreased injection effects indicate that the rate of memory
consolidation declines as learning proceeds and that the neural traces of sequential movements in
M1 are gradually stabilized through repetitive practice to support a high level of performance.

Growing evidence demonstrates that M1's activity changes with the learning of
sequential movements both in the short term (within a training session to several months) (*1-9,*
12-15) and with years of training in humans and non-human primates (*9, 11, 16-33*). The
changes in M1 after years of sequence training were reported in studies comparing musicians to
non-musicians or comparing monkeys performing extensively trained sequential movements to
those performing visually guided movements. However, it remains unclear from this work how
learning-related plasticity occurs in M1, whether M1 is critically involved in sequence learning,
and in which learning stage it may be. Changes of M1 activity after extensive training could be
the outcome of plasticity during the early phase of the training. Our study using continuous long-
term training and the staggered introduction of sequences covered learning stages within
individual monkeys. Our results demonstrated that the protein synthesis inhibition disrupted the
performance of memory-guided sequential reaching tasks at all stages of learning examined. This
suggests that plasticity in M1 occurs at all stages of learning during repetitive practice over an
extended period, continuously supporting learning of sequential movements.

The supplementary motor area (SMA) has long been considered responsible for the
execution and maintenance of learned sequential movements (*5, 9, 50-57*). The SMA neurons
exhibit sequence specific activity (*52, 58-61*). Inactivation or lesion of the SMA disrupted the
performance of memorized sequential movements and inactivation or lesion of the pre-SMA

disrupted the learning of a new sequence structure (*53, 55, 62-66*). In this view, M1 has been
considered to relay the signal from the SMA to generate muscle commands for sequential
movements. Additionally, inactivation of the dorsal premotor cortex (PMd) disrupted the
performance of memorized sequential movements (*67*) and PMd neurons exhibit sequence
preferential activity (*10, 67*). We cannot rule out the possibility that the SMA or PMd input to
M1 (*68-72*) drives learning-related changes in M1. However, our results showed that protein
synthesis inhibition disrupted the sequence performance at all stages of learning tested and
suggest that aspects of learned sequences are consolidated within M1 at all times. M1's
contribution could be to enhance synaptic efficacy, which then leads to improving fluency of
transitions between movements and the execution of individual movement components within a
sequence. Further studies are needed to explore these possibilities.

Studies in rodents using protein synthesis inhibitors suggest that memory may be
dynamically modified or reconsolidated upon retrieval (*41, 42, 44, 73-76*). Specifically, upon
memory retrieval, synapses within the memory network are first destabilized by protein
degradation during reactivation, followed by stabilization to update or strengthen the memory
through protein synthesis (*41, 42, 44, 75*). Supporting this view, the inhibition of protein
degradation by *clasto*-lactacystin-*β*-lactone (*β*lac) has been shown to prevent anisomycin-
induced memory impairment (*77*). Moreover, human studies using TMS and functional imaging
suggest that memory reconsolidation may occur in humans as well (*78-80*). Several factors,
including the age and strength of a memory (*47, 73, 81*), the novelty of the experience (*82-84*),
and duration of memory reactivation necessary for reconsolidation to occur (*47, 48*) are proposed
as boundary conditions, which are considered to influence whether memory reconsolidation
occurs.

In our study, we observed that the performance of the Repeating task was disrupted with
a delayed onset of deficits after the anisomycin injection. The performance deficits may result
from interference with the reconsolidation occurring during repetitive practice. The delayed
onset of deficits (Fig. 7) may be linked to the duration of memory reactivation necessary for
reconsolidation to occur (*47, 48*), specifically the duration for protein degradation to destabilize
the memory network for reconsolidation. In fact, when we tested the injection effect outside of
the drug's effective window, three days after the injection without training, there was no
significant effect on performance (Fig. 7C). The results suggest that protein synthesis inhibition
alone may not be sufficient to disrupt memory-guided sequential movements. Consistent with
rodent studies, memory reactivation during practice may be required to reveal the
reconsolidation impairment due to protein synthesis inhibition.

Previous studies have shown that memory could be updated through reconsolidation
when new information needs to be integrated (*85-87*), and that both the age and strength of a
memory influence whether reconsolidation occurs or not (*47, 48*). Specifically, older and
stronger memories (i.e., those reinforced through extended training) tend to be less sensitive to
protein synthesis inhibition. In our study, the effect of protein synthesis inhibition on memory-
guided sequence performance were smaller during the 'Long-term and 'Extended' training
phases in which RT variability was smaller (Fig. 1C). In our task, performance fluctuations due

to movement variability may represent a form of "novelty" that serves as a boundary condition
for reconsolidation. Our findings suggest that the memory network in M1 may be destabilized
prior to update (or reconsolidation) at every practice rather than accumulating changes purely
incrementally. This is consistent with the 'lingering consolidation' hypothesis, in which 'the
reactivation and reconsolidation cycle progressively stabilizes a memory' (*42*). Furthermore, this
would align with the gradual and continuous improvement of motor skill performance observed
during extensive, repetitive practice. Recent studies in rodents have suggested that the motor
cortex of rodents may not be essential for maintaining certain types of motor skills after "long-
term" training (*88-90*). In contrast, our results demonstrate that protein synthesis inhibition in
M1 of monkeys significantly impaired sequence performance even after more than 300 days of
training (Figs. 3, 4 and 6). These differences may reflect several factors, including differences in
whether the animals' performance reached the boundary conditions of memory updating, the
higher complexity of the sequential task used in our study, and species-specific anatomical
differences (*72*). Further investigation will be needed to elucidate these possibilities.

Two-photon imaging studies offer insights into the possible mechanisms underlying
memory-update in the motor cortex. In rodents, two photon imaging of the motor cortex during
motor skill learning has shown the formation and growth of new dendritic spines and the
elimination of old spines (*91, 92*). Additionally, two-photon imaging of cortical slices and
electron microscopic studies revealed that protein synthesis is required for long-lasting synaptic
plasticity (*43*) and spine-head enlargement and growth during learning (*93, 94*). The inhibition of
protein synthesis resulted in a significant reduction in synapse number and synapse size in motor
cortex of rodents in vivo (*39*). These observations suggest that, during motor skill training,
unnecessary spines are eliminated through protein degradation, while the formation and growth
of new spines are supported by protein synthesis which leads to enhanced neural efficacy. Thus,
deficits in spine growth and formation may be associated with impairments in memory updates
(i.e., consolidation), as observed in behavioral tests. Together with our findings, these results
suggest that the neural traces for motor skills are continuously updated during repetitive practice
through protein synthesis. The delayed onset of performance deficit, as shown in Figure 7, may
reflect the time required for protein degradation to eliminate a sufficient number of spines to
impact task performance (*41, 42, 44*).

Our study suggests that motor skill is gradually consolidated in M1 every time a subject
practice and that the rate of consolidation (i.e., memory strengthening) becomes smaller as the
learning proceeds. The lingering consolidation process implies that consolidation reaches a
theoretical limit when the memory trace has become old and stable. Nevertheless, our results
suggest that lingering consolidation still occurs in M1 in our monkeys after 377 sessions of
practice on simple sequences. M1 may well make a steady contribution to the development of
expertise, serving as the neural substrate underlying "Practice makes perfect".

Materials and Methods

    Behavioral task: We trained three monkeys (*Cebus apella*, male or female, weighing 1~5 kg; > 2 years old) on the Random and Repeating tasks (Fig. 1) (*32, 33, 45, 67, 95, 96*). Each monkey was required to make sequential reaching movements to targets on a touch sensitive monitor with their right arms. In the Random task, the reaching movements were guided by visual targets displayed on a touch sensitive monitor in a pseudo-random order. Contact of the correct target triggered display of the next target after a 100 ms delay. After the monkey became proficient in performing the Random task (~50 days of practice), we introduced the Repeating task. In the Repeating task, new targets were presented according to a repeating sequence of 3 elements (e.g., 5-3-1-5-3-1- ..., Fig. 1A, B) 400 msec after contact of the preceding target. If the monkey made a correct response during the delay, the target is not shown and the task increments to the next target in the sequence. With practice, a monkey memorized and performed the sequence of the Repeating task using predictive responses instead of visual cues to direct their movements in more than 80% of trials. The two tasks were performed continuously in alternating blocks of 200-500 trials. The monkey received a liquid reward after every 4-5 correct responses. After the monkey became proficient in the performance of the first sequence (Fig. 1C), additional sequences (e.g., 1-2-4-1-2-4...) were introduced. The new sequences were introduced in a staggered design with 50-200 days intervals (Fig. 1D). This training design enabled us to compare the effect of pharmacological injection between the 'Long-term' trained and 'Short-term' trained sequences in a single injection session. Each monkey learned two to three sequences (Monkey N: '5-3-1', '1-2-4', '2-3-4'; Monkey R: '1-2-4', '2-3-4'; Monkey S: '5-3-1', '1-2-4'). When the monkey frequently stopped working during a training session, the session was omitted and the behavioral data were excluded from further analysis because an insufficient number of trials and conditions were collected. Control of the behavioral task and collection of movement data including videos of task performance were performed by PC computers running TEMPO software (Reflective Computing, Olympia, WA) and Matlab. Touch screen signals, task related events and neural activity were collected on-line at 1 kHz.

    Behavioral estimates of learning phases: We used "Response Time" (RT) as the principal measure of sequence learning. We defined RT during the Random task as the time between the presentation of a new target and contact of that target. We defined RT during the Repeating task as the time between two targets touches minus the delay time, 400 msec (*33, 45, 67, 96*). We subtracted 400 ms to account for the delay in the cue presentation. This could result in a negative RT if the monkey moved quickly to the next target in the sequence before the presentation of a cue. RTs less than 150 ms were considered to be predictive. RTs less than 150 ms were chosen as a conservative cut-off for predictive responses as it is too fast for a simple reaction time to the visual cue. During the learning, an animal's RT became shorter with practice (Fig. 7c, d). A negative RT indicates that the animal was internally generating the response before the visual cue was presented. After about 50 days of training on the Repeating sequence, the monkeys made predictive responses (RT < 150 ms) without using visual cues in more than 80% of the trials (Fig. 1C, D).

Surgical Procedures: All experimental procedures were conducted according to NIH guidelines and were approved by the IACUC of the University of Pittsburgh. We implanted a head restraint device, along with a MR compatible chamber for micro-injections and neural recording, on an animal's skull using small screws and dental acrylic (*45, 67*). All surgical procedures were performed under general anesthesia using aseptic techniques. Anesthesia was induced with ketamine (10 mg/kg, IM) and maintained to surgical levels with 1-2.5% isoflurane. The animal received fluids throughout the surgical procedure (6-10 cc/hr, IV) and glycopyrolate (0.01 mg/kg, IM) to reduce secretions. Body temperature was maintained at 37-38 °C with a heating pad. We monitored body temperature, breathing, heart rate, end-tidal CO_2, and O_2 saturation. After the surgery, the animal received broad spectrum antibiotics (ceftriaxone, 75 mg/kg/day, IM, ~5 days) and analgesics (buprenorphine, 0.01 mg/kg, IM, ~3 days). The chamber's placement over M1 was verified using structural MR images taken prior to and after the surgery (Fig. 2B). MR procedures for pretreatment, anesthesia, monitoring of vital signs and post-anesthesia monitoring were identical to surgical procedures, except that analgesic and antibiotic were not be given. After more than 2 weeks of recovery, we accustomed the animal to perform the tasks with its head restrained. When task performance returned to the pre-surgical level, we performed a craniotomy to expose the dura matter overlying M1 in the chamber using the same anesthetic techniques and safeguards used in the implant surgery. Animals were given dexamethasone (0.5 mg/kg divided into 3 doses, IM) on the day of the craniotomy surgery to prevent brain swelling.

Intracortical stimulation: We used intracortical microstimulation to identify the body part represented at each site in M1 and to physiologically define the border between M1 and the PMd (*45, 67, 71*). We used glass-coated micro-electrodes (0.6-2 MΩ at 1 kHz) to deliver intracortical microstimuli (*45, 67, 71*). A constant-current stimulator was used to deliver cathodal pulses (10–20 pulses, 0.2 ms duration, 333 Hz, 1- 40 µA) at a depth of 1500 µm below the cortical surface. Stimulus intensity was measured with a current monitor (Ion Physics). The motor response evoked by stimulation was determined by visual observation and muscle palpation. The response threshold was defined as the lowest stimulus intensity necessary to evoke a response on more than 50% of the trials (*71, 97, 98*). We systematically mapped M1 with micro-electrode penetrations spaced ~1.0 mm apart (except to avoid blood vessels) (Fig. 2).

Micro-injections: We made micro-injections of pharmacological agents (anisomycin, cycloheximide or muscimol) at selected sites in the forelimb area of M1 of a trained monkey (*45, 67*). We prepared solutions of anisomycin (100 µg/µl in ACSF, pH 7.2-7.4), cycloheximide (40 µg/µl in 20% ethanol/ACSF) and muscimol (5 mg/ml in saline) from commercially available powders (Sigma-Aldrich, MO). We injected the test solution at 1.5 mm below the cortical surface using a 30-gauge cannula connected to a 10 µl Hamilton syringe by applying slow pressure (e.g., over 10 minutes). Injection sites were placed more than 2 mm away from the border between M1 and PMd identified by microstimulation (Fig. 2). Previous studies using monkeys have shown that injection of 3 µl of muscimol into the cortex inhibited neural activity within a diameter of approximately 2-3 mm (*66*). Given that M1 in Cebus monkeys is more than 5 mm in width (Fig. 2), the effect of an injection of less than 3 µl of a pharmacological agent 2

mm away from the border was expected to remain confined within M1 (*45*). The cannula was left in place for more than 5 min to allow diffusion of the solution and prevent its reflux. For each anisomycin injection session, we injected a total of 5 µl of anisomycin solution into M1 of two monkeys. The solution was divided into two aliquots (2.6 µl and 2.4 µl) and injected at sites within M1 where microstimulation elicited elbow or shoulder movements. The injection volume was divided to minimize potential tissue damage and to ensure broader coverage of the cortical regions representing the elbow and shoulder. Task performance was assessed 20–24 hours after the injection and daily thereafter. Measurement of performance on days prior to an injection and/or on trial blocks preceding an injection was used as the baseline for comparison with post-injection performance. We repeated the anisomycin injections with behavioral testing five times in monkey N and four times in monkey R to test its effect on learning. To verify that the effect on behavior is due to the common effect of the agents (i.e., protein synthesis inhibition), we injected 3 µl of cycloheximide (40 µg/µl) at a site within M1 where microstimulation elicited elbow or shoulder movements. Cycloheximide blocks protein synthesis by a different mechanism than anisomycin. The effect of cycloheximide injection on tasks was tested 20-24 hours after the injection. Measurement of performance on a day prior to an injection and/or on trial blocks preceding an injection was used as the baseline for comparison with post-injection performance. Muscimol, a GABA$_A$ agonist, (1-3 µl) was injected at a site within M1 of the monkey S where microstimulation elicited elbow or shoulder movements immediately after the collection of the baseline behavioral data and the effect of injection on task performance was tested 20 min after the injection to allow diffusion of the chemical in the brain tissue (*n*=2). After each injection session, the task performance was monitored every day until it recovered to the baseline to track the emergence of injection effects and to assess recovery from the drug's effects. Injections were separated with more than 7 days and 5 training sessions. We examined whether the injection of the pharmacological agent impaired performance of the Random and Repeating tasks. During a post-injection test session, blocks of Random and Repeating trials alternated at frequent intervals to sample the animal's performance evenly as the effect of the test substance emerged and intensified.

Analysis of performance: For every trial of each task session, we recorded various task parameters and measures of performance. The effect of an injection was assessed by examining the following: percentage of correct responses, types of incorrect responses, percentage of predictive responses (defined as RT < 150 msec), Response Time (RT) and Movement Time (MT) (*45, 67*). The errors recorded were: no hit, wrong target hit, background hit, or a corrective response (a correct response that immediately follows an error). MT is defined as the interval between the release of contact from one target to touch of the next target. We defined RT during the Random task as the time between the presentation of a new target and contact of that target. We defined RT during the Repeating task as the time between targets touches minus the delay time, 400 msec (see the section of Behavioral estimates of learning phases above)(*33, 45, 67, 96*). RT less than 150 ms was chosen as a cut-off for predictive responses as it is too fast for a simple reaction time to the visual cue. A wrong target hit during the Repeating task was categorized as two types: *accuracy errors* and *direction errors*. An *accuracy error* is defined as a reach performed in the correct direction, but to an endpoint outside of the correct target. This type of

error suggests a deficit in motor production. A *direction error* was defined as a reach performed in the direction opposite to the correct target. This type of error suggests a deficit in selecting the movement component in the sequence. Direction errors could only occur for movements starting from targets on the center of the touch monitor (target 2, 3, 4), since movements in the opposite direction from targets at the edge of the touch monitor would fall outside the touch monitor. For statistical analysis, both types of errors were included in the calculation of performance error. An increase in RT and decrease in the number of predictive responses suggest an increase in the time for movement selection. Corrective responses were removed from analysis because the target is predictable after an error. We also monitored movement kinematics during the task using high speed video recording (100 Hz, Basler Inc., PA) in the frontal plane. For statistical analysis, we used χ^2 tests with Holm–Bonferroni's correction to examine the significance of changes in success rate and predictive responses in a session. We used *t-tests* with Holm–Bonferroni's correction to examine MT and RT changes. To test the reproducibility of the task performance results, we repeated the injections five times in monkey N and four times in monkey R. In monkey R, we performed three additional injections to examine the effect of anisomycin injection on neural activity (see the Electrophysiology section of the Methods). To compare the strength of the injection effect between the sequences in the different learning phases, we fitted a generalized linear model (GLM), using either performance or predictive responses as the variable and the learning phase as a predictor for analysis shown in Fig. 4A and B. A binomial distribution with a logit link function was applied. For analysis in Fig. 4C, we used two-way ANOVA with RT and the learning phases as factors. To compare the strength of the injection effect between the sequences in the different learning phases as a population, the most affected movement of each Repeating sequence from two monkeys was classified into one of three groups based on the training duration: Short-term (less than 100 days, large RT decrease and large RT variability), Long-term (100- 200 days, smaller RT decrease and large RT variability), Extended (more than 200 days, small RT decrease and small RT variability) (Fig. 6A-C). These groupings were defined according to the learning curves shown in Fig. 1. In Fig 6C, we compared changes in performance measures of the most affected movements of each sequence across the Short, Long, and Extended groups using pairwise statistical tests (5 injections for monkey N and 4 injections for monkey R). RT was analyzed using a linear mixed-effects models with Condition (pre vs post) and Group (Short, Long, Extended) as fixed effects and InjectionID as a random effect to account for repeated measurements within each injection session. Pairwise comparisons of the Condition effect (post-pre) between groups were performed by testing linear contrasts on the interaction coefficients. Performance (correct/error) was analyzed using a logistic generalized linear mixed-effects model (logit GLME) with Condition (pre vs post) and Group (Short, Long, Extended) as fixed effects and InjectionID as a random effect. Pairwise comparisons for the Condition effect between groups were performed same as the RT analysis. For the *Pearson's correlation coefficient* analysis between the training duration and the effect of the injection on RT (Fig. 6D), data from all the movements performed in the Repeating sequences were included. For the RT analysis examining changes within a training session (Fig. 7A, B), the first 20 trials and the last 20 trials of each movement in each session were compared using a *t-test*. Matlab (Mathworks) was used for analysis.

Electrophysiological recording and analysis of neural activity: To evaluate the effect of pharmacological agents on neural activity, we recorded extracellular activity of single neurons in M1 using glass-coated Elgiloy electrodes (0.6-1.5 MΩ). We recorded neural activity before and after the injection of anisomycin while the monkey S performed the tasks. As shown in Fig. 3, an anisomycin injection disrupted performance of the Repeating task on up to 80% of trials. Therefore, to evaluate the effect of injection on neural activity, we limited analysis on the data collected during the Random task. Neural data after the anisomycin injection were obtained during four injection sessions. Of these, performance data of three injection sessions were excluded from behavioral analysis because of insufficient number of trials in the Repeating task. The sample of neurons was taken 20-24 hours after the injection within 1-2 mm of the injection site from regions of M1 where intracortical stimulation evoked shoulder, elbow or wrist movements at thresholds lower than 25 µA. For each neuron, we used maximum firing rate and a selectivity index (SI) to assess the effect of injections. For each Random move, we measured the mean firing rate of a neuron in a 200 msec interval centered on target contact or release. We used this value to calculate a SI for each neuron: SI = [Max - Min]/ [Max + Min]. For statistical analysis, we used *t-test* to examine changes in SI and max firing rate between the group of M1 neurons recorded before and after the injections.

References

1. H. Shibasaki *et al.*, Both primary motor cortex and supplementary motor area play an
important role in complex finger movement. *Brain : a journal of neurology* **116** (Pt 6),
1387-1398 (1993).
2. S. T. Grafton, R. P. Woods, M. Tyszka, Functional imaging of procedural motor learning:
Relating cerebral blood flow with individual subject performance. *Human brain mapping*
1, 221-234 (1994).
3. S. T. Grafton, E. Hazeltine, R. Ivry, Functional mapping of sequence learning in normal
humans. *J Cogn Neurosci* **7**, 497-510 (1995).
4. A. Karni *et al.*, Functional MRI evidence for adult motor cortex plasticity during motor
skill learning. *Nature* **377**, 155-158 (1995).
5. O. Hikosaka, K. Nakamura, K. Sakai, H. Nakahara, Central mechanisms of motor skill
learning. *Current opinion in neurobiology* **12**, 217-222 (2002).
6. L. G. Ungerleider, J. Doyon, A. Karni, Imaging brain plasticity during motor skill
learning. *Neurobiology of learning and memory* **78**, 553-564 (2002).
7. J. Doyon, V. Penhune, L. G. Ungerleider, Distinct contribution of the cortico-striatal and
cortico-cerebellar systems to motor skill learning. *Neuropsychologia* **41**, 252-262 (2003).
8. J. Doyon, H. Benali, Reorganization and plasticity in the adult brain during learning of
motor skills. *Current opinion in neurobiology* **15**, 161-167 (2005).
9. E. Dayan, L. G. Cohen, Neuroplasticity subserving motor skill learning. *Neuron* **72**, 443-
454 (2011).
10. R. M. Hardwick, C. Rottschy, R. C. Miall, S. B. Eickhoff, A quantitative meta-analysis
and review of motor learning in the human brain. *NeuroImage* **67**, 283-297 (2013).
11. M. Ohbayashi, The roles of the cortical motor areas in sequential movements. *Frontiers*
in behavioral neuroscience **15**, 640659 (2021).
12. A. Floyer-Lea, P. M. Matthews, Changing brain networks for visuomotor control with
increased movement automaticity. *Journal of neurophysiology* **92**, 2405-2412 (2004).
13. S. Lehericy *et al.*, Distinct basal ganglia territories are engaged in early and advanced
motor sequence learning. *Proceedings of the National Academy of Sciences of the United*
States of America **102**, 12566-12571 (2005).
14. N. G. Hatsopoulos, L. Paninski, J. P. Donoghue, Sequential movement representations
based on correlated neuronal activity. *Experimental brain research* **149**, 478-486 (2003).
15. X. Lu, J. Ashe, Anticipatory activity in primary motor cortex codes memorized
movement sequences. *Neuron* **45**, 967-973 (2005).
16. M. Hund-Georgiadis, D. Y. von Cramon, Motor-learning-related changes in piano
players and non-musicians revealed by functional magnetic-resonance signals.
Experimental brain research **125**, 417-425 (1999).
17. T. Krings *et al.*, Cortical activation patterns during complex motor tasks in piano players
and control subjects. A functional magnetic resonance imaging study. *Neuroscience*
letters **278**, 189-193 (2000).
18. L. Jancke, N. J. Shah, M. Peters, Cortical activations in primary and secondary motor
areas for complex bimanual movements in professional pianists. *Brain Res Cogn Brain*
Res **10**, 177-183 (2000).
19. B. Haslinger *et al.*, Reduced recruitment of motor association areas during bimanual
coordination in concert pianists. *Human brain mapping* **22**, 206-215 (2004).

20.   I. Meister *et al.*, Effects of long-term practice and task complexity in musicians and
nonmusicians performing simple and complex motor tasks: implications for cortical
motor organization. *Human brain mapping* **25**, 345-352 (2005).
21.   K. Amunts *et al.*, Motor cortex and hand motor skills: structural compliance in the human
brain. *Human brain mapping* **5**, 206-215 (1997).
22.   C. Gaser, G. Schlaug, Brain structures differ between musicians and non-musicians. *The
Journal of neuroscience* **23**, 9240-9245 (2003).
23.   B. Draganski *et al.*, Neuroplasticity: changes in grey matter induced by training. *Nature*
427, 311-312 (2004).
24.   B. Draganski, A. May, Training-induced structural changes in the adult human brain.
Behavioural brain research **192**, 137-142 (2008).
25.   S. C. Herholz, R. J. Zatorre, Musical training as a framework for brain plasticity:
behavior, function, and structure. *Neuron* **76**, 486-502 (2012).
26.   R. J. Zatorre, R. D. Fields, H. Johansen-Berg, Plasticity in gray and white: neuroimaging
changes in brain structure during learning. *Nature neuroscience* **15**, 528-536 (2012).
27.   E. Wenger, C. Brozzoli, U. Lindenberger, M. Lovden, Expansion and renormalization of
human brain structure during skill acquisition. *Trends in cognitive sciences* **21**, 930-939
(2017).
28.   C. Sampaio-Baptista, H. Johansen-Berg, White matter plasticity in the adult brain.
Neuron **96**, 1239-1251 (2017).
29.   B. J. Anderson *et al.*, Glial hypertrophy is associated with synaptogenesis following
motor-skill learning, but not with angiogenesis following exercise. *Glia* **11**, 73-80 (1994).
30.   S. L. Bengtsson *et al.*, Extensive piano practicing has regionally specific effects on white
matter development. *Nature neuroscience* **8**, 1148-1150 (2005).
31.   Y. Han *et al.*, Gray matter density and white matter integrity in pianists' brain: a
combined structural and diffusion tensor MRI study. *Neuroscience letters* **459**, 3-6 (2009).
32.   Y. Matsuzaka, N. Picard, P. L. Strick, Skill representation in the primary motor cortex
after long-term practice. *Journal of neurophysiology* **97**, 1819-1832 (2007).
33.   N. Picard, Y. Matsuzaka, P. L. Strick, Extended practice of a motor skill is associated
with reduced metabolic activity in M1. *Nature neuroscience* **16**, 1340-1347 (2013).
34.   H. P. Davis, L. R. Squire, Protein synthesis and memory: a review. *Psychological bulletin*
96, 518-559 (1984).
35.   K. Nader, G. E. Schafe, J. E. Le Doux, Fear memories require protein synthesis in the
amygdala for reconsolidation after retrieval. *Nature* **406**, 722-726 (2000).
36.   K. Nader, G. E. Schafe, J. E. LeDoux, The labile nature of consolidation theory. *Nature
reviews. Neuroscience* **1**, 216-219 (2000).
37.   E. R. Kandel, The molecular biology of memory storage: a dialogue between genes and
synapses. *Science* **294**, 1030-1038 (2001).
38.   K. Nader, Memory traces unbound. *Trends in neurosciences* **26**, 65-72 (2003).
39.   J. A. Kleim *et al.*, Functional organization of adult motor cortex is dependent upon
continued protein synthesis. *Neuron* **40**, 167-176 (2003).
40.   A. R. Luft, M. M. Buitrago, T. Ringer, J. Dichgans, J. B. Schulz, Motor skill learning
depends on protein synthesis in motor cortex after training. *The Journal of neuroscience*
24, 6515-6520 (2004).
41.   Y. Dudai, The neurobiology of consolidations, or, how stable is the engram? *Annual
review of psychology* **55**, 51-86 (2004).

42. Y. Dudai, M. Eisenberg, Rites of passage of the engram: reconsolidation and the
lingering consolidation hypothesis. *Neuron* **44**, 93-100 (2004).
43. R. J. Kelleher, 3rd, A. Govindarajan, S. Tonegawa, Translational regulatory mechanisms
in persistent forms of synaptic plasticity. *Neuron* **44**, 59-73 (2004).
44. J. W. Rudy, Destroying memories to strengthen them. *Nature neuroscience* **11**, 1241-
1242 (2008).
45. M. Ohbayashi, Inhibition of protein synthesis in M1 of monkeys disrupts performance of
sequential movements guided by memory. *Elife* **9**, (2020).
46. D. J. Linden, A protein synthesis-dependent late phase of cerebellar long-term depression.
Neuron **17**, 483-490 (1996).
47. A. Suzuki *et al.*, Memory reconsolidation and extinction have distinct temporal and
biochemical signatures. *The Journal of neuroscience* **24**, 4787-4795 (2004).
48. N. Mamiya *et al.*, Brain region-specific gene expression activation required for
reconsolidation and extinction of contextual fear memory. *The Journal of neuroscience*
29, 402-413 (2009).
49. J. L. Lee, Memory reconsolidation mediates the strengthening of memories by additional
learning. *Nature neuroscience* **11**, 1264-1266 (2008).
50. P. E. Roland, B. Larsen, N. A. Lassen, E. Skinhoj, Supplementary motor area and other
cortical areas in organization of voluntary movements in man. *Journal of*
neurophysiology **43**, 118-136 (1980).
51. C. Gerloff, B. Corwell, R. Chen, M. Hallett, L. G. Cohen, Stimulation over the human
supplementary motor area interferes with the organization of future elements in complex
motor sequences. *Brain : a journal of neurology* **120 (Pt 9)**, 1587-1602 (1997).
52. K. Nakamura, K. Sakai, O. Hikosaka, Neuronal activity in medial frontal cortex during
learning of sequential procedures. *Journal of neurophysiology* **80**, 2671-2687 (1998).
53. K. Shima, J. Tanji, Both supplementary and presupplementary motor areas are crucial for
the temporal organization of multiple movements. *Journal of neurophysiology* **80**, 3247-
3260 (1998).
54. N. Picard, P. L. Strick, Imaging the premotor areas. *Current opinion in neurobiology* **11**,
663-672 (2001).
55. J. Tanji, Sequential organization of multiple movements: involvement of cortical motor
areas. *Annual review of neuroscience* **24**, 631-651 (2001).
56. P. Nachev, C. Kennard, M. Husain, Functional role of the supplementary and pre-
supplementary motor areas. *Nature reviews. Neuroscience* **9**, 856-869 (2008).
57. G. Cona, C. Semenza, Supplementary motor area as key structure for domain-general
sequence processing: A unified account. *Neuroscience and biobehavioral reviews* **72**, 28-
42 (2017).
58. J. Tanji, K. Shima, H. Mushiake, Multiple cortical motor areas and temporal sequencing
of movements. *Brain Res Cogn Brain Res* **5**, 117-122 (1996).
59. W. T. Clower, G. E. Alexander, Movement sequence-related activity reflecting numerical
order of components in supplementary and presupplementary motor areas. *Journal of*
neurophysiology **80**, 1562-1566 (1998).
60. K. Shima, J. Tanji, Neuronal activity in the supplementary and presupplementary motor
areas for temporal organization of multiple movements. *Journal of neurophysiology* **84**,
2148-2160 (2000).

61.   D. Lee, S. Quessy, Activity in the supplementary motor area related to learning and
performance during a sequential visuomotor task. *Journal of neurophysiology* **89**, 1039-
1056 (2003).
62.   C. Brinkman, Supplementary motor area of the monkey's cerebral cortex: short- and
long-term deficits after unilateral ablation and the effects of subsequent callosal section.
The Journal of neuroscience **4**, 918-929 (1984).
63.   G. Goldberg, Supplementary motor area structure and function: Review and hypotheses.
Behavioral and Brain Sciences **8**, 567-588 (1985).
64.   J. P. Dick, R. Benecke, J. C. Rothwell, B. L. Day, C. D. Marsden, Simple and complex
movements in a patient with infarction of the right supplementary motor area. *Mov*
Disord **1**, 255-266 (1986).
65.   U. Halsband, N. Ito, J. Tanji, H. J. Freund, The role of premotor cortex and the
supplementary motor area in the temporal control of movement in man. *Brain : a journal*
of neurology **116 (Pt 1)**, 243-266 (1993).
66.   K. Nakamura, K. Sakai, O. Hikosaka, Effects of local inactivation of monkey medial
frontal cortex in learning of sequential procedures. *Journal of neurophysiology* **82**, 1063-
1068 (1999).
67.   M. Ohbayashi, N. Picard, P. L. Strick, Inactivation of the dorsal premotor area disrupts
internally generated, but not visually guided, sequential movements. *The Journal of*
neuroscience **36**, 1971-1976 (2016).
68.   K. F. Muakkassa, P. L. Strick, Frontal lobe inputs to primate motor cortex: evidence for
four somatotopically organized 'premotor' areas. *Brain research* **177**, 176-182 (1979).
69.   S. Ghosh, C. Brinkman, R. Porter, A quantitative study of the distribution of neurons
projecting to the precentral motor cortex in the monkey (M. fascicularis). *J Comp Neurol*
259, 424-444 (1987).
70.   R. P. Dum, P. L. Strick, Motor areas in the frontal lobe of the primate. *Physiol Behav* **77**,
677-682 (2002).
71.   R. P. Dum, P. L. Strick, Frontal lobe inputs to the digit representations of the motor areas
on the lateral surface of the hemisphere. *The Journal of neuroscience* **25**, 1375-1386
(2005).
72.   P. L. Strick, R. P. Dum, J. A. Rathelot, The cortical motor areas and the emergence of
motor skills: A Neuroanatomical Perspective. *Annual review of neuroscience* **44**, 425-447
(2021).
73.   M. Eisenberg, T. Kobilo, D. E. Berman, Y. Dudai, Stability of retrieved memory: inverse
correlation with trace dominance. *Science* **301**, 1102-1104 (2003).
74.   S. H. Wang, R. G. Morris, Hippocampal-neocortical interactions in memory formation,
consolidation, and reconsolidation. *Annual review of psychology* **61**, 49-79, C41-44
(2010).
75.   Y. Dudai, The restless engram: consolidations never end. *Annual review of neuroscience*
35, 227-247 (2012).
76.   J. L. C. Lee, K. Nader, D. Schiller, An Update on Memory Reconsolidation Updating.
Trends in cognitive sciences **21**, 531-545 (2017).
77.   S. H. Lee *et al.*, Synaptic protein degradation underlies destabilization of retrieved fear
memory. *Science* **319**, 1253-1256 (2008).
78.   N. Censor, E. R. Buch, K. Nader, L. G. Cohen, Altered human memory modification in
the presence of normal consolidation. *Cerebral cortex* **26**, 3828-3837 (2016).

79. M. B. Silva, A. B. Soares, Reconsolidation of human motor memory: From boundary
conditions to behavioral interventions-How far are we from clinical applications?
Behavioural brain research **353**, 83-90 (2018).
80. J. Herszage, H. Sharon, N. Censor, Reactivation-induced motor skill learning.
Proceedings of the National Academy of Sciences of the United States of America **118**,
(2021).
81. M. H. Milekic, C. M. Alberini, Temporally graded requirement for protein synthesis
following memory reactivation. *Neuron* **36**, 521-525 (2002).
82. M. E. Pedreira, L. M. Perez-Cuesta, H. Maldonado, Mismatch between what is expected
and what actually occurs triggers memory reconsolidation or extinction. *Learning &*
memory **11**, 579-585 (2004).
83. S. Tronel, M. H. Milekic, C. M. Alberini, Linking new information to a reactivated
memory requires consolidation and not reconsolidation mechanisms. *PLoS biology* **3**,
e293 (2005).
84. J. Debiec, V. Doyere, K. Nader, J. E. Ledoux, Directly reactivated, but not indirectly
reactivated, memories undergo reconsolidation in the amygdala. *Proceedings of the*
National Academy of Sciences of the United States of America **103**, 3428-3433 (2006).
85. C. J. Rodriguez-Ortiz, V. De la Cruz, R. Gutierrez, F. Bermudez-Rattoni, Protein
synthesis underlies post-retrieval memory consolidation to a restricted degree only when
updated information is obtained. *Learning & memory* **12**, 533-537 (2005).
86. R. G. Morris *et al.*, Memory reconsolidation: sensitivity of spatial memory to inhibition
of protein synthesis in dorsal hippocampus during encoding and retrieval. *Neuron* **50**,
479-489 (2006).
87. C. J. Rodriguez-Ortiz, P. Garcia-DeLaTorre, E. Benavidez, M. A. Ballesteros, F.
Bermudez-Rattoni, Intrahippocampal anisomycin infusions disrupt previously
consolidated spatial memory only when memory is updated. *Neurobiology of learning*
and memory **89**, 352-359 (2008).
88. E. J. Hwang *et al.*, Disengagement of motor cortex from movement control during long-
term learning. *Sci Adv* **5**, eaay0001 (2019).
89. E. J. Hwang, J. E. Dahlen, M. Mukundan, T. Komiyama, Disengagement of Motor
Cortex during Long-Term Learning Tracks the Performance Level of Learned
Movements. *The Journal of neuroscience* **41**, 7029-7047 (2021).
90. K. G. C. Mizes, J. Lindsey, G. S. Escola, B. P. Olveczky, The role of motor cortex in
motor sequence execution depends on demands for flexibility. *Nature neuroscience* **27**,
2466-2475 (2024).
91. T. Xu *et al.*, Rapid formation and selective stabilization of synapses for enduring motor
memories. *Nature* **462**, 915-919 (2009).
92. G. Yang, F. Pan, W. B. Gan, Stably maintained dendritic spines are associated with
lifelong memories. *Nature* **462**, 920-924 (2009).
93. O. Steward, E. M. Schuman, Compartmentalized synthesis and degradation of proteins in
neurons. *Neuron* **40**, 347-359 (2003).
94. J. Tanaka *et al.*, Protein synthesis and neurotrophin-dependent structural plasticity of
single dendritic spines. *Science* **319**, 1683-1687 (2008).
95. W. B. Verwey, T. Eikelboom, Evidence for lasting sequence segmentation in the discrete
sequence-production task. *Journal of motor behavior* **35**, 171-181 (2003).

96.   M. Ohbayashi, N. Picard, Sequential Reaching Task for the Study of Motor Skills in
Monkeys. *Bio Protoc* **10**, e3719 (2020).
97.   P. L. Strick, J. B. Preston, Two representations of the hand in area 4 of a primate. II.
Somatosensory input organization. *Journal of neurophysiology* **48**, 150-159 (1982).
98.   J. W. Holsapple, J. B. Preston, P. L. Strick, The origin of thalamic inputs to the "hand"
representation in the primary motor cortex. *The Journal of neuroscience* **11**, 2644-2654
(1991).

898
899
900

901 **Acknowledgments**

902   We thank Dr. Peter L. Strick for support, discussions and suggestions; Dr. Floh Thiels for
903   discussions; Moya Carrier for animal training and assistance; Mike Page for task
904   programming.
905

Funding:
This work was supported by
National Institutes of Health grant R01NS129551 (MO)
National Institutes of Health grant R21NS101499 (MO)
The Brain Sciences Project of the CNSI & NINS BS291006 (MO)

Author contributions:
Conceptualization: MO, NP
Methodology: MO
Investigation: MO
Visualization: MO
Supervision: MO
Writing—original draft: MO
Writing—review & editing: MO, NP
920

921 **Competing interests:** Authors declare that they have no competing interests.
922

923 **Data and materials availability:** All data needed to evaluate the conclusions in the paper are
924   present in the paper. The data in other format can be provided by M.O. pending scientific
925   review and a completed material transfer agreement. Requests for the data should be
926   submitted to: machiko@pitt.edu.
927
928

Figures and Tables

Fig. 1 Sequential Reaching Task. A. Sequential reaching task. B. Repeating sequences in the
Repeating task. C. Response Time (RT) of sequence '5-3-1' of the Repeating task. We arbitrary
divided the learning curve on the Repeating task into 3 periods based on the slope of RT changes.
The first period corresponds to the 'Fast Learning' phase where a large RT reduction occurs in a
short time frame. The second and third periods are defined as the early and late epochs of the
Slow Learning phase and correspond to 'Skill Acquisition' and the 'Development of Expert
Performance'. D. The introductions of the new sequences were staggered at ~ 100-200 days
intervals. This staggered training design enabled us to compare the effect of pharmacological
injections between the 'Long-term' trained and 'Short-term' trained sequences in a single
injection session. Gray circle: RT of the Random task; Blue triangle: RT of the Repeating
sequence "5-3-1"; blue lines in C: mean +/- s.d.; Green square: RT of the Repeating sequence "1-
2-4"; Red diamond: RT of the Repeating sequence "2-3-4": Arrow: injection of anisomycin.

Fig. 2 M1 location and the microstimulation maps of M1. A. Lateral view of cebus brain. Dashed lines indicate the M1-PMd border and the pre-PMd-PMd border. PS: principal sulcus; ArS: arcuate sulcus; CS: central sulcus; IPS: intra parietal sulcus; LS: lateral sulcus; pre-PMd: pre-dorsal premotor cortex; PMd: dorsal premotor cortex; M1: primary motor cortex. R: rostral; L: lateral. B. MRI image after the chamber implantation for monkey N. The dotted circle indicates the chamber outline. C, D. Intracortical stimulation map from monkey N and monkey R. Letters indicate the movements evoked at each site. S: Shoulder; E: Elbow; W: Wrist; D: Digit; Gray dots: representative injection sites.

Fig. 3 Effect of anisomycin injection on performance of the Random and Repeating tasks.
A–C. Effect on movements "3 to 4" in the sequence "2-3-4" of the Repeating task after 42 days of training. A. Reaching end points during Random and Repeating tasks (pre- and post-injection). Gray dots: correct responses; black dots: error responses. E_D: direction errors. Percentages of reaches ending at each target are shown below the targets. Touches between targets were assigned to the closest target. B. Errors increased only in the Repeating task after injection (Random: pre 96.96%, post 92.63%; χ^2 test, p=0.09; Repeating: pre 77.24%, post 40.20%; χ^2 test, $p < 0.0001$). C. Predictive responses decreased significantly after injection (pre 76.55%, post 12.74%; χ^2 test, $p < 0.0001$). D–F. Effect on movement "4 to 1" in the sequence "1-2-4" of the Repeating task after 171 days of training, and corresponding movements in the Random task. D. Reaching end points for movements "4-1" during Random and Repeating tasks (pre- and post-injection). Gray dots: correct responses; black dots: error responses. E_A: accuracy errors. E. Error rate significantly increased only in the Repeating task (Random: pre 89.47%, post 88.57%; χ^2 test, $p = 0.92$; Repeating: pre 84.54%, post 50.38%; χ^2 test, $p < 0.0001$). F. Predictive responses significantly decreased after injection (pre 84.54%, post 42.11%; χ^2 test, $p < 0.0001$). G–I. Effect on movement "1 to 5" in the sequence "5-3-1" of the Repeating task after 377 days of training, and corresponding movements in the Random task. G. Reaching end points for movements "1-5" during Random and Repeating tasks (pre- and post-injection). Gray: correct; black: error. E_A: accuracy errors. H. Errors significantly increased only in the Repeating task after injection (Random: pre 80.85%, post 100%; χ^2 test, $p = 0.006$; Repeating: pre 96.51%, post 66.33%; χ^2 test, $p < 0.001$). I. Predictive responses significantly decreased after injection (pre 95.58%, post 88.65%; χ^2 test, $p = 0.006$). Note: Asterisks indicate $p < 0.05$.

Fig. 4 Performance in the Repeating task of the injection session shown in Fig. 3. A-C.
Differences in performance measures of the most affected movement within each sequence,
calculated as post-injection performance minus the pre-injection performance. A. Increase in
error rates across the three sequences: '2-3-4' (37.0%), '1-2-4' (34.1%), and '5-3-1' (30.2%).
Significant differences were observed among all sequence pairs (generalized linear model
[GLM] analysis, $p < 0.001$). B. Decrease in percentage of predictive trials across sequences: '2-
3-4' (63.9%), '1-2-4' (42.4%), and '5-3-1' (35.3%). Significant differences were observed
among all sequence pairs (GLM analysis, $p < 0.001$). C. Increase in response times (RTs) across
sequences: '2-3-4' (385.6 ms), '1-2-4' (193.4 ms), and '5-3-1' (147.8 ms). Significant
differences were observed among all sequence pairs (two-way ANOVA, $p < 0.001$). D–L. Time
course of performance in the Repeating task from 4 days before to 7 days after anisomycin
injection (arrows indicate injection). Anisomycin injection in M1 impaired task performance for
2–3 days post-injection, even after extended training. Performance recovered to baseline levels
after 2–3 days of retraining. Asterisks denote significant differences from baseline (χ^2 *test* for
performance and predictive trials; *t-test* for RT; $p < 0.05$).

Fig. 5. The observed behavioral effects (Fig. 3) were specifically due to protein synthesis
inhibition affecting memory. A-C, The effect of muscimol injection in M1 on movements from
target "3 to 1" of the Random and Repeating tasks. A. Left: pre-injection; Right: post-injection.
gray: correct: black: error. B. Error rates. The number of error trials increased both in the
Random and Repeating tasks after the injection (Random: pre 100%, post 55.4 %: $\chi 2$ test, $p <$
0.001; Repeating: pre 81.0%, post 61. 8%; $\chi 2$ test, $p = 0.001$). C. Number of predictive responses
significantly decreased after the muscimol injection (pre 71.4%, post 37.8%; $\chi 2$ test, $p < 0.001$).
D. Activity of two individual M1 neurons. Neural activity was aligned at the monkey's release
of the touch screen. Left: Activity of a neuron recorded before the anisomycin injection (neuron
33). Right: Activity of a neuron recorded 24 hours after the anisomycin injection (neuron 56).
Upper panels: neural activity during the movements that had maximum firing rates (Neuron 33:
during movement from target 4 to 1; Neuron 56: during movement from target 3 to 1). Lower
panels: neural activity during the movements that had minimum firing rates (Neuron 33: during
movement from target 2 to 4; Neuron 56: during movement from target 4 to 1). Both neurons
were recorded 1 mm away from the injection site. Both neurons had movement selectivity in the
Random task. E. Selectivity index of M1 neurons recorded before and after anisomycin
injections (pre: n=62; post n=45). Selectivity Index, pre: mean=0.76, sd=0.24; post: mean=0.82,
sd=0.22 (*t-test*, *p*=0.22, *df*=74). There was no significant difference in selectivity indexes
between pre- and post- injection conditions. F. Maximum firing rate of M1 neurons recorded
before and after anisomycin injections (pre: n=62; post: n=45). Maximum firing rate, pre:
mean=21.73, sd=15.37, post: mean=21.59, sd=13.71 (*t-test*, *p*=0.96, *df*=104). Asterisks indicate
p < 0.05.

1018

Fig. 6 Effect of anisomycin injection on performance of the Repeating task (population data). A.B. Mean error rates (A) and mean response time (RT, B) of the most affected movements from 3 days before to 8 days after anisomycin injection (monkey N, n=5; monkey R, n=4). Arrows indicate injection timing. Light and dark gray dots denote the means for monkeys 1 and 2, respectively; error bars show se. Anisomycin injection in M1 affected both error rate and RT in the Repeating task for 2-3 days post-injection, even after 'Extended' training and performance returned to baseline after 2-3 days of training. C. Mean change in performance measures for Short-term, Long-term, and Extended trained sequences, calculated for the most affected movement within each sequence. Light and dark gray dots represent monkeys 1 and 2, respectively; error bars show s.e. Top: Increase in error rate: Short = 41.03%, Long=26.52%, Extended=36.98%. (*pairwise comparisons*: Short vs Long, $p = 7.9 \times 10^{-19}$, Short vs Extended, $p=0.48$, Long vs Extended, $p = 6.1 \times 10^{-19}$,). Bottom: Increase in RT: Short =280.47ms, Long=198.66ms, Extended=134.82ms. (*pairwise comparisons*: Short vs Long, $p = 2.1 \times 10^{-4}$; Short vs Extended, $p = 7.2 \times 10^{-42}$; Long vs Extended, $p = 2.9 \times 10^{-25}$). D. Effect size of RT change was negatively correlated with training duration (*Pearson's correlation coefficient, r*= -0.4964, *p*=5.4791e-05). The x-axis represents the number of training days, while the y-axis indicates the effect size of RT change. Asterisks indicate $p < 0.05$.

1036

Fig. 7 Effect of anisomycin injection on Response Time (RT). A. RTs in Random (black) and
Repeating tasks (sequence '5-3-1': blue triangles; '1-2-4': green squares; '2-3-4': red circles)
before (top) and after (bottom) an anisomycin injection. RTs increased for all three sequences
post-injection. B. Mean RTs for the first and last 20 trials of movement '3-1' in Random (top)
and Repeating ('5-3-1'; bottom) tasks during pre- and post-injection sessions (*t-test*, early vs late
trials: Random: pre $p=0.55$, post $p=0.37$; Repeating pre $p=0.36$, post $p=0.002$). C. Performance
test 3 days after the anisomycin injection. There was no significant effect of injection on
performance of the tasks and RTs. Top: Mean Error rate ($\chi 2$ test, $p>0.05$ for Random and
Repeating). Bottom: Mean RT (*t-test*, $p>0.05$ for Random and Repeating). Error bars, s.e.
Asterisks indicate $p<0.05$.