Decoding the biogenesis of HIV-induced CPSF6 puncta and their fusion with the nuclear speckle.

Chiara Tomasini[1*], Celine Cuche[1*], Selen Ay[1], Maxence Collard[1], Bin Cui[2], Mohammad Rashid[2], Shaoni Bhattacharjee[1], Bruno Tello-Rubio[1], Julian Buchrieser[3], Charlotte Luchsinger[2], Cinzia Bertelli[2], Vladimir N. Uversky[4], Felipe Diaz-Griffero[2#], Francesca Di Nunzio[1#]

[1] *Institut Pasteur, Advanced Molecular Virology Unit, Department of Virology, Université Paris Cité, 75015 Paris, France*

[2] *Albert Einstein College of Medicine, Department of Immunology and Microbiology, New York 10461, USA*

[3] *Institut Pasteur,* Virus and Immunity Unit*, Department of Virology, Université Paris Cité, 75015 Paris, France.*

[4] *Department of Molecular Medicine and USF Health Byrd Alzheimer's Research Institute, Morsani College of Medicine, University of South Florida, Tampa, FL 33612, USA*

 # Address correspondence to Francesca Di Nunzio, dinunzio@pasteur.fr, or Felipe Diaz-Griffero, felipe.diaz-griffero@einsteinmed.edu

* These authors have equally contributed

Highlights

- The formation of HIV-induced CPSF6 puncta is critical for restoring HIV-1 nuclear reverse transcription.
- CPSF6 protein lacking the FG peptide cannot bind to the viral core, thereby failing to form HIV-induced CPSF6 puncta.
- The FG peptide, rather than low-complexity regions (LCRs) or the mixed charge domains (MCDs) of the CPSF6 protein, drives the formation of HIV-induced CPSF6 puncta.
- The presence of the FG peptide plays a pivotal role in facilitating viral replication within macrophage-like cells.
- HIV-induced CPSF6 puncta form individually and later fuse with nuclear speckles (NS) via the intrinsically disordered region (IDR) of SRRM2.

Summary

Viruses rely on host cellular machinery for replication. After entering the nucleus, the HIV genome accumulates in nuclear niches where it undergoes reverse transcription and integrates into neighboring chromatin, promoting high transcription rates and new virus

progeny. Despite antiretroviral treatment, viral genomes can persist in these nuclear niches

and reactivate upon treatment interruption, raising the possibility that they could play a role in

the establishment of viral reservoirs. The post-nuclear entry dynamics of HIV remain unclear,

and understanding these steps is critical for revealing how viral reservoirs are established.

In this study, we elucidate the formation of HIV-induced CPSF6 puncta and the domains of

CPSF6 essential for this process. We also explore the roles of nuclear speckle scaffold factors,

SON and SRRM2, in the biogenesis of these puncta. Through genetic manipulation and

depletion experiments, we demonstrate the key role of the intrinsically disordered region of

SRRM2 in enlarging nuclear speckles in the presence of the HIV capsid.

We identify the FG domain of CPSF6 as essential for both puncta formation and binding to the

viral core, which serves as the scaffold for CPSF6 puncta. While the low-complexity regions

(LCRs) modulate CPSF6 binding to the viral capsid, they do not contribute to puncta formation,

nor do the disordered mixed charge domains (MCDs) of CPSF6. Interestingly, the FG peptide

facilitates viral replication. These results demonstrate how HIV evolved to hijack host nuclear

factors, enabling its persistence in the host.

Of note, this study provides new insights into the underlying interactions between host factors

and viral components, advancing our understanding of HIV nuclear dynamics and offering

potential therapeutic targets for preventing viral persistence.

Introduction

Since the discovery of HIV (Barre-Sinoussi et al., 1983), the initial stages of the viral life cycle, such as the reverse transcription (the conversion of the viral RNA in DNA) and the uncoating (loss of the capsid), have been understood to mainly occur within the host cytoplasm. Only the pre-integration complex, carrying the fully reverse-transcribed viral DNA, was believed to enter the nucleus for integration into the host chromatin (Suzuki and Craigie, 2007). Recent studies highlighted that the viral genome is transported in the nucleus via a shuttle that shields it from the hostile cellular environment (Rasaiyaah et al., 2013). This shuttle is constituted by the viral capsid (Blanco-Rodriguez and Di Nunzio, 2021; Blanco-Rodriguez et al., 2020; Chen et al., 2016; Selyutina et al., 2020; Yamashita and Emerman, 2004; Zila et al., 2021), which comprises 250 hexamers (depending on the size and shape of the capsid) and 12 pentamers (Pornillos et al., 2009). Within the structure of the capsid, hydrophobic pockets exist between hexamers, which serve as targets for various nucleoporins, especially the ones carrying FG repeats, facilitating the translocation of the capsid through the nuclear pore complex (NPC) (Buffone et al., 2018; Di Nunzio, 2013; Di Nunzio et al., 2012; Lelek et al., 2015; Matreyek et al., 2013; Price et al., 2014). Recent studies suggest that HIV uses multiple FG regions of several nucleoporins to translocate through the NPC, acting as a chaperone by itself (Dickson et al., 2024; Fu et al., 2024). Using a reductionist system of Nup98 condensates, it has been demonstrated that FG-mediated phase partitioning identifies specific sites on the capsid that allow it to interact autonomously with these phases. These findings complement the evolving understanding of the early stages of HIV infection, which has been revisited in recent years. They unveil that crucial stages of early viral infection occur within the host nucleus (Burdick et al., 2020; Dharan et al., 2020; Francis et al., 2020; Scoca et al., 2023; Selyutina et al., 2020). Notably, it has been shown that the pre-integration complex forms within the host nucleus (Muller et al., 2021; Scoca et al., 2023), and incoming viral RNA genomes accumulate in nuclear niches containing, cleavage and polyadenylation specificity factor subunit 6 (CPSF6), RNA-binding protein SON and Splicing component, 35 kDa (SC35, also known as serine/arginine-rich splicing factor 2 (SRSF2) (Francis et al., 2020; Rensen et al., 2021; Scoca et al., 2023). However, the mechanistic requirements that govern post-nuclear entry phases, which are crucial for a successful viral infection and the establishment of viral reservoirs, remain enigmatic.

Specifically, the mechanism underlying the formation of nuclear niches containing viral components and nuclear speckle (NS) factors, such as SC35—traditionally used as a marker for NS remains unclear. However, recent studies have revealed that SON and serine/arginine repetitive matrix protein 2 (SRRM2) are essential for NS biogenesis (Fu and Maniatis, 1990). Particularly, intrinsically disordered regions (IDRs) play a critical role in the NS formation (Ilik

et al., 2020). These membraneless organelles (MLOs) fulfil various cellular functions besides splicing. Recent evidence indicates a direct role of NSs in cellular transcription regulation, as their spatial proximity correlates with gene expression amplification, as demonstrated by live-cell imaging of heat-shock responsive genes (Chen et al., 2018; Zhang et al., 2021). HIV, being a virus capable of generating new particles through splicing and integrating into active host genes, finds NSs highly conducive for viral replication. HIV particles, along with their RNA genome, accumulate within nuclear MLOs enriched in NS factors (Ay et al., 2025; Rensen et al., 2021; Scoca et al., 2023). Notably, CPSF6, a paraspeckle factor first identified as a viral partner by KewalRamani's laboratory (Lee et al., 2010), has been clearly detected in HIV-induced CPSF6 puncta (Ay et al., 2025; Francis et al., 2020; Lee et al., 2010; Luchsinger et al., 2023; Rensen et al., 2021; Scoca et al., 2023). These puncta serve as hubs for nuclear reverse transcription and the formation of pre-integration complexes, which generate active proviruses detected outside but in close proximity to NSs (Ay et al., 2025; Li et al., 2021; Scoca et al., 2023).

In our study, we aim to elucidate how HIV-induced CPSF6 puncta form and identify the NS factors involved in their formation. CPSF6, along with NS factors, contains intrinsically disordered regions that can guide HIV-1 to the correct nuclear location for successful infection or allow the virus to remain sequestered during drug treatment, forming reservoirs. Importantly, viral reservoirs are the major bottleneck for curing the infection.

In this study, we investigate which disordered domain of CPSF6 is responsible for the recruitment of the viral core and generation HIV-induced CPSF6 puncta in the host nucleus. Simultaneously, we elucidate the key component of NSs that, through its IDRs, enable fusion with HIV-induced CPSF6 puncta, likely stabilizing them. Notably, we observed that the virus rebounds when anti-reverse transcription drugs are removed, but only if nuclear niches containing HIV, NS factors, and CPSF6 are present. If these niches are pharmacologically dismantled, viral rebound does not occur.

Overall, studying the biogenesis of HIV-induced nuclear niches is crucial for understanding how the virus navigates and persists in the nucleus of infected cells, and for designing new antiretroviral strategies.

Results

Critical Role of HIV-Induced CPSF6 Puncta in Restoring Nuclear reverse transcription After Anti-Reverse transcriptase (RT) Therapy Discontinuation.

Upon nuclear entry, HIV enhances the formation of CPSF6 puncta, where reverse transcription

ends. The treatment with the reversible RT inhibitor nevirapine (NVP) can trap the viral RNA

genome in these nuclear niches (Ay et al., 2025; Rensen et al., 2021; Scoca et al., 2023).

Once NVP is removed, the trapped vRNA can resume RT entirely within the nucleus, a process

we term nuclear reverse transcription. Here we demonstrate that this phenomenon depends

on the presence of CPSF6 puncta, which are mainly composed of CPSF6 proteins and viral

cores. Disruption of these puncta by high doses of PF74 (25 µM) significantly impairs the

rebound of the RT, as evidenced by the absence of luciferase expression, similar to what is

observed with complete NVP treatment (Figure 1A-B).

CPSF6 FG Domain is Required for HIV-Induced Puncta Formation.

Formation of CPSF6 puncta upon HIV-1 infection hinges on two key events: the entry of the

HIV-1 core into the nucleus and the binding of CPSF6 to the HIV-1 core (Blanco-Rodriguez

and Di Nunzio, 2021; Blanco-Rodriguez et al., 2020; Buffone et al., 2018; Zila et al., 2021). To

determine the contribution of CPSF6's disordered domains for the formation of CPSF6 puncta

upon HIV-1 infection, we correlated the binding of CPSF6 to the HIV-1 core with the formation

of CPSF6 puncta. To this end, we first generated CPSF6 knockout (KO) THP-1 cells (Figure

1C-D) to eliminate the interference from the endogenous protein, which could affect the

interpretation of results regarding the role of the analyzed CPSF6 domains. CPSF6 depletion

in THP-1 cells was performed by CRISPR Cas9 technology. To completely eliminate the

expression of the CPSF6 gene we selected single clones by limiting dilution. We identified a

clone that was completely KO for CPSF6, confirmed through western blot and

immunofluorescence (Fig. 1C-D) and we infected this clone and the control clone with HIV.

CPSF6 puncta were detected only in the control-infected cells and not in the KO clone (Fig.

1D; Suppl. Fig.1A). The viral integrase (IN) was observed within CPSF6 puncta, consistent

with previous studies (Francis et al., 2020; Rensen et al., 2021; Scoca et al., 2023), but absent

in CPSF6 KO cells where viral IN was observed in the cytoplasm (Fig. 1D; Suppl. Fig.1A).

Thus, we used KO cells for CPSF6 to assess the role of selected CPSF6 domains in HIV-

induced condensates (Suppl. Fig.1B). We designed various CPSF6 deletion mutants (Figure

2A) to specifically assess the significance of the main disordered regions of CPSF6 protein

(Suppl. Figure 2A) such as the FG (phenylalanine and glycine) motif, the low complexity

regions (LCRs), and the mixed charge domain (MCD), in the ability of CPSF6 to bind to the

core and facilitate the formation of CPSF6 puncta. We investigated the role of the FG peptide

by generating a mutant that exclusively lacks the FG peptide (ΔFG). Previous *in vitro* studies

have shown that the FG peptide binds to the hydrophobic pocket formed between capsid

hexamers (Buffone et al., 2018; Price et al., 2014). Here, we want to investigate the role of FG

peptide in the context of the protein during the viral infection.

To further explore this, we developed an alternative plasmid by expanding the FG peptide

deletion to include surrounding prion-like LCRs (ΔFG ΔLCR). These regions, outside the

CPSF6 context, have been identified as crucial for facilitating strong CPSF6 binding to capsid

lattices (Wei et al., 2022). In our study, we aim to evaluate their role within a more physiological

setting. Additionally, we assessed a CPSF6 variant that carries the 15-mer FG peptide flanked

by non-LCR sequences, such as those derived from Beta-adducin (ADD2), kindly provided by

Mamuka Kvaratskhelia (ΔLCR+ADD2). These protein segments are known for their high

flexibility, akin to the LCR of CPSF6. Furthermore, to elucidate the contribution of the LCRs of

CPSF6 in the formation of CPSF6 puncta, we generated a mutant lacking both LCRs (ΔLCR).

Analysis of the MCD contribution to both the ability of CPSF6 to bind to the core and formation

of CPSF6 puncta was achieved by deleting the MCD and adding 3 nuclear localization signals

(3xNLS ΔMCD) since the deletion of the MCD results in a protein that localizes mainly into the

cytoplasm (Figure 2A-B; Suppl. Figure 2B).

To correlate the ability of CPSF6 to bind to the HIV-1 core with formation of CPSF6 puncta,

we expressed wild type and mutant CPSF6 constructs in THP-1 cells knockout for CPSF6.

Subsequently, we infected these cells with HIV-1 and analyzed the presence or absence of

CPSF6 puncta at 24 hours post-infection. Importantly, for the imaging experiment we

expressed CPSF6 WT and mutants without tags to avoid the formation of aggregates that

could interfere with our conclusions. Our data show that HIV-induced CPSF6 puncta can form

extremely rarely with the deletion mutant CPSF6 ADD2ΔLCR and with the mutant lacking the

FG or both the FG peptide and the LCRs (Fig. 2B-D; Suppl. Fig.1B). However, when we

analyzed the role of the MCD domain in CPSF6 puncta formation, which was indicated to be

important for condensing CPSF6 in NS (Greig et al., 2020), comparing the number of CPSF6

WT puncta induced by HIV infection with CPSF6 mutants revealed that the MCD domain does

not play a critical role in HIV-induced CPSF6 puncta formation (Fig. 2B-D). In addition, we

observed that the majority of analysed CPSF6 3xNLSΔMCD puncta contain vRNA inside,

similar to CPSF6 WT puncta (Figure 2E), thus corroborating the lack of a role for this

intrinsically disordered domain in HIV-induced CPSF6 puncta. Since the NLS domain from

SV40, which replaces the MCD, is highly basic and could potentially induce condensates, we

fused CPSF6 with a non-basic NLS (PY-NLS) or removed the NLS entirely. Even though these

two proteins do not efficiently enter the nucleus, the few that do manage to reach the nucleus

can host viral particles, as evidenced by the presence of IN. Many viruses are typically blocked

in the cytoplasm due to the presence of these mutants that are mainly cytoplasmic. However,

because we used a high viral dose, the blockage in the cytoplasm was not complete. As a

result, the viruses that successfully entered the nucleus induced the formation of puncta
associated with CPSF6-deleted mutants, indicating that the MCD is not critical for the formation
of HIV-induced CPSF6 puncta (Fig.2F). Similar to the MCD, when we compared CPSF6
truncated for the LCRs with CPSF6 WT, we observed that the LCRs do not contribute to CPSF6
puncta formation. Therefore, the FG peptide alone, without the LCRs, is the only CPSF6
domain required for their formation (Fig. 2B-F; Suppl. Fig.1B).

Next, we tested the ability of the different CPSF6 deletion mutants for their ability to bind the
viral core using a previously described capsid binding assay (Selyutina et al., 2018). Wild type
and mutant CPSF6 proteins were expressed in human 293T cells at similar levels (INPUT)
(Figure 3A). Extracts containing wild type and mutant CPSF6 proteins were incubated with
stabilized HIV-1 capsid tubes for 1 h at 25° C in the presence of 10 µM of PF74, which is a
small molecule that competes with CPSF6 for binding to the hydrophobic pocket formed
between hexamers that constitute the viral core (Buffone et al., 2018; Price et al., 2014). HIV-
1 capsid stabilized tubes were washed, and the bound proteins were eluted using Laemli buffer
(BOUND). For every construct, the percentage of bound protein relative to input in the
presence or absence of PF74 is shown (Figure 3B). Our results revealed that the absence of
the FG peptide (ΔFG) entirely abolished CPSF6's ability to bind to the viral core. In agreement,
simultaneous deletion of the FG motif and LCRs (ΔFG ΔLCR) resulted in a construct unable to
bind to the viral core. Similar outcomes were observed when the LCRs were replaced with
sequences derived from ADD2, even if the FG was present.

LCR-FG is notably more disordered than ADD2-FG, containing a high proportion of prolines
(48 out of 98 residues), which makes it mostly non-foldable (Figure 4A-H). Since proline is a
structure-disrupting residue, LRC-FG is not expected to adopt any secondary structure. In
contrast, ADD2-FG contains fewer prolines (15 out of 98 residues) but has many charged
residues. It is predicted to form two short α helices and a ß strand, arranged as: α helix - FG –
ß strand - α helix (Figure 4E). ADD2-FG may form a flexible collapsed state, as its oppositely
charged residues are evenly distributed, potentially allowing polyelectrostatic compaction. This
suggests that FG within ADD2-FG may be less accessible for the interaction with the viral
core's hydrophobic pocket due to its involvement in this collapsed conformational ensemble
(Figure 4A-E, suppl. Fig.3). This aligns with the inability of CPSF6 carrying ADD2 in place of
the LCRs to induce CPSF6 puncta (Figure 2B). On the other hand, the deletion of only the
two LCRs, while keeping the FG peptide intact, resulted in unexpected findings. The ΔLCR
mutant exhibited a stronger binding affinity for the viral core when compared to the wild-type
protein (Figure 3B). These results suggest that the LCRs surrounding the FG motif are
modulating the affinity of CPSF6 to the viral core, which might be important for function. By
contrast, deletion of the MCD (ΔMCD) but retention of other regions, such as the FG peptide

and the LCRs, demonstrated a binding affinity to the viral core similar to that of the wild-type
protein. These results suggest that the MCD domain is not involved in the binding of CPSF6
to the viral core, which is not surprising since the CPSF6 (1-358), which does not have an
MCD, binds to the viral core (Lee et al., 2010).

Thus, the viral capsid, through the FG peptide of CPSF6, constitutes the scaffold of HIV-
induced CPSF6 puncta.

In summary, our results suggest that the FG peptide is the main determinant involved in the
binding of CPSF6 to the viral capsid. Interestingly, our work implies that the LCRs may be
modulating the affinity of the FG motif for the viral core (Figure 3B). Recognition motifs that
mediate protein-protein interactions, such as the FG motif of CPSF6, are usually embedded
within longer intrinsically disordered regions that can modulate affinity of the interaction
(Karlsson et al., 2022). Taken together, our data show that the FG peptide coordinates both
the binding to the viral core and the induction of CPSF6 puncta. This coordination suggests
that the FG peptide plays a critical dual role in recognizing the viral capsid and facilitating the
cellular clustering of CPSF6, which may be part of the cellular response to viral entry.

Role of FG domain in viral replication.

We aimed to investigate the role of FG domain of CPSF6 in viral infection. To correctly perform
the infectivity assay, we generated stable cell clones expressing protein levels comparable to
WT cells (Fig.5 A-B). Importantly, we assessed the reproducibility of our experiments by
freezing and thawing these clones. Thus, we performed infectivity assays in THP-1
macrophage-like cells WT, KO for CPSF6 or complemented with CPSF6ΔFG, using both a
single-round virus and a replication-competent virus. The results, shown in Figure 5 C-D,
indicate that complete depletion of CPSF6 reduces infectivity, as measured by luciferase
expression in a single-round infection (KO: ~65%; ΔFG: ~74%; compared to WT: 100% on
average). Notably, a more pronounced defect in viral particle production was observed when
WT virus was used for infection (KO: ~21%; ΔFG: ~16%; compared to WT: 100% on average).
According to our results, depletion of full-length CPSF6, or expression of a CPSF6 variant
lacking only the FG domain, both reduce viral infection compared to cells expressing full-length
CPSF6, indicating that the FG domain plays an important role in HIV-1 infection.

HIV-induced CPSF6 mutants puncta and Nuclear Speckles.

The NS factor SC35, commonly used as a marker of NS, has been detected in HIV-induced
CPSF6 puncta (Figure 6A)**.** In this study, we investigated whether HIV-induced CPSF6 mutant

puncta are associated with SC35. We infected cells expressing CPSF6 WT, CPSF6 3xNLS ΔMCD, CPSF6 ΔLCR, CPSF6 ΔMCD, and CPSF6 PY NLS ΔMCD, and examined whether the nuclear puncta formed by the various CPSF6 proteins associate with SC35. Imaging analysis revealed no significant difference in the association of SC35 with CPSF6 WT or the CPSF6 mutants (Figure 6B), confirming that the disordered domains, LCRs and MCDs, are dispensable for the formation of HIV-induced CPSF6 puncta that localize in the canonical nuclear niches marked by SC35.

Biogenesis of HIV-Induced CPSF6 Puncta carrying Nuclear Speckle factors.

Nuclear speckle factors, particularly those involved in their biogenesis, such as SON and SRRM2 (Ilik et al., 2020), have been identified as constituents of HIV-induced CPSF6 puncta. However, the specific role of NSs in these puncta remains unclear. Most of the existing results have been obtained through immunofluorescence experiments conducted several days post-infection. In this study, we conducted a time course experiment to investigate the biogenesis of HIV-induced CPSF6 puncta, which contain NS factors. Our goal was to capture the fusion event between the HIV-induced CPSF6 puncta and NS, providing insights into the dynamics of how HIV manipulates host nuclear structures during infection. We hypothesized that NS factors might either be recruited during the initial formation of HIV-induced CPSF6 puncta, shortly after the virus is released from the nuclear basket of the NPC, or later via the fusion between NSs and HIV-induced CPSF6 puncta. To investigate this, we performed live imaging to track CPSF6-mNeonGreen and NSs in cells expressing endogenous SRRM2 fused with a Halo tag, using CRISPR Paint (courtesy of Roy Parker) (Lester et al., 2021).We fixed and labeled samples at different time points post-infection, ranging from 6 h.p.i. to 30 h.p.i. (Figure 7A, Suppl. Fig.4A-B). At 6 h.p.i., 27% of HIV-induced CPSF6 puncta were still individual, compared to only 9% at 30 h.p.i. (Figure 7B). Concurrently, 61% of HIV-induced CPSF6 puncta were fused with NS at 6 h.p.i., rising to 75% at 30 h.p.i. (Figure 7B). This indicates a progressive increase in the number of HIV-induced CPSF6 puncta fusing with NS over time (Figure 7C). Overall, we detected individual CPSF6 puncta (green) and NSs (red) that quickly fused, confirming that this fusion occurs within the two independent puncta, CPSF6 and SRRM2, rather than during the formation of HIV-induced CPSF6 puncta (Figure 7A-C, movies 1A, 1B).

Taken together, these results suggest that HIV-induced CPSF6 puncta first form independently of NS and later fuse with NS, causing an enlargement of NS as part of the hijacking process by HIV-1.

Role of SON and SRRM2 in the Fusion and Stabilization of HIV-Induced CPSF6 Puncta within Nuclear Speckles.

Macrophage-like cells, THP-1, were depleted for SON or SRRM2 using AUM*silence*™ ASO technology (Gao et al., 2024; Marasca et al., 2022; Mazzeo et al., 2024; Zhang et al.). The level of depletion of SON and SRRM2 was evaluated by immunofluorescence and western blot using antibodies against SON and SRRM2 (Figure 8A). Both depleted cells were analyzed also for the presence of NS, labelled by SC35 (Suppl. Fig.6A). However, recent findings suggest that the primary target of the SC35 mAb is SRRM2 (Ilik et al., 2020). To confirm this, cells depleted of SRRM2 were labelled with antibodies against both SRRM2 and SC35 (Suppl. Figure 6 A-B). We have observed that the reduction of SRRM2 resulted in a slight decrease in the mean intensity of SC35, whereas the depletion of SON (Suppl. Fig.6A) did not have the same effect.

Subsequently, we infected THP-1 depleted cells for SRRM2 and SON or control cells with HIV-1 and we fixed them at 48 h post-infection for immunofluorescence. We calculated the percentage of CPSF6 puncta in HIV infected THP-1 control cells (approximately 78%) and in HIV infected THP-1 cells depleted for SRRM2 (about 43%) and SON (around 66%). Results from our experimental conditions indicate that the partial depletion of SRRM2 affects the formation of HIV-induced CPSF6 puncta, while the depletion of SON slightly reduces their establishment (Figure 8B).

The IDR of SRRM2 is a crucial element for the fusion of HIV-Induced CPSF6 puncta to the nuclear speckles.

CPSF6 contains several disordered regions (Di Nunzio et al., 2023) as well as NS factors (Ilik et al., 2020). Previous studies have established the role of IDR of SRRM2 in the biogenesis of NS (Ilik et al., 2020).

Here, we investigate the role of IDRs within NS factors, with a specific emphasis on SRRM2, in the fusion and stabilization of HIV-induced CPSF6 puncta. To address this inquiry, we utilized previously published HEK293 cell lines generated using the CRISPaint system (Lester et al., 2021), comprising two distinct lines: HaloTag SRRM2, SRRM2 full-length (FL) (1-2748 aa) fused with Halo tag(insertion at amino acid [aa] 2,708), and a cell line lacking the C-terminal IDR of SRRM2, known as ΔIDR HaloTag SRRM2(1-429 aa, with halo insertion at aa 430) .

As anticipated, the truncated form of SRRM2 displayed a more diffuse distribution within the nucleus, without recruitment to NS (suppl. Fig.5 A-E), and consequently lacked nuclear puncta

(Figure 8C top panels). On the other hand, the number of SON puncta was highly similar

between the two cell lines (Figure 8C, panels on the bottom, Suppl. Fig.7).

Subsequently, we quantified the formation of CPSF6 puncta in both cell lines infected with HIV-

1. No significant difference in HIV-induced CPSF6 puncta formation was observed between

HEK293 and HEK293 carrying the SRRM2 halo tag (~27%). However, a substantial reduction

in HIV-induced CPSF6 puncta was evident in the cell line carrying the SRRM2 form that lacks

the IDR (~11%) (Figure 8D). Collectively, these results underscore the pivotal role of the

SRRM2 IDR in the stabilization of HIV-induced CPSF6 puncta through their fusion with NSs

(Figure 7A-C).

Discussion

HIV-1 capsid has transformed long-standing assumptions in the field. Previously considered an undruggable viral target due to the belief that it disassembled shortly after HIV entered target cells, whereas recent findings have revealed its essential role in nuclear import (Ay and Di Nunzio, 2023; Blanco-Rodriguez and Di Nunzio, 2021; Blanco-Rodriguez et al., 2020; Chen et al., 2016; Taylor and Fassati, 2024; Yamashita and Emerman, 2004; Zila et al., 2021). Notably, the first-in-class antiretroviral capsid inhibitor, Lenacapavir, has shown remarkable results in patients, demonstrating that the capsid, contrary to previous belief, can indeed be a viable therapeutic target. While Lenacapavir improves patients' quality of life with only two injections per year (Link et al., 2020; Segal-Maurer et al., 2022), no current antiretroviral drugs provide a cure. This may be due to our incomplete understanding of certain aspects of HIV biology. In this study, we shed light on the post-nuclear entry steps, a critical phase for the establishment of viral reservoirs, which represent the main barrier to a cure. Recent findings have shown that not only the HIV viral capsid translocates through the NPC, but that the viral nuclear entry also enhances the formation of CPSF6 puncta. Additionally, it has been revealed that RT is completed within the nucleus (Burdick et al., 2020; Dharan et al., 2020; Rensen et al., 2021; Scoca et al., 2023; Selyutina et al., 2020). Furthermore, incoming viral RNA has been observed to be sequestered in nuclear niches in cells treated with the reversible reverse transcriptase inhibitor, NVP. When macrophage-like cells are infected in the presence of NVP, the incoming viral RNA is held within the nucleus (Ay et al., 2025; Rensen et al., 2021; Scoca et al., 2023). This scenario could also occur in patients receiving antiretroviral therapy. Notably, we found that pharmacological dismantling of CPSF6 puncta prevents the restoration of nuclear reverse transcription, indicating that HIV-induced CPSF6 puncta are critical for the viral life cycle. Conversely, THP-1 cells either depleted of CPSF6 or expressing a CPSF6 variant lacking the FG domain exhibit impaired viral replication. Understanding the biogenesis of these puncta could be a significant step towards deepening our knowledge of HIV biology and providing additional tools to combat this pandemic virus.

Here we identify the disordered FG peptide essential for the binding with the viral capsid as the inducer of HIV-induced CPSF6 puncta. Notably, CPSF6 protein lacking the FG peptide is incapable of forming nuclear puncta. Moreover, we discovered that the two major intrinsically disordered regions of CPSF6, the MCDs and the LCRs, are dispensable for the formation of viral nuclear puncta. The MCDs of CPSF6 have been shown to provide cohesion for NS condensation (Greig et al., 2020). A study performed in HEK293T-derived CPSF6 knockout (CKO) cells (clone B8) complemented with a tagged CPSF6 reported that the MCD is critical for CPSF6 cluster formation (Jang et al., 2024). In contrast, we chose to work with a more physiologically relevant cell line for HIV infection, macrophage-like cells, and we employed

untagged CPSF6 proteins (WT and deletion mutants) to avoid potential artifacts introduced by
protein tagging. Under our experimental conditions, the MCD did not appear to play a critical
role in the formation of HIV-induced CPSF6 puncta. CPSF6 puncta induced by CPSF6
depleted for the MCD upon infection are highly similar to those formed by wild-type CPSF6,
suggesting that MCDs do not play a significant role in this process. Additionally, the LCRs of
CPSF6 also do not appear to influence puncta formation, as their depletion does not reduce
the number of CPSF6 puncta. This indicates that neither the MCDs nor the LCRs are involved
in CPSF6 puncta formation during HIV infection. Surprisingly, when we assessed the ability of
CPSF6 domains to bind to the viral capsid, we observed that the deletion of LCRs increases
CPSF6's ability to bind to the viral capsid. We hypothesize that a change in charges may alter
the binding mechanism of CPSF6 when LCRs are absent. In scenarios where the FG motif is
depleted, we observed a dramatic inhibition of HIV-induced CPSF6 puncta formation and a
lack of binding to the viral core *in vitro*. On the flip side, the linkage between CPSF6 entities is
facilitated by the FG peptides' interaction with certain hydrophobic CA pockets along adjoining
hexamers (Wei et al., 2022). Therefore, it's conceivable that FG peptides, not involved in the
capsid's binding, could coalesce similarly to FG-Nups. These undergo phase separation,
forming condensates with nuclear pore complex-like permeability barrier features (Hülsmann
et al., 2012).

We have also identified HIV-induced CPSF6 puncta formation independent of NS at early
stages post-infection, with a progressive increase in CPSF6 puncta co-localizing with NS over
time. These results were obtained by focusing our studies on two scaffold proteins involved in
NS biogenesis: SON and SRRM2. Over the past approximately 0.6–1.2 billion years of
metazoan evolution, these two factors have undergone significant lengthening, unlike many
other proteins involved in splicing. This extension primarily occurred within their IDRs, which
are commonly associated with liquid-liquid phase separation (LLPS) and the formation of
biomolecular condensates (Rai et al., 2018). Co-depleting SRRM2 with SON, or depleting SON
in a cell line where it is deleted the intrinsically disordered C-terminus of SRRM2, abolished
the formation of NSs (Ilik et al., 2020). However, the depletion of only one factor does not
abolish NSs. Consistent with this finding, we observed that the depletion of SRRM2 does not
affect the presence of SON nuclear puncta. Similar results were obtained with cells genetically
modified to express the truncated form of SRRM2 lacking the IDRs. When the ΔIDR SRRM2-
halo tag was detected by the halo ligand, there was no recruitment of the truncated SRRM2
form in NSs. However, if SC35 is used as a target for antibodies, it can still be detected, albeit
with a much lower intensity signal than in cells expressing the full-length SRRM2. Additionally,
we observed a significant reduction in the detection of HIV-induced CPSF6 puncta. The few
CPSF6 puncta detected in these cells colocalized with the weak SC35 signal. These results

suggest that the IDR of SRRM2 plays an important role in HIV-induced CPSF6 puncta
stabilization, but rare puncta can appear, likely induced by redundant NS factors. This indicates
that further investigation is needed to better understand the hijacking of NS by HIV.

Taken together, our results reveal the intricate interplay between individual CPSF6 domains
and the viral capsid in dictating the nuclear fate of the virus. Concurrently, the IDR domain of
SRRM2 contributes to the enlargement of particular nuclear speckles with the fusion of HIV-
induced CPSF6 puncta. Lastly, these HIV-induced CPSF6 puncta necessitate of FG peptide
to engage the viral core.

Overall, this study could provide insights into the understanding of viral invasion and
persistence within the host.

Materials and methods

Cell lines

THP-1 cells are immortalized monocytic cells, which, once seeded, differentiate into
macrophage-like cells under phorbol 12-myristate 13-acetate (PMA) treatment (160 nM). THP-
1 cells were also engineered knocking out CPSF6. These cells were cultivated in RPMI 1640
medium supplemented with 10% fetal bovine serum (FBS) and 1% penicillin–streptomycin
solution (100 U/ml). HEK293T cells are human embryonic kidney cells used to produce LVs.
For figure 2, we used two engineered HEK293 strains, Halo tagged SRRM2 HEK293 cells and
Halo tagged SRRM2 ∆IDR HEK293 cells, kind gifts from Roy Parker's lab (Lester et al., 2021),
characterized by the Halo tagged SRRM2 protein. In Halo tagged SRRM2 ∆IDR HEK293 cells
Parker's lab deleted also the SRRM2 sequence encoding for aa 430-2748. All the HEK293
strains were cultivated in Dulbecco's modified Eagle medium supplemented with 10% FBS and
1% penicillin–streptomycin (100 U/ml).

Bacteria strains

All E. coli bacteria strains were grown in Luria-Bertani (LB) medium at 37°C. DHα competent
cells and Stellar Competent Cells were used for molecular cloning, while E.coli One-Shot
BL21star (DE3) cells were exploited for protein production.

Plasmids

To express the WT CPSF6/WT CPSF6-mNeonGreen and the mutant CPSF6/mutant CPSF6-
mNeonGreen clones, the correspondent coding sequences were engineered in pSICO
plasmids. The two original plasmids used were pSICO CPSF6-mNeonGreen and pLPCX
CPSF6 ADD2, gift from Mamuka's lab. HIV-1ΔEnvIN$_{HA}$ LAI (BRU) plasmid encodes the
ΔEnvHIV-1 LAI (BRU) viral genome where the IN protein is fused to the HA tag while pNL4.3
Δenv ΔNef IRES GFP plasmid encodes the ΔEnvHIV-1 NL4.3 viral genome and contains also
a GFP sequence headed by an IRES. The pNL4.3 Δenv ΔNefLuc has the Luciferase cDNA as
reporter gene.

CRISPR-Cas9 knockout in THP-1 cells

To target CPSF6, three different crRNAs were used simultaneously (specific sequence: 5'-
TCGGGCAAATGGCCAGTCAAAGG-3', 5'-AGGACGGGGCCGTTTTCCAGGGG-3', and 5'-
CATGTAATCTCGGTCTTCTGGGG-3', all ordered from Integrated DNA Technologies, IDT).
Pre-designed unspecific crRNA was used as control (IDT). crRNA and tracrRNA were
resuspended in IDT Duplex Buffer according to the manufacturer's instructions. On the day of
the nucleofection, duplexes were formed by mixing equimolar concentration of crRNA and
tracrRNA, followed by 5-min annealing at 95°C. RNA duplexes were then mixed (1:2) with
TrueCut^{TM} Cas9 Protein v2 for 10 min at room temperature to generate ribonucleoprotein
(RNP) complexes. 2×10^5 THP1 cells were resuspended in P3 Primary Cell Nucleofector^{TM}
Solution, mixed with RNP and Alt-R$^®$ Cas9 Electroporation Enhancer (90 pmol, IDT), and
nucleofected in a 4D-Nucleofector^{TM} System using the P3 Primary Cell 4D-Nucleofector^{TM}X Kit
S (program FI-110). After nucleofection, cells were seeded in complete RPMI medium with
20% FBS. Three days after nucleofection, cells were plated for clonal selection.

Clonal selection of KO cell lines

479 Seventy-two hours post nucleofection, cells were diluted in RPMI medium containing 20% FBS
480 and plated in five 96-well plates at 1 and 5 cells/well condition. After one-month, selected
481 microcolonies (50-100) are placed into 24- wells plate. Once that wells were near confluence,
482 cells were transferred into the well of a 6- well plate. After growing for another one-month, cells
483 were proceeded for western blot.

Western blot

Proteins were extracted on ice from THP-1 cells using RIPA buffer (20 mM HEPES, pH 7.6,
150 mM NaCl, 1% sodium deoxycholate, 1% Nonidet P-40, 0.1% SDS, 2mM EDTA, complete
protease inhibitor), and protein concentration was quantified using a detergent-compatible
(DC) protein assay (Pierce^{TM} BCA Protein Assay Kit) with bovine serum albumin (BSA) as a
standard. 90 µg of total protein lysate were loaded onto SDS-PAGE 4- 12% Bis-Tris gels
(Invitrogen); an Ab rabbit anti-CPSF6 (1:500) and an anti-rabbit HRP-conjugated (1:5000)
were used for the detection of CPSF6, whereas the normalization was done by an Ab anti-
actin HRP-conjugated (1:3000). Visualization was carried out using an ECL solution .

AUMsilence^{TM} ASO

AUMsilenceTM 352 ASOs were synthesized by AUM BioTech, LLC (Philadelphia, USA). THP-
1 negative control, THP-1 KD SRRM2 and THP-1 KD SON cells were differentiated with PMA
(160 nM) for 48 h then incubated 72 h with 10 µM of a AUMsilenceTM ASOs complementary
to the mRNA of SRRM2 and SON, respectively (scramble control AUMscrambleTM). All cells
were kept in incubator at 37°C and 5% CO2. These cells were cultivated in RPMI 1640 medium
supplemented with 10% fetal bovine serum (FBS) and 1% penicillin–streptomycin solution (100
U/ml).

Cloning

pSICO CPSF6-mNeonGreen was used to generate deletion mutants by exceeding different
regions from the original sequence according to Table2. The mutants were produced with and
without the mNeonGreen tag, except for pSICO CPSF6-ΔMCD and pSICO CPSF6-ΔMCD PY
NLS. All primers are specified in Table1.

For pSICO CPSF6-mNeonGreen ΔLCR ADD2, CPSF6 ΔLCR ADD2 sequence was amplified
by PCR from pLPCX CPSF6 ADD2 and specific primers were designed to add also BamHI
restriction sites at the extremities. Phusion Flash High-Fidelity PCR Master Mix was used and
the reaction was performed according to the manufacture datasheet (100 ng template DNA
and 56° annealing temperature). PCR products were treated with DpnI for 1h at 37°C and then
digested with BamHI restriction enzyme at 37°C for 1h. 2 ul of the backbone pSICO CPSF6-
mNeonGreen were digested with BamHI at 37°C for 3h and further treated with CIP. After gel
extraction of the DNA and purification, insert and backbone were ligated with T4 Ligase for 2
h at 22°C. 5 ul of the product were used to transform 50 ul of DH5α bacteria (30 min at 4°C,
45 sec at 42°C, 2 min at 4°C, incubation in SOC medium for 1h at 37°C and plating on LB agar
dishes).

All the other mutants were obtained using In-Fusion®Snap Assembly protocol, which allowed
to amplify the original plasmid deleting the small region of interest. The different primers used
are reported in Table1. As reported in the In-Fusion®Snap Assembly protocol

, 5 ng of the original plasmid were amplified with PrimeSTARMax

DNA polymerase in 35 PCR cycles (10 sec at 98°C, 15 sec at 55°C, 5sec/kb at 72°C). The

PCR products were then digested for 1h at 37°C using DpnI to get rid of the original plasmid.

After plasmid cleaning up, it was circularized through a ligation of 15 min at 50°C. 2.5 ul of

products were used in Stellar™ Competent Cells' transformation, following the same

procedure used for DH5α transformation, previously explained.

Lentiviral vectors and viral productions

LVs and HIV-1 viruses were produced by transient transfection of HEK293T cells through

calcium chloride co-precipitation. Co-transfection was performed as follows: for LVs, 10 µg of

transfer vector, 10 µg of packaging plasmid (gag-pol-tat-rev), and 2.5 µg of pHCMV-VSV-G

envelope plasmid; for VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) -vpx viruses and VSV-G/pNL4.3

Δenv ΔNef IRES GFP-vpx, 10 µg HIV-1ΔEnvIN$_{HA}$ LAI (BRU) plasmid or pNL4.3 Δenv ΔNef

IRES GFP or pNL4.3 ΔenvΔRΔNef Luc, 2.5 µg of pHCMV-VSV-G plasmid and 3 ug of SIVMAC

vpx (Durand et al., 2013) or full length virus 10 µg HIV-1 NL4.3AD8 and 3 ug of SIVMAC vpx.

After the collection of the supernatant 48 h post-transfection, lentiviral particles were

concentrated by ultracentrifugation for 1 h at 22000 rpm at 4°C and stored at −80°C. LVs and

viruses were tittered by qPCR in HEK293T cells 3 days post-transduction.

Cell transduction and infection

THP-1 ctrl CRISPR clone 2 cells and THP-1 (duplex1-2-3 CRISPR) KO clone 4 cells were

differentiated with PMA (160 nM) for 72 h then transduced with different mutants of CPSF6 for

72h (MOI=1) and then infected for 30h with VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) -vpx (MOI=10)

in presence of Nevirapine (10uM). The medium was always supplemented with PMA (160 nM)

.

For Halo tagged SRRM2 HEK 293 cells and Halo tagged SRRM2 ΔIDR HEK 293 cells, 2×10^5

cells were seeded on coverslips coated with polylysine in complete growth medium (DMEM,

GlutaMAX^{TM}-I, 10% FBS, and 1% P/S) and incubated at 37°C (5% CO_2) for 24 h. Cells were

then infected with the VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) (MOI 10) in complete growth medium

supplemented with Nevirapine (10μM) for 24h.

THP-1 control (scramble), THP-1 KD SRRM2 Cells and THP-1 KD SON cells were seeded on

coverslips and differentiated with PMA (160 nM) for 72 h. Then incubated 48 h with 10 µM of

a FANA ASOs (scramble control FANA (SCR-FANA), SRRM2-FANA and SON-FANA). ASOs

used in this study were designed and synthesized by AUM LifeTech (Philadelphia, PA, USA).

Next, cells were infected with the pNL4.3 Δenv ΔNef IRES GFP-VPX (MOI 25) in complete
growth medium and incubated at 37°C in 5% CO_2 for 4 days.

In the four timepoints experiment, THP-1 cells were differentiated with PMA (160 nM) for 72 h.
Cells were then infected with VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) -vpx (MOI=10) in presence
of Nevirapine (10uM), in complete growth medium supplemented with PMA (160 nM), and
incubated at 37°C in 5% CO_2 for 6 h, 9 h, 12 h, 30 h post infection.

THP-1 infection with HIV-1 replication competent virus and viral RNA real-time
quantification.

THP-1 cells differentiate into macrophage-like cells within 48 h under phorbol 12-myristate 13-
acetate (PMA) treatment (160 nM). THP1 cells were cultured at 37 °C in 5% CO2 in RPMI
Medium 1640 supplemented with GlutaMAXTM-I, 10% FBS, 1% P/S (100 U/mL), MEM Non-
essential Amino Acid Solution (0.1 mM), Sodium Pyruvate (1 mM), HEPES buffer solution (10
mM). THP1 cells differentiated into macrophages are infected with HIV-1 NL4.3AD8 with Vpx
(MOI 6). 24 hours after infection, the cells were washed and RPMI is added to the cells. Three
days post- infection, the supernatant is collected and centrifuged. Total viral RNA was extracted
from the supernatant via the QIAamp Viral RNA Mini Kit (Qiagen, Hilden Germany) following
the manufacturer's protocol. Total viral RNA was eluted into 60 µL of DEPC. vRNA is reverse
transcribed into DNA with the Maxima H minus reverse transcriptase (Thermoscientific,
EP0752) following the manufacturer's protocol and the following primers were used (forward:
GCCTCAATAAAGCTTGCCTTGA, reverse: TGACTAAAAGGGTCTGAGGGATCT). DNA
copies for each sample were determined by qPCR and the SYBR green dye. The absolute
DNA copies were calculated from a standard curve run in parallel using HIV-1 NL4.3AD8
plasmid. The qPCR cycle was as follows: 15 min at 95 °C, 40 cycles (15 s at 95 °C, 30 s at 58
°C, 30 s at 72 °C), 15 s at 95 °C, 15 s at 55 °C, 20 min melting curve until reaching 95 °C, 15
s at 90 °C.

Immunofluorescense Microscopy

Immunostaining

On the day of fixation, the cells were washed with PBS and fixed with 4% PFA for 15 min. Cells
were treated with glycine 0.15% for 10 min, permeabilized with 0.5% Triton X-100 for 30 min,
and blocked with 1% BSA for 30 min. All antibody incubations were carried out at room
temperature in a dark humid chamber, for 1 h with primary antibodies and for 45 min with
secondary antibodies. Washes between antibody incubations and antibody dilutions were
done in 1% BSA.

Primary antibodies were diluted as follows: anti-HA 1:500, anti-CPSF6 1:400, anti-SC35 1:200,

anti-SON 1:200, anti-SRRM2 1:200, Secondary antibodies used were goat anti-rabbit Alexa-

488 1:300 and donkey anti-Rabbit Cy3 1:1000 for CPSF6, SON and SRRM2, goat anti-rat

Alexa-647 1:100 for IN-HA, goat anti-mouse Alexa-647 1:300 or goat anti-mouse Alexa-488

1:300 for SC35, donkey anti-mouse Alexa-647 1:300 for SRRM2.

Finally, cells were stained with Hoechst 33342 1:10000 for 5 min. Coverslips were mounted

on glass slides (Star Frost) with Prolong Diamond Antifade Mountant.

HaloTag Labelling

To detect the HaloTag in Halo tagged SRRM2 HEK 293 cells and Halo tagged SRRM2 ΔIDR

HEK 293 cells, we used HaloTag®TMR Ligand following the Technical Manual available at

www.promega.com/protocols. Specifically, cells were incubated in DMEM supplemented with

HaloTag®TMR Ligand (5 µM) for 15 min at 37°C. The ligand-containing medium was then

removed and replaced with an equal volume of 1X PBS, repeating the step twice and ending

with warm complete medium. Cells were incubated in complete culture medium for 30 min at

37°C. The medium was then removed and replaced with an equal volume of fresh warm culture

medium.

Immuno-RNA FISH

On the day of fixation, fixed cells were incubated in Permeabilization/Blocking buffer (1% BSA,

0.3% Triton X-100, 2mM Vanadyl Ribonucleoside complexes (VRCs) in RNase-free PBS) for

1 h before the antibodies' incubations. Antibodies were diluted in Permeabilization/Blocking

buffer. After the primary and secondary antibody staining (respectively of 1h and 45 min) with

the respective washes, cells were fixed for a second time in PFA 4% (in RNase-free PBS) for

10 min at room temperature with subsequent washes with RNase-free PBS. In the meantime,

40 pmol of primary smiFISH probes (Tsanov et al., 2016) (24 smiFISH probes designed against

HIV-1 pol sequence (Rensen et al., 2021) were hybridized with 50 pmol of secondary FLAP

probe   conjugated   to   a   Cy5   fluorophore

(Cy5/AATGCATGTCGACGAGGTCCGAGTGTAA/Cy5Sp/) in 1X NEBuffer 3 (diluted in RNase-

free H_2O) using a thermocycler. The program setting follows: 3 min at 85°C, 3 min at 65°C and

5 min at 25°C. FISH-probe solution was then diluted 1:50 in Hybridization buffer (90%

Stellaris® RNA-FISH Hybridization Buffer, 10% Deionized Formamide). After the samples were

washed in Wash A buffer (20% Stellaris® RNA-FISH Wash Buffer A, 10% Deionized

Formamide, in RNase-free H_2O) at room temperature for 5 min, they were placed on parafilm,

covered with 50 ul of FISH-probe solution in Hybridization buffer and incubated overnight at

37°C. The next day, cells were washed with Wash A buffer in the dark at 37°C for 30 min.

Afterwards, the samples were incubated for 10 min in Hoechst 333342 diluted 1:10000 in

RNase-free H_2O and then washed with Wash B buffer for 5 min at room temperature in the

dark. Finally, the cells were washed in RNase-free H_2O before the coverslips were mounted

on microscopy slides using ProLongTM Diamond Antifade mounting medium. The mounting

medium was cured overnight at room temperature under the chemical hood and away from

light.

Images acquisition

Images were acquired using a Zeiss LSM700 confocal inverted microscope, with a 63×

objective (Plan Apochromat, oil immersion, NA = 1.4), using diode lasers at 405 nm, 488 nm,

555 nm and 639 nm for the respective fluorophores. A pixel size of 0.07 μm was used.

Live imaging on HIV-1 infected Halo tagged SRRM2 HEK 293 cells

Halo tagged SRRM2 HEK293 cells were seeded and transduced with CPSF6-mNeonGreen

lentiviral vector (MOI 0.5) for 24 h. 0.3×10^6 Cells were then transferred on poly-L-Lysin coated

Ibidi-dishes and infected with VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) (MOI = 40) in presence of

Nevirapine (10 uM) for 2 h. After having changed the medium and labeled with HaloTag®TMR

Ligand (5 μM) (see previous "HaloTag labelling" section) and Hoechst 333342 diluted 1:80000

in complete DMEM medium, 4D movies were acquired.

3D movies were acquired using a Nikon Ti2-E Confocal Inverted Spinning Disk microscope,

using a 63x objective (Plan Apochromat, oil immersion, NA = 1.4) and a sCMOS Hamamatsu

camera, Orca Flash 4. Pixel size 6.5 μm, 2048 x 2044 pixel, QE 82%. For the Z stuck a Z piezo

stage was used, with 0.33 μm interval. For live imaging acquisitions, cells were placed in an

environmental chamber with 37°C temperature, 5% CO_2 and 21% O_2.

Protein expression and purification.

pET-11a vectors were used to express the HIV-1 capsid protein. Point mutations, A14C and

E45C were introduced using the QuikChange II site-directed mutagenesis kit (Stratagene)

according to the manufacturer's instructions. All proteins were expressed in *Escherichia coli*

one-shoot BL21star™ (DE3) cells (Invitrogen). Briefly, LB medium was inoculated with

overnight cultures, which were grown at 30°C until mid log-phase (A_{600}, 0.6-0.8). Protein

expression was induced with 1 mM isopropyl-β-d-thiogalactopyranoside (IPTG) overnight at

18°C. Cells were harvested by centrifugation at 5,000 x g for 10 min at 4°C, and pellets were

stored at −80°C until purification. Purification of capsid was carried out as follows. Pellets from

2 L of bacteria were lysed by sonication (Qsonica microtip: 4420; A=45; 2 minutes; 2 seconds

on; 2 seconds off for 12 cycles), in 40 ml of lysis buffer (50 mM Tris pH=8, 50mM NaCl, 100mM

β-mercaptoethanol and Complete EDTA-free protease inhibitor tablets). Cell debris were

removed by centrifugation at 40,000g for 20 min at 4°C. Proteins from the supernatant were

precipitated by incubation with 1/3 of volume of saturated ammonium sulfate containing 100

mM β-mercaptoethanol for 20 min at 4°C and centrifugation at 8,000g for 20 min at 4°C.

Precipitated proteins were resuspended in 30 ml of buffer A (25mM MES pH6.5, 100 mM β-

mercaptoethanol) and sonicated 2-3 times (Qsonica microtip: 4420; A=45; 2 minutes; 1 second

on; 2 seconds off). Sample was dialyzed 3 times in buffer A (2 h, overnight, 2 h). The sample

was sonicated and diluted in 500 ml of buffer A and was chromatographed sequentially on a 5

ml HiTrap™ Q HP column and on a 5 ml HiTrap™ SP FF column (GE Healthcare), both pre-

equilibrated with buffer A. The capsid protein was eluted from HiTrap™ SP FF column using a

linear gradient from 0-2 M of NaCl. Absorbance at 280 nm was checked to take the eluted

fraction that had higher protein levels. Pooled fractions were dialyzed 3 times (2 h, overnight,

2 h) in storage buffer (25mM MES, 2M NaCl, 20 mM β-mercaptoethanol). Sample was

concentrated using centricons to a concentration of 20 mg/ml and stored at -80°C.

Assembly of stabilized HIV-1 capsid tubes

1 mL of monomeric capsid (3 mg/mL or 1 mg/mL) was dialyzed in SnakeSkin dialysis tubing

10,000 MWCO (Thermo Scientific) against a buffer that is high in salt and contains a reducing

agent (buffer 1: 50 mM Tris, pH 8, 1 M NaCl, 100 mM β-mercaptoethanol) at 4°C for 8 hours.

Subsequently the protein was dialyzed against the same buffer without the reducing agent β-

mercaptoethanol (buffer 2: 50 mM Tris, pH 8, 1 M NaCl) at 4°C for 8 hours. The absence of β-

mercaptoethanol in the second dialysis allows formation of disulfide bonds between Cysteine

14 and 43 inter-capsid monomers in the hexamer. Finally the protein is dialyzed against buffer

3 (20 mM Tris, pH 8,0, 40 mM NaCl) at 4°C for 8 hours. Assembled complexes were kept at

4°C up to 1 month.

Capsid binding assay protocol

Human HEK293T cells were transfected for 24 h with a plasmid expressing the specified

CPSF6 variant tagged with mNeonGreen. Cell media was completely removed and cells were

lysed in 300 μL of capsid binding buffer (CBB: 10 mM Tris, pH 8,0, 1,5 mM MgCl2, 10 mM

KCl) by scrapping off the plate. Cells were rotated at 4°C for 15 min and then centrifuged to

remove cellular debris (21,000 *x g* ,15 minutes, 4 °C). Cell lysates were incubated with

stabilized HIV-1 capsid tubes for 1 h at 25 °C. Subsequently, stabilized HIV-1 capsid tubes

were washed by pelleting the complexes by centrifugation at 21,000 x g for 2 min. Pellets were

washed using by resuspension in CBB or PBS. Pellets were resuspended in Laemmli buffer

1X and analyzed by Western blotting using anti-p24 or anti-mNeonGreen antibodies.

Bioinformatics analysis of CPSF6

Intrinsic disorder propensity of CPSF6 was evaluated using the Rapid Intrinsic Disorder

Analysis Online platform (RIDAO) (https://ridao.app/) designed to predict disordered residues

and regions in a query protein based on its amino acid sequence (Dayhoff and Uversky, 2022).

RIDAO yields results by combining the outputs of several commonly used per-residue disorder

predictors, such as PONDR® VLXT (Romero et al., 2001), PONDR® VL3(Peng et al., 2006),

PONDR® VLS2B(Peng et al., 2005), PONDR® FIT (Xue et al., 2010), as well as IUPred2

(Short) and IUPred2 (Long)(Dosztanyi et al., 2005a, b). RIDAO also computes a mean disorder

score for each residue based on these. In the resulting intrinsic disorder profile, the disorder

score of 0.5 is the threshold between order and disorder, where residues/regions above 0.5

are disordered, and residues/regions below 0.5 are ordered. The disorder score of 0.15 is the

threshold between order and flexibility, where residues/regions with the disorder scores above

0.15 but below 0.5 are flexible, and residues/regions below 0.15 are highly ordered.

Amino acid compositions of the intrinsically disordered C-terminal domain (residues 261-358)

of human CPSF6 and its different variants (CPSF6 ΔFG ΔLCR, CPSF6 ΔLCR, CPSF6 ΔFG,

and CPSF6 ADD2 ΔLCR) were analyzed to evaluate the relative abundance of prion-like low

complexity region (LCR) defining uncharged, charged, and Pro residues in these protein

regions. The corresponding values of the relative abundance of these residue groups were

calculated by dividing numbers of prion-like LCR defining uncharged (Ala, Gly, Val, Phe, Tyr,

Leu, Ile, Ser, Thr, Pro, Asn, Gln), charged (Asp, Glu, Lys, Arg), and Pro residues by total

number of amino acids in the corresponding protein fragments. As references, we used the

corresponding data for protein sequences deposited to the UniProtKB/Swiss-Prot database

that provides information on the overall distribution of amino acids in nature (Bairoch et al.,

2005); PDB Select 25 (Berman et al., 2000), which is a subset of structures from the Protein

Data Bank with less than 25% sequence identity, biased towards the composition of proteins

amenable to crystallization studies; and DisProt 3.4 that is comprised of a set of consensus

sequences of experimentally determined disordered regions (Sickmeier et al., 2007). Per-

residue intrinsic disorder propensities of the LCR-FG and ADD2-FG sequences were

evaluated by PONDR® VLXT (Romero et al., 2001), which is sensitive to local peculiarities of the amino acid sequences freely available at http://www.pondr.com/ (accessed on August 03, 2024). Linear distribution of the net charge per residue (NCPR) within the LCR-FG and ADD2-FG sequences were evaluated by CIDER (Holehouse et al., 2017), which is a webserver for the analysis of a wide range of the physicochemical properties encoded by IDP sequences freely available at http://pappulab.wustl.edu/CIDER (accessed on August 03, 2024). Secondary structure propensities of the LCR-FG and ADD2-FG sequences were evaluated by PSIPRED (McGuffin et al., 2000), which is a highly accurate secondary structure prediction method freely available to non-commercial users at http://globin.bio.warwick.ac.uk/psipred/ (accessed on August 03, 2024).

IMAGING AND STATISTICAL ANALYSIS

All images were analysed using Fiji Software. More in detail, for the count of the nuclear speckles a macro was computed to segment cellular nuclei and to select and count the nuclear speckles. For the nuclei segmentation a Gaussian Blur with sigma=2 was used and for the channel related to the nuclear speckles the set threshold was at 7000-46012 (min and max). For the counting, a "size =10 – Infinity summarize" was used for the nuclei, whereas for the nuclear speckles the size was reduced to "size=0-Infinity summarize".

For live imaging analysis, Arivis software was used to reconstruct the 3D movies.

All data were statistically analyzed with GraphPad Prism 9 (GraphPad Software, La Jolla California USA, www.graphpad.com). Calculations were performed and figures were drawn using Excel 365 or GraphPad Prism 8.0. Statistical analysis was performed, with Wilcoxon matched paired t-tests or Mann–Whitney unpaired t-tests. Spearman correlation coefficients (r) were calculated using GraphPad Prism.

Figure legends

Figure 1. Role of HIV-induced CPSF6 puncta in the nuclear reverse transcription upon removal of NVP. A) A) THP-1 cells, infected with VSV-G Δenv HIV-1 (NL4.3) ΔR LUC (MOI 10) in presence or not of Nevirapine (10 μM) for 5 days, or in presence of Nevirapine (10 μM) for 2 days and then the remaining 3 days without drug or in presence of Nevirapine (10 μM) for 2 days then in presence of PF74 (25 μM). Confocal microscopy images, to verify the presence of CPSF6 puncta, the cells are stained with anti-CPSF6 antibody (green). Nuclei are

stained with Hoechst (blue). Scale bar 10 µm. **B)** Luciferase Assay, to verify luciferase expression in the aforementioned samples. Luciferase values were normalized by total proteins revealed with the Bradford kit. One-way ANOVA statistical test with multiple comparison was performed (****=$p<0.0001$; *=$p<0.05$; ns=$p>0.05$). Data are representative of two independent experiments. **C)** Western blots demonstrate CPSF6 depletion using a specific antibody against CPSF6 in THP-1 cells subjected to CRISPR Cas9 methods: CRISPR Cas9 bulk (left), and CRISPR Cas9 clones selected by limiting dilution (right). Each condition is normalized for actin labeling. The ratio between the intensity signal of CPSF6 and actin was analyzed via ImageJ and is plotted below each western blot. **D)** Confocal microscopy images of THP-1 ctrl CRISPR clone 2 cells (Ctrl 2) and THP-1 duplex1-2-3 CRISPR clone 4 cells (CPSF6 KO 4) infected with VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) -vpx (MOI 10) in the presence of Nevirapine (10 µM). The cells are stained 30 h p.i. with anti-CPSF6 antibody and anti-HA antibodies to detect HA tagged integrase (IN).

Figure 2. Role of CPSF6 Domains in HIV-Induced CPSF6 Puncta. A) Schema of CPSF6 isoform 588aa deletion mutants. **B)** Confocal microscopy images of THP-1 CPSF6 KO cells, transduced with different mutants of CPSF6, infected with VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) -vpx (MOI 10) in presence of Nevirapine (10 µM). The cells are stained with CPSF6 and HA antibody 30 h p.i.. Scale bar 5µm. **C)** Analysis of the number of CPSF6 puncta in THP-1 CPSF6 KO cells transduced with different mutants of CPSF6, not infected or infected in presence of Nevirapine (10 µM) (the number of analyzed cells is shown under the x-axis). Statistical test: ordinary one-way ANOVA (****=$p<0.0001$; ***=$p<0.001$; *=$p<0.05$; ns=$p>0.05$). **D)** The plot compares the number of CPSF6 puncta per cell in THP-1 CPSF6 KO cells transduced with different mutants of CPSF6, infected with HIV-1 in the presence of Nevirapine (10 µM). Statistical test: ordinary one-way ANOVA (****=$p<0.0001$; ns=$p>0.05$). **E)** Confocal microscopy images of THP-1 CPSF6 KO clone 4, non-transduced and non-infected or transduced with WT CPSF6 and CPSF6 3xNLSΔMCD and infected with VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) -vpx (MOI 10) in presence of Nevirapine (10 µM). Immuno-RNA FISH: the cells are stained with CPSF6 (green) antibody and with 24 probes against HIV-1 Pol sequence (gray)(RNA-FISH) 25 h p.i. Nuclei are stained with Hoechst (blue). Scale bar 10 µm. Violin plot presenting the percentage of CPSF6 puncta colocalizing with the viral RNA in THP-1 CPSF6 KO clone 4 cells transduced with LVs expressing CPSF6 WT or CPSF6 3xNLSΔMCD (respectively n=73 and n=103) and infected with VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) -vpx (MOI 10) in presence of Nevirapine (10 µM). A total of 198 CPSF6 WT puncta and 264 CPSF6 3xNLSΔMCD puncta were counted. Statistical test: unpaired t-test, ns=$p>0.05$. **F)** Confocal microscopy images of THP-1 KO CPSF6 cells transduced with WT CPSF6 and CPSF6 ΔMCD without NLS, with

3xNLS or with PY NLS, respectively. Cells were differentiated for 3 days, transduced with
CPSF6 lentiviral vectors (MOI 1) for 3 days and infected for 24 h with VSV-G / HIV-1ΔEnvIN$_{HA}$
LAI (BRU) -vpx (MOI 10) in presence of Nevirapine (10 uM) (left panels). The panels on the
right show transduced and uninfected cells. CPSF6 and the IN tagged with the HA are labeled
with anti-CPSF6 (green) and anti-HA (white) antibodies, respectively. Nuclei are stained with
Hoechst (blue). The arrows show CPSF6 puncta in colocalization with IN-HA. Scale bar 10μm.

Figure 3. Evaluation of CPSF6 Deletion Mutants' Binding Capacity to the Viral Core. A)
Ability of wild type and mutant CPSF6 proteins to bind to the HIV-1 core. Cellular extracts
derived from human 293T cells expressing similar levels of the indicated CPSF6 proteins
(INPUT) were incubated with HIV-1 capsid stabilized tubes for 1 hour at room temperatures in
the presence and absence of 10 μM PF74, as described in materials and methods. As a carrier
control, we utilized DMSO. Subsequently, HIV-1 capsid stabilized tubes were washed, and the
bound proteins eluted 1X Laemmli buffer 1X. The BOUND fractions were analysed by Western
blotting using antibodies against neon-GFP and the HIV-1 capsid. **B)** Experiments were
repeated at least three times and the average BOUND fraction relative to the INPUT fraction
normalized to wild type binding are shown for the different CPSF6 mutants.   *** indicates a p-
value < 0.001; **** indicates a p-value < 0.0001; and ns indicates no significant difference as
determined by unpaired t-tests.

Figure 4. Comparison of second structures of ADD2 and LCR. A) Physicochemical
characteristics of the LCR-FG and ADD2-FG sequences. Intrinsic disorder predispositions
evaluated by PONDR® VLXT. Position of the FR segment within the LCR-FG and ADD2-FG
sequences is shown as gray shaded area. **B)** Linear distribution of the net charge per residue
(NCPR) within the LCR-FG sequence evaluated by CIDER. **C)** Linear distribution of the net
charge per residue (NCPR) within the ADD2-FG sequence evaluated by CIDER. **D)** Secondary
structure propensity of the LCR-FG sequence evaluated by PSIPRED. **E)** Secondary structure
propensity of the ADD2-FG sequence evaluated by PSIPRED. **F)** Analysis of the peculiarities
of the amino acid compositions of the intrinsically disordered C-terminal domain (residues 261-
358) of human CPSF6 and its different mutants. Relative abundance of prion-like LCR defining
uncharged residues in analyzed protein segments. **G)** Relative abundance of proline residues
in analyzed protein segments. **H)** Relative abundance of charged residues in analyzed protein
segments. The values were calculated by dividing numbers of prion-like LCR defining
uncharged (Ala, Gly, Val, Phe, Tyr, Leu, Ile, Ser, Thr, Pro, Asn, Gln), Pro, and charged (Asp,
Glu, Lys, Arg) residues by the total number of amino acids in the respective protein fragments.

Corresponding values for all protein sequences deposited in the UniProtKB/Swiss-Prot
database, PDB Select25, and DisProt are shown for comparison.

Figure 5. Role of FG motif in viral replication. A) WB showing CPSF6 protein from several
single clones derived from CPSF6 KO clone obtained upon complementation with CPSF6 ΔFG
and normalized with beta-actin. **B)** Quantification of the expression of CPSF6 ΔFG protein in
different single clones compared to CPSF6 WT (value 1). **C)** Infectivity essay using a single
round infectious virus carrying the cDNA of Luciferase as reporter gene. Values are expressed
as% of RLU compared to WT cells. **D)** Infectivity essay of a replication competent virus: vRNA
from new viruses produced after infection of WT, CPSF6 KO and CPSF6ΔFG cells was
analyzed and showed in the histograms as % of vRNA copies compared to vRNA in WT THP-
1 cells considered 100%.

Figure 6. Depletion of MCD or LCR does not affect the formation of HIV-induced CPSF6
puncta. A) Epifluorescence microscopy images of both infected and non-infected
differentiated THP-1 cells showing the presence of CPSF6 puncta only in the infected
condition. CPSF6 and SC35 are labelled with anti-CPSF6 (green) and anti-SC35 (red)
antibodies, respectively. Nuclei are stained with Hoechst (blue). Scale bar 10μm. **B)** Confocal
microscopy images of THP-1 KO CPSF6 cells, differentiated for 3 days, transduced with
CPSF6 lentiviral vector (MOI 1) (specifically WT CPSF6, CPSF6 ΔLCRs, CPSF6 ΔMCD with
3xNLS, without NLS, or with PY NLS) for 3 days and infected for 24 h with VSV-G / HIV-
1ΔEnvIN$_{HA}$ LAI (BRU) -vpx MOI 10) in presence of Nevirapine (10 uM). CPSF6 and nuclear
speckles were labelled with anti-CPSF6 (green) and anti-SC35 (red) antibodies, respectively.
Nuclei are stained with Hoechst (blue). Scale bar 10μm. The percentage of CPSF6 puncta
associated with SC35 per field of view is shown in the graph. N cells were counted in each
condition and a one-way ANOVA statistical test with multiple comparison was performed; ns=
p value > 0.05.

Figure 7. Dynamics of the HIV-induced CPSF6 puncta formation and their fusion with
NSs. A) Time course of infection of THP-1, 6h.p.i., 9h.p.i., 12h.p.i., 30h.p.i. or non infected.
Cells were stained with antibodies against CPSF6 (green) and SC35 (red). **B)** The graph
shows the percentage of CPSF6 puncta associated with NS or adjacent to NS or isolated from
NS at different time post-infection. **C)** The graph shows the progression of CPSF6 puncta

associated to NS during the time post-infection. N indicates the number of cells analyzed. One-way ANOVA statistical test with multiple comparison was performed; ns= p value > 0.05; **** indicates a p-value < 0.0001.

Figure 8. Role of SRRM2 and SON in the Formation of HIV-Induced CPSF6 puncta. A) Depletion of SON and SRRM2 in THP-1 cells using AUM*silence*™ ASO technology. The degree of depletion is quantified by WB and the mean intensity through immunofluorescence using antibodies against SON and SRRM2, respectively. Scale bar 5µm. **B)** The percentage of CPSF6 puncta formation is quantified by IF in THP-1 cells knocked down for SON, SRRM2, and control (Ctrl) infected with HIV-1 (MOI 25) for 48 h. CPSF6 is stained with an antibody against CPSF6 (green), and nuclei are stained with Hoechst (blue). The graph on the right reports the percentage of CPSF6 puncta calculated from more than 100 cells. Scale bar 10µm. Experiments were performed at least twice. **C)** (Top panels) confocal microscopy images of ΔIDR HaloTag SRRM2 HEK293 and HaloTag SRRM2 HEK293 cells stained with the halo tag ligand (red), and nuclei (blue). Scale bar 10µm. (Bottom panels) confocal microscopy images of HaloTag SRRM2 HEK293 and ΔIDR HaloTag SRRM2 HEK293 cells, both labeled with anti-SRRM2 (red) and anti-SON (gray) antibodies. Nuclei are stained with Hoechst (blue). Scale bar 10µm. Statistical studies are summarized in the violin plot which displays the distribution of the number of SON puncta per cell in the two conditions. N cells were counted and Kolmogorov Smirnov test was performed, ns= p>0.05. **D)** Confocal microscopy images of HaloTag SRRM2 HEK293 and ΔIDR HaloTag SRRM2 HEK293 cells, either non-infected or infected for 24 h with VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) (MOI 10) in the presence of Nevirapine (10 µM). CPSF6 and SC35 are labeled with anti-CPSF6 (red) and anti-SC35 (gray) antibodies, respectively. Nuclei are stained with Hoechst (blue). The plot shows the mean ± SD of the percentage of cells with CPSF6 puncta calculated in n fields of view (n=24, 29, 32); N is the number of cells analyzed for each of the three different cell lines; an unpaired t-test was performed, ****=p<0.0001, ns=p>0.05. Scale bar 10µm. Experiments were performed at list twice.

Movies 1 A and B. CPSF6 MLOs can form independently from SRRM2 MLOs in HEK293 SRRM2 HaloTag cells. Spinning Disk confocal images of HEK293 SRRM2 HaloTag cells acquired 9 hours after VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) (MOI 10) infection, occurred in presence of Nevirapine (10 µM). Cells were previously transduced for 24h with CPSF6-mNeonGreen lentiviral vector (MOI 0.5) to detect CPSF6 (visualized in green). SRRM2 MLOs were detected using the TMR-HaloTag Ligand (in red) and the nuclei were stained with

896 Hoechst (blue). Acquisitions were performed continuously for 27 minutes, which allowed to see

897 the independent formation of CPSF6 puncta, their migration toward the SRRM2 MLOs and the

898 final fusion of CPSF6 puncta and SRRM MLOs.

Supplementary Figures

Suppl. Figure 1. A) Multiple examples of confocal microscopy images of THP-1 ctrl CRISPR

clone 2 cells and THP-1 CPSF6 KO 4 cells infected with VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) -

vpx (MOI 10) in the presence of Nevirapine (10 µM). The cells are stained 30 h p.i. with CPSF6

(green) and HA (red) antibodies to detect integrase (IN). **B)** Multiple examples of THP-1 CPSF6

KO clone 4 cells transduced with different LVs carrying CPSF6 WT or mutants and stained

with CPSF6 and HA antibody 30 h p.i.. Scale bar 5µm.

Suppl. Figure 2. A) Per-residue intrinsic disorder propensity of the CPSF6 isoform 588 aa

evaluated by the Rapid Intrinsic Disorder Analysis Online platform (RIDAO) (Dayhoff and

Uversky, 2022) that yields results for IU-Pred_short (yellow line), IUPred_long (blue line),

PONDR® VL3 (green line), PONDR® VLXT (black line), PONDR® VSL2 (red line), and

PONDR® FIT (pink line) and computes a mean disorder score for each residue based on these

predictors (MDP, thick, dark pink, dashed line). Light pink shadow represents MDP error

distribution. The thin black line at the disorder score of 0.5 is the threshold between order and

disorder, where residues/regions with disorder scores above 0.5 are disordered, and

residues/regions below 0.5 are ordered. The dashed line at the disorder score of 0.15 is the

threshold between order and flexibility, where residues/regions above 0.15 are flexible, and

residues/regions below 0.15 are highly ordered (upper). Schema of the deletion mutants of

CPSF6 (bottom). **B)** Lentiviral Vector Transduction of PMA-Differentiated THP-1 Cells

expressing CPSF6 ΔMCD fused to mNeonGreen (left), CPSF6 NLSΔMCD fused to

mNeonGreen (center), CPSF6 3xNLSΔMCD fused to mNeonGreen (right). CPSF6 is

represented in green, and nuclei are stained in blue. Scale bar 5µm.

Suppl. Figure 3. Sequences of FG and LCRs or substituted amino acids sequences analyzed

in figure 4.

Suppl. Figure 4. A) THP-1 infected with VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) -vpx were labelled

after 6h, 9h, 12h, 30h p.i. with specific antibodies against CPSF6 (green) and SC35 (red),

nuclei are stained with Hoechst (blue). Scale bar 10µm. **B)** Timelapse of Movie B which

recapitulates the key steps of the dynamics of HIV induced-CPSF6 puncta biogenesis. CPSF6

was visualized transducing HEK293 SRRM2 HaloTag cells for 24h with CPSF6-mNeonGreen

lentiviral vector (MOI 0.5). SRRM2-HaloTag allowed the use of TMR-Halo Tag Ligand to

visualize the SRRM2 MLOs. After 9 hours after VSV-G / ΔEnvHIV-1 LAI (BRU) (MOI 10)

infection in presence of Nevirapine, CPSF6 puncta formed independently (9h 01m), migrating

afterwards to the SRRM2 MLOs and leading to a final fusion between CPSF6 puncta and
SRRM2 MLOs (9h14m).

Suppl. Figure 5. Multiple examples (**A-E**) of confocal microscopy images of HaloTag SRRM2
HEK293 and ΔIDR HaloTag SRRM2 HEK293 cells, either non-infected or infected for 24 h with
VSV-G / HIV-1ΔEnvIN$_{HA}$ LAI (BRU) (MOI 10) in the presence of Nevirapine (10 µM). CPSF6
and SC35 are labeled with anti-CPSF6 (red) and anti-SC35 (gray) antibodies, respectively.
Nuclei are stained with Hoechst (blue). Scale bar 10µm.

Suppl. Figure 6. SRRM2 depletion is confirmed using antibodies against both SRRM2 and
SC35 by IF. Statistical analysis: One-Way ANOVA (****=$p<0.0001$; *=$p<0.05$; ns=$p>0.05$).

Suppl. Figure 7. HEK 293 HaloTag SRRM2 or ΔIDR were labelled with Halo ligand (red) and
an anti-SON antibody (gray), nuclei are stained by Hoechst (blue). Scale bar 10µm.

ACKNOWLEDGMENTS

F.D.N. is supported by the Institut Pasteur and ANRS grants (ECTZ192036, ECTZ137593),
ANR-PRCI grant, Sidaction grant. C.T. is supported by fellowships Sidaction, ANR-PRCI. S.A.
is supported by ANRS fellowship ECTZ204694. F.D.-G., B.C., M.R., C.L., C.B. are supported
by NIH Grants R01AI087390 and R01AI150455. We gratefully acknowledge the UtechS
Photonic BioImaging platform (Imagopole), C2RT at Institut Pasteur. We thank the NIH AIDS
Reagents program to support us with precious reagents and Addgene.

Bibliography

Ay, S., Burlaud-Gaillard, J., Gazi, A., Tatirovsky, Y., Cuche, C., Diana, J.S., Scoca, V., Di Santo, J.P.,
Roingeard, P., Mammano, F*., et al.* (2025). In vivo HIV-1 nuclear condensates safeguard against cGAS
and license reverse transcription. EMBO J *44*, 166-199.
Ay, S., and Di Nunzio, F. (2023). HIV-Induced CPSF6 Condensates. J Mol Biol *435*, 168094.
Bairoch, A., Apweiler, R., Wu, C.H., Barker, W.C., Boeckmann, B., Ferro, S., Gasteiger, E., Huang, H.,
Lopez, R., Magrane, M*., et al.* (2005). The Universal Protein Resource (UniProt). Nucleic Acids Res *33*,
D154-159.
Barre-Sinoussi, F., Chermann, J.C., Rey, F., Nugeyre, M.T., Chamaret, S., Gruest, J., Dauguet, C., Axler-
Blin, C., Vezinet-Brun, F., Rouzioux, C*., et al.* (1983). Isolation of a T-lymphotropic retrovirus from a
patient at risk for acquired immune deficiency syndrome (AIDS). Science *220*, 868-871.
Berman, H.M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T.N., Weissig, H., Shindyalov, I.N., and
Bourne, P.E. (2000). The Protein Data Bank. Nucleic Acids Res *28*, 235-242.

Blanco-Rodriguez, G., and Di Nunzio, F. (2021). The Viral Capsid: A Master Key to Access the Host
Nucleus. Viruses *13*.

Blanco-Rodriguez, G., Gazi, A., Monel, B., Frabetti, S., Scoca, V., Mueller, F., Schwartz, O., Krijnse-
Locker, J., Charneau, P., and Di Nunzio, F. (2020). Remodeling of the Core Leads HIV-1 Preintegration
Complex into the Nucleus of Human Lymphocytes. J Virol *94*.

Buffone, C., Martinez-Lopez, A., Fricke, T., Opp, S., Severgnini, M., Cifola, I., Petiti, L., Frabetti, S.,
Skorupka, K., Zadrozny, K.K*., et al.* (2018). Nup153 Unlocks the Nuclear Pore Complex for HIV-1
Nuclear Translocation in Nondividing Cells. J Virol *92*.

Burdick, R.C., Li, C., Munshi, M., Rawson, J.M.O., Nagashima, K., Hu, W.S., and Pathak, V.K. (2020).
HIV-1 uncoats in the nucleus near sites of integration. Proc Natl Acad Sci U S A *117*, 5486-5493.

Chen, N.Y., Zhou, L., Gane, P.J., Opp, S., Ball, N.J., Nicastro, G., Zufferey, M., Buffone, C., Luban, J.,
Selwood, D*., et al.* (2016). HIV-1 capsid is involved in post-nuclear entry steps. Retrovirology *13*, 28.

Chen, Y., Zhang, Y., Wang, Y., Zhang, L., Brinkman, E.K., Adam, S.A., Goldman, R., van Steensel, B., Ma,
J., and Belmont, A.S. (2018). Mapping 3D genome organization relative to nuclear compartments
using TSA-Seq as a cytological ruler. J Cell Biol *217*, 4025-4048.

Dayhoff, G.W., 2nd, and Uversky, V.N. (2022). Rapid prediction and analysis of protein intrinsic
disorder. Protein Sci *31*, e4496.

Dharan, A., Bachmann, N., Talley, S., Zwikelmaier, V., and Campbell, E.M. (2020). Nuclear pore
blockade reveals that HIV-1 completes reverse transcription and uncoating in the nucleus. Nature
Microbiology *5*, 1088-1095.

Di Nunzio, F. (2013). New insights in the role of nucleoporins: a bridge leading to concerted steps
from HIV-1 nuclear entry until integration. Virus Res *178*, 187-196.

Di Nunzio, F., Danckaert, A., Fricke, T., Perez, P., Fernandez, J., Perret, E., Roux, P., Shorte, S., Charneau,
P., Diaz-Griffero, F*., et al.* (2012). Human nucleoporins promote HIV-1 docking at the nuclear pore,
nuclear import and integration. PLoS One *7*, e46037.

Di Nunzio, F., Uversky, V.N., and Mouland, A.J. (2023). Biomolecular condensates: insights into early
and late steps of the HIV-1 replication cycle. Retrovirology *20*, 4.

Dickson, C.F., Hertel, S., Tuckwell, A.J., Li, N., Ruan, J., Al-Izzi, S.C., Ariotti, N., Sierecki, E., Gambin, Y.,
Morris, R.G*., et al.* (2024). The HIV capsid mimics karyopherin engagement of FG-nucleoporins.
Nature.

Dosztanyi, Z., Csizmok, V., Tompa, P., and Simon, I. (2005a). IUPred: web server for the prediction of
intrinsically unstructured regions of proteins based on estimated energy content. Bioinformatics *21*,
3433-3434.

Dosztanyi, Z., Csizmok, V., Tompa, P., and Simon, I. (2005b). The pairwise energy content estimated
from amino acid composition discriminates between folded and intrinsically unstructured proteins. J
Mol Biol *347*, 827-839.

Durand, S., Nguyen, X.N., Turpin, J., Cordeil, S., Nazaret, N., Croze, S., Mahieux, R., Lachuer, J., Legras-
Lachuer, C., and Cimarelli, A. (2013). Tailored HIV-1 vectors for genetic modification of primary human
dendritic cells and monocytes. J Virol *87*, 234-242.

Francis, A.C., Marin, M., Singh, P.K., Achuthan, V., Prellberg, M.J., Palermino-Rowland, K., Lan, S.,
Tedbury, P.R., Sarafianos, S.G., Engelman, A.N*., et al.* (2020). Publisher Correction: HIV-1 replication
complexes accumulate in nuclear speckles and integrate into speckle-associated genomic domains.
Nat Commun *11*, 6165.

Fu, L., Weiskopf, E.N., Akkermans, O., Swanson, N.A., Cheng, S., Schwartz, T.U., and Görlich, D. (2024).
HIV-1 capsids enter the FG phase of nuclear pores like a transport receptor. Nature *626*, 843-851.

Fu, X.D., and Maniatis, T. (1990). Factor required for mammalian spliceosome assembly is localized to
discrete regions in the nucleus. Nature *343*, 437-441.

Gao, L., Zhang, W., Zhang, L., Gromova, B., Chen, G., Csizmadia, E., Cagle, C., Nastasio, S., Ma, Y.,
Bonder, A*., et al.* (2024). Silencing of aryl hydrocarbon receptor repressor restrains Th17 cell
immunity in autoimmune hepatitis. Journal of Autoimmunity *143*, 103162.

Greig, J.A., Nguyen, T.A., Lee, M., Holehouse, A.S., Posey, A.E., Pappu, R.V., and Jedd, G. (2020).
Arginine-Enriched Mixed-Charge Domains Provide Cohesion for Nuclear Speckle Condensation. Mol
Cell *77*, 1237-1250.e1234.
Holehouse, A.S., Das, R.K., Ahad, J.N., Richardson, M.O., and Pappu, R.V. (2017). CIDER: Resources to
Analyze Sequence-Ensemble Relationships of Intrinsically Disordered Proteins. Biophys J *112*, 16-21.
Hülsmann, B.B., Labokha, A.A., and Görlich, D. (2012). The permeability of reconstituted nuclear
pores provides direct evidence for the selective phase model. Cell *150*, 738-751.
Ilik, I.A., Malszycki, M., Lubke, A.K., Schade, C., Meierhofer, D., and Aktas, T. (2020). SON and SRRM2
are essential for nuclear speckle formation. eLife *9*.
Jang, S., Bedwell, Gregory J., Singh, Satya P., Yu, Hyun J., Arnarson, B., Singh, Parmit K.,
Radhakrishnan, R., Douglas, AidanDarian W., Ingram, Zachary M., Freniere, C.*, et al.* (2024). HIV-1
usurps mixed-charge domain-dependent CPSF6 phase separation for higher-order capsid binding,
nuclear entry and viral DNA integration. Nucleic Acids Research.
Karlsson, E., Schnatwinkel, J., Paissoni, C., Andersson, E., Herrmann, C., Camilloni, C., and Jemth, P.
(2022). Disordered Regions Flanking the Binding Interface Modulate Affinity between CBP and NCOA.
J Mol Biol *434*, 167643.
Lee, K., Ambrose, Z., Martin, T.D., Oztop, I., Mulky, A., Julias, J.G., Vandegraaff, N., Baumann, J.G.,
Wang, R., Yuen, W.*, et al.* (2010). Flexible use of nuclear import pathways by HIV-1. Cell Host Microbe
7, 221-233.
Lelek, M., Casartelli, N., Pellin, D., Rizzi, E., Souque, P., Severgnini, M., Di Serio, C., Fricke, T., Diaz-
Griffero, F., Zimmer, C.*, et al.* (2015). Chromatin organization at the nuclear pore favours HIV
replication. Nat Commun *6*, 6483.
Lester, E., Ooi, F.K., Bakkar, N., Ayers, J., Woerman, A.L., Wheeler, J., Bowser, R., Carlson, G.A.,
Prusiner, S.B., and Parker, R. (2021). Tau aggregates are RNA-protein assemblies that mislocalize
multiple nuclear speckle components. Neuron *109*, 1675-1691 e1679.
Li, C., Burdick, R.C., Nagashima, K., Hu, W.S., and Pathak, V.K. (2021). HIV-1 cores retain their integrity
until minutes before uncoating in the nucleus. Proc Natl Acad Sci U S A *118*.
Link, J.O., Rhee, M.S., Tse, W.C., Zheng, J., Somoza, J.R., Rowe, W., Begley, R., Chiu, A., Mulato, A.,
Hansen, D.*, et al.* (2020). Clinical targeting of HIV capsid protein with a long-acting small molecule.
Nature *584*, 614-618.
Luchsinger, C., Lee, K., Mardones, G.A., KewalRamani, V.N., and Diaz-Griffero, F. (2023). Formation of
nuclear CPSF6/CPSF5 biomolecular condensates upon HIV-1 entry into the nucleus is important for
productive infection. Sci Rep *13*, 10974.
Marasca, F., Sinha, S., Vadalà, R., Polimeni, B., Ranzani, V., Paraboschi, E.M., Burattin, F.V., Ghilotti, M.,
Crosti, M., Negri, M.L.*, et al.* (2022). LINE1 are spliced in non-canonical transcript variants to regulate
T cell quiescence and exhaustion. Nature Genetics *54*, 180-193.
Matreyek, K.A., Yucel, S.S., Li, X., and Engelman, A. (2013). Nucleoporin NUP153 phenylalanine-
glycine motifs engage a common binding pocket within the HIV-1 capsid protein to mediate lentiviral
infectivity. PLoS Pathog *9*, e1003693.
Mazzeo, L., Ghosh, S., Di Cicco, E., Isma, J., Tavernari, D., Samarkina, A., Ostano, P., Youssef, M.K.,
Simon, C., and Dotto, G.P. (2024). ANKRD1 is a mesenchymal-specific driver of cancer-associated
fibroblast activation bridging androgen receptor loss to AP-1 activation. Nature Communications *15*,
1038.
McGuffin, L.J., Bryson, K., and Jones, D.T. (2000). The PSIPRED protein structure prediction server.
Bioinformatics *16*, 404-405.
Muller, T.G., Zila, V., Peters, K., Schifferdecker, S., Stanic, M., Lucic, B., Laketa, V., Lusic, M., Muller, B.,
and Krausslich, H.G. (2021). HIV-1 uncoating by release of viral cDNA from capsid-like structures in the
nucleus of infected cells. Elife *10*.
Peng, K., Radivojac, P., Vucetic, S., Dunker, A.K., and Obradovic, Z. (2006). Length-dependent
prediction of protein intrinsic disorder. BMC Bioinformatics *7*, 208.
Peng, K., Vucetic, S., Radivojac, P., Brown, C.J., Dunker, A.K., and Obradovic, Z. (2005). Optimizing long
intrinsic disorder predictors with protein evolutionary information. J Bioinform Comput Biol *3*, 35-60.

Pornillos, O., Ganser-Pornillos, B.K., Kelly, B.N., Hua, Y., Whitby, F.G., Stout, C.D., Sundquist, W.I., Hill,
C.P., and Yeager, M. (2009). X-ray structures of the hexameric building block of the HIV capsid. Cell
137, 1282-1292.
Price, A.J., Jacques, D.A., McEwan, W.A., Fletcher, A.J., Essig, S., Chin, J.W., Halambage, U.D., Aiken, C.,
and James, L.C. (2014). Host cofactors and pharmacologic ligands share an essential interface in HIV-1
capsid that is lost upon disassembly. PLoS Pathog *10*, e1004459.
Rai, A.K., Chen, J.X., Selbach, M., and Pelkmans, L. (2018). Kinase-controlled phase transition of
membraneless organelles in mitosis. Nature *559*, 211-216.
Rasaiyaah, J., Tan, C.P., Fletcher, A.J., Price, A.J., Blondeau, C., Hilditch, L., Jacques, D.A., Selwood, D.L.,
James, L.C., Noursadeghi, M.*, et al.* (2013). HIV-1 evades innate immune recognition through specific
cofactor recruitment. Nature *503*, 402-405.
Rensen, E., Mueller, F., Scoca, V., Parmar, J.J., Souque, P., Zimmer, C., and Di Nunzio, F. (2021).
Clustering and reverse transcription of HIV-1 genomes in nuclear niches of macrophages. EMBO J *40*,
e105247.
Romero, P., Obradovic, Z., Li, X., Garner, E.C., Brown, C.J., and Dunker, A.K. (2001). Sequence
complexity of disordered protein. Proteins *42*, 38-48.
Scoca, V., Morin, R., Collard, M., Tinevez, J.Y., and Di Nunzio, F. (2023). HIV-induced membraneless
organelles orchestrate post-nuclear entry steps. J Mol Cell Biol *14*.
Segal-Maurer, S., DeJesus, E., Stellbrink, H.J., Castagna, A., Richmond, G.J., Sinclair, G.I., Siripassorn,
K., Ruane, P.J., Berhe, M., Wang, H.*, et al.* (2022). Capsid Inhibition with Lenacapavir in Multidrug-
Resistant HIV-1 Infection. N Engl J Med *386*, 1793-1803.
Selyutina, A., Bulnes-Ramos, A., and Diaz-Griffero, F. (2018). Binding of host factors to stabilized HIV-1
capsid tubes. Virology *523*, 1-5.
Selyutina, A., Persaud, M., Lee, K., KewalRamani, V., and Diaz-Griffero, F. (2020). Nuclear Import of the
HIV-1 Core Precedes Reverse Transcription and Uncoating. Cell Rep *32*, 108201.
Sickmeier, M., Hamilton, J.A., LeGall, T., Vacic, V., Cortese, M.S., Tantos, A., Szabo, B., Tompa, P., Chen,
J., Uversky, V.N.*, et al.* (2007). DisProt: the Database of Disordered Proteins. Nucleic Acids Res *35*,
D786-793.
Suzuki, Y., and Craigie, R. (2007). The road to chromatin — nuclear entry of retroviruses. Nature
Reviews Microbiology *5*, 187-196.
Taylor, I.A., and Fassati, A. (2024). The capsid revolution. J Mol Cell Biol *15*.
Tsanov, N., Samacoits, A., Chouaib, R., Traboulsi, A.M., Gostan, T., Weber, C., Zimmer, C., Zibara, K.,
Walter, T., Peter, M.*, et al.* (2016). smiFISH and FISH-quant - a flexible single RNA detection approach
with super-resolution capability. Nucleic Acids Res *44*, e165.
Wei, G., Iqbal, N., Courouble, V.V., Francis, A.C., Singh, P.K., Hudait, A., Annamalai, A.S., Bester, S.,
Huang, S.W., Shkriabai, N.*, et al.* (2022). Prion-like low complexity regions enable avid virus-host
interactions during HIV-1 infection. Nat Commun *13*, 5879.
Xue, B., Dunbrack, R.L., Williams, R.W., Dunker, A.K., and Uversky, V.N. (2010). PONDR-FIT: a meta-
predictor of intrinsically disordered amino acids. Biochim Biophys Acta *1804*, 996-1010.
Yamashita, M., and Emerman, M. (2004). Capsid is a dominant determinant of retrovirus infectivity in
nondividing cells. J Virol *78*, 5670-5678.
Zhang, L., Zhang, Y., Chen, Y., Gholamalamdari, O., Wang, Y., Ma, J., and Belmont, A.S. (2021). TSA-seq
reveals a largely conserved genome organization relative to nuclear speckles with small position
changes tightly correlated with gene expression changes. Genome Res *31*, 251-264.
Zhang, Z.-Y., Harischandra, D.S., Wang, R., Ghaisas, S., Zhao, J.Y., McMonagle, T.P., Zhu, G., Lacuarta,
K.D., Song, J., Trojanowski, J.Q.*, et al.* TRIM11 protects against tauopathies and is down-regulated in
Alzheimer's disease. Science *381*, eadd6696.
Zila, V., Margiotta, E., Turonova, B., Muller, T.G., Zimmerli, C.E., Mattei, S., Allegretti, M., Borner, K.,
Rada, J., Muller, B.*, et al.* (2021). Cone-shaped HIV-1 capsids are transported through intact nuclear
pores. Cell *184*, 1032-1046 e1018.

1120

Figure 1

Figure 2

A

B

Figure 3

Figure 4

A

B

C

D

Replication competent viral infection

Figure 5

A

B

Figure 6

Figure 7

Figure 8