1 **Primosomal protein PriC rescues replication initiation stress by bypassing the DnaA-DnaB**
2 **interaction step for DnaB helicase loading at *oriC***
3

4 Ryusei Yoshida, Kazuma Korogi, Qinfei Wu, Shogo Ozaki*, and Tsutomu Katayama*
5

6 Department of Molecular Biology, Kyushu University Graduate School of Pharmaceutical Sciences,
7 Fukuoka, Japan
8 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan
9

10 *Corresponding authors: Shogo Ozaki, shogo.ozaki@phar.kyushu-u.ac.jp; Tsutomu Katayama,
11 katayama@phar.kyushu-u.ac.jp

12
13

Abstract
In *Escherichia coli*, replisome and replication fork assembly are initiated by DnaB helicase
loading at the chromosomal origin *oriC* via its interactions with the DnaA initiator and the DnaC
helicase loader. Upon replication fork arrest, the replisome including DnaB dissociates from the stalled
fork. Replication fork progression is rescued by primosomal protein PriA- or PriC-dependent pathway
in which PriA and PriC promote reloading of DnaB in different mechanisms. However, the mechanism
responsible for rescue of blocked replication initiation at *oriC* remains unclear. Here, we found that
PriC rescued blocked replication initiation in cells expressing an initiation-specific DnaC mutant, in
mutant cells defective in DnaA-DnaB interactions, and in cells containing truncated *oriC* sequence
variants. PriC rescued DnaB loading at *oriC* even in the absence of Rep helicase, a stimulator of the
PriC-dependent replication fork restart pathway. These and results of *in vitro* reconstituted assays
concordantly suggest that this initiation-specific rescue mechanism provides a bypass of the DnaA-
DnaB interaction for DnaB loading by PriC-promoted loading of DnaB to the unwound *oriC* region.
These findings expand understanding of mechanisms sustaining the robustness of replication initiation
and specific roles for PriC in the genome maintenance.

29
30

Introduction
Chromosome replication is strictly regulated to ensure successful duplication of the genetic
material (Costa & Diffley, 2022; Grimwade & Leonard, 2021; Kasho et al., 2023). In *Escherichia coli*,
the ATP-bound form of the replication initiator protein DnaA (ATP-DnaA) forms a higher-order
complex with DNA-bending protein IHF at the unique chromosomal origin *oriC*. This complex
promotes local unwinding of *oriC* and recruits a pair of DnaB helicases, which are successively loaded
to the single-stranded (ss) DNA regions in a bidirectional manner with the aid of DnaC helicase loader.

ssDNA-loaded DnaB interacts with DnaG primase and DNA polymerase III holoenzyme, resulting in replisome (Arias-Palomo et al., 2019; Chodavarapu & Kaguni, 2016; Hayashi et al., 2020; O'Donnell et al., 2013; Sakiyama et al., 2017, 2018, 2022; Wegrzyn & Konieczny, 2023). Replisomes disassemble when the replication fork stalls in front of an obstacle, such as protein-DNA roadblock or single strand breaks. PriA recognizes the abandoned fork structure, which triggers reloading of DnaB with its partner proteins, such as PriB, PriC, DnaT, and DnaC (Heller & Marians, 2005a; Masai et al., 1994; Michel & Sandler, 2017; Windgassen et al., 2018). In addition, independently of PriA, PriC binds to the abandoned fork structure, triggering reloading of DnaB with the aid of DnaC, which is supported by Rep helicase depending on the fork structure (Heller & Marians, 2005b; Sandler et al., 1999). Also, homologous recombination mechanisms can rescue the abandoned replication fork during SOS responses (Asai & Kogomal, 1994; Kogoma, 1997; Né Dicte Michel et al., 2001). In contrast to these well characterized mechanisms at abandoned forks, little is known about what happens when replication initiation is impeded at *oriC*.

E. coli oriC comprises a Duplex-Unwinding Element (DUE) and a DnaA-Oligomerization Region (DOR) (Figure 1A) (Grimwade & Leonard, 2021; Kasho et al., 2023; Wegrzyn & Konieczny, 2023). The DUE contains 13-mer AT-rich sequence repeats known as L-, M-, and R-DUEs. The M- and R-DUEs are essential for stable DUE unwinding with the specific sequences (TT[A/G]T(T)) used for DnaA-ssDNA binding (Ozaki & Katayama, 2012; Sakiyama et al., 2017). The L-DUE containing the TTATT sequence promotes efficient DnaB loading by expanding the unwound *oriC* region (Sakiyama et al., 2022). The DOR is divided into three subregions: the Left-, Middle-, and Right-DORs, which contain asymmetric 9-mer DnaA binding sequences (DnaA box) with the consensus sequence TTA[T/A]NCACA (Figure 1A) (Noguchi et al., 2015; Ozaki & Katayama, 2012; Rozgaja et al., 2011; Sakiyama et al., 2017; Shimizu et al., 2016). The Left-DOR contains a cluster of unidirectionally aligned DnaA boxes including high-affinity DnaA box R1, and low-affinity boxes R5M, τ1−2, and I1-2, with an IHF-binding region between R1 and R5M boxes. The Right-DOR contains an oppositely oriented DnaA box cluster including high-affinity DnaA box R4 and low-affinity boxes I3 and C1-3. These DnaA box clusters form frameworks for the Left- and Right-DnaA subcomplexes, respectively. DnaA bound to the R2 box, which solely resides in the Middle-DOR, stabilizes these DnaA subcomplexes (Miller et al., 2009; Rozgaja et al., 2011; Shimizu et al., 2016). In addition, the AT-cluster (TATTAAAAGAA) region, which connects to the L-DUE, stimulates DnaB loading in the absence of the Right-DnaA subcomplex (Sakiyama et al., 2022).

DnaA comprises four functional domains (Figure 1B). Domain I binds to multiple proteins such as DnaB helicases and the DnaA-assembly stimulator DiaA (Abe et al., 2007; Hayashi et al., 2020; Keyamura et al., 2007, 2009). Domain II is a flexible linker (Abe et al., 2007; Nozaki & Ogawa, 2008).

Domain III contains AAA+ (ATPases Associated with various cellular Activities) motifs, which are involved in tight ATP/ADP binding, ATP hydrolysis, and domain III-III interactions (Duderstadt et al., 2011; Erzberger et al., 2006; Felczak & Kaguni, 2004; Kawakami et al., 2005; Ozaki et al., 2012). The arginine finger motif (Arg285) in domain III interacts with ATP bound to domain III of adjacent DnaA protomers, stimulating DnaA complex formation in a cooperative manner (Kawakami et al., 2005; Noguchi et al., 2015). In addition, H/B motifs (Val 211 and Arg245) in this domain bind to ssDNA in a sequence-specific manner (Ozaki et al., 2008; Sakiyama et al., 2017). Domain IV, which comprises the helix-turn-helix motif, recognizes the DnaA box (Fujikawa et al., 2003).

For replication initiation, ATP-DnaA molecules cooperatively oligomerize at *oriC* to form the Left- and Right-DnaA subcomplexes (Figure 1C). Formation of these complexes is stimulated by homotetrameric DiaA protein, which binds up to four DnaA molecules via domain I, including Phe46 (Keyamura et al., 2007, 2009). In concert with DNA superhelicity and thermal energy, the DUE undergoes initial unstable unwinding, which is stabilized through specific interaction between H/B motifs of domain III in the Left-DnaA subcomplex and the upper strand of M- and R-DUEs. This interaction is facilitated by sharp DNA bending by IHF, resulting in stable DUE unwinding (Figure 1C) (Ozaki et al., 2008; Sakiyama et al., 2017). When the single-stranded region is expanded to the L-DUE, the resultant single-stranded L-DUE binds the H/B motifs within the Right-DnaA subcomplex, maximizing the efficiency of the DnaB loading process (Sakiyama et al., 2022).

Subsequently, two DnaB helicases are loaded onto each ssDUE strand through interactions with DnaC and DnaA (Figure 1D). DnaC binding to DnaB changes the closed ring structure of the DnaB hexamer to an open spiral form to allow it to encircle ssDNA (Arias-Palomo et al., 2019; Nagata et al., 2020). Each Left- and Right-DnaA subcomplex binds a DnaB-DnaC complex via high-affinity interaction between DnaA Phe46 and DnaB Leu160 and guides the DnaB-DnaC complexes to unwound ssDUE strands via a low-affinity interaction between DnaA domain III His136 and DnaB (Figure 1B and 1D). Upon interaction with ssDNA, DnaC dissociates from DnaB, which enables migration of DnaB hexamer with helicase action. The pair of DnaB helicases loaded onto the ssDNA strands progress in opposite directions, forming replisomes with DnaG and Pol III holoenzyme (Chodavarapu & Kaguni, 2016; O'Donnell et al., 2013).

Replication restart pathways ensure fork progression of the entire chromosome under conditions that trigger replisome disassembly (Heller & Marians, 2005a; Lopper et al., 2007; Michel & Sandler, 2017). PriA helicase-dependent pathways predominantly facilitate DnaB reloading onto abandoned forks *in vivo* (Flores et al., 2002). PriA is a 3'-5' DNA helicase and has specific affinity for forked DNA structures comprising one parental dsDNA and two newly synthesized sister dsDNA strands. This permits PriA recognition of the abandoned replication fork and the subsequent unwinding of the nascent lagging

strand (Duckworth et al., 2023; Windgassen et al., 2018; Windgassen & Keck, 2016). The resultant unwound ssDNA associates with PriB and DnaT to promote reloading of DnaB helicase with the aid of DnaC (Duckworth et al., 2023; Heller & Marians, 2005a; Lopper et al., 2007). In the absence of PriB, PriC participates in a PriA-dependent pathway (Sandler et al., 1999). In addition, PriA is required for *oriC*/DnaA-independent chromosomal replication called stable DNA replication (SDR) (Masai et al., 1994), which is promoted by UV irradiation (inducible iSDR) and loss of *rnhA* or *recG* (constitutive cSDR) (Kogoma, 1997). In both mechanisms, replication is initiated at triple-stranded structures comprising a dsDNA-ssDNA hybrid (D-loop) or a dsDNA-ssRNA hybrid (R-loop) generated at specific chromosomal loci. These structures mimic the abandoned fork structure, thereby allowing helicase loading to occur similarly to that during PriA-dependent fork restart (Masai et al., 1994).

Independently of PriA, PriC can restart replication from abandoned forks through its interaction with the ssDNA region and SSB (Single-Stranded DNA Binding protein) (Figure 1E) (Heller & Marians, 2005a; Wessel et al., 2013, 2016). PriC consists of N-terminal and C-terminal domains, which are composed of α helices and are connected by a short linker. The PriC C-terminal domain remodels the SSB-ssDNA complex, which is a prerequisite for the recruitment of DnaB helicase (Heller & Marians, 2005a; Wessel et al., 2013, 2016). PriC also interacts with Rep helicase. When the length of the ssDNA gap within the abandoned fork is short, Rep helps expand the ssDNA gap, promoting PriC-mediated remodeling of the SSB-ssDNA complex (Heller & Marians, 2005a, 2005b; Nguyen et al., 2021). Notably, in addition to these functions at the abandoned forks, PriC is suggested to play a role in DnaA-*oriC*-dependent replication initiation under challenging conditions, as evidenced by the synthetic lethality between *priC303*::*kan* and a subgroup of temperature-sensitive *dnaA* mutant alleles (*dnaA46* and *dnaA508*): i.e. *priC303*::*kan* mutants bearing *dnaA46* or *dnaA508*, but not wild-type (WT) *dnaA*, can not grow even at 30°C (Hinds & Sandler, 2004). However, the mechanisms via which PriC rescues blocked replication initiation remain elusive.

In this study, we analyzed the PriC-dependent promotion of replication initiation from *oriC* in the context of various replication initiation stresses. We found that PriC stimulated replication initiation in *dnaC2* cells, in which DnaB helicase loading at *oriC* is defective, indicating that it has a role in the rescue of blocked replication initiation. Moreover, PriC stimulated the growth of cells defective in DnaA-DnaB interactions in addition to the growth of cells with *oriC* sequence deletions that inhibit DnaB loading. Consistent with these results, in an *in vitro* reconstituted system, PriC stimulated DnaB loading when the DnaA-DnaB interaction was inhibited at *oriC*. Furthermore, we found that PriC did not stimulate initiation of cSDR, demonstrating that PriC functions specificity in the DnaA-*oriC* system. Taken together, we suggest that PriC rescues blocks in replication initiation by bypassing DnaB loading by *oriC*-DnaA complexes.

<div align="center">**Results**</div>

PriC stimulates DNA replication initiation in *dnaA46* and *dnaC2* cells.

The temperature-sensitive *dnaA46* and *dnaA508* mutants exhibit synthetic lethality with *priC303*::*kan* even at 30°C (Hinds & Sandler, 2004), suggesting that PriC can rescue blocks in DnaA-dependent replication initiation. However, the mechanism underlying PriC-promoted rescue of blocks in replication initiation remains a mystery. Based on previous reports demonstrating PriC binding to short oligo-ssDNA, SSB, and DnaB and its role in DnaB loading at stalled replication forks (Wessel et al., 2013, 2016), we considered the possibility that PriC permits bypass of DnaA-dependent stable DUE unwinding and/or DnaB loading at the *oriC* ssDUE. To genetically test these ideas, we analyzed the impact of PriC on the growth of a temperature-sensitive *dnaC2* mutant, which is competent for ongoing replisome progression at 37–42°C, but defective in DnaB loading specifically at the *oriC* ssDUE (Withers & Bernander, 1998).

Colony formation of *dnaA46* cells was moderately inhibited at 37°C (Figure 2A). Consistent with a previous report, when crossed with the Δ*priC* allele, colony formation of *dnaA46* mutant cells but not WT *dnaA* cells was severely inhibited at 37°C (Figure 2A). Unlike the *dnaA46 priC303*::*kan* mutant, *dnaA46* Δ*priC* cells formed colonies at 30°C, although colony growth was slightly retarded. This difference may stem from the different strain backgrounds. Notably, colony formation of *dnaC2* cells at 35°C was moderately inhibited, whereas it was severely inhibited in *dnaC2* Δ*priC* cells (Figure 2A). At 30°C, colony formation by *dnaC2* cells and *dnaC2* Δ*priC* cells was similar. The observed requirement for PriC for colony formation by *dnaC2* cells at 35°C supports the idea that PriC assists in the DnaC-dependent DnaB loading step at *oriC*.

To investigate replication initiation from *oriC*, we used flow cytometry of synchronized cells bearing *dnaA46* or *dnaC2* with or without Δ*priC*. In these experiments, cells were incubated at 42°C or 37°C for 80–90 min to synchronize the replication cycle at the pre-initiation step, after which incubation was continued at 30°C for 5 min to allow initiation and then 4 h in the presence of rifampicin and cephalexin to inhibit both replication initiation and cell division and allow replication run-out of the whole chromosome. The resultant number of chromosomes per cell, measured by flow cytometry, is known to correspond to the number of *oriC* copies per cell at the time of drug addition, an indicator of initiation activity.

A majority of the synchronized *dnaA46* and *dnaC2* cells had only a single *oriC* copy per cell after incubation at 42°C or 37°C regardless of the presence of Δ*priC* (Figure 2B and 2C). In WT *priC* cells, after release of initiation at 30°C for 5 min, most cells had two *oriC* copies. However, in *dnaA46* Δ*priC* cells, only about half of the cells had two *oriC* copies after the release (Figure 2B), indicating

<div align="center">5</div>

moderate inhibition of initiation. In *dnaC2 ΔpriC* cells, the number of two-*oriC* cells was slightly lower

than that of *dnaC2* cells (Figure 2C). These observations are consistent with the colony formation data

and with our hypothesis that PriC contributes to the rescue of blocked replication initiation by assisting

DnaB loading onto the *oriC* ssDNA.

PriC rescues the cell growth of DnaA mutants defective in DnaB interactions.

DnaA contains two regions for interactions with DnaB during DnaB loading at *oriC* ssDNA:

one is a high-affinity region in domain I containing Phe46, and the other is a low-affinity region in

domain III containing His136. To analyze the mechanism of PriC-promoted DnaB loading at *oriC*, we

introduced a series of pING1 vector-based plasmids encoding WT DnaA (pKA234) or the DnaA F46A

H136A double mutant (pFH) into *dnaA46* cells with or without WT *priC*. Growth of *dnaA46* cells

containing pING1 was inhibited at 42°C, but the leaky expression of *dnaA* from pKA234 enabled the

growth of these cells, as previously reported (Sakiyama et al., 2018) (Figure 3). Introduction of pFH

moderately supported colony formation by *dnaA46* cells at 42°C. By contrast, *dnaA46 ΔpriC* cells

carrying pFH exhibited little growth at 42°C (Figure 3). These results suggest that PriC is crucial for the

growth of cells defective in specific DnaA-DnaB interactions, possibly because it can bypass the DnaA-

DnaB interaction step required for DnaB loading onto *oriC* ssDNA.

PriC rescues DNA replication inhibition by DiaA overexpression *in vivo*.

DiaA, a homotetrameric protein, binds to multiple DnaA molecules and stimulates DnaA

multimer assembly at *oriC* (Keyamura et al., 2007). The Phe46 residue in DnaA domain I is part of the

DiaA-binding site and DnaB binding site. Oversupply of DiaA inhibits timely initiation of replication in

cells growing in tryptone medium at 30°C, probably because of binding competition between DiaA and

DnaB for DnaA domain I (Ishida et al., 2004; Keyamura et al., 2009). If PriC can bypass the DnaA-

DnaB interaction requirement for DnaB loading onto *oriC*, PriC should suppress inhibition of initiation

by DiaA oversupply. To test this possibility, we introduced pBR322 or its derivative encoding *diaA*

(pNA135) into cells with or without WT *priC* (Figure 4). WT *priC* cells carrying pBR322 or pNA135

similarly formed colonies on LB agar plates containing ampicillin between 25°C and 42°C (Figure 4A

and Figure 4-figure supplement 1). However, in *ΔpriC* cells carrying pNA135, but not pBR322, colony

formation was moderately inhibited at 25°C (Figure 4A), supporting our conjecture that PriC can

bypass the requirement for the DnaA-DnaB interaction.

Rep helicase stimulates PriC-mediated rescue of arrested replication forks containing nascent

strands (Heller & Marians, 2005b, 2005a; Sandler, 2000). To test the contribution of Rep helicase to

PriC-promoted rescue of blocked replication initiation, we introduced pBR322 or pNA135 into Δ*rep*

cells. Unlike Δ*priC* cells, Δ*rep* cells carrying pNA135 grew similarly to Δ*rep* cells carrying pBR322

(Figure 4A), suggesting no involvement of Rep in the PriC-promoted rescue of blocked replication

initiation. This is consistent with the features of unwound *oriC*, which has a fork structure similar to

that of the replication fork but without nascent strands, and the specific role of Rep in rescuing

abandoned replication forks with nascent strands in the PriC-dependent pathway. Rep is specifically

required for the unwinding of the nascent leading strand of abandoned replication forks, which is

needed to expand the ssDNA region required for entry of PriC (Heller & Marians, 2005b, 2005a).

We further analyzed replication initiation in Δ*priC* cells carrying pNA135 at 30°C by flow

cytometry (Figure 4B). Exponentially growing cells were further incubated for run-out replication after

the addition of rifampicin and cefalexin. The volume (mass) of cells at the time of drug addition was

also quantified. Under these conditions, WT *priC* cells bearing pBR322 mainly had eight or four *oriC*

copies per cell with eight-*oriC* cells predominating (Figure 4B). The number of *oriC* copies per cell was

similar to that in WT *priC* cells bearing pNA135, with only slight increase of asynchronous initiation,

as shown by the slight increase in the number of cells with non-2^n *oriC* copies, which is basically

consistent with our previous data (Ishida et al., 2004). However, in Δ*priC* cells bearing pNA135, severe

inhibition of initiation occurred, as shown by the substantial decrease in the number of eight-*oriC* cells

and the large increase of the number of cells with only one to three *oriC* copies (Figure 4B). This result

supports the proposed function of PriC at *oriC* mentioned above. In addition, the observed

asynchronous initiations in Δ*priC* cells bearing pBR322 is worthy of attention. Indeed, the fraction of

cells with non-2^n *oriC* copies was higher among Δ*priC* cells than among WT *priC* cells (Figure 4B),

suggesting that even in normal cells, inhibition of replication initiation occurs at low frequency and

requires PriC function to rescue replication initiation.

To test whether PriC rescues replication initiation inhibited by deletion of the *diaA* gene, we

conducted similar flow cytometry analysis of Δ*priC*, Δ*diaA*, and Δ*priC* Δ*diaA* cells (Figure 4C). Δ*diaA*

cells showed severe asynchrony of replication initiation, as shown by the increase in the number of five-

to-seven-*oriC* cells, which supports the stimulatory role of DiaA in DnaA-*oriC* complex formation

(Figure 4C). However, Δ*priC* Δ*diaA* cells did not show more asynchronization of replication initiation

than Δ*diaA* cells, as shown by the similar asynchrony index (A.I.). These results suggest that PriC does

not support or bypass DnaA-*oriC* complex formation.

PriC stimulates replication initiation in cells with *oriC* sequence truncations

Right DnaA-subcomplex plays a stimulatory role in DnaB helicase loading (Sakiyama et al.,

2022). The loading of two DnaB hexamers at *oriC* ssDNA regions requires multiple steps that depend

on the distinct functions of two DnaA subcomplexes (Figure 1): the Left-DnaA subcomplex stably

unwinds M-R regions of the DUE and the Right-DnaA subcomplex expands the unwound region to the

AT-L region, supporting efficient DnaB helicase loading (Sakiyama et al., 2017, 2022; Yoshida et al.,

2023). The expanded ssDNA regions in this open *oriC* complex efficiently promote the loading of one

DnaB hexamer onto the lower (A-rich) strand M-R region and a second DnaB hexamer onto the upper

(T-rich) strand of the DUE (Fang et al., 1999; Sakiyama et al., 2022). In the absence of the Right-DnaA

subcomplex, the AT region of the AT-L region assists in DnaB loading (Sakiyama et al., 2022;

Stepankiw et al., 2009)

To determine whether PriC rescues defects in DnaB loading when the *oriC* sequence is altered

to prevent Right-DnaA subcomplex formation, we first assessed the colony formation of cells lacking

the Middle- and Right-DOR of *oriC* (Left-*oriC*) (Figure 5A). The Left-*oriC* mutant formed colonies at

37°C but showed severe growth defects at 25°C and 30°C (Figure 5B). At 37°C, thermal energy might

be sufficient to expand DUE unwinding into the AT region and allow DnaB loading in the absence of

the Right-DnaA subcomplex (Sakiyama et al., 2022), but not at 25°C or 30°C. By contrast, colony

formation of the Left-*oriC* Δ*priC* double mutant was markedly compromised at 37˚C (Figure 5B) and

the growth defects of the Left-*oriC* mutant 25°C and 30°C were aggravated. These findings support the

idea that PriC functions to rescue defective processes in DnaB loading caused by deletions of the *oriC*

sequence that prevent DnaA subcomplex formation.

Next, similar experiments were performed using a mutant in which the R4 DnaA box within

Right-DOR was substituted with a sequence (R4*Tma*) defective in binding of *E. coli* DnaA (Figure 5A

and 5C) (Noguchi et al., 2015; Sakiyama et al., 2022). We reasoned that formation of the Right-DnaA

subcomplex would be inhibited by the introduction of R4*Tma* and PriC would be required for robust

initiation in this context. As in the Left-*oriC* strain, the R4*Tma* strain showed normal colony formation

in the presence of WT *priC*. By contrast, in the R4*Tma* Δ*priC* strain, colony formation was impaired

moderately at 25°C and slightly at 30°C (Figure 5C), indicating that PriC can rescue the blocked

replication initiation caused by the absence of the R4 box. The reduced colony formation of R4*Tma*

Δ*priC* double mutant cells was alleviated at higher temperatures, similar to the phenotype of the Left-

oriC mutant.

Furthermore, we assessed replication initiation in R4*Tma* cells at 30°C using flow cytometry

(Figure 5D). R4*Tma* cells grew slightly slower than WT *oriC* cells and showed clear inhibition of

initiation; in contrast to the WT *oriC* strain, the R4*Tma* mutant cell population contained more four-

oriC cells and fewer eight-*oriC* cells, and showed severe asynchronous initiation. Notably, these

negative effects of the R4*Tma* mutation were amplified by deletion of *priC*, i.e., the number of one to

three-*oriC* cells increased and the ori/mass ratio was further reduced.

We next analyzed the role of PriC in cells lacking the AT-L region of *oriC* (subATL *oriC*)

(Figure 6A). L-DUE and the flanking AT region assist in DnaB helicase loading by stimulating *oriC*

unwinding, which is essential in a strain defective in Right-DnaA subcomplex formation (Sakiyama et

al., 2022). SubATL *oriC* cells and WT *oriC* cells formed colonies with similar efficiency on LB agar

plates between 25°C and 37°C, regardless of the presence of Δ*priC* (Figure 6B). However, flow

cytometry analysis revealed that deletion of *priC* specifically inhibited initiation in subATL *oriC* cells

(Figure 6C); the number of eight-*oriC* cells was lower while the number of four-to-seven *oriC* cells was

higher in subATL *oriC* Δ*priC* cells than in subATL *oriC* cells with WT *priC*. Taken together, these

results are consistent with the idea that PriC restores initiation and DnaB loading at such truncated *oriC*

sequences.

PriC loads DnaB at *oriC* unwound by the DnaA complex in an *in vitro* reconstituted system.

To analyze the mechanism of PriC rescue of blocked initiation at *oriC*, we assessed PriC activity

in an *in vitro* reconstituted system for DnaB loading and DnaB-promoted DNA unwinding using the

supercoiled circular form (form I) of the *oriC* plasmid pBS*oriC* and purified proteins, namely DnaA, N-

terminally histidine-tagged DnaB (His-DnaB), DnaC, IHF, SSB, and gyrase. In this system, DnaA and

IHF unwind the DUE and stimulate His-DnaB loading to the ssDUE region with the aid of DnaC. The

loaded His-DnaB expands the ssDNA regions in concert with the ssDNA-binding activities of SSB.

Concomitantly, the positive supercoiling generated through DNA unwinding by His-DnaB is resolved by

DNA gyrase, resulting in the formation of plasmid DNA topoisomers (form I*). Form I* can be separated

from form I by agarose gel electrophoresis (Sakiyama et al., 2018, 2022).

First, we assessed the effect of PriC on WT DnaA. WT DnaA stimulated form I* formation in

the absence of PriC (Figure 7A, lanes 1 and 5) and increasing concentrations of PriC moderately

inhibited WT DnaA-mediated form I* formation (Figure 7A, lanes 5 to 8), probably because of

competition between DnaA and PriC for binding to DnaB. Next, we focused on the DnaA F46A H136A

double mutant. The DnaA F46A and DnaA H136A mutant proteins alone are reported not to support

form I* formation (Keyamura et al., 2009; Sakiyama et al., 2018). Consistently, the DnaA F46A H136A

double mutant protein was inactive in form I* formation (Figure 7A, lane 9). However, when PriC was

added to the assay, DnaA F46A H136A promoted form I* formation to a level comparable to that of WT

DnaA in the presence of PriC (Figure 7A, lanes 5-12 and 7B), indicating that PriC rescues defective

DnaB loading by DnaA F46A H136A-*oriC* complexes. These findings are consistent with the idea that

PriC can bypass the strict reliance on DnaA-DnaB interactions for DnaB loading at *oriC*.

To corroborate this idea, we also determined whether PriC rescues blocked initiation resulting

from unstable DUE unwinding. *oriC* complexes containing DnaA V211A cannot bind to the ssDUE and

consequently fails in stable DUE unwinding (Ozaki et al., 2008). The DnaA V211A mutant was

virtually inactive for form I * formation irrespective of the presence or absence of PriC (Figure 7A,

lanes 13-16, and 7B), indicating that PriC cannot efficiently rescue a defect of DUE unwinding.

We further evaluated the DnaB loading activity of PriC in the presence of DiaA using a similar

in vitro reconstituted system. Previously, DiaA was shown to stimulate *oriC* DUE unwinding, but to

inhibit form I* formation, because DiaA-DnaA binding competitively inhibits DnaB-DnaA binding; DiaA

and DnaB share the same binding site in DnaA domain I (Figures 7C lanes 1, 3, 5, and 7 and 7D)

(Keyamura et al., 2009). PriC moderately stimulated form I* formation specifically in the presence of

both DnaA and DiaA (Figures 7C lanes 7 and 8, and 7D). These results further support the idea that PriC

can bypass the specific requirement for the DnaA-DnaB interaction in DnaB loading at *oriC*. DiaA bound

to *oriC*-DnaA complexes might be a physical obstacle reducing efficiency of DnaB loading by PriC.

In addition, we analyzed the DUE unwinding activities of DnaA mutants using the *in vitro*

reconstituted system. As previously reported, DnaA V211A is largely inactive in DUE unwinding, but

DnaA F46A H136A unwinds the DUE at a level comparable to that of WT DnaA (Figure 7-figure

supplement 2A and 2B). PriC did not stimulate DUE unwinding irrespective of the presence of WT

DnaA (Figure 7-figure supplement 2C and 2D). These results demonstrate the functional specificity of

PriC in rescuing failed DnaB loading at *oriC* ssDNA and are consistent with the PriC mechanism

mentioned above.

Role for PriC in cSDR

Finally, to analyze the role of PriC in other types of replication initiation, we examine whether

PriC contributes to the initiation of cSDR, which does not require *dnaA* or *oriC* and is activated in cells

lacking *rnhA* or *recG*. Although the PriA-PriB-DnaT primosome complex has previously been reported

to be involved in the initiation of cSDR (Heller & Marians, 2006), it is unclear whether PriC has any

role in this type of initiation.

Unlike *dnaA46* cells, *dnaA46 rnhA*::*cat* double mutant cells formed colonies even at 40°C

(Figure 8A), indicating that these cells were engaged in cSDR, as previously reported (Hinds & Sandler,

2004). The formation of colonies by Δ*priC dnaA46 rnhA*::*cat* triple mutant cells was severely inhibited

at 40°C (Figure 8A), suggesting that PriC contributes to the growth of *dnaA46 rnhA*::*cat* mutant cells.

To determine whether PriC contributes to the initiation of *dnaA46-oriC* replication or cSDR,

we calculated the relative ratios of *oriC* (84.6 min) and *ter* (32.4 min) copy numbers to the refence

locus *yapB* (50.5 min) copy number using real-time quantitative PCR (qPCR). To calculate the copy

number of the chromosomal *ter* region, the preferential initiation site of cSDR (Maduike et al., 2014),

dnaA46 rnhA::*cat* cells growing exponentially at 30°C were shifted to 40°C for 90 min and the copy

number before and after incubation at 40°C was calculated. Consistent with the colony formation data (Figure 8A), the *ter* copy number ratio of the *dnaA46 rnhA*::*cat* double mutant increased after 40°C incubation, whereas the copy number ratio of the *dnaA46* mutant expressing WT *rnhA* remained the same. The *ter* copy number ratio of the *dnaA46 rnhA*::*cat* Δ*priC* triple mutant strain was also increased, indicating that cSDR could occur even in the absence of PriC.

To monitor initiation of DnaA-*oriC* replication, we determined the *oriC* copy number ratio (Figure 8B). At the permissive temperature, the *oriC* copy number ratio of the *dnaA46* Δ*priC* double mutant strain was lower than that of the *dnaA46* single mutant, confirming the importance of PriC for replication initiation in *dnaA46* cells, as shown in Figure 2. In the *dnaA46 rnhA*::*cat* double mutant expressing WT *priC*, the *oriC* copy number ratio was reduced, which is explained by activation of cSDR and increase of the relative copy number of the reference *yapB* locus (50.5 min chromosomal map position). Together, these results suggest the idea that PriC stimulates replication fork progression during cSDR.

Discussion

In growing cells, replication initiation can be blocked under challenging conditions. However, the mechanisms that rescue blocked initiations are largely unknown. To elucidate such mechanisms, we investigated PriC-promoted replication initiation under various challenging conditions. We showed that PriC was necessary for the optimal growth of *dnaC2* cells at the semi-restrictive temperature and for replication initiation even at the permissive temperature (Figure 2), suggesting that PriC contributes to replication initiation rescue in *dnaC2* cells by facilitating DnaB helicase loading at *oriC*. Furthermore, we observed that PriC was necessary for the optimal growth of cells in which the DnaA-DnaB interaction was inhibited by DiaA overexpression or DnaA F46A H136A double mutations (Figures 3 and 4). In addition, PriC was stimulatory for the colony formation and replication initiation bearing *oriC* mutations that impair efficient DnaB loading (Figures 5 and 6). These results, together with results of form I* formation assays (Figures 7), suggest that PriC bypasses the need for the DnaA-DnaB interaction by binding to unwound *oriC* DNA and recruiting DnaB to facilitate its loading onto ssDNA (Figure 9). This mechanism is similar to that proposed for the rescue of abandoned replication forks by PriC (Figure 1E).

The finding that PriC failed to stimulate form I* formation in the presence of DnaA V211A, which lacks DUE unwinding activity (Figures 7), provides mechanistic insight into PriC-dependent replication initiation at *oriC*. We propose that PriC-dependent replication initiation possibly begins with the recognition of a stably unwound ssDNA region by PriC (Figure 9). Consistent with this perspective, PriC ssDNA-binding activity and ssDNA regions have been reported to be important for PriC-promoted

replication fork rescue (Heller & Marians, 2006; McMillan & Keck, 2024). The failure of PriC to

stimulate replication initiation in Δ*diaA* cells (Figure 4C) is also potentially explained by the reduced

DUE unwinding caused by destabilization of the DnaA-*oriC* complex in Δ*diaA* cells. Based on this, we

suggest that the stably unwound ssDNA region in *oriC* serves as a distinctive feature for PriC-

promoting DnaB loading at *oriC* for replication initiation rescue, and distinguishes it from PriA-

dependent replication restart, which requires the structure of the replication fork DNA consisting of

sister dsDNA strands (Heller & Marians, 2005a; Windgassen et al., 2018).

In this study, we found that cells with DnaA F46A H136A double mutations grew well, but

only when the cells expressed WT PriC, whereas the growth of cells expressing the DnaA H136A single

mutant was severely inhibited even when the cells expressed WT PriC (Sakiyama et al., 2018). The

His136 residue is located within the weak, secondary DnaB interaction region in DnaA, and is crucial

for DnaB loading onto *oriC* ssDNA. Although domain I in DnaA H136A can stably tether DnaB-DnaC

complexes to DnaA complexes on *oriC* (Sakiyama et al., 2018), the complexes fail to load DnaB onto

oriC ssDNA even in the presence of PriC. It is possible that the interaction between PriC and DnaB is

inhibited by DnaB binding to DnaA domain I. Conversely, the PriC-DnaB interaction may inhibit the

interaction between DnaA domain I and DnaB, as suggested by the inhibitory effect of PriC on DnaB

loading *in vitro* in the presence of WT DnaA (Figure 7). However, this inhibitory effect of PriC was not

observed *in vivo*, suggesting that PriC-DnaB binding could involve still unknown factors and regulatory

mechanisms.

PriC supported cSDR-dependent growth in *dnaA46 rnhA*::*cat* double mutant cells without

stimulating the initiation of cSDR (Figure 8). In this mutant, cSDR predominantly initiates from the *ter*

region, resulting in head-on collisions between the replisome and transcription complexes in *rrn*

operons (Maduike et al., 2014). This suggests that in cSDR cells PriC specifically rescues abandoned

forks including those generated during head-on conflicts between the replisome and transcription

complexes.

Bacillus subtilis (*B. subtilis*), which expresses a PriA homolog but not a PriC homolog,

employs a PriA-independent fork rescue pathway (Bruand et al., 2001). In this pathway, *B. subtilis*

DnaC (*Bsu*DnaC) helicase is reloaded using the helicase loader DnaI together with *B. subtilis* DnaB

(*Bsu*DnaB) and DnaD coloaders. The ssDNA-binding activity of *Bsu*DnaB has a crucial role in this

PriA-independent pathway (Bruand et al., 2005). SsDNA-bound *Bsu*DnaB remodels the SSB-ssDNA

complex with DnaD and recruits *Bsu*DnaC. These functions of *Bsu*DnaB are similar to the proposed

functions of PriC in replication fork rescue in *E. coli*. In addition, *Bsu*DnaC helicase is recruited to *oriC*

by concerted actions of *B. subtilis* DnaA and the DnaD-*Bsu*DnaB complex, which has an essential role

in chromosome replication initiation (Jameson & Wilkinson, 2017), suggesting a similar role of

*Bsu*DnaB to PriC in *E. coli* cells expressing DnaA F46A H136A. These similarities between PriC and

*Bsu*DnaB in these two evolutionarily distant bacterial species suggest that the mechanism of PriC-

promoted helicase loading is conserved among bacterial species despite the absence of sequence

conservation of PriC and *Bsu*DnaB homologs.

Based on our results, we propose the abnormal competition between DnaB and DiaA for

DnaA domain I could represent a form of intrinsic replication initiation stress in bacteria with conserved

DiaA homologs. This type of stress could also occur in ε-proteobacterial species, such as *Helicobacter*

pylori, because they express HobA, a DiaA-functional homolog (Natrajan et al., 2007; Zawilak-Pawlik

et al., 2011). Similarly, YfdR, encoded by *E. coli* prophages, inhibits replication initiation by competing

with DnaB for binding to DnaA domain I (Noguchi & Katayama, 2016), suggesting that such

extraneously introduced inhibitors could trigger replication initiation stress.

Even in human, ORC1, ORC2, and ORC5, the essential components of the eukaryotic

replication initiation complex, are not essential in some cancer cell-lines (Shibata & Dutta, 2020; Struhl

et al., 2016), suggesting that mechanisms of replication initiation rescue may also operate beyond the

bacterial kingdom. Therefore, further investigation of the initiation rescue processes and the factors

involved in diverse organisms from bacteria to human will be important for a full understanding of the

common principles and diverse mechanisms that ensure robust initiation of chromosomal DNA

replication.

430   **Materials and Methods**

Plasmids, proteins, and strains

The plasmids used in this study are listed in Key Resource Table. pKA234, pKW44-1, pNA135,

pBS*oriC*, pET22b(+)-priC, and pTKM601 were described previously (Aramaki et al., 2015; Ishida et

al., 2004; Kawakami et al., 2005; Keyamura et al., 2009; Kubota et al., 1997; Ozaki et al., 2008). For

the construction of pFH, an alanine substitution was introduced into pH136A with specific mutagenic

primer sets using the QuikChange site-directed mutagenesis protocol (Stratagene [Agilent], Agilent, La

Jolla, CA, United States) as described previously (Keyamura et al., 2007; Sakiyama et al., 2018).

WT DnaA and its derivative proteins were overproduced in *E. coli* strain KA450 from pKA234,

pKW44-1, or pFH and purified as described previously (Noguchi et al., 2015; Ozaki et al., 2008;

Sakiyama et al., 2017).

PriC protein was prepared as previously reported with minor modifications (Aramaki et al., 2013).

Briefly, PriC protein was overproduced in *E. coli* strain BL21-codonPlus-RIL by inducing its

expression from pET22b(+)-priC using 0.2 mM isopropyl-β-D-1-thiogalactopyranoside (IPTG). The

resulting cells were suspended in chilled buffer A (50 mM HEPES-NaOH [pH7.0], 10% sucrose, 1 mM

EDTA, 2 mM dithiothreitol (DTT), and 1 mM PMSF) and disrupted by sonication. Proteins in the

soluble fraction were precipitated with 0.24 g/mL ammonium sulfate, resuspended in a separation buffer

(50 mM imidazole [pH 7.0], 20% glycerol, 2 mM DTT, 1 mM EDTA, and 40 mM ammonium sulfate)

and loaded onto a 1 mL HiTrap® SP HP column (Cytiva, Uppsala, Sweden). Bound proteins were

eluted with a linear gradient from 0 to 1 M sodium chloride.

All *E. coli* strains used in this study are listed in Key Resource Table. KYA018, MIT125,

MIT162, SA103, NY20 and NY21 were described previously (Kasho & Katayama, 2013; Noguchi et

al., 2015; Noguchi & Katayama, 2016). Strains bearing mutant *oriC* (NY20L and NY20ATL) were

constructed using the λRed site-directed recombination system as previously described (Noguchi et al.,

2015; Sakiyama et al., 2017, 2022). NY20-frt strain and *oriC* mutant strains (NY21-frt, NY20L-frt, and

NY20ATL-frt) were constructed by eliminating *kan* from the NY20 strain and strains NY21, NY20L,

and NY20ATL. To remove the kanamycin-resistant cassette (*kan*) from *oriC*, FLP recombinase encoded

on pCP20 was used. Elimination of *kan* was verified by checking sensitivity to 50 µg/ml kanamycin in

LB agar plates.

Δ*priC*::*frt-kan* from JW0456-KC and Δ*rep*::*frt-kan* from JW5604-KC were introduced into

MG1655 by P1 transduction, resulting in the construction of KRC002 and FF001, respectively. For

construction of other Δ*priC*::*frt-kan* strains (KRC004, KRC005, NY20-*priC*, NY21-*priC*, NY20L-*priC*,

NY20ATL-*priC*), P1 phage lysates prepared from KRC002 were used for transduction of strains

MIT125, KYA018, NY20-frt, SYM21-frt, NY20L-frt, and NY20ATL-frt. Transductants were screened

on LB agar plates containing 50 µg/mL kanamycin. Δ*diaA*::*frt-kan* or *rnhA*::*cat* derivatives (KRC006,

MIT125c, and KRC004c) were constructed using a similar protocol except that P1 phage lysates from

SA103 (Δ*diaA*::*frt-kan*) or MIT162 (*rnhA*::*cat*) were used. KRC003 was generated by removal of *kan*

from KRC002.

<u>Buffers</u>

Buffer P contained 60 mM HEPES–KOH (pH 7.6), 0.1 mM zinc acetate, 8 mM magnesium

acetate, 30% [v/v] glycerol, and 0.32 mg/mL bovine serum albumin (BSA). Buffer N contained 50 mM

HEPES–KOH (pH 7.6), 2.5 mM magnesium acetate, 0.3 mM EDTA, 7 mM DTT, 0.007% [v/v] Triton

X-100, and 20% [v/v] glycerol. Form I* buffer contained 20 mM Tris-HCl (pH 7.5), 125 mM potassium

glutamate, 10 mM magnesium acetate, 8 mM DTT, and 0.5 mg/mL BSA.

<u>DUE unwinding assay</u>

DUE unwinding assays were performed essentially as described with minor modifications

(Sakiyama et al., 2022; Yoshida et al., 2023). Briefly, DnaA was incubated with ATP in buffer N on ice

for 3 min to generate ATP-DnaA. The indicated amount of PriC and ATP-DnaA or its mutant derivatives

was incubated for 9 min at 30°C in 10 μL buffer P containing 5 mM ATP, 125 mM potassium glutamate,

1.6 nM pBS*oriC*, and 42 nM IHF, followed by further incubation with 1.5 units of P1 nuclease (Wako)

for 5 min. The reaction was stopped by the addition of 1% SDS and 25 mM EDTA, and DNA was

purified by phenol-chloroform extraction and ethanol precipitation. One-third of each purified DNA

was digested with AlwNI (NEB), which yielded 2.6 kb and 1.0 kb fragments after DUE unwinding of

pBS*oriC*. The resultant DNA fragments were analyzed by 1% agarose gel electrophoresis with 1xTris-

acetate-EDTA buffer for 30 min at 100 V, followed by ethidium bromide staining. Gel images were

taken using the GelDoc GO imaging system (Bio-Rad Laboratories, Hercules, CA), and products

derived from unwound plasmids were quantified using ImageJ software.

Form I* assay

This assay was performed as previously described with minor modifications (Noguchi et al.,

2015; Sakiyama et al., 2022). The indicated amount of ATP-DnaA was incubated for 15 min at 30°C in

12.5 μL of Form I* buffer containing 3 mM ATP, 1.6 nM pBS*oriC*, 42 nM IHF, 400 nM His-DnaB, 400

nM DnaC, 76 nM GyrA, 100 nM His-GyrB, and 760 nM SSB. The reaction was stopped by the

addition of 0.5% SDS, and DNA was purified by phenol-chloroform extraction. Samples were analyzed

by 0.65% agarose gel electrophoresis with 0.5xTris-borate-EDTA buffer for 15 h at 23 V, followed by

ethidium bromide staining.

Flow cytometry analysis

Flow cytometry analysis was performed essentially as described (Kasho et al., 2014). Briefly,

cells were grown at 30°C in LB medium until the absorbance of the culture (A_{600}) reached 0.1. Portions

of the cultures were diluted a thousand-fold into 5 ml LB medium and incubated at 30°C until the

absorbance of the culture (A600) reached 0.1. The remaining portions were further incubated to

determine the doubling time (Td) by measuring A_{600} every 20 min. At A_{600} 0.1, aliquots of the cultures

were fixed in 70% ethanol to analyze cell mass using the Multisizer 3 Coulter counter (Beckman

Coulter, Brea, CA). The remaining cultures were further incubated for 4 h with 0.3 mg/mL rifampicin

and 0.01 mg/mL cephalexin to allow run-out replication of chromosomal DNA. The resultant cells were

fixed in 70% ethanol. After DNA staining with SYTOX Green (Thermo Fisher Scientific, Waltham,

MA), cellular DNA contents were analyzed on a FACS Calibur flow cytometer (BD Bioscience,

Franklin Lakes, NJ). The number of the origins/cell (ori/cell) was determined from the histograms of

flow cytometry analysis. The number of ori/cell was divided by the mean cell mass determined by cell

mass analysis, resulting in ori/mass ratios. The numbers of cells containing non-2^n copy number of *oriC*

were divided by the number of cells containing 2^n copy number of *oriC* to calculate the asynchrony

index (A.I.). For the analysis of cells bearing plasmids, LB medium was supplemented with 100 μg/mL

ampicillin.

Replication initiation frequency test using synchronized cultures

MIT125 (*dnaA46, tnaA*::Tn10) cells or its Δ*priC* derivative KRC004 were grown at 30°C, a

permissive temperature, in LB medium until the absorbance of the culture (A_{600}) reached 0.04, followed

by further incubation for 90 min at 42°C, a restrictive temperature. Aliquots of the cultures were fixed

in 70% ethanol as "Synchronization" samples. The remaining cultures were further incubated for 5 min

to initiate DNA replication, followed by further incubation for 4 h at 42°C with 0.3 mg/mL rifampicin

and 0.01 mg/mL cephalexin to allow run-out replication of chromosomal DNA. The resultant cells were

fixed in 70% ethanol as "5 min release" samples. After DNA staining with SYTOX Green (Life

Technologies), cellular DNA contents were analyzed on a FACS Calibur flow cytometer (BD

Bioscience). Ratios of the origins/cell (ori/cell) were determined from the histograms of flow cytometry

analysis.

For synchronization of *dnaC2* mutant cells (KYA018 and KRC005), cells were grown in LB

medium at 30°C until the A_{600} reached 0.04, followed by further incubation for 80 min at 37°C, at the

restrictive temperature. The resulting synchronized cultures were released for 5 min at 30°C and

incubated in the presence of 0.3 mg/mL rifampicin and 0.01 mg/mL cephalexin at 30°C to allow run-out

of chromosomal DNA replication.

qPCR for analysis of cSDR

Cells of MIT125 (*dnaA46, tnaA*::Tn10), MIT125c (*dnaA46, tnaA*::Tn10, *rnhA*::*cat*) and its

Δ*priC* derivatives KRC004 (*dnaA46, tnaA*::Tn10, Δ*priC*::*frt-kan*) and KRC004c (*dnaA46, tnaA*::Tn10,

rnhA::*cat*, Δ*priC*::*frt-kan*) were grown at 30°C in LB medium until the absorbance of the culture (A_{600})

reached 0.04 and aliquots of the cultures were withdrawn. The remaining samples were further

incubated for 90 min at 40°C and aliquots of the cultures were withdrawn. These samples were boiled

for 5 min at 95°C and the genome DNA was extracted. The levels of *oriC* (84.6 min), *ter* (32.4 min),

and *yapB* (50.5 min) were quantified by real-time qPCR using TB Green Premix Ex TaqII (Tli RNaseH

Plus) (TaKaRa, Shiga, Japan) and the following primers: ORI_1 and KWoriCRev for *oriC*, qoriK fw and

qoriK rev for *ter*, STM419 and STM420 for *yapB*.

Acknowledgments

We are grateful to Dr. Yoshio Abe for discussion and *priC*-overproducing plasmid, to Drs. Yusuke

Akama and Yukari Sakiyama for initial exploratory study related to this work, and to NRBP-E. coli at

NIG for bacterial strains.

Fundings

553 Japan Society for the Promotion of Science (JSPS KAKENHI) [JP17H03656, JP20H03212 and

554 JP23K27131].

References

Abe, Y., Jo, T., Matsuda, Y., Matsunaga, C., Katayama, T., & Ueda, T. (2007). Structure and function of

DnaA N-terminal domains: Specific sites and mechanisms in inter-DnaA interaction and in DnaB

helicase loading on *oriC*. *Journal of Biological Chemistry*, *282*(24), 17816–17827.

https://doi.org/10.1074/jbc.M701841200

Aramaki, T., Abe, Y., Furutani, K., Katayama, T., & Ueda, T. (2015). Basic and aromatic residues in the C-

terminal domain of PriC are involved in ssDNA and SSB binding. *Journal of Biochemistry*, *157*(6),

529–537. https://doi.org/10.1093/jb/mvv014

Aramaki, T., Abe, Y., Ohkuri, T., Mishima, T., Yamashita, S., Katayama, T., & Ueda, T. (2013). Domain

separation and characterization of PriC, a replication restart primosome factor in *Escherichia coli*.

Genes to Cells, *18*(9), 723–732. https://doi.org/10.1111/gtc.12069

Arias-Palomo, E., Puri, N., O'Shea Murray, V. L., Yan, Q., & Berger, J. M. (2019). Physical Basis for the

Loading of a Bacterial Replicative Helicase onto DNA. *Molecular Cell*, *74*(1), 173-184.e4.

https://doi.org/10.1016/j.molcel.2019.01.023

Asai, T., & Kogomal, T. (1994). MINIREVIEW D-Loops and R-Loops: Alternative Mechanisms for the

Initiation of Chromosome Replication in *Escherichia coli*. *JOURNAL OF BACTERIOLOGY*, *176*(7),

1807–1812. https://journals.asm.org/journal/jb

Bruand, C., Farache, M., McGovern, S., Ehrlich, S. D., & Polard, P. (2001). DnaB, DnaD and DnaI proteins

are components of the *Bacillus subtilis* replication restart primosome. *Molecular Microbiology*, *42*(1),

245–256. https://doi.org/10.1046/j.1365-2958.2001.02631.x

Bruand, C., Velten, M., McGovern, S., Marsin, S., Sérèna, C., Dusko Ehrlich, S., & Polard, P. (2005).

Functional interplay between the *Bacillus subtilis* DnaD and DnaB proteins essential for initiation and

re-initiation of DNA replication. *Molecular Microbiology*, *55*(4), 1138–1150.

https://doi.org/10.1111/j.1365-2958.2004.04451.x

Chodavarapu, S., & Kaguni, J. M. (2016). Replication Initiation in Bacteria. *Enzymes*, *39*, 1–30.

https://doi.org/10.1016/bs.enz.2016.03.001

Costa, A., & Diffley, J. F. X. (2022). The Initiation of Eukaryotic DNA Replication. *Annu. Rev. Biochem*, *91*,

107–131. https://doi.org/10.1146/annurev-biochem-072321

Duckworth, A. T., Ducos, P. L., McMillan, S. D., Satyshur, K. A., Blumenthal, K. H., Deorio, H. R., Larson,

J. A., Sandler, S. J., Grant, T., & Keck, J. L. (2023). Replication fork binding triggers structural

changes in the PriA helicase that govern DNA replication restart in *E. coli*. *Nature Communications*,

14(1), 1–13. https://doi.org/10.1038/s41467-023-38144-x

Duderstadt, K. E., Chuang, K., & Berger, J. M. (2011). DNA stretching by bacterial initiators promotes

replication origin opening. *Nature*, *478*(7368), 209–213. https://doi.org/10.1038/nature10455

Erzberger, J. P., Mott, M. L., & Berger, J. M. (2006). Structural basis for ATP-dependent DnaA assembly and

replication-origin remodeling. *Nature Structural and Molecular Biology*, *13*(8), 676–683.

https://doi.org/10.1038/nsmb1115

Fang, L., Davey, M. J., & O'donnell, M. (1999). Replisome Assembly at *oriC*, the Replication Origin of *E.*

coli, Reveals an Explanation for Initiation Sites outside an Origin. *Molecular Cell*, *4*, 541–553.

Felczak, M. M., & Kaguni, J. M. (2004). The box VII motif of *Escherichia coli* DnaA protein is required for

DnaA oligomerization at the *E. coli* replication origin. *Journal of Biological Chemistry*, *279*(49),

51156–51162. https://doi.org/10.1074/jbc.M409695200

Flores, M. J., Ehrlich, S. D., & Michel, B. (2002). Primosome assembly requirement for replication restart in

the *Escherichia coli holDG10* replication mutant. *Molecular Microbiology*, *44*(3), 783–792.

https://doi.org/10.1046/j.1365-2958.2002.02913.x

Fujikawa, N., Kurumizaka, H., Nureki, O., Terada, T., Shirouzu, M., Katayama, T., & Yokoyama, S. (2003).

Structural basis of replication origin recognition by the DnaA protein. *Nucleic Acids Research*, *31*(8),

2077–2086. https://doi.org/10.1093/nar/gkg309

Grimwade, J. E., & Leonard, A. C. (2021). Blocking, Bending, and Binding: Regulation of Initiation of

Chromosome Replication During the *Escherichia coli* Cell Cycle by Transcriptional Modulators That

Interact With Origin DNA. In *Frontiers in Microbiology* (Vol. 12, p. 732270). Frontiers Media S.A.

https://doi.org/10.3389/fmicb.2021.732270

Hayashi, C., Miyazaki, E., Ozaki, S., Abe, Y., & Katayama, T. (2020). DnaB helicase is recruited to the

replication initiation complex via binding of DnaA domain I to the lateral surface of the DnaB N-

terminal domain. *Journal of Biological Chemistry*, *295*(32), 1131–1143.

https://doi.org/10.1074/jbc.ra120.014235

Heller, R. C., & Marians, K. J. (2005a). The disposition of nascent strands at stalled replication forks dictates

the pathway of replisome loading during restart. *Molecular Cell*, *17*(5), 733–743.

https://doi.org/10.1016/j.molcel.2005.01.019

Heller, R. C., & Marians, K. J. (2005b). Unwinding of the nascent lagging strand by Rep and PriA enables

the direct restart of stalled replication forks. *Journal of Biological Chemistry*, *280*(40), 34143–34151.

https://doi.org/10.1074/jbc.M507224200

Heller, R. C., & Marians, K. J. (2006). Replication fork reactivation downstream of a blocked nascent

leading strand. *Nature*, *439*(7076), 557–562. https://doi.org/10.1038/nature04329

Hinds, T., & Sandler, S. J. (2004). Allele specific synthetic lethality between *priC* and *dnaAts* alleles at the

permissive temperature of 30°C in *E. coli* K-12. *BMC Microbiology*, *4*, 1–9.

https://doi.org/10.1186/1471-2180-4-47

Ishida, T., Akimitsu, N., Kashioka, T., Hatano, M., Kubota, T., Ogata, Y., Sekimizu, K., & Katayama, T. (2004). DiaA, a novel DnaA-binding protein, ensures the timely initiation of *Escherichia coli* chromosome replication. *Journal of Biological Chemistry*, *279*(44), 45546–45555. https://doi.org/10.1074/jbc.M402762200

Jameson, K. H., & Wilkinson, A. J. (2017). Control of initiation of DNA replication in *Bacillus subtilis* and *Escherichia coli*. In *Genes* (Vol. 8, Issue 1, p. 22). MDPI AG. https://doi.org/10.3390/genes8010022

Kasho, K., Fujimitsu, K., Matoba, T., Oshima, T., & Katayama, T. (2014). Timely binding of IHF and Fis to *DARS2* regulates ATP-DnaA production and replication initiation. *Nucleic Acids Research*, *42*(21), 13134–13149. https://doi.org/10.1093/nar/gku1051

Kasho, K., & Katayama, T. (2013). DnaA binding locus *datA* promotes DnaA-ATP hydrolysis to enable cell cycle-coordinated replication initiation. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(3), 936–941. https://doi.org/10.1073/pnas.1212070110

Kasho, K., Ozaki, S., & Katayama, T. (2023). IHF and Fis as *Escherichia coli* Cell Cycle Regulators: Activation of the Replication Origin *oriC* and the Regulatory Cycle of the DnaA Initiator. In *International Journal of Molecular Sciences* (Vol. 24, Issue 14, p. 11572). Multidisciplinary Digital Publishing Institute (MDPI). https://doi.org/10.3390/ijms241411572

Kawakami, H., Keyamura, K., & Katayama, T. (2005). Formation of an ATP-DnaA-specific initiation complex requires DnaA arginine 285, a conserved motif in the AAA+ protein family. *Journal of Biological Chemistry*, *280*(29), 27420–27430. https://doi.org/10.1074/jbc.M502764200

Keyamura, K., Abe, Y., Higashi, M., Ueda, T., & Katayama, T. (2009). DiaA dynamics are coupled with changes in initial origin complexes leading to helicase loading. *Journal of Biological Chemistry*, *284*(37), 25038–25050. https://doi.org/10.1074/jbc.M109.002717

Keyamura, K., Fujikawa, N., Ishida, T., Ozaki, S., Su'etsugu, M., Fujimitsu, K., Kagawa, W., Yokoyama, S., Kurumizaka, H., & Katayama, T. (2007). The interaction of DiaA and DnaA regulates the replication cycle in *E. coli* by directly promoting ATP-DnaA-specific initiation complexes. *Genes and Development*, *21*(16), 2083–2099. https://doi.org/10.1101/gad.1561207

Kogoma, T. (1997). Stable DNA Replication: Interplay between DNA Replication, Homologous Recombination, and Transcription. *MICROBIOLOGY AND MOLECULAR BIOLOGY REVIEWS*, *61*(2), 212–238. https://journals.asm.org/journal/mmbr

Kubota, T., Katayama, T., Ito, Y., Mizushima, T., & Sekimizu, K. (1997). Conformational Transition of DnaA Protein by ATP: Structural Analysis of DnaA Protein, the Initiator of *Escherichia coli* Chromosome Replication. *BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS*, *232*, 130–135.

Lopper, M., Boonsombat, R., Sandler, S. J., & Keck, J. L. (2007). A Hand-Off Mechanism for Primosome Assembly in Replication Restart. *Molecular Cell*, *26*(6), 781–793. https://doi.org/10.1016/j.molcel.2007.05.012

Maduike, N. Z., Tehranchi, A. K., Wang, J. D., & Kreuzer, K. N. (2014). Replication of the *Escherichia coli* chromosome in RNase HI-deficient cells: Multiple initiation regions and fork dynamics. *Molecular Microbiology*, *91*(1), 39–56. https://doi.org/10.1111/mmi.12440

Masai, H., Asai, T., Kubota, Y., Arai, K. I., & Kogoma, T. (1994). *Escherichia coli* PriA protein is essential

for inducible and constitutive stable DNA replication. *EMBO Journal*, *13*(22), 5338–5345.

https://doi.org/10.1002/j.1460-2075.1994.tb06868.x

McMillan, S. D., & Keck, J. L. (2024). Biochemical characterization of *Escherichia coli* DnaC variants that

alter DnaB helicase loading onto DNA. *Journal of Biological Chemistry*, *300*(5), 107275.

https://doi.org/10.1016/j.jbc.2024.107275

Michel, B., & Sandler, S. J. (2017). Replication restart in bacteria. In *Journal of Bacteriology* (Vol. 199,

Issue 13, pp. e00102-17). American Society for Microbiology. https://doi.org/10.1128/JB.00102-17

Miller, D. T., Grimwade, J. E., Betteridge, T., Rozgaja, T., Torgue, J.-C., Leonard, A. C., & Helinski, D. R.

(2009). Bacterial origin recognition complexes direct assembly of higher-order DnaA oligomeric

structures. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(44),

18479–18484. www.pnas.orgcgidoi10.1073pnas.0909472106

Nagata, K., Okada, A., Ohtsuka, J., Ohkuri, T., Akama, Y., Sakiyama, Y., Miyazaki, E., Horita, S., Katayama,

T., Ueda, T., & Tanokura, M. (2020). Crystal structure of the complex of the interaction domains of

Escherichia coli DnaB helicase and DnaC helicase loader: Structural basis implying a distortion-

accumulation mechanism for the DnaB ring opening caused by DnaC binding. *Journal of*

Biochemistry, *167*(1), 1–14. https://doi.org/10.1093/jb/mvz087

Natrajan, G., Hall, D. R., Thompson, A. C., Gutsche, I., & Terradot, L. (2007). Structural similarity between

the DnaA-binding proteins HobA (HP1230) from *Helicobacter pylori* and DiaA from *Escherichia coli*.

Molecular Microbiology, *65*(4), 995–1005. https://doi.org/10.1111/j.1365-2958.2007.05843.x

Né Dicte Michel, B., Flores, M.-J., Viguera, E., Grompone, G., Seigneur, M., & Bidnenko, V. (2001). Rescue

of arrested replication forks by homologous recombination. *Proceedings of the National Academy of*

Sciences of the United States of America, *17*(15), 8181–8188.

www.pnas.orgcgidoi10.1073pnas.111008798

Nguyen, B., Shinn, M. K., Weiland, E., & Lohman, T. M. (2021). Regulation of *E. coli* Rep helicase activity

by PriC. *Journal of Molecular Biology*, *433*(15), 167072. https://doi.org/10.1016/j.jmb.2021.167072

Noguchi, Y., & Katayama, T. (2016). The *Escherichia coli* cryptic prophage protein YfdR binds to DnaA and

initiation of chromosomal replication is inhibited by overexpression of the gene cluster *yfdQ-yfdR-*

yfdS-yfdT. *Frontiers in Microbiology*, *7*(MAR), 1–21. https://doi.org/10.3389/fmicb.2016.00239

Noguchi, Y., Sakiyama, Y., Kawakami, H., & Katayama, T. (2015). The Arg fingers of key DnaA protomers

are oriented inward within the replication origin *oriC* and stimulate DnaA subcomplexes in the

initiation complex. *Journal of Biological Chemistry*, *290*(33), 20295–20312.

https://doi.org/10.1074/jbc.M115.662601

Nozaki, S., & Ogawa, T. (2008). Determination of the minimum domain II size of *Escherichia coli* DnaA

protein essential for cell viability. *Microbiology*, *154*(11), 3379–3384.

https://doi.org/10.1099/mic.0.2008/019745-0

O'Donnell, M., Langston, L., & Stillman, B. (2013). Principles and concepts of DNA replication in bacteria,

archaea, and eukarya. *Cold Spring Harbor Perspectives in Biology*, *5*(7), a010108.

https://doi.org/10.1101/cshperspect.a010108

Ozaki, S., & Katayama, T. (2012). Highly organized DnaA-*oriC* complexes recruit the single-stranded DNA for replication initiation. *Nucleic Acids Research*, *40*(4), 1648–1665. https://doi.org/10.1093/nar/gkr832

Ozaki, S., Kawakami, H., Nakamura, K., Fujikawa, N., Kagawa, W., Park, S. Y., Yokoyama, S., Kurumizaka, H., & Katayama, T. (2008). A common mechanism for the ATP-DnaA-dependent formation of open complexes at the replication origin. *Journal of Biological Chemistry*, *283*(13), 8351–8362. https://doi.org/10.1074/jbc.M708684200

Ozaki, S., Noguchi, Y., Hayashi, Y., Miyazaki, E., & Katayama, T. (2012). Differentiation of the DnaA-*oriC* subcomplex for DNA unwinding in a replication initiation complex. *Journal of Biological Chemistry*, *287*(44), 37458–37471. https://doi.org/10.1074/jbc.M112.372052

Rozgaja, T. A., Grimwade, J. E., Iqbal, M., Czerwonka, C., Vora, M., & Leonard, A. C. (2011). Two oppositely oriented arrays of low-affinity recognition sites in *oriC* guide progressive binding of DnaA during *Escherichia coli* pre-RC assembly. *Molecular Microbiology*, *82*(2), 475–488. https://doi.org/10.1111/j.1365-2958.2011.07827.x

Sakiyama, Y., Kasho, K., Noguchi, Y., Kawakami, H., & Katayama, T. (2017). Regulatory dynamics in the ternary DnaA complex for initiation of chromosomal replication in *Escherichia coli*. *Nucleic Acids Research*, *45*(21), 12354–12373. https://doi.org/10.1093/nar/gkx914

Sakiyama, Y., Nagata, M., Yoshida, R., Kasho, K., Ozaki, S., & Katayama, T. (2022). Concerted actions of DnaA complexes with DNA-unwinding sequences within and flanking replication origin *oriC* promote DnaB helicase loading. *Journal of Biological Chemistry*, *298*(6), 102051. https://doi.org/10.1016/j.jbc.2022.102051

Sakiyama, Y., Nishimura, M., Hayashi, C., Akama, Y., Ozaki, S., & Katayama, T. (2018). The DnaA AAA+ Domain His136 residue directs DnaB replicative helicase to the unwound region of the replication origin, *oriC*. *Frontiers in Microbiology*, *9*(AUG), 1–17. https://doi.org/10.3389/fmicb.2018.02017

Sandler, S. J. (2000). Multiple Genetic Pathways for Restarting DNA Replication Forks in *Escherichia coli* K-12. *Genetics*, *155*, 487–497. https://academic.oup.com/genetics/article/155/2/487/6048048

Sandler, S. J., Marians, K. J., Zavitz, K. H., Coutu, J., Parent, M. A., & Clark, A. J. (1999). *dnaC* mutations suppress defects in DNA replication- and recombination-associated functions in *priB* and *priC* double mutants in *Escherichia coli* K-12. *Molecular Microbiology*, *34*(1), 91–101. https://doi.org/10.1046/j.1365-2958.1999.01576.x

Shibata, E., & Dutta, A. (2020). A human cancer cell line initiates DNA replication normally in the absence of ORC5 and ORC2 proteins. *Journal of Biological Chemistry*, *295*(50), 16949–16959. https://doi.org/10.1074/jbc.RA120.015450

Shimizu, M., Noguchi, Y., Sakiyama, Y., Kawakami, H., Katayama, T., & Takada, S. (2016). Near-atomic structural model for bacterial DNA replication initiation complex and its functional insights. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(50), E8021–E8030. https://doi.org/10.1073/pnas.1609649113

Stepankiw, N., Kaidow, A., Boye, E., & Bates, D. (2009). The right half of the *Escherichia coli* replication origin is not essential for viability, but facilitates multi-forked replication. *Molecular Microbiology*,

74(2), 467–479. https://doi.org/10.1111/j.1365-2958.2009.06877.x

Struhl, K., Shibata, E., Kiran, M., Shibata, Y., Singh, S., Kiran, S., & Dutta, A. (2016). Two subunits of

human ORC are dispensable for DNA replication and proliferation. *ELife*, *5*, e19084.

https://doi.org/10.7554/eLife.19084.001

Wegrzyn, K., & Konieczny, I. (2023). Toward an understanding of the DNA replication initiation in bacteria.

In *Frontiers in Microbiology* (Vol. 14, p. 1328842). Frontiers Media SA.

https://doi.org/10.3389/fmicb.2023.1328842

Wessel, S. R., Cornilescu, C. C., Cornilescu, G., Metz, A., Leroux, M., Hu, K., Sandler, S. J., Markley, J. L.,

& Keck, J. L. (2016). Structure and function of the PriC DNA replication restart protein. *Journal of*

Biological Chemistry, *291*(35), 18384–18396. https://doi.org/10.1074/jbc.M116.738781

Wessel, S. R., Marceau, A. H., Massoni, S. C., Zhou, R., Ha, T., Sandler, S. J., & Keck, J. L. (2013). PriC-

mediated DNA replication restart requires PriC complex formation with the single-stranded DNA-

binding protein. *Journal of Biological Chemistry*, *288*(24), 17569–17578.

https://doi.org/10.1074/jbc.M113.478156

Windgassen, T. A., & Keck, J. L. (2016). An aromatic-rich loop couples DNA binding and ATP hydrolysis in

the PriA DNA helicase. *Nucleic Acids Research*, *44*(20), 9745–9757.

https://doi.org/10.1093/nar/gkw690

Windgassen, T. A., Leroux, M., Satyshur, K. A., Sandler, S. J., & Keck, J. L. (2018). Structure-specific DNA

replication-fork recognition directs helicase and replication restart activities of the PriA helicase.

Proceedings of the National Academy of Sciences of the United States of America, *115*(39), E9075–

E9084. https://doi.org/10.1073/pnas.1809842115

Withers, H. L., & Bernander, R. (1998). Characterization of *dnaC2* and *dnaC28* Mutants by Flow Cytometry.

JOURNAL OF BACTERIOLOGY, *180*(7), 1624–1631. https://journals.asm.org/journal/jb

Yoshida, R., Ozaki, S., Kawakami, H., & Katayama, T. (2023). Single-stranded DNA recruitment mechanism

in replication origin unwinding by DnaA initiator protein and HU, an evolutionary ubiquitous nucleoid

protein. *Nucleic Acids Research*, *51*(12), 6286–6306. https://doi.org/10.1093/nar/gkad389

Zawilak-Pawlik, A., Donczew, R., Szafrański, S., MacKiewicz, P., Terradot, L., & Zakrzewska-Czerwińska,

J. (2011). DiaA/HobA and DnaA: A pair of proteins co-evolved to cooperate during bacterial orisome

assembly. *Journal of Molecular Biology*, *408*(2), 238–251. https://doi.org/10.1016/j.jmb.2011.02.045

Figure legend

Figure 1 Schematic structures of *oriC* and DnaA and the replication initiation mechanism.

(A) Overall structure of *oriC* with the AT-rich cluster. *oriC* (245 bp) includes the Duplex-Unwinding Element (DUE, purple bar) and the DnaA-Oligomerization Region (DOR, gray bar). DUE is composed of three AT-rich 13-mer repeats termed L, M, and R. Dark purple boxes indicate the specific sequences involved in DnaA-ssDNA binding (TT[A/G]T(T)). The AT-cluster (purple dots) flanking the DUE outside *oriC* is a supplementary unwinding region. The DOR contains 12 DnaA boxes (filled and open arrowheads representing sites with the full consensus TTA[A/T]NCACA sequence and sites with mismatches, respectively) and an IHF-binding region (IBR, green box). The DOR is subdivided into Left, Middle, and Right subregions. (B) Domains of DnaA. Domains I-IV are shown schematically, with amino acid residue numbers shown in each bracket. H/B motifs (V211 and R245, squares) and Arginine finger (R285, triangle) are indicated. The major functions of each domain are described on the right side of the structure. (C) DUE unwinding by the ssDUE recruitment mechanism. DnaA and IHF are indicated by red and green diagrams, respectively. When ATP-DnaA oligomerizes, DUE is unwound unstably by thermal motion and torsional stress. The M/R region of the DUE-upper strand (purple line) binds to R1-DnaA and R5M-DnaA via IHF-induced DNA bending, resulting in stable DUE unwinding. In addition to M/R-DUE, L-DUE is moderately unwound and interacts with the Right-DnaA subcomplex. (D) Mechanism for DnaB loading. Each Left-DnaA and Right-DnaA subcomplex binds to a DnaB hexamer (blue ovals) complexed with DnaC (yellow circles). (E) PriC-dependent fork rescue pathways. PriC (vivid red circle) promotes DnaB reloading to the abandoned fork, which exposes ssDNA regions. PriC binds to ssDNA at the replication fork, which remodels the SSB-ssDNA complex (for simplicity, SSB is omitted) for DnaC-dependent reloading of DnaB (*upper panel*). If the ssDNA region is accompanied with a nascent lagging strand, Rep helicase (green shape) is recruited by PriC and unwinds the lagging strand to expose the ssDNA region for DnaB loading (the lower case).

Figure 2. Requirement of *priC* for DNA replication initiation in *dnaA46* and *dnaC2* strains.

(A) Colony forming abilities. MG1655 and its derivatives MIT125 (MG1655 *dnaA46 tnaA*::Tn*10*), KYA018 (MG1655 *dnaC2 zjj18*::*cat*), KRC002 (MG1655 Δ*priC*::*frt-kan*), KRC004 (MG1655 *dnaA46 tnaA*::Tn*10* Δ*priC*::*frt-kan*), and KRC005 (MG1655 *dnaC2 zjj18*::*cat* Δ*priC*::*frt-kan*) were grown overnight. 10-fold-serial dilutions of the overnight cultures (~10^9 cells/mL) were incubated for 16 h at 30℃, or for 14 h at 35/37/42℃ on LB agar medium. Three independent experiments were performed. +, WT *priC*; −, Δ*priC*::*frt-kan*. The images of representative plates are shown.

(B) DNA replication initiation in synchronized *dnaA46* cells with or without WT *priC*. Exponentially growing MIT125 (MG1655 *dnaA46 tnaA*::Tn*10*) (+) and KRC004 (MG1655 *dnaA46 tnaA*::Tn*10*

Δ*priC*::*frt-kan*) (−) cells were synchronized to the pre-initiation stage by incubation for 90 min at 42°C

in LB medium, followed by incubation for 5 min at 30°C to initiate DNA replication, which was further

followed by incubation with rifampicin and cephalexin for 4 h at 42°C to allow run-out of ongoing DNA

replication. Samples were withdrawn after synchronization and run-out replication. The DNA contents of

these samples were measured by flow cytometry. Representative histograms of three independent

experiments are shown. For each sample, the mean number of cells with more than one chromosome

equivalent after a 5 min release was quantified and is shown as a percentage with standard deviations.

(C) DNA replication initiation in synchronized *dnaC2* cells with or without WT *priC*. Exponentially

growing KYA018 (MG1655 *dnaC2 zjj18*::*cat*) (+) and KRC005 (MG1655 *dnaC2 zjj18*::*ca* Δ*priC*::*frt-*

kan) (−) cells were synchronized to the pre-initiation stage by incubation for 80 min at 37°C in LB

medium, followed by incubation for 5 min at 30°C and further incubation with rifampicin and cephalexin

as described above. Samples were withdrawn and analyzed as described above.

Figure 3. Role of PriC in growth of DnaA mutant defective in interactions with DnaB.

Colony forming abilities of MIT125 (MG1655 *dnaA46 tnaA*::Tn*10*) and KRC004 (MG1655 *dnaA46*

tnaA::Tn*10* Δ*priC*::*frt-kan*) cells bearing pING1 plasmid vector (−) or its derivatives pKA234 expressing

WT DnaA (WT) or pFH expressing DnaA F46A H136A (F46A H136A). Cells were grown at 30°C

overnight and 10-fold-serial dilutions of the cultures (~10^9 cells/mL) were incubated on LB agar plates

containing ampicillin for 16 h at 30°C and for 14 h at 42°C, respectively. Four independent experiments

were performed. +, WT *priC*; −, Δ*priC*::*frt-kan*.

Figure 4. PriC stimulation of DNA replication initiation inhibited by DiaA oversupply.

(A) Colony forming abilities of MG1655, KRC002 (MG1655 Δ*priC*::*frt-kan*), and FF001 (MG1655

Δ*rep*::*frt-kan*) cells bearing pBR322 or pNA135 (pBR322 derivatives carrying *diaA*). Overnight cultures

(~10^9 cells/mL) of MG1655 and KRC002 cells bearing pBR322 or pNA135 were 10-fold serially diluted

and incubated on LB agar plates containing ampicillin for 40 h at 25°C. The results of three independent

experiments were consistent, and one which is shown.

(B) Flow cytometry analyses of MG1655 and KRC002 (MG1655 Δ*priC*::*frt-kan*) cells bearing pBR322

or pNA135. Cells were exponentially grown at 30°C in LB medium with ampicillin, followed by further

incubation with rifampicin and cephalexin to allow run-out of chromosomal DNA replication. DNA

contents were quantified by flow cytometry, and cell size (mass) at the time of drug addition was

measured by a Coulter counter. Mean mass, ori/mass ratio, doubling time (Td), and asynchrony index

(A.I.: the percentage of cell numbers with non-2^n copies of *oriC* per the cell numbers with 2^n copies of

oriC) of each strain are indicated at the top right of each panel with standard deviations. Three to four

independent experiments were performed. +, WT *priC*; −, Δ*priC*::*frt-kan*.

(C) Flow cytometry analyses of MG1655, KRC003 (MG1655 Δ*priC*::*frt*), SA103 (MG1655 Δ*diaA*::*frt-*

kan), and KRC006 (MG1655 Δ*priC*::*frt*, Δ*diaA*::*frt-kan*) cells. Cells were grown and analyzed as

described above. Three independent experiments were performed. For "*priC*" column, +, WT *priC*;

−, Δ*priC*::*frt*. For "*diaA*" column, +, WT *diaA*; −, Δ*diaA*::*frt-kan.*

Figure 5. Stimulation of replication initiation by PriC in Right-DnaA subcomplex-defective cells

(A) Schematic structure of Left-*oriC* and R4*Tma oriC*. AT-rich region, DUE, and DOR are indicated by

a gray dotted box, a gray box, and an open box, respectively. Filled and open arrowheads in the DOR

show DnaA boxes with the full consensus sequence or mismatches. In addition, a striped box indicates

the IBR. The regions of each mutant *oriC* are indicated below the structure as open boxes. The position

of the *Tma*DnaA box substitution is indicated by a yellow arrowhead.

(B) Colony forming abilities of WT *oriC* and Left-*oriC* cells with or without WT *priC*. Overnight cultures

(~10^9 cells/mL) of NY20-frt (WT *oriC*; WT) and NY20L-frt (Left-*oriC*; Left) cells with (+) or without

(−) WT *priC* were 10-fold serially diluted and incubated on LB agar plates for 24 h at 25°C, for 16 h at

30°C, and for 14 h at 37°C. +, WT *priC*; −, Δ*priC*::*frt-kan*. Three independent experiments were

performed.

(C) Colony forming abilities of WT *oriC* and *oriC* R4-box mutant cells with or without WT *priC*. NY20-

frt (WT *oriC*) and NY24-frt (*oriC* with R4*Tma* substitution, R4*Tma*) cells with (+) or without (−) WT

priC were grown as described in panel A. +, WT *priC*; −, Δ*priC*::*frt-kan*. Four independent experiments

were performed.

(D) Flow cytometry analyses of WT and R4*Tma* cells with or without WT *priC*. Cells were exponentially

grown at 30°C in LB medium with ampicillin and analyzed as described in Figure 4. Representative

histograms from four independent experiments are shown. Mean mass, ori/mass, and Td of each strain

are indicated at the top right of each panel with standard deviations.

Figure 6. Stimulation of replication initiation by PriC in cells with a deletion in the ssDNA-

expanding region of *oriC*

(A) Schematic structure of subATL *oriC* as shown in Figure 5. The *oriC* region including the subATL

oriC is indicated below the structure as a white box.

(B) Colony forming abilities of WT *oriC* and *oriC* subATL mutant cells with or without WT *priC*. NY20-

frt (WT *oriC*) and NY20ATL-frt (*oriC* lacking AT and L sequences, subATL) with (+) or without (−) WT

priC were grown as described in Figure 5. +, WT *priC*; −, Δ*priC*::*frt-kan.* Three independent experiments

were performed.

(C) Flow cytometry analyses of WT *oriC* and *oriC* subATL mutant cells with or without WT *priC*. Cells were exponentially grown at 30°C in LB medium and analyzed as described in Figure 4. Representative histograms from three independent experiments are shown. Mean mass, ori/mass, and Td of each strain are indicated at the top right of each panel with standard deviations.

Figure 7. DnaB loading onto the ssDUE by PriC.

(A and B) *In vitro* reconstituted system for DnaB loading. Form I of *oriC* plasmid (pBS*oriC*; 1.6 nM) was incubated for 15 min at 30°C with the indicated amount of PriC in the presence of ATP-DnaA or its mutant derivatives, DnaB, DnaC, IHF, gyrase, and SSB. The resulting plasmids were purified and analyzed using agarose gel electrophoresis. A representative image of three independent experiments is shown in the black/white-inverted mode (A). Band intensities of each lane in the gel image were analyzed by densitometric scanning. The percentages of form I* *oriC* plasmid per input DNA are shown as 'Form I* (%)'. Mean and standard deviations are shown (B). Abbreviations in panel B: −, no addition of DnaA; WT, wild-type DnaA; FH, DnaA F46A H136A; VA, DnaA V211A.

 (C and D) *In vitro* reconstituted system containing DiaA. Form I of pBS*oriC* plasmid was used as described above except for the addition of the indicated amounts of DiaA. A representative image of three independent experiments is shown in black/white-inverted mode (C). Band intensities of each lane in the gel image were analyzed by densitometric scanning. The percentages of form I**oriC* plasmid together with the mean and standard deviations are shown as described above (D).

Figure 8. PriC stimulation of cSDR-dependent cell growth.

(A) Colony forming abilities. Cells of MIT125 (MG1655 *dnaA46*), MIT125c (MG1655 *dnaA46 rnhA*::*cat*), KRC004 (MG1655 *dnaA46* Δ*priC*::*frt-kan*), and KRC004c (MG1655 *dnaA46* Δ*priC*::*frt-kan rnhA*::*cat*) were grown at 30°C overnight and 10-fold-serial dilutions of the overnight cultures (~10^9 cells/mL) were incubated for 16 h at 30°C and for 36 h at 40°C on LB agar plates. Three independent experiments were performed. +, WT; −, deletion.

(B) *oriC* and *ter* copy numbers. MIT125, MIT125c, KRC004, and KRC004c cells growing exponentially at 30°C were further incubated for 90 min at 40°C in LB medium. Samples were withdrawn before and after the 40°C incubation. The genome DNA of each sample was extracted by boiling the cells for 5 min at 95°C. The relative copy numbers of *oriC* (84.6 min) and *ter* (32.4 min) to that of *yapB* (50.5 min) were quantified using real-time qPCR. The data and averages of three independent experiments are shown with standard deviations.

Figure 9. Model for PriC-promoted replication initiation.

PriC-promoted rescue of blocked DnaA-*oriC* dependent replication initiation. *oriC*, DnaA, IHF, DnaB, and DnaC are illustrated as depicted in Figure 1C. When tight DnaA binding of DiaA (dark gray circle) (or DnaA F46A H136A double mutations) inhibits DnaB recruitment, PriC (Brilliant red circle) binds to a stably unwound DUE strand, recruiting a DnaB-DnaC complex to lower strand of DUE instead of Left-DnaA subcomplex. When a sufficient ssDNA region is exposed, PriC then recruits the DnaB-DnaC complex to the opposite strand.

Figure 4-figure supplement 1. Colony formation of cells carrying pBR322 or pNA135 with WT *priC* or Δ*priC*::*frt-kan*.

Colony forming abilities of MG1655 or KRC002 (MG1655 Δ*priC*::*frt-kan*) cells bearing pBR322 or pNA135 (pBR322 derivatives carrying *diaA*). Overnight cultures (~10^9 cells/mL) of MG1655 and KRC002 cells carrying pBR322 or pNA135 were 10-fold serially diluted and incubated on LB agar plates containing ampicillin for 14 h at 30–42°C. +, WT *priC*; −, Δ*priC*::*frt-kan*. The results of three independent experiments were consistent, and one representative result is shown.

Figure 7-figure supplement 1. DUE unwinding activities of DnaA mutant deriviatives in the presence and absence of PriC.

(A and B) *in vitro* DUE unwinding assays using DnaA mutant derivatives. Form I of *oriC* (pBS*oriC*; 1.6 nM) was incubated for 9 min at 30°C with the indicated amount of DnaA or its mutant derivatives in the presence of IHF (42 nM), followed by incubation with P1 nuclease (1.5 units) and further incubation with AlwNI before agarose gel electrophoresis. Note that 2.6 kb and 1.0 kb fragments are observed with restriction enzyme digestion when the ssDNA-specific P1 nuclease cleaves 3.6 kb pBS*oriC* at the DUE. A representative image of two independent experiments is shown in black/white-inverted mode (A). The sizes of each band are indicated at the left side of the image. Bands intensities of each lane in the gel image were analyzed by densitometric scanning. The percentages of P1 nuclease-digested *oriC* DNA per input DNA molecules are shown as 'DUE unwinding (%)'. Mean and data are shown (B). Abbreviations in panel B: −, no addition of DnaA; WT, wild-type DnaA; FH, DnaA F46A H136A; VA, DnaA V211A. (C and D) *In vitro* DUE unwinding assay with PriC. pBS*oriC* was used as described above except for PriC addition. A representative image of two independent experiments is shown in black/white-inverted mode (C). Data were analyzed as described above (D). n.t., not tested.

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
strain, strain background (*Escherichia coli*)	BW25113 Δ*priC*::*frt-kan*	Keio collection	JW0456-KC	
strain, strain background (*Escherichia coli*)	BW25113 Δ*rep*::*frt-kan*	Keio collection	JW5604-KC	
strain, strain background (*Escherichia coli*)	MG1655 *dnaC2 zjj18*::*cat*	Kasho and Katayama, 2013	KYA018	
strain, strain background (*Escherichia coli*)	MG1655 *dnaA46 tnaA*::Tn10	Noguchi, *et al.*, 2016	MIT125	
strain, strain background (*Escherichia coli*)	MG1655 *rnhA*::*cat*	Noguchi, *et al.*, 2016	MIT162	
strain, strain background (*Escherichia coli*)	MG1655 Δ*diaA*::*frt-kan*	Noguchi, *et al.*, 2016	SA103	
strain, strain background (*Escherichia coli*)	MG1655 Δ*priC*::*frt-kan*	This work	KRC002	MG1655 x P1 JW0456-KC
strain, strain background (*Escherichia coli*)	MG1655 Δ*priC*::*frt*	This work	KRC003	KRC002 x pCP20

strain, strain background (*Escherichia coli*)	MIT125 Δ*priC*::*frt-kan*	This work	KRC004	MIT125 x P1 JW0456-KC
strain, strain background (*Escherichia coli*)	KYA018 Δ*priC*::*frt-kan*	This work	KRC005	KYA018 x P1 JW0456-KC
strain, strain background (*Escherichia coli*)	KRC003 Δ*diaA*::*frt-kan*	This work	KRC006	KRC003 x P1 SA103
strain, strain background (*Escherichia coli*)	MG1655 Δ*rep*::*frt-kan*	This work	FF001	MG1655 x P1 JW5604-KC
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt-kan*	Noguchi, *et al*, 2015	NY20	
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt-kan* *oriC*ΔR4 box::*Tma*DnaA box	Noguchi, *et al*, 2015	NY21	
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt-kan* *oriC*ΔMiddle-Right DOR	This work	NY20L	
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt-kan* *oriC*ΔAT-L region	This work	NY20ATL	

strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt*	Yoshida, *et al*, 2023	NY20-frt	
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt* *oriC*ΔR4 box::*Tma*DnaA box	This work	NY21-frt	NY21 x pCP20
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt* *oriC*ΔMiddle-Right DOR	This work	NY20L-frt	NY20L x pCP20
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt* *oriC*ΔAT-L region	This work	NY20ATL-frt	NY20ATL x pCP20
strain, strain background (*Escherichia coli*)	NY20-*frt* Δ*priC*::*frt-kan*	This work	NY20-*priC*	NY20-frt x P1 KRC002
strain, strain background (*Escherichia coli*)	NY21-*frt* Δ*priC*::*frt-kan*	This work	NY21-*priC*	NY21-frt x P1 KRC002
strain, strain background (*Escherichia coli*)	NY20L-*frt* Δ*priC*::*frt-kan*	This work	NY20L-*priC*	NY20L-frt x P1 KRC002
strain, strain background (*Escherichia coli*)	NY20ATL-*frt* Δ*priC*::*frt-kan*	This work	NY20ATL-*priC*	NY20ATL-frt x P1 KRC002

strain, strain background (*Escherichia coli*)	MIT125 Δ*rnhA*::*cat*	This work	MIT125c	MIT125 x P1 MIT162
strain, strain background (*Escherichia coli*)	KRC004 Δ*rnhA*::*cat*	This work	KRC004c	KRC004 x P1 MIT162
Recombinant DNA reagent	pING1 (Plasmid)	Katayama, *et al.*, 1994		Vector bearing arabinose-inducible promoter
Recombinant DNA reagent	pKA234 (Plasmid)	Kubota, *et al.,* 1997		pING1 encoding *dnaA*
Recombinant DNA reagent	pH136A (Plasmid)	Sakiyama, *et al.*, 2018		pKA234 *dnaA* H136A
Recombinant DNA reagent	pFH (Plasmid)	This study		pH136A mutated by Quick Change site-directed mutagenesis by primers F46A 1/2
Recombinant DNA reagent	pKW44-1 (Plasmid)	Ozaki, *et al.,* 2008		pKA234 *dnaA* V211A
Recombinant DNA reagent	pNA135 (Plasmid)	Ishida, *et al.*, 2004		pBR322 bearing *diaA* gene
Recombinant DNA reagent	pBS*oriC* (Plasmid)	Kawakami, *et al.* 2005		pBluescript bearing a 678 bp chromosome-derived region including *oriC*
Recombinant DNA reagent	pET22b(+)-priC (Plasmid)	Aramaki, *et al.*, 2015		pET22b(+) bearing *priC* under the T7 promoter

Recombinant DNA reagent	pTKM601 (Plasmid)	Keyamura, *et al.*, 2007		pBAD/HisB bearing *diaA*
Sequence-based reagent	F46A 1	Keyamura *et al.*, 2009	PCR primer	GTACGCGCCAAACCGC GCGGTCCTTCGATTG GGTACG
Sequence-based reagent	F46A 2	Keyamura *et al.*, 2009	PCR primer	CGTACCCAATCGAAGG ACCGCGCGGTTTGGCG CGTAC
Sequence-based reagent	ORI_1	Kasho *et al.*, 2014	qPCR primer	CTGTGAATGATCGGTG ATC
Sequence-based reagent	KW*oriC*Rev	Kasho *et al.*, 2014	qPCR primer	GTGGATAACTCTGTCA GGAAGCTTG
Sequence-based reagent	qoriK fw	Brochu *et al.*, 2014	qPCR primer	CGAGACTTCAGCGACA GTTAAG
Sequence-based reagent	qoriK rev	Brochu *et al.*, 2014	qPCR primer	CCTGCGGATATTTGCG ATACA
Sequence-based reagent	STM419	This work	qPCR primer	CGGACACCTTGTCTGA CCTAAG
Sequence-based reagent	STM420	This work	qPCR primer	AGTGTGAAAATGACCC TCTTGC

Figure 1

A *oriC* with AT-cluster

DUE
(DNA-Unwinding Element)

DOR
(DnaA-Oligomerization Region)

ssDNA binding motifs of DnaA

Left   Middle   Right

AT | L . M . R | R1 | IBR | τ1 R5M τ2 I1 I2 | R2 | C3 C2 I3 C1 R4 :DnaA box name

B   ATP-DnaA

Domain
I (1-86)   Binding to DiaA and DnaB by F46
II (87-134)   Flexible linker
III (135-374)   AAA+ motif
Arginine finger (R285)
H/B motifs (V211 and R245)
Low-affinity DnaB binding (H136)
IV (375-467)   DnaA box binding

H/B motifs
ATP
Arginine finger

C

IHF

D

DnaC
DnaB

E

Lagging strand
Leading strand

PriC

Rep

Fork restart

Figure 2

Figure 3

Host strain			*dnaA46*							
priC			+					−		
Number of cell [log10]	5	4	3	2	1	5	4	3	2	1

(n=4)

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 4-figure supplement 1

Number of cell [log10]   5   4   3   2   1

priC

30℃
+ pBR322
+ pNA135
− pBR322
− pNA135

37℃
+ pBR322
+ pNA135
− pBR322
− pNA135

42℃
+ pBR322
+ pNA135
− pBR322
− pNA135

(n=3)

Figure 7-figure supplement 1