1

Primosomal protein PriC rescues replication initiation stress by bypassing the DnaA-DnaB

interaction step for DnaB helicase loading at *oriC*

Ryusei Yoshida, Kazuma Korogi[1], Qinfei Wu[1], Shogo Ozaki*, and Tsutomu Katayama*

Department of Molecular Biology, Kyushu University Graduate School of Pharmaceutical Sciences,

Fukuoka, Japan

[1] These authors equally contributed to this study.

*Corresponding authors

Abstract

In *Escherichia coli*, replisome and replication fork assembly is initiated by DnaB helicase

loading at the chromosomal origin *oriC* via its interactions with the DnaA initiator and the DnaC

helicase loader. Upon replication fork arrest, the replisome including DnaB dissociates from the stalled

fork. Replication fork progression is rescued by primosomal protein PriA- or PriC-dependent pathway

in which PriA and PriC promote reloading of DnaB in different mechanisms. However, the mechanism

responsible for rescue of blocked replication initiation at *oriC* remains unclear. Here, we found that

PriC rescued blocked replication initiation in cells expressing an initiation-specific DnaC mutant, in

mutant cells defective in DnaA-DnaB interactions, and in cells containing truncated *oriC* sequence

variants. PriC rescued DnaB loading at *oriC* even in the absence of Rep helicase, a stimulator of the

PriC-dependent replication fork restart pathway. These and results of *in vitro* reconstituted assays

concordantly suggest that this initiation-specific rescue mechanism provides a bypass of the DnaA-

DnaB interaction for DnaB loading by PriC-promoted loading of DnaB to the unwound *oriC* region.

These findings expand understanding of mechanisms sustaining the robustness of replication initiation

and specific roles for PriC in the genome maintenance.

                                    Introduction

Chromosome replication is strictly regulated to ensure successful duplication of the genetic

material (Costa & Diffley, 2022; Grimwade & Leonard, 2021; Kasho et al., 2023). In *Escherichia coli*,

the ATP-bound form of the replication initiator protein DnaA (ATP-DnaA) forms a higher-order

complex with DNA-bending protein IHF at the unique chromosomal origin *oriC*. This complex

promotes local unwinding of *oriC* and recruits a pair of DnaB helicases, which are successively loaded

to the single-stranded (ss) DNA regions in a bidirectional manner with the aid of DnaC helicase loader.

ssDNA-loaded DnaB interacts with DnaG primase and DNA polymerase III holoenzyme, resulting in

assembly of the replisome (Arias-Palomo et al., 2019; Chodavarapu & Kaguni, 2016; Hayashi et al.,

2020; O'Donnell et al., 2013; Sakiyama et al., 2017, 2018, 2022; Wegrzyn & Konieczny, 2023).

Replisomes disassemble when the replication fork stalls in front of an obstacle, such as protein-DNA

roadblock or single strand breaks. PriA recognizes the abandoned fork structure, which triggers

reloading of DnaB with its partner proteins, such as PriB, PriC, DnaT, and DnaC (Heller & Marians,

2005a; Masai et al., 1994; Michel & Sandler, 2017; Windgassen et al., 2018). In addition, independently of

PriA, PriC binds to the abandoned fork structure, triggering reloading of DnaB with the aid of DnaC,

which is supported by Rep helicase depending on the fork structure (Heller & Marians, 2005b; Sandler

et al., 1999). Also, homologous recombination mechanisms can rescue the abandoned replication fork

during SOS responses (Asai & Kogomal, 1994; Kogoma, 1997; Né Dicte Michel et al., 2001). In

contrast to these well characterized mechanisms at abandoned forks, little is known about what happens

when replication initiation is impeded at *oriC*.

E. coli oriC comprises a Duplex-Unwinding Element (DUE) and a DnaA-Oligomerization

Region (DOR) (Figure 1A) (Grimwade & Leonard, 2021; Kasho et al., 2023; Wegrzyn & Konieczny,

2023). The DUE contains 13-mer AT-rich sequence repeats known as L-, M-, and R-DUEs. The M- and

R-DUEs are essential for stable DUE unwinding with the specific sequences (TT[A/G]T(T)) used for

DnaA-ssDNA binding (Ozaki & Katayama, 2012; Sakiyama et al., 2017). The L-DUE containing the

TTATT sequence promotes efficient DnaB loading by expanding the unwound *oriC* region (Sakiyama et

al., 2022). The DOR is divided into three subregions: the Left-, Middle-, and Right-DORs, which contain

asymmetric 9-mer DnaA binding sequences (DnaA box) with the consensus sequence TTA[T/A]NCACA

(Figure 1A) (Noguchi et al., 2015; Ozaki & Katayama, 2012; Rozgaja et al., 2011; Sakiyama et al., 2017;

Shimizu et al., 2016). The Left-DOR contains a cluster of unidirectionally aligned DnaA boxes including

high-affinity DnaA box R1, and low-affinity boxes R5M, τ1−2, and I1-2, with an IHF-binding region

between R1 and R5M boxes. The Right-DOR contains an oppositely oriented DnaA box cluster including

high-affinity DnaA box R4 and low-affinity boxes I3 and C1-3. These DnaA box clusters form

frameworks for the Left- and Right-DnaA subcomplexes, respectively. DnaA bound to the R2 box, which

solely resides in the Middle-DOR, stabilizes these DnaA subcomplexes (Miller et al., 2009; Rozgaja et al., 2011; Shimizu et al., 2016). In addition, the AT-cluster (TATTAAAAAGAA) region, which connects to the L-DUE, stimulates DnaB loading in the absence of the Right-DnaA subcomplex (Sakiyama et al., 2022).

DnaA comprises four functional domains (Figure 1B). Domain I binds to multiple proteins such as DnaB helicases and the DnaA-assembly stimulator DiaA (Abe et al., 2007; Hayashi et al., 2020; Keyamura et al., 2007, 2009). Domain II is a flexible linker (Abe et al., 2007; Nozaki & Ogawa, 2008). Domain III contains AAA+ (ATPases Associated with various cellular Activities) motifs, which are involved in tight ATP/ADP binding, ATP hydrolysis, and domain III-III interactions (Duderstadt et al., 2011; Erzberger et al., 2006; Felczak & Kaguni, 2004; Kawakami et al., 2005; Ozaki et al., 2012). The arginine finger motif (Arg285) in domain III interacts with ATP bound to domain III of adjacent DnaA protomers, stimulating DnaA complex formation in a cooperative manner (Kawakami et al., 2005; Noguchi et al., 2015). In addition, H/B motifs (Val 211 and Arg245) in this domain bind to ssDNA in a sequence-specific manner (Ozaki et al., 2008; Sakiyama et al., 2017). Domain IV, which comprises the helix-turn-helix motif, recognizes the DnaA box (Fujikawa et al., 2003).

For replication initiation, ATP-DnaA molecules cooperatively oligomerize at *oriC* to form the Left- and Right-DnaA subcomplexes (Figure 1C). Formation of these complexes is stimulated by homotetrameric DiaA protein, which binds up to four DnaA molecules via domain I, including Phe46 (Keyamura et al., 2007, 2009). In concert with DNA superhelicity and thermal energy, the DUE undergoes initial unstable unwinding, which is stabilized through specific interaction between H/B motifs of domain III in the Left-DnaA subcomplex and the upper strand of M- and R-DUEs. This interaction is facilitated by sharp DNA bending by IHF, resulting in stable DUE unwinding (Figure 1C) (Ozaki et al., 2008; Sakiyama et al., 2017). When the single-stranded region is expanded to the L-DUE, the resultant single-stranded L-DUE binds the H/B motifs within the Right-DnaA subcomplex, maximizing the efficiency of the DnaB loading process (Sakiyama et al., 2022).

Subsequently, two DnaB helicases are loaded onto each ssDUE strand through interactions with DnaC and DnaA (Figure 1D). DnaC binding to DnaB changes the closed ring structure of the DnaB hexamer to an open spiral form to allow it to encircle ssDNA (Arias-Palomo et al., 2019; Nagata et al., 2020). Each Left- and Right-DnaA subcomplex binds a DnaB-DnaC complex via high-affinity interaction between DnaA Phe46 and DnaB Leu160 and guides the DnaB-DnaC complexes to unwound ssDUE strands via a low-affinity interaction between DnaA domain III His136 and DnaB (Figure 1B and 1D). Upon interaction with ssDNA, DnaC dissociates from DnaB, which enables migration of DnaB hexamer with helicase action. The pair of DnaB helicases loaded onto the ssDNA strands progress in opposite directions, forming replisomes with DnaG and Pol III holoenzyme (Chodavarapu & Kaguni, 2016;

O'Donnell et al., 2013).

Replication restart pathways ensure fork progression of the entire chromosome under conditions

that trigger replisome disassembly (Heller & Marians, 2005a; Lopper et al., 2007; Michel & Sandler,

2017). PriA helicase-dependent pathways predominantly facilitate DnaB reloading onto abandoned forks

in vivo (Flores et al., 2002). PriA is a 3'-5' DNA helicase and has specific affinity for forked DNA

structures comprising one parental dsDNA and two newly synthesized sister dsDNA strands. This permits

PriA recognition of the abandoned replication fork and the subsequent unwinding of the nascent lagging

strand (Duckworth et al., 2023; Windgassen et al., 2018; Windgassen & Keck, 2016). The resultant

unwound ssDNA associates with PriB and DnaT to promote reloading of DnaB helicase with the aid of

DnaC (Duckworth et al., 2023; Heller & Marians, 2005a; Lopper et al., 2007). In the absence of PriB,

PriC participates in a PriA-dependent pathway (Sandler et al., 1999). In addition, PriA is required for

oriC/DnaA-independent chromosomal replication called stable DNA replication (SDR) (Masai et al.,

1994), which is promoted by UV irradiation (inducible iSDR) and loss of *rnhA* or *recG* (constitutive

cSDR) (Kogoma, 1997). In both mechanisms, replication is initiated at triple-stranded structures

comprising a dsDNA-ssDNA hybrid (D-loop) or a dsDNA-ssRNA hybrid (R-loop) generated at specific

chromosomal loci. These structures mimic the abandoned fork structure, thereby allowing helicase

loading to occur similarly to that during PriA-dependent fork restart (Masai et al., 1994).

Independently of PriA, PriC can restart replication from abandoned forks through its interaction

with the ssDNA region and SSB (Single-Stranded DNA Binding protein) (Figure 1E) (Heller & Marians,

2005a; Wessel et al., 2013, 2016). PriC consists of N-terminal and C-terminal domains, which are

composed of α helices and are connected by a short linker. The PriC C-terminal domain remodels the

SSB-ssDNA complex, which is a prerequisite for the recruitment of DnaB helicase (Heller & Marians,

2005a; Wessel et al., 2013, 2016). PriC also interacts with Rep helicase. When the length of the ssDNA

gap within the abandoned fork is short, Rep helps expand the ssDNA gap, promoting PriC-mediated

remodeling of the SSB-ssDNA complex (Heller & Marians, 2005a, 2005b; Nguyen et al., 2021). Notably,

in addition to these functions at the abandoned forks, PriC is suggested to play a role in DnaA-*oriC*-

dependent replication initiation under challenging conditions, as evidenced by the synthetic lethality

between *priC303*::*kan* and a subgroup of temperature-sensitive *dnaA* mutant alleles (*dnaA46* and

dnaA508): i.e. *priC303*::*kan* mutants bearing *dnaA46* or *dnaA508*, but not wild-type (WT) *dnaA*, can not

grow even at 30°C (Hinds & Sandler, 2004). However, the mechanisms via which PriC rescues blocked

replication initiation remain elusive.

In this study, we analyzed the PriC-dependent promotion of replication initiation from *oriC* in

the context of various replication initiation stresses. We found that PriC stimulated replication initiation

in *dnaC2* cells, in which DnaB helicase loading at *oriC* is defective, indicating that it has a role in the

rescue of blocked replication initiation. Moreover, PriC stimulated the growth of cells defective in

DnaA-DnaB interactions in addition to the growth of cells with *oriC* sequence deletions that inhibit

DnaB loading. Consistent with these results, in an *in vitro* reconstituted system, PriC stimulated DnaB

loading when the DnaA-DnaB interaction was inhibited at *oriC*. Furthermore, we found that PriC did

not stimulate initiation of cSDR, demonstrating that PriC functions specifically in the DnaA-*oriC*

system. Taken together, we suggest that PriC rescues blocks in replication initiation by bypassing DnaB

loading at *oriC*-DnaA complexes.

                                                    Results

    PriC stimulates DNA replication initiation in *dnaA46* and *dnaC2* cells

The temperature-sensitive *dnaA46* and *dnaA508* mutants exhibit synthetic lethality with

priC303::*kan* even at 30°C (Hinds & Sandler, 2004), suggesting that PriC can rescue blocks in DnaA-

dependent replication initiation. However, the mechanism underlying PriC-promoted rescue of blocks in

replication initiation remains a mystery. Based on previous reports demonstrating PriC binding to short

oligo-ssDNA, SSB, and DnaB and its role in DnaB loading at stalled replication forks (Wessel et al., 2013,

2016), we considered the possibility that PriC permits bypass of DnaA-dependent stable DUE unwinding

and/or DnaB loading at the *oriC* ssDUE. To genetically test these ideas, we analyzed the impact of PriC

on the growth of a temperature-sensitive *dnaC2* mutant, which is competent for ongoing replisome

progression at 37–42°C, but defective in DnaB loading specifically at the *oriC* ssDUE (Withers &

Bernander, 1998).

In spotting assays performed using serial dilutions of the cell cultures and LB agar plates, the

cell growth of *dnaA46* cells was slightly inhibited at 37°C (Figure 2A). Consistent with the previous

report, when crossed with the Δ*priC* allele, the cell growth of *dnaA46* cells but not WT *dnaA* cells was

severely inhibited at 37°C (Figure 2A). Unlike the *dnaA46 priC303*::*kan* cells, the *dnaA46* Δ*priC* cells

grew at 30°C without prominent inhibition (Figure 2A). This difference may stem from the different

strain backgrounds. Notably, the cell growth of *dnaC2* mutant at 35°C was moderately inhibited,

whereas it was severely inhibited in *dnaC2* Δ*priC* double mutant (Figure 2A). Also, the cell growth of

dnaC2 cells and *dnaC2* Δ*priC* cells was similar at 30°C. The observed requirement of PriC for cell

growth of *dnaC2* cells at 35°C supports the idea that PriC assists in the DnaC-dependent DnaB loading

step at *oriC*.

Introduction of a *priC*-encoded plasmid (pBR*priC*), but not an empty vector (pBR322),

restored the cell growth of *dnaA46* Δ*priC* double mutant at 37°C (Figure 2-figure supplement 1AB) and

that of *dnaC2* Δ*priC* double mutant at 35°C. Also, introduction of pBR*priC* to *dnaA46* and *dnaC2*

mutants with intact *priC* (*priC*⁺) might only slightly enhance the cell growth at these temperatures

(Figure 2- figure supplement 1AB), suggesting that the intrinsic level of PriC is functionally nearly

sufficient.

To investigate replication initiation from *oriC*, we used flow cytometry of synchronized cells

bearing *dnaA46* or *dnaC2* with or without Δ*priC*. In these experiments, cells were incubated at 42°C or

37°C for 80–90 min to synchronize the replication cycle at the pre-initiation step, after which

incubation was continued at 30°C for 5 min to allow initiation and then 4 h in the presence of

rifampicin and cephalexin to inhibit both replication initiation and cell division and allow replication

run-out of the whole chromosome. The resultant number of chromosomes per cell, measured by flow

cytometry, is known to correspond to the number of *oriC* copies per cell at the time of drug addition, an

indicator of initiation activity.

A majority of the synchronized *dnaA46* and *dnaC2* cells had only a single *oriC* copy per cell

after incubation at 42°C or 37°C regardless of the presence of Δ*priC* (Figure 2B and 2C). In cells with

intact *priC* (*priC*$^+$), after release of initiation at 30°C for 5 min, most cells had two *oriC* copies.

However, in *dnaA46* Δ*priC* cells, only about half of the cells had two *oriC* copies after the release

(Figure 2B), indicating moderate inhibition of initiation. In *dnaC2* Δ*priC* cells, the number of two-*oriC*

cells was slightly lower than that of *dnaC2* cells (Figure 2C). These observations are consistent with the

cell growth ability (Figure 2A) and with our hypothesis that PriC contributes to the rescue of blocked

replication initiation by assisting DnaB loading onto the *oriC* ssDNA.

We further performed flow cytometry analysis to assess the role for PriC in replication

initiation of the WT *dnaA/dnaC* cells. To estimate the number of *oriC* copies per unit cell mass

(ori/mass) as a proxy for initiation activity (Sakiyama et al., 2017, 2022), exponentially growing cells

were subjected to run-out replication in the presence of rifampicin and cefalexin (Figure2-figure

supplement 2). When cells were grown in LB medium at 25˚C, 30˚C or 42˚C, the ori/mass values for

Δ*priC* mutant were comparable to those of *priC*$^+$ cells. Similar results were observed in cells grown in

M9 minimal medium at 30˚C. However, when grown in M9 minimal medium at 37˚C, Δ*priC* mutant

cells exhibited slightly reduced ori/mass values. Also, it is noteworthy that in all conditions we tested,

the fraction of cells with non-2^n *oriC* copies was slightly higher in Δ*priC* cells compared to *priC*$^+$ cells.

The asynchronous initiations would be caused by partial inhibition of initiation of multiple origins in

Δ*priC* cells, as representatively seen for the case of Δ*priC* cells growing in M9 minimal medium at

37°C where four-*oriC* cells reduced and three-*oriC* cells increased, compared to *priC*$^+$ cells. A similar

case might be shown for Δ*priC* cells growing in LB medium at 25°C in that eight-*oriC* cells reduced

and asynchronous initiations (A.I. %) as well as four-*oriC* cells increased. As such, inhibition of

replication initiation would occur intrinsically at low frequency in the WT *dnaA*/*dnaC* background. PriC

function would be effective to suppress the inhibition, thereby ensuring replication initiation of multiple

origins.

PriC rescues the cell growth of DnaA mutants defective in DnaB interactions

DnaA contains two regions for interactions with DnaB during DnaB loading at *oriC* ssDNA:

one is a high-affinity region in domain I containing Phe46, and the other is a low-affinity region in

domain III containing His136. To analyze the mechanism of PriC-promoted DnaB loading at *oriC*, we

introduced a series of pING1 vector-based plasmids encoding WT DnaA (pKA234) or the DnaA F46A

H136A double mutant (pFH) into *dnaA46* cells with or without *priC*. Growth of *dnaA46* cells

containing pING1 was inhibited at 42°C, but the leaky expression of *dnaA* from pKA234 enabled the growth of these cells, as previously reported (Sakiyama et al., 2018) (Figure 3). Introduction of pFH moderately supported the cell growth of *dnaA46* cells at 42°C. By contrast, *dnaA46 ΔpriC* cells carrying pFH exhibited little growth at 42°C (Figure 3).

Western blotting analysis supported that the expression levels of WT DnaA and DnaA F46A H136A proteins in *dnaA46* cells bearing pKA234 or pFH were comparable, regardless of the presence or absence of *priC⁺* (Figure 3-figure supplement 1). As DnaA46 is unstable and thermolabile (Shimuta et al., 2004), these cells were grown at 30°C, and were incubated at 42°C for 90 min before Western blotting analysis. The DnaA level in *dnaA46* cells bearing the vector pING1 plasmid was minimum.

These results suggest that PriC is crucial for the growth of cells defective in specific DnaA-DnaB interactions, possibly because it can bypass the DnaA-DnaB interaction step required for DnaB loading onto *oriC* ssDNA.

PriC rescues DNA replication inhibition by DiaA overexpression *in vivo*

DiaA, a homotetrameric protein, binds to multiple DnaA molecules and stimulates DnaA multimer assembly at *oriC* (Keyamura et al., 2007). The Phe46 residue in DnaA domain I is part of the DiaA-binding site and DnaB binding site. Oversupply of DiaA inhibits timely initiation of replication in cells growing in tryptone medium at 30°C, probably because of binding competition between DiaA and DnaB for DnaA domain I (Ishida et al., 2004; Keyamura et al., 2009). If PriC can bypass the DnaA-DnaB interaction requirement for DnaB loading onto *oriC*, PriC should suppress inhibition of initiation by DiaA oversupply. To test this possibility, we introduced pBR322 or its derivative encoding *diaA* (pNA135) into cells with or without *priC* (Figure 4). *priC⁺* cells carrying pBR322 or pNA135 similarly grew on LB agar plates containing ampicillin between 25°C and 42°C (Figure 4A and Figure 4-figure supplement 1). However, cell growth of *ΔpriC* cells carrying pNA135, but not pBR322, was moderately inhibited at 25°C (Figure 4A), supporting our conjecture that PriC can bypass the requirement for the DnaA-DnaB interaction.

Rep helicase stimulates PriC-mediated rescue of arrested replication forks containing nascent strands (Heller & Marians, 2005b, 2005a; Sandler, 2000). To test the contribution of Rep helicase to PriC-promoted rescue of blocked replication initiation, we introduced pBR322 or pNA135 into *Δrep* cells. Unlike *ΔpriC* cells, *Δrep* cells carrying pNA135 grew similarly to *Δrep* cells carrying pBR322 (Figure 4A), suggesting no involvement of Rep in the PriC-promoted rescue of blocked replication initiation. This is consistent with the features of unwound *oriC*, which has a fork structure similar to that of the replication fork but without nascent strands, and the specific role of Rep in rescuing abandoned replication forks with nascent strands in the PriC-dependent pathway. Rep is specifically

required for the unwinding of the nascent leading strand of abandoned replication forks, which is

needed to expand the ssDNA region required for entry of PriC (Heller & Marians, 2005b, 2005a).

We further analyzed replication initiation in Δ*priC* cells carrying pNA135 at 30°C by flow

cytometry (Figure 4B). Under these conditions, *priC*[+] cells bearing pBR322 mainly had eight or four

oriC copies per cell with eight-*oriC* cells predominating (Figure 4B). The number of *oriC* copies per

cell was similar to that in *priC*[+] cells bearing pNA135, with only slight increase of asynchronous

initiation, as shown by the slight increase in the number of cells with non-2^n *oriC* copies, which is

basically consistent with our previous data (Ishida et al., 2004). However, in Δ*priC* cells bearing

pNA135, severe inhibition of initiation occurred, as shown by the substantial decrease in the number of

eight-*oriC* cells and the large increase of the number of cells with only one to three *oriC* copies (Figure

4B). This result supports the proposed function of PriC at *oriC* mentioned above.

To test whether PriC rescues replication initiation inhibited by deletion of the *diaA* gene, we

conducted similar flow cytometry analysis of Δ*priC*, Δ*diaA*, and Δ*priC* Δ*diaA* cells (Figure 4C). Δ*diaA*

cells showed severe asynchrony of replication initiation, as shown by the increase in the number of five-

to-seven-*oriC* cells, which supports the stimulatory role of DiaA in DnaA-*oriC* complex formation

(Figure 4C). However, Δ*priC* Δ*diaA* cells did not show more asynchronization of replication initiation

than Δ*diaA* cells, as shown by the similar asynchrony index (A.I.). These results suggest that PriC does

not support or bypass DnaA-*oriC* complex formation.

PriC stimulates replication initiation in cells with *oriC* sequence truncations

Right DnaA-subcomplex plays a stimulatory role in DnaB helicase loading (Sakiyama et al.,

2022; Stepankiw et al., 2009). The loading of two DnaB hexamers at *oriC* ssDNA regions requires

multiple steps that depend on the distinct functions of two DnaA subcomplexes (Figure 1): the Left-

DnaA subcomplex stably unwinds M-R regions of the DUE and the Right-DnaA subcomplex expands

the unwound region to the AT-L region, supporting efficient DnaB helicase loading (Sakiyama et al.,

2017, 2022; Yoshida et al., 2023). The expanded ssDNA regions in this open *oriC* complex efficiently

promote the loading of one DnaB hexamer onto the lower (A-rich) strand M-R region and a second

DnaB hexamer onto the upper (T-rich) strand of the DUE (Fang et al., 1999; Sakiyama et al., 2022). In

the absence of the Right-DnaA subcomplex, the AT region of the AT-L region assists in DnaB loading

(Sakiyama et al., 2022)

To determine whether PriC rescues defects in DnaB loading when the *oriC* sequence is altered

to prevent Right-DnaA subcomplex formation, we first assessed the cell growth of cells lacking the

Middle- and Right-DOR of *oriC* (Left-*oriC*) (Figure 5A). The Left-*oriC* mutant formed colonies at

37°C but showed severe growth defects at 25°C and 30°C (Figure 5B). At 37°C, thermal energy might

be sufficient to expand DUE unwinding into the AT region and allow DnaB loading in the absence of

the Right-DnaA subcomplex (Sakiyama et al., 2022), but not at 25°C or 30°C. By contrast, the cell

growth of the Left-*oriC* Δ*priC* double mutant was markedly compromised at 37˚C and moderately

reduced at 25°C and 30°C (Figure 5B). These findings support the idea that PriC functions to rescue

defective processes in DnaB loading caused by deletions of the *oriC* sequence that prevent DnaA

subcomplex formation.

Next, similar experiments were performed using a mutant in which the R4 DnaA box within

Right-DOR was substituted with a sequence (R4*Tma*) defective in binding of *E. coli* DnaA (Figure 5A

and 5C) (Noguchi et al., 2015; Sakiyama et al., 2022). We reasoned that formation of the Right-DnaA

subcomplex would be inhibited by the introduction of R4*Tma* and PriC would be required for robust

initiation in this context. As the Left-*oriC* strain, the *priC*[+] R4*Tma* strain showed normal cell growth.

By contrast, cell growth of the R4Tma Δ*priC* double mutant was impaired moderately at 25°C and

slightly at 30°C (Figure 5C), indicating that PriC can rescue the blocked replication initiation caused by

the absence of the R4 box. The reduced cell growth of R4*Tma* Δ*priC* strain was alleviated at higher

temperatures, similar to the phenotype of the Left-*oriC* mutant.

Furthermore, we assessed replication initiation in R4*Tma* cells at 30°C using flow cytometry

(Figure 5D). R4*Tma* cells grew slightly slower than WT *oriC* cells and showed clear inhibition of

initiation; in contrast to the WT *oriC* strain, the R4*Tma* mutant cell population contained more four-

oriC cells and fewer eight-*oriC* cells, and showed severe asynchronous initiation. Notably, these

negative effects of the R4*Tma* mutation were amplified by deletion of *priC*, i.e., the number of one to

three-*oriC* cells increased and the ori/mass ratio was further reduced.

We next analyzed the role of PriC in cells lacking the AT-L region of *oriC* (subATL *oriC*)

(Figure 6A). L-DUE and the flanking AT region assist in DnaB helicase loading by stimulating *oriC*

unwinding, which is essential in a strain defective in Right-DnaA subcomplex formation (Sakiyama et

al., 2022). SubATL *oriC* cells and WT *oriC* cells grew similarly on LB agar plates between 25°C and

37°C, regardless of the presence of Δ*priC* (Figure 6B). However, flow cytometry analysis revealed that

deletion of *priC* specifically inhibited initiation in subATL *oriC* cells (Figure 6C); the number of eight-

oriC cells was lower while the number of four-to-seven *oriC* cells was higher in subATL *oriC* Δ*priC*

cells than in *priC*[+] subATL *oriC* cells. Taken together, these results are consistent with the idea that PriC

restores initiation and DnaB loading at such truncated *oriC* sequences.

PriC loads DnaB at *oriC* unwound by the DnaA complex in an *in vitro* reconstituted system

To analyze the mechanism of PriC rescue of blocked initiation at *oriC*, we assessed PriC activity

in an *in vitro* reconstituted system for DnaB loading and DnaB-promoted DNA unwinding using the

supercoiled circular form (form I) of the *oriC* plasmid pBS*oriC* and purified proteins, namely DnaA, N-terminally histidine-tagged DnaB (His-DnaB), DnaC, IHF, SSB, and gyrase. In this system, DnaA and IHF unwind the DUE and stimulate His-DnaB loading to the ssDUE region with the aid of DnaC. The loaded His-DnaB expands the ssDNA regions in concert with the ssDNA-binding activities of SSB. Concomitantly, the positive supercoiling generated through DNA unwinding by His-DnaB is resolved by DNA gyrase, resulting in the formation of plasmid DNA topoisomers (form I*). Form I* can be separated from form I by agarose gel electrophoresis (Baker et al., 1986; Sakiyama et al., 2022).

First, we assessed the effect of PriC on WT DnaA. WT DnaA stimulated form I* formation in the absence of PriC (Figure 7A, lanes 1 and 5) and increasing concentrations of PriC moderately inhibited WT DnaA-mediated form I* formation (Figure 7A, lanes 5 to 8), probably because of competition between DnaA and PriC for binding to DnaB. Next, we focused on the DnaA F46A H136A double mutant. The DnaA F46A and DnaA H136A mutant proteins alone are reported not to support form I* formation (Keyamura et al., 2009; Sakiyama et al., 2018). Consistently, the DnaA F46A H136A double mutant protein was inactive in form I* formation (Figure 7A, lane 9). However, when PriC was added to the assay, DnaA F46A H136A promoted form I* formation to a level comparable to that of WT DnaA in the presence of PriC (Figure 7A, lanes 5-12 and 7B), indicating that PriC rescues defective DnaB loading by DnaA F46A H136A-*oriC* complexes. These findings are consistent with the idea that PriC can bypass the strict reliance on DnaA-DnaB interactions for DnaB loading at *oriC*.

To corroborate this idea, we also determined whether PriC rescues blocked initiation resulting from unstable DUE unwinding. *oriC* complexes containing DnaA V211A cannot bind to the ssDUE and consequently fails in stable DUE unwinding (Ozaki et al., 2008). The DnaA V211A mutant was virtually inactive for form I * formation irrespective of the presence or absence of PriC (Figure 7A, lanes 13-16, and 7B), indicating that PriC cannot efficiently rescue a defect of DUE unwinding.

We further evaluated the DnaB loading activity of PriC in the presence of DiaA using a similar *in vitro* reconstituted system. Previously, DiaA was shown to stimulate *oriC* DUE unwinding, but to inhibit form I* formation, because DiaA-DnaA binding competitively inhibits DnaB-DnaA binding; DiaA and DnaB share the same binding site in DnaA domain I (Figures 7C lanes 1, 3, 5, and 7 and 7D) (Keyamura et al., 2009). PriC moderately stimulated form I* formation specifically in the presence of both DnaA and DiaA (Figures 7C lanes 7 and 8, and 7D). These results further support the idea that PriC can bypass the specific requirement for the DnaA-DnaB interaction in DnaB loading at *oriC*. DiaA bound to *oriC*-DnaA complexes might be a physical obstacle reducing efficiency of DnaB loading by PriC.

In addition, we analyzed the DUE unwinding activities of DnaA mutants using the *in vitro* reconstituted system. As previously reported, DnaA V211A is largely inactive in DUE unwinding, but DnaA F46A H136A unwinds the DUE at a level comparable to that of WT DnaA (Figure 7-figure

supplement 2A and 2B). PriC did not stimulate DUE unwinding irrespective of the presence of WT

DnaA (Figure 7-figure supplement 2C and 2D). These results demonstrate the functional specificity of

PriC in rescuing failed DnaB loading at *oriC* ssDNA and are consistent with the PriC mechanism

mentioned above.

Role for PriC in cSDR

Finally, to analyze the role of PriC in other types of replication initiation, we examine whether

PriC contributes to the initiation of cSDR, which does not require *dnaA* or *oriC* and is activated in cells

lacking *rnhA* or *recG*. Although the PriA-PriB-DnaT primosome complex has previously been reported

to be involved in the initiation of cSDR (Heller & Marians, 2006), it is unclear whether PriC has any

role in this type of initiation.

Unlike *dnaA46* cells, *dnaA46 rnhA*::*cat* double mutant cells grew even at 40°C (Figure 8A),

indicating that these cells were engaged in cSDR, as previously reported (Hinds & Sandler, 2004). The

cell growth of Δ*priC dnaA46 rnhA*::*cat* triple mutant was severely inhibited at 40°C (Figure 8A),

suggesting that PriC contributes to the growth of *dnaA46 rnhA*::*cat* mutant cells.

To determine whether PriC contributes to the initiation of *dnaA46-oriC* replication or cSDR,

we calculated the relative copy numbers of *oriC* (84.6 min) and *ter* (32.4 min: corresponding to *insQ*)

using real-time quantitative PCR (qPCR). The *ypaB* (50.5 min) located at the middle position between

oriC and *ter* was used as a reference. To calculate the copy number of the chromosomal *ter* region, the

preferential initiation site of cSDR (Brochu et al., 2018; Maduike et al., 2014), *dnaA46 rnhA*::*cat* cells

growing exponentially at 30°C were shifted to 40°C for 90 min and the copy number before and after

incubation at 40°C was calculated. Consistent with the cell growth data (Figure 8A), the *ter* copy

number ratio of the *dnaA46 rnhA*::*cat* double mutant increased after 40°C incubation, whereas the copy

number ratio of the *dnaA46* mutant expressing WT *rnhA* remained the same (Figure 8B). The *ter* copy

number ratio of the *dnaA46 rnhA*::*cat* Δ*priC* triple mutant strain was also increased, indicating that

cSDR could occur even in the absence of PriC.

To monitor initiation of DnaA-*oriC* replication, we determined the *oriC* copy number ratio

(Figure 8B). At the permissive temperature, despite having similar growth rates, the *oriC* copy number

ratio of the *dnaA46* Δ*priC* double mutant cells was lower than that of the *dnaA46* single mutant,

confirming the importance of PriC for replication initiation in *dnaA46* cells, as shown in Figure 2. In the

priC[+] *dnaA46 rnhA*::*cat* mutant cells, the *oriC* copy number ratio was reduced, which is explained by

activation of cSDR and increase of the relative copy number of the reference *ypaB* locus (50.5 min

chromosomal map position). Together, these results suggest the idea that PriC stimulates replication

fork progression during cSDR.

To consolidate these findings, we performed the chromosome loci copy-number analysis using

whole-genome sequencing. The *dnaA46* cells and its derivative cells were grown exponentially at 30°C,

followed by incubation at 40°C for 90 min prior to DNA extraction. In growing WT cells, the highest

copy numbers are observed near the *oriC*-proximal region and the lowest are near the *ter*-proximal

region, resulting in a coverage bias from *oriC* to *ter* (Brochu et al., 2018; Maduike et al., 2014).

However, in *dnaA46* cells incubated at 40˚C, the coverage bias was largely diminished due to the run-

out replication (Figure 8-figure supplement 1A). Notably, *dnaA46 rnhA*::*cat* cells incubated at 40˚C

exhibited discrete coverage peaks at the preferential initiation sites of cSDR near the *ter* (or *insQ*)

region with a decreasing coverage bias to *oriC*. Similar profiles were observed in the Δ*priC* mutant

background, supporting the idea that cSDR can occur even in the absence of PriC (Figure 8-figure

supplement 1A). Moreover, in the *dnaA46 rnhA*::*cat* background, the absence of *priC* appeared to

decrease the coverage level at genomic positions near the 1.3 MB and 4 MB map-positions, which are

slightly outside of *ter* and *oriC*, respectively (Figure 8-fugure supplement 1B). This observation may

reflect the role for *priC* in the replication fork progression.

Discussion

In growing cells, replication initiation can be blocked under challenging conditions. However, the mechanisms that rescue blocked initiations are largely unknown. To elucidate such mechanisms, we investigated PriC-promoted replication initiation under various challenging conditions. We showed that PriC was necessary for the optimal growth of *dnaC2* cells at the semi-restrictive temperature and for replication initiation even at the permissive temperature (Figure 2), suggesting that PriC contributes to replication initiation rescue in *dnaC2* cells by facilitating DnaB helicase loading at *oriC*. Furthermore, we observed that PriC was necessary for the optimal growth of cells in which the DnaA-DnaB interaction was inhibited by DiaA overexpression or DnaA F46A H136A double mutations (Figures 3 and 4). In addition, PriC was stimulatory for the cell growth and replication initiation bearing *oriC* mutations that impair efficient DnaB loading (Figures 5 and 6). These results, together with results of form I* formation assays (Figures 7), suggest that PriC bypasses the need for the DnaA-DnaB interaction by binding to unwound *oriC* DNA and recruiting DnaB to facilitate its loading onto ssDNA (Figure 9). This mechanism is similar to that proposed for the rescue of abandoned replication forks by PriC (Figure 1E).

Also, PriC contributes to replication initiation rescue in *dnaA46* cells of 37°C. It is presumable that DnaA46 protein becomes partially denatured at the sub-permissive temperature of 37°C. This partial denaturation should impair both origin unwinding and helicase loading, though not to the extent that cell viability is lost. The *priC* deletion should further exacerbate helicase loading defects by inhibiting the bypass mechanism (Figure 9), resulting in the lethality of *dnaA46* cells at this temperature.

There might remain an alternative possibility that PriC could stabilize the DnaA-DnaB interaction for recusing the impaired DnaB loading process. However, we believe that this possibility is not very likely. Given that interactions between DnaA and DnaB during DnaB loading to *oriC* are highly dynamic and involve regulated multiple steps, the stabilization of the DnaA-DnaB interaction by PriC, even if it occurs, carries a considerable risk of inhibiting the DnaB loading by constructing abortive complexes. In addition, DnaA-DiaA binding is very tight and stable (Keyamura et al.,2007, 2009). Even if WT DnaA and WT DnaB are present, PriC can rescue the initiation defects of *oriC* mutants. Based on these facts and the known characteristics of PriC, it is more reasonable to infer that PriC provides a bypass of DnaB loading at *oriC*, similar to the mechanism at the stalled replication fork (Figures 1E and 9).

The finding that PriC failed to stimulate form I* formation in the presence of DnaA V211A, which lacks DUE unwinding activity (Figures 7), provides mechanistic insight into PriC-dependent replication initiation at *oriC*. We propose that PriC-dependent replication initiation possibly begins with

the recognition of a stably unwound ssDNA region by PriC (Figure 9). Consistent with this perspective,

PriC ssDNA-binding activity and ssDNA regions have been reported to be important for PriC-promoted

replication fork rescue (Heller & Marians, 2006; McMillan & Keck, 2024). The failure of PriC to

stimulate replication initiation in Δ*diaA* cells (Figure 4C) is also potentially explained by the reduced

DUE unwinding caused by destabilization of the DnaA-*oriC* complex in Δ*diaA* cells. Based on this, we

suggest that the stably unwound ssDNA region in *oriC* serves as a distinctive feature for PriC-

promoting DnaB loading at *oriC* for replication initiation rescue, and distinguishes it from PriA-

dependent replication restart, which requires the specific structure of the replication fork DNA

consisting of sister dsDNA strands (Heller & Marians, 2005a; Windgassen et al., 2018). Unlike PriA,

PriC can interact with the simple ssDNA to reload DnaB (Figure 1E).

In this study, we found that cells with DnaA F46A H136A double mutations grew well, but

only when the cells expressed PriC. As for the DnaA F46A single mutant, our previous studies show that

DnaA F46A has a limited residual activity *in vivo* (unlike *in vitro*), and allows slow growth of cells (Abe et

al., 2007). The growth of cells expressing the DnaA H136A single mutant was severely inhibited even

when the cells expressed PriC (Sakiyama et al., 2018). The His136 residue is located within the weak,

secondary DnaB interaction region in DnaA, and is crucial for DnaB loading onto *oriC* ssDNA.

Although domain I in DnaA H136A can stably tether DnaB-DnaC complexes to DnaA complexes on

oriC (Sakiyama et al., 2018), the complexes fail to load DnaB onto *oriC* ssDNA even in the presence of

PriC. It is possible that the interaction between PriC and DnaB is inhibited by stable DnaB binding to

DnaA domain I. This idea is consistent with the *in vivo* feature of DnaA F46A single mutant.

Conversely, the PriC-DnaB interaction may inhibit the interaction between DnaA domain I and DnaB,

as suggested by the inhibitory effect of PriC on DnaB loading *in vitro* in the presence of WT DnaA

(Figure 7). However, this inhibitory effect of PriC was not observed *in vivo*, suggesting that PriC-DnaB

binding could involve still unknown factors and regulatory mechanisms.

PriC supported cSDR-dependent growth in *dnaA46 rnhA*::*cat* double mutant cells without

stimulating the initiation of cSDR (Figure 8). In this mutant, cSDR predominantly initiates from the *ter*

region, resulting in head-on collisions between the replisome and transcription complexes in *rrn*

operons (Maduike et al., 2014). This suggests that in cSDR cells PriC specifically rescues abandoned

forks including those generated during head-on conflicts between the replisome and transcription

complexes.

Bacillus subtilis (*B. subtilis*), which expresses a PriA homolog but not a PriC homolog,

employs a PriA-independent fork rescue pathway (Bruand et al., 2001). In this pathway, *B. subtilis*

DnaC (*Bsu*DnaC) helicase is reloaded using the helicase loader DnaI together with *B. subtilis* DnaB

(*Bsu*DnaB) and DnaD coloaders. The ssDNA-binding activity of *Bsu*DnaB has a crucial role in this

PriA-independent pathway (Bruand et al., 2005). SsDNA-bound *Bsu*DnaB remodels the SSB-ssDNA

complex with DnaD and recruits *Bsu*DnaC. These functions of *Bsu*DnaB are similar to the proposed

functions of PriC in replication fork rescue in *E. coli*. In addition, *Bsu*DnaC helicase is recruited to *oriC*

by concerted actions of *B. subtilis* DnaA and the DnaD-*Bsu*DnaB complex, which has an essential role

in chromosome replication initiation (Jameson & Wilkinson, 2017), suggesting a similar role of

*Bsu*DnaB to PriC in *E. coli* cells expressing DnaA F46A H136A. These similarities between PriC and

*Bsu*DnaB in these two evolutionarily distant bacterial species suggest that the mechanism of PriC-

promoted helicase loading is conserved among bacterial species despite the absence of sequence

conservation of PriC and *Bsu*DnaB homologs.

Based on our results, we propose the abnormal competition between DnaB and DiaA for

DnaA domain I could represent a form of intrinsic replication initiation stress in bacteria with conserved

DiaA homologs. This type of stress could also occur in ε-proteobacterial species, such as *Helicobacter*

pylori, because they express HobA, a DiaA-functional homolog (Natrajan et al., 2007; Zawilak-Pawlik

et al., 2011). Similarly, YfdR, encoded by *E. coli* prophages, inhibits replication initiation by competing

with DnaB for binding to DnaA domain I (Noguchi & Katayama, 2016), suggesting that such

extraneously introduced inhibitors could trigger replication initiation stress.

Even in human, ORC1, ORC2, and ORC5, the essential components of the eukaryotic

replication initiation complex, are not essential in some cancer cell-lines (Shibata & Dutta, 2020; Struhl

et al., 2016), suggesting that mechanisms of replication initiation rescue may also operate beyond the

bacterial kingdom. Therefore, further investigation of the initiation rescue processes and the factors

involved in diverse organisms from bacteria to human will be important for a full understanding of the

common principles and diverse mechanisms that ensure robust initiation of chromosomal DNA

replication.

 Materials and Methods

 Plasmids, proteins, and strains

The plasmids used in this study are listed in Key Resource Table. pKA234, pKW44-1, pNA135,

pBS*oriC*, pET22b(+)-priC, and pTKM601 were described previously (Aramaki et al., 2015; Ishida et

al., 2004; Kawakami et al., 2005; Keyamura et al., 2009; Kubota et al., 1997; Ozaki et al., 2008). For

the construction of pFH, an alanine substitution was introduced into pH136A with specific mutagenic

primer sets using the QuikChange site-directed mutagenesis protocol (Stratagene [Agilent], Agilent, La

Jolla, CA, United States) as described previously (Keyamura et al., 2007; Sakiyama et al., 2018).

WT DnaA and its derivative proteins were overproduced in *E. coli* strain KA450 from pKA234,

pKW44-1, or pFH and purified as described previously (Noguchi et al., 2015; Ozaki et al., 2008;

Sakiyama et al., 2017).

PriC protein was prepared as previously reported with minor modifications (Aramaki et al., 2013).

Briefly, PriC protein was overproduced in *E. coli* strain BL21-codonPlus-RIL by inducing its

expression from pET22b(+)-priC using 0.2 mM isopropyl-β-D-1-thiogalactopyranoside (IPTG). The

resulting cells were suspended in chilled buffer A (50 mM HEPES-NaOH [pH7.0], 10% sucrose, 1 mM

EDTA, 2 mM dithiothreitol (DTT), and 1 mM PMSF) and disrupted by sonication. Proteins in the

soluble fraction were precipitated with 0.24 g/mL ammonium sulfate, resuspended in a separation buffer

(50 mM imidazole [pH 7.0], 20% glycerol, 2 mM DTT, 1 mM EDTA, and 40 mM ammonium sulfate)

and loaded onto a 1 mL HiTrap® SP HP column (Cytiva, Uppsala, Sweden). Bound proteins were

eluted with a linear gradient from 0 to 1 M sodium chloride.

All *E. coli* strains used in this study are listed in Key Resource Table. KYA018, MIT125,

MIT162, SA103, NY20 and NY21 were described previously (Kasho & Katayama, 2013; Noguchi et

al., 2015; Noguchi & Katayama, 2016). Strains bearing mutant *oriC* (NY20L and NY20ATL) were

constructed using the λRed site-directed recombination system as previously described (Noguchi et al.,

2015; Sakiyama et al., 2017, 2022). NY20-frt strain and *oriC* mutant strains (NY21-frt, NY20L-frt, and

NY20ATL-frt) were constructed by eliminating *kan* from the NY20 strain and strains NY21, NY20L,

and NY20ATL. To remove the kanamycin-resistant cassette (*kan*) from *oriC*, FLP recombinase encoded

on pCP20 was used. Elimination of *kan* was verified by checking sensitivity to 50 μg/ml kanamycin in

LB agar plates. Δ*priC*::*frt-kan* from JW0456-KC and Δ*rep*::*frt-kan* from JW5604-KC were introduced

into MG1655 by P1 transduction, resulting in the construction of KRC002 and FF001, respectively. For

construction of other Δ*priC*::*frt-kan* strains (KRC004, KRC005, NY20-*priC*, NY21-*priC*, NY20L-*priC*,

NY20ATL-*priC*), P1 phage lysates prepared from KRC002 were used for transduction of strains

MIT125, KYA018, NY20-frt, SYM21-frt, NY20L-frt, and NY20ATL-frt. Transductants were screened

on LB agar plates containing 50 μg/mL kanamycin. Δ*diaA*::*frt-kan* or *rnhA*::*cat* derivatives (KRC006,

MIT125c, and KRC004c) were constructed using a similar protocol except that P1 phage lysates from SA103 (Δ*diaA*::*frt-kan*) or MIT162 (*rnhA*::*cat*) were used. KRC003 was generated by removal of *kan* from KRC002.

Buffers

Buffer P contained 60 mM HEPES–KOH (pH 7.6), 0.1 mM zinc acetate, 8 mM magnesium acetate, 30% [v/v] glycerol, and 0.32 mg/mL bovine serum albumin (BSA). Buffer N contained 50 mM HEPES–KOH (pH 7.6), 2.5 mM magnesium acetate, 0.3 mM EDTA, 7 mM DTT, 0.007% [v/v] Triton X-100, and 20% [v/v] glycerol. Form I* buffer contained 20 mM Tris-HCl (pH 7.5), 125 mM potassium glutamate, 10 mM magnesium acetate, 8 mM DTT, and 0.5 mg/mL BSA.

DUE unwinding assay

DUE unwinding assays were performed essentially as described with minor modifications (Sakiyama et al., 2022; Yoshida et al., 2023). Briefly, DnaA was incubated with ATP in buffer N on ice for 3 min to generate ATP-DnaA. The indicated amount of PriC and ATP-DnaA or its mutant derivatives was incubated for 9 min at 30°C in 10 µL buffer P containing 5 mM ATP, 125 mM potassium glutamate, 1.6 nM pBS*oriC*, and 42 nM IHF, followed by further incubation with 1.5 units of P1 nuclease (Wako) for 5 min. The reaction was stopped by the addition of 1% SDS and 25 mM EDTA, and DNA was purified by phenol-chloroform extraction and ethanol precipitation. One-third of each purified DNA was digested with AlwNI (NEB), which yielded 2.6 kb and 1.0 kb fragments after DUE unwinding of pBS*oriC*. The resultant DNA fragments were analyzed by 1% agarose gel electrophoresis with 1xTris-acetate-EDTA buffer for 30 min at 100 V, followed by ethidium bromide staining. Gel images were taken using the GelDoc GO imaging system (Bio-Rad Laboratories, Hercules, CA), and products derived from unwound plasmids were quantified using ImageJ software.

Form I* assay

This assay was performed as previously described with minor modifications (Baker et al., 1986; Noguchi et al., 2015; Sakiyama et al., 2022). The indicated amount of ATP-DnaA was incubated for 15 min at 30°C in 12.5 µL of Form I* buffer containing 3 mM ATP, 1.6 nM pBS*oriC*, 42 nM IHF, 400 nM His-DnaB, 400 nM DnaC, 76 nM GyrA, 100 nM His-GyrB, and 760 nM SSB. The reaction was stopped by the addition of 0.5% SDS, and DNA was purified by phenol-chloroform extraction. Samples were analyzed by 0.65% agarose gel electrophoresis with 0.5xTris-borate-EDTA buffer for 15 h at 23 V, followed by ethidium bromide staining.

Flow cytometry analysis

Flow cytometry analysis was performed essentially as described (Kasho et al., 2014). Briefly, cells were grown at 30°C in LB medium until the absorbance of the culture (A_{600}) reached 0.1. Portions of the cultures were diluted a thousand-fold into 5 ml LB medium and incubated at 30°C until the absorbance of the culture (A_{600}) reached 0.1. The remaining portions were further incubated to determine the doubling time (Td) by measuring A_{600} every 20 min. At A_{600} 0.1, aliquots of the cultures were fixed in 70% ethanol to analyze cell mass using the Multisizer 3 Coulter counter (Beckman Coulter, Brea, CA). The remaining cultures were further incubated for 4 h with 0.3 mg/mL rifampicin and 0.01 mg/mL cephalexin to allow run-out replication of chromosomal DNA. The resultant cells were fixed in 70% ethanol. After DNA staining with SYTOX Green (Thermo Fisher Scientific, Waltham, MA), cellular DNA contents were analyzed on a FACS Calibur flow cytometer (BD Bioscience, Franklin Lakes, NJ). The number of the origins/cell (ori/cell) was determined from the histograms of flow cytometry analysis. The number of ori/cell was divided by the mean cell mass determined by cell mass analysis, resulting in ori/mass ratios. The numbers of cells containing non-2^n copy number of *oriC* were divided by the number of cells containing 2^n copy number of *oriC* to calculate the asynchrony index (A.I.). For the analysis of cells bearing plasmids, LB medium was supplemented with 100 µg/mL ampicillin.

Replication initiation frequency test using synchronized cultures

MIT125 (*dnaA46 tnaA*::Tn*10*) cells or its Δ*priC* derivative KRC004 were grown at 30°C, a permissive temperature, in LB medium until the absorbance of the culture (A_{600}) reached 0.04, followed by further incubation for 90 min at 42°C, a restrictive temperature. Aliquots of the cultures were fixed in 70% ethanol as "Synchronization" samples. The remaining cultures were further incubated for 5 min to initiate DNA replication, followed by further incubation for 4 h at 42°C with 0.3 mg/mL rifampicin and 0.01 mg/mL cephalexin to allow run-out replication of chromosomal DNA. The resultant cells were fixed in 70% ethanol as "5 min release" samples. After DNA staining with SYTOX Green (Life Technologies), cellular DNA contents were analyzed on a FACS Calibur flow cytometer (BD Bioscience). Ratios of the origins/cell (ori/cell) were determined from the histograms of flow cytometry analysis.

For synchronization of *dnaC2* mutant cells (KYA018 and KRC005), cells were grown in LB medium at 30°C until the A_{600} reached 0.04, followed by further incubation for 80 min at 37°C, at the restrictive temperature. The resulting synchronized cultures were released for 5 min at 30°C and incubated in the presence of 0.3 mg/mL rifampicin and 0.01 mg/mL cephalexin at 30°C to allow run-out of chromosomal DNA replication.

qPCR for analysis of cSDR

Cells of MIT125 (*dnaA46 tnaA*::Tn*10*) cells and MIT125c (*dnaA46 tnaA*::Tn*10 rnhA*::*cat*) or its Δ*priC* derivative KRC004 (*dnaA46 tnaA*::Tn*10* Δ*priC*::*frt-kan*) and KRC004c (*dnaA46 tnaA*::Tn*10 rnhA*::*cat* Δ*priC*::*frt-kan*) were grown at 30°C in LB medium until the absorbance of the culture (A_{600}) reached 0.04 and aliquots of the cultures were withdrawn. The remaining samples were further incubated for 90 min at 40°C and aliquots of the cultures were withdrawn. These samples were boiled for 5 min at 95°C and the genome DNA was extracted. The levels of *oriC* (84.6 min), *ter* (32.4 min), and *ypaB* (50.5 min) were quantified by real-time qPCR using TB Green Premix Ex TaqII (Tli RNaseH Plus) (TaKaRa, Shiga, Japan) and the following primers: ORI_1 and KWoriCRev for *oriC*, qoriK fw and qoriK rev for *ter*, STM419 and STM420 for *ypaB*.

Chromosome loci copy-number analysis by whole-genome sequencing

The analysis was performed based on a previous paper with modifications (Brochu et al., 2018). Cells of the *dnaA46* mutant derivatives MIT125 (*dnaA46 tnaA*::Tn*10*), MIT125c (*dnaA46 tnaA*::Tn*10 rnhA*::*cat*), KRC004 (*dnaA46 tnaA*::Tn*10* Δ*priC*::*frt-kan*) and KRC004c (*dnaA46 tnaA*::Tn*10 rnhA*::*cat* Δ*priC*::*frt-kan*) were grown exponentially at 30°C in LB medium (50 mL). When the absorbance of the culture (A_{600}) reached 0.04, cells were incubated for 90 min at 40°C and collected by centrifugation. Cell pellets were dissolved by incubation at 37°C for 1h in TE buffer (0.6 mL) containing 0.5% SDS and 25 µg/ml proteinase K, and were mixed with 0.2 mL of buffer containing 4% cetyltrimethylammonium bromide, followed by further incubated for 10 minutes at 65°C, phenol-chloroform extraction and isopropanol precipitation. Precipitates were resuspended in Tris (pH7.5) buffer containing 0.1 mg/ml RNaseA, followed by incubation for 1 hour at 37°C and DNA purification by FastGene Gel/PCR Extraction Kit (Nippon Genetics, Tokyo, Japan). The quality control of genomic DNA, library preparation, and whole-genome sequencing were performed by Rhelixa, Inc. using NEBNext Ultra II DNA Library Prep Kit and Illumina NovaSeq X Plus. The number of reads (150 bp x 2 pair-ended) was at least 6.7 M (3.3 M pairs) per sample. The sequence data were analyzed using Galaxy (https://usegalaxy.org) and the coverage plot was generated essentially as described (Ivanova et al., 2015). Briefly, paired-end reads were mapped using the MG1655 reference genome (NC_000913) and the BWA-MEM algorithm, followed by calculation of the number of reads per genome coverage. DNA-sequence data are deposited in the NCBI Sequence Read Archive (Accession number: PRJNA1222470).

Immunoblot analysis

The assay was performed as described previously with minor modifications (Kawakami et al., 2005; Sakiyama et al., 2017; Yoshida et al., 2023). Briefly, MIT125 (*dnaA46 tnaA*::Tn*10*) and KRC004 (*dnaA46 tnaA*::Tn*10* Δ*priC*::*frt-kan*) cells bearing pING1, pKA234, or pFH were grown at 30°C in LB

medium including 100 µg/ml ampicillin until the absorbance of the culture (A_{600}) reached 0.05,

followed by incubation at 42°C for 90 min. Cells were harvested by centrifugation and dissolved in

SDS sample buffer to adjust the calculated A_{600} of 0.5. Proteins were separated using SDS-12%

polyacrylamide gel electrophoresis and were transferred to polyvinylidene difluoride membranes

(Millipore) using a semi-dry blotting method (Bio-Rad). After blocking at 30°C for 30 min in 3%

gelatin-TBS (3% gelatin, 20 mM Tris–HCl, 500 mM NaCl, pH 7.5), the membrane was incubated

overnight at 4°C in 1% gelatin-TTBS (20 mM Tris–HCl, 500 mM NaCl, 0.05% Tween-20, pH 7.5)

containing anti-DnaA rabbit antiserum (1:3000). After washing, the membrane was incubated at 30°C

for 1 h in 1% gelatin-TTBS containing goat anti-rabbit IgG antibody conjugated to alkaline phosphatase

(Bio-Rad), followed by development using AP Conjugate substrate Kit (Bio-Rad).

Acknowledgments

We are grateful to Drs. Yusuke Akama and Yukari Sakiyama for initial exploratory study related to this

work.

Fundings

Japan Society for the Promotion of Science (JSPS KAKENHI) [JP17H03656, JP20H03212 and

JP23K27131].

References

Abe, Y., Jo, T., Matsuda, Y., Matsunaga, C., Katayama, T., & Ueda, T. (2007). Structure and function of DnaA N-terminal domains: Specific sites and mechanisms in inter-DnaA interaction and in DnaB helicase loading on *oriC*. *Journal of Biological Chemistry*, *282*(24), 17816–17827. https://doi.org/10.1074/jbc.M701841200

Aramaki, T., Abe, Y., Furutani, K., Katayama, T., & Ueda, T. (2015). Basic and aromatic residues in the C-terminal domain of PriC are involved in ssDNA and SSB binding. *Journal of Biochemistry*, *157*(6), 529–537. https://doi.org/10.1093/jb/mvv014

Aramaki, T., Abe, Y., Ohkuri, T., Mishima, T., Yamashita, S., Katayama, T., & Ueda, T. (2013). Domain separation and characterization of PriC, a replication restart primosome factor in *Escherichia coli*. *Genes to Cells*, *18*(9), 723–732. https://doi.org/10.1111/gtc.12069

Arias-Palomo, E., Puri, N., O'Shea Murray, V. L., Yan, Q., & Berger, J. M. (2019). Physical Basis for the Loading of a Bacterial Replicative Helicase onto DNA. *Molecular Cell*, *74*(1), 173-184.e4. https://doi.org/10.1016/j.molcel.2019.01.023

Asai, T., & Kogomal, T. (1994). MINIREVIEW D-Loops and R-Loops: Alternative Mechanisms for the Initiation of Chromosome Replication in *Escherichia coli*. *Journal of Bacteriology*, *176*(7), 1807–1812. https://journals.asm.org/journal/jb

Baker, T. A., Sekimizu, K., Funnell, B. E., & Kornberg, A. (1986). Extensive Unwinding of the Plasmid Template during Staged Enzymatic Initiation of DNA Replication from the Origin of the Escherichia coli Chromosome. *Cell*, *45*(1), 53–64. https://doi.org/10.1016/0092-8674(86)90537-4.

Brochu, J., Vlachos-Breton, É., Sutherland, S., Martel, M., & Drolet, M. (2018). Topoisomerases I and III inhibit R-loop formation to prevent unregulated replication in the chromosomal *Ter* region of *Escherichia coli*. *PLoS Genetics*, *14*(9). https://doi.org/10.1371/journal.pgen.1007668

Bruand, C., Farache, M., McGovern, S., Ehrlich, S. D., & Polard, P. (2001). DnaB, DnaD and DnaI proteins are components of the *Bacillus subtilis* replication restart primosome. *Molecular Microbiology*, *42*(1), 245–256. https://doi.org/10.1046/j.1365-2958.2001.02631.x

Bruand, C., Velten, M., McGovern, S., Marsin, S., Sérèna, C., Dusko Ehrlich, S., & Polard, P. (2005). Functional interplay between the *Bacillus subtilis* DnaD and DnaB proteins essential for initiation and re-initiation of DNA replication. *Molecular Microbiology*, *55*(4), 1138–1150. https://doi.org/10.1111/j.1365-2958.2004.04451.x

Carr, K. M., & Kaguni, J. M. (1996). The A184V missense mutation of the *dnaA5* and *dnaA46* alleles confers a defect in ATP binding and thermolability in initiation of *Escherichia coli* DNA replication. *Molecular Microbiology*, *20*(6), 1307–1318. https://doi.org/10.1111/j.1365-2958.1996.tb02649.x

Chodavarapu, S., & Kaguni, J. M. (2016). Replication Initiation in Bacteria. *Enzymes*, *39*, 1–30. https://doi.org/10.1016/bs.enz.2016.03.001

Costa, A., & Diffley, J. F. X. (2022). The Initiation of Eukaryotic DNA Replication. *Annu. Rev. Biochem*, *91*, 107–131. https://doi.org/10.1146/annurev-biochem-072321

Duckworth, A. T., Ducos, P. L., McMillan, S. D., Satyshur, K. A., Blumenthal, K. H., Deorio, H. R., Larson, J. A., Sandler, S. J., Grant, T., & Keck, J. L. (2023). Replication fork binding triggers structural

changes in the PriA helicase that govern DNA replication restart in *E. coli*. *Nature Communications*, *14*(1), 1–13. https://doi.org/10.1038/s41467-023-38144-x

Duderstadt, K. E., Chuang, K., & Berger, J. M. (2011). DNA stretching by bacterial initiators promotes replication origin opening. *Nature*, *478*(7368), 209–213. https://doi.org/10.1038/nature10455

Erzberger, J. P., Mott, M. L., & Berger, J. M. (2006). Structural basis for ATP-dependent DnaA assembly and replication-origin remodeling. *Nature Structural and Molecular Biology*, *13*(8), 676–683. https://doi.org/10.1038/nsmb1115

Fang, L., Davey, M. J., & O'donnell, M. (1999). Replisome Assembly at *oriC*, the Replication Origin of *E. coli*, Reveals an Explanation for Initiation Sites outside an Origin. *Molecular Cell*, *4*, 541–553. https://doi.org/10.1016/s1097-2765(00)80205-1

Felczak, M. M., & Kaguni, J. M. (2004). The box VII motif of *Escherichia coli* DnaA protein is required for DnaA oligomerization at the *E. coli* replication origin. *Journal of Biological Chemistry*, *279*(49), 51156–51162. https://doi.org/10.1074/jbc.M409695200

Flores, M. J., Ehrlich, S. D., & Michel, B. (2002). Primosome assembly requirement for replication restart in the *Escherichia coli holDG10* replication mutant. *Molecular Microbiology*, *44*(3), 783–792. https://doi.org/10.1046/j.1365-2958.2002.02913.x

Fujikawa, N., Kurumizaka, H., Nureki, O., Terada, T., Shirouzu, M., Katayama, T., & Yokoyama, S. (2003). Structural basis of replication origin recognition by the DnaA protein. *Nucleic Acids Research*, *31*(8), 2077–2086. https://doi.org/10.1093/nar/gkg309

Grimwade, J. E., & Leonard, A. C. (2021). Blocking, Bending, and Binding: Regulation of Initiation of Chromosome Replication During the *Escherichia coli* Cell Cycle by Transcriptional Modulators That Interact With Origin DNA. *Frontiers in Microbiology*, *12*, 732270. https://doi.org/10.3389/fmicb.2021.732270

Hayashi, C., Miyazaki, E., Ozaki, S., Abe, Y., & Katayama, T. (2020). DnaB helicase is recruited to the replication initiation complex via binding of DnaA domain I to the lateral surface of the DnaB N-terminal domain. *Journal of Biological Chemistry*, *295*(32), 1131–1143. https://doi.org/10.1074/jbc.ra120.014235

Heller, R. C., & Marians, K. J. (2005a). The disposition of nascent strands at stalled replication forks dictates the pathway of replisome loading during restart. *Molecular Cell*, *17*(5), 733–743. https://doi.org/10.1016/j.molcel.2005.01.019

Heller, R. C., & Marians, K. J. (2005b). Unwinding of the nascent lagging strand by Rep and PriA enables the direct restart of stalled replication forks. *Journal of Biological Chemistry*, *280*(40), 34143–34151. https://doi.org/10.1074/jbc.M507224200

Heller, R. C., & Marians, K. J. (2006). Replication fork reactivation downstream of a blocked nascent leading strand. *Nature*, *439*(7076), 557–562. https://doi.org/10.1038/nature04329

Hinds, T., & Sandler, S. J. (2004). Allele specific synthetic lethality between *priC* and *dnaAts* alleles at the permissive temperature of 30°C in *E. coli* K-12. *BMC Microbiology*, *4*, 1–9. https://doi.org/10.1186/1471-2180-4-47

Hwang, D. S., & Kaguni, J. M. (1988). Purification and Characterization of the *dnaA46* Gene Product.

726   *Journal of Biological Chemistry* 263(22), 10625–10632.

727 Ivanova, D., Taylor, T., Smith, S.L. et al. (2015) Shaping the landscape of the *Escherichia coli* chromosome:

728   replication-transcription encounters in cells with an ectopic replication origin. *Nucleic Acids Research,*

729   43:7865–7877. https://doi.org/10.1093/nar/gkv704

730 Ishida, T., Akimitsu, N., Kashioka, T., Hatano, M., Kubota, T., Ogata, Y., Sekimizu, K., & Katayama, T.

731   (2004). DiaA, a novel DnaA-binding protein, ensures the timely initiation of *Escherichia coli*

732   chromosome replication. *Journal of Biological Chemistry*, *279*(44), 45546–45555.

733   https://doi.org/10.1074/jbc.M402762200

734 Jameson, K. H., & Wilkinson, A. J. (2017). Control of initiation of DNA replication in *Bacillus subtilis* and

735   *Escherichia coli*. *Genes*, *8*(1), 22. https://doi.org/10.3390/genes8010022

736 Kasho, K., Fujimitsu, K., Matoba, T., Oshima, T., & Katayama, T. (2014). Timely binding of IHF and Fis to

737   *DARS2* regulates ATP-DnaA production and replication initiation. *Nucleic Acids Research*, *42*(21),

738   13134–13149. https://doi.org/10.1093/nar/gku1051

739 Kasho, K., & Katayama, T. (2013). DnaA binding locus *datA* promotes DnaA-ATP hydrolysis to enable cell

740   cycle-coordinated replication initiation. *Proceedings of the National Academy of Sciences of the United*

741   *States of America*, *110*(3), 936–941. https://doi.org/10.1073/pnas.1212070110

742 Kasho, K., Ozaki, S., & Katayama, T. (2023). IHF and Fis as *Escherichia coli* Cell Cycle Regulators:

743   Activation of the Replication Origin *oriC* and the Regulatory Cycle of the DnaA Initiator. *International*

744   *Journal of Molecular Sciences*, *24*(14), 11572. https://doi.org/10.3390/ijms241411572

745 Kawakami, H., Keyamura, K., & Katayama, T. (2005). Formation of an ATP-DnaA-specific initiation

746   complex requires DnaA arginine 285, a conserved motif in the AAA+ protein family. *Journal of*

747   *Biological Chemistry*, *280*(29), 27420–27430. https://doi.org/10.1074/jbc.M502764200

748 Keyamura, K., Abe, Y., Higashi, M., Ueda, T., & Katayama, T. (2009). DiaA dynamics are coupled with

749   changes in initial origin complexes leading to helicase loading. *Journal of Biological Chemistry*,

750   *284*(37), 25038–25050. https://doi.org/10.1074/jbc.M109.002717

751 Keyamura, K., Fujikawa, N., Ishida, T., Ozaki, S., Su'etsugu, M., Fujimitsu, K., Kagawa, W., Yokoyama, S.,

752   Kurumizaka, H., & Katayama, T. (2007). The interaction of DiaA and DnaA regulates the replication

753   cycle in *E. coli* by directly promoting ATP-DnaA-specific initiation complexes. *Genes and*

754   *Development*, *21*(16), 2083–2099. https://doi.org/10.1101/gad.1561207

755 Kogoma, T. (1997). Stable DNA Replication: Interplay between DNA Replication, Homologous

756   Recombination, and Transcription. *MICROBIOLOGY AND MOLECULAR BIOLOGY REVIEWS*,

757   *61*(2), 212–238. https://journals.asm.org/journal/mmbr

758 Kubota, T., Katayama, T., Ito, Y., Mizushima, T., & Sekimizu, K. (1997). Conformational Transition of DnaA

759   Protein by ATP: Structural Analysis of DnaA Protein, the Initiator of *Escherichia coli* Chromosome

760   Replication. *Biochemical and Biophysical Research Communications*, *232*, 130–135.

761   https://doi.org/10.1006/bbrc.1997.6244

762 Lopper, M., Boonsombat, R., Sandler, S. J., & Keck, J. L. (2007). A Hand-Off Mechanism for Primosome

763   Assembly in Replication Restart. *Molecular Cell*, *26*(6), 781–793.

764   https://doi.org/10.1016/j.molcel.2007.05.012

Maduike, N. Z., Tehranchi, A. K., Wang, J. D., & Kreuzer, K. N. (2014). Replication of the *Escherichia coli*
chromosome in RNase HI-deficient cells: Multiple initiation regions and fork dynamics. *Molecular*
Microbiology, *91*(1), 39–56. https://doi.org/10.1111/mmi.12440

Masai, H., Asai, T., Kubota, Y., Arai, K. I., & Kogoma, T. (1994). *Escherichia coli* PriA protein is essential
for inducible and constitutive stable DNA replication. *EMBO Journal*, *13*(22), 5338–5345.
https://doi.org/10.1002/j.1460-2075.1994.tb06868.x

McMillan, S. D., & Keck, J. L. (2024). Biochemical characterization of *Escherichia coli* DnaC variants that
alter DnaB helicase loading onto DNA. *Journal of Biological Chemistry*, *300*(5), 107275.
https://doi.org/10.1016/j.jbc.2024.107275

Michel, B., & Sandler, S. J. (2017). Replication restart in bacteria. *Journal of Bacteriology*, *199*(13), e00102-
17. https://doi.org/10.1128/JB.00102-17

Miller, D. T., Grimwade, J. E., Betteridge, T., Rozgaja, T., Torgue, J.-C., Leonard, A. C., & Helinski, D. R.
(2009). Bacterial origin recognition complexes direct assembly of higher-order DnaA oligomeric
structures. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(44),
18479–18484. www.pnas.orgcgidoi10.1073pnas.0909472106

Nagata, K., Okada, A., Ohtsuka, J., Ohkuri, T., Akama, Y., Sakiyama, Y., Miyazaki, E., Horita, S., Katayama,
T., Ueda, T., & Tanokura, M. (2020). Crystal structure of the complex of the interaction domains of
Escherichia coli DnaB helicase and DnaC helicase loader: Structural basis implying a distortion-
accumulation mechanism for the DnaB ring opening caused by DnaC binding. *Journal of*
Biochemistry, *167*(1), 1–14. https://doi.org/10.1093/jb/mvz087

Natrajan, G., Hall, D. R., Thompson, A. C., Gutsche, I., & Terradot, L. (2007). Structural similarity between
the DnaA-binding proteins HobA (HP1230) from *Helicobacter pylori* and DiaA from *Escherichia coli*.
Molecular Microbiology, *65*(4), 995–1005. https://doi.org/10.1111/j.1365-2958.2007.05843.x

Né Dicte Michel, B., Flores, M.-J., Viguera, E., Grompone, G., Seigneur, M., & Bidnenko, V. (2001). Rescue
of arrested replication forks by homologous recombination. *Proceedings of the National Academy of*
Sciences of the United States of America, *17*(15), 8181–8188.
www.pnas.orgcgidoi10.1073pnas.111008798

Nguyen, B., Shinn, M. K., Weiland, E., & Lohman, T. M. (2021). Regulation of *E. coli* Rep helicase activity
by PriC. *Journal of Molecular Biology*, *433*(15), 167072. https://doi.org/10.1016/j.jmb.2021.167072

Noguchi, Y., & Katayama, T. (2016). The *Escherichia coli* cryptic prophage protein YfdR binds to DnaA and
initiation of chromosomal replication is inhibited by overexpression of the gene cluster *yfdQ-yfdR-*
yfdS-yfdT. *Frontiers in Microbiology*, *7*(MAR), 1–21. https://doi.org/10.3389/fmicb.2016.00239

Noguchi, Y., Sakiyama, Y., Kawakami, H., & Katayama, T. (2015). The Arg fingers of key DnaA protomers
are oriented inward within the replication origin *oriC* and stimulate DnaA subcomplexes in the
initiation complex. *Journal of Biological Chemistry*, *290*(33), 20295–20312.
https://doi.org/10.1074/jbc.M115.662601

Nozaki, S., & Ogawa, T. (2008). Determination of the minimum domain II size of *Escherichia coli* DnaA
protein essential for cell viability. *Microbiology*, *154*(11), 3379–3384.
https://doi.org/10.1099/mic.0.2008/019745-0

O'Donnell, M., Langston, L., & Stillman, B. (2013). Principles and concepts of DNA replication in bacteria,

archaea, and eukarya. *Cold Spring Harbor Perspectives in Biology*, *5*(7), a010108.

https://doi.org/10.1101/cshperspect.a010108

Ozaki, S., & Katayama, T. (2012). Highly organized DnaA-*oriC* complexes recruit the single-stranded DNA

for replication initiation. *Nucleic Acids Research*, *40*(4), 1648–1665.

https://doi.org/10.1093/nar/gkr832

Ozaki, S., Kawakami, H., Nakamura, K., Fujikawa, N., Kagawa, W., Park, S. Y., Yokoyama, S., Kurumizaka,

H., & Katayama, T. (2008). A common mechanism for the ATP-DnaA-dependent formation of open

complexes at the replication origin. *Journal of Biological Chemistry*, *283*(13), 8351–8362.

https://doi.org/10.1074/jbc.M708684200

Ozaki, S., Noguchi, Y., Hayashi, Y., Miyazaki, E., & Katayama, T. (2012). Differentiation of the DnaA-*oriC*

subcomplex for DNA unwinding in a replication initiation complex. *Journal of Biological Chemistry*,

287(44), 37458–37471. https://doi.org/10.1074/jbc.M112.372052

Rozgaja, T. A., Grimwade, J. E., Iqbal, M., Czerwonka, C., Vora, M., & Leonard, A. C. (2011). Two

oppositely oriented arrays of low-affinity recognition sites in *oriC* guide progressive binding of DnaA

during *Escherichia coli* pre-RC assembly. *Molecular Microbiology*, *82*(2), 475–488.

https://doi.org/10.1111/j.1365-2958.2011.07827.x

Sakiyama, Y., Kasho, K., Noguchi, Y., Kawakami, H., & Katayama, T. (2017). Regulatory dynamics in the

ternary DnaA complex for initiation of chromosomal replication in *Escherichia coli*. *Nucleic Acids*

Research, *45*(21), 12354–12373. https://doi.org/10.1093/nar/gkx914

Sakiyama, Y., Nagata, M., Yoshida, R., Kasho, K., Ozaki, S., & Katayama, T. (2022). Concerted actions of

DnaA complexes with DNA-unwinding sequences within and flanking replication origin *oriC* promote

DnaB helicase loading. *Journal of Biological Chemistry*, *298*(6), 102051.

https://doi.org/10.1016/j.jbc.2022.102051

Sakiyama, Y., Nishimura, M., Hayashi, C., Akama, Y., Ozaki, S., & Katayama, T. (2018). The DnaA AAA+

Domain His136 residue directs DnaB replicative helicase to the unwound region of the replication

origin, *oriC*. *Frontiers in Microbiology*, *9*(AUG), 1–17. https://doi.org/10.3389/fmicb.2018.02017

Sandler, S. J. (2000). Multiple Genetic Pathways for Restarting DNA Replication Forks in *Escherichia coli*

K-12. *Genetics*, *155*, 487–497. https://academic.oup.com/genetics/article/155/2/487/6048048

Sandler, S. J., Marians, K. J., Zavitz, K. H., Coutu, J., Parent, M. A., & Clark, A. J. (1999). *dnaC* mutations

suppress defects in DNA replication- and recombination-associated functions in *priB* and *priC* double

mutants in *Escherichia coli* K-12. *Molecular Microbiology*, *34*(1), 91–101.

https://doi.org/10.1046/j.1365-2958.1999.01576.x

Shibata, E., & Dutta, A. (2020). A human cancer cell line initiates DNA replication normally in the absence

of ORC5 and ORC2 proteins. *Journal of Biological Chemistry*, *295*(50), 16949–16959.

https://doi.org/10.1074/jbc.RA120.015450

Shimizu, M., Noguchi, Y., Sakiyama, Y., Kawakami, H., Katayama, T., & Takada, S. (2016). Near-atomic

structural model for bacterial DNA replication initiation complex and its functional insights.

Proceedings of the National Academy of Sciences of the United States of America, *113*(50), E8021–

E8030. https://doi.org/10.1073/pnas.1609649113

Shimuta, T. R., Nakano, K., Yamaguchi, Y., Ozaki, S., Fujimitsu, K., Matsunaga, C., Noguchi, K., Emoto, A.,

& Katayama, T. (2004). Novel heat shock protein HspQ stimulates the degradation of mutant DnaA

protein in *Escherichia coli*. *Genes to Cells*, *9*(12), 1151–1166. https://doi.org/10.1111/j.1365-

2443.2004.00800.x

Stepankiw, N., Kaidow, A., Boye, E., & Bates, D. (2009). The right half of the *Escherichia coli* replication

origin is not essential for viability, but facilitates multi-forked replication. *Molecular Microbiology*,

74(2), 467–479. https://doi.org/10.1111/j.1365-2958.2009.06877.x

Struhl, K., Shibata, E., Kiran, M., Shibata, Y., Singh, S., Kiran, S., & Dutta, A. (2016). Two subunits of

human ORC are dispensable for DNA replication and proliferation. *eLife*, *5*, e19084.

https://doi.org/10.7554/eLife.19084.001

Wegrzyn, K., & Konieczny, I. (2023). Toward an understanding of the DNA replication initiation in bacteria.

Frontiers in Microbiology, *14*, 1328842. https://doi.org/10.3389/fmicb.2023.1328842

Wessel, S. R., Cornilescu, C. C., Cornilescu, G., Metz, A., Leroux, M., Hu, K., Sandler, S. J., Markley, J. L.,

& Keck, J. L. (2016). Structure and function of the PriC DNA replication restart protein. *Journal of

Biological Chemistry*, *291*(35), 18384–18396. https://doi.org/10.1074/jbc.M116.738781

Wessel, S. R., Marceau, A. H., Massoni, S. C., Zhou, R., Ha, T., Sandler, S. J., & Keck, J. L. (2013). PriC-

mediated DNA replication restart requires PriC complex formation with the single-stranded DNA-

binding protein. *Journal of Biological Chemistry*, *288*(24), 17569–17578.

https://doi.org/10.1074/jbc.M113.478156

Windgassen, T. A., & Keck, J. L. (2016). An aromatic-rich loop couples DNA binding and ATP hydrolysis in

the PriA DNA helicase. *Nucleic Acids Research*, *44*(20), 9745–9757.

https://doi.org/10.1093/nar/gkw690

Windgassen, T. A., Leroux, M., Satyshur, K. A., Sandler, S. J., & Keck, J. L. (2018). Structure-specific DNA

replication-fork recognition directs helicase and replication restart activities of the PriA helicase.

Proceedings of the National Academy of Sciences of the United States of America, *115*(39), E9075–

E9084. https://doi.org/10.1073/pnas.1809842115

Withers, H. L., & Bernander, R. (1998). Characterization of *dnaC2* and *dnaC28* Mutants by Flow Cytometry.

JOURNAL OF BACTERIOLOGY, *180*(7), 1624–1631. https://journals.asm.org/journal/jb

Yoshida, R., Ozaki, S., Kawakami, H., & Katayama, T. (2023). Single-stranded DNA recruitment mechanism

in replication origin unwinding by DnaA initiator protein and HU, an evolutionary ubiquitous nucleoid

protein. *Nucleic Acids Research*, *51*(12), 6286–6306. https://doi.org/10.1093/nar/gkad389

Zawilak-Pawlik, A., Donczew, R., Szafrański, S., MacKiewicz, P., Terradot, L., & Zakrzewska-Czerwińska,

J. (2011). DiaA/HobA and DnaA: A pair of proteins co-evolved to cooperate during bacterial orisome

assembly. *Journal of Molecular Biology*, *408*(2), 238–251. https://doi.org/10.1016/j.jmb.2011.02.045

 Figure legend

Figure 1 Schematic structures of *oriC* and DnaA and the replication initiation mechanism.

(A) Overall structure of *oriC* with the AT-rich cluster. *oriC* (245 bp) includes the Duplex-Unwinding

Element (DUE, purple bar) and the DnaA-Oligomerization Region (DOR, gray bar). DUE is composed

of three AT-rich 13-mer repeats termed L, M, and R. Dark purple boxes indicate the specific sequences

involved in DnaA-ssDNA binding (TT[A/G]T(T)). The AT-cluster (purple dots) flanking the DUE

outside *oriC* is a supplementary unwinding region. The DOR contains 12 DnaA boxes (filled and open

arrowheads representing sites with the full consensus TTA[A/T]NCACA sequence and sites with

mismatches, respectively) and an IHF-binding region (IBR, green box). The DOR is subdivided into

Left, Middle, and Right subregions. (B) Domains of DnaA. Domains I-IV are shown schematically,

with amino acid residue numbers shown in each bracket. H/B motifs (V211 and R245, squares) and

Arginine finger (R285, triangle) are indicated. The major functions of each domain are described on the

right side of the structure. (C) DUE unwinding by the ssDUE recruitment mechanism. DnaA and IHF

are indicated by red and green diagrams, respectively. When ATP-DnaA oligomerizes, DUE is unwound

unstably by thermal motion and torsional stress. The M/R region of the DUE-upper strand (purple line)

binds to R1-DnaA and R5M-DnaA via IHF-induced DNA bending, resulting in stable DUE unwinding.

In addition to M/R-DUE, L-DUE is moderately unwound and interacts with the Right-DnaA

subcomplex. (D) Mechanism for DnaB loading. Each Left-DnaA and Right-DnaA subcomplex binds to

a DnaB hexamer (blue ovals) complexed with DnaC (yellow circles). (E) PriC-dependent fork rescue

pathways. PriC (vivid red circle) promotes DnaB reloading to the abandoned fork, which exposes

ssDNA regions. PriC binds to ssDNA at the replication fork, which remodels the SSB-ssDNA complex

(for simplicity, SSB is omitted) for DnaC-dependent reloading of DnaB (*upper panel*). If the ssDNA

region is accompanied with a nascent lagging strand, Rep helicase (green shape) is recruited by PriC

and unwinds the lagging strand to expose the ssDNA region for DnaB loading (the lower case).

Figure 2. Requirement of *priC* for DNA replication initiation in *dnaA46* and *dnaC2* strains.

(A) Cell growth abilities. MG1655 and its derivatives MIT125 (MG1655 *dnaA46 tnaA*::Tn*10*), KYA018

(MG1655 *dnaC2 zjj18*::*cat*), KRC002 (MG1655 Δ*priC*::*frt-kan*), KRC004 (MG1655 *dnaA46 tnaA*::Tn*10*

Δ*priC*::*frt-kan*), and KRC005 (MG1655 *dnaC2 zjj18*::*cat* Δ*priC*::*frt-kan*) were grown overnight. 10-fold-

serial dilutions of the overnight cultures (~10^9 cells/mL) were incubated for 16 h at 30°C, or for 14 h at

35/37/42°C on LB agar medium. Three independent experiments were performed. +, WT *priC*; -,

Δ*priC*::*frt-kan*. The images of representative plates are shown.

(B) DNA replication initiation in synchronized *dnaA46* cells with or without *priC*. Exponentially growing

MIT125 (MG1655 *dnaA46 tnaA*::Tn*10*) (+) and KRC004 (MG1655 *dnaA46 tnaA*::Tn*10* Δ*priC*::*frt-kan*)

(-) cells were synchronized to the pre-initiation stage by incubation for 90 min at 42°C in LB medium, followed by incubation for 5 min at 30°C to initiate DNA replication, which was further followed by incubation with rifampicin and cephalexin for 4 h at 42°C to allow run-out of ongoing DNA replication. Samples were withdrawn after synchronization and run-out replication. The DNA contents of these samples were measured by flow cytometry. Representative histograms of three independent experiments are shown. For each sample, the mean number of cells with more than one chromosome equivalent after a 5 min release was quantified and is shown as a percentage with standard deviations.

(C) DNA replication initiation in synchronized *dnaC2* cells with or without *priC*. Exponentially growing KYA018 (MG1655 *dnaC2 zjj18*::*cat*) (+) and KRC005 (MG1655 *dnaC2 zjj18*::*ca* Δ*priC*::*frt-kan*) (-) cells were synchronized to the pre-initiation stage by incubation for 80 min at 37°C in LB medium, followed by incubation for 5 min at 30°C and further incubation with rifampicin and cephalexin as described above. Samples were withdrawn and analyzed as described above.

Figure 3. Role of PriC in growth of DnaA mutant defective in interactions with DnaB.

Cell growth of MIT125 (MG1655 *dnaA46 tnaA*::Tn*10*) and KRC004 (MG1655 *dnaA46 tnaA*::Tn*10* Δ*priC*::*frt-kan*) cells bearing pING1 plasmid vector (-) or its derivatives pKA234 expressing WT DnaA (WT) or pFH expressing DnaA F46A H136A (F46A H136A). Cells were grown at 30°C overnight and 10-fold-serial dilutions of the cultures (~10^9 cells/mL) were incubated on LB agar plates containing ampicillin for 16 h at 30°C and for 14 h at 42°C, respectively. Four independent experiments were performed. +, WT *priC*; -, Δ*priC*::*frt-kan*.

Figure 4. PriC stimulation of DNA replication initiation inhibited by DiaA oversupply.

(A) Cell growth of MG1655, KRC002 (MG1655 Δ*priC*::*frt-kan*), and FF001 (MG1655 Δ*rep*::*frt-kan*) cells bearing pBR322 or pNA135 (pBR322 derivatives carrying *diaA*). Overnight cultures (~10^9 cells/mL) of MG1655 and KRC002 cells bearing pBR322 or pNA135 were 10-fold serially diluted and incubated on LB agar plates containing ampicillin for 40 h at 25°C. The results of three independent experiments were consistent, and one which is shown.

(B) Flow cytometry analyses of MG1655 and KRC002 (MG1655 Δ*priC*::*frt-kan*) cells bearing pBR322 or pNA135. Cells were exponentially grown at 30°C in LB medium with ampicillin, followed by further incubation with rifampicin and cephalexin to allow run-out of chromosomal DNA replication. DNA contents were quantified by flow cytometry, and cell size (mass) at the time of drug addition was measured by a Coulter counter. Mean mass, ori/mass ratio, doubling time (Td), and asynchrony index (A.I.: the percentage of cell numbers with non-2^n copies of *oriC* per the cell numbers with 2^n copies of *oriC*) of each strain are indicated at the top right of each panel with standard deviations. Three to four

independent experiments were performed. +, WT *priC*; -, Δ*priC*::*frt-kan*.

(C) Flow cytometry analyses of MG1655, KRC003 (MG1655 Δ*priC*::*frt*), SA103 (MG1655 Δ*diaA*::*frt-*

kan), and KRC006 (MG1655 Δ*priC*::*frt*, Δ*diaA*::*frt-kan*) cells. Cells were grown and analyzed as

described above. Three independent experiments were performed. For "*priC*" column, +, WT *priC*; -,

Δ*priC*::*frt.* For "*diaA*" column, +, WT *diaA*; -, Δ*diaA*::*frt-kan.*

Figure 5. Stimulation of replication initiation by PriC in Right-DnaA subcomplex-defective cells

(A) Schematic structure of Left-*oriC* and R4*Tma oriC.* AT-rich region, DUE, and DOR are indicated by

a gray dotted box, a gray box, and an open box, respectively. Filled and open arrowheads in the DOR

show DnaA boxes with the full consensus sequence or mismatches. In addition, a striped box indicates

the IBR. The regions of each mutant *oriC* are indicated below the structure as open boxes. The position

of the *Tma*DnaA box substitution is indicated by a yellow arrowhead.

(B) Cell growth of WT *oriC* and Left-*oriC* cells with or without *priC*. Overnight cultures (~10^9 cells/mL)

of NY20-frt (WT *oriC*; WT) and NY20L-frt (Left-*oriC*; Left) cells with (+) or without (-) *priC* were 10-

fold serially diluted and incubated on LB agar plates for 24 h at 25°C, for 16 h at 30°C, and for 14 h at

37°C. +, WT *priC*; -, Δ*priC*::*frt-kan*. Three independent experiments were performed.

(C) Cell growth of WT *oriC* and *oriC* R4-box mutant cells with or without *priC*. NY20-frt (WT *oriC*)

and NY24-frt (*oriC* with R4*Tma* substitution, R4*Tma*) cells with (+) or without (-) *priC* were grown as

described in panel A. +, WT *priC*; -, Δ*priC*::*frt-kan*. Four independent experiments were performed.

(D) Flow cytometry analyses of WT and R4*Tma* cells with or without *priC*. Cells were exponentially

grown at 30°C in LB medium with ampicillin and analyzed as described in Figure 4. Representative

histograms from four independent experiments are shown. Mean mass, ori/mass, and Td of each strain

are indicated at the top right of each panel with standard deviations.

Figure 6. Stimulation of replication initiation by PriC in cells with a deletion in the ssDNA-

expanding region of *oriC*

(A) Schematic structure of subATL *oriC* as shown in Figure 5. The *oriC* region including the subATL

oriC is indicated below the structure as a white box.

(B) Cell growth of WT *oriC* and *oriC* subATL mutant cells with or without WT *priC*. NY20-frt (WT

oriC) and NY20ATL-frt (*oriC* lacking AT and L sequences, subATL) with (+) or without (-) WT *priC*

were grown as described in Figure 5. +, WT *priC*; -, Δ*priC*::*frt-kan.* Three independent experiments were

performed.

(C) Flow cytometry analyses of WT *oriC* and *oriC* subATL mutant cells with or without WT *priC*. Cells

were exponentially grown at 30°C in LB medium and analyzed as described in Figure 4. Representative

histograms from three independent experiments are shown. Mean mass, ori/mass, and Td of each strain

are indicated at the top right of each panel with standard deviations.

Figure 7. DnaB loading onto the ssDUE by PriC.

(A and B) *In vitro* reconstituted system for DnaB loading. Form I of *oriC* plasmid (pBS*oriC*; 1.6 nM) was

incubated for 15 min at 30°C with the indicated amount of PriC in the presence of ATP-DnaA or its mutant

derivatives, DnaB, DnaC, IHF, gyrase, and SSB. The resulting plasmids were purified and analyzed using

agarose gel electrophoresis. A representative image of three independent experiments is shown in the

black/white-inverted mode (A). Band intensities of each lane in the gel image were analyzed by

densitometric scanning. The percentages of form I* *oriC* plasmid per input DNA are shown as 'Form I*

(%)'. Mean and standard deviations are shown (B). Abbreviations in panel B: -, no addition of DnaA;

WT, wild-type DnaA; FH, DnaA F46A H136A; VA, DnaA V211A.

(C and D) *In vitro* reconstituted system containing DiaA. Form I of pBS*oriC* plasmid was used as

described above except for the addition of the indicated amounts of DiaA. A representative image of three

independent experiments is shown in black/white-inverted mode (C). Band intensities of each lane in the

gel image were analyzed by densitometric scanning. The percentages of form I**oriC* plasmid together

with the mean and standard deviations are shown as described above (D).

Figure 8. PriC stimulation of cSDR-dependent cell growth.

(A) Cell growth abilities. Cells of MIT125 (MG1655 *dnaA46*), MIT125c (MG1655 *dnaA46 rnhA*::*cat*),

KRC004 (MG1655 *dnaA46* Δ*priC*::*frt-kan*), and KRC004c (MG1655 *dnaA46* Δ*priC*::*frt-kan rnhA*::*cat*)

were grown at 30°C overnight and 10-fold-serial dilutions of the overnight cultures (~10^9 cells/mL) were

incubated for 16 h at 30°C and for 36 h at 40°C on LB agar plates. Three independent experiments were

performed. +, WT; -, deletion.

(B) *oriC* and *ter* copy numbers. MIT125, MIT125c, KRC004, and KRC004c cells growing exponentially

at 30°C were further incubated for 90 min at 40°C in LB medium. Samples were withdrawn before and

after the 40°C incubation. The genome DNA of each sample was extracted by boiling the cells for 5 min

at 95°C. The relative copy numbers of *oriC* (84.6 min) and *ter* (32.4 min) to that of *ypaB* (50.5 min) were

quantified using real-time qPCR. The data and averages of three independent experiments are shown with

standard deviations.

1012

Figure 9. Model for PriC-promoted replication initiation.

PriC-promoted rescue of blocked DnaA-*oriC* dependent replication initiation. *oriC*, DnaA, IHF, DnaB,

and DnaC are illustrated as depicted in Figure 1C. When tight DnaA binding of DiaA (dark gray circle)

(or DnaA F46A H136A double mutations) inhibits DnaB recruitment, PriC (Brilliant red circle) binds to

a stably unwound DUE strand, recruiting a DnaB-DnaC complex to lower strand of DUE instead of

Left-DnaA subcomplex. When a sufficient ssDNA region is exposed, PriC then recruits the DnaB-DnaC

complex to the opposite strand.

1020

Figure 2-figure supplement 1. Role of PriC in cell growth of *dnaA46* and *dnaC2* cells.

pBR322 vector or pBR322 with *priC* (pBR*priC*) was introduced into cells of MG1655 (*dnaA/dnaC*

WT), KRC002 (MG1655 Δ*priC*::*frt-kan*), MIT125 (MG1655 *dnaA46 tnaA*::Tn*10*), or KRC004

(MG1655 *dnaA46 tnaA*::Tn*10* Δ*priC*::*frt-kan*) (A). The same plasmids were also introduced into cells

of MG1655, KRC002 , KYA018 (MG1655 *dnaC2 zjj18*::*cat*) or KRC005 (MG1655 *dnaC2 zjj18*::*cat*

Δ*priC*::*frt-kan*) (B). Independent transformants were assessed for the cell growth by spotting assays

with incubation at the indicated temperatures for 24 hrs on LB agar plates supplemented with 100 μg/ml

ampicillin as described for Figure 2A.

1029

Figure 2-figure supplement 2. Role of PriC in cell growth of cells with WT DnaA and DnaC.

(A) Flow cytometry analyses of MG1655 and KRC002 (MG1655 Δ*priC*::*frt-kan*) cells in LB medium.

Cells were exponentially grown at 25°C, 30°C, and 42°C in LB medium, followed by further incubation

with rifampicin and cephalexin to allow run-out replication of chromosomal DNA. DNA contents were

quantified by flow cytometry, and cell size (mass) at the time of drug addition was measured by a Coulter

counter. Mean mass, ori/mass ratio, doubling time (Td), and asynchrony index (A.I.: the percentage of

cell numbers with non-2^n copies of *oriC* per the cell numbers with 2^n copies of *oriC*) of each strain are

indicated at the top right of each panel with standard deviations. Three to four independent experiments

were performed. +, WT *priC*; -, Δ*priC*::*frt-kan*.

(B) Flow cytometry analyses of MG1655 and KRC002 (MG1655 Δ*priC*::*frt-kan*) cells in M9 minimal

medium including 0.2% glucose and 0.2% casamino acids. Cells were grown and analyzed as described

above. Three independent experiments were performed. For "*priC*" column, +, WT *priC*; -, Δ*priC*::*frt*.

1042

Figure 3-figure supplement 1. Immunoblot analysis for plasmid encoded DnaA proteins in *dnaA46*

cells.

(A and B) Cells of MIT125 (MG1655 *dnaA46 tnaA*::Tn*10*) and KRC004 (MG1655 *dnaA46 tnaA*::Tn*10*

Δ*priC*::*frt-kan*) bearing pING1 (Vector), pKA234 (WT) or pFH (FH) were grown in LB mediium

including ampicillin at 30°C until the absorbance of the culture (A$_{600}$) reached 0.05, followed by

incubation at 42°C for 90 minutes. Immunoblot analysis was performed using anti-DnaA serum. Purified

DnaA was also used as a quantitative standard. MW, molecular weight markers. A representative

membrane image from three independent experiments is shown (A). Band intensities of each lane in the

image were analyzed by densitometric scanning and the amounts of DnaA were deduced using the

standard and shown as 'DnaA (ng)' (B). Means and standard deviations (SDs) are shown with each data.

1053

Figure 4-figure supplement 1. Cell growth ability of cells carrying pBR322 or pNA135 with WT

***priC* or Δ*priC*::*frt-kan*.**

Cell growth of MG1655 or KRC002 (MG1655 Δ*priC*::*frt-kan*) cells bearing pBR322 or pNA135

(pBR322 derivatives carrying *diaA*). Overnight cultures (~10^9 cells/mL) of MG1655 and KRC002 cells

carrying pBR322 or pNA135 were 10-fold serially diluted and incubated on LB agar plates containing

ampicillin for 14 h at 30–42℃. +, WT *priC*; -, Δ*priC*::*frt-kan*. The results of three independent

experiments were consistent, and one representative result is shown.

1061

Figure 7-figure supplement 1. DUE unwinding activities of DnaA mutant derivatives in the presence

and absence of PriC.

(A and B) *in vitro* DUE unwinding assays using DnaA mutant derivatives. Form I of *oriC* (pBS*oriC*; 1.6

nM) was incubated for 9 min at 30℃ with the indicated amount of DnaA or its mutant derivatives in the

presence of IHF (42 nM), followed by incubation with P1 nuclease (1.5 units) and further incubation with

AlwNI before agarose gel electrophoresis. Note that 2.6 kb and 1.0 kb fragments are observed with

restriction enzyme digestion when the ssDNA-specific P1 nuclease cleaves 3.6 kb pBS*oriC* at the DUE.

A representative image of two independent experiments is shown in black/white-inverted mode (A). The

sizes of each band are indicated at the left side of the image. Bands intensities of each lane in the gel

image were analyzed by densitometric scanning. The percentages of P1 nuclease-digested *oriC* DNA per

input DNA molecules are shown as 'DUE unwinding (%)'. Mean and data are shown (B). Abbreviations

in panel B: -, no addition of DnaA; WT, wild-type DnaA; FH, DnaA F46A H136A; VA, DnaA V211A.

(C and D) *In vitro* DUE unwinding assay with PriC. pBS*oriC* was used as described above except for

PriC addition. A representative image of two independent experiments is shown in black/white-inverted

mode (C). Data were analyzed as described above (D). n.t., not tested.

1077

Figure 8-figure supplement 1. Chromosome loci copy-number analysis of *dnaA46* mutant cells and

its derivative cells.

(A) Coverage of aligned sequence reads. MIT125, MIT125c, KRC004, and KRC004c cells were

exponentially grown in LB medium at 30℃, followed by incubation for 90 min at 40℃. Chromosomal

DNA from each sample was extracted and analyzed using whole-genome sequencing. Coverage was

calculated for every 250-bp window and plotted, with the coverage at the *oriC* region normalized to 1.0.

Dotted splines represent coverage profiles based on the combined data from two biological replicates,

shown in different colors (red and blue).

(B) Relative coverage. The coverage for each 250-bp window in $priC^+$ cells was divided by that in

$\Delta priC$ cells, and the resulting values were plotted as "relative coverage." Yellow lines indicate a value

of 1.0.

1089

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
strain, strain background (*Escherichia coli*)	BW25113 Δ*priC*::*frt-kan*	Keio collection	JW0456-KC	
strain, strain background (*Escherichia coli*)	BW25113 Δ*rep*::*frt-kan*	Keio collection	JW5604-KC	
strain, strain background (*Escherichia coli*)	MG1655 *dnaC2* *zjj18*::*cat*	Kasho and Katayama, 2013	KYA018	
strain, strain background (*Escherichia coli*)	MG1655 *dnaA46* *tnaA*::Tn10	Noguchi, *et al.*, 2016	MIT125	
strain, strain background (*Escherichia coli*)	MG1655 *rnhA*::*cat*	Noguchi, *et al.*, 2016	MIT162	
strain, strain background (*Escherichia coli*)	MG1655 Δ*diaA*::*frt-kan*	Noguchi, *et al.*, 2016	SA103	
strain, strain background (*Escherichia coli*)	MG1655 Δ*priC*::*frt-kan*	This work	KRC002	MG1655 x P1 JW0456-KC
strain, strain background (*Escherichia coli*)	MG1655 Δ*priC*::*frt*	This work	KRC003	KRC002 x pCP20

strain, strain background (*Escherichia coli*)	MIT125 Δ*priC*::*frt-kan*	This work	KRC004	MIT125 x P1 JW0456-KC
strain, strain background (*Escherichia coli*)	KYA018 Δ*priC*::*frt-kan*	This work	KRC005	KYA018 x P1 JW0456-KC
strain, strain background (*Escherichia coli*)	KRC003 Δ*diaA*::*frt-kan*	This work	KRC006	KRC003 x P1 SA103
strain, strain background (*Escherichia coli*)	MG1655 Δ*rep*::*frt-kan*	This work	FF001	MG1655 x P1 JW5604-KC
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt-kan*	Noguchi, *et al*, 2015	NY20	
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt-kan* *oriC*ΔR4 box::*Tma*DnaA box	Noguchi, *et al*, 2015	NY21	
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt-kan* *oriC*ΔMiddle-Right DOR	This work	NY20L	
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt-kan* *oriC*ΔAT-L region	This work	NY20ATL	

strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt*	Yoshida, *et al*, 2023	NY20-frt	
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt* *oriC*ΔR4 box::*Tma*DnaA box	This work	NY21-frt	NY21 x pCP20
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt* *oriC*ΔMiddle-Right DOR	This work	NY20L-frt	NY20L x pCP20
strain, strain background (*Escherichia coli*)	MG1655 *asnA*::*frt* *oriC*ΔAT-L region	This work	NY20ATL -frt	NY20ATL x pCP20
strain, strain background (*Escherichia coli*)	NY20-*frt* Δ*priC*::*frt-kan*	This work	NY20-*priC*	NY20-frt x P1 KRC002
strain, strain background (*Escherichia coli*)	NY21-*frt* Δ*priC*::*frt-kan*	This work	NY21-*priC*	NY21-frt x P1 KRC002
strain, strain background (*Escherichia coli*)	NY20L-*frt* Δ*priC*::*frt-kan*	This work	NY20L-*priC*	NY20L-frt x P1 KRC002
strain, strain background (*Escherichia coli*)	NY20ATL-*frt* Δ*priC*::*frt-kan*	This work	NY20ATL -*priC*	NY20ATL-frt x P1 KRC002

strain, strain background (*Escherichia coli*)	MIT125 Δ*rnhA*::*cat*	This work	MIT125c	MIT125 x P1 MIT162
strain, strain background (*Escherichia coli*)	KRC004 Δ*rnhA*::*cat*	This work	KRC004c	KRC004 x P1 MIT162
antibody	anti-DnaA (Rabbit polyclonal)	Kawakami, *et al.*, 2005		Immunoblot (1:3000)
antibody	Goat Anti-Rabbit IgG (H+L)-AP Conjugate	Bio-Rad	Cat. #1706518	Immunoblot (1:3000)
recombinant DNA reagent	pBR*priC* (Plasmid)	This work		pBR322 encoding *priC*
recombinant DNA reagent	pING1 (Plasmid)	Katayaam, *et al.*, 1994		Vector bearing arabinose-inducible promoter
recombinant DNA reagent	pKA234 (Plasmid)	Kubota, *et al.*, 1997		pING1 encoding *dnaA*
recombinant DNA reagent	pH136A (Plasmid)	Sakiyama, *et al.*, 2018		pKA234 *dnaA* H136A
recombinant DNA reagent	pFH (Plasmid)	This study		pH136A mutated by Quick Change site-directed mutagenesis by primers F46A 1/2
recombinant DNA reagent	pKW44-1 (Plasmid)	Ozaki, *et al.*, 2008		pKA234 *dnaA* V211A

recombinant DNA reagent	pNA135 (Plasmid)	Ishida, *et al.*, 2004		pBR322 bearing *diaA* gene
recombinant DNA reagent	pBS*oriC* (Plasmid)	Kawakami, *et al.* 2005		pBluescript bearing a 678 bp chromosome-derived region including *oriC*
recombinant DNA reagent	pET22b(+)-priC (Plasmid)	Aramaki, *et al.*, 2015		pET22b(+) bearing *priC* under the T7 promoter
recombinant DNA reagent	pTKM601 (Plasmid)	Keyamura, *et al.*, 2007		pBAD/HisB bearing *diaA*
sequence-based reagent	F46A 1	Keyamura *et al.*, 2009	PCR primer	GTACGCGCCAAACCGC GCGGTCCTTCGATTG GGTACG
sequence-based reagent	F46A 2	Keyamura *et al.*, 2009	PCR primer	CGTACCCAATCGAAGG ACCGCGCGGTTTGGCG CGTAC
sequence-based reagent	ORI_1	Kasho *et al.*, 2014	qPCR primer	CTGTGAATGATCGGTG ATC
sequence-based reagent	KW*oriC*Rev	Kasho *et al.*, 2014	qPCR primer	GTGGATAACTCTGTCA GGAAGCTTG
Sequence-based reagent	qoriK fw	Brochu *et al.*, 2014	qPCR primer	CGAGACTTCAGCGACA GTTAAG
sequence-based reagent	qoriK rev	Brochu *et al.*, 2014	qPCR primer	CCTGCGGATATTTGCG ATACA
sequence-based reagent	STM419	This work	qPCR primer	CGGACACCTTGTCTGA CCTAAG

sequence-based reagent	STM420	This work	qPCR primer	AGTGTGAAAATGACCCTCTTGC
commercial assay or kit	AP Conjugate Substrate Kit	Bio-Rad	Cat #1706432	

Figure 1

A *oriC* with AT-cluster

DUE (DNA-Unwinding Element)

DOR (DnaA-Oligomerization Region)

ssDNA binding motifs of DnaA

Left   Middle   Right

AT   L : M : R   IBR

R1   τ1 R5M τ2 I1 I2   R2   C3 C2 I3 C1 R4 :DnaA box name

B ATP-DnaA

Domain

I (1-86)   Binding to DiaA and DnaB by F46

II (87-134)   Flexible linker

III (135-374)   AAA+ motif
Arginine finger (R285)
H/B motifs (V211 and R245)
Low-affinity DnaB binding (H136)

IV (375-467)   DnaA box binding

H/B motifs
ATP
Arginine finger

C

IHF

D

DnaC
DnaB

E

Lagging strand

Leading strand

PriC

Rep

Fork restart

Figure 2

Figure 3

Host strain			*dnaA46*							
priC			+					−		
Number of cell [log10]	5	4	3	2	1	5	4	3	2	1

30℃ / 42℃
pdnaA: Vector, WT, F46A H136A

(n=4)

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9