Loss of the Inv(16) Oncogene *CBFB::MYH11* Eliminates Leukemia from the Blood and Spleen, but not the Bone Marrow

Sipra Panda[1], Yiqian Wang[1], Michelle Becker[1], Arjun Dhir[1], Calvin Lam[1], Cecilia Rivas[2], Lemlem Alemu[3], Lisa Garrett[2], Samantha Swenson[1], and R. Katherine Hyde[1]

[1]Department of Biochemistry and Molecular Biology, Fred and Pamela Buffett Cancer Center, University of Nebraska Medical Center, Omaha, NE, [2]Transgenic Mouse Core [3]Translational and Functional Genomics Branch, National Human Genome Research Institute, NIH, Bethesda, MD.

This work was supported by R00 CA148963 and R01 CA244900 to R.K.H and T32 CA009467 to support C.L. C.R., L.A., and L.G. were supported by the Intramural Research Programs of the National Human Genome Research Institute, National Institutes of Health. Core facilities used for this project are supported by state fund from the Nebraska Research Initiative and the Fred and Pamela Buffett Cancer Center's National Cancer Institute Cancer Support Grant.

Corresponding Author
R. Katherine Hyde, PhD
985879 Nebraska Medical Center
Omaha, NE 68198
kate.hyde@unmc.edu

Abstract

Inversion of chromosome 16 [inv(16)] is one of the most common chromosomal rearrangements in Acute Myeloid Leukemia (AML) and generates the fusion gene *CBFB::MYH11 (CM),* which initiates leukemogenesis. Patients with inv(16) at diagnosis invariably have the rearrangement at relapse, leading to the assumption that *CM* is required after leukemic transformation. However, this has yet to be shown experimentally. Using a knock-in mouse that allows for deletion of *CM* after leukemia development, we found that loss of the fusion gene increased apoptosis and decreased colony growth *in vitro*. Interestingly, 5-20% of the colonies had successfully deleted *CM*. To test the role of *CM in vivo*, we used an inducible shRNA knockdown (KD) construct against the fusion gene. We found that decreased *CM* expression eliminated leukemia cells from the peripheral blood and spleen, but not the bone marrow, despite these cells showing significant knockdown of *CM* at the mRNA and protein levels. Furthermore, with prolonged KD of *CM*, ~40% of mice re-established disease while maintaining KD of the fusion gene. Our work indicates that *CM* is required by leukemia cells in the spleen and blood, but that cells in the bone marrow can survive and re-establish disease independent of the fusion protein.

Introduction

Acute Myeloid Leukemia (AML) is a heterogeneous clonal malignancy arising from the immature cells in the bone marrow, resulting in a more than 20% increase in immature blast cells, reduced mature blood cell production and ultimately BM failure (1-3). It develops as a neoplastic disease, often with specific chromosomal rearrangements, including the inversion of chromosome 16(inv(16))(4, 5). Inv(16)] AML accounts for ~10% of human AML, making it one of the most common chromosomal abnormalities in AML (6-10). On the molecular level, the inv(16) generates a fusion between the transcription factor gene Core Binding Factor β (*CBFB*) and the gene for Smooth Muscle Myosin Heavy Chain (*MYH11*), generating the oncogene *CBFB::MYH11 (CM),* which encodes CBFβ::SMMHC (11, 12). Expression of *CM* induces a block in myeloid differentiation and is required but not sufficient for leukemogenesis (13). Additional cooperating mutations are required for transformation to a frank leukemia (14-16). However, it is not clear if this oncogenic fusion gene continues to be important during leukemia maintenance or if the cooperating mutations are sufficient to maintain leukemia. With current frontline treatments, inv(16) AML has a favorable prognosis due to high rates of initial response; however, ~40% of patients relapse, at which point there are limited curative options (10, 17, 18). Because *CM* is the initiating mutation in this subtype of AML, it is expressed in all leukemia cells but not in healthy cells, making it an attractive target for drug development. *CM* is assumed to be required for leukemia maintenance due to the near universal presence of Inv(16) at relapse. However, the knockdown of the fusion gene in the inv(16) AML cell line, ME-1, induces signs of differentiation but does not affect cell survival(19). Moreover, two different CBFβ-SMMHC inhibitors have been developed and in initial testing were able to decrease leukemic burden and increase survival in mouse models, but all mice eventually died of the disease (20, 21). For both compounds, it is unclear whether their limited pre-clinical effect was due to incomplete inhibition of CBFβ-SMMHC activity, or because the fusion protein is not strictly required after leukemic transformation. This issue has important implications for the further developed of targeted therapies for patients with inv(16) AML.

In this study, we address the role of *CM* in non-immortalized mouse leukemia cells. We demonstrate that *CM* loss eliminates leukemia from the peripheral blood and spleen, but not the bone marrow, indicating that loss of CBFβ::SMMHC activity is sufficient to control, but not cure, the disease.

Methods

Mice

All animal experiments were approved by the University of Nebraska Medical Center IACUC in accordance with NIH guidelines. 4-6-week-old mice of both sexes were used. To generate knock-in mice with the *flMYH11* allele, we started with a previously described pPNT vector containing the *CBFB Exon 5::MYH11* cassette, and added flanking *LoxP* sites by site directed mutagenesis (QuikChange, Agilent) (22). Mouse ES cells were electroporated, selected, screened for homologous recombination and injected into blastocysts, as described previously (22, 23). To confirm the *Cbfb^flMYH11* allele functions as predicted, chimeric mice were bred to mice expressing *Cre* from the β-Actin promoter (*ActB Cre^+*), purchased from Jackson Laboratory (Bar Harbor, ME) and embryos harvested at embryonic day 12.5(24). To induce leukemia, chimeric *Cbfb^+/flMYH11*mice were treated with N-ethyl-N-nitrosourea (ENU) and monitored for leukemia development (13, 25, 26). Leukemia cells were transplanted into sub-lethally irradiated congenic recipient mice as described previously (27, 28). Knockdown studies were performed using leukemia cells from *Mx1-Cre, Cbfb^+/56M* mice, which allows for inducible expression of *CM* (29). To induce shRNA expression, mice were given 5% sucrose supplemented drinking water with or without doxycycline at 2mg/ml. Mice were sacrificed on the indicated day of treatment and tissue collected.

97 qRT-PCR Analysis
98 RNA was extracted using Trizol method and cDNA synthesized using Luna RT Supermix (New
99 England Biolabs, MA) and amplified using Luna Universal qPCR kit (New England Biolabs, MA).
100 Primer sequences available upon request.
101
Western Blot Analysis
Cell lysates were prepared, subjected to SDS-PAGE and transferred to PVDF, as described
previously (27, 28). Membranes were blocked in 5% milk and then probed using primary
antibodies against MYH11 (Novus Biologicals, CO, and Origene, MD), GAPDH (Ambion, Texas),
β-Actin (Santa Cruz Biotechnology, TX), MCL-1(Cell Signaling Technology, MA), BCL-2 (Cell
Signaling Technology, MA), and GFP (Invitrogen, CA) and with secondary HRP-linked antibodies
anti-rabbit or anti-mouse. Protein bands were visualized with SuperSignal West Pico PLUS
(Thermo Scientific, MA). Protein levels were quantified using ImageJ.
110
Flow Cytometry Analysis
Cells were stained with fluorophore coupled antibodies against B220 (RA3-6B2, Becton
Dickinson, NJ), CD3 (17A2, BD Biosciences, NJ), Mac-1/CD11b (M1/70, BD Biosciences, NJ),
GR-1 (RM4-5, BD Biosciences, NJ), CKIT (53-6.7, BD Biosciences, NJ), Csf2rb/CD131 (JORO50,
BD Biosciences, NJ), IL1RL1 (U29-93, BD Biosciences, NJ), CD41 (MWReg30, BD Biosciences,
NJ), CD123 (5B11, BD Biosciences, NJ) and analyzed by flow cytometry (BD LSR II). Apoptosis
was assayed using Annexin V (BD Pharmigen, NJ). Proliferation was determined using the BrdU
Flow Kit (BD Pharmigen, NJ). Mice were injected with BrdU at dose of 200mg/kg body weight.
After 1 hr, mice were sacrificed and tissue harvested. All flow cytometry data was analyzed using
FlowJo 10.0 (BD, OR).
121
Viral Transduction
To excise the *Cbfb* ^{flMYH11} allele, lentiviral plasmids co-expressing *Cre Recombinase* and either
GFP or DsRED were used. To knockdown *Cbfb-MYH11*, a lentiviral plasmid expressing GFP and
an shRNA against *MYH11* was kindly provided by Dr. J. Martens, Nijmegen, NL(19). Lentiviral
plasmids were transfected into HEK293T cells. Mouse leukemia cells were incubated with viral
supernatant, cytokines and 8ug/ml polybrene for transduction. Transduction was performed by
spinfection centrifugation for 90 minutes at 2000 rpm, 30 °C, followed by a 6-hour incubation and
a second spinfection. Transduced cells were sorted for GFP or dsREd 48hrs post transduction.
130
131 Colony Assay
132 Transduced cells were sorted, and equal numbers were suspended in 35mm dishes in Methocult
133 3434 (StemCell Technologies, Vancouver, BC) in triplicate, according to the manufacturer's
134 protocol. Colonies were counted after 14 days in culture.
135
136 Statistical Analysis
137 Statistical significance was analyzed using either Student's T-Test or ANOVA, as appropriate
138 (GraphPad Prism, version 10, Dotmatics, CA) with p-values ≤0.05 deemed significant.
139
140 **Results**
141
Generation and characterization of *Cbfb*^{floxMYH11} leukemia cells
To test the role of *Cbfb::MYH11* in leukemia maintenance, we generated and characterized a new
knock-in mouse model, where the *Cbfb exon5::MYH11* fusion exon is flanked by *LoxP* sites and
knocked into the endogenous *Cbfb* locus (*Cbfb*$^{+/flMYH11}$) (Fig. 1A). We identified embryonic stem
cell (ES) clones with the knock-in allele by southern blot and confirmed CBFβ::SMMHC
expression by western blot (Supplementary Fig. S1A, B). Three independent *Cbfb*$^{+/ floxMYH11}$ ES

clones were used to generate chimeric mice. To test that the *Cbfb* ^{flMYH11} allele can be fully excised, chimeric mice were crossed with mice expressing *Cre Recombinase* from the *Actin B* promoter (*ActB Cre*), which should allow for excision of the fusion gene in all cells of an embryo. We harvested embryos at day 12.5 in development. In *Cbfb* $^{+/flMYH11}$ embryos lacking the *ActB Cre* transgene, we observed central nervous system hemorrhaging, indicating expression of *Cbfb::MYH11*, as in previous knock-in models (Fig. 1B) (22). Importantly, this phenotype was not observed in *Cbfb* $^{+/flMYH11}$; *ActB Cre*+, indicating loss of the fusion gene in the presence of *Cre Recombinase*. PCR using primers specific for the unexcised and excised alleles confirmed that the *Cbfb* ^{flMYH11} was efficiently excised in *Cbfb* $^{+/flMYH11}$; *ActB Cre*+ mice. To generate leukemia cells expressing the *Cbfb* ^{flMYH11} allele, chimeric mice were treated with N-ethyl-N-nitrosourea and monitored for disease onset. Chimeric *Cbfb* $^{+/flMYH11}$ mice from each of the three ES clones developed transplantable leukemia with a similar immunophenotype and histological appearance as previous *CM knock-in* mouse models (Fig. 1C, D; Supplementary Fig. S1C, D) (29, 30). These results indicate that *Cbfb* $^{+/flMYH11}$ mice develop a frank leukemia with a similar phenotype to previous models and that the fusion gene can be deleted by *Cre Recombinase*.

Loss of *Cbfb::MYH11 (CM)* impairs the growth and survival of leukemia cells *in vitro*, but rare *Cbfb::MYH11* deleted cells form colonies

To test the effect of *CM* deletion *in vitro*, we transduced *Cbfb* $^{+/flMYH11}$ leukemia cells with lentivirus expressing *Cre* and GFP from an IRES (Fig. 2A). Transduced cells were sorted, and *CM* excision was confirmed by qPCR and western blot (Fig. 2B-C). We found that *CM* deletion caused a statistically significant increase in Annexin V, indicating induction of apoptosis (Fig. 1D). To test the effect of *CM* deletion on colony forming ability, we plated equal numbers of control and *Cre* transduced *Cbfb* $^{+/flMYH11}$ leukemia cells in methylcellulose. After 14 days, colonies were counted, resuspended as single cells, and equal numbers replated in methylcellulose. At both the first and 2nd plating, we found that *Cbfb* $^{+/flMYH11}$ cells transduced with *Cre* generated significantly fewer colonies (Fig. 1E). To determine if the colonies that were able to grow from *Cre*-infected cells had successfully deleted *CM*, individual colonies were picked and qPCR for the *Cbfb* ^{flMYH11} allele was performed. We found that most colonies retained the unexcised *CM* allele; however, 5-20% of the colonies showed successful deletion of *CM* (Supplementary Fig. S2A, B). These findings imply that the majority of leukemia cells require continued expression of the fusion gene after leukemia initiation, but that some cells may be able to survive without *CM*.

Knockdown of *Cbfb::MYH11* impairs leukemia survival *in vivo*

Due to the high rate of apoptosis seen *in vitro* with genetic deletion of *Cbfb* ^{flMYH11} allele, we used a different mouse model and a shRNA knockdown (KD) construct to test the requirement for *Cbfb::MYH11* in leukemia maintenance *in vivo*. We generated leukemia cells from mice with the *Cbfb* ^{56M} allele, which allows inducible expression of *Cbfb::MYH11* and the *Mx1-Cre* transgene, which is inducibly expressed in nearly all hematopoietic cells (29). Leukemia cells from *Cbfb* $^{+/56M}$, *Mx1-Cre*+ were transduced with a doxycycline (Dox) inducible shRNA against *MYH11* (*shMYH11*) that also expresses GFP constitutively (19). To test the feasibility of this approach *in vitro*, we sorted GFP+ cells and cultured them with Dox for 48 hours. After 48 hrs, *CM* expression was determined by qRT-PCR and western blot. We observed that KD resulted in 70-95% decrease in expression at mRNA levels and 51-98% decrease of the fusion protein *in vitro* (Supplementary Fig. S3A-C). We also found that *CM* KD induced a similar phenotype to the genetic deletion of the fusion gene, with increased apoptosis and decreased colony formation *in vitro* (Supplementary Fig. S3D, E).

To determine the effect of decreased *CM* expression *in vivo*, we transplanted *shMYH11*
transduced cells into congenic mice, allowing them to engraft. Once mice showed at least 20%
GFP$^+$ leukemia cells in the peripheral blood, mice were treated with control (5% sucrose) or Dox
(5% sucrose and Dox) drinking water (Supplementary Fig. S4A). Mice were sacrificed after 4, 7,
14, 21, and 28 days, and the presence of GFP$^+$ leukemia cells was determined. We observed that
CM KD induced a significant decrease in GFP$^+$ leukemic cells in the peripheral blood and spleen
starting at day 7, with near complete elimination by day 14 (Fig. 3A, 3B). In addition, we observed
a notable decrease in spleen size with *CM* KD (Supplementary Fig. S4B, C). In the bone marrow,
there was a significant decrease in leukemia cells starting at day 14 of Dox treatment. However,
20-30% of leukemia cells persisted through day 28 (Fig. 3C). To determine if the persistence of
leukemia cells in the bone marrow was due to failure of Dox to penetrate the tissue and adequately
KD the fusion gene, we sorted GFP$^+$ leukemia cells from the bone marrow of control and Dox
treated mice at days 7 and 21. qRT-PCR and western blot of bone marrow GFP$^+$ cells showed
significant *CM* KD at both time points (Fig. 3D-F). These findings indicate that cells with reduced
CM can survive in the bone marrow, but not the peripheral blood and spleen.
212
To investigate potential mechanisms for the persistence of *CM* KD cells in the bone marrow, we
analyzed apoptosis and proliferation in the leukemic cells of Dox and control treated mice at Day
21. We found a significantly lower percentage of leukemia cells undergoing apoptosis in the bone
marrow of Dox treated mice, as compared to control (Fig. 4A). To determine if the decrease in
apoptosis is related to expression of anti-apoptotic proteins, we sorted leukemia cells from the
spleen and bone marrow of control and Dox treated mice and performed western blots. We found
that KD of *CM* induced decreased BCL-2 protein in the spleen but caused no change in MCL-1.
In leukemic cells from the bone marrow of both control and Dox treated mice, BCL-2 was barely
detectable. In contrast, MCL-1 expression was higher in leukemia cells from the bone marrow of
both control and dox treated mice, as compared to cells from the spleen (Fig. 4B, C). These
findings imply that MCL-1, but not BCL-2, may contribute to the survival of CM knockdown cells
in the bone marrow.
225
We used BrdU incorporation to determine if a difference in proliferation also contributed to the
persistence of *CM* KD cells in the bone marrow. Leukemia cells from both the spleen and bone
marrow of day 21 Dox treated mice had significantly lower proliferation compared to the control
(Fig. 4D). To test if the decrease in proliferation was due to a change in the differentiation state
of the leukemia cells, we examined the expression of Mac-1 and Gr-1, markers of mature myeloid
cells. In the spleen of Dox treated mice, we observed an increase in Mac1$^+$, GR-1$^+$ cells,
consistent with differentiation. In contrast, there was no difference in the expression of Mac1 or
GR-1 on the leukemia cells in the bone marrow (Fig. 4E). To test whether the *CM* KD cells in the
bone marrow were instead enriched for Leukemia Stem Cell (LSC) markers, we examined
expression of KIT, CD59a, IL1RL1, CD123. Work by us and others previously showed that KIT,
CD59a and IL1RL1 are enriched on the LSC population driven by *CM* expression, and CD123 is
an established LSC marker in human AML (27, 31, 32). We observed a small yet statistically
significant increase in CD123 and decrease in IL1RL1 in the leukemia cells in the bone marrow
of Dox treated mice (Fig. 4F). These results indicate that the decreased proliferation observed in
the *CM* KD cells in the bone marrow is likely not due to differentiation to a post-mitotic mature
myeloid cell or enrichment for quiescent LSCs.
242
The persistence of a *CM* KD leukemic population in the bone marrow raises the possibility that
these cells may be able to cause relapse. To test this possibility, we maintained mice transplanted
with *shMYH11* transduced leukemia cells on Dox or control drinking water until they showed signs
of lethal disease. Five out of 12 Dox treated mice showed re-emergence of GFP$^+$ cells in the
blood (Fig. 5A). To determine if the relapse cells in the blood expressed *CM*, we sorted for GFP

and performed qRT-PCR (Fig. 5B) and western blot (Fig. 5C, D). We found that relapsed samples
had reduced expression of *CM* at both the mRNA and protein level, as compared to control treated
mice. In addition, we found that the immunophenotype of the *CM* KD relapse leukemia cells was
similar to leukemia cells from control mice (Supplementary Fig. S5A and B). These results imply
that despite the reduction in *CM* expression, relapse cells acquired additional mutations that
promote survival outside the bone marrow microenvironment, and re-establishment of fatal
disease.
255
Discussion
CBFβ::SMMHC is a tempting target for drug development, as it expressed in all leukemia cells in
patients with inv(16) AML, but not in healthy cells. However, work in the immortalized cell line ME-
1 raises the possibility that the fusion protein may not be strictly required after leukemic
transformation (19). To our knowledge, this is the first study to test this possibility using non-
immortalized leukemia cells. We found that genetic deletion of *CM* induced significant apoptosis
and impaired colony growth. However, a small number of colonies were able to grow after *CM*
deletion, implying that there is a population of leukemia cells that don't strictly require the fusion
gene. *In vivo*, we found that KD of *CM* induced a near complete elimination of leukemic cells in
the blood and spleen. Surprisingly, the leukemic burden in the bone marrow did not decrease
below 20-30% despite almost complete loss of CBFβ-SMMHC protein, implying that the bone
marrow microenvironment can sustain leukemia cells despite the loss of *CM* expression.
268
Our results concerning the role of CBFβ::SMMHC during leukemia maintenance are strikingly
similar to what has been demonstrated with the BCR::ABL fusion protein in Chronic Myeloid
Leukemia (CML). Like CBFβ::SMMHC, BCR::ABL is the product of a chromosomal
rearrangement characteristic of the disease, and its expression is the initiating event.
Pharmacological inhibitors of BCR::ABL, such as imatinib, are remarkably successful at inducing
long-term, stable cytogenetic remission (33, 34). However, these kinase inhibitors generally do
not eliminate the LSC population in the bone marrow. Similarly, near complete knockdown of
BCR::ABL expression reduces the LSC population but does not eliminate it, and the remaining
LSCs are able to re-establish disease when BCR::ABL activity is restored (35). In our model, we
also observed a population of LSC-like cells that can survive in the bone marrow and re-establish
disease, despite near complete loss of the CBFβ::SMMHC protein. In our case, relapse did not
require re-expression of the fusion protein. However, given the period of latency before relapse,
it is likely that additional mutations developed to compensate for the lack of CBFβ::SMMHC
activity.
283
Our observation that *CM* knockdown cells can persist only in the bone marrow implies that there
are factors present in the microenvironment critical for their survival. This is consistent with other
work demonstrating that the bone marrow microenvironment is unique in promoting leukemia cell
survival and preventing apoptosis (35-37). In fact, we observed that *CM* KD cells in the bone
marrow, but not the spleen, had decreased rates of apoptosis as compared to control. Thus, to
identify potential therapeutically targetable factors promoting the survival of *CM KD* cells, we
performed westerns for the anti-apoptotic proteins BCL-2 and MCL-1 (38, 39). We observed that
both control and *CM KD* had had robust expression of MCL-1 but nearly undetectable levels of
BCL-2. This finding is consistent with recent reports indicating that the BCL-2 inhibitor venetoclax
is less effective in patients with French-American-British (FAB) M4 and M5 monocytic
differentiation AML, which includes inv(16) AML (40).
295
Collectively, this study supports the development of inhibitors targeting the *CM* fusion protein for
the treatment of inv(16) AML. To date, drugs targeting CBFβ-SMMHC show the feasibility of such
an approach, even if the current inhibitors are not suitable for use in humans (20, 21). Our data

shows that as the next generation of such inhibitors are developed, they would likely be effective
at managing leukemia burden, and when used in combination with other drugs such as MCL-1
inhibitors, have the potential to cure the disease.
302

303   **Acknowledgments**
304   The authors would like to thank Paul Liu, M.D, Ph.D (NHGRI/NIH) for his support and
305   encouragement of this project, as well as the members of the UNMC flow cytometry core facility
306   for their expertise.
307
308   This work was supported by R00 CA148963 and R01 CA244900 to R.K.H and T32 CA009467 to
309   support C.L. C.R., L.A., and L.G. were supported by the Intramural Research Programs of the
310   National Human Genome Research Institute, National Institutes of Health. Core facilities used for
311   this project are supported by state fund from the Nebraska Research Initiative and the Fred and
312   Pamela Buffett Cancer Center's National Cancer Institute Cancer Support Grant.
313

314   **Author Contributions**
315   S.P. designed and performed experiments, analyzed data and wrote the paper. Y.W. designed
316   and performed experiments and analyzed data. M.B., C.L., A.D., and S.S. performed
317   experiments. C.R., L.A., and L.G. generated, bred, and mutagenized $Cbfb^{+/floxMYH11}$ knock-in mice.
318   R.K.H designed experiments, analyzed data and wrote the paper.
319
320   **Disclosure of Conflicts of Interest**
321   The authors have no conflicts of interest to disclose.
322
323

References

1.   Akinduro O, Weber TS, Ang H, Haltalli MLR, Ruivo N, Duarte D, et al. Proliferation

dynamics of acute myeloid leukaemia and haematopoietic progenitors competing for bone

marrow space. Nature Communications. 2018;9(1):519.

2.   Thomas D, Majeti R. Biology and relevance of human acute myeloid leukemia stem cells.

Blood, The Journal of the American Society of Hematology. 2017;129(12):1577-85.

3.   Lane SW, Scadden DT, Gilliland DG. The leukemic stem cell niche: current concepts and

therapeutic opportunities. Blood, The Journal of the American Society of Hematology.

2009;114(6):1150-7.

4.   Kumar CC. Genetic abnormalities and challenges in the treatment of acute myeloid

leukemia. Genes Cancer. 2011;2(2):95-107.

5.   Speck NA, Gilliland DG. Core-binding factors in haematopoiesis and leukaemia. Nature

Reviews Cancer. 2002;2(7):502-13.

6.   Khoury JD, Solary E, Abla O, Akkari Y, Alaggio R, Apperley JF, et al. The 5th edition of

the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and

Histiocytic/Dendritic Neoplasms. Leukemia. 2022;36(7):1703-19.

7.   Balgobind BV, Hollink IH, Arentsen-Peters ST, Zimmermann M, Harbott J, Beverloo HB,

et al. Integrative analysis of type-I and type-II aberrations underscores the genetic heterogeneity

of pediatric acute myeloid leukemia. Haematologica. 2011;96(10):1478-87.

8.   Kihara R, Nagata Y, Kiyoi H, Kato T, Yamamoto E, Suzuki K, et al. Comprehensive

analysis of genetic alterations and their prognostic impacts in adult acute myeloid leukemia

patients. Leukemia. 2014;28(8):1586-95.

9.   Swerdlow S. World Health Organization classification of tumours of the haematopoietic

and lymphoid tissues. Eds. SH Swerdlow, E. Campo, NL Harris, ES Jaffa, SA Pileri, H. Stein. J

Thiele 4th ed Postgraduate Haematology. 2017:986-8.

10.   Mancini M, Cedrone M, Diverio D, Emanuel B, Stul M, Vranckx H, et al. Use of dual-color

interphase FISH for the detection of inv(16) in acute myeloid leukemia at diagnosis, relapse and

during follow-up: a study of 23 patients. Leukemia. 2000;14(3):364-8.

11.   Liu P, Tarlé SA, Hajra A, Claxton DF, Marlton P, Freedman M, et al. Fusion between

transcription factor CBF beta/PEBP2 beta and a myosin heavy chain in acute myeloid leukemia.

Science. 1993;261(5124):1041-4.

12.   Liu PP, Wijmenga C, Hajra A, Blake TB, Kelley CA, Adelstein RS, et al. Identification of

the chimeric protein product of the CBFB-MYH11 fusion gene in inv(16) leukemia cells. Genes

Chromosomes Cancer. 1996;16(2):77-87.

13.   Castilla LH, Garrett L, Adya N, Orlic D, Dutra A, Anderson S, et al. The fusion gene Cbfb-

MYH11 blocks myeloid differentiation and predisposes mice to acute myelomonocytic leukaemia

[letter]. Nat Genet. 1999;23(2):144-6.

14.   Castilla LH, Perrat P, Martinez NJ, Landrette SF, Keys R, Oikemus S, et al. Identification

of genes that synergize with Cbfb-MYH11 in the pathogenesis of acute myeloid leukemia.

Proceedings of the National Academy of Sciences of the United States of America.

2004;101(14):4924-9.

15.   Sood R, Hansen NF, Donovan FX, Carrington B, Bucci D, Maskeri B, et al. Somatic

mutational landscape of AML with inv(16) or t(8;21) identifies patterns of clonal evolution in

relapse leukemia. Leukemia. 2016;30(2):501-4.

16.   Duployez N, Marceau-Renaut A, Boissel N, Petit A, Bucci M, Geffroy S, et al.

Comprehensive mutational profiling of core binding factor acute myeloid leukemia. Blood.

2016;127(20):2451-9.

17.   Marcucci G, Mrozek K, Ruppert AS, Maharry K, Kolitz JE, Moore JO, et al. Prognostic

factors and outcome of core binding factor acute myeloid leukemia patients with t(8;21) differ from

those of patients with inv(16): a Cancer and Leukemia Group B study. Journal of clinical oncology

: official journal of the American Society of Clinical Oncology. 2005;23(24):5705-17.

18.   Appelbaum FR, Kopecky KJ, Tallman MS, Slovak ML, Gundacker HM, Kim HT, et al. The

clinical spectrum of adult acute myeloid leukaemia associated with core binding factor

translocations. British journal of haematology. 2006;135(2):165-73.

19.   Mandoli A, Singh AA, Jansen PW, Wierenga AT, Riahi H, Franci G, et al. CBFB-

MYH11/RUNX1 together with a compendium of hematopoietic regulators, chromatin modifiers

and basal transcription factors occupies self-renewal genes in inv(16) acute myeloid leukemia.

Leukemia. 2013.

20.   Cunningham L, Finckbeiner S, Hyde RK, Southall N, Marugan J, Yedavalli VR, et al.

Identification of benzodiazepine Ro5-3335 as an inhibitor of CBF leukemia through quantitative

high throughput screen against RUNX1-CBFbeta interaction. Proceedings of the National

Academy of Sciences of the United States of America. 2012;109(36):14592-7.

21.   Illendula A, Gilmour J, Grembecka J, Tirumala VS, Boulton A, Kuntimaddi A, et al. Small

Molecule Inhibitor of CBFbeta-RUNX Binding for RUNX Transcription Factor Driven Cancers.

EBioMedicine. 2016;8:117-31.

22.   Castilla LH, Wijmenga C, Wang Q, Stacy T, Speck NA, Eckhaus M, et al. Failure of

embryonic hematopoiesis and lethal hemorrhages in mouse embryos heterozygous for a

knocked-in leukemia gene CBFB-MYH11. Cell. 1996;87(4):687-96.

23.   Kamikubo Y, Zhao L, Wunderlich M, Corpora T, Hyde RK, Paul TA, et al. Accelerated

leukemogenesis by truncated CBF beta-SMMHC defective in high-affinity binding with RUNX1.

Cancer Cell. 2010;17(5):455-68.

24.   Lewandoski M, Meyers EN, Martin GR. Analysis of Fgf8 gene function in vertebrate

development. Cold Spring Harb Symp Quant Biol. 1997;62:159-68.

25.   Kamikubo Y, Hyde RK, Zhao L, Alemu L, Rivas C, Garrett LJ, et al. The C-terminus of

CBFbeta-SMMHC is required to induce embryonic hematopoietic defects and leukemogenesis.

Blood. 2013;121(4):638-42.

26.   Hyde RK, Zhao L, Alemu L, Liu PP. Runx1 is required for hematopoietic defects and

leukemogenesis in Cbfb-MYH11 knock-in mice. Leukemia. 2015;29(8):1771-8.

27.   Wang Y, Richter L, Becker M, Amador C, Hyde RK. IL1RL1 is dynamically expressed on

Cbfb-MYH11+ leukemia stem cells and promotes cell survival. Scientific reports. 2019;9(1):1729.

28.   Richter LE, Wang Y, Becker ME, Coburn RA, Williams JT, Amador C, et al. HDAC1 is a

Required Cofactor of CBFbeta-SMMHC and a Potential Therapeutic Target in Inversion 16 Acute

Myeloid Leukemia. Molecular cancer research : MCR. 2019.

29.   Kuo YH, Landrette SF, Heilman SA, Perrat PN, Garrett L, Liu PP, et al. Cbf beta-SMMHC

induces distinct abnormal myeloid progenitors able to develop acute myeloid leukemia. Cancer

Cell. 2006;9(1):57-68.

30.   Hyde RK, Kamikubo Y, Anderson S, Kirby M, Alemu L, Zhao L, et al. Cbfb/Runx1

repression-independent blockage of differentiation and accumulation of Csf2rb-expressing cells

by Cbfb-MYH11. Blood. 2010;115(7):1433-43.

31.   Diemer JL, Yu K, Kelly M, Zhen T, Anderson S, Lopez G, et al. Characterization of

leukemia progression in the Cbfb-MYH11 knockin mice by single cell RNA sequencing. Leukemia.

2023;37(7):1549-53.

32.   Arnone M, Konantz M, Hanns P, Paczulla Stanger AM, Bertels S, Godavarthy PS, et al.

Acute Myeloid Leukemia Stem Cells: The Challenges of Phenotypic Heterogeneity. Cancers

(Basel). 2020;12(12).

33.   Druker BJ, Guilhot F, O'Brien SG, Gathmann I, Kantarjian H, Gattermann N, et al. Five-

year follow-up of patients receiving imatinib for chronic myeloid leukemia. N Engl J Med.

2006;355(23):2408-17.

34.   Hughes TP, Hochhaus A, Branford S, Muller MC, Kaeda JS, Foroni L, et al. Long-term

prognostic significance of early molecular response to imatinib in newly diagnosed chronic

myeloid leukemia: an analysis from the International Randomized Study of Interferon and STI571

(IRIS). Blood. 2010;116(19):3758-65.

35.   Hamilton A, Helgason GV, Schemionek M, Zhang B, Myssina S, Allan EK, et al. Chronic

myeloid leukemia stem cells are not dependent on Bcr-Abl kinase activity for their survival. Blood.

2012;119(6):1501-10.

36.   Wang A, Zhong H. Roles of the bone marrow niche in hematopoiesis, leukemogenesis,

and chemotherapy resistance in acute myeloid leukemia. Hematology. 2018;23(10):729-39.

37.   Weinberg OK, Hasserjian RP, Baraban E, Ok CY, Geyer JT, Philip JKSS, et al. Clinical,

immunophenotypic, and genomic findings of acute undifferentiated leukemia and comparison to

acute myeloid leukemia with minimal differentiation: a study from the bone marrow pathology

group. Modern Pathology. 2019;32(9):1373-85.

38.   Fowler-Shorten DJ, Hellmich C, Markham M, Bowles KM, Rushworth SA. BCL-2 inhibition

in   haematological   malignancies: Clinical   application   and   complications.   Blood   Rev.

2024;65:101195.

39.   Griffioen MS, de Leeuw DC, Janssen J, Smit L. Targeting Acute Myeloid Leukemia with

Venetoclax;   Biomarkers   for   Sensitivity   and   Rationale   for   Venetoclax-Based   Combination

Therapies. Cancers (Basel). 2022;14(14).

40.   Bruserud O, Selheim F, Hernandez-Valladares M, Reikvam H. Monocytic Differentiation

in Acute Myeloid Leukemia Cells: Diagnostic Criteria, Biological Heterogeneity, Mitochondrial

Metabolism, Resistance to and Induction by Targeted Therapies. Int J Mol Sci. 2024;25(12).

Figure Legends

Figure 1: Generation of the novel model and *in vitro* analysis. **A.** Schematic representation of the ES-cell targeting strategy. The probe (0.2C) and restriction enzyme sites (NcoI) used for screening are indicated. The black arrows represent PCR primers for the excised allele, and the purple arrows represent primers for the unexcised allele. **B.** Pictures of representative embryos harvested at embryonic day 12.5 generated by crossing *Cbfb+/floxMYH11* chimeras with transgenic mice expressing *Cre recombinase* (Cre) from the *β–Actin* promoter (*ActB Cre*). The right panel shows a representative PCR for the excised and unexcised *Cbfb+/floxMYH11* allele in DNA from embryos of the indicated genotype. **C.** Kaplan-Meyer survival curve of recipient mice transplanted with *Cbfb+/flMYH11* leukemic cells from the chimeric donor mice. **D.** Representative Wright-Giemsa-stained peripheral blood smear and hematoxylin and eosin-stained spleen and liver section from a leukemic *Cbfb+/flMYH11* recipient mouse.

Figure 2: Deletion of *Cbfb::MYH11* induces apoptosis and decreased colony formation *in vitro* **A.** Schematic representation of the experimental design to test the role of *Cbfb::MYH11* in leukemia maintenance *in vitro*. **B.** Quantitative Reverse Transcription PCR (qRT-PCR) analysis for the unexcised and excised *Cbfb+/floxMYH11* alleles in leukemia cells transduced with a lentivirus expressing Cre or Control (CTRL) and sorted for GFP expression. **C.** Western blot analysis of CBFβ::SMMHC in *Cbfb+/flMYH11* leukemia cells infected with Cre and CTRL virus. **D.** Graph of the percent (%) Annexin V+ Cre and CTRL transduced *CbfbflMYH11* leukemia cells from 3 independent mice. **E.** Bar graph showing the relative number of colonies from equal numbers of Cre and control transduced *Cbfb+/flMYH11* leukemia cells from 3 independent mice, each plated in triplicate plated in methylcellulose. Colonies were scored on day 14. N=3. ***p <0.001 as compared to CTRL.

Figure 3: *CM* knockdown decreases leukemia burden in *vivo*. **A.** Line graph showing percent (%) GFP+ *shCM* transduced leukemia cells (left) and representative flow cytometry plots (right) in the peripheral blood (PB), **B.** spleen (SP) and **C.** bone marrow (BM) of mice treated with Dox or control for 7, 14, 21, or 28 days. **D.** Bar graph showing relative expression of *CM* in sorted GFP+ leukemic cells from the BM of mice treated with Dox or control for 7 and **E.** 2 days. **F.** Representative western blot (left) and graph of relative CBFB::MYH11 protein expression in sorted GFP+ cells from the BM of mice treated with either control or Dox for 7 days. For all experiments N≥ 3 biological replicates, ns = not significant, *p=<0.05.

Figure 4: Leukemia cells in the bone marrow with decreased *CM* have a similar immunophenotype to control leukemia cells.
A. Bar graph showing the percentage (%) of Annexin V+, GFP+ from the spleen (SP) and bone marrow (BM) of mice treated with Dox or control for 21 days. **B.** Representative western blot for the indicated proteins in leukemic cells sorted from SP and BM after 7 days of treatment with control or Dox **C.** Graph of the relative protein expression from the westerns blots in B. **D.** Graph of the % of BrdU+, GFP+ leukemia cells harvested from the SP and BM of mice treated with Dox or control for 21 days. **E.** Bar graph showing % of GFP+ cells expressing the mature myeloid markers Mac-1 and Gr-1. Leukemic cells were harvested from the BM of mice treated as in A. **F.** Bar graph showing the % of GFP+ cells expressing the indicated leukemia stem cell markers in the BM from of mice treated for 21 days with Dox or vehicle. For all experiments N≥ 3 biological replicates, *p=<0.05, ** = p<0.01, *** = p<0.001

Figure 5: The remaining *CM* knockdown cells in the bone marrow can give rise to relapse.
A. Line graph showing the percentage (%) of GFP+, *shCM* transduced cells in peripheral blood (PB) of mice treated for up to 42 days with either control or Dox**. B.** Graph of relative expression of *CM* mRNA in GFP+ sorted leukemia cells from dox treated mice after relapse. **C.**

497   Representative western blot of the indicated proteins in sorted GFP$^+$ cells harvested from control
498   and Dox treated mice. **D.** Graph of relative CBFβ::SMMHC protein expression in samples treated
499   as in C. N≥ 3 biological replicates, *p=<0.05
500
501

Figure 1

Figure 2

A.

Selection of knockin clones

Generate leukemic mice

Cre & GFP

Test requirement for *CM in vitro*

B.

C.

D.

E.

Figure 3

Figure 4

Figure 5