Tissue Specific Requirement for the Inv(16) Oncogene *CBFB::MYH11* in Acute

Myeloid Leukemia

Sipra Panda[1,2], Yiqian Wang[1,2], Venkatasai Rahul Dogiparthi[2,3], Michelle Becker[1,2], Arjun Dhir[1,2],

Calvin Lam[1,2], Cecilia Rivas[4], Lemlem Alemu[5], Lisa Garrett[4], Peng Xiao[3], Samantha A.

Swenson[1,2], Kyle J. Hewitt[2,3], and R. Katherine Hyde[1,2]

[1]Department of Biochemistry and Molecular Biology, [2]Fred and Pamela Buffett Cancer Center

[3]Department of Genetics, Cell Biology and Anatomy, University of Nebraska Medical

Center, Omaha, NE, [4]Transgenic Mouse Core, [5]Translational and Functional Genomics Branch,

National Human Genome Research Institute, NIH, Bethesda, MD.

Corresponding Author

R. Katherine Hyde, PhD

985879 Nebraska Medical Center

Omaha, NE 68198

kate.hyde@unmc.edu

Running Title: Tissue Specific Requirement for *CBFB::MYH11* in AML

This work was supported by R00 CA148963 and R01 CA244900 to R.K.H and T32 CA009467 to

support C.L. C.R., L.A., and L.G. were supported by the Intramural Research Programs of the

National Human Genome Research Institute, National Institutes of Health. Core facilities used for

this project are supported by a state fund from the Nebraska Research Initiative, the Fred and

Pamela Buffett Cancer Center's National Cancer Institute Cancer Support Grant (P30

CA036727), and the Nebraska Center for Molecular Target Discovery and Development COBRE

(P20 GM121316).

Abstract

Inversion of chromosome 16 [inv(16)] generates the fusion gene *CBFB::MYH11 (CM)* and is one

of the most common chromosomal rearrangements in Acute Myeloid Leukemia (AML).

Expression of *CM* is required for leukemia initiation. Patients with inv(16) at diagnosis invariably

have the rearrangement at relapse, leading to the assumption that *CM* is also required after

leukemic transformation. However, a role for *CM* in leukemia maintenance has yet to be shown

experimentally. To address this, we used an inducible *CM* knockdown (KD) mouse model and

found that decreased *CM* eliminated leukemia cells from the peripheral blood and spleen, but not

the bone marrow, despite all populations exhibiting significantly decreased *CM* mRNA and

protein. The surviving *CM* KD cells in the bone marrow showed decreased apoptosis and

proliferation, and increased expression of autophagy related genes. Surprisingly, with prolonged

KD of *CM*, ~40% of mice re-established disease despite maintaining decreased *CM*. Our work

indicates that *CM* is required for leukemia survival in the spleen and peripheral blood, but in the

bone marrow *CM* KD leukemia cells can survive and re-establish disease independent of the

fusion protein. These findings imply that targeting *CM* alone has potential to reduce leukemic

burden but not cure the disease.

Introduction

Acute Myeloid Leukemia (AML) is a heterogeneous clonal malignancy arising from the hematopoietic stem and progenitor cells of the bone marrow, as the result of a block in differentiation that gives rise to a more than 20% increase in immature blast cells, reduced mature blood cell production and ultimately BM failure (1-3). AML develops as a neoplastic disease, often with specific chromosomal rearrangements, including the inversion of chromosome 16[inv(16)] (4, 5). These rearrangements are considered the initiating mutation, and with the acquisition of additional cooperating mutations, these abnormal myeloid cells can transform into a full-blown, frank leukemia (6-8). Inv(16) accounts for ~10% of all AMLs, making it one of the most common chromosomal abnormalities in AML (9-12). At the molecular level, inv(16) generates a fusion between the genes encoding the transcription factor Core Binding Factor β (*CBFB*) and the contractile protein Smooth Muscle Myosin Heavy Chain (*MYH11*). The resulting oncogene, *CBFB::MYH11 (CM),* encodes the protein CBFβ::SMMHC (13-15). With current treatment protocols, inv(16) AML has a favorable prognosis due to high rates of initial response to chemotherapy. However, ~40% of patients will eventually relapse, at which point there are limited curative options (12, 16, 17).

Because *CM* is the initiating mutation in this subtype of AML and is expressed in all leukemia cells but not in healthy cells, it is an ideal drug target. However, the role and requirement for *CM* after malignant transformation is poorly understood. In Chronic Myeloid Leukemia (CML), another fusion protein driven leukemia, expression of BCR::ABL initiates the disease, but once established, BCR::ABL activity is largely dispensable for maintenance of the leukemia stem cell population. Currently, it is not known if this is a unique feature of BCR::ABL and CML, or if the existence of a population of cells that no longer require the activity of the initiating mutation is a more common phenomenon. Resolving this issue is critical to the development and eventual clinical use of inhibitors targeting other leukemia fusion proteins like CBFβ::SMMHC in AML.

In this study, we investigate the role of *CM* in primary mouse leukemia cells after leukemia

initiation. We found that decreased *CM* expression eliminates leukemia from the peripheral blood

and spleen, but not the bone marrow, indicating that loss of CBFβ::SMMHC activity is sufficient

to control, but not cure the disease.

Methods

<u>Mice</u>

All animal experiments were approved by the University of Nebraska Medical Center IACUC in

accordance with NIH guidelines. 4-6-week-old mice of both sexes were used.

Mx1-Cre, Cbfb$^{+/56M}$ mice

To generate *CM* expressing leukemia cells, we used conditional knock-in mice expressing

Cbfb::MYH11 from the endogenous promoter, (*Cbfb$^{+/56M}$*) paired with the inducible *Mx1-Cre*

transgene: *Mx1-Cre, Cbfb$^{+/56M}$* mice (18). To induce leukemia, *CM* mice were treated with 250μg

Polyinosine-polycytidylic acid (pI:pC) 3 times a week, for 3 rounds of treatment.

Cbfb$^{+/flMYH11}$ mice

To generate knock-in mice with the *flMYH11* allele, *LoxP* sites were added by site directed

mutagenesis (QuikChange, Agilent) flanking the *Cbfb Exon 5::MYH11* cassette in the *pPNT*

vector. Mouse embryonic stem (ES) cells were electroporated, selected, screened for

homologous recombination and injected into blastocysts, as described previously (19, 20). To

confirm the *Cbfb^{flMYH11}* allele functions as predicted, chimeric mice were bred to mice expressing

Cre from the *β-Actin* promoter (*ActB-Cre$^+$*), purchased from Jackson Laboratory (Bar Harbor, ME,

USA) and embryos harvested at embryonic day 12.5 (21). To induce leukemia, chimeric

*Cbfb[+/flMYH11]*mice were treated with a single dose of N-ethyl-N-nitrosourea (ENU, 100mg/kg) and

monitored for leukemia development, as described previously (22-24).

Transplantation of leukemia cells

CM expressing leukemia cells were transplanted into sub-lethally irradiated congenic recipient

mice by retro-orbital injection, as described previously (25, 26).

<u>Viral Transduction</u>

To excise the *Cbfb[flMYH11]* allele, lentiviral plasmids co-expressing *Cre Recombinase* and either

GFP or DsRED were used. To knockdown *Cbfb::MYH11*, a doxycycline (Dox) inducible lentiviral

plasmid expressing GFP and a shRNA against *MYH11* (*shMYH11*) was kindly provided by Dr. J.

Martens, Nijmegen, NL (27). Virus was produced in 293KEK cells and used to transduce mouse

leukemia cells, as previously described (25, 26, 28). Transduced cells were sorted for GFP or

dsRED using Aria II Cell Sorter (BD Biosciences, San Diego, CA, USA) 48hrs post transduction

and transplanted into congenic mice as above. To induce *shMYH11* expression, mice were given

5% sucrose supplemented drinking water with or without (CTRL) Dox at 2mg/ml. Mice were

sacrificed on the indicated day of treatment and tissue collected.

<u>qRT-PCR Analysis</u>

RNA was extracted using Trizol (Thermo Scientific, Waltham, MA, USA) as described previously

(25, 26). cDNA was synthesized and amplified using Luna RT Supermix and Luna Universal

qPCR kit (New England Biolabs, Ipswich, MA, USA) per manufacturer's protocol. Primer

sequences available upon request.

<u>Western Blot Analysis</u>

Cell lysates were prepared, subjected to SDS-PAGE and transferred to PVDF, as described

previously (25, 26). Membranes were blocked in 5% milk in TBST and then probed using primary

antibodies and then with secondary HRP-linked antibodies as indicated in (Supplementary Table

1). Protein bands were visualized with SuperSignal West Pico PLUS (Thermo Scientific).

Quantification was performed using ImageJ software.

<u>Flow Cytometry Analysis</u>

Cells were stained with fluorophore coupled antibodies against B220 (RA3-6B2), CD3 (17A2),

Mac-1/CD11b (M1/70), GR-1 (RM4-5), CKIT (53-6.7), Csf2rb/CD131 (JORO50), IL1RL1 (U29-

93), CD41 (MWReg30), CD123 (5B11) (BD Bioscience) and analyzed by flow cytometry (BD,

LSR II). Apoptosis was assayed using Annexin V (BD). Proliferation was determined using the

BrdU Flow Kit (BD). Mice were injected with BrdU at dose of 200mg/kg body weight. After 1 hr,

mice were sacrificed and tissue harvested. All flow cytometry data was analyzed using FlowJo

10.0 (BD, Ashland, OR, USA).

<u>Colony Assays</u>

Equal numbers of cells were suspended in Methocult (m3434, StemCell Technologies,

Vancouver, BC) in triplicate and plated in 35mm dishes, according to manufacturer's protocol.

Colonies were counted after 14 days.

<u>Whole Transcriptome sequencing</u>

Total RNA was extracted as above. RNA was quantified using Qubit 2.0 Fluorometer (Thermo

Scientific) and integrity was checked using Agilent TapeStation 4200 (Agilent Technologies, Palo

Alto, CA, USA). Library preparation and sequencing was carried out by Azenta Life Sciences

(Burlington, MA, USA). Further details about analysis provided in (Supplementary Methods).

<u>Single cell RNA sequencing</u>

Leukemia cells were enriched using the MagniSort Mouse Hematopoietic Lineage Depletion kit

(Invitrogen) then single cell RNA sequencing (scRNA-Seq) was performed using the Seq-well,

nanowell-based high throughput platform, as described previously (29, 30). Details provided in

the (Supplementary Methods).

<u>Statistical Analysis</u>

Statistical significance was determined using either Student's T-Test or ANOVA, as appropriate

(GraphPad Prism, version 10, Dotmatics, Boston, MA, USA) with p-values ≤ 0.05 deemed

significant. Sample size given in the figure legends.

Results

Knockdown of *Cbfb::MYH11* (*CM*) decreases leukemic burden in the spleen and peripheral

blood, but not the bone marrow

To test the role of *Cbfb::MYH11 (CM)* in leukemia maintenance *in vitro*, we used leukemia cells

from *Cbfb$^{+/56M}$*, *Mx1-Cre^{+}* mice which conditionally express *CM* from the endogenous *Cbfb*

promoter and develop leukemia that faithfully recapitulates human inv(16) AML (18). We collected

and transduced these leukemia cells with a doxycycline (Dox) inducible shRNA against *MYH11*

(*shMYH11*) which also constitutively expresses GFP (27). We sorted GFP^{+} cells and cultured

them with or without Dox for 48 hours (hrs). As expected, we observed that Dox induced *CM* KD

resulted in a significant decrease of *CM* expression at both the mRNA and protein levels (Fig. 1A-

B). We also found that *CM* KD induced increased apoptosis and decreased colony formation (Fig.

1C, D). These findings indicate that *CM* is required for the growth and survival of frank leukemia

cells *in vitro*.

To determine the effect of decreased *CM* expression *in vivo*, we transplanted *shMYH11*

transduced cells into congenic mice and allowed them to engraft. Once mice showed at least 20%

GFP⁺ cells in the peripheral blood (Supplementary Fig. S1A), we induced *CM* KD and determined

the percentage of GFP⁺ leukemia cells after 4, 7, 14, 21, and 28 days. We observed that *CM* KD

induced a significant decrease in GFP⁺ leukemia cells in the peripheral blood and spleen starting

at day 7, with near complete elimination by day 14 (Fig. 2A, B). In addition, we observed a notable

decrease in spleen size with *CM* KD (Supplementary Fig. S1B, C). In the bone marrow, there

was a significant decrease in leukemia cells starting at day 7 of Dox treatment. However, leukemia

cells continued to make up 20-30% of the bone marrow through day 28 (Fig. 2C). To determine if

the persistence of leukemia cells in the bone marrow was due to a failure of Dox to penetrate the

tissue, we sorted GFP⁺ leukemia cells from the bone marrow of control and Dox treated mice at

days 7 and 21. Quantitative real time PCR (qRT-PCR) and western blot analysis of bone marrow

GFP⁺ cells showed significant *CM* KD at both time points (Fig. 2D-F).

To confirm that the loss of leukemia cells in the Dox treated mice is due to decreased *CM*

expression and not the effect of Dox, we used *CM* expressing leukemia cells transduced with a

non-targeted shRNA construct. We observed no difference in the percentage of GFP⁺ leukemia

cells between Dox and control treated mice (Supplementary Fig. S1D, E). These findings indicate

that cells with reduced *CM* can survive in bone marrow, but not the peripheral blood or spleen.

Deletion of *Cbfb::MYH11* impairs the growth and survival, but does not eliminate colony

formation, of leukemia cells *in vitro*

To validate our finding that a population of leukemia cells can survive without the fusion gene, we

generated a new knock-in mouse model that allows deletion of the *Cbfb::MYH11* fusion gene by

Cre Recombinase. We used homologous recombination to knock-in the *Cbfb exon5::MYH11*

fusion flanked by *LoxP* sites into the endogenous *Cbfb* locus (*Cbfb⁺/flMYH11*) (Fig. 3A). We identified

embryonic stem cell (ES) clones with the knock-in allele by southern blot and confirmed

CBFβ::SMMHC expression by western blot (Supplementary Fig. S2A, B). Three independent

Cbfb^{+/ floxMYH11} ES clones were used to generate chimeric mice. To test that the *Cbfb* *fIMYH11* allele

can be fully excised, chimeric mice were crossed with mice expressing *Cre Recombinase* from

the *Actin B* promoter (*ActB Cre*), which should allow for excision of the fusion gene in all cells of

an embryo. We harvested embryos at day 12.5 of development. In *Cbfb*^{+/ fIMYH11} embryos lacking

the *ActB-Cre* transgene, we observed central nervous system hemorrhaging, indicating

expression of *Cbfb::MYH11*, as in previous knock-in models (Fig. 3B) (19). Importantly, this

phenotype was not observed in *Cbfb*^{+/fIMYH11}; *ActB Cre*⁺, indicating excision of the fusion gene in

the presence of Cre Recombinase. PCR using primers specific for the unexcised and excised

alleles confirmed that the *Cbfb* *fIMYH11* was efficiently excised in *Cbfb*^{+/fIMYH11}; *ActB-Cre*⁺ mice. To

generate leukemia cells expressing the *Cbfb* *fIMYH11* allele, chimeric mice were treated with N-ethyl-

N-nitrosourea and monitored for disease onset. Once leukemia was detected, cells were collected

and transplanted into congenic recipient mice and monitored for survival. Recipient mice

transplanted with leukemia cells from 3 different ES cell clones developed disease with a similar

immunophenotype and histological appearance as previous *CM* knock-in mouse models (Fig. 3C,

D; Supplementary Fig. S2C, D) (18, 31). These results indicate that *Cbfb*^{+/fIMYH11} mice develop a

frank leukemia with a similar phenotype to previous models and that the fusion gene can be

deleted by Cre Recombinase.

To test the effect of *CM* deletion *in vitro*, we transduced *Cbfb*^{+/fIMYH11} leukemia cells with a lentivirus

expressing *Cre* and GFP from an IRES (Fig. 4A). Transduced cells were sorted, and *CM* excision

was confirmed by qPCR and western blot (Fig. 4B-C). We found that *CM* deletion caused a

statistically significant increase in Annexin V, indicating induction of apoptosis (Fig. 4D). To test

the effect of *CM* deletion on colony forming ability, we plated equal numbers of control and *Cre*

transduced *Cbfb*^{+/fIMYH11} leukemia cells in methylcellulose. After 14 days, colonies were counted,

resuspended as single cells, and equal numbers of cells replated in methylcellulose**.** At both the

1st and 2nd plating, we found that *Cbfb*^{+/fIMYH11} cells transduced with *Cre* generated significantly

fewer colonies (Fig. 4E). To determine if the colonies that were able to grow from *Cre*-infected

cells had successfully deleted *CM*, individual colonies were picked and qPCR for the *Cbfb*^*flMYH11*^

allele was performed. We found that most colonies retained the unexcised *CM* allele. However,

5-20% of the colonies showed successful deletion of *CM* (Supplementary Fig. S3A, B). These

findings are consistent with our *in vivo* knockdown data in which the majority of leukemia cells

require continued expression of the fusion gene after leukemia initiation, but that there is a

population of cells that are able to survive without *CM.*

KD of *CM* alters apoptosis and proliferation

To investigate potential mechanisms for the persistence of *CM* KD cells in the bone marrow, we

analyzed apoptosis and proliferation in the leukemic cells of Dox and control treated mice at Day

7 and Day 21. At Day 7, we found that Dox, but not control, treated mice had a significant increase

in apoptosis in the spleen, but not the bone marrow (Fig. 5A left). However, at day 21, a time point

in which control mice have high leukemic burden, we found high levels of apoptotic leukemia cells

in the bone marrow of control treated mice, but not Dox treated mice (Fig. 5A right). To determine

if the failure to induce apoptosis in the bone marrow with *CM* KD is related to the levels of anti-

apoptotic proteins, we sorted leukemia cells from the spleen and bone marrow of control and Dox

treated mice on day 7 and performed western blots. We found that KD of *CM* induced decreased

BCL-2 in the spleen but caused no change in MCL-1. In leukemia cells from the bone marrow of

both control and Dox treated mice, BCL-2 was barely detectable. In contrast, MCL-1 was higher

in leukemia cells from the bone marrow of both control and Dox treated mice, as compared to

cells from the spleen (Fig. 5B, C). These findings imply that MCL-1, but not BCL-2, may contribute

to the survival of *CM* KD cells in the bone marrow.

To determine if a difference in proliferation could also contribute to the persistence of *CM* KD cells

in the bone marrow, we used BrdU incorporation. Leukemia cells from both the spleen and bone

marrow of day 21 Dox treated mice had significantly lower proliferation compared to controls (Fig.

5D). To test if this decrease was due to a change in differentiation of the leukemia cells, we

examined markers of mature myeloid cells, Mac-1 and Gr-1. In the spleen of Dox treated mice,

we observed an increase in Mac1$^+$, GR-1$^+$ leukemia cells, consistent with differentiation. In

contrast, there was no difference in the presence of Mac1 or GR-1 on the bone marrow leukemia

cells (Fig. 5E). To test whether bone marrow *CM* KD cells were instead enriched for leukemia

stem cell (LSC) markers, we examined expression of KIT, CD59a, IL1RL1, and CD123. Work by

us and others previously showed that KIT, CD59a and IL1RL1 are enriched on the LSC population

driven by *CM* expression, and CD123 is an established LSC marker in human AML (25, 32, 33).

We observed a small yet statistically significant increase in CD123 and decrease in IL1RL1 in the

leukemia cells in the bone marrow of Dox treated mice (Fig. 5F). These results indicate that the

decreased proliferation observed in the bone marrow *CM* KD cells is not caused by differentiation

to a post-mitotic mature myeloid cell or enrichment for quiescent LSCs.

KD of *CM* is associated with distinct changes in gene expression in the spleen and bone

marrow

To investigate the transcriptomic differences associated with the survival of *CM* KD leukemic cells

in the bone marrow, but not the spleen, we performed bulk RNA sequencing (RNA-Seq) on GFP$^+$

leukemia cells from both populations on day 7 of control or Dox treated mice. We identified a total

of 348 differentially expressed genes (DEGs); (221 upregulated and 127 downregulated) in

leukemia cells from the spleen of Dox treated animals, as compared to control. In leukemia cells

from the bone marrow, we identified a total of 4650 DEGs (2328 upregulated and 2322

downregulated) in cells from Dox treated mice, as compared with control. (Supplementary Table

2). Hierarchical clustering shows the bone marrow replicates from control and Dox treated mice

have similar DEG patterns (Supplementary Fig. S4A). While spleen control replicates cluster

together, one Dox treated sample clustered more closely with the control spleen samples,

suggesting an intermediate level of transcriptional changes, perhaps related to inefficient KD of

the fusion protein (Supplementary Fig. S4B, C). In the bone marrow, several known targets of the

CM fusion protein, including *Cebpe, Cebpd, Runx1 and Il1rl1*, are downregulated with *CM* KD. In

addition, genes associated with an abnormal myeloid progenitor (AMP) phenotype, including

Rhag, Cldn13, Add2, Gata2 were found to be upregulated (32). Overall, these changes in genes

expression are consistent with loss of CM activity but imply that *CM* KD cells retain transcriptional

changes associated with leukemic blasts (Fig. 6A).

To identify biological functions that promote survival of bone marrow *CM* KD cells, we compared

DEGs in spleen and bone marrow leukemia cells and performed pathway analysis on only those

genes altered in the bone marrow. Of the 2162 unique DEGs upregulated in bone marrow

leukemia cells we identified autophagy, a known promoter of AML survival, as well as mTOR

signaling, AMPK signaling, and mitophagy which have been implicated in promoting autophagy,

survival, and stress adaptation in AML (Fig 6C) (34-38). Analysis of individual autophagy related

genes showed significantly increased expression of *Atf6*, *Atg14*, *Atg2A*, *Becn1* and *mTor* (Fig

6D). These results imply a role for autophagy in the survival of *CM* KD cells in the bone marrow.

Pathway analysis of leukemia cells from the spleen indicated that the top deregulated pathway

with KD of *CM* is apoptosis, consistent with the Annexin V staining and western blot analysis in

Figure 5 (Supplementary Fig. 4C).

To determine if *CM* KD induces distinct gene expression changes in different subpopulations of

leukemia cells*,* we performed single-cell RNA sequencing (scRNA-Seq). We harvested cells from

the bone marrow and spleen of control and dox treated mice at day 7 of treatment. Single cell

suspensions were lineage depleted and analyzed by flow cytometry to confirm enrichment of

leukemia cells. All samples showed >60% leukemia cells, with the bone marrow approaching 80%

in both control and Dox treated samples (Supplementary Fig. 5A). Single-cell transcript libraries

were prepared and sequenced.

Unbiased clustering analysis revealed multiple subpopulations of leukemia cells in both the spleen

and bone marrow. Interestingly, *CM* KD affected the relative proportion of these clusters, with the

number of cells increased in some clusters while others decreased in samples from Dox treated

mice (Fig 7A, B). In the spleen, Clusters 2 and 4 appeared to be the most sensitive to *CM* KD

(Supplementary Fig. 5B,C). In the bone marrow, Clusters 3, 4, 5, 6, 9 and 10 showed marked

reduction with *CM* KD while the number of cells in Clusters 0, 1, 2, 7 and 8 increased, indicating

these clusters don't require *CM* expression for survival (Fig. 7B). This analysis shows that there

is dramatic cellular heterogeneity among the leukemia cells in both the bone marrow and spleen,

with different subpopulations differentially dependent on *CM* expression.

To test if there are inherent transcriptional programs promoting the survival of the *CM*-

independent cells in the bone marrow, we compared the transcripts expressed in the clusters that

don't require CM expression. We identified 456 genes expressed in all *CM*-independent clusters

(Fig 7C). Among these, several genes including *Ccl5, Ccr1, S100a4, CD74, CD27, Ccnd2* and

Cdkn1a are all associated with leukemia survival and proliferation (39-43). Consistent with this

finding, pathway analysis indicates that these genes are significantly enriched for molecular

functions involving cytokine and chemokine activity (Fig. 7D). In contrast, the common genes

expressed in *CM*-dependent clusters were associated with non-immune molecular functions

(Supplementary Fig. 5D). Collectively these results imply that cytokine and chemokine signaling

within the bone marrow microenvironment may be a critical mechanism promoting survival of *CM*-

independent leukemia cells.

_CM_ KD leukemia cells can re-establish disease

The persistence of a *CM* KD leukemic population in the bone marrow raises the possibility that

these cells may be able to cause relapsed disease. To test this possibility, we maintained mice

transplanted with *shMYH11* transduced leukemia cells on Dox or control drinking water until they

showed signs of lethal disease. 11 out of 18 Dox treated mice showed re-emergence of GFP$^+$

cells in the blood (Fig. 8A). To determine if these cells expressed *CM*, we sorted for GFP$^+$ cells

and performed qRT-PCR and western blot (Fig. 8 B-D). We found that GFP$^+$ relapse cells had

reduced *CM* at both the mRNA and protein levels, as compared to parental leukemia cells from

control treated mice. In addition, we found that the immunophenotype of relapsed *CM* KD

leukemia cells was similar to parental leukemia cells from control mice (Supplementary Fig. S6A

and B). To understand the molecular mechanism allowing *CM* KD cells to re-emerge into the

blood, we performed whole transcriptome sequencing on relapse and parental leukemia cells.

The leukemic cells from relapsed mice were harvested from the spleen and compared with either

spleen or bone marrow from control mice. Principal component analysis (PCA) showed that

relapsed samples did not cluster with parental leukemia cells harvested from either the bone

marrow or spleen indicating that relapse is accompanied by widespread alterations in transcription

(Fig. 8E). In addition, only 3 of the 4 relapsed samples clustered together, implying that there may

be multiple transcriptional programs that promote re-emergence of *CM* KD cells from the bone

marrow. Comparing the transcriptome of relapsed samples to control, we observed a total of 930

DEGs (359 upregulated and 571 downregulated) in relapsed samples compared to parental

leukemia cells (Supplementary Table 3). Further, pathway analysis indicates that the DEGs are

associated with multiple pathways implicated in leukemia survival and relapse, including

interferon regulator factor (IRF) signaling (Fig. 8F). Specifically, *Irf1*, *Irf7*, and *Irf9* are decreased

in relapse samples compared to controls (Fig. 8G). Notably, loss of these genes is associated

with relapse in AML patients (46-48). This implies that suppression of IRF signaling allows *CM*

KD leukemia cells to survive outside the bone marrow.

353 Collectively, this study indicates that while *CM* is required for the survival of most frank leukemia

354 cells, a population of leukemia cells in the bone marrow do not strictly require *CM* activity and can

355 re-establish fatal disease.

357 **Discussion**

358 Given that CBFβ::SMMHC is expressed in all leukemia cells of inv(16) AML patients, but not in

359 healthy cells it is a tempting target for drug development. However, most studies have focused

360 on the role of the fusion protein during leukemia initiation and less is known about its role in frank

361 leukemia cells in the context of cooperating mutations. Previous work in the inv(16) immortalized

362 cell line, ME-1, reported that decreased expression of *CM* caused changes in gene expression

363 consistent with differentiation, but did not cause any changes in viability at the time point analyzed,

364 raising the possibility that continued *CM* expression may not be strictly required for leukemia

365 maintenance (27). To our knowledge, ours is the first study to test whether *CM* activity is required

366 for leukemia cell survival using non-immortalized leukemia cells. We found that KD of *CM* induced

367 apoptosis and decreased colony forming ability *in vitro*. *CM* KD *in vivo* caused a near complete

368 elimination of leukemia cells in the blood and spleen. Surprisingly, the leukemic burden in the

369 bone marrow did not decrease below 20-30% despite almost complete loss of the CBFβ::SMMHC

370 protein, implying that the bone marrow microenvironment can sustain leukemia cells despite the

371 loss of *CM* expression.

373 Our observation that bone marrow *CM* KD cells , but not spleen, had decreased apoptosis as

374 compared to controls suggests tissue specific effects on the survival of leukemic cells *in vivo*.

375 Thus, to identify potential factors promoting the survival of bone marrow *CM* KD cells, we analyzed

376 the anti-apoptotic proteins BCL-2 and MCL-1 (44, 45). However, we observed that both control

377 and *CM* KD leukemic cells had robust expression of MCL-1 but nearly undetectable levels of BCL-

378 2, indicating that differential expression of anti-apoptotic proteins does not explain their survival.

Rather, our bulk RNA-seq indicates that the upregulation of autophagy may contribute to the

survival of bone marrow *CM* KD cells. These findings are consistent with recent work showing a

role for autophagy in leukemia progression and that overexpression of the autophagy genes

BECLIN1 and *ATG* are associated with shorter overall survival in AML patients (46, 47).

To understand the cellular heterogeneity in *CM* sufficient and KD cells, we performed scRNA-

Seq. We found that there are clusters of leukemia cells with distinct transcriptional profiles and

different requirements for *CM* expression. Some clusters had decreased cellularity while others

were increased, implying the presence of *CM*-dependent and *CM*-independent leukemic

populations in the bone marrow. This is consistent with our *in vivo* results where we observed an

initial decrease in bone marrow leukemic burden of dox treated mice, but eventual stabilization of

the leukemia population at 20-30%. Interestingly, the population of *CM*-independent leukemia

cells did not continue to increase over time, implying that there is a maximum number of *CM*-

independent leukemia cells that the bone marrow can support. Of note, the *CM*-independent

clusters in the bone marrow shared a transcriptional profile associated with cytokine and

chemokine activity, which is consistent with a model in which the bone marrow microenvironment

provides a limited amount of survival factors, keeping the *CM*-independent population from

expanding indefinitely in the bone marrow.

Our results concerning the role of CBFβ::SMMHC during leukemia maintenance are strikingly

similar to what has been demonstrated with the BCR::ABL fusion protein in CML. Pharmacological

inhibitors of BCR::ABL, such as imatinib, are remarkably successful at inducing long-term, stable

cytogenetic remission in CML (48, 49). However, these kinase inhibitors generally do not eliminate

the LSC population in the bone marrow. Further, transgenic models have demonstrated that with

near complete knockdown of *BCR::ABL* expression, LSCs remain in the bone marrow and are

able to re-establish disease when BCR::ABL activity is restored (50, 51). In our model, we also

observed a similar population of LSC-like cells that can survive in the bone marrow and re-establish disease, despite near complete loss of the CBFβ::SMMHC protein. In our case, relapse did not require re-expression of the fusion protein. However, given the period of latency before relapse, it is likely that additional mutations developed to compensate for the lack of CBFβ::SMMHC activity.

Collectively, this study demonstrates that *CM* expression continues to be required after leukemic transformation and supports the development of inhibitors targeting the fusion protein for the treatment of inv(16) AML. Our data indicates such inhibitors would be effective at managing leukemic burden and, when used in combination with other therapeutics, have the potential to cure the disease.

Acknowledgments

The authors would like to thank Paul Liu, M.D, Ph.D (NHGRI/NIH) for his support and encouragement of this project, as well as the members of the UNMC flow cytometry and Bioinformatics and Systems Biology Core Facilities, and the Nebraska Center for Molecular Target Discovery and Development for their expertise.

This work was supported by R00 CA148963 and R01 CA244900 to R.K.H and T32 CA009467 to support C.L. C.R., L.A., and L.G. were supported by the Intramural Research Programs of the National Human Genome Research Institute, National Institutes of Health. Core facilities used for this project are supported by a state fund from the Nebraska Research Initiative, the Fred and Pamela Buffett Cancer Center's National Cancer Institute Cancer Support Grant (P30 CA036727), and the Nebraska Center for Molecular Target Discovery and Development COBRE (P20 GM121316).

Author Contributions

S.P. designed and performed experiments, analyzed data and wrote the paper. Y.W. designed and performed experiments and analyzed data. M.B., C.L., A.D., and S.A.S. performed experiments. R.D. and P.W analyzed single cell and bulk RNA-Seq data, respectively. C.R., L.A., and L.G. generated, bred, and mutagenized $Cbfb^{+/floxMYH11}$ knock-in mice. S.A.S., K.J.H. and R.K.H designed experiments, analyzed data and wrote the paper.

Disclosure of Conflicts of Interest

The authors have no conflicts of interest to disclose.

References

1.   Akinduro O, Weber TS, Ang H, Haltalli MLR, Ruivo N, Duarte D, et al. Proliferation dynamics of acute myeloid leukaemia and haematopoietic progenitors competing for bone marrow space. Nature communications. 2018;9(1):519.

2.   Thomas D, Majeti R. Biology and relevance of human acute myeloid leukemia stem cells. Blood. 2017;129(12):1577-85.

3.   Lane SW, Scadden DT, Gilliland DG. The leukemic stem cell niche: current concepts and therapeutic opportunities. Blood. 2009;114(6):1150-7.

4.   Kumar CC. Genetic abnormalities and challenges in the treatment of acute myeloid leukemia. Genes Cancer. 2011;2(2):95-107.

5.   Speck NA, Gilliland DG. Core-binding factors in haematopoiesis and leukaemia. Nat Rev Cancer. 2002;2(7):502-13.

6.   Castilla LH, Perrat P, Martinez NJ, Landrette SF, Keys R, Oikemus S, et al. Identification of genes that synergize with Cbfb-MYH11 in the pathogenesis of acute myeloid leukemia. Proceedings of the National Academy of Sciences of the United States of America. 2004;101(14):4924-9.

7.   Sood R, Hansen NF, Donovan FX, Carrington B, Bucci D, Maskeri B, et al. Somatic mutational landscape of AML with inv(16) or t(8;21) identifies patterns of clonal evolution in relapse leukemia. Leukemia. 2016;30(2):501-4.

8.   Duployez N, Marceau-Renaut A, Boissel N, Petit A, Bucci M, Geffroy S, et al. Comprehensive mutational profiling of core binding factor acute myeloid leukemia. Blood. 2016;127(20):2451-9.

9.   Khoury JD, Solary E, Abla O, Akkari Y, Alaggio R, Apperley JF, et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. Leukemia. 2022;36(7):1703-19.

10.   Balgobind BV, Hollink IH, Arentsen-Peters ST, Zimmermann M, Harbott J, Beverloo HB, et al. Integrative analysis of type-I and type-II aberrations underscores the genetic heterogeneity of pediatric acute myeloid leukemia. Haematologica. 2011;96(10):1478-87.

11.   Kihara R, Nagata Y, Kiyoi H, Kato T, Yamamoto E, Suzuki K, et al. Comprehensive analysis of genetic alterations and their prognostic impacts in adult acute myeloid leukemia patients. Leukemia. 2014;28(8):1586-95.

12.   Mancini M, Cedrone M, Diverio D, Emanuel B, Stul M, Vranckx H, et al. Use of dual-color interphase FISH for the detection of inv(16) in acute myeloid leukemia at diagnosis, relapse and during follow-up: a study of 23 patients. Leukemia. 2000;14(3):364-8.

13.   Kundu M, Liu PP. Function of the inv(16) fusion gene CBFB-MYH11. Current opinion in hematology. 2001;8(4):201-5.

488   14.   Liu P, Tarle SA, Hajra A, Claxton DF, Marlton P, Freedman M, et al. Fusion between
489   transcription factor CBF beta/PEBP2 beta and a myosin heavy chain in acute myeloid leukemia.
490   Science. 1993;261(5124):1041-4.
491

492   15.   Liu PP, Wijmenga C, Hajra A, Blake TB, Kelley CA, Adelstein RS, et al. Identification of
493   the chimeric protein product of the CBFB-MYH11 fusion gene in inv(16) leukemia cells. Genes
494   Chromosomes Cancer. 1996;16(2):77-87.
495

496   16.   Marcucci G, Mrozek K, Ruppert AS, Maharry K, Kolitz JE, Moore JO, et al. Prognostic
497   factors and outcome of core binding factor acute myeloid leukemia patients with t(8;21) differ
498   from those of patients with inv(16): a Cancer and Leukemia Group B study. J Clin Oncol.
499   2005;23(24):5705-17.
500

501   17.   Appelbaum FR, Kopecky KJ, Tallman MS, Slovak ML, Gundacker HM, Kim HT, et al.
502   The clinical spectrum of adult acute myeloid leukaemia associated with core binding factor
503   translocations. British journal of haematology. 2006;135(2):165-73.
504

505   18.   Kuo YH, Landrette SF, Heilman SA, Perrat PN, Garrett L, Liu PP, et al. Cbf beta-
506   SMMHC induces distinct abnormal myeloid progenitors able to develop acute myeloid leukemia.
507   Cancer Cell. 2006;9(1):57-68.
508

509   19.   Castilla LH, Wijmenga C, Wang Q, Stacy T, Speck NA, Eckhaus M, et al. Failure of
510   embryonic hematopoiesis and lethal hemorrhages in mouse embryos heterozygous for a
511   knocked-in leukemia gene CBFB-MYH11. Cell. 1996;87(4):687-96.
512

513   20.   Kamikubo Y, Zhao L, Wunderlich M, Corpora T, Hyde RK, Paul TA, et al. Accelerated
514   leukemogenesis by truncated CBF beta-SMMHC defective in high-affinity binding with RUNX1.
515   Cancer Cell. 2010;17(5):455-68.
516

517   21.   Lewandoski M, Meyers EN, Martin GR. Analysis of Fgf8 gene function in vertebrate
518   development. Cold Spring Harb Symp Quant Biol. 1997;62:159-68.
519

520   22.   Castilla LH, Garrett L, Adya N, Orlic D, Dutra A, Anderson S, et al. The fusion gene
521   Cbfb-MYH11 blocks myeloid differentiation and predisposes mice to acute myelomonocytic
522   leukaemia [letter]. Nat Genet. 1999;23(2):144-6.
523

524   23.   Kamikubo Y, Hyde RK, Zhao L, Alemu L, Rivas C, Garrett LJ, et al. The C-terminus of
525   CBFbeta-SMMHC is required to induce embryonic hematopoietic defects and leukemogenesis.
526   Blood. 2013;121(4):638-42.
527

528   24.   Hyde RK, Zhao L, Alemu L, Liu PP. Runx1 is required for hematopoietic defects and
529   leukemogenesis in Cbfb-MYH11 knock-in mice. Leukemia. 2015;29(8):1771-8.
530

531   25.   Wang Y, Richter L, Becker M, Amador C, Hyde RK. IL1RL1 is dynamically expressed on
532   Cbfb-MYH11+ leukemia stem cells and promotes cell survival. Scientific Reports.
533   2019;9(1):1729.
534

26.   Richter LE, Wang Y, Becker ME, Coburn RA, Williams JT, Amador C, et al. HDAC1 is a
Required Cofactor of CBFbeta-SMMHC and a Potential Therapeutic Target in Inversion 16
Acute Myeloid Leukemia. Mol Cancer Res. 2019.

27.   Mandoli A, Singh AA, Jansen PW, Wierenga AT, Riahi H, Franci G, et al. CBFB-
MYH11/RUNX1 together with a compendium of hematopoietic regulators, chromatin modifiers
and basal transcription factors occupies self-renewal genes in inv(16) acute myeloid leukemia.
Leukemia. 2013.
542
543 28.   Dhir A, Ethell A, Watkins R, Lam C, Tur-Rodriguez K, Persinger J, et al. The splicing
544 factor PTBP1 interacts with RUNX1 and is required for leukemia cell survival. Leukemia. 2025.
545
546 29.   Gierahn TM, Wadsworth MH, 2nd, Hughes TK, Bryson BD, Butler A, Satija R, et al. Seq-
547 Well: portable, low-cost RNA sequencing of single cells at high throughput. Nature methods.
548 2017;14(4):395-8.
549
550 30.   Zhou Y, Dogiparthi VR, Ray S, Schaefer MA, Harris HL, Rowley MJ, et al. Defining a
551 cohort of anemia-activated cis elements reveals a mechanism promoting erythroid precursor
552 function. Blood Adv. 2023;7(20):6325-38.
553
554 31.   Hyde RK, Kamikubo Y, Anderson S, Kirby M, Alemu L, Zhao L, et al. Cbfb/Runx1
555 repression-independent blockage of differentiation and accumulation of Csf2rb-expressing cells
556 by Cbfb-MYH11. Blood. 2010;115(7):1433-43.
557
558 32.   Diemer JL, Yu K, Kelly M, Zhen T, Anderson S, Lopez G, et al. Characterization of
559 leukemia progression in the Cbfb-MYH11 knockin mice by single cell RNA sequencing.
560 Leukemia. 2023;37(7):1549-53.
561
562 33.   Arnone M, Konantz M, Hanns P, Paczulla Stanger AM, Bertels S, Godavarthy PS, et al.
563 Acute Myeloid Leukemia Stem Cells: The Challenges of Phenotypic Heterogeneity. Cancers
564 (Basel). 2020;12(12).
565
566 34.   Saito Y, Chapple RH, Lin A, Kitano A, Nakada D. AMPK Protects Leukemia-Initiating
567 Cells in Myeloid Leukemias from Metabolic Stress in the Bone Marrow. Cell stem cell.
568 2015;17(5):585-96.
569
570 35.   Glytsou C, Chen X, Zacharioudakis E, Al-Santli W, Zhou H, Nadorp B, et al. Mitophagy
571 Promotes Resistance to BH3 Mimetics in Acute Myeloid Leukemia. Cancer Discov.
572 2023;13(7):1656-77.
573
574 36.   Chen Y, Chen J, Zou Z, Xu L, Li J. Crosstalk between autophagy and metabolism:
575 implications for cell survival in acute myeloid leukemia. Cell Death Discov. 2024;10(1):46.
576
577 37.   Santos de Macedo BG, Albuquerque de Melo M, Pereira-Martins DA, Machado-Neto JA,
578 Traina F. An updated outlook on autophagy mechanism and how it supports acute myeloid
579 leukemia maintenance. Biochim Biophys Acta Rev Cancer. 2024;1879(6):189214.
580
581 38.   Nepstad I, Hatfield KJ, Gronningsaeter IS, Reikvam H. The PI3K-Akt-mTOR Signaling
582 Pathway in Human Acute Myeloid Leukemia (AML) Cells. Int J Mol Sci. 2020;21(8).
583
39.   Menssen AJ, Hudson CA, Alonzo T, Gerbing R, Pardo L, Leonti A, et al. CD74 is
expressed in a subset of pediatric acute myeloid leukemia patients and is a promising target for
therapy: a report from the Children's Oncology Group. Haematologica. 2024;109(10):3182-93.

40.   Alanazi B, Munje CR, Rastogi N, Williamson AJK, Taylor S, Hole PS, et al. Integrated
nuclear proteomics and transcriptomics identifies S100A4 as a therapeutic target in acute
myeloid leukemia. Leukemia. 2020;34(2):427-40.
590
591   41.   Waldeck S, Rassner M, Keye P, Follo M, Herchenbach D, Endres C, et al. CCL5
592   mediates target-kinase independent resistance to FLT3 inhibitors in FLT3-ITD-positive AML.
593   Mol Oncol. 2020;14(4):779-94.
594
595   42.   Mou J, Huang Q, Liu X, Liu W, Liu Y, Mei Y, et al. AML1-ETO and CCND2
596   overexpression cooperate to drive acute myeloid leukemia initiation and progression. J Leukoc
597   Biol. 2025;117(6).
598
599   43.   Kittang AO, Hatfield K, Sand K, Reikvam H, Bruserud O. The chemokine network in
600   acute myelogenous leukemia: molecular mechanisms involved in leukemogenesis and
601   therapeutic implications. Curr Top Microbiol Immunol. 2010;341:149-72.
602
603   44.   Fowler-Shorten DJ, Hellmich C, Markham M, Bowles KM, Rushworth SA. BCL-2
604   inhibition in haematological malignancies: Clinical application and complications. Blood Rev.
605   2024;65:101195.
606
607   45.   Griffioen MS, de Leeuw DC, Janssen J, Smit L. Targeting Acute Myeloid Leukemia with
608   Venetoclax; Biomarkers for Sensitivity and Rationale for Venetoclax-Based Combination
609   Therapies. Cancers (Basel). 2022;14(14).
610
611   46.   Lian Y, Xie Y, Hong M, Zhu Y, Zhao H, Zhao X, et al. Clinical significance of BECLIN1
612   and ATG5 expression in acute myeloid leukemia patients. Int J Clin Exp Pathol.
613   2018;11(3):1529-37.
614
615   47.   Seo W, Silwal P, Song I-C, Jo E-K. The dual role of autophagy in acute myeloid
616   leukemia. Journal of hematology & oncology. 2022;15(1):51.
617
618   48.   Druker BJ, Guilhot F, O'Brien SG, Gathmann I, Kantarjian H, Gattermann N, et al. Five-
619   year follow-up of patients receiving imatinib for chronic myeloid leukemia. N Engl J Med.
620   2006;355(23):2408-17.
621
622   49.   Hughes TP, Hochhaus A, Branford S, Muller MC, Kaeda JS, Foroni L, et al. Long-term
623   prognostic significance of early molecular response to imatinib in newly diagnosed chronic
624   myeloid leukemia: an analysis from the International Randomized Study of Interferon and
625   STI571 (IRIS). Blood. 2010;116(19):3758-65.
626
627   50.   Hamilton A, Helgason GV, Schemionek M, Zhang B, Myssina S, Allan EK, et al. Chronic
628   myeloid leukemia stem cells are not dependent on Bcr-Abl kinase activity for their survival.
629   Blood. 2012;119(6):1501-10.
630
631   51.   Chen Y, Zou J, Cheng F, Li W. Treatment-Free Remission in Chronic Myeloid Leukemia
632   and New Approaches by Targeting Leukemia Stem Cells. Frontiers in oncology.
633   2021;11:769730.
634

Figure Legends

Figure 1: Inducible knockdown of *CM* causes apoptosis and impairs colony formation *in vitro*. A. Bar graph showing quantitative reverse transcription PCR (qRT-PCR) of *CBFB::MYH11* in *Cbfb^{+/56M}* leukemia cells transduced with an inducible shRNA against *MYH11* (shMYH11) treated *in vitro* with either control (CTRL) or doxycycline (Dox) for 48 hrs **B.** Representative western blot (left) and bar graph of relative CBFβ::SMMHC protein levels (right) in *shMYH11* transduced leukemia cells treated as in A. **C**. Bar graph showing the percentage (%) of *shMYH11* transduced leukemia cells treated with control or Dox that are positive for Annexin V. **D.** Bar graph showing relative colony numbers generated by cells treated as in A. N ≥ 3 biological and technical replicates, * = p ≤ 0.05 compared to CTRL.

Figure 2: *CM* knockdown decreases leukemic burden *in vivo*. A. Line graph showing % GFP^+ *shMYH11* transduced leukemia cells (left) and representative flow cytometry plots (right) in the peripheral blood (PB), **B.** spleen (SP) and **C.** bone marrow (BM) of mice treated with Dox or CTRL for 7, 14, 21, or 28 days. **D.** Bar graph showing relative expression of *CM* in sorted GFP^+ leukemia cells from the BM of mice treated with Dox or CTRL for 7 and **E.** 21 days. **F.** Representative western blot (left) and bar graph (right) of relative CBFβ::SMMHC protein levels in sorted GFP^+ cells from the BM of mice treated with either CTRL or Dox for 7 days. N ≥ 3 biological replicates, ns = not significant, * = p ≤ 0.05.

Figure 3: Generation of flox*Cbfb::MYH11* mice. A. Schematic representation of the embryonic stem (ES) cell targeting strategy. The probe (0.2C) and restriction enzyme sites (NcoI) used for screening are indicated. The black arrows represent PCR primers for the excised allele, and the purple arrows represent primers for the unexcised allele. **B.** Pictures of representative embryos harvested at embryonic day 12.5 generated by crossing *Cbfb^{+/floxMYH11}* chimeras with transgenic

mice expressing *Cre recombinase* (Cre) from the *β–Actin* promoter (*ActB Cre*). The right panel

shows a representative PCR for the excised and unexcised *Cbfb$^{+/floxMYH11}$* allele in DNA from

embryos of the indicated genotype. **C.** Kaplan-Meyer survival curve of recipient mice transplanted

with *Cbfb$^{+/flMYH11}$* leukemic cells from three different chimeric donor mice. **D.** Representative

Wright-Giemsa-stained peripheral blood smear and hematoxylin & eosin-stained spleen and liver

section from a leukemic *Cbfb$^{+/flMYH11}$* recipient mouse.

Figure 4. Deletion of *Cbfb::MYH11* induces apoptosis and reduces colony formation *in*

***vitro* A.** Schematic representation of the experimental design to test the effect of *Cbfb::MYH11*

deletion *in vitro*. **B.** Bar graph of qRT-PCR of the unexcised or excised *Cbfb$^{+/floxMYH11}$* alleles in

leukemia cells transduced with a lentivirus expressing Cre Recombinase (Cre) or Control (CTRL)

and sorted for GFP. **C.** Representative western blot of CBFβ::SMMHC in *Cbfb$^{+/flMYH11}$* leukemia

cells infected with CTRL or Cre virus. **D.** Bar graph of % Annexin V in CTRL or Cre transduced

Cbfb^{flMYH11} leukemia cells. **E.** Bar graph showing the relative number of colonies from equal

numbers of CTRL or Cre transduced *Cbfb$^{+/flMYH11}$* leukemia cells. Colonies were scored on day

14. N = 3 biological and technical replicates, * = $p \leq 0.05$, *** = $p \leq 0.001$ as compared to CTRL.

Figure 5: *CM* Knockdown induces apoptosis in leukemia cells in the spleen, but not the

bone marrow. A. Bar graph showing the % Annexin V$^+$, GFP$^+$ cells from the SP and BM of mice

treated with CTRL or Dox for 7 (left) or 21 days (right). **B.** Representative western blot and **C.** Bar

graph of quantification for the indicated proteins from leukemia cells sorted for GFP from the SP

and BM after 7 days of treatment with CTRL or Dox **D.** Bar graph of the % BrdU$^+$, GFP$^+$ leukemia

cells harvested from the SP and BM of mice after 21 days of treatment with CTRL or Dox. **E.** Bar

graph showing % GFP$^+$ cells expressing the mature myeloid markers Mac-1 and Gr-1 from the

SP and BM of mice treated as in D. **F.** Bar graph showing the % GFP$^+$ cells expressing the

indicated leukemia stem cell markers in the BM from of mice treated as in D. N ≥ 3 biological

replicates, ns = not significant, * = p ≤ 0.05, ** = p ≤ 0.01, *** = p ≤ 0.001 as compared to CTRL.

Figure 6: Increased expression of autophagy related genes in *CM* knockdown leukemia

cells from the bone marrow. A. Volcano plot showing distribution of the statistically significant

differentially expressed genes (DEGs) in the GFP$^+$ sorted leukemia cells from the BM of mice

treated for 7 days with CTRL or Dox. A few of the key genes with a $\log_2$ fold change ≥ 0.5 and

adjusted p value ≤ 0.05 are highlighted. **B**. Venn diagram showing overlap of significantly

upregulated DEGs in *CM* KD leukemia cells from the SP and BM, as compared to CTRL. **C.**

Pathway analysis of upregulated DEGs in *CM* KD cells from the BM showing the top KEGG

pathways affected. **D.** Box plot depicting relative expression of indicated autophagy associated

genes in control and *CM* KD leukemia cells from the BM. N = 3 biological replicates, * = padj ≤

0.05 as compared to CTRL.

Figure 7: Single cell transcriptomics reveals distinct *CM*-dependent and *CM*-independent

subpopulations in the bone marrow. A. Umap projection of leukemia cells from the BM of CTRL

or Dox treated mice. Each dot represents a single cell and is colored according to the assigned

transcriptional cluster. **B.** Bar graph of the number of cells in each cluster in BM samples from

CTRL or Dox treated mice **C.** Venn diagram representing the shared genes expressed by cells in

the *CM* independent clusters in BM samples from CTRL mice with 456 genes common among

the five groups. **D.** Top 5 molecular functions associated with the 456 shared genes from C.

Figure 8: The surviving *CM* knockdown cells in the bone marrow can give rise to relapse.

A. Line graph showing the % GFP$^+$, *shMYH11* transduced cells in PB of mice treated for up to 42

days with either CTRL or Dox**. B.** Bar graph of relative *CM* expression in GFP$^+$ sorted leukemia

cells from Dox treated mice after relapse. **C.** Representative western blot and **D.** Bar graph of

relative CBFβ::SMMHC levels from sorted GFP$^+$ cells harvested from CTRL or Dox treated mice.

Quantification to GFP. **E.** Principal Component Analysis (PCA) plots of whole transcriptome

sequencing data from GFP sorted relapse or CTRL leukemic cells harvested from the SP or BM.

F. Pathway analysis of the differentially expressed genes in relapse samples compared to the

combined SP and BM CTRL leukemia samples. **G.** Bar graph depicting the *Irf* genes deregulated

in Dox treated mice after relapse. N ≥ 3 biological replicates, * = p ≤ 0.05 as compared to CTRL.

Figure 1

Figure 2

Figure 3

A. Targeting construct

Wildtype *Cbfb⁺* allele

Cbfb^flMYH11 allele

B.

Cbfb^+/flMYH11

Cbfb^+/flMYH11, ActB Cre⁺

Unexcised

Excised

C.

Chimera A
Chimera B
Chimera C

D.

Peripheral Blood

Spleen

Liver

Figure 4

A.

B.

C.

D. **E.**

Figure 5

Figure 6

Figure 7

Figure 8