The biogeography of evolutionary radiations on oceanic archipelagos

Baptiste Brée[1], Thomas J. Matthews[2,3], José María Fernández-Palacios[4], Christian Paroissin[5], Kostas A. Triantis[6], Robert J. Whittaker[7,8], François Rigal[1,3]

Université de Pau et des Pays de l'Adour (UPPA), Energy Environment Solutions (E2S), Centre National de la Recherche Scientifique (CNRS), Institut des Sciences Analytiques et de Physico-Chimie pour l'Environnement et les Matériaux (IPREM), Pau, France

GEES (School of Geography, Earth and Environmental Sciences) and Birmingham Institute of Forest Research, University of Birmingham, Birmingham, UK

cE3c—Centre for Ecology, Evolution and Environmental Changes/Azorean Biodiversity Group/CHANGE—Global Change and Sustainability Institute and Universidade dos Açores—Faculty of Agricultural Sciences and Environment, Angra do Heroismo, Azores, Portugal

Island Ecology and Biogeography Group, Instituto Universitario de Enfermedades Tropicales y Salud Pública de Canarias (IUETSPC), Universidad de La Laguna (ULL), La Laguna, Spain

Université de Pau et des Pays de l'Adour (UPPA), Energy Environment Solutions (E2S), Centre National de la Recherche Scientifique (CNRS), Laboratoire de Mathématiques et de leurs Applications de Pau (LMAP), Pau, France

Faculty of Biology, Department of Ecology and Taxonomy, National and Kapodistrian University of Athens, Athens, Greece

School of Geography and the Environment, University of Oxford, Oxford, UK

Center for Macroecology, Evolution and Climate, GLOBE Institute, University of Copenhagen, Copenhagen, Denmark

Abstract

Evolutionary radiations on oceanic archipelagos (ROAs) have long served as models for understanding evolutionary and ecological processes underlying species diversification. Yet, diversity patterns emerging from ROAs have received relatively little attention from biogeographers, even though characterizing the effect of key geo-environmental factors on island clades species distribution could be important for unraveling diversification dynamics. In this study, we conducted a comparative analysis using island-specific species richness values for approximately one hundred ROAs across major oceanic archipelagos (mostly Hawaii, Canary Islands, Galapagos and Fiji) and taxa (vascular plants, invertebrates and vertebrates). Our aim was to determine whether (1) ROA species richness patterns scale as a function of key geo-environmental factors including island area, geological age, environmental heterogeneity (elevation and topographic complexity) and inter-island isolation, and (2) whether the magnitude of the effects of these factors varies across archipelagos and taxa. Our results identified elevation as a key driver of ROA species richness patterns on islands, supporting existing theoretical and empirical work that highlighted the central role of environmental heterogeneity in driving diversification on oceanic islands. As importantly, we found that the influence of geo-environmental factors varies across archipelagos and taxa, suggesting that unique archipelagic dynamics and biological traits together shape diversification differently. Our findings emphasize the value of applying biogeographical modeling at resolution of individual radiations to improve our understanding of evolutionary processes on oceanic archipelagos.

1.Introduction

Evolutionary radiations on oceanic archipelagos (hereafter ROAs) have served as invaluable models for understanding the eco-evolutionary processes underlying species diversification, due to the high-magnitude geological and environmental changes over evolutionary timescales that characterize volcanic islands (Losos & Ricklefs 2009, Gillespie 2016, Shaw & Gillespie 2016, Whittaker et al. 2023). Here, we define a ROA as a monophyletic clade endemic to an oceanic archipelago or a group of oceanic archipelagos that originates from a single colonization event. Since the initial description of the Galápagos finch radiation more than 150 years ago (Darwin 1845, Grant & Grant 2007), many ROAs have been described for multiple taxa and archipelagos, including the iconic Hawaiian Lobeliads (Givnish et al. 2009) and *Drosophila* (O'Grady & DeSalle 2018), and *Naesiotus* land snails of the Galápagos (Parent & Crespi 2006).

Although ROA research has developed considerably in recent decades (e.g. Emerson 2002, Cerca et al. 2023), the diversity patterns emerging from ROAs have attracted surprisingly little attention from island biogeographers. Yet, an evaluation of the geo-environmental factors influencing the island species richness pattern emerging from radiating processes could prove crucial for our understanding of the diversification dynamics taking place in oceanic archipelagos (Whittaker et al. 2008, Economo et al. 2015). In other words, it might be informative to address why, within a given archipelago, some islands have accumulated more species than others during the radiation process. However, despite recent work on oceanic and other island-like systems (e.g. Roell et al. 2021, Wagner et al. 2014, Economo et al. 2015, Lim & Marshall 2017, Tao et al. 2021), evolutionary radiations have rarely been studied from this perspective.

Since the publication of the Equilibrium Theory of Island Biogeography (ETIB) by MacArthur and Wilson (1963, 1967), many strong and predictable relationships between island characteristics and species

richness have been identified (Warren et al. 2015). Typically, models applied include island area and

isolation as well as environmental heterogeneity and geological age as primary predictors (e.g. Whittaker

et al. 2008, 2023, Triantis et al. 2012, Weigelt & Kreft 2013, Barajas-Barbosa et al. 2020). Most of these

factors have also been reported as direct or indirect drivers of species diversification on oceanic islands

(e.g. Funk & Wagner 1995, Juan et al. 2000, Lim & Marshall 2017, Roell et al. 2021). Therefore, it is

straightforward to hypothesise that they will also affect the species richness on islands for a given ROA

(hereafter Island S_{ROA}). For example, a positive relationship between area and Island S_{ROA} is expected (see

Parent & Crespi 2006, Whittaker et al. 2008, Beatty et al. 2017) because larger islands in general offer

both more ecological (e.g. diversity of habitats) and geographical (intra-island subdivision of habitats and

populations) opportunities for speciation (Losos & Schluter 2000).

For isolation, it is primarily the way in which islands are isolated from each other within the archipelago

(inter-island isolation) that is expected to determine Island S_{ROA} patterns. Hence, as the ROA spreads

throughout an archipelago, species are more likely to colonize the least-isolated islands leading to a

negative relationship between within-archipelago inter-island isolation and Island S_{ROA} (Parent & Crespi

2006). Isolation between islands can also enhance or limit diversification by modulating gene flow,

promoting or preventing between-island speciation (Hamilton & Rubinoff 1963, Grant & Grant 2007).

Environmental heterogeneity is expected to positively affect the radiation processes as it links speciation

rate to two critical island features: (1) the elevation gradient, which serves as a proxy for climatic

gradients and habitat diversity, and often promotes speciation by providing ecological opportunities that

drive species adaptation (Stuessy et al. 2006), and (2) the topographic complexity that promote within-

island allopatry (Whittaker et al. 2008, Kisel & Barraclough 2010). A positive relationship between

environmental heterogeneity and Island S_{ROA} is therefore expected. Island geological age might also play a

fundamental role in controlling Island S_{ROA}. Age has a direct effect on Island S_{ROA} as older islands should

(all else being equal) host more species because they have had more time for speciation (Parent & Crespi

2006). However, factors such as area and environmental heterogeneity vary over time following the

ontogeny of the islands comprising an archipelago (Whittaker et al. 2008). As they emerge, oceanic

islands following typical hotspot dynamics rapidly reach their maximum size and elevation. They then

begin to erode and subside, until they become smaller and eventually founder. Topographic complexity

will also generally peak during the middle age of the island. As we expect Island S_{ROA} to correlate with

elevation and topographic complexity, age should also have an effect on Island S_{ROA}, which should rise

and fall throughout the ontogeny of each island, as predicted by the General Dynamic Model of oceanic

island biogeography (GDM) (Whittaker et al. 2008, Borregaard et al. 2017).

ROAs are found across a wide array of taxa and archipelagos, raising the question of how the effects of

geo-environmental factors on Island S_{ROA} patterns vary across ROAs. A growing body of evidence has

substantiated the prevalence of the geo-environmental context in shaping diversification patterns of groups

that radiated on oceanic archipelagos (Funk & Wagner 1995, Roman-Palacios & Wiens 2018). It is

therefore reasonable to hypothesize that Island S_{ROA} patterns of ROAs belonging to the same archipelago

should respond to the distinct combination of island properties in broadly similar ways, thus generating

differences in predictors that best explain Island S_{ROA} in different archipelagos. On the other hand, the

intrinsic biological characteristics of the ROAs themselves (such as body size and mobility) could also

explain why some factors hold greater significance than others. For instance, elevation gradients have

been suggested as a primary driver of diversification for several vascular plant ROAs (Itow 1995, Baldwin

& Sanderson 1998), for which a positive correlation between Island S_{ROA} and elevation is expected, as

have been found for Hawaiian Lobeliads (Givnish et al. 2009). However, such a correlation might be less

prevalent, for example, for a number of arthropod ROAs in Hawaii that appear to have diversified within

narrow elevation belts (Hiller et al. 2019).

In this study, we carried out an extensive analysis using a unique dataset that combines Island S_{ROA} values

for a hundred ROAs distributed across several oceanic archipelagos and taxa. The aims of our study were

twofold, to test whether: (1) Island S_{ROA} scales as a function of different island geo-environmental factors,

and (2) the effects of the factors vary depending on the archipelago and taxon to which the ROAs belong.

Specifically, we hypothesized that island geo-environmental factors will scale with the Island S_{ROA} of

ROAs as outlined in our above predictions (i.e., positive relationships with area, elevation and topographic

complexity, negative with inter-island isolation and hump-shaped with geological age). We also

hypothesized that, overall, environmental heterogeneity will be the predominant driver of Island S_{ROA} as it

is known to trigger diversification across archipelagos and taxonomic groups (Givnish 1997, Losos and

Ricklefs 2009, Roell et al. 2021). Lastly, we anticipated that the variation in the contribution of factors

would be explained equally by both archipelagos and taxa, as the effect of a factor on Island S_{ROA} stems

from complex interactions between the dynamics of the archipelago and the biological characteristics of

the ROA itself.

2. Material and Methods

2.1. Data collection

We have restricted our study to oceanic archipelagos of exclusively volcanic origin that have never been

connected to any other landmasses (Triantis et al. 2016). We excluded island systems composed of both

volcanic islands and continental fragments such as the Philippines (Brown et al. 2013). We then

established a list of criteria to determine whether a radiation should be included in our study. First, we

selected terrestrial ROAs with a minimum of 10 species. Although arbitrary, this threshold ensures that

diversification within the selected ROAs has been sufficiently important for different species to

accumulate on several islands within the archipelago, thus providing a minimum of variation in Island

S_{ROA} values for the subsequent analyses (see below). For ROAs that diversified across more than one

oceanic archipelago, we treated each archipelago independently, only keeping those that harbour a

monophyletic clade of at least 10 species (e.g. the *Tarphius* beetle radiation on the archipelagos of

Macaronesia, Amorim et al. 2012). To facilitate comparison between archipelagos, we excluded clades

that reach 10 species only at the scale of the meta-archipelago (e.g., South Pacific, Macaronesia). Second,

within a given archipelago, the ROA needed to be distributed on at least six islands as any lower threshold

is problematic for the regression analyses (Matthews et al. 2023). This threshold also allows for the

inclusion of Hawaiian ROAs, as most of them are distributed on the six main islands of the archipelago

(i.e. on Kauaʻi, Oʻahu, Molokaʻi, Lānaʻi, Maui and Hawaiʼi; see further details on the island selection

procedure for Hawaii in **Appendix 1**). On the other hand, oceanic archipelagos such as Madeira and the

Mascarenes, both comprising only 3 islands, were discarded. Third, the ROA should be taxonomically

coherent, i.e. all species are from the same genera or from multiple genera that together formed a clade

from which the monophyly has been validated by a phylogenetic reconstruction.

To guide our literature search, we first made a list of the major oceanic archipelagos that contain at least

six main islands (**Table S1**) and then performed a targeted search of ROA studies with ISI Web of

Knowledge and Google Scholar indexing systems (last access December 2022), using archipelago names

in combination with "island", "phylog*", "radiation" and "endemic" strings. Additionally, we also

checked the references list of both the studies we retrieved and some key reviews on the topic (e.g. Shaw

& Gillespie 2016, Gillespie et al. 2020, de La Harpe et al. 2017, Florencio et al. 2021, Hembry et al. 2021)

to make sure that no relevant ROA had been overlooked. Overall, a total of 105 ROAs distributed across

nine oceanic archipelagos were recovered from the literature, Hawaii being by far the most represented of

the four with 53 ROAs, followed by the Canary Islands (29), Galápagos (7) and Fiji (6), Society (4),

Marquesas (3) and one for Azores, Cabo Verde and Austral Islands (**Figure 1A**). Details on the discarded

archipelagos and ROAs are provided in **Table S1** and **Table S2**.

For each selected ROA, we created an occurrence matrix with islands as rows and species as columns for

all the species known to belong to the ROA in the focal archipelago, using the most recent taxonomy and

sampling published works we found in the literature. When the information was available, we included

globally and locally (i.e., species extirpated from individual islands) extinct species. We also created an

alternative version of the matrix by including sub-species where they are distinguished, since some

authors consider subspecies to be a more appropriate evolutionary unit than species, particularly in the

island context (Lasky et al. 2017). The number of species per island was then calculated for each ROA

separately to form the Island S_{ROA} vector. For each ROA, we recorded four distinct features: (1) the

archipelago it belongs to, (2) taxonomic information (Order, Family, Genus), (3) ROA size (i.e. number of

species) and (4) proportion of Single Island Endemic (SIE) species in the ROA. Furthermore, we created a

variable for Taxa where each ROA was classified as either invertebrate, vertebrate or plant. This

classification is used here as a coarse proxy for biological characteristics, as previously done in similar

contexts (Triantis et al. 2012, Matthews et al. 2016, 2019). Although variability within each taxon may

add noise to the analyses, we chose not to use lower ranks in order to maintain sufficient sample sizes. All

occurrence matrices are available in **Supplementary Online Data**. The list of the selected ROAs, along

with their characteristics and original literature resources is available in **Table S3**.

Figure 1. (A) World map indicating the location of the nine archipelagos for which at least one ROA (radiation on oceanic archipelago) fitting our criteria was retrieved. Pie charts showing the proportion of the main orders to which the radiations belong. The four archipelagos analysed in the main text are those with coloured, thick borders and depicted by a map. All together, these four archipelagos represent 90% (95) of all the ROAs retrieved. (B to E). Distribution of (B) the size of the radiation ($\log_{10}$-transformed number of species, without subspecies) and (C) the proportion of SIE between archipelagos. The results of a Kruskal-Wallis test of the difference in the size of the radiation and the proportion of SIE between archipelagos is given at the top of each panel. Distribution of (D) the size of the radiation and (E) the proportion of SIE for invertebrates (blue) and plants (green) for both Hawaii and the Canary Islands. The results of Wilcoxon Tests of the difference in the size of the radiation and the proportion of SIE between invertebrates and plants within Hawaii and Canary Islands are given at the top of each panel.

2.2. Island characteristics

We used the database of Global Administrative Areas (GADM http://www.gadm.org/version1) to obtain high-resolution present coastlines for all islands included in our dataset. We then gathered information for island area, environmental heterogeneity, geological age and intra-archipelagic island isolation as potential determinants of Island S_{ROA}. Environmental heterogeneity was assessed using two distinct variables: (1) the maximum elevation – used here as a proxy of the climatic gradient and habitat diversity, and; (2) the variation in the rate of elevational change over the horizontal surface – quantified here as the standard deviation of slope (hereafter SDS) – which is a measure of the topographic complexity of an island. The maximum elevation and SDS were calculated based on the 30-m digital elevation model (DEM) Advanced Land Observing Satellite (ALOS) from the OpenTopography Portal (https://opentopography.org). Inter-island isolation was calculated using the neighbor index (NI) of Kalmar & Currie (2006), which postulates that the importance of an island as a potential source of colonization to the focal island is proportional to its area and to the coast-to-coast distance to the focal island. Since the way NI is calculated can give more or less weight to area and distance, we identified three distinct configurations [i.e. (1) involving only distance, (2) area and distance and (3) area and squared distance] and selected the one that maximized the correlation with Island S_{ROA} of each ROA (see further details in **Appendix 1**). Finally, we obtained the geological age of the islands from a literature search. All island characteristics and the associated literature sources are available in **Table S4**. For each archipelago, pairwise correlation between geo-environmental factors was assessed using Pearson's correlation coefficient (**Figure S1**).

2.3. Statistical analyses

All statistical analyses were carried out within the R programming environment (R Development Core Team, 2023). To ensure meaningful between-archipelago comparisons, the analyses below were restricted to the four archipelagos with five or more ROAs (i.e. Hawaii, Canary Islands, Galapagos and Fiji).

Collectively, these four archipelagos account for 90% (95) of all the retrieved ROAs. For the five

remaining archipelagos (i.e. Society, Marquesas, Austral Islands, Azores, and Cabo Verde), which

together represented only 10 ROAs, basic descriptive statistics are given in **Table S5**.

2.3.1 Testing the relationships between geo-environmental factors and Island S_{ROA}

We first aimed to investigate univariate relationships between geo-environmental factors and Island S_{ROA}

for each ROA separately using the 95 ROAs retrieved from Hawaii, Canary Islands, Galapagos and Fiji.

Prior to the analyses, area and elevation were $\log_{10}$ transformed to approximate normality. Univariate

relationships were assessed using generalized linear models (GLM) using a Poisson distribution or a

Negative Binomial distribution (both with a log-link) when overdispersion was detected (Zuur et al. 2009,

O'Hara & Kotze 2010). We classified each relationship as (i) being positive or negative and (ii) significant

or not. Because a total of 95 GLMs were performed for each factor, we also corrected the P-values using

the False Discovery Rate method (FDR, Benjamini & Hochberg 1995) to guard against inflation of Type-I

errors. Applying such a correction could, however, hamper the detection of important but less strong

relationships due to the small number of data points (Borges & Hortal 2009). Therefore, we present the

results both with and without the FDR correction. To evaluate the presence of a hump-shaped relationship

between age and Island S_{ROA}, we performed the GLM with age and its quadratic term (age^2) and

considered the relationship as hump-shaped when it met the three following criteria: (1) the sign of the

slope is positive for the linear term and negative for the quadratic term, (2) the two slopes are significant,

and (3) the inflection point predicted by the fitted model is present within the range of age values

contained in the data (Hortal et al. 2009). To provide an overview of our univariate GLMs, we summed

the number of significant positive or negative univariate relationships across archipelagos and taxa for

each individual geo-environmental factor. For age $+$ age^2, we simply counted how many ROAs presented

a hump-shaped pattern meeting our three criteria.

2.3.2 Testing whether the magnitude of the effect of geo-environmental factors on Island S_{ROA} varies between archipelagos and taxa.

For each geo-environmental factor, we calculated the standardized β coefficient for each relationship established by means of the GLM (above) as a normalized measure of the strength of the effect of each individual factor on Island S_{ROA}. The higher or lower the value of the coefficient, the greater the positive or negative effect, respectively. We further computed analyses of variance (ANOVA) for each factor using the standardized β coefficient as the response variable, the archipelago and the taxon to which the ROA belongs to as predictors along with the proportion of SIEs in the ROA and the size of the ROA. For the specific case of age $+$ age^2, we used instead a binomial GLM with a binary response variable with 1 if S_{ROA} follows a hump-shaped relationship with age and 0 if not. All models were subsequently reduced using backward selection. Initially, we did not include any interaction terms, as the archipelago and taxon variables were both highly unbalanced. Therefore, to test for interactions, we reduced our dataset to only Hawaii and the Canary Islands and to plants and invertebrates in order to obtain a more balanced design. We then reran the ANOVAs as detailed above. We then reran the ANOVAs, testing for interactions between archipelago and taxon only, because preliminary tests for the other potential interactions did not indicate significance.

2.3.3 Quantifying the relative contribution of the geo-environmental factors in explaining Island S_{ROA} variations between archipelagos and taxa

Given that none of the factors is expected to fully account for Island S_{ROA} variation on its own, and that a combination of several (if not all) factors is probably necessary (Roell et al. 2021), we computed Generalized Linear Mixed Models (GLMM) to identify the best set of factors for Island S_{ROA} across ROAs for each of the four archipelagos separately. For Hawaii and the Canary Islands only, we also reran the GLMMs for plant and invertebrate subsets. GLMMs were performed with a Poisson distribution or a negative binomial distribution when overdispersion was detected. The GLMMs were implemented with all

geo-environmental factors, including the quadratic term for age to account for potential hump-shaped

relationships. We constructed each GLMM by combining all the Island S_{ROA} values belonging to an

archipelago (or taxon within an archipelago) in such a way that if, for a given archipelago of six islands,

we have recovered 10 ROAs that each radiated over the six islands, the length of the response variable in

our GLMM will be 60. ROA identity was added as the random intercept as well as a random slope for all

the factors to allow the intercepts and the slopes of the relationships to vary between ROAs. Because the

same island appears multiple times in the model, island identity was also included as a separate random

effect (intercept only).

The GLMMs were fitted using the R package *glmmTMB* (Brooks et al. 2017), with all factors scaled to

zero mean and unit variance. We used a top-down strategy for model selection (Zuur et al. 2009). First,

the best random effect structures, with all fixed effects considered, were selected using the *buildglmmTMB*

function in the R package *buildmer* (Voeten 2023). Next, we used the *dredge* function in the *MuMIn*

package (Bartoń 2023) to fit all possible combinations of fixed effects and ranked the models using their

value of AIC corrected for sample size (henceforth AIC_c). We considered all models with $\Delta AIC_c < 2$ (the

difference between each model's AIC_c and the lowest AIC_c) as best models. Note that the term age^2 was

included in a model only where age was included. For each best model we used the *squaredGLMM*

function to compute the conditional R^2 (R^2c) (Nakagawa & Schielzeth 2013), partitioned into the marginal

R^2 (R^2m), representing the variance explained by fixed factors and the random R^2 ($R^2r = R^2c - R^2m$),

representing the variance explained by the random factors. Finally, we derived a measure of variable

importance by summing up AIC_c-weighted standardized regression coefficients for all best models and by

scaling them to sum to one for comparison across archipelagos and taxa (Weigelt et al. 2016).

3.Results

None of the analyses carried out using subspecies data showed any specific differences from our main results. We therefore present the former only in the Supporting Information (specifically, **Tables S6-S8** and **Figure S2)**.

3.1. Radiation characteristics

The 95 retrieved ROAs from Hawaii, Canary Islands, Galapagos and Fiji covered a large taxonomic range including 30 orders and 64 families (**Figure 1A** and **Table S3**). They were largely dominated by invertebrates (60%: 58 ROAs) and plants (37%: 35 ROAs, all being vascular plants), with only three vertebrates ROAs including two birds (Hawaiian Honeycreepers and Galápagos finches) and one reptile (Galápagos Chelonoidis tortoises) **(Figure 1A)**. 73 ROAs (about 77%) had fewer than 50 described species. Conversely, only six had a hundred or more species, five being in Hawaii and one in the Canary Islands (**Figure 1B** and **Table S3**). Overall, the difference in ROA size between archipelagos was marginally non-significant (**Figure 1B**). The majority of ROA constituent species were SIE (median = 71%) and this proportion did not vary significantly between archipelagos (**Figure 1C**). For both Hawaii and the Canary Islands, where comparisons between plants and invertebrates were possible (**Figure 1D, E**), we found that invertebrate ROAs in Hawaii were on average larger and with a higher proportion of SIEs than plant ROAs, while no specific differences were found between taxa for the Canary Islands.

3.2. Univariate relationships between geo-environmental factors and Island S_{ROA}

The results of the univariate GLMs before applying the FDR correction revealed significant linear relationships between Island S_{ROA} and elevation for about 44% of the ROAs, followed by inter-island isolation and area (both 39%), SDS (14%) and age (12%). We also identified 14% of ROAs showing a hump-shaped relationship with age fitting our criteria (**Figure 2**). Notably, most of the observed significant relationships with area, elevation, SDS and inter-island isolation had a sign aligned with our predictions (see a selection of relationships in **Figure 3**; all univariate plots are shown in **Supplementary**

Online Data). Specifically, 100% of the significant relationships with area, 98% with elevation, and 70%

with SDS were positive, while 92% of the significant relationships with inter-island isolation were

negative. For age, eight of the 11 significant relationships recovered were positive. As importantly, the

proportion of significant relationships between Island S_{ROA} and the factors appears to vary considerably

from one archipelago to another (**Figure 2**). For example, in Hawaii, a higher proportion of ROAs

exhibited significant relationships with area compared to the Canary Islands (37% and 20% of the

relationships respectively), while all Fiji's ROAs scaled significantly with area. Furthermore, the

proportion of ROAs in the Canary Islands showing significant relationships with elevation was higher

than in Hawaii (62% and 32% respectively). In addition, significant relationships with inter-island

isolation and a hump-shaped relationship with age were more prevalent among ROAs in the Canary

Islands than in all other archipelagos (**Figure 2**). In contrast, differences between taxa within each

archipelago were less pronounced (**Figure 2**). We noticed a higher proportion of significant relationship

with area for Invertebrates in Hawaii and higher proportion of significant relationships with elevation for

plants in the Canary Islands (**Figure 2**). Overall, similar trends were obtained after FDR correction of the

P-values, but with a substantial decrease in the proportion of significant relationships from a 15%

decrease for Elevation to 85% for SDS (**Figure 2**).

325
326
Figure 2. Summary of the results of the univariate GLMs between Island S_{ROA} and each geo-
environmental factor separately. The coloured part within each pie chart shows the proportion of
radiations having a significant relationship between Island S_{ROA} and a given factor. Because of multiple
testing, the *P*-values of each individual GLM was corrected using the False Discovery Rate method (FDR,
Benjamini & Hochberg 1995) to guard against inflation of Type-I errors. Therefore, the coloured part
represents the proportion of relationships that were significant before FDR correction (the first number in
parentheses above the pie chart) while the dark-shaded part shows the proportion of significant
relationship that remained significant after FDR (false discovery rate) correction (i.e. second number in
parentheses above the pie chart). The light-shaded part therefore represents the proportion of relationships
that were significant only before FDR correction. Specifically for age + age^2, only the proportion of
radiations having a hump-shaped relationship with age that met our three criteria is shown (See Materials
and Methods). For each factor, the four pie charts on the right show the proportion of significant
relationships for each archipelago and for the plant and invertebrate subsets for Hawaii and the Canary
Islands.

Figure 3. Selection of significant relationships between Island S_{ROA} (species richness of radiation on oceanic archipelago) and the geo-environmental factors. One radiation was selected for each factor, namely log_{10}(area), age, age+age^2, log_{10}(elevation), SDS (standard deviation of slope) and isolation. Dots indicate islands and the black curves show the fitted values extracted from the Poisson / Negative binomial generalized linear model. All relationships were significant at 5% after the FDR correction. Plots for all ROAs and all factors are available in **Supplementary Online Data**.

Table 1. Results of the ANOVAs testing whether the strength of the effect of the geo-environmental factors on the Island S_{ROA} can be explained by
archipelago, taxa, radiation size ($\log_{10}$-tranformed), and proportion of SIE. The effect of a given geo-environmental factor on the Island S_{ROA} of a
given ROA is quantified by its standardized β coefficient for the factors area, age, elevation, standard deviation of slope (SDS) and inter-island
isolation while the effect of age + age^2 is coded as a binary variable with a value of 1 if S_{ROA} follows a hump-shaped relationship with age
according to our three criteria (See main text) and 0 if not. For the age + age^2 model, the model is fitted with a binomial GLM. The R^2 is given for
each model and the *P*-value is provided for each predictor remaining in the model after backward elimination procedure. Analyses were initially
performed for all radiations without testing the interaction between Archipelago and Taxa owing to the design being unbalanced. The models with
the interaction term were performed for a subset of the data including only the radiations of Hawaii and the Canary Islands and for plants and
invertebrates. Note that archipelago and taxa were kept in the model when their interaction was significant even if their independent effect was not.
Results of the post-hoc tests are given in **Table S9**. For the predictors Size and Proportion of SIE, the sign of the relationship is given in
parentheses. Significant effects are marked in bold.
358

				Predictors			
	Factors	Archipelago	Taxa	A x T	Radiation Size	Proportion of SIE	R^2
All radiations							
Standardized β coefficient	Area	**<0.001**	**0.026**	-	**<0.001 (-)**	**<0.001 (+)**	0.541
	Age	**0.031**		-			0.090
	Age + age2	**0.001**		-			0.288
	Elevation	**<0.001**		-	**<0.001 (-)**	**0.007 (+)**	0.425
	SDS	**<0.001**	**<0.001**	-			0.527
	Inter-island isolation	**<0.001**	**0.018**	-			0.227
Subset							
Standardized β coefficient	Area		**0.037**		**<0.001 (-)**	**0.004 (+)**	0.231
	Age	0.561	0.901	**0.003**		**0.031 (+)**	0.138
	Age + age2	**0.007**					0.146
	Elevation	**0.015**	0.730	**<0.001**	**<0.001 (-)**		0.318
	SDS	**0.003**	**<0.001**				0.267
	Inter-island isolation	**<0.001**	**0.044**	**0.034**			0.314

3.3. Testing whether the magnitude of the effect of geo-environmental factors on Island S_{ROA} varies between archipelagos and taxa.

The differences observed between archipelagos in the proportion of significant relationships between factors and Island S_{ROA} were confirmed by the ANOVAs carried out to identify predictors of the strength of the effect of the factors on Island S_{ROA} (here quantified by the standardized β coefficient). Indeed, the ANOVAs explicitly demonstrated the importance of archipelago in explaining the strength of the effect of all factors on Island S_{ROA} (**see Table 1**), with post hoc tests revealing consistent pairwise differences between archipelagos as previously mentioned (see **Table S9**). To a lesser extent, the taxon also predicted the strength of the effects of area, SDS and isolation, with the effect of area and inter-island isolation being more pronounced for invertebrates than for plants while the opposite was true for SDS. Interestingly, we also found a negative effect of radiation size on the strength of effects of area and elevation, while the proportion of SIE exhibited a positive relationship with the same two factors (see **Table 1**). Similar outcomes were obtained when we conducted the analyses with a subset including only Hawaii and the Canary Islands and restricted the analyses to plants and invertebrates (**Table S9**). As importantly, these last analyses unveiled significant interactions between archipelagos and taxa for age, elevation and isolation, supporting the fact that the difference between taxa in the effect of these factors was archipelago dependent (see **Table 1** and **Table S9**).For instance, for Canarian ROAs, the effect of elevation was stronger for plants than for invertebrates but did not differ between the two taxa in Hawaii (**Table S9**).

3.4 Quantifying the relative contribution of the geo-environmental factors in explaining Island S_{ROA} variations between archipelagos and taxa

Figure 4 presents the results of the GLMMs (run separately for each archipelago) including factor contribution (**Figure 4A**), the R^2 (**Figure 4B**) and averaging standardized estimates (**Figure 4C**). The GLMMs shed further light on the role of the archipelago and the taxon in relation to the effects of geo-

environmental factors on Island S_{ROA}. For Hawaii, we recovered a positive effect of area and SDS, with an

averaged R^2m (fixed effects only) accounting for approximately 40% of the variation in Island S_{ROA}. For

the Canary Islands, best models explained 32% of the variance (R^2m) and included a humped-shaped

relationship with age, a positive effect of elevation and a negative effect of isolation. In contrast, no

specific factor emerged as significant in the Galápagos GLMMs and the model selection procedure

exhibited high uncertainty (**Table S10**) with only a small fraction of the variance explained ($R^2m = 19\%$).

For Fiji, we identified a humped-shape relationship with age and a positive effect of elevation with an

R^2m of 70%. When performed solely for plant ROAs, the analyses highlighted a strong positive effect of

elevation for both Hawaii and the Canary Islands while a humped-shape relationship with age was

detected for Hawaiian plant ROAs. Overall, the plant ROA models exhibited a R^2m of 46% for both

archipelagos. For invertebrate ROAs, the results for Hawaii emphasised the importance of area and SDS

with an average R^2m of 50%, while for the Canary Islands, humped-shape relationship with age, a positive

effect of elevation and a negative effect of inter-island isolation were identified with an average R^2m of

30%. Importantly, for most archipelagos and subsets, the variance captured by the random structure was

high, reaching 47% for Galápagos. Further details of these analyses, including the best random structures

and family distribution are given in **Table S6.**

Figure 4. (A) The relative importance of each fixed effect as a predictor of Island S_{ROA} for all ROAs in each archipelago, followed by the plant and invertebrate subsets for Hawaii and the Canary Islands; (B) decomposition of the variance explained by the best set of models (conditional R^2c) into its components assigned to the fixed effects (marginal R^2m, dark grey) and random effects (R^2c - R^2m, light grey) for each subset; (C) standardized coefficients of the effect of the factors on Island S_{ROA} variation and their associated confidence intervals, extracted from the model constructed by averaging over the best models identified for each archipelago and for the plant and invertebrate subsets. Solid dots indicate a significant effect, while empty dots indicate no effect. Details for the best models are in **Table S10**.

4. Discussion

Despite the large amount of work undertaken on ROAs in recent decades, little research has focused on

the emergent diversity patterns, with the few studies that have done so focusing on the effect of island area

and geological age (e.g. Lim & Marshall 2017, Gillespie & Baldwin 2010; but see Parent & Crespi 2006

and Roell et al. 2021). In this study, we have examined how the Island S_{ROA} of almost a hundred ROAs

scale with five major geo-environmental factors (area, age, elevation, topographic complexity and intra-

archipelago isolation), and whether the magnitude of their effects was archipelago and taxon specific.

4.1 A brief overview of the selected ROAs

Our dataset reveals a clear dominance of ROAs from Hawaii and the Canary Islands in the literature that

qualify for our basic criteria for inclusion (e.g. purely volcanic, oceanic islands; minimum of 6 islands,

etc.). This was unsurprising since these two archipelagos are known to host a tremendous number of

successful ROAs (e.g., Machado et al. 2017, Hembry et al. 2021, Cerca et al. 2023). These archipelagos

are also two of the best studied volcanic island systems in the world, both in terms of their natural history

and the phylogenetic origin of their biota (e.g., Caujapé-Castells et al. 2017, Price & Wagner 2018). On

the other hand, the Galápagos, which are more ecologically uniform and geologically younger (Naciri &

Linder 2020), are home to smaller ROAs, many of which number fewer than 10 species (Parent et al.

2008) and were thus excluded from our study. The few ROAs retrieved for Fiji can be explained by the

overall less advanced phylogenetic understanding of Fijian endemic groups compared to the other three

archipelagos (Rudbeck et al. 2022).

The study of evolutionary radiations invariably raises the question of whether they are adaptive or not.

Despite recent, albeit debated, classifications underlining the dominance of adaptive radiation within

ROAs (Schenk 2021, Cerca et al. 2023), we decided to not apply such a classification in our data.

Actually, for several ROAs, no indication of the mode of speciation is available in the literature and more

importantly, the lack of consensus on what an adaptive radiation is prevents us from drawing robust

conclusions on this issue (e.g., Hernández-Hernández 2019). In particular, it seems that the role of non-

adaptive processes is often neglected, even though they could potentially explain the bulk of ROA

speciation events (Dimitrov et al. 2008, Benavides et al. 2008, Groom et al. 2013, Vitales et al. 2015,

Chiba & Cowie 2016, Barajas-Barbosa et al. 2023, Illera et al. 2024; see also discussion in Whittaker et al.

2023).

4.2. Island geo-environmental factors, and specifically elevation, are effective drivers of Island S$_{ROA}$

Our study confirms our first hypothesis that the selected geo-environmental factors were overall good predictors of Island S$_{ROA}$, but they were not all equally important. Indeed, most of the significant relationships recovered were attributed, in a decreasing order, to elevation, inter-island isolation and area with a sign in line with our predictions, i.e. positive for area and elevation and negative for inter-island isolation. In this respect, these results corroborate what is usually reported in studies on island species richness of larger groups (vascular plants, Kreft et al. 2008; spiders, Cardoso et al. 2010; birds, Kalmar & Currie 2006). The predominance of elevation in our analyses confirms our prediction that environmental heterogeneity is a key driver of Island S$_{ROA}$. This is further supported by the high proportion of SIE detected in the ROAs (median 71%), confirming the role of within-island speciation and underscoring the importance of environmental heterogeneity in the radiation process (Cowie & Holland 2008). The elevation gradient often serves as a proxy of habitat diversity and is closely associated with adaptive processes in ROA diversification (e.g., Givnish 1997, Schluter 2000). Indeed, several radiated species in many different ROAs have developed into locally adapted ecotypes in response to elevational gradients (Jorgensen & Olesen 2001 and see example below). High-elevation zones can also provide refugia during periods of climate change, allowing relict populations of ancient species to persist (Irl et al. 2015). Elevation can also act as a direct surrogate of topographic complexity, as high elevation is a prerequisite for the presence of a complex landscape structure promoting within-island allopatry (Brown et al. 2006). In this context, it is worth highlighting that the ROAs exhibiting a strong relationship between Island S$_{ROA}$ and elevation were also the one with the highest proportion of SIE (**Table 1**). Note that the SDS factor, which quantifies topographical complexity *per se* in our analyses, exhibited very few significant relationships with Island S$_{ROA}$. This was somewhat anticipated, since its effect seems to become apparent only after controlling for other factors such as area or age, as reported by Roell et al. (2021) and Barajas-Barbosa et al. (2020). Most significant relationships between Island S$_{ROA}$ and inter-island isolation

revealed that, as expected, the most remote islands in the archipelago were the least diverse. However,

inter-island isolation did not affect Island S_{ROA} for 60% of the ROAs. In some cases, the effects of inter-

island isolation may have been obscured by subsequent within-island allopatry or back-colonization

following inter-island speciation (Cowie & Holland 2008). Additionally, the high vagility of certain ROA

species may have prevented geographic speciation between islands, with radiation in these cases resulting

primarily from sympatric speciation (Losos & Ricklefs 2009). It is important to note that we focused on a

single scale of isolation—island isolation within the archipelago—because our primary goal was to test

the effect of island-specific characteristics on Island S_{ROA}. However, we should not exclude the possibility

that isolation may operate at other scales. Isolation of the archipelago itself could indirectly impact Island

S_{ROA} by affecting the overall size of the ROA (Tao et al. 2021). Within-island isolation, i.e., the isolation

between populations on the same island, will also play a crucial role in determining Island S_{ROA} by driving

allopatric speciation, although we expect topographic complexity to capture such mechanisms to some

extent (Barajas-Barbosa et al. 2020).

The weak effect of age on Island S_{ROA}, whether considered through a linear or hump-shaped relationship,

may stem from the fact that age tends to have its strongest influence when interacting with area (Whittaker

et al. 2008). This aligns with the idea that the impact of time on speciation cannot be fully decoupled from

the amount of ecological space available for it as postulated by the GDM (Whittaker et al. 2008,

Borregaard et al. 2017). Moreover, our results may also show the limitation of assigning a single age to an

island, given that many of the oceanic islands we considered are formed by different basaltic shields of

varying ages (Irl et al. 2015, Otto et al. 2016). It may also be noted that for much of the last 1M years, the

islands of Lanzarote and Fuerteventura have been united by lower eustatic sea-levels to form a single

island, thus arguably conveying the biological legacies of Fuerteventura's greater age (23 Ma) on the

island of Lanzarote (15 Ma). Were the age of 23 Ma used for both islands this would make a stronger case

for a humped-relationship between Island S_{ROA}

Finally, the above results should be viewed with caution due to the substantial correlation between geo-

environmental variables observed in most of the archipelagos studied (Figure S1), which may lead to

confounding effects.

4.3. Archipelago and taxon determine the strength of the effect of geo-environmental

factors on Island S_{ROA}

The Island S_{ROA} of a ROA on a particular island is the outcome of many processes affecting the

colonization, evolution, and persistence of island populations. One important finding of our study is to

show that the expression of these processes is likely to differ among archipelagos because of differences

in their geo-environmental dynamics (e.g., island chronosequence, habitat diversity, spatial configuration)

but also among taxa due to the variation in biological attributes among ROAs (e.g., dispersion, adaptive

capacity, population density) (**Table 1** and **Figure 2** and **4**). While other authors have already highlighted

an archipelagic and/or taxonomic dependence in the factors driving diversity patterns of ROAs (e.g.,

Parent 2012, Lim & Marshall 2017), our study is the first to quantify such effects across multiple

archipelagos and taxa via a comparative statistical approach.

Taken together, variation in Island S_{ROA} of Hawaiian ROA were mostly driven by area and topographic

complexity (SDS). The explanation for this likely lies in the role of the Big Island in driving diversity

patterns in Hawaii. Indeed, our results and several other studies (e.g. Price & Wagner 2011, Eckstut et al.

2011) show that for most ROAs, Big Island has a diversity value lower than predicted by the ISAR with

the other Hawaiian Islands (**Supplementary Online Data**), reflecting that the island is both the largest

and the youngest island (0.53 Ma, **Table S4**) in origin. In our case, this deviation is captured by the

comparatively lower topographic complexity of Big Island, indicative of its smooth and young terrain,

compared to the older and more rugged islands (Price & Wagner 2011, Barajas-Barbosa et al. 2020).

Given that Hawaii hosts some of the largest ROAs, and with the highest proportion of SIEs (Figure S3),

we cannot rule out that the positive effect of SDS on Island S_{ROA} may reflect the importance of species

accumulation through within-island allopatry during the radiating process (Carson & Kaneshiro 1976,

Hembry et al. 2021). In contrast to Hawaii, isolation, elevation and age (hump-shaped) appear to shape the

Island S_{ROA} patterns for the Canary Islands (**Figure 4**). The predominance of a negative effect of inter-

island isolation reveals that species accumulation was higher in the central, least isolated islands within

the archipelago, namely Tenerife and Gran Canaria. This aligns with previous studies that have positioned

these two islands as primary centers of diversification (e.g. Francisco-Ortega et al. 2002, Allan 2004,

Fernández-Mazuecos & Vargas 2011), making them the main sources of dispersal for the peripheral

western islands. Using a Bayesian framework to infer dispersion processes within the archipelago,

Sanmartín et al. (2008) indeed identified a stepwise dispersal pattern from the central islands to the islands

of La Palma and El Hierro in the west. This distinctive feature of these two central islands relies also on

their intrinsic characteristics, such as elevation (3,715 and 1,956 m.a.s.l respectively) and topographic

complexity, both resulting from their intermediate age (12 and 14.5 Myr respectively) relative to the age

of the archipelago (i.e., 23 Myr). According to the GDM, intermediate-age islands reach 'maturity' through

the slowing down of island-building processes in favour of erosive processes, generating a more complex

topography (Borregaard et al. 2017). In that respect, we found that two-thirds of the ROAs of Canary

Islands had their highest values of Island S_{ROA} and SIE in Tenerife (see **Figure S4**). This relationship with

age was reflected in the importance of the hump-shaped relationship detected for the Canary Islands

(**Figure 4**), with young and old islands having low Island S_{ROA} values compared to the central ones.

However, the low Island S_{ROA} of the two oldest islands (i.e., Lanzarote and Fuerteventura) are not simply

or exclusively the result of the island aging process as predicted by the GDM. It also reflects that these

islands underwent climatic fluctuations during the Pleistocene followed by a recent onset of an arid

climate that wiped out species not adapted to these novel conditions, thereby reinforcing the hump-shaped

relationship with age (García-Verdugo et a. 2019).

For both Hawaii and the Canary Islands, the two archipelagos for which a comparison between vascular

plants and invertebrates ROAs was possible, taxon also appears to wield a significant role in driving the

effect of the geo-environmental factors on Island S_{ROA}. Probably the most striking finding here was the

strong effect of elevation detected for the plant subset of both archipelagos. Indeed, several emblematic

plant adaptive ROAs span the entire elevation gradient of islands, with different species each adapted to narrow habitats, ranging from dry coastal areas to mesic environments and wet forests [e.g. Lobeliads (Givnish et al. 2009), Silversword alliance (Baldwin & Sanderson 1998), *Schiedea* (Sakai et al. 2006) and *Euphorbia* (Yang et al. 2018) in Hawaii; *Aeonium* Alliance (Kim et al. 2008, Messerschmid et al. 2023), *Argyranthemum* (White et al. 2020), *Echium* (García-Maroto et al. 2009), *Sonchus* (Kim et al. 2008) in the Canary Islands]. In contrast, the relative contribution of elevation for invertebrates (**Figure 4**) was very limited in both archipelagos. Several studies suggest that diversification in many arthropod ROAs on oceanic islands has been primarily driven by niche partitioning resulting from biotic interactions, such as microhabitat choice or host plant specialization, rather than by the island ecological/climatic gradients (Bennet & O'Grady 2012, Machado et al. 2017, Johns et al. 2018, Cotoras et al. 2018). A recent study on Hawaiian radiations even indicated that strong environmental conservatism dominates for arthropod ROAs as they largely occur within restricted environmental envelopes (Hiller et al. 2019, Dorey et al. 2020).

The best models for Fiji highlighted the pivotal roles of age and elevation in shaping the diversification of Fijian ROAs. Firstly, all Fijian ROAs included in our study reached their maximum Island S_{ROA} on Viti Levu, the oldest island of Fiji (**Figures S4** and **Supplementary Online Data**). The observed effect of age + age² detected in our GLMM was mainly due to a non-linear increase of Island S_{ROA} with age, rather than a true hump-shaped relationship (**Figure 4** and **Supplementary Online Data**). Secondly, studies examining the presence of a taxon cycle in Fijian ants, in particular the *Strumigenys* and *Pheidole* ROAs, supported the hypothesis that the radiation process was primarily associated with a niche shift towards higher elevations (Sarnat & Economo 2012, Liu et al. 2020), with species accumulation being more pronounced on high-elevation islands. Moreover, diversification of *Cyrtandra* in Fiji is thought to have been triggered by the pronounced elevational gradient with several small-range species of the rainforest understory being distributed from the lowlands to the montane zone (Johnson 2017, 2023). Conversely, the absence of significant signal for Galápagos ROAs could be due to the wide taxonomic range covered

by the seven retrieved ROAs (two beetles, one moth, one bird, one reptile, one land snail and one

flowering plant ROA), which might have hindered the identification of common drivers.

Finally, it should be noted that the random structure of our GLMMs, mostly the random effect of radiation

identity for slope and intercept (**Table S6**), occasionally demonstrated an explanatory power greater than

the geo-environmental factors in explaining Island S_{ROA} variation (**Figure 4B** and **Table S6**). Analytically,

this suggests that while a given geo-environmental factor may influence most radiations similarly within

an archipelago (e.g. a positive relationship between elevation and Island S_{ROA} in the Canary Islands,

Figure 4), the magnitude of its effect can vary significantly between radiations. This finding is important

because this variance likely reflects finer differences between ROAs, particularly in terms of their

ecological and evolutionary characteristics (e.g., dispersal abilities, speciation rates), which are not

accounted for by our fixed effects. Overall, this highlights the need for a deeper understanding of how

ROA life history traits and geo-environmental factors interact to influence the radiation process.

4.4. Limitations

Our approach undoubtedly comes with some limitations. First, our values of Island S_{ROA} may be

potentially biased due to unrecorded human mediated species extinctions or local (islands) extirpations

(Paulay 1994, Lim & Marshall 2017, Whittaker et al. 2023). Moreover, we cannot exclude the possibility

that inconsistencies among taxonomists and the varying degrees of completeness of these endemic clades

may have generated noise in our analyses. However, the analyses that included subspecies did not show

any specific differences with the analyses run without subspecies, suggesting that our results may be

robust to marginal variation in species number (**Table S7, S8** and **Figure S2**). Second, we acknowledge

that the limited number of islands within each archipelago poses a challenge in terms of statistical power

and restricts the use of complex models. Third, the ten species threshold we chose to ensure sufficient

variance in Island S_{ROA} between islands introduces a bias toward "successful" ROAs, and our conclusions

therefore might not apply for more modest ROAs.

4.5. Conclusion & Perspectives

Our study confirms that geo-environmental factors are effective predictors of Island S_{ROA}, aligning with previous biogeographic expectations. Therefore, it seems that the main abiotic forces driving overall island biodiversity also govern the diversification of endemic clades. Elevation emerged as a key factor, complementing existing theoretical and empirical research that has highlighted the central role of environmental heterogeneity in driving diversification on oceanic islands (e.g., Cabral et al. 2019, Roell et al. 2021). The influence of geo-environmental factors also varied by archipelago and taxon, indicating that unique archipelagic dynamics and biological characteristics channel diversification differently. Future analyses should explore the contribution of additional factors such as climate change and variability (e.g., Barajas-Barbosa et al. 2020) as well as direct measurements of habitat diversity (Yu et al. 2015). Moreover, accounting for palaeo-configuration (Norder et al. 2018) may be particularly relevant in the context of ROAs, as variation in archipelago configuration over geological time is likely to have left detectable evolutionary legacies in contemporary distributions of endemism (Weigelt et al. 2016, Fernández-Palacios et al. 2016).

Acknowledgments

We thank Delphine Montagne for her help with QGIS. We also thank the organisers of the Island Biology 2023 conference in Lipari, which enabled us to have fruitful live discussions on this work.

Author Contributions

B.B. and F.R. designed the study. K.A.T. helped compile the list of oceanic archipelagos. B.B. collected the data on evolutionary radiations and F.R. collected the data on island characteristics. B.B. and F.R. analysed the data. B.B. and F.R. drafted the manuscript in collaboration with T.J.M., J.M.F.P, C.P., K.A.T. and R. J.W.. All authors contributed to and approved the submitted version.

Funding

Baptiste Brée was funded by a PhD grant Energy Environment Solutions (E2S) supported by the Agence Nationale de la Recherche (ANR-16-IDEX-0002).

Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the content of this article.

Data and code availability

All the data used in this manuscript are available in Supplementary Information and in Supplementary Online Data.

References

Allan, G. J., Francisco-Ortega, J., Santos-Guerra, A., Boerner, E., & Zimmer, E. A. (2004). Molecular phylogenetic evidence for the geographic origin and classification of Canary Island *Lotus* (Fabaceae : Loteae). *Molecular Phylogenetics and Evolution*, *32*(1), 123-138.

Amorim, I. R., Emerson, B. C., Borges, P. A. V., & Wayne, R. K. (2012). Phylogeography and molecular phylogeny of Macaronesian island *Tarphius* (Coleoptera : Zopheridae) : Why are there so few species in the Azores? *Journal of Biogeography*, *39*(9), 1583-1595. https://doi.org/10.1111/j.1365-2699.2012.02721.x

Baldwin, B. G., & Sanderson, M. J. (1998). Age and rate of diversification of the Hawaiian silversword alliance (Compositae). *Proceedings of the National Academy of Sciences*, *95*(16), 9402-9406.

Barajas Barbosa, M. P., Craven, D., Weigelt, P., Denelle, P., Otto, R., Díaz, S., … Kreft, H. (2023). Assembly of functional diversity in an oceanic island flora. *Nature*, *619*(7970), 545-550.

Barajas-Barbosa, M. P., Weigelt, P., Borregaard, M. K., Keppel, G., & Kreft, H. (2020). Environmental heterogeneity dynamics drive plant diversity on oceanic islands. *Journal of Biogeography*, *47*(10), 2248-2260. https://doi.org/10.1111/jbi.13925

Bartoń, K. (2023). *MuMIn : Multi-Model Inference. R package version 1.47.5, <https://CRAN.R-project.org/package=MuMIn>*.

Beatty, C. D., Sánchez Herrera, M., Skevington, J. H., Rashed, A., Van Gossum, H., Kelso, S., &
Sherratt, T. N. (2017). Biogeography and systematics of endemic island damselflies : The
Nesobasis and *Melanesobasis* (Odonata: Zygoptera) of Fiji. *Ecology and Evolution*, *7*(17),
7117-7129. https://doi.org/10.1002/ece3.3175
Benavides, E., Baum, R., Snell, H. M., Snell, H. L., & Sites, Jr., J. W. (2008). Island biogeography
of Galápagos lava lizards (Tropiduridae : *Microlophus*): species diversity and colonization of
the archipelago. *Evolution*, *63*(6), 1606-1626. https://doi.org/10.1111/j.1558-
5646.2009.00617.x
Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate : A practical and powerful
approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*,
57(1), 289-300.
Bennett, G. M., & O'Grady, P. M. (2012). Host–plants shape insect diversity : Phylogeny, origin,
and species diversity of native Hawaiian leafhoppers (Cicadellidae: *Nesophrosyne*). *Molecular*
Phylogenetics and Evolution, *65*(2), 705-717. https://doi.org/10.1016/j.ympev.2012.07.024
Borges, P. A. V., & Hortal, J. (2009). Time, area and isolation : Factors driving the diversification of
Azorean arthropods. *Journal of Biogeography*, *36*(1), 178-191. https://doi.org/10.1111/j.1365-
2699.2008.01980.x
Borregaard, M. K., Amorim, I. R., Borges, P. A. V., Cabral, J. S., Fernández-Palacios, J. M., Field,
R., … Whittaker, R. J. (2017). Oceanic island biogeography through the lens of the general
dynamic model : Assessment and prospect. *Biological Reviews*, *92*(2), 830-853.
https://doi.org/10.1111/brv.12256
Brooks, M. E., Kristensen, K., Van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., …
Bolker, B. M. (2017). glmmTMB balances speed and flexibility among packages for zero-
inflated generalized linear mixed modeling. *The R journal*, *9*(2), 378-400.
Brown, R. M., Siler, C. D., Oliveros, C. H., Esselstyn, J. A., Diesmos, A. C., Hosner, P. A., … Alcala,
A. C. (2013). Evolutionary Processes of Diversification in a Model Island Archipelago. *Annual*
Review of Ecology, Evolution, and Systematics, *44*(1), 411-435.
https://doi.org/10.1146/annurev-ecolsys-110411-160323
Brown, R. P., Hoskisson, P. A., Welton, J.-H., & Baez, M. (2006). Geological history and within-
island diversity : A debris avalanche and the Tenerife lizard *Gallotia galloti*. *Molecular Ecology*,
15(12), 3631-3640.
Cabral, J. S., Whittaker, R. J., Wiegand, K., & Kreft, H. (2019). Assessing predicted isolation effects
from the general dynamic model of island biogeography with an eco-evolutionary model for
plants. *Journal of Biogeography*, jbi.13603. https://doi.org/10.1111/jbi.13603
Cardoso, P., Arnedo, M. A., Triantis, K. A., & Borges, P. A. V. (2010). Drivers of diversity in
Macaronesian spiders and the role of species extinctions : Diversity of Macaronesian spiders.
Journal of Biogeography, *37*(6), 1034-1046. https://doi.org/10.1111/j.1365-2699.2009.02264.x
Carson, H. L., & Kenneth, K. Y. (1976). *Drosophila* of Hawaii : Systematics and Ecological
Genetics. *Annual Review of Ecology and Systematics*, 311-345.
Caujapé-Castells, J., García-Verdugo, C., Marrero-Rodríguez, Á., Fernández-Palacios, J. M.,
Crawford, D. J., & Mort, M. E. (2017). Island ontogenies, syngameons, and the origins and

evolution of genetic diversity in the Canarian endemic flora. *Perspectives in Plant Ecology,*
Evolution and Systematics, 27, 9-22. https://doi.org/10.1016/j.ppees.2017.03.003

Cerca, J., Cotoras, D. D., Bieker, V. C., De-Kayne, R., Vargas, P., Fernández-Mazuecos, M., …
Martin, M. D. (2023). Evolutionary genomics of oceanic island radiations. *Trends in Ecology*
& Evolution, *38*(7), 631-342. https://doi.org/10.1016/j.tree.2023.02.003

Chiba, S., & Cowie, R. H. (2016). Evolution and Extinction of Land Snails on Oceanic Islands.
Annual Review of Ecology, Evolution, and Systematics, *47*(1), 123-141.
https://doi.org/10.1146/annurev-ecolsys-112414-054331

Cotoras, D. D., Bi, K., Brewer, M. S., Lindberg, D. R., Prost, S., & Gillespie, R. G. (2018). Co-
occurrence of ecologically similar species of Hawaiian spiders reveals critical early phase of
adaptive radiation. *BMC Evolutionary Biology*, *18*(1), 100. https://doi.org/10.1186/s12862-018-
1209-y

Cowie, R. H., & Holland, B. S. (2008). Molecular biogeography and diversification of the endemic
terrestrial fauna of the Hawaiian Islands. *Philosophical Transactions of the Royal Society B:*
Biological Sciences, *363*(1508), 3363-3376.

Darwin, C. (1845). *Journal of researches into the natural history and geology of the countries visited*
during the voyage of HMS Beagle round the world (Vol. 7). London: John Murray.

de la Harpe, M., Paris, M., Karger, D. N., Rolland, J., Kessler, M., Salamin, N., & Lexer, C. (2017).
Molecular ecology studies of species radiations : Current research gaps, opportunities and
challenges. *Molecular Ecology*, *26*(10), 2608-2622. https://doi.org/10.1111/mec.14110

Dimitrov, D., Arnedo, M. A., & Ribera, C. (2008). Colonization and diversification of the spider
genus *Pholcus* Walckenaer, 1805 (Araneae, Pholcidae) in the Macaronesian archipelagos :
Evidence for long-term occupancy yet rapid recent speciation. *Molecular Phylogenetics and*
Evolution, *48*(2), 596-614. https://doi.org/10.1016/j.ympev.2008.04.027

Dorey, J. B., Groom, S. V. C., Freedman, E. H., Matthews, C. S., Davies, O. K., Deans, E. J., …
Schwarz, M. P. (2020). Radiation of tropical island bees and the role of phylogenetic niche
conservatism as an important driver of biodiversity. *Proceedings of the Royal Society B:*
Biological Sciences, *287*(1925), 20200045. https://doi.org/10.1098/rspb.2020.0045

Eckstut, M. E., McMahan, C. D., Crother, B. I., Ancheta, J. M., McLennan, D. A., & Brooks, D. R.
(2011). PACT in practice : Comparative historical biogeographic patterns and species-area
relationships of the Greater Antillean and Hawaiian Island terrestrial biotas: Comparative PACT
analyses. *Global Ecology and Biogeography*, *20*(4), 545-557. https://doi.org/10.1111/j.1466-
8238.2010.00626.x

Economo, E. P., Klimov, P., Sarnat, E. M., Guénard, B., Weiser, M. D., Lecroq, B., & Knowles, L.
L. (2015). Global phylogenetic structure of the hyperdiverse ant genus *Pheidole* reveals the
repeated evolution of macroecological patterns. *Proceedings of the Royal Society B: Biological*
Sciences, *282*(1798), 20141416.

Emerson, B. C. (2002). Evolution on oceanic islands : Molecular phylogenetic approaches to
understanding pattern and process. *Molecular ecology*, *11*(6), 951-966.

Fernández-Mazuecos, M., & Vargas, P. (2011). Genetically depauperate in the continent but rich in
oceanic islands : *Cistus monspeliensis* (Cistaceae) in the Canary Islands. *PLoS ONE*, *6*(2),
e17172.

Fernández-Palacios, J. M., Rijsdijk, K. F., Norder, S. J., Otto, R., Nascimento, L. de, Fernández-
Lugo, S., … Whittaker, R. J. (2016). Towards a glacial-sensitive model of island biogeography.
Global Ecology and Biogeography, 25(7), 817-830. https://doi.org/10.1111/geb.12320

Florencio, M., Patiño, J., Nogué, S., Traveset, A., Borges, P. A. V., Schaefer, H., … Santos, A. M.
C. (2021). Macaronesia as a Fruitful Arena for Ecology, Evolution, and Conservation Biology.
Frontiers in Ecology and Evolution, *9*, 718169. https://doi.org/10.3389/fevo.2021.718169

Francisco-Ortega, J., Fuertes-Aguilar, J., Kim, S., Santos-Guerra, A., Crawford, D. J., & Jansen, R.
K. (2002). Phylogeny of the Macaronesian endemic *Crambe* section *Dendrocrambe*
(Brassicaceae) based on internal transcribed spacer sequences of nuclear ribosomal DNA.
American Journal of Botany, *89*(12), 1984-1990. https://doi.org/10.3732/ajb.89.12.1984

Funk, V. A., & Wagner, W. L. (1995). Biogeographic patterns in the Hawaiian Islands. In W. L.
Wagner & V. A. Funk (Éds.), *Hawaiian biogeography : Evolution on a hot spot archipelago*.
Smithsonian Institution Press.

García-Maroto, F., Mañas-Fernández, A., Garrido-Cárdenas, J. A., Alonso, D. L., Guil-Guerrero, J.
L., Guzmán, B., & Vargas, P. (2009). Δ6-Desaturase sequence evidence for explosive Pliocene
radiations within the adaptive radiation of Macaronesian *Echium* (Boraginaceae). *Molecular*
Phylogenetics and Evolution, *52*(3), 563-574. https://doi.org/10.1016/j.ympev.2009.04.009

García-Verdugo, C., Caujapé-Castells, J., Illera, J. C., Mairal, M., Patiño, J., Reyes-Betancort, A., &
Scholz, S. (2019). Pleistocene extinctions as drivers of biogeographical patterns on the
easternmost Canary Islands. *Journal of Biogeography*, *46*(5), 845-859.

Gillespie, R. G. (2016). Island time and the interplay between ecology and evolution in species
diversification. *Evolutionary Applications*, *9*(1), 53-73. https://doi.org/10.1111/eva.12302

Gillespie, R. G., & Baldwin, B. G. (2010). Island biogeography of remote archipelagoes. In *The*
theory of island biogeography revisited (p. 358-387). Princeton University Press Princeton.

Gillespie, R. G., Bennett, G. M., De Meester, L., Feder, J. L., Fleischer, R. C., Harmon, L. J., …
Wogan, G. O. U. (2020). Comparing Adaptive Radiations Across Space, Time, and Taxa.
Journal of Heredity, *111*(1), 1-20. https://doi.org/10.1093/jhered/esz064

Givnish, T. J. (1997). Adaptive radiation and molecular systematics : Issues and approaches. In T. J.
Givnish & K. J. Systema (Éds.), *Molecular evolution and adaptive radiation* (p. 1-54).
Cambridge University Press.

Givnish, Thomas J, Millam, K. C., Mast, A. R., Paterson, T. B., Theim, T. J., Hipp, A. L., … Sytsma,
K. J. (2009). Origin, adaptive radiation and diversification of the Hawaiian lobeliads (Asterales :
Campanulaceae). *Proceedings of the Royal Society B: Biological Sciences*, *276*, 407-416.
https://doi.org/10.1098/rspb.2008.1204

Grant, P. R., & Grant, B. R. (2007). *How and why species multiply : The radiation of Darwin's*
finches. Princeton: Princeton University Press.

Groom, S. V. C., Stevens, M. I., & Schwarz, M. P. (2013). Diversification of Fijian halictine bees : Insights into a recent island radiation. *Molecular Phylogenetics and Evolution*, *68*(3), 582-594. https://doi.org/10.1016/j.ympev.2013.04.015

Hamilton, T. H., & Rubinoff, I. (1963). Isolation, endemism, and multiplication of species in the Darwin finches. *Evolution*, *17*(4), 388-403. https://doi.org/10.1111/j.1558-5646.1963.tb03296.x

Hembry, D. H., Bennett, G., Bess, E., Cooper, I., Jordan, S., Liebherr, J., … O'Grady, P. M. (2021). Insect Radiations on Islands : Biogeographic Pattern and Evolutionary Process in Hawaiian Insects. *The Quarterly Review of Biology*, *96*(4), 247-296. https://doi.org/10.1086/717787

Hernández-Hernández, T. (2019). Evolutionary rates and adaptive radiations. *Biology & Philosophy*, *34*(4), 41. https://doi.org/10.1007/s10539-019-9694-y

Hiller, A. E., Koo, M. S., Goodman, K. R., Shaw, K. L., O'Grady, P. M., & Gillespie, R. G. (2019). Niche conservatism predominates in adaptive radiation : Comparing the diversification of Hawaiian arthropods using ecological niche modelling. *Biological Journal of the Linnean Society*, *127*(2), 479-492. https://doi.org/10.1093/biolinnean/blz023

Hortal, J., Triantis, K. A., Meiri, S., Thébault, E., & Sfenthourakis, S. (2009). Island Species Richness Increases with Habitat Diversity. *The American Naturalist*, *174*(6), E205-E217. https://doi.org/10.1086/645085

Illera, J. C., Rando, J. C., Melo, M., Valente, L., & Stervander, M. (2024). Avian Island Radiations Shed Light on the Dynamics of Adaptive and Nonadaptive Radiation. In C. L. Peichel, D. I. Bolnick, Å. Brännström, U. Dieckmann, & J. S. Rebecca (Éds.), Additional Perspectives on Speciation (p.1-29). Cold Spring Harbor Laboratory Press.

Irl, S. D., Harter, D. E., Steinbauer, M. J., Gallego Puyol, D., Fernández-Palacios, J. M., Jentsch, A., & Beierkuhnlein, C. (2015). Climate vs. Topography–spatial patterns of plant species diversity and endemism on a high-elevation island. *Journal of Ecology*, *103*(6), 1621-1633.

Itow, S. (1995). Phytogeography and ecology of *Scalesia* (Compositae) endemic to the Galápagos Islands. *Pacific Science*, *49*(1), 17-30.

Johns, C. A., Toussaint, E. F. A., Breinholt, J. W., & Kawahara, A. Y. (2018). Origin and macroevolution of micro-moths on sunken Hawaiian Islands. *Proceedings of the Royal Society B: Biological Sciences*, *285*, 20181047. https://doi.org/10.1098/rspb.2018.1047

Johnson, M. A. (2017). Four new species of *Cyrtandra* (Gesneriaceae) from the South Pacific islands of Fiji. *PhytoKeys*, *91*, 105-124. https://doi.org/10.3897/phytokeys.91.21623

Johnson, M. A. (2023). *Phylogenetic and functional trait-based community assembly within Pacific* Cyrtandra *(Gesneriaceae) : Evidence for clustering at multiple spatial scales*. *13*(5), e10048.

Jorgensen, T. H., & Olesen, J. M. (2001). Adaptive radiation of island plants : Evidence from Aeonium (Crassulaceae) of the Canary Islands. *Perspectives in Plant Ecology, Evolution and Systematics*, *4*(1), 29-42.

Juan, C., Emerson, B. C., Oromí, P., & Hewitt, G. M. (2000). Colonization and diversification : Towards a phylogeographic synthesis for the Canary Islands. *Trends in Ecology & Evolution*, *15*(3), 104-109.

Kalmar, A., & Currie, D. J. (2006). A global model of island biogeography. *Global Ecology and*
Biogeography, *15*(1), 72-81. https://doi.org/10.1111/j.1466-822X.2006.00205.x
Kim, S.-C., McGowen, M. R., Lubinsky, P., Barber, J. C., Mort, M. E., & Santos-Guerra, A. (2008).
Timing and Tempo of Early and Successive Adaptive Radiations in Macaronesia. *PLoS ONE*,
3(5), e2139. https://doi.org/10.1371/journal.pone.0002139
Kisel, Y., & Barraclough, T. G. (2010). Speciation Has a Spatial Scale That Depends on Levels of
Gene Flow. *The American Naturalist*, *175*(3), 316-334. https://doi.org/10.1086/650369
Kreft, H., Jetz, W., Mutke, J., Kier, G., & Barthlott, W. (2008). Global diversity of island floras from
a macroecological perspective. *Ecology letters*, *11*(2), 116-127.
Lasky, J. R., Keitt, T. H., Weeks, B. C., & Economo, E. P. (2017). A hierarchical model of whole
assemblage island biogeography. *Ecography*, *40*(8), 982-990.
Lim, J. Y., & Marshall, C. R. (2017). The true tempo of evolutionary radiation and decline revealed
on the Hawaiian archipelago. *Nature*, *543*(7647), 710-713. https://doi.org/10.1038/nature21675
Liu, C., Sarnat, E. M., Friedman, N. R., Hita Garcia, F., Darwell, C., Booher, D., … Economo, E. P.
(2020). Colonize, radiate, decline : Unraveling the dynamics of island community assembly
with Fijian trap-jaw ants. *Evolution*, *74*(6), 1082-1097. https://doi.org/10.1111/evo.13983
Losos, J. B., & Ricklefs, R. E. (2009). Adaptation and diversification on islands. *Nature*, *457*(7231),
830-836. https://doi.org/10.1038/nature07893
Losos, J. B., & Schluter, D. (2000). Analysis of an evolutionary species–area relationship. *Nature*,
408(6814), 847-850. https://doi.org/10.1038/35048558
MacArthur, R. H., & Wilson, E. O. (1963). An Equilibrium Theory of Insular Zoogeography.
Evolution, *17*(4), 373. https://doi.org/10.2307/2407089
MacArthur, R. H., & Wilson, E. O. (1967). *The theory of island biogeography*. Princeton: Princeton
university press.
Machado, A., Rodríguez-Expósito, E., López, M., & Hernández, M. (2017). Phylogenetic analysis
of the genus *Laparocerus*, with comments on colonisation and diversification in Macaronesia
(Coleoptera, Curculionidae, Entiminae). *ZooKeys*, *651*, 1-77.
https://doi.org/10.3897/zookeys.651.10097
Matthews, T. J., Guilhaumon, F., Triantis, K. A., Borregaard, M. K., & Whittaker, R. J. (2016). On
the form of species–area relationships in habitat islands and true islands. *Global Ecology and*
Biogeography, *25*(7), 847-858.
Matthews, T. J., Rigal, F., Triantis, K. A., & Whittaker, R. J. (2019). A global model of island
species–area relationships. *Proceedings of the National Academy of Sciences*, *116*(25),
12337-12342. https://doi.org/10.1073/pnas.1818190116
Matthews, T. J., Wayman, J. P., Whittaker, R. J., Cardoso, P., Hume, J. P., Sayol, F., … Rigal, F.
(2023). A global analysis of avian island diversity–area relationships in the Anthropocene.
Ecology Letters, *26*(6), 965-982. https://doi.org/10.1111/ele.14203
Messerschmid, T. F. E., Abrahamczyk, S., Bañares-Baudet, Á., Brilhante, M. A., Eggli, U., Hühn,
P., … Kadereit, G. (2023). Inter- and intra-island speciation and their morphological and
ecological correlates in *Aeonium* (Crassulaceae), a species-rich Macaronesian radiation. *Annals*
of Botany, *131*(4), 697-721. https://doi.org/10.1093/aob/mcad033

Naciri, Y., & Linder, H. P. (2020). The genetics of evolutionary radiations. *Biological Reviews*,
95(4), 1055-1072. https://doi.org/10.1111/brv.12598

Nakagawa, S., & Schielzeth, H. (2013). *A general and simple method for obtaining R2 from*
generalized linear mixed-effects models. *4*(2), 133-142.

Norder, S. J., Proios, K., Whittaker, R. J., Alonso, M. R., Borges, P. A. V., Borregaard, M. K., …
Rijsdijk, K. F. (2018). Beyond the Last Glacial Maximum : Island endemism is best explained
by long-lasting archipelago configurations. *Global Ecology and Biogeography*, *28*(2), 184-197.
https://doi.org/10.1111/geb.12835

O'Grady, P., & DeSalle, R. (2018). Hawaiian *Drosophila* as an Evolutionary Model Clade : Days of
Future Past. *BioEssays*, *40*(5), 1700246. https://doi.org/10.1002/bies.201700246

O'Hara, R., & Kotze, J. (2010). Do not log-transform count data. *Nature Precedings*, 1-1.

Otto, R., Whittaker, R. J., von Gaisberg, M., Stierstorfer, C., Naranjo-Cigala, A., Steinbauer, M. J.,
… Fernández-Palacios, J. M. (2016). Transferring and implementing the general dynamic model
of oceanic island biogeography at the scale of island fragments : The roles of geological age and
topography in plant diversification in the Canaries. *Journal of Biogeography*, *43*(5), 911-922.
https://doi.org/10.1111/jbi.12684

Parent, C. E. (2012). Biogeographical and Ecological Determinants of Land Snail Diversification on
Islands [*]. *American Malacological Bulletin*, *30*(1), 207-215.
https://doi.org/10.4003/006.030.0118

Parent, C. E., Caccone, A., & Petren, K. (2008). Colonization and diversification of Galápagos
terrestrial fauna : A phylogenetic and biogeographical synthesis. *Philosophical Transactions of*
the Royal Society B: Biological Sciences, *363*(1508), 3347-3361.
https://doi.org/10.1098/rstb.2008.0118

Parent, C. E., & Crespi, B. J. (2006). Sequential colonization and diversification of Galápagos
endemic land snail genus *Bulimulus* (Gastropoda, Stylommatophora). *Evolution*, *60*(11),
2311-2328. https://doi.org/10.1111/j.0014-3820.2006.tb01867.x

Paulay, G. (1994). Biodiversity on oceanic islands : Its origin and extinction[1]. *American zoologist*,
34(1), 134-144.

Price, J. P., & Wagner, W. L. (2011). A phylogenetic basis for species-area relationships among three
Pacific Island floras. *American Journal of Botany*, *98*(3), 449-459.
https://doi.org/10.3732/ajb.1000388

Price, J. P., & Wagner, W. L. (2018). Origins of the Hawaiian flora : Phylogenies and biogeography
reveal patterns of long-distance dispersal. *Journal of Systematics and Evolution*, 56(6), 600-620.
https://doi.org/10.1111/jse.12465

R Core Team. (2023). *R: A Language and Environment for Statistical Computing. R Foundation for*
Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

Roell, Y. E., Phillips, J. G., & Parent, C. E. (2021). Effect of topographic complexity on species
richness in the Galápagos Islands. *Journal of Biogeography*, *48*(10), 2645-2655.
https://doi.org/10.1111/jbi.14230

Román-Palacios, C., & Wiens, J. J. (2018). The Tortoise and the Finch : Testing for island effects on
diversification using two iconic Galápagos radiations. *Journal of Biogeography*, *45*(8),
1701-1712. https://doi.org/10.1111/jbi.13366

Rudbeck, A. V., Sun, M., Tietje, M., Gallagher, R. V., Govaerts, R., Smith, S. A., … Eiserhardt, W.
L. (2022). The Darwinian shortfall in plants : Phylogenetic knowledge is driven by range size.
Ecography, *2022*(8), e06142.

Sakai, A. K., Weller, S. G., Wagner, W. L., Nepokroeff, M., & Culley, T. M. (2006). Adaptive
radiation and evolution of breeding systems in *Schiedea* (Caryophyllaceae), an endemic
hawaiian genus. *Annals of the Missouri Botanical Garden*, *93*(1), 49-63.

Sanmartín, I., Van Der Mark, P., & Ronquist, F. (2008). Inferring dispersal : A Bayesian approach to
phylogeny-based island biogeography, with special reference to the Canary Islands. *Journal of
Biogeography*, *35*(3), 428-449. https://doi.org/10.1111/j.1365-2699.2008.01885.x

Sarnat, E. M., & Economo, E. P. (2012). *The ants of Fiji*. Univ of California Press.

Schenk, J. J. (2021). The Next Generation of Adaptive Radiation Studies in Plants. *International
Journal of Plant Sciences*, *182*(4), 245-262. https://doi.org/10.1086/713445

Schluter, D. (2000). *The ecology of adaptive radiation*. Oxford: Oxford University Press.

Shaw, K. L., & Gillespie, R. G. (2016). Comparative phylogeography of oceanic archipelagos :
Hotspots for inferences of evolutionary process. *Proceedings of the National Academy of
Sciences*, *113*(29), 7986-7993. https://doi.org/10.1073/pnas.1601078113

Stuessy, T. F., Jakubowsky, G., Gómez, R. S., Pfosser, M., Schlüter, P. M., Fer, T., … Kato, H.
(2006). Anagenetic evolution in island plants. *Journal of Biogeography*, *33*(7), 1259-1265.
https://doi.org/10.1111/j.1365-2699.2006.01504.x

Tao, R., Sack, L., & Rosindell, J. (2021). Biogeographic Drivers of Evolutionary Radiations.
Frontiers in Ecology and Evolution, *9*, 644328. https://doi.org/10.3389/fevo.2021.644328

Triantis, K. A., Guilhaumon, F., & Whittaker, R. J. (2012). The island species-area relationship :
Biology and statistics: The island species-area relationship. *Journal of Biogeography*, *39*(2),
215-231. https://doi.org/10.1111/j.1365-2699.2011.02652.x

Triantis, K. A., Whittaker, R. J., Fernández-Palacios, J. M., & Geist, D. J. (2016). Oceanic
archipelagos : A perspective on the geodynamics and bi- ogeography of the World's smallest
biotic provinces. *Frontiers of Biogeography*.

Vitales, D., Pellicer Moscardó, J., Vallès Xirau, J., & Garnatje i Roca, T. (2015). Molecular insights
into the diversification of *Cheirolophus* (Asteraceae) in Macaronesia. In *Recent Advances in
Pharmaceutical Sciences V* (p. 85-100). Research Signpost.

Voeten, C. (2023). *Buildmer : Stepwise Elimination and Term Reordering for Mixed-Effects
Regression. R package version 2.8, <https://CRAN.R-project.org/package=buildmer>*.

Wagner, C. E., Harmon, L. J., & Seehausen, O. (2014). Cichlid species-area relationships are shaped
by adaptive radiations that scale with area. *Ecology Letters*, *17*(5), 583-592.
https://doi.org/10.1111/ele.12260

Warren, B. H., Simberloff, D., Ricklefs, R. E., Aguilée, R., Condamine, F. L., Gravel, D., …
Thébaud, C. (2015). Islands as model systems in ecology and evolution : Prospects fifty years
after MacArthur-Wilson. *Ecology Letters*, *18*(2), 200-217. https://doi.org/10.1111/ele.12398

Weigelt, P., & Kreft, H. (2013). Quantifying island isolation—Insights from global patterns of insular plant species richness. *Ecography*, *36*(4), 417-429. https://doi.org/10.1111/j.1600-0587.2012.07669.x

Weigelt, P., Steinbauer, M. J., Cabral, J. S., & Kreft, H. (2016). Late Quaternary climate change shapes island biodiversity. *Nature*, *532*(7597), 99-102. https://doi.org/10.1038/nature17443

White, O. W., Reyes-Betancort, J. A., Chapman, M. A., & Carine, M. A. (2020). Geographical isolation, habitat shifts and hybridisation in the diversification of the Macaronesian endemic genus *Argyranthemum* (Asteraceae). *New Phytologist*, *228*(6), 1953-1971.

Whittaker, R. J., Fernández-Palacios, J. M., & Matthews, T. J. (2023). *Island Biogeography : Geo-environmental Dynamics, Ecology, Evolution, Human Impact, and Conservation*. Oxford University Press.

Whittaker, R. J., Triantis, K. A., & Ladle, R. J. (2008). A general dynamic theory of oceanic island biogeography : A general dynamic theory of oceanic island biogeography. *Journal of Biogeography*, *35*(6), 977-994. https://doi.org/10.1111/j.1365-2699.2008.01892.x

Yang, Y., Morden, C. W., Sporck-Koehler, M. J., Sack, L., Wagner, W. L., & Berry, P. E. (2018). Repeated range expansion and niche shift in a volcanic hotspot archipelago : Radiation of C 4 Hawaiian *Euphorbia* subgenus *Chamaesyce* (Euphorbiaceae). *Ecology and Evolution*, *8*(16), 8523-8536. https://doi.org/10.1002/ece3.4354

Yu, F., Wang, T., Groen, T. A., Skidmore, A. K., Yang, X., Geng, Y., & Ma, K. (2015). Multi-scale comparison of topographic complexity indices in relation to plant species richness. *Ecological complexity*, *22*, 93-101.

Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects models and extensions in ecology with R* (Vol. 574). Springer.