Species-specific loss of genetic diversity and accumulation of genetic load following agricultural intensification

Zachary J. Nolen[1*], Patrycja Jamelska[1,2], Ana Sofia Torres Lara[1,3], Niklas Wahlberg[1], Anna Runemark[1]

[1] Department of Biology, Lund University, Lund, Sweden 22362

[2] Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland 2000

[3] Faculty 2 Biology/Chemistry, University of Bremen, Bremen, Germany 28359

* Corresponding author: Zachary J. Nolen; Kontaktvägen 10, Lund 22362, Sweden; zachary.nolen@biol.lu.se

Keywords: Conservation, agricultural intensification, inbreeding, genetic load, genomic erosion, historical DNA

Abstract

Agriculture driven land-use change is a major contributor to insect declines globally. As a result of population decline and subdivision, species lose genetic diversity, potentially reducing adaptive potential and increasing genetic load, reinforcing decline. We examine how genetic diversity has changed over the past century in three Polyommatinae (Lepidoptera: Lycaenidae) butterfly species in southern Sweden, leveraging genome sequencing of specimens from museum collections. Based on 63 historical and 92 contemporary genomes, we document a general decline in genetic diversity and reveal that the extent of decline depends on habitat specialization. The habitat generalist has retained most of its genetic diversity and connectivity. In contrast, the two specialists have declined sufficiently that even common variation is being eroded by drift, with up to a 3.3% reduction in heterozygosity per decade in the strongly grassland associated *Cyaniris semiargus*. While this sharp decline has reduced total genetic load in the two specialist species, increased population isolation has led to inbreeding in *Cy. semiargus*, exposing a greater proportion of load in homozygous state. This realized load likely reduces mean fitness and reinforces population decline, highlighting the need for well-connected grassland landscapes to maintain grassland specialist functional connectivity and genetic diversity. We complement these analyses with data on seven additional Polyommatinae species, uncovering that declines are common across the group, with alarming rates of genetic decline in species listed as viable. This study highlights the urgency of incorporating genetic data into conservation planning, as threats from genetic erosion are not captured by current conservation assessment.

Introduction

Insect decline has received increased attention in recent years as reports uncover alarming declines in insect assemblages and biomass (1). Efforts to quantify this decline estimate that terrestrial insects are declining at an average rate of 11% per decade globally, with this trend being primarily driven by even stronger regional declines in Europe and North America (2). This stark decline is the result of a variety of anthropogenic pressures, primarily climate change and land use changes such as agricultural intensification and urbanization (1, 3–7). Intensive agricultural landscapes place a particularly strong pressure on grassland insect species (6), especially in Europe where formerly biodiverse semi-natural grasslands have been abandoned or converted into intensively cultivated land (8).

While these declines already have alarming implications for natural communities and ecosystem services in which insects play key roles, they can also leave insect populations with genetic challenges to long term persistence. Declining populations are expected to lose genetic diversity as the relative strength of genetic drift increases, reducing their ability to adapt to changing environments and potentially unmasking genetic load through inbreeding and weakened selection (9). These processes can reduce mean fitness within the population, reinforcing demographic decline, a process referred to as mutational meltdown (10). An increasing number of studies document that this process can continue even upon demographic recovery, with persisting loss of genetic diversity and accumulation of genetic load due to drift debt (11–14).

How to assess the threat genetic factors pose to natural populations is still an active area of research, with several genetic monitoring approaches being suggested and implemented (15–18). Interpretation of genetic diversity is complex, as it is influenced by a multitude of factors (19), such as ecology and life history (20), genomic architecture (21), and demographic trends over thousands to millions of years (22). Low genetic diversity is thus not always a cause for alarm, as populations with a long history of small population sizes may have effectively purged deleterious variation that threatens persistence (23–25). Rather, reductions in genetic diversity in historically

large populations, which carry a high masked potential load, can increase extinction risk

as genetic load becomes stochastically realized (9, 26).

The rate of genetic decline, rather than current levels of genetic diversity, has become a

focus for genetic monitoring, with programs in several countries using genetic metrics

measured over time or that can directly measure ongoing genetic decline (16, 27),

e.g. contemporary effective population size (28). Temporal sampling is, however, limited

to estimating genetic decline from the first sampled time point, with considerable erosion

likely to have occurred several decades before. To overcome this, genomic resources

from historic samples preserved in natural history museums can be an invaluable

resource, extending monitoring further back in time. The direct measures of pre- and

early bottleneck genetic parameters museum samples provide are crucial to fully

understand the extent and character of genetic decline (18).

Here, we use museomics to address to what extent genetic diversity has been lost and

functional connectivity reduced in grassland associated butterflies during the second

wave of agricultural modernization in Sweden starting in the 1940s (29, 30). During this

modernization, small farms were abandoned and their associated semi-natural

grasslands began to decline considerably (30). We examine three focal butterfly species

in this landscape that vary in habitat specialization, ranging from grassland generalist to

specialist, enabling us to examine if patterns of genetic decline are stronger in more

specialized species, in line with expectations from greater demographic declines in

specialist insect abundance compared to generalists (31–33). By comparing pairs of

historical population samples collected in southern Sweden in the early 20th century

with their contemporary counterparts, we document species-specific losses of genetic

diversity and functional connectivity, as well as increased inbreeding depression in the

most specialized species. We further examine if patterns of loss of genetic diversity are

general, and if loss of genetic diversity is reflected in current conservation status, by

extending the dataset to include historical and modern individuals of seven additional

related butterfly species with Swedish Red List (34) conservation statuses ranging from

least concern to critically endangered.

Results

We analyzed changes in genetic diversity, differentiation, and genetic load in three focal Polyommatinae (Lepidoptera: Lycaenidae) butterfly species: *Polyommatus icarus,* a grassland generalist, *Plebejus argus*, a heathland specialist, and *Cyaniris semiargus*, a grassland specialist (Figure 1A). We generated whole genome resequencing data from pinned museum specimens from the entomological collections of the Biological Museum at Lund University (35) collected from 1917-1956 (Table S1; N = 28 - *Po. icarus*; 12 - *Pl. argus*; 14 - *Cy. semiargus*) in Southern Sweden. These individuals were sampled in populations generally collected at the same locality during the same year, and compared with WGS data from samples collected at corresponding localities in 2020-2023 by Nolen et al (36) (Figure 1B; Table S1; N = 27 - *Po. icarus*; 29 - *Pl. argus*; 29 - *Cy. semiargus*). As DNA extracted from historical specimens is fragmented and subject to post-mortem DNA damage (37, 38) (Figure S1), we confirmed shorter fragments did not induce a reference bias in the historical samples (Figure S2). To address DNA damage, we tested post-mortem DNA damage correction I) using MapDamage2 (39) and II) removing all transition sites, which are the most common changes caused by DNA damage (37, 38). Rescaling base quality scores with MapDamage2 biased historical samples towards the reference in two of the three focal species, whereas transition removal did not (Figure S2). All subsequent results were therefore generated without MapDamage2 correction and with transitions removed. To account for lower sequencing depth in historical samples, we performed analyses both at full depth and with samples subsampled to equivalent mean sequencing depths, presenting the subsampled results in the main text, with full depth results included in the Appendix.

General reductions in genetic diversity

To assess if agricultural intensification and the associated reduction in semi-natural grassland habitat has resulted in reduced genetic diversity, we compared individual genome-wide heterozygosity of historical and modern samples within each species. In all three species, heterozygosity is lower in modern individuals, but considerably more so in the two specialist species compared to the generalist (Welch two-sample t-test: P = 0.006 in *Po. icarus*, < 0.001 in *Pl. argus*, and < 0.001 in *Cy. semiargus;* Figure 1C;

Tables S2-3). In the specialist species, *Pl. argus* and *Cy. semiargus*, population-specific

estimates of genetic diversity are lower in all modern populations compared to their

historical counterparts, both for nucleotide diversity (π; Wilcoxon signed-rank test:

median effect size range [-0.52, -0.40] in *Pl. argus* and [-0.76, -0.51] in *Cy. semiargus*)

and Watterson's θ (median effect size range [-0.77, -0.68] in *Pl. argus* and [-0.86, 0.70]

in *Cy. semiargus*), even after subsampling to account for variation in sample size

(Figure S3; Tables S4-5). In the generalist, *Po. icarus*, only Watterson's θ shows

declines robust to variation in sample size, and only in two of the three populations

(median effect size range [-0.29, -0.24]; Figure S3; Table S4-5). Tajima's D is higher in

modern populations of all three species compared to their historical counterparts,

consistent with population bottlenecks across the past century (median effect size range

[0.63, 0.68] in *Po. icarus*, [0.82, 0.84] in *Pl. argus*, and [0.73, 0.85] in *Cy. semiargus*;

Figure S3; Table S4-5).

Reduced connectivity may accelerate loss of genetic diversity

While genetic diversity decline is an expected product of demographic decline, it can

additionally be accelerated by reductions in gene flow (40). We reveal a loss of

functional connectivity among modern populations, reflected in stronger contemporary

population structure, supported by both genetic differentiation and clustering

approaches, likely contributing to the loss of genetic diversity. *Po. icarus* and *Cy.*

semiargus have stronger genetic differentiation (F_{ST}) between modern than historical

population pairs (Paired t-test: ΔF_{ST} = 0.014, *P* = 0.007 in *Po. icarus* and ΔF_{ST} = 0.053,

P = 0.036 in *Cy. semiargus*; Figure 1F; Table 6-7). F_{ST} increased in *Pl. argus* in the

single population pair present in both time periods (ΔF_{ST} = 0.021; Table S7). Including

all modern population pairs in the analysis estimates mean F_{ST} increases of 0.027 and

0.071 in *Pl. argus* and *Cy. semiargus*, respectively (Welch two-sample t-test: *P* = 0.001

in *Cy. semiargus*;Table S7). Principal component analyses (PCAs) and admixture

analyses also reveal generally stronger population structure in modern samples than

historical, and strongest in the most specialized *Cy. semiargus* (Figure 2A). Along PC1,

the greatest axis of variation, both *Po. icarus* and *Pl. argus* are clustered primarily into

historical and modern samples, consistent with the best fit admixture analyses for these

species (K=2; Figure 2B; Table S8). In these two species, modern samples show
greater among population differences than historical along PC2, however, consistent
with increases in genetic differentiation between localities. Greater than two clusters are
visible on the PCA for these two species, which, along with the correlation of residuals
for the admixture analyses (Figure S4), highlights that greater genetic structure likely
exists in these two species, but from low enough variation that we cannot reliably
uncover it with our sample sizes. Each modern population forms a distinct cluster in the
specialized species, *Cy. semiargus*, while historical populations cluster together in their
own separate cluster (Figure 2B). The stronger contemporary population structure is
especially driven by isolation of populations in the agricultural landscapes within
Sweden's southernmost province, Skåne (Figure 1D; Figure 2B; Figure S5).

Species-specific increases in inbreeding

The pronounced losses of functional connectivity and genetic diversity have resulted in
inbreeding within the specialist *Cy. semiargus*. This contemporary inbreeding is
reflected by an increase in the inbreeding coefficient F_{RoH} from historical to modern *Cy.*
semiargus samples (Welch two-sample t-test: ΔF_{RoH} = 0.065, *P* < 0.001; Figure 1D;
Table S3&9). F_{RoH} measures the proportion of the autosomal genome in runs of
homozygosity greater than a certain length (41), which we set to 100kb, roughly
corresponding to identical by descent tracks with shared ancestry in the past 200
generations (42). Across species, no historical individual has a coefficient greater than
0.04, and historical species averages are all below 0.01. In *Cy. semiargus*, modern
samples have a maximum coefficient of 0.33 and an average coefficient of 0.072, an
order of magnitude higher than the mean for historical samples. In modern *Cy.*
semiargus, individuals with higher inbreeding coefficients have fewer runs of
homozygosity relative to their cumulative length, which may indicate they stem from
close-kin matings (Figure 3A, Figure S6). For other individuals, there is a largely linear
relationship between the number of runs of homozygosity and their cumulative length,
suggesting that close-kin mating is not common overall and inbreeding is primarily due
to increases in background relatedness, consistent with patterns expected of a
population bottleneck (42).

Purging of total genetic load

Population decline and subdivision can result in increased exposure of genetic load, in the form of rare recessive deleterious variants in a homozygous state, to selection (43). This may facilitate purging, but can also result in greater realization of minorly deleterious variation in homozygous state, reducing mean population fitness and reinforcing decline. We find evidence consistent with purging in both specialist species *Pl. argus* and *Cy. semiargus*, as total genetic load is lower in modern compared to historical samples both for high (e.g. loss of function; Welch two-sample t-test: $P =$ 0.024 in *Pl. argus* and 0.021 in *Cy. semiargus*) and moderate (e.g. missense; $P < 0.001$ in *Pl. argus* and $= 0.003$ in *Cy. semiargus*) impact variants (Figure 3B; Table S10). In generalist *Po. icarus*, total genetic load from high and moderate impact variants is similar in modern and historical samples. Total genetic load from derived variants at highly genomically conserved positions has remained stable in all three species.

Increased realized genetic load in *Cy. semiargus*

Modern individuals have greater realized genetic load at moderate impact sites compared to historical individuals in *Cy. semiargus* ($P = 0.013$; Figure 3C; Table S10), despite evidence for purging of the genetic load in this species. In contrast, there is no increase in realized load for high impact sites. Realized genetic load incorporates only putatively deleterious alleles in a homozygous state, reflecting the proportion of total load that is expressed in an individual. We find no evidence for increases in realized genetic load at moderate or high impact sites in *Po. icarus* or *Pl. argus*, with realized load even decreasing at moderate sites in *Pl. argus* ($P < 0.001$; Figure 3C; Table S10). However, derived variants at conserved regions are realized to a greater extent in modern samples of both *Pl. argus* and *Cy. semiargus* ($P < 0.001$ in *Pl. argus* and $=$ 0.009 in *Cy. semiargus*; Figure 3C; Table S10), despite total load in these regions remaining stable. In *Cy. semiargus*, individuals with higher inbreeding coefficients had higher realized, but not total, genetic load across all variant categories (linear regression: $R^2 = 0.46, 0.37, 0.77$ for conserved, high, and moderate impact variants, respectively; Figure S7), indicative of inbreeding depression.

Genetic decline not reflected in current conservation status

We utilized the genetic decline indicator thresholds outlined by Andersson et al (15) to determine if current conservation status relates to the rate of genetic decline. These thresholds suggest categorizations of 'Acceptable', 'Warning', and 'Alarm' rates of genetic decline, reflecting >=95%, 75-94%, and <75% retention of genetic diversity over 100 years, respectively. Based on changes in heterozygosity, nucleotide diversity, and Watterson's θ, *Po. icarus* largely exhibits 'Acceptable' or 'Warning' declines, while *Pl. argus* and *Cy. semiargus* exhibit 'Warning' or 'Alarm' declines, depending on the metric of genetic diversity used (Figure 4&S8; Tables S5&11), which largely are in line with the concerns raised by our genetic load analyses. This is in stark contrast with the 2020 Swedish Red List of species (34), where all three focal species share a conservation status of 'Least Concern'.

To assess if the loss of genetic diversity in the three focal species is representative of other grassland butterfly species, we supplemented the dataset by generating whole genome resequencing data for historical-modern pairs of seven additional species of Polyommatinae butterfly (Table S1). We assessed if the conservation status of these species, ranging from of 'Least Concern' to 'Critically Endangered' (34) (Table S12) capture losses of genetic diversity uncovered in this data set. Comparing individual heterozygosity, we find that current conservation status generally fails to capture the levels of genetic erosion in these species (Figure 4A&S8; Table S11). Utilizing estimates at full individual sequencing depth, least concern species predominantly fell into the warning category, with only *Celastrina argiolus* and Öland *Po. icarus* falling in the acceptable category. Of species classified as least concern, only *Cy. semiargus* fell into the alarm category. Two endangered species, *Phengaris alcon* and *Scolitantides orion,* fell into the acceptable and alarm categories, respectively, with *Ph. alcon* even increasing slightly in genetic diversity over the time frame. *Plebejus argyrognomon*, a critically endangered species that is now presumed extinct in Sweden (44), had already experienced genetic decline categorized as 'alarm' by the early 2000s, when our modern sample for this species was collected. Results from our focal species illustrate that categorization can depend considerably on the individuals/populations sampled

240　(Figure 4). Categorization is largely consistent for each species both at full sequencing
241　depth and when subsampled to 2-3X sequencing depth, but some species/populations
242　bordering adjacent categories shift between them upon subsampling (Table S11).

Discussion

Our analyses of 63 historical and 92 modern genomes uncover a decline in butterfly
genetic diversity over the past century. This decline is greater in the specialist species,
especially *Cy. semiargus*, where the once highly connected populations have become
isolated into distinct genetic clusters. This subdivision and resulting loss of diversity has
increased rates of inbreeding and exposed genetic load to natural selection, resulting in
purging of both high and moderate impact mutations. Importantly, we uncover a higher
realization of genetic load in conserved regions and at moderate impact variants in
modern *Cy. semiargus*, likely reducing fitness. The rates of decline uncovered in the
three focal species examined are also common in seven additional species of blue
butterflies, despite most of these species having conservation statuses of 'Least
Concern'. Our findings reveal that recent genetic erosion is likely to be ubiquitous
across grassland butterflies in southern Sweden. These results highlight the need for
incorporation of genetic data into conservation assessments of insect threat status, as
census-based estimates fail to detect threats posed by genetic decline.

Our findings of reduced genetic diversity across all three focal species are consistent
with widespread demographic decline in arthropod species in Europe over the past
century, with the modernization of agriculture as a major driver (5, 45). The highest
average decline in genetic diversity we document, an up to 3.3% reduction in
heterozygosity per decade in *Cy. semiargus,* appears to be accelerated by
strengthened isolation between populations. However, both *Po. icarus* and *Pl. argus*,
where increases in population subdivision are more moderate, have experienced
declines of up to 0.7% and 2.7% per decade, respectively. Population structure and
genetic differentiation analyses suggest modest increases in population isolation in
these species, potentially reflecting that declines in genetic diversity predate genomic
signals of population subdivision. Population differentiation accumulates over time,
lagging behind barriers to gene flow (46, 47), with the rate at which genetic

differentiation accumulates after subdivision depending upon the underlying effective
population sizes of the populations (48).

Importantly, the finding that these three species have historically had low levels of
genetic differentiation across southern Sweden highlights that currently isolated
populations may be made up of partitions of historically well-connected
metapopulations. Hence, concerns about genetic rescue (49) wiping out locally adapted
ancestry (50, 51) are not likely to be relevant for these and other grassland associated
insects in regions with historically abundant habitat. Instead, restoring functional
connectivity to reduce inbreeding and associated accumulation of deleterious genetic
variation (10, 52) is likely key to improve prospects of long-term persistence. Previous
work on *Cy. semiargus* has shown that the inbreeding in these modern populations is
related to low grassland land cover in the surrounding landscape (36). Expanding green
infrastructure and restoration of semi-natural grassland habitat, enabling dispersal and
buffering genetic decline, is likely needed to remedy future isolation and inbreeding in
grassland associated butterflies.

Understanding to what extent genetic diversity decline actually reduces the fitness of
individuals and populations is crucial for determining conservation priorities. The
relationship between genetic diversity and extinction risk is highly contextual, and may
be confounded by demography and dispersal dynamics (25). To determine if extinction
risk has increased without having direct estimates of fitness, we assess genetic load.
Decreases in genetic load may reflect purging of recessive, highly deleterious variation
during and after a bottleneck, as the increased strength of genetic drift and higher rates
of inbreeding expose recessive variants to selection in homozygous state (43).
However, while selection may be strong enough to purge highly deleterious alleles,
moderately deleterious alleles may become expressed more often due to increased
homozygosity, with insufficient selective pressure to overcome the increased strength of
drift (9, 26, 53). We reveal that both specialist species have experienced purging of both
loss of function and missense mutations. Despite this, in *Cy. semiargus*, increases in
inbreeding have led to higher realized genetic load from persisting missense mutations,
largely driven by greater expression of load in individuals with higher inbreeding

coefficients. Reduced heterozygosity has been shown to increase extinction risk in

small, isolated, recently declined populations belonging to a larger meta-population of

the Glanville fritillary butterfly (*Melitaea cinxia*) (25, 54), underscoring the risks

associated with increased expression of genetic load in the recently isolated, inbred

populations of *Cy. semiargus*.

Genetic load estimates must be interpreted with consideration. Functional effect based

genetic load estimates are reliant upon correct underlying gene models in the

annotation and assume the reference allele is less deleterious than the alternate. We

define conserved positions by selecting those with the top 1% of Genomic Evolutionary

Rate Profiling (GERP) scores, however, these scores may not confidently distinguish

between weakly and strongly deleterious mutations (55, 56). Our score thresholds are

notably lower than thresholds utilized to assess genetic load in vertebrates (55, 57–59),

and lower than estimates for other Lepidopterans estimated from a multi-species

alignment (60), the latter likely due to our use of a time tree to estimate scores

compared to a neutral rate tree by Bortoluzzi et al (60). However, our identification of

conserved sites should be robust for within-species genetic load comparisons, as they

represent the upper extreme of the distribution in our data. Moreover, our findings are

consistent with predictions from population genetic theory, and hence likely reflect true

impacts on genetic load and can inform management for these species. Genetics

informed conservation action to increase long-term viability must focus on reducing both

masked and realized genetic load (26). This is most pressing in *Cy. semiargus*, where

translocations are likely necessary if populations with high levels of inbreeding are not

to suffer continued decline from inbreeding depression.

None of the three focal species are currently demographically declining at a rate to

qualify for the Swedish Red List based on demographic data. Systematic monitoring in

Sweden from 2010-2023 reveal declines in both *Po. icarus* and *Cy. semiargus*, while *Pl.*

argus appears to have stabilized (61). Demographic stability in *Pl. argus* does not

necessarily mean that genetic diversity has stabilized in the species, as populations

may continue to lose genetic diversity even after demographic recovery, a phenomenon

termed drift debt (11). In *Cy. semiargus*, genetic factors are likely already contributing to

its demographic decline. Our findings illustrate the importance of incorporating genetic

monitoring into conservation assessment (62, 63) to identify genetic decline before it

results in expressed genetic load. Temporal monitoring with historical samples provides

an opportunity to detect species facing recent declines in genetic diversity, separating

them from those with longer term low genetic diversity (18). Using this approach on

several species of Polyommatinae blues in southern Sweden, we document a general

loss of genetic diversity in grassland butterflies and uncover that additional species

currently not on the Red List likely face threats from reduced genetic diversity.

Estimations of genetic decline along with complementary estimates of genetic load

provide a promising way to assess genetic conservation concerns in the wild (14), but

these analyses are limited by the availability of high quality reference genomes. This

underscores the importance of efforts to generate high quality references for wild

species, as they will improve our ability to accurately assess these metrics (64).

In conclusion, we document a general decline in genetic diversity in grassland

butterflies in southern Sweden, which is likely mirrored in other regions that have lost

grasslands to intensive agriculture and forestry. While demographic declines largely

inform current conservation status of these species, we show that this is insufficient to

capture declines in genetic diversity and resulting inbreeding, highlighting the acute

need for genetic monitoring of wild insects to prevent extirpation and extinction. *Cyaniris*

semiargus, a once widespread species in Sweden, now persists in segmented

populations experiencing inbreeding depression and requiring prompt conservation

action to reduce the risk of local extinction, providing a strong example of this.

Methods

Study System and Sampling

To assess changes in genetic diversity, differentiation, inbreeding, and genetic load, we

selected three blue butterfly species (Polyommatinae) as focal species, *Polyommatus*

icarus, *Plebejus argus*, and *Cyaniris semiargus* Figure 1A. These three species were

selected as they vary in habitat specialization, being grassland generalist, heathland

specialist, and grassland specialist, respectively, in order to assess how specialization

may impact genetic decline. For these three species, we used population level genetic

data we previously generated (36) as our modern dataset. We selected pinned

specimens from the entomological collections of the Biological Museum, Lund

University (35) that corresponded to roughly the same geographic area as the modern

samples for each species. These historical specimens were sampled between 1917-

1956, during the early stages of the second wave of agricultural modernization in

Sweden (29). For each historical locality, we selected 4-11 specimens sampled in the

same year to estimate population level metrics. The historical and modern population

level pairs were sampled in four broad geographic areas: I) Western and II) Eastern

Skåne, III) Småland, and IV) Öland (Figure 1B), with historical localities ranging

approximately 12-40km from their modern counterparts. This resulted in historical-

modern population level pairs for three regions in *Po. icarus* (Western and Eastern

Skåne and Öland), two regions in *Pl. argus* (Småland and Öland), and three regions in

Cy. semiargus (Western and Eastern Skåne and Småland). While most regions consist

of only one historical-modern population pair, we additionally compared a modern *Cy.*

semiargus population in Southeastern Skåne to the historical Eastern Skåne population.

We were unable to sample population level historical data for *Pl. argus* in Skåne,

however we did sample one historical specimen each from Western and Eastern Skåne,

which we include in individual level analyses, pairing them with the modern samples

from Southwestern and Eastern Skåne, respectively.

To more broadly assess the variation in rates of genetic decline in Polyommatinae of

differing conservation statuses, we sampled historical and modern individuals of seven

additional species: *Agriades optilete, Aricia agestis, Celastrina argiolus, Polyommatus*

amandus, Phengaris alcon, Scolitantides orion, and *Plebejus argyrognomon.* Modern

specimens of *Ag. optilete, Ar. agestis, Ce. argiolus,* and *Po. amandus* were collected

across the same geographic regions as the focal species in 2021 (Table S1). Pinned

specimens to be used as modern samples of *Ph. alcon* and *S. orion,* collected in 2003

in Western Skåne and 2016 in Södermanland, respectively, were donated to this study

by Håkan Elmqvist. Legs from a pinned speciemen of *Pl. argyrognomon*, collected in

2003 in Western Skåne, were donated by Göran Engqvist and used as the modern

sample for this species. The historical samples for these eight species all were selected

to be proximal to the modern specimen sampling locality with collection dates ranging

from 1932-1967, all provided as pinned specimens from the entomological collections of

the Biological Museum, Lund University (35).

DNA Extraction and Sequencing

DNA from fresh specimens was extracted and sequenced following the methods in

Nolen et al (36). We extracted DNA and prepared sequencing libraries from pinned

museum specimens in a facility dedicated to historical DNA processing at Lund

University, following the protocol described by Twort et al (65). Libraries prepared for

pinned specimens were pooled into two equimolar pools containing 37-38 libraries

each, sequencing each pool on a single NovaSeq 6000 S4 lane. DNA was extracted

from the abdomens or legs of male specimens without homogenization to preserve

anatomical structures. Abdomens were then stored in glycerol together with the original

specimen.

Genomic Analyses

We performed the majority of our analyses using the PopGLen pipeline (66), a

Snakemake (67) pipeline we have built that combines several genotype likelihood

based population genomic analyses. Genotype likelihood based analyses use a

probabilistic framework to account for the uncertainty that comes with low coverage

sequencing data, reducing biases related to lower sequencing depth in our historical

samples. We utilized genotype likelihood based analyses for all results, except for the

estimation of inbreeding coefficients and genetic load, where the chosen methods

required called genotypes.

Sequence Processing

We pre-processed both modern and historical raw sequencing data using fastp v0.23.4

(68), using default settings to trim low quality base pairs and poly-G tails from the ends

of reads in modern samples and additionally collapsing overlapping read pairs in

historical samples, where we discarded reads with less than 30bp overlap. This

processing of historical samples discards short (<30bp) fragments that contain little data

usable for unique mapping, as well as long fragments (>270bp) that are more likely to

be modern contaminants.

We mapped the trimmed, paired modern reads separately per sequencing run to the

respective species-specific reference genomes, or to the closest relative which an

available chromosome level assembly (Table S12), using BWA mem v0.7.17 (69). We

mapped the collapsed historical reads to the same reference genomes using BWA aln,

which is the recommended algorithm for reads shorter than 70bp (70) using settings

optimized for ancient/historical DNA (`-l 16500 -n 0.01 -o 2`) (71).

We merged mapped sequencing runs per sample using Samtools v1.17 (72) and

removed PCR and optical duplicates using Picard v2.27 (73) with default settings

except for an optical duplicate pixel distance of 2500 as recommended for patterned

flow cell data. We removed PCR duplicates for our historical samples using dedup

v0.12.8 (74), which identifies duplicates using both ends of the collapsed sequencing

read.

We realigned around indels using the GATK IndelRealigner (75). As the primary tool we

are using, ANGSD, counts overlapping reads twice (76), we additionally clipped

overlapping read pairs in the modern samples using BamUtil v1.0.15 (77). We

estimated post-mortem DNA damage in historical samples using DamageProfiler v1.1

(78).

Site and read filtering

We utilized a site based filtering scheme similar to that implemented in (79) in which we

developed multiple independent sets of filtered sites, generating a global list of filtered

sites from all analyses that excludes sites filtered by any of the sets. We performed this

filtering for each species dataset separately.

We excluded sex linked chromosomes, mitochondria, and scaffolds less than 1 Mb in

length. We excluded repeats identified by RepeatMasker v4.1.2 (80), using the

reference specific repeat libraries provided by the Darwin Tree of Life Project

(https://projects.ensembl.org/darwin-tree-of-life/). We estimated regions in the lower 1%

and upper 99% of global sequencing depth separately across all samples, historical

samples, and modern samples, excluding sites that appeared in any of the three sets.

The number and proportion of sites passing each of these filters, as well as passing the

final intersected filter set, are reported for each species in Table S13. We estimated

mean sequencing depth without filtering, as well as within the filtered site set, for each

sample using ANGSD (Table S1). All subsequent analyses were performed on the

species-specific filtered site sets.

Reference bias and post-mortem DNA damage

Due to DNA degradation, historical samples will have fragmented DNA with post-

mortem damage that alters base composition (81). We investigated two potential

methods to account for post-mortem DNA damage: I) recalibration of putatively

damaged bases using MapDamage2 (39), and II) removal of transitions across the

dataset, where most DNA damage is expected to occur (37, 38). To assess the

performance of these methods, we compared the identity by state (IBS) similarity to the

reference at filtered sites between these two correction methods and the uncorrected

data (Figure S2). We estimated the identity by state similarity to the reference using

ANGSD's `-doIBS 1` function with the `-doMajorMinor 4` and `-output01 1` options, which

together sample a single base at each position, assigning 1 to positions that sample the

reference base and 0 to positions sampling the non-reference. We calculated individual

identity by state similarity to the reference by averaging these values, where a value of

1 indicates complete similarity to the reference and 0 indicates complete dissimilarity.

Historical samples without correction had less similarity to the reference than modern

(Figure S2), which is expected for historical samples without substantial mapping bias

due to drift from the sampling time of the historical samples and the modern reference

as well as differences introduced by post-mortem DNA damage (82, 83), suggesting

shorter fragment size does not introduce substantial reference bias. After MapDamage2

rescaling, historical samples showed higher identity by state similarity to the reference

than modern samples in *Po. icarus* and *Pl. argus*. As removal of transitions did not

introduce this reference bias, we proceeded using transition removal in all analyses to

account for post-mortem DNA damage.

Genotype Likelihood Calculations

We estimated genotype likelihoods in ANGSD v0.940 (84) using the GATK model (`-GL`

`2`), using reads with a mapping quality greater than 30 (`-mapMapQ 30`) and base quality

greater than 20 (`-minQ 20`). We filtered our analyses to sites present in the species-
specific filtered sites list using the `-sites` option.

Heterozygosity and genetic diversity metrics

We estimated individual genome-wide heterozygosity for all samples using the single
sample site frequency spectrum (SFS) produced by realSFS, included with ANGSD.
First, we generated a sample allele frequency (SAF) file using (`-doSaf 1`) for each
sample, setting the ancestral state (`-anc`) to the reference sequence. We then
generated a folded (`-fold 1`) SFS in realSFS using this SAF file and calculated
heterozygosity as the second value (count of heterozygous sites) divided by the total
sum of all values (count of total analyzed sites). To test for differences in mean
individual heterozygosity across time periods, we performed an unequal variances *t*-test
comparing all samples within each time category for each of the focal species.

For the three focal species with population level samples, we estimated nucleotide
diversity, Watterson's θ, and Tajima's D per sampled population using thetaStat,
included with ANGSD. As the estimates of these metrics, especially Watterson's θ and
Tajima's D, can depend on sample size, we made our estimates using 100 subsamples
of each population to four individuals. For each subsample, we estimated a SAF file in
the same fashion as the heterozygosity estimates, additionally limited to sites for which
three of the four subsampled individuals had data (`-minInd 3`). We then used realSFS
to generate a folded SFS and thetaStat to estimate these three diversity metrics from
the SAF file and SFS in 10kb sliding windows with a step size of 5kb, excluding
windows with less than 1000 usable sites. The per site estimate of these three metrics
was obtained for each window by dividing the window estimate by the number of usable
sites (`nsites`). We report the median estimate across subsamples as the population
estimate of each metric. To statistically assess differences between historical and
modern populations, we performed a paired difference Wilcoxon signed-rank test each
corresponding historical and modern subsamples, pairing estimates by genomic
window. We estimated the effect size using the following equation:

$$\frac{Z}{\sqrt{N_{win}}}$$

where Z is the *Z*-score from the Wilcoxon signed-rank test and N_{win} is the number of

paired windows. We considered modern populations to have significantly different

estimates of nucleotide diversity, Watterson's θ, or Tajima's D if the distribution of effect

sizes across all 100 subsample tests did not overlap 0.

Genetic Differentiation

We estimated genetic differentiation using pairwise population F_{ST} using realSFS. We

generated population SAF files using the same parameters as for the genetic diversity

metrics and generated a two dimensional folded SFS for each population pair using

realSFS. We then used this SFS and the population SAF files to estimate a global

estimate of F_{ST} using the Bhatia estimator (`-whichFst 1`), which is less sensitive to

small or uneven sample sizes (85). Where reported, F_{ST} estimates refer to the weighted

F_{ST}. To statistically assess differences in F_{ST} between modern and historical

populations, we performed two tests: I) a paired *t*-test pairing corresponding population

pairs in historical and modern population sets and II) an unequal variances unpaired *t*-

test incorporating all possible population pairs in each set, comparing the historical to

the modern population set.

Relatedness

As the analyses we utilize to assess population structure can be sensitive to the

presence of close relatives in the dataset, we estimated pairwise relatedness between

all samples within each species dataset. We utilized the IBSrelate SFS-based method

(86), as implemented in NGSrelate (87), to estimate R0, R1, and KING-robust kinship

coefficients (88) for all pairs of individuals in the dataset. For input, we generated a

Beagle file using ANGSD for each focal species dataset (`-doGlf 2`), utilizing only

positions where greater than 80% of the individuals had data, calling SNPs (`-SNP_pval`

`1e-6`) and assuming known major and minor alleles inferred from the genotype

likelihoods (`-doMajorMinor 1 -doMaf 1`) (89, 90).

Population Structure

We explored population structure using principal component analysis (PCA) as

implemented in PCAngsd v1.10 (91). As the population structure methods we use

assume independence of sites, we first estimated linkage disequilibrium for SNPs less

than 50kb apart in the species dataset Beagle files using ngsLD v1.2.0 (92). We pruned

SNPs using the recommended prune_graph tool

(https://github.com/fgvieira/prune_graph; v0.3.2), assuming independence of SNPs

>50kb apart and with a coefficient of correlation (r^2) < 0.1 and subset the Beagle file to

include only pruned sites. We then removed one individual from each pair of third-

degree or closer relatives in the dataset (expected KING-kinship coefficient >= 0.0625;

Table S14) before running the analysis.

We formally tested for the presence of population structure using an admixture analysis

for 1-10 ancestral cluster (K) using the pruned, relative removed Beagle file in

NGSadmix v33 (93). We performed at least 20 replicate runs for each value of K, up to

100 total runs, stopping if the three highest likelihood replicates converged within 2 log-

likelihood units of each other. The 'best-fit' value for K was then determined to be the

highest K for which replicates met this convergence criteria, following Pečnerová et al

(79). We further assessed the quality of the model fit for each value of K using

evalAdmix v0.961 (94).

Genotype Calling

To estimate runs of homozygosity and genetic load in our three focal species, we

utilized methods that require genotype calls. As genetic load estimates can be biased

by variation in sequencing depth (14), we subsampled all samples to a mean 5X filtered

sequencing depth using Samtools, excluding samples with a mean filtered sequencing

depth below 5X from genotype call based analyses. We called genotypes per sample

using BCFtools multi-allelic caller v1.19 (72). Calling was performed only for regions in

our filtered sites file, using reads with a minimum mapping quality of 30 and bases with

a minimum quality of 20. We set genotypes with a read depth <4 to missing. We

additionally filtered heterozygous calls for allelic balance, setting heterozygous

genotypes where one allele was supported by <20% of the reads to missing. We then

merged the samples into a single BCF per species, filtering SNPs within 5bp of indels,

removing indels, and including only biallelic SNPs that were segregating in the dataset.

Finally, we only included sites with called genotypes in 80% of samples and excluded all

transition variants.

Runs of Homozygosity

We assessed rates of recent and background inbreeding by estimating the inbreeding

coefficient F_{RoH}, the proportion of the autosomal genome existing in runs of

homozygosity over a specified length, for each individual from the genotype calls. We

incorporated runs of homozygosity >100kb into our estimates of F_{RoH}. We estimated

runs of homozygosity using BCFtools/RoH v1.19 (95), setting a constant species-

specific recombination rate (`-M`; estimated assuming 50 centimorgans per autosomal

chromosome as in Mackintosh et al (21), estimates in Table S12) and running the

analysis on genotype calls only, with the PL of unseen genotypes set to 30 (`-G30`). As

we did not have sufficient samples to confidently estimate population allele frequencies,

we used a default minor allele frequency of 0.4 (`--AF-dflt 0.4`). We additionally

ignored homozygous reference positions (`--ignore-homref`), as including these can

inflate estimates of F_{RoH} when using a default allele frequency based on our testing

(Figure S9). We calculated F_{RoH} for each individual by dividing the total sum of runs of

homozygosity greater than 100kb in length by the total length of the autosomes.

Genetic load

We estimated individual levels of genetic load from the genotype calls using two

complementary methods, one targeting conserved regions of the genome (96) and the

other utilizing gene annotations to infer functional effects (97). We filtered the input

genotype calls for all genetic load analyses to include only positions with genotype calls

for 80% of the individuals in both the historical and the modern subsets.

Genomic conservation based methods utilize multiple species alignments to determine

genomic regions that are highly conserved across a taxonomic group, with deviations

from the ancestral state being likely to be deleterious. To estimate to which extent the

focal species had accumulated deleterious variants in conserved regions of the

genome, we used the GenErode pipeline v0.6.0 (98) to estimate Genomic Evolutionary

Rate Profiling (GERP) scores (99) across each of the three focal species genomes. The

method implemented in this pipeline does not require a multi-species alignment, instead

fragmenting the outgroup genomes and mapping the fragments to the target to generate

alignments which are then used to estimate scores. We used a dataset of 106

Lepidoptera chromosome level outgroup genomes (Table S15) to estimate the scores

and ancestral states, using divergence times estimated with TimeTree 5 (100). We

selected the highest 1% of GERP scores as the most conserved positions using the

`get_gerp_score_percentile.py` script included with GenErode. We estimated the total

relative genetic load at conserved sites for each individual by multiplying the derived

allele count at each position by its GERP score, summing these values across the

genome and dividing by the total derived allele count for that individual. We estimate

realized relative genetic load by multiplying the derived allele count for homozygous

derived genotypes by the GERP score, summing these values across the genome and

dividing by the total derived allele count for that individual. While estimating realized

load in this fashion neglects the realized effects of variants with partial dominance in

heterozygous state (9), the majority of mildly and highly deleterious mutations are likely

to be recessive (101) and realized only in homozygous state (82).

For the functional annotation based method, we used the annotations by the Darwin

Tree of Life Project (https://projects.ensembl.org/darwin-tree-of-life/). We assigned

impacts to each variant using Ensembl's Variant Effect Predictor (VEP) v112.0 (97).

VEP categorizes variants into four classes: 'High', 'Moderate', 'Low', or 'Modifier', which

largely correspond to loss of function, missense, synonymous, and intergenic variants.

As variants often overlap multiple transcripts, multiple impacts will be assigned, so we

required VEP to pick one effect per variant (`--flag_pick`), using the default selection

order. After assignment, we estimated total and realized genetic load at high and

moderate impact variants, treating the alternate allele as the putatively deleterious

variant. We estimated total genetic load as the alternative allele count for the given

impact category, divided by the total count of alternate alleles. We estimated realized

genetic load as the alternative allele count for homozygous alternate positions for the

given impact category, divided by the total count of alternate alleles.

Categorizing rates of genetic decline

To assess the changes in genetic diversity in a framework developed for conservation

focused genetic monitoring, we used the indicator thresholds suggested by Andersson

et al (15), which are based upon indicators applied for national use in Sweden (102).

Using these thresholds, genetic decline within populations is measured as the amount

of genetic diversity retained over a period of 100 years, assuming a constant rate of

change. Three threat categories are suggested: retention >=95% is considered

'Acceptable', 75-94% as 'Warning', and <75% as 'Alarm'. As different population genetic

estimates of genetic diversity are influenced by different components of genetic

diversity, such as rare and common variants, this metric can be assessed for each

estimate separately.

We estimated the predicted retention of genetic diversity over 100 years for each

population using individual heterozygosity, nucleotide diversity, and Watterson's θ.

Genetic diversity retention per year was estimated using the following equation:

$$e^{\frac{\ln(H_{mod}/H_{hist})}{Y_{mod}-Y_{hist}}}$$

where H is the estimate of genetic diversity for a given sample/population and Y is the

sampling year for that sample/population. As different populations had different sample

sizes, we used repeated subsamples to generate estimates of decline. For individual

heterozygosity, we performed all pairwise comparisons of historical and modern

individual estimates of heterozygosity within corresponding historical-modern sample

pairs, generating a single estimate for species with only one specimen per time period

and a range for ones with more (see Figure 4). For nucleotide diversity and Watterson's

θ, we utilized the previously generated 100 subsamples (N=4) for each population,

estimating H_{mod}/H_{hist} for paired historical and modern populations for each subsample,

utilizing the median value of H_{mod}/H_{hist} across subsamples to calculate the predicted

retention.

Acknowledgements

We would like to thank Melina Eberhagen, Amanda Görnerup, Kajsa Svensson, Aske Brydegaard Runemark, and Mikkel Brydegaard for assistance with collecting butterflies, Håkan Elmqvist for providing pinned specimens of *Ph. alcon* and *S. orion*, Göran Engqvist for providing tissue from the pinned specimen of *Pl. argyrognomon*, Emma Kärrnäs and Isolde van Riemsdijk for assistance with DNA extraction of samples, Jadranka Rota for assistance with the entomological collections of the Biological Museum, Lund University (https://doi.org/10.15468/dahk2a), Verena Kutschera and the Swedish Bioinformatics Advisory Program for advice regarding the bioinformatic analyses, Anna Penna for comments on the manuscript, and Isolde van Riemsdijk, Rachel Steward, Kalle Tunström, Simon Jacobsen Ellestrand, and Magne Friberg for helpful discussions regarding the study species, methods and interpretations. The computations were enabled by resources provided by the National Academic Infrastructure for Supercomputing in Sweden (NAISS) and the Swedish National Infrastructure for Computing (SNIC) at UPPMAX partially funded by the Swedish Research Council through grant agreements no. 2022-06725 and no. 2018-05973. This work was supported by grants from the Strategic Research Area "Biodiversity and Ecosystem services in a Changing Climate" (BECC), funded by the Swedish government, the Foundation Oscar och Lili Lamms Minne, the Jörgen Lindström's Foundation, and the Swedish Research Council for Sustainable Development (grant 2021-01163).

Data Availability Statement

The genetic data that support these findings will be made available upon publication at NCBI in BioProjects PRJNA1068054 (focal species, modern) & PRJNA1168917 (focal species, historical; additional species). The main population genomic pipeline is available at https://doi.org/10.5281/zenodo.13785563. Workflow configurations, extensions, and resources specific to this study are available at https://github.com/zjnolen/polyommatini-temporal-genomics.

Author Contributions

Zachary J. Nolen: Conceptualization, Methodology, Formal analysis, Investigation (Lead), Data curation, Writing – original draft, Visualization, Funding acquisition. **Patrycja Jamelska**: Investigation (Supporting), Writing - review & editing. **Ana Sofía Torres Lara**: Investigation (Supporting), Writing - review & editing. **Niklas Wahlberg:** Methodology, Writing – review & editing, Supervision. **Anna Runemark:** Conceptualization, Methodology, Writing – review & editing, Supervision, Funding acquisition.

References

References

1.   D. L. Wagner, E. M. Grames, M. L. Forister, M. R. Berenbaum, D. Stopak, Insect
decline in the anthropocene: Death by a thousand cuts. *Proceedings of the National*
Academy of Sciences **118**, e2023989118 (2021).

2.   R. van Klink, *et al.*, Meta-analysis reveals declines in terrestrial but increases in
freshwater insect abundances. *Science* **368**, 417–420 (2020).

3.   R. Dirzo, *et al.*, Defaunation in the Anthropocene. *Science* **345**, 401–406 (2014).

4.   C. A. Halsch, *et al.*, Insects and recent climate change. *Proceedings of the*
National Academy of Sciences **118**, e2002543117 (2021).

5.   P. H. Raven, D. L. Wagner, Agricultural intensification and climate change are
rapidly decreasing insect biodiversity. *Proceedings of the National Academy of*
Sciences **118**, e2002548117 (2021).

6.   S. Seibold, *et al.*, Arthropod decline in grasslands and forests is associated with
landscape-level drivers. *Nature* **574**, 671–674 (2019).

7.   D. H. Boyes, D. M. Evans, R. Fox, M. S. Parsons, M. J. O. Pocock, Is light
pollution driving moth population declines? A review of causal mechanisms across the
life cycle. *Insect Conservation and Diversity* **14**, 167–187 (2021).

8.   J. R. Shipley, *et al.*, Agricultural practices and biodiversity: Conservation policies
for semi-natural grasslands in europe. *Current Biology* **34**, R753–R761 (2024).

9.   G. Bertorelle, *et al.*, Genetic load: genomic estimates and applications in non-
model animals. *Nature Reviews Genetics* **23**, 492–503 (2022).

10.   M. Lynch, W. Gabriel, Mutation Load and the Survival of Small Populations.
Evolution **44**, 1725–1737 (1990).

11.   D. L. Gilroy, K. P. Phillips, D. S. Richardson, C. van Oosterhout, Toll-like receptor
variation in the bottlenecked population of the Seychelles warbler: computer simulations
see the 'ghost of selection past' and quantify the 'drift debt'. *Journal of Evolutionary*
Biology **30**, 1276–1287 (2017).

12.   A. V. Pinto, B. Hansson, I. Patramanis, H. E. Morales, C. van Oosterhout, The
impact of habitat loss and population fragmentation on genomic erosion. *Conservation*
Genetics **25**, 49–57 (2024).

13.   H. A. Jackson, *et al.*, Genomic erosion in a demographically recovered bird
species during conservation rescue. *Conservation Biology* **36**, e13918 (2022).

14.   M. Dehasque, *et al.*, Temporal dynamics of woolly mammoth genome erosion
prior to extinction. *Cell* **187**, 3531–3540.e13 (2024).

15.   A. Andersson, S. Karlsson, N. Ryman, L. Laikre, Monitoring genetic diversity with
new indicators applied to an alpine freshwater top predator. *Molecular Ecology* **31**,
6422–6439 (2022).

16.   D. O'Brien, *et al.*, Bringing together approaches to reporting on within species
genetic diversity. *Journal of Applied Ecology* **59**, 2227–2233 (2022).

17.   K. Theissinger, *et al.*, How genomics can help biodiversity conservation. *Trends
in Genetics* **39**, 545–559 (2023).

18.   D. Díez-del-Molino, F. Sánchez-Barreiro, I. Barnes, M. T. P. Gilbert, L. Dalén,
Quantifying temporal genomic erosion in endangered species. *Trends in Ecology &
Evolution* **33**, 176–185 (2018).

19.   H. Ellegren, N. Galtier, Determinants of genetic diversity. *Nature Reviews
Genetics* **17**, 422–433 (2016).

20.   J. Romiguier, *et al.*, Comparative population genomics in animals uncovers the
determinants of genetic diversity. *Nature* **515**, 261–263 (2014).

21.   A. Mackintosh, *et al.*, The determinants of genetic diversity in butterflies. *Nature
Communications* **10**, 3466 (2019).

22.   G. Hewitt, The genetic legacy of the Quaternary ice ages. *Nature* **405**, 907–913
(2000).

23.   J. A. Robinson, *et al.*, Genomic flatlining in the endangered island fox. *Current
Biology* **26**, 1183–1189 (2016).

24.   J. A. Robinson, *et al.*, The critically endangered vaquita is not doomed to
extinction by inbreeding depression. *Science* **376**, 635–639 (2022).

25.   M. F. DiLeo, A. Nair, M. Kardos, A. Husby, M. Saastamoinen, Demography and
environment modulate the effects of genetic diversity on extinction risk in a butterfly
metapopulation. *Proceedings of the National Academy of Sciences* **121**, e2309455121
(2024).

26.   N. Dussex, H. E. Morales, C. Grossen, L. Dalén, C. van Oosterhout, Purging and
accumulation of genetic load in conservation. *Trends in Ecology & Evolution* **38**, 961–
969 (2023).

27.   M. K. Schwartz, G. Luikart, R. S. Waples, Genetic monitoring as a promising tool
for conservation and management. *Trends in Ecology & Evolution* **22**, 25–33 (2007).

28.   S. Hoban, *et al.*, Effective population size remains a suitable, pragmatic indicator
of genetic diversity for all species, including forest trees. *Biological Conservation* **253**,
108906 (2021).

29.   M. Ihse, Swedish agricultural landscapes patterns and changes during the last
50 years, studied by aerial photos. *Landscape and Urban Planning* **31**, 21–37 (1995).

30.   R. Lindborg, O. Eriksson, Historical Landscape Connectivity Affects Present
Plant Species Diversity. *Ecology* **85**, 1840–1845 (2004).

31.   D. L. Wagner, R. Fox, D. M. Salcido, L. A. Dyer, A window to the world of global
insect declines: Moth biodiversity trends are complex and heterogeneous. *Proceedings*
of the National Academy of Sciences **118**, e2002549117 (2021).

32.   M. S. Warren, *et al.*, The decline of butterflies in europe: Problems, significance,
and possible solutions. *Proceedings of the National Academy of Sciences* **118**,
e2002551117 (2021).

33.   J. C. Habel, *et al.*, Butterfly community shifts over two centuries. *Conservation*
Biology **30**, 754–762 (2016).

34.   SLU Artdatabanken, *Rödlistade arter i Sverige 2020* (SLU, 2020).

35.   C. Fägerström, Lund University Biological Museum - Insect collections Inventory.
(2024). https://doi.org/10.15468/dahk2a.

36.   Z. J. Nolen, M. Rundlöf, A. Runemark, Species-specific erosion of genetic
diversity in grassland butterflies depends on landscape land cover. *Biological*
Conservation **296**, 110694 (2024).

37.   M. T. P. Gilbert, *et al.*, Recharacterization of ancient DNA miscoding lesions:
Insights in the era of sequencing-by-synthesis. *Nucleic Acids Research* **35**, 1–10
(2007).

38.   M. Stiller, *et al.*, Patterns of nucleotide misincorporations during enzymatic
amplification and direct large-scale sequencing of ancient DNA. *Proceedings of the*
National Academy of Sciences **103**, 13578–13584 (2006).

39.   H. Jónsson, A. Ginolhac, M. Schubert, P. L. F. Johnson, L. Orlando,
mapDamage2.0: Fast approximate bayesian estimates of ancient DNA damage
parameters. *Bioinformatics* **29**, 1682–1684 (2013).

40.   P. W. Hedrick, *Genetics of Populations* (Jones & Bartlett Publishers, 2009).

41.   R. McQuillan, *et al.*, Runs of homozygosity in european populations. *The*
American Journal of Human Genetics **83**, 359–372 (2008).

42.   E. A. Thompson, Identity by descent: Variation in meiosis, across genomes, and
in populations. *Genetics* **194**, 301–326 (2013).

43.   S. Glémin, How are deleterious mutations purged? Drift versus nonrandom
mating. *Evolution* **57**, 2678–2687 (2003).

44.   SLU Artdatabanken, Artfakta: kronärtsblåvinge (Plebejus argyrognomon).

45.   J. C. Habel, M. J. Samways, T. Schmitt, Mitigating the precipitous decline of
terrestrial European insects: Requirements for a new strategy. *Biodiversity and*
Conservation **28**, 1343–1360 (2019).

46.   E. L. Landguth, *et al.*, Quantifying the lag time to detect barriers in landscape
genetics. *Molecular Ecology* **19**, 4179–4191 (2010).

47.   C. W. Epps, N. Keyghobadi, Landscape genetics in a changing world:
disentangling historical and contemporary influences and inferring change. *Molecular*
Ecology **24**, 6021–6040 (2015).

48.   W. H. Lowe, F. W. Allendorf, What can genetics tell us about population
connectivity? *Molecular Ecology* **19**, 3038–3051 (2010).

49.   P. W. Hedrick, J. R. Adams, J. A. Vucetich, Reevaluating and Broadening the
Definition of Genetic Rescue. *Conservation Biology* **25**, 1069–1070 (2011).

50.   P. W. Hedrick, A. Garcia-Dorado, Understanding inbreeding depression, purging,
and genetic rescue. *Trends in Ecology & Evolution* **31**, 940–952 (2016).

51.   D. A. Bell, *et al.*, The Exciting Potential and Remaining Uncertainties of Genetic
Rescue. *Trends in Ecology & Evolution* **34**, 1070–1079 (2019).

52.   R. Lande, Risk of population extinction from fixation of new deleterious
mutations. *Evolution* **48**, 1460–1469 (1994).

53.   S. Mathur, J. A. DeWoody, Genetic load has potential in large populations but is
realized in small inbred populations. *Evolutionary Applications* **14**, 1540–1557 (2021).

54.   I. Saccheri, *et al.*, Inbreeding and extinction in a butterfly metapopulation. *Nature*
392, 491–494 (1998).

55.   C. D. Huber, B. Y. Kim, K. E. Lohmueller, Population genetic models of GERP
scores suggest pervasive turnover of constrained sites across mammalian evolution.
PLOS Genetics **16**, e1008827 (2020).

56.   C. C. Kyriazis, J. A. Robinson, K. E. Lohmueller, Using computational simulations
to model deleterious variation and genetic load in natural populations. *The American*
Naturalist **202**, 737–752 (2023).

57.   N. Dussex, *et al.*, Population genomics of the critically endangered kākāpō. *Cell*
Genomics **1**, 100002 (2021).

58.   T. van der Valk, D. Díez-del-Molino, T. Marques-Bonet, K. Guschanski, L. Dalén,
Historical Genomes Reveal the Genomic Consequences of Recent Population Decline
in Eastern Gorillas. *Current Biology* **29**, 165–170.e6 (2019).

59.   J. von Seth, *et al.*, Genomic insights into the conservation status of the world's
last remaining Sumatran rhinoceros populations. *Nature Communications* **12**, 2393
(2021).

60.   C. Bortoluzzi, *et al.*, Lepidoptera genomics based on 88 chromosomal reference
sequences informs population genetic parameters for conservation. *bioRxiv* (2023).
https://doi.org/10.1101/2023.04.14.536868.

61.   L. B. Pettersson, H. Arnberg, *Svensk dagfjärilsövervakning, årsrapport för 2023*
(Biologiska institutionen, Lunds universitet, 2024).

62.   S. Hoban, *et al.*, Genetic diversity targets and indicators in the CBD post-2020
Global Biodiversity Framework must be improved. *Biological Conservation* **248**, 108654
(2020).

63.   L. Laikre, *et al.*, Post-2020 goals overlook genetic diversity. *Science* **367**, 1083–
1085 (2020).

64.   G. Formenti, *et al.*, The era of reference genomes in conservation genomics.
Trends in Ecology & Evolution **37**, 197–202 (2022).

65.   V. G. Twort, J. Minet, C. W. Wheat, N. Wahlberg, Museomics of a rare taxon:
placing Whalleyanidae in the Lepidoptera Tree of Life. *Systematic Entomology* **46**, 926–
937 (2021).

66.   Z. J. Nolen, Zjnolen/PopGLen: V0.3.0. (2024).
https://doi.org/10.5281/zenodo.13785563. Deposited September 2024.

67.   F. Mölder, *et al.*, Sustainable data analysis with snakemake [version 2; peer
review: 2 approved]. *F1000Research* **10** (2021).

68.   S. Chen, Y. Zhou, Y. Chen, J. Gu, Fastp: An ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

69.   H. Li, Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. (2013). https://doi.org/10.48550/arXiv.1303.3997.

70.   H. Li, R. Durbin, Fast and accurate long-read alignment with BurrowsWheeler
transform. *Bioinformatics* **26**, 589–595 (2010).

71.   E. Palkopoulou, *et al.*, Complete genomes reveal signatures of demographic and
genetic declines in the woolly mammoth. *Current biology: CB* **25**, 1395–1400 (2015).

72.   P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *GigaScience* **10**
(2021).

73.   Picard toolkit. *Broad Institute, GitHub Repository* (2019). Deposited 2019.

74.   A. Peltzer, *et al.*, EAGER: efficient ancient genome reconstruction. *Genome*
Biology **17**, 1–14 (2016).

75.   G. A. van der Auwera, B. D. O'Connor, *Genomics in the Cloud: Using Docker,*
GATK, and WDL in Terra (O'Reilly Media, Inc., 2020).

76.   R. N. Lou, A. Jacobs, A. P. Wilder, N. O. Therkildsen, A beginner's guide to low-
coverage whole genome sequencing for population genomics. *Molecular Ecology* **30**,
5966–5993 (2021).

77.   G. Jun, M. K. Wing, G. R. Abecasis, H. M. Kang, An efficient and scalable
analysis framework for variant extraction and refinement from population-scale DNA
sequence data. *Genome Research* **25**, 918–925 (2015).

78.   J. Neukamm, A. Peltzer, K. Nieselt, DamageProfiler: Fast damage pattern
calculation for ancient DNA. *Bioinformatics* **37**, 3652–3653 (2021).

79.   P. Pečnerová, *et al.*, High genetic diversity and low differentiation reflect the
ecological versatility of the African leopard. *Current Biology* **31**, 1862–1871.e5 (2021).

80.   A. F. A. Smit, R. Hubley, P. Green, RepeatMasker open-4.0. (2013). Deposited
2013.

81.   S. Sawyer, J. Krause, K. Guschanski, V. Savolainen, S. Pääbo, Temporal
Patterns of Nucleotide Misincorporations and DNA Fragmentation in Ancient DNA.
PLOS ONE **7**, e34131 (2012).

82.   G. Femerling, *et al.*, Genetic Load and Adaptive Potential of a Recovered Avian
Species that Narrowly Avoided Extinction. *Molecular Biology and Evolution* **40**,
msad256 (2023).

83.   S. Gopalakrishnan, *et al.*, The population genomic legacy of the second plague
pandemic. *Current Biology* **32**, 4743–4751.e6 (2022).

84.   T. S. Korneliussen, A. Albrechtsen, R. Nielsen, ANGSD: Analysis of Next
Generation Sequencing Data. *BMC Bioinformatics* **15**, 1–13 (2014).

85.   G. Bhatia, N. Patterson, S. Sankararaman, A. L. Price, Estimating and
interpreting FST: The impact of rare variants. *Genome Research* **23**, 1514–1521 (2013).

86.   R. K. Waples, A. Albrechtsen, I. Moltke, Allele frequency-free inference of close
familial relationships from genotypes or low-depth sequencing data. *Molecular Ecology*
28, 35–48 (2019).

87.   K. Hanghøj, I. Moltke, P. A. Andersen, A. Manica, T. S. Korneliussen, Fast and
accurate relatedness estimation from high-throughput sequencing data in the presence
of inbreeding. *GigaScience* **8**, giz034 (2019).

88.   A. Manichaikul, *et al.*, Robust relationship inference in genome-wide association
studies. *Bioinformatics* **26**, 2867–2873 (2010).

89.   L. Skotte, T. S. Korneliussen, A. Albrechtsen, Association testing for next-
generation sequencing data using score statistics. *Genetic Epidemiology* **36**, 430437
(2012).

90.   S. Y. Kim, *et al.*, Estimation of allele frequency and association mapping using
next-generation sequencing data. *BMC Bioinformatics* **12**, 231 (2011).

91.   J. Meisner, A. Albrechtsen, Inferring population structure and admixture
proportions in low-depth NGS data. *Genetics* **210**, 719–731 (2018).

92.   E. A. Fox, A. E. Wright, M. Fumagalli, F. G. Vieira, ngsLD: Evaluating linkage
disequilibrium using genotype likelihoods. *Bioinformatics* **35**, 3855–3856 (2019).

93.   L. Skotte, T. S. Korneliussen, A. Albrechtsen, Estimating individual admixture
proportions from next generation sequencing data. *Genetics* **195**, 693–702 (2013).

94.   G. Garcia-Erill, A. Albrechtsen, Evaluation of model fit of inferred admixture
proportions. *Molecular Ecology Resources* **20**, 936–949 (2020).

95.   V. Narasimhan, *et al.*, BCFtools/RoH: A hidden markov model approach for
detecting autozygosity from next-generation sequencing data. *Bioinformatics* **32**, 1749–
1751 (2016).

96.   E. V. Davydov, *et al.*, Identifying a High Fraction of the Human Genome to be
under Selective Constraint Using GERP++. *PLOS Computational Biology* **6**, e1001025
(2010).

97.   W. McLaren, *et al.*, The Ensembl Variant Effect Predictor. *Genome Biology* **17**,
122 (2016).

98.   V. E. Kutschera, *et al.*, GenErode: a bioinformatics pipeline to investigate
genome erosion in endangered and extinct species. *BMC Bioinformatics* **23**, 1–17
(2022).

99.   G. M. Cooper, *et al.*, Distribution and intensity of constraint in mammalian
genomic sequence. *Genome Research* **15**, 901–913 (2005).

100.   S. Kumar, *et al.*, TimeTree 5: An expanded resource for species divergence
times. *Molecular Biology and Evolution* **39**, msac174 (2022).

101.   D. Charlesworth, J. H. Willis, The genetics of inbreeding depression. *Nature
Reviews Genetics* **10**, 783–796 (2009).

102.   K. Johannesson, L. Laikre, Monitoring of genetic diversity in environmental
monitoring [in swedish]. *Report to the Swedish Agency for Marine and Water
Management (dnr. HaV 3642-2018, 3643-2018)* (2020).

Figure Captions

Figure 1. Sampling design and changes in genetic diversity, differentiation, and inbreeding in the focal species. (A) Images of the three focal species. (B) Map of the sampling area in southern Sweden, sampling locations are approximate. Circles represent areas with modern samples in at least one of the species and triangles represent areas with historical samples, see Table S1 for complete sampling information. (C) Change in individual heterozygosity over time, measured as heterozygous sites per 1000bp, excluding transitions. Each point represents an individual sample, grouped into historical and modern samples per species. Coloring refers to sampling region and corresponds to the map in panel B. (D) Change in inbreeding coefficients over time, estimated using runs of homozygosity larger than 100kb (F_{RoH}). (E) Change in pairwise population genetic differentiation, F_{ST} over time between equivalent comparisons in each time period. Color of the points and line refer to the geographic distance between populations. Populations with no connecting line refer to pairs of modern sample populations that have no comparable historic pair. In C & D, results of unequal variance t-tests between historical and modern samples for each species is shown using the following scale, where panels without * are not significant ($P > 0.05$): * $P <= 0.05$, ** $P <= 0.01$, *** $P <= 0.001$, **** $P <= 0.0001$. Statistical assessment of genetic differentiation is not shown, as it could not be assessed for all three species (Table S7).

Figure 2. Population structure of the focal species. (A) Principal Component Analysis (PCA) of the three focal species. Shapes refer to sampling period, with each circle representing a modern and each triangle a historical individual, with colors corresponding to the sampling locations in Figure 1B. Percentages on each axis refer to the proportion of total variance explained by the component. (B) Best fit admixture analyses for the three focal species. Each bar represents an individual and each color one of K genetic clusters. Each bar is colored in proportion to the probability of the individual belonging to each cluster.

Figure 3. Changes in genetic load of the focal species (A) Individual inbreeding coefficients (F_{RoH}) partitioned by *RoH* size class. Longer *RoH* indicate more recent common ancestry of the identical haplotypes. (B) Total genetic load measured for conserved regions, high impact variants, and moderate impact variants. Total genetic load is measured for conserved region as the derived allele counts scaled by their genomic conservation (GERP) scores per 1000 conserved derived alleles. For high and moderate impact variants, total load is measured as alternate allele counts for the given impact category per 1000 alternate alleles. (C) Realized genetic load measured for the same three variant categories. Realized genetic load is measured for conserved region as the derived homozygous allele counts scaled by their conservation (GERP) scores per 1000 conserved derived alleles. For high and moderate impact variants, realized

load is measured as alternate allele counts in homozygous state for the given impact category per 1000 alternate alleles. In B & C, results of unequal variance t-tests between historical and modern samples are shown in each panel, where panels without * are not significant ($P > 0.05$): * $P <= 0.05$, ** $P <= 0.01$, *** $P <= 0.001$, **** $P <= 0.0001$.

Figure 4. Changes in heterozygosity predicted over 100 years based on the rate of change estimated from modern and historical samples for each of 11 species. Species are grouped by conservation status, which follows the species name on the x-axis. Background coloration refers to the three genetic decline thresholds outlined by Andersson et al (15), where a retention of >=95% over 100 year is considered 'acceptable' (green), a retention of 75-94% as 'warning' (yellow), and a retention of <75% as 'alarm' (red). Points represent mean estimates for change in heterozygosity across all sample pairs for a species. Species with bars have more than one possible sample pair, with error bars illustrating the spread of estimates across all sample pairs. Species with a * next to their species name on the x-axis have been mapped to a related species' reference genome as they lack a species-specific genome (Table S12).