Species-specific loss of genetic diversity and inbreeding following
agricultural intensification

Zachary J. Nolen[1*], Patrycja Jamelska[1], Ana Sofia Torres Lara[1,2], Niklas Wahlberg[1], Anna Runemark[1]

[1] Department of Biology, Lund University, Lund, Sweden 22362
[2] Faculty 2 Biology/Chemistry, University of Bremen, Bremen, Germany 28359

*Corresponding author: Zachary J. Nolen

Email: zachary.nolen@biol.lu.se

Abstract

Land-use change from agricultural intensification is a major driver of global insect declines. We

investigated the genetic consequences of grassland decline in Sweden by sequencing museum and

modern specimens of three focal and five additional Polyommatinae butterfly species. From 59 historical

and 90 contemporary genomes, we find a 6% decline in genetic diversity and increased isolation and

inbreeding in the grassland specialist *Cyaniris semiargus* over the past 70 years. In contrast, generalist

Polyommatus icarus and heathland specialist *Plebejus argus* maintained genetic diversity and

connectivity. Although currently genomic erosion is mild, we infer it lags considerably larger declines in

effective population sizes. Using simulations, we demonstrate that only a small portion of genetic diversity

is lost in early stages of decline, creating a genetic extinction debt. Two of five additional species show

similar reductions, underscoring that restoration of grassland habitat is necessary to restore gene flow

and halt further genomic erosion in grassland insects.

Introduction

Insect decline has received increased attention in recent years as reports uncover alarming declines in insect assemblages and biomass (*1*). Efforts to quantify this decline estimate that terrestrial insects are declining at an average rate of 11% per decade globally, with this trend being primarily driven by even stronger regional declines in Europe and North America (*2*). This stark decline is the result of a variety of anthropogenic pressures, primarily climate change and land use changes such as agricultural intensification and urbanization (*1*, *3–7*). In Europe, formerly biodiverse semi-natural grasslands have been abandoned or converted into intensively cultivated land (*8*), placing a strong pressure on grassland insect species (*6*).

While insect declines have alarming implications for the natural communities and ecosystem services they play key roles in, declines also lead to genomic erosion within insect populations themselves. As populations decline, the relative strength of drift increases (*9*), eroding genetic diversity, thereby reducing adaptive potential (*10*, *11*). Additionally, in small, isolated populations, deleterious mutations are more likely to accumulate through drift and become unmasked through inbreeding (*12–17*), further reinforcing population decline (*18*, *19*). Mitigation of genomic erosion that threatens population persistence has been made a part of Goal A of the Kunming-Montréal Global Biodiversity Framework, which stipulates maintenance of genetic diversity to preserve the adaptive potential of wild populations (*20*).

Monitoring approaches to assess genomic erosion are under development. While plans to monitor genetic diversity in the wild have been implemented in several countries (*21–24*), interpreting measures of genetic diversity remains a challenge. First, genomic erosion lags disturbance events, and current estimates of genetic diversity will therefore not capture the genetic extinction debt of populations (*25–30*). Therefore, identifying measures that enable early detection of genomic erosion is a priority. Second, absolute levels of genetic diversity are influenced by a multitude of factors not related to threat level and vary considerably across taxa and populations (*31–34*). To control for variation in baseline levels of genetic diversity, programs in several countries supplement estimates of contemporary effective population sizes with measures of genetic diversity metrics across multiple time points (*22*, *35*).

Temporal estimates of genetic diversity enable direct quantification of genomic erosion but are limited to detecting changes after the first sampled time point. To overcome this, genomic resources from historic samples preserved in natural history museums provide an opportunity to establish baseline estimates of genetic diversity from before major anthropogenically driven insect declines (*24*, *36*). Using comparisons to museum specimens, populations already undergoing genomic erosion can be identified with only one contemporary sampling point, reducing time to detection of populations at risk.

Here, we use museomics to address to what extent genetic diversity and connectivity has been lost in grassland associated butterflies during the second wave of agricultural modernization in Sweden (*37*, *38*). Starting in the 1940s, small farms were abandoned, and their associated semi-natural grasslands began

to decline considerably (*38*). We examine three focal butterfly species with Swedish Red List statuses of
'least concern' (*39*) in this landscape. These species vary in habitat specialization, ranging from grassland
generalist to specialist, enabling us to examine if patterns of genetic diversity decline are stronger in more
specialized species, in line with expectations from greater demographic declines in specialist insect
abundance compared to generalists (*40–42*). We assess genetic change by comparing museum and
contemporary estimates of genetic diversity and use simulations to assess if the lag in loss of diversity is
expected given the scale of demographic change. We further examine if changes in genetic diversity are
common across grassland butterfly species by extending the dataset to include historical and modern
individuals of five additional related butterfly species with Swedish Red List statuses ranging from least
concern to endangered (*39*). In the face of ongoing mass extinction, identifying species and populations
at risk at an early stage of decline is paramount. Understanding early signals of genomic erosion is key
for intervention before substantial diversity is irreversibly lost. Here, by combining a large set of historical
and contemporary specimens, we uncover a case of ongoing genomic erosion in response to recent
anthropogenic change and highlight metrics that respond quickly to population decline, providing insights
into genomic erosion at its early stages when intervention is most effective.

Results

We analyzed changes in genetic diversity, differentiation, and proxies for genetic load in three focal
Polyommatinae (Lepidoptera: Lycaenidae) butterfly species: *Polyommatus icarus,* a grassland generalist,
Plebejus argus, a heathland specialist, and *Cyaniris semiargus*, a grassland specialist (Figure 1A). We
generated whole genome resequencing data from pinned museum specimens from the entomological
collections of the Biological Museum at Lund University (*43*) collected from 1917-1956 (Table S1; $N = 28$ -
Po. icarus; 12 - *Pl. argus*; 14 - *Cy. semiargus*) in southern Sweden. These individuals were sampled as
populations by grouping specimens generally collected at the same locality during the same year and
compared with WGS data from samples collected at corresponding localities in 2020-2023 by Nolen et al
(*44*) (Figure 1B; Table S1; $N = 27$ - *Po. icarus*; 29 - *Pl. argus*; 29 - *Cy. semiargus*). As DNA extracted from
historical specimens is fragmented and subject to post-mortem DNA damage (*45, 46*) (Figure S1), we
removed transitions from all analyses. As the shorter fragment size of historical samples can result in
poor mapping of reads containing non-reference alleles, we constructed variation graphs for each
reference genome containing high quality variants called in the modern samples. Mapping both sample
types to these variation graphs reduced reference bias compared to aligning to the linear reference
genome, as well as reduced the disparity in reference bias between historical and modern samples,
especially when calling genotypes (Figure S2&3; Supplementary Methods). The reduction in reference
bias and its influence on downstream analyses is clearly visible in population structure analyses, where
using the linear reference alignments resulted in samples clustering primarily by sample type (historical or
modern), while using the variation graph alignments resulted in samples clustering primarily by sampling
locality (Figure S4). While the variation graph alignment method reduced reference bias in historical
samples, it was not eliminated, and results should be considered with this in mind, especially for *Po.*

icarus which exhibits the largest residual bias (Figure S2&4). Median sequencing depth was 6.4x (range =

2.2 to 16.7x) for historical samples and 19.3x (range = 7.0 to 56.7x; Table S1) for modern samples. To

account for variation in sequencing depth, all analyses were performed with data subsampled to between

6 and 8x sequencing depth.

Figure 1. Sampling design and changes in genetic diversity, differentiation, and inbreeding in the focal species. (A) Images of the three focal species. (B) Map of the sampling area in southern Sweden, sampling locations are approximate. Circles represent areas with modern samples in at least one of the species and triangles represent areas with historical samples, see Table S1 for complete sampling information. (C) Change in individual heterozygosity over time, measured as heterozygous sites per 1000bp, excluding transitions. Each point represents an individual sample, grouped into historical and modern samples per species. Coloring refers to sampling region and corresponds to the map in panel B. (D) Change in inbreeding coefficients over time, estimated using runs of homozygosity larger than 100kb (F_{RoH}). (E) Change in pairwise population genetic differentiation, $_{ST}$ over time between equivalent comparisons in each time period. Color of the points and line refer to the geographic distance between populations. Populations with no connecting line refer to pairs of modern sample populations that have no comparable historic pair. In C & D, results of linear mixed models assessing the effect of time period on the response are shown using the following scale, where panels without * are not significant (P > 0.05): * P <= 0.05, ** P <= 0.01, *** P <= 0.001.

Reduced genetic diversity and effective population size in Cy. semiargus

To assess if agricultural intensification and the associated reduction in semi-natural grassland habitat has

resulted in reduced genetic diversity, we compared individual genome-wide heterozygosity of historical

and modern samples within each species. Individual heterozygosity is significantly reduced in the

specialist *Cy. semiargus* (percent difference: -6%; LMM: *P* = 0.008; Figure 1C; Table S2&3), but not in

Po. icarus or *Pl. argus*. Population specific changes in genetic diversity estimated both from mean

heterozygosity and nucleotide diversity (Table S4) suggest that genetic diversity has mostly been reduced

in populations in Skåne, where arable land cover is greatest, with only a 1% change in nucleotide

diversity in Småland (Tables S5&6).

To understand the recent demographic patterns contributing to contemporary genetic diversity, we

assessed changes in effective population size (N_e) over the past century using the modern genomes with

a linkage disequilibrium (LD) based approach. We find stable and very high effective population sizes in

Po. icarus and fluctuating or only recently declined sizes in *Pl. argus*, in line with their maintenance of

genetic diversity (Figure S5; Table S6). In *Cy. semargus*, we find that all populations in Skåne have gone

through population declines within the past 30 generations, with current effective population sizes of 35

(SE Skåne), 202 (W Skåne), and 520 (E Skåne) (Figure S5; Table S6).

Reduced connectivity may drive loss of genetic diversity

While genetic diversity decline is an expected product of demographic decline, it can also be a product of

reductions in gene flow (*9*). Modern *Cy. semiargus* populations exhibit stronger population structure than

historical ones, based on both genetic differentiation and clustering approaches. For population pairs

represented in both modern and historical samplings, modern estimates of genetic differentiation (F_{ST}) in

Cy. semiargus have increased from an average of 0.041 to 0.065, with the increase being largest

between W Skåne and Småland (Figure 1E; Table S7). Including population pairs in the modern sampling

that are not represented in the historical, but that include similarly geographically separated populations,

increases the average F_{ST} of modern *Cy. semiargus* pairs to 0.086. For *Po. icarus* and *Pl. argus*, average

F_{ST} between pairs represented in both time periods has changed from 0.018 to 0.011 and 0.034 to 0.038

(Figure 1E; Table S7). When including additional modern pairs in *Pl. argus*, the modern average is slightly

lower, 0.035, though this includes pairs at short distances that are not represented in historical pairs.

Historical population pairs are sampled at different time points, with pairs including populations sampled

up to 20 years apart (Table S7), which could inflate historical F_{ST} estimates.

Principal component analyses (PCAs) and admixture analyses also reveal stronger population structure

in modern *Cy. semiargus* samples than historical (Figure 2A). Each modern population forms a distinct

cluster in the specialist *Cy. semiargus*, with modern Småland samples clustering with their historical

counterparts (Figure 2B). Historical samples from Western and Eastern Skåne cluster together in a

separate cluster from the respective modern samples, highlighting that the stronger genetic structure

found in modern samples is largely driven by isolation of populations in the agricultural landscapes within

Sweden's southernmost province, Skåne (Figure 1D; Figure 2A&B; Figure S6). Populations of both *Po.*

icarus and *Pl. argus* from Skåne separate from those sampled elsewhere in southern Sweden along the

major axis of variation, with historical samples clustering with geographically proximal modern samples.

Admixture analyses support only one genetic cluster in each of these species (Table S8), but the

correlation of residuals suggest additional genetic clusters in *Pl. argus* (Figure S7), consistent with the

PCA. This highlights that while genetic structure likely exists, the variation is too low for reliable clustering
at these sample sizes. We additionally find that historical and modern samples separate out on PC2 in
Po. icarus, potentially an artifact of residual reference bias in historical samples of this species (Figure
S2). The general pattern of historical populations clustering in the PCA with at least one geographically
proximal modern population suggests that historical samples are representative of ancestral populations
to the modern ones.

Figure 2. Population structure of the focal species. (A) Principal Component Analysis (PCA) of the
three focal species. Shapes refer to sampling period, with each circle representing a modern and each
triangle a historical individual, with colors corresponding to the sampling locations in Figure 1B.
Percentages on each axis refer to the proportion of total variance explained by the component. (B)
Admixture analyses for the three focal species. Each color represents one of K genetic clusters, with bars
representing individuals, colored by their probability of belonging to a given cluster. For Po. icarus and Pl.
argus, the best fit was at K = 1, but K = 2 is shown for illustrative purposes. For Cy. semiargus, the best fit
K = 5 is shown.

Increased inbreeding in specialist Cy. semiargus

To investigate changes in rates of inbreeding, we estimated the inbreeding coefficient F_{RoH}, which is the

proportion of the autosomal genome in runs of homozygosity (*RoH*) greater than a certain length (*47*). We

quantified F_{RoH} for *RoH* longer than 100kb, corresponding roughly to *RoH* with shared ancestry in the past

200 generations (*48*). We found a significant increase in F_{RoH} from historical to modern *Cy. semiargus*

samples (difference: 0.065, LMM: $P < 0.01$; Figure 1D; Table S9). Across species, the highest inbreeding

coefficient for a historical sample is 0.031, with historical species averages all below 0.01 (Tables S2&9).

In *Cy. semiargus*, modern samples have a maximum coefficient of 0.329 and an average coefficient of

0.073, an order of magnitude higher than the mean for historical samples. Individual inbreeding

coefficients were largely determined by runs of homozygosity >= 1.5 Mb in length (Figure 3A),

corresponding to low effective population sizes in the past ~12 generations, in line with estimates of

recently low effective population size in our demographic analyses. In *Po. icarus* and *Pl. argus*, both

modern and historical estimates of F_{RoH} are low, with averages <= 0.01 and no evidence of significant

increases (Table S9).

To assess if reductions in heterozygosity explained by increases in F_{RoH} match theoretical expectations,

we performed coalescent simulations parameterized on the recent demographic histories inferred from

the modern genomic data, estimating heterozygosity and F_{RoH} for individuals sampled at time points

corresponding to our modern and historical samplings. In all simulations, genome-wide heterozygosity

decline in this time frame was primarily due to the presence of long runs of homozygosity, rather than an

even reduction in heterozygosity across the genome (Table S10). To assess if we observed this same

pattern in the empirical genomic data, we adjusted estimates of individual heterozygosity by the

inbreeding coefficient F_{RoH} per individual. Individual heterozygosity outside *RoH* is not different between

time periods (Figure S8), agreeing with the simulation predictions. Additionally, for all simulated

populations except for Småland *Cy. semiargus*, the estimated ratios of modern/historical heterozygosity

from the genomic data fell within ranges predicted by the simulations, as did average estimates of F_{RoH}

(Figure S9). In the case of Småland *Cy. semiargus*, we estimated a reduction in heterozygosity and

increase in F_{RoH} from the genomic data, but no change was predicted by the simulations. As the *RoH*

estimated from the modern sample suggest some level of recent demographic decline that is absent from

the inferred demographic history, this may reflect a limitation of the inferred history for this population

used to parameterize the simulations.

Figure 3. Temporal changes in derived count of putatively deleterious mutations in the focal species (A) Individual inbreeding coefficients (F_{RoH}) partitioned by RoH size class. Longer RoH indicate more recent common ancestry of the identical haplotypes. (B) Total count of derived alleles per 1000 called genotypes for individuals of the three focal species at positions classifying three categories of putatively deleterious mutations: I) missense (i.e. weakly deleterious; upper), II) loss of function (LoF, i.e. strongly deleterious; middle), and III) conserved positions (presumed strongly deleterious; lower). C) Count of derived alleles in homozygous state per 1000 called genotypes for the same mutation categories as in B. In B & C, results of linear mixed models with burden as a response to time period are shown in each panel, where panels without * are not significant (P > 0.05): * P <= 0.05, ** P <= 0.01, *** P <= 0.001.

Increased homozygosity of weakly deleterious mutations in Cy. semiargus

Population decline and subdivision can increase the exposure of partially recessive, deleterious mutations to selection due to increased homozygosity, facilitating purging of strongly deleterious mutations (*15, 49,*

50). However, as populations decline and subdivide, selection weakens relative to drift, allowing weakly
deleterious mutations to drift to higher frequencies and even fix, increasing genetic load (*14*, *51*, *52*).
Here, as a proxy for genetic load, we estimated deleterious mutation burden per individual using derived
counts of putatively deleterious mutations (as in e.g. *53–57*). Deleterious mutations were identified based
off the functional annotation or the sequence conservation score of the position and divided into three
categories: (I) missense, i.e. weakly deleterious; (II) loss of function (LoF), i.e. strongly deleterious; and
(III) conserved, i.e. with high sequence conservation across Lycaenidae, presumed to be strongly
deleterious. We partitioned these counts in to total derived counts, approximating load if mutations are
primarily additive, and homozygous derived counts, reflecting load if mutations are primarily recessive.

We find no evidence of changes in the total derived count for any of the three mutation categories in *Po.*
icarus or *Cy. semiargus*. However, in *Pl. argus*, total derived count at missense positions is reduced
(LMM: $P < 0.001$; Figure 3B; Table S10), but there is no significant change in any other mutation category
or at either intergenic or synonymous genic positions (Table S11). However, homozygous derived count
has increased at missense and conserved positions in *Cy. semiargus* (LMM: $P = 0.002$ and 0.006;
Figure 3C; Table S10), and at conserved positions in *Pl. argus* (LMM: $P = 0.006$; Figure 3B; Table S10),
but not changed over time in *Po. icarus. Cyaniris semiargus* individuals with higher inbreeding coefficients
also had higher homozygous derived counts across all categories of deleterious mutations (LM: $R^2 =$
0.83, 0.40, 0.76 for missense, LoF, and conserved positions, respectively; Figure S10). There is also a
weaker increase of total derived count with inbreeding coefficient at LoF and conserved positions (LM: R^2
= 0.12, 0.12, respectively). There is no change in homozygous derived count at LoF positions over time in
any species.

Genetic diversity indicators

We utilized the genetic diversity indicator thresholds outlined by Andersson et al. (*21*) to assess if
genomic erosion is reflected in conservation status in the three focal species as well as historical-modern
pairs of five additional species of Polyommatinae butterfly (Table S1). The thresholds suggest
categorizations of 'Acceptable', 'Warning', and 'Alarm' rates of genetic erosion, reflecting >=95%, 75-
94%, and <75% retention of genetic diversity over 100 years, respectively (*21*). Based on changes in
mean population heterozygosity and nucleotide diversity, populations of all three species largely fall in the
'Acceptable' category, except for *Cy. semiargus* in Southeast and Western Skåne, which fall into
'Warning' category (Figure 4; Tables S5&6). The same indicator framework in Sweden categorizes
effective population sizes >= 500 as 'Acceptable', 51 - 499 as 'Warning', and <= 50 as 'Alarm' (*21*, *58*).
Under this indicator, all *Po. icarus* and *Pl. argus* populations and Småland *Cy. semiargus* would be
classed as 'Acceptable'. The low effective population sizes in Skåne for *Cy. semiargus* would place
Eastern Skåne as 'Acceptable', but bordering on 'Warning', Western Skåne as 'Warning', and
Southeastern Skåne as 'Alarm' (Table S6).

We also assessed if the conservation status of the five additional species, ranging from 'Least Concern'

to 'Endangered' (*39*) (Table S12) are associated with genetic erosion. As a single pair of individuals were

sampled from each additional species, we reassessed the focal three species, comparing all possible

combinations of historical and modern individuals for each population. We find that 'Least Concern'

species largely show 'Acceptable' changes in genetic diversity, except for Southeastern and Western

Skåne *Cy. semiargus*, which fall into the 'Warning' category. Of the Red Listed species, *Cupido minimus*

('Near Threatened') falls into the 'Acceptable' category, *Plebejus argyrognomon* ('Vulnerable' in 2003

when the modern individual in this study was collected; 'Critically Endangered' and presumed extinct in

Sweden presently (*59*)) into 'Warning', and *Scolitantides orion* ('Endangered') into 'Alarm' (Figure 4; Table

S6). Results from our focal species illustrate that estimates from single individuals are challenging to

interpret, as estimates can vary considerably depending on the individual pair (Figure 4).

Figure 4. Changes in heterozygosity projected over 100 years based on the rate of change
estimated from modern and historical samples for each of 8 species. Species are grouped by
conservation status, which follows the species name on the x-axis. Background coloration refers to the
three genetic decline thresholds outlined by Andersson et al (21), where a retention of >=95% over 100
year is considered 'acceptable' (green), a retention of 75-94% as 'warning' (yellow), and a retention of
<75% as 'alarm' (red). Points represent median estimates for change in heterozygosity across all
historical/modern individual pairs for a species. Species with bars have more than one possible sample
pair, with error bars illustrating the spread of estimates across all sample pairs. *Sc. orion and Pl.
argyrognomon include pinned museum specimens for both the modern and historical individuals. The
modern individuals for Pl. argyrognomon and Sc. orion were sampled in 2003 and 2016, respectively, and
the conservation statuses shown here reflect the statuses at those times. Plebejus argyrognomon is
additionally the only species aligned to a reference genome for a different species (Table S12).

Discussion

Our analyses of 59 historical and 90 modern genomes uncover a decline in genetic diversity in the

specialist grassland butterfly *Cyaniris semiargus* over the past century. The previously connected

populations of this species in southern Sweden have become isolated into distinct genetic clusters

following agricultural intensification. Increased rates of inbreeding have begun to unmask weakly

deleterious mutations in *Cy. semiargus*, potentially reducing individual and mean fitness. This contrasts to
the generalist *Polyommatus icarus*, where we detect minimal changes in genetic diversity, inbreeding,
and genetic differentiation over the focal period. While the decline in genetic diversity and increased
homozygosity of deleterious mutations in *Cy. semiargus* is relatively mild, such genomic erosion lags
demographic change (*25, 28, 30*). We infer that effective population sizes have decreased dramatically in
some *Cy. semiargus* populations and that genetic diversity will continue to decline even if these
populations stabilize at their current sizes. All three species share a status of 'Least Concern' in Sweden,
highlighting that assessing genetic erosion can identify genetic extinction debts in species and
populations not captured by census-based conservation assessments.

The low genetic differentiation and absence of significant increases in population subdivision in *Po. icarus*
and *Pl. argus* suggest gene flow maintains cohesive metapopulations in these species. Consistently,
levels of heterozygosity remain stable, and inbreeding remains low in all populations. In contrast, we
uncover a loss of population connectivity in *Cy. semiargus*, evidenced by increased population structure
in the clustering analysis and higher genetic differentiation among modern populations. The historically
lower levels of genetic differentiation across southern Sweden in *Cy. semiargus* suggest that currently
isolated populations in Skåne are subsets of a historically well-connected metapopulation. The similarly
high estimates of historical effective population size in West- and Eastern Skåne provide an independent
line of evidence supporting this interpretation. The concurrent abrupt declines in effective population sizes
may reflect the cessation of gene flow, shifting effective size estimates from those of the metapopulation
to estimates for each subpopulation. This pattern is consistent with theoretical expectations of high
genetic diversity being maintained by even low levels of gene flow (*60, 61*), as documented in another
Lycaenid butterfly (*62*). Hence, concerns about genetic rescue (*63*) wiping out locally adapted ancestry
(*64, 65*) are not likely to be relevant for *Cy. semiargus* and other grassland associated insects in regions
with historically abundant habitat. In general, estimates of historical population structure may prove
valuable for distinguishing whether contemporary population structure reflects species ecology or is a
consequence of anthropogenic habitat isolation.

In the specialist *Cy. semiargus* we document an on average 6% reduction in heterozygosity. This change
is connected to increases in the rate of inbreeding, with differences between modern and historical
individual heterozygosity largely explained by changes in the abundance of long runs of homozygosity.
Thus, F_{RoH} can provide a powerful metric for detecting genomic erosion even from only contemporary
samples. Additionally, trajectories of recent effective population sizes, inferred from LD, have the potential
to uncover recent demographic declines before changes in genetic diversity begin. For instance, temporal
changes in effective population size suggest that *Pl. argus* has experienced very recent declines, though
not to sizes as low as *Cy. semiargus*, but with no changes in heterozygosity or F_{RoH}. Hence, there is a
possibility that there have been recent reductions in gene flow also in *Pl. argus*, despite stable estimates
of differentiation, as population differentiation lags reductions in gene flow (*26, 66*). Ultimately, while the

observed reductions in genetic diversity in these species are relatively mild, genetic diversity loss lags
demographic change and will continue even if population sizes stabilize.

Understanding to what extent genetic diversity decline, resulting from habitat loss and isolation, reduces
the fitness of individuals and prospects for maintaining long-term viable populations remains a challenge.
As we have no direct fitness estimates available, we quantified counts of putatively deleterious mutations
as a proxy for genetic load (*67*). We found little evidence for either purging or accumulation of deleterious
mutations in any of the three species, as total derived counts did not differ significantly for most mutation
categories. We found a small decrease (3.7%) in the total derived count of missense mutations in *Pl.*
argus, which is surprising, as theory predicts weakly deleterious mutations to be purged less efficiently
than more strongly deleterious ones (*15*, *49*). As there is no change in the derived count at loss of
function or conserved positions, which are expected to be more strongly deleterious, this decrease is
potentially an artifact.

In *Cy. semiargus*, increased rates of inbreeding have increased the homozygous derived count at
missense and conserved positions. Assuming at least partial recessivity for these mutations, this would
increase the realized load in more inbred populations, even if the total count of derived alleles has not
changed. However, a small increase in homozygosity at missense positions may not contribute to
substantial fitness effects, especially if these alleles tend to be additive. Surprisingly, the increased
homozygous derived count at conserved positions, expected to have relatively strong deleterious effects
(*12*, *68*), more closely mirrors patterns at missense than loss of function positions. This could reflect the
limited ability of sequence conservation scores to distinguish between weakly and strongly deleterious
mutations (*69*, *70*). In addition, the metric we utilize to assess sequence conservation, Genomic
Evolutionary Rate Profiling (GERP) scores (*68*), depends on estimates of the rate of neutral evolution,
approximating the rejected substitution rate. Our GERP score estimates are based on a time tree (similar
to *54*, *71*, *72*) and hence differ from scores estimated based on neutral rates (e.g. *73*). Nevertheless, our
identification of conserved sites selects for the most conserved variants in each dataset and should be
robust for within-species comparisons.

While it is unclear to what extent the genomic erosion we estimate in *Cy. semiargus* populations affects
fitness in the current generation, the patterns we observe are in line with a lag between population decline
and subsequent genomic erosion. While we measure only an average 6% reduction in individual
heterozygosity across the study period, we infer greater than 10-fold declines in effective population sizes
in recent decades. Using coalescent simulations, we demonstrate that this disparity is to be expected for
such recent declines, implying that genetic erosion will continue to 'catch up' to the scale of demographic
decline in future generations. Recent empirical and simulation studies (*27–29, 74*) have highlighted a time
lag between disturbance events and genomic erosion referred to as drift debt (*16, 75*) or genetic
extinction debt (*25*). Importantly, such a lag is not only expected in the loss of genome-wide genetic
diversity, but also in increases in the rate of inbreeding and accumulation of genetic load (*28*). It can take

many generations for drift and inbreeding to bring low frequency deleterious mutations to high enough
frequency for purging and accumulation of deleterious variants to be noticeable (*28*), so it is not surprising
we do not detect these patterns. Given time, populations will lose a portion of their deleterious burden
through drift and purging, but a portion also will drift to higher frequency and fixation in subsequent
generations, reducing mean fitness of the population in the long term (*12–14*, *16*, *76*). Simultaneously,
erosion of genome-wide genetic diversity will continuously reduce adaptive potential (*77*), further reducing
the prospects of long-term persistence. Stabilization of populations at current effective population sizes is
insufficient to reduce the genetic extinction debt they now carry. Increasing the levels of gene flow to
historic levels to increase effective population sizes is needed to reduce this debt.

We assessed if the reduction in heterozygosity and increases in inbreeding found in *Cy. semiargus* are
also found in other species of Polyommatinae blues in this region. We find that genetic diversity is
maintained in five of the seven additional species assessed. Under the threat categories used in Sweden
for genetic monitoring, we uncover 'Warning' and 'Alarm' level classifications for two additional species,
both endangered grassland specialists, *Pl. argyrognomon* and *Sc. orion*. The estimated level of
heterozygosity decline in *Pl. argyrognomon* across a period from 1967-2003 is comparable to the decline
in *Cy. semiargus*. This is concerning, as when the modern *Pl. argyrognomon* was sampled, the species
already had a 'Vulnerable' conservation status and is presumed extinct in Sweden since 2018 (*59*).

None of the three focal species are currently demographically declining at a rate to qualify for the
Swedish Red List based on census estimates. Systematic monitoring in Sweden from 2010-2023 reveal
declines in both *Po. icarus* and *Cy. semiargus*, while *Pl. argus* appears to have stabilized (*78*). In *Cy.*
semiargus, we find that demographic decline has been sufficient to begin reducing genetic diversity, but
that most genomic erosion has yet to be realized. Genetic monitoring, especially monitoring incorporating
historical samples, can enable early identification of signals of populations experiencing genetic erosion
to inform conservation actions to mitigate loss of genetic diversity (*24*, *25*). We have previously shown
that individual heterozygosity in these *Cy. semiargus* populations is positively related to grassland area
on the landscape level (5-20km from the local patch, *44*). Combined with the findings from the current
study, these results show that preservation of current grassland area and configuration will be insufficient
for maintaining genetic diversity in *Cy. semiargus* populations in southern Sweden. Rather, restoration of
suitable habitat at a scale that facilitates gene flow between populations is vital to prevent further genetic
erosion in this species, and potentially other insect taxa that dependent on semi-natural grassland habitat.

In conclusion, we document the early stages of genomic erosion in a specialist grassland butterfly in
southern Sweden. While census size-based estimates of demographic declines largely inform current
conservation status of these species, the isolation and inbreeding of *Cyaniris semiargus* highlights that
even formerly common grassland insects may now carry a substantial genetic extinction debt, forming the
basis for an extinction debt at the community level (*25*). Our findings suggest that prompt restoration of
grassland connectivity, especially in agricultural regions, is needed to restore gene flow and maintain

genetic diversity in specialist grassland butterflies, reducing their risk of extirpation and extinction. This
work emphasizes the urgency of semi-natural grassland restoration efforts to meet the goal of maintaining
genetic diversity under Goal A of the Kunming-Montreal Global Biodiversity Framework (*20*). Additionally,
the inbreeding we observe in a formerly abundant grassland butterfly points to the importance of
considering population connectivity during the restoration process if restoration is to successfully halt the
erosion of genetic diversity. Broadly, our findings underscore that genetic monitoring is a valuable tool in
conservation assessment of insects, as early detection of genomic erosion can highlight populations that
may have stabilized but still carry a genetic extinction debt, enabling timely action.

Materials and Methods

Study System and Sampling

To assess changes in genetic diversity, differentiation, inbreeding, and deleterious mutation burden, we
compared modern and historical genomes from three focal blue butterfly species (Polyommatinae),
Polyommatus icarus, *Plebejus argus*, and *Cyaniris semiargus* (Figure 1A). These species vary in habitat
specialization, being grassland generalist, heathland specialist, and grassland specialist, respectively. We
used previously generated population-level genetic data (*44*) for the modern genomes. For the historical
genomes, we selected pinned specimens from the entomological collections of the Biological Museum,
Lund University (*43*) sampled between 1917-1956 in roughly the same geographic area as the modern
samples. For each historical locality, we selected 4-11 specimens sampled in the same year to estimate
population-level metrics. The historical and modern population-level pairs were sampled in four broad
geographic areas: I) Western and II) Eastern Skåne, III) Småland, and IV) Öland (Figure 1B), with
historical localities ranging approximately 12-40km from their modern counterparts. This resulted in three
historical-modern population pairs for *Po. icarus* and two for *Pl. argus*. For *Cy. semiargus*, we included
three pairs and an additional comparison of a modern Southeastern Skåne population paired with the
historical Eastern Skåne population. For *Pl. argus*, only one historical specimen was available from
Western and Eastern Skåne, respectively, and these were compared to the closest modern populations in
individual-level analyses only.

To more broadly assess the variation in rates of genetic decline in Polyommatinae of differing
conservation statuses, we sampled historical and modern individuals of five additional species: *Agriades*
optilete, Aricia agestis, Cupido minimus, Plebejus argyrognomon and *Scolitantides orion.* Modern
specimens of *Ag. optilete, Ar. agestis,* and *Cu. minimus* were collected across the same geographic
regions as the focal species in 2021 (Table S1). A pinned specimen used as the modern sample of *Sc.*
orion was collected in 2016 in Södermanland and donated to this study by Håkan Elmqvist. Legs from a
pinned speciemen of *Pl. argyrognomon*, collected in 2003 in Western Skåne, were donated by Göran
Engqvist and used as the modern sample for this species. Historical samples for these five species were
selected to be proximal to the modern specimen sampling locality with collection dates ranging from

1939-1967, all provided as pinned specimens from the entomological collections of the Biological
Museum, Lund University (*43*).

DNA Extraction and Sequencing

DNA from fresh specimens was extracted and sequenced following Nolen et al (*44*). We extracted DNA
and prepared sequencing libraries from pinned museum specimens in a facility dedicated to historical
DNA processing at Lund University, following the protocol described by Twort et al (*79*). Libraries
prepared for pinned specimens were pooled into two equimolar pools containing 37-38 libraries each,
sequencing each pool on a single NovaSeq 6000 S4 lane. DNA was extracted from the abdomens or legs
of male specimens without homogenization to preserve anatomical structures.

Sequence Processing

Sequence processing and analyses were largely performed using Snakemake v8.25.5 (*80*), incorporating
and extending the PopGLen v0.4.1 (*81*) workflow. We trimmed raw sequencing reads using fastp v0.23.4
(*82*) with default settings, additionally collapsing overlapping read pairs in historical samples, discarding
those with less than 30bp overlap. As historical DNA is expected to be fragmented, mapping can be
biased towards reads containing the reference allele. We investigated two potential alignment options to
reduce this impact: (I) BWA aln (*83*) with relaxed parameters, as is used in most historical DNA studies,
and (II) alignment to a variation graph incorporating known alternative variants from the modern samples,
which has been shown to reduce reference biases when mapping short fragments (*84*). As we found that
mapping to the variation graph reduced the amount of reference bias in the historical samples (Figure
S2&3), we performed all analyses with both modern and historical samples aligned to the species-specific
variation graphs. Detailed methods of the processing for this comparison and the final alignment method
are available in the Supplementary Methods, with a brief outline of the final method here.

To construct variation graphs for each species, we first aligned the trimmed, paired-end modern reads to
the respective reference genome (Table S12) (*85–87*) using BWA mem v0.7.17 (*88*), removed duplicates
using Picard v2.27 (*89*), and realigned around indels using GATK (*90*). We called genotypes in these
samples using the BCFtools multi-allelic caller v1.21 (*91*) and constructed a variation graph from these
variants and the linear reference genome using vg v1.63.1 (*92*). To produce the final alignments, we
mapped modern and historical samples to the variant graph using vg with default settings, additionally
setting -k 15 for historical samples as recommended by Martiniano et al. (*84*). We removed duplicates
with Picard and dedup v0.12.8 (*93*) for modern and historical samples, respectively, and clipped
overlapping read pairs using BamUtil v1.0.15 (*94*). We estimated post-mortem DNA damage in historical
samples using DamageProfiler v1.1 (*95*).

Genotype calling and filtering

To account for variation in sequencing depth, we subsampled all individuals to a mean depth of 6x using
Samtools v1.20 (*91*). We excluded individuals with a mean coverage of less than 5x from genotype

calling. We implemented a sites-based filtering scheme (see Supplementary Methods) and called
genotypes for each species jointly across samples using the BCFtools multi-allelic caller at sites passing
the species filter sets (Table S13), including reads with minimum mapping and base qualities of 30 and
20, respectively. We filtered to biallelic and invariant positions, excluding those within 5bp of indels or a
variant quality less than 30. We set genotypes with a depth less than 6 and heterozygous genotypes with
one allele supported by less than 20% of the reads to missing. We removed transition positions across all
samples and filtered out positions with more than 40% missing data in either the historical or modern
sample subsets.

Heterozygosity, nucleotide diversity, and genetic differentiation

We estimated individual genome-wide heterozygosity for all samples from the called genotypes by
dividing the count of heterozygous calls by the total count of called genotypes, including invariant sites.
We estimated nucleotide diversity using Pixy v2.0.0.beta2 (*96*), which utilizes a sample-size weighted
calculation. Nucleotide diversity estimates in these populations are robust to low (N = 4) sample sizes
(*44*), so we included estimates for all populations with multiple individuals. We estimated this summary
statistic in 50kb non-overlapping windows for populations grouped by sampling locality, estimating the
genome-wide mean across windows, and confidence intervals from 10,000 bootstraps, resampling
windows with replacement. We estimated genetic differentiation using pairwise population F_{ST} with Pixy
on the same windows with the Hudson estimator (*97*), which is robust to small and variable sample sizes
(*98*). We estimated mean F_{ST} and confidence intervals with the same approach as nucleotide diversity.

Relatedness

As analyses of population structure can be sensitive to the presence of close relatives in the dataset, we
estimated pairwise relatedness between all samples within each species dataset. We utilized the
IBSrelate SFS-based method (*99*), as implemented in NGSrelate (*100*), to estimate R0, R1, and KING-
robust kinship coefficients (*101*) for all pairs of individuals in the dataset. For input, we generated a
Beagle file for each focal species dataset, using all sampled individuals, but subsampling individuals with
higher coverage to 8X to reduce the influence of the large variation in coverage in our modern samples.
We estimated genotype likelihoods in ANGSD v0.940 (*102*) using the GATK model from reads with a
mapping quality >= 30, base quality >= 20, at positions with a minimum depth of three that passed
species-specific site filters. We called SNPs with a p-value < 1e-6, a minor allele frequency >= 0.05, and
with data in at least 90% of the individuals.

Population Structure

We explored population structure using principal component analysis (PCA) as implemented in PCAngsd
v1.10 (*103*). To obtain independent sites for population structure analyses, we estimated linkage
disequilibrium in the Beagle file from the relatedness analyses using ngsLD v1.2.0 (*104*) and pruned
SNPs using prune_graph (https://github.com/fgvieira/prune_graph; v0.3.2), assuming independence of

SNPs >50kb apart and with a coefficient of correlation (r^2) < 0.1. We removed one individual from each

pair of third-degree or closer relatives in the dataset (expected KING-kinship coefficient >= 0.0625; Table

S14) before running the PCA.

We formally tested for population structure using admixture analyses for 1-8 ancestral clusters (K) using

the pruned, relative removed Beagle file in NGSadmix v33 (*105*). We performed at least 20 replicate runs

for each value of K, up to 100 total runs, stopping if the three highest likelihood replicates converged

within 2 log-likelihood units of each other. The 'best-fit' value for K was then determined to be the highest

K for which replicates met this convergence criteria, following Pečnerová et al (*106*). We further assessed

the model fit for each value of K using evalAdmix v0.961 (*107*).

Runs of Homozygosity

We assessed rates of recent and background inbreeding by estimating the inbreeding coefficient F_{RoH}, the

proportion of the autosomal genome existing in runs of homozygosity over a specified length,

incorporating runs of homozygosity >100kb in our estimates of F_{RoH}. Per individual runs of homozygosity

were estimated using BCFtools/RoH v1.19 (*108*). We set a constant species-specific recombination rate,

assuming 50 centimorgans per autosomal chromosome as in Mackintosh et al (*33*) (Table S12), and ran

the analysis on genotype calls at variable sites, setting the PL of unseen genotypes set to 30 as

suggested by the documentation. To ensure F_{RoH} estimates were inferred equivalently in species with

population and individual level sampling, we did not estimate allele frequencies, instead using a default

minor allele frequency of 0.4 as suggested in the documentation and ignored homozygous reference

positions, as including these can inflate estimates of F_{RoH} based on our testing (Figure S11). We

calculated F_{RoH} for each individual by dividing the cumulative length of runs of homozygosity by the total

length of the autosomes.

Recent demography and coalescent simulations of heterozygosity loss

To infer timing and magnitude of demographic declines that may contribute to reduced genetic diversity,

we inferred changes in effective population size over the past 100 years from the modern samples using

the linkage disequilibrium-based approach implemented in GONE (*109*). We inferred these histories in

modern samples by sampling locality, as this method performs best when input samples do not have

additional genetic substructure (*110*). We performed this analysis on the called modern variants used to

construct the variation graphs, which included transitions. As we observed a frequent sudden drop in the

most recent four generations in most of the GONE analyses, we visualized trajectories both including and

excluding these generations (Figure S5).

To assess if our empirical estimates of heterozygosity decline and accumulation of runs of homozygosity

fell within expected ranges for the characterized demographic changes, we performed coalescent

simulations using Slendr (*111*) with the msprime (*112*) backend. For each modern locality with a paired

historical locality, we constructed demographic models that followed size changes inferred for the most

recent 100 generations in the GONE analyses. As most sampled populations showed a roughly constant

historical size by this time point (Figure S5), we set the ancestral size to that of generation 100. For

localities where we inferred a constant history of very large size (>100,000), we modeled a simple

constant size model with N = 100,000. Due to the sharp drop in the four most recent generations, we set

these generations in the model to the estimate of generation five. For each sampled locality, we ran 100

replicate simulations of a single, 10 Mb chromosome with a recombination rate corresponding to 50cM

(*33*) and a mutation rate of 2.9e-8 (*113*). At time points corresponding to the collection of the historical

samples for a given locality and the most recent generation, we sampled 100 individuals from each

simulation and calculated mean heterozygosity, F_{RoH}, and heterozygosity outside of RoH to infer a range

of expected values to compare our empirical estimates to.

Deleterious mutation burden

As proxies for genetic load, we estimated the burden of deleterious mutations per individual using the

count of derived alleles with putatively deleterious consequences. We inferred these consequences using

two complementary methods, one based on gene annotations to infer functional effects (*114*) and the

other identifying positions with high sequence conservation (*115*). As variants were called across the

combined historical and modern sample set for each species, alleles that are fixed within one time

category, but variable or fixed for the opposing allele in the other, are included.

To identify which alleles were likely ancestral, we co-inferred ancestral states and sequence conservation

scores using the GenErode pipeline v0.6.2 (*116*), which estimates Genomic Evolutionary Rate Profiling

(GERP) scores (*68*) from a set of outgroup species assemblies. We utilized 24 Lycaenidae chromosome-

level genomes from distinct genera (Table S15) (*85–87*, *117–120*) as outgroups, modifying the default

fragment size for GenErode's alignment step from 35 to 100bp, and using divergence times estimated

with TimeTree 5 (*121*). We then subset the variable sites to include only those with GERP > 0 and where

the ancestral allele matched the reference to avoid underestimating derived counts in historical samples.

We identified deleterious mutations based on functional effects using the genome annotations provided

by the Darwin Tree of Life Project (https://projects.ensembl.org/darwin-tree-of-life/). We assigned impacts

to each variant using Ensembl's Variant Effect Predictor (VEP) v112.0 (*114*). VEP categorizes variants

into four classes: 'High', 'Moderate', 'Low', or 'Modifier', which largely correspond to loss of function,

missense, synonymous, and intergenic variants. We forced VEP to pick one effect per variant, using the

default selection order (*114*). To identify conserved positions, we selected the top 1% of GERP scores as

the most conserved positions using the get_gerp_score_percentile.py script included with GenErode. We

inferred deleterious burden at positions inferred as loss of function ('High' impact), missense ('Moderate'

impact), or conserved based on GERP scores, using the derived allele count at these positions. Total

deleterious burden is the total derived allele count of a variant category per 1000 called genotypes.

Homozygous deleterious burden is the homozygous derived allele count per 1000 called genotypes. We

additionally estimated these counts at synonymous ('low' impact) and intergenic positions to assess

biases in derived counts across sample types.

Categorizing rates of genetic decline

To assess the changes in genetic diversity in a framework developed for conservation focused genetic

monitoring, we used the indicator thresholds suggested by Andersson et al (*21*), which are based upon

indicators applied for national use in Sweden (*58*). Using these thresholds, genetic decline within

populations is measured as the amount of genetic diversity retained over a period of 100 years, assuming

a constant rate of change. Three threat categories are suggested: retention >=95% is considered

'Acceptable', 75-94% as 'Warning', and <75% as 'Alarm'. We estimated the predicted retention of genetic

diversity over 100 years for each population using individual heterozygosity and nucleotide diversity.

Genetic diversity retention per year was estimated using the following equation:

$$e^{\frac{ln(H_{mod}/H_{hist})}{Y_{mod}-Y_{hist}}}$$

where H is the estimate of genetic diversity for a given sample/population and Y is the sampling year for

that sample/population. We estimated this metric for nucleotide diversity in the focal species using

population mean estimates. For individual heterozygosity, we estimated this metric using population

means in the focal species and using all pairwise comparisons of historical and modern individuals in both

focal and supplemental species. For the effective population size indicator, we utilized the estimate of

effective population size at generation five in the GONE analyses.

References

1. D. L. Wagner, E. M. Grames, M. L. Forister, M. R. Berenbaum, D. Stopak, Insect decline in the
Anthropocene: Death by a thousand cuts. *Proceedings of the National Academy of Sciences* **118**,
e2023989118 (2021).
2. R. van Klink, D. E. Bowler, K. B. Gongalsky, A. B. Swengel, A. Gentile, J. M. Chase, Meta-analysis reveals
declines in terrestrial but increases in freshwater insect abundances. *Science* **368**, 417–420 (2020).
3. R. Dirzo, H. S. Young, M. Galetti, G. Ceballos, N. J. B. Isaac, B. Collen, Defaunation in the Anthropocene.
Science **345**, 401–406 (2014).
4. C. A. Halsch, A. M. Shapiro, J. A. Fordyce, C. C. Nice, J. H. Thorne, D. P. Waetjen, M. L. Forister, Insects
and recent climate change. *Proceedings of the National Academy of Sciences* **118**, e2002543117 (2021).
5. P. H. Raven, D. L. Wagner, Agricultural intensification and climate change are rapidly decreasing insect
biodiversity. *Proceedings of the National Academy of Sciences* **118**, e2002548117 (2021).
6. S. Seibold, M. M. Gossner, N. K. Simons, N. Blüthgen, J. Müller, D. Ambarlı, C. Ammer, J. Bauhus, M.
Fischer, J. C. Habel, K. E. Linsenmair, T. Nauss, C. Penone, D. Prati, P. Schall, E.-D. Schulze, J. Vogt, S.
Wöllauer, W. W. Weisser, Arthropod decline in grasslands and forests is associated with landscape-level
drivers. *Nature* **574**, 671–674 (2019).
7. D. H. Boyes, D. M. Evans, R. Fox, M. S. Parsons, M. J. O. Pocock, Is light pollution driving moth population
declines? A review of causal mechanisms across the life cycle. *Insect Conservation and Diversity* **14**, 167–
187 (2021).
8. J. R. Shipley, E. R. Frei, A. Bergamini, S. Boch, T. Schulz, C. Ginzler, M. Barandun, P. Bebi, J. Bolliger, K.
Bollmann, N. Delpouve, M. M. Gossner, C. Graham, F. Krumm, M. Marty, N. Pichon, A. Rigling, C. Rixen,
Agricultural practices and biodiversity: Conservation policies for semi-natural grasslands in Europe. *Current
Biology* **34**, R753–R761 (2024).
9. R. C. Lacy, Loss of Genetic Diversity from Managed Populations: Interacting Effects of Drift, Mutation,
Immigration, Selection, and Population Subdivision. *Conservation Biology* **1**, 143–158 (1987).
10. R. Lande, Mutation and Conservation. *Conservation Biology* **9**, 782–791 (1995).

11. R. Lande, S. Shannon, The Role of Genetic Variation in Adaptation and Population Persistence in a Changing
Environment. *Evolution* **50**, 434–437 (1996).
12. G. Bertorelle, F. Raffini, M. Bosse, C. Bortoluzzi, A. Iannucci, E. Trucchi, H. E. Morales, C. van Oosterhout,
Genetic load: genomic estimates and applications in non-model animals. *Nat Rev Genet* **23**, 492–503 (2022).
13. C. van Oosterhout, Mutation load is the spectre of species conservation. *Nat Ecol Evol* **4**, 1004–1006 (2020).
14. M. Lynch, J. Conery, R. Burger, Mutation Accumulation and the Extinction of Small Populations. *The*
American Naturalist **146**, 489–518 (1995).
15. J. Wang, W. G. Hill, D. Charlesworth, B. Charlesworth, Dynamics of inbreeding depression due to deleterious
mutations in small populations: mutation parameters and inbreeding rate. *Genetics Research* **74**, 165–178
(1999).
16. N. Dussex, H. E. Morales, C. Grossen, L. Dalén, C. van Oosterhout, Purging and accumulation of genetic
load in conservation. *Trends in Ecology & Evolution* **38**, 961–969 (2023).
17. R. Lande, RISK OF POPULATION EXTINCTION FROM FIXATION OF NEW DELETERIOUS MUTATIONS.
Evolution **48**, 1460–1469 (1994).
18. M. E. Gilpin, M. E. Soule, "Minimum viable populations: Processes of species extinction" in *Conservation*
Biology: The Science of Scarcity and Diversity (Sinauer Associates, Sunderland, Massachusetts, 1986), pp.
19–34.
19. J. Höglund, "The extinction vortex, is genetic variation related to extinction?" in *Evolutionary Conservation*
Genetics, J. Höglund, Ed. (Oxford University Press, 2009;
https://doi.org/10.1093/acprof:oso/9780199214211.003.0001), p. 0.
20. Convention on Biological Diversity, "Decision Adopted by the Conference of the Parties to the Convention on
Biological Diversity 15/4. Kunming-Montreal Global Biodiversity Framework" (Montreal, Canada, 2022;
https://www.cbd.int/doc/decisions/cop-15/cop-15-dec-04-en.pdf).
21. A. Andersson, S. Karlsson, N. Ryman, L. Laikre, Monitoring genetic diversity with new indicators applied to an
alpine freshwater top predator. *Molecular Ecology* **31**, 6422–6439 (2022).
22. D. O'Brien, L. Laikre, S. Hoban, M. W. Bruford, R. Ekblom, M. C. Fischer, J. Hall, C. Hvilsom, P. M.
Hollingsworth, F. Kershaw, C. S. Mittan, T. A. Mukassabi, R. Ogden, G. Segelbacher, R. E. Shaw, C. Vernesi,
A. J. MacDonald, Bringing together approaches to reporting on within species genetic diversity. *Journal of*
Applied Ecology **59**, 2227–2233 (2022).
23. K. Theissinger, C. Fernandes, G. Formenti, I. Bista, P. R. Berg, C. Bleidorn, A. Bombarely, A. Crottini, G. R.
Gallo, J. A. Godoy, S. Jentoft, J. Malukiewicz, A. Mouton, R. A. Oomen, S. Paez, P. J. Palsbøll, C.
Pampoulie, M. J. Ruiz-López, S. Secomandi, H. Svardal, C. Theofanopoulou, J. de Vries, A.-M. Waldvogel,
G. Zhang, E. D. Jarvis, M. Bálint, C. Ciofi, R. M. Waterhouse, C. J. Mazzoni, J. Höglund, S. A. Aghayan, T. S.
Alioto, I. Almudi, N. Alvarez, P. C. Alves, I. R. A. do Rosario, A. Antunes, P. Arribas, P. Baldrian, G.
Bertorelle, A. Böhne, A. Bonisoli-Alquati, L. L. Boštjančić, B. Boussau, C. M. Breton, E. Buzan, P. F. Campos,
C. Carreras, L. Fi. C. Castro, L. J. Chueca, F. Čiampor, E. Conti, R. Cook-Deegan, D. Croll, M. V. Cunha, F.
Delsuc, A. B. Dennis, D. Dimitrov, R. Faria, A. Favre, O. D. Fedrigo, R. Fernández, G. F. Ficetola, J.-F. Flot,
T. Gabaldón, D. R. Agius, A. M. Giani, M. T. P. Gilbert, T. Grebenc, K. Guschanski, R. Guyot, B. Hausdorf, O.
Hawlitschek, P. D. Heintzman, B. Heinze, M. Hiller, M. Husemann, A. Iannucci, I. Irisarri, K. S. Jakobsen, P.
Klinga, A. Kloch, C. F. Kratochwil, H. Kusche, K. K. S. Layton, J. A. Leonard, E. Lerat, G. Liti, T. Manousaki,
T. Marques-Bonet, P. Matos-Maraví, M. Matschiner, F. Maumus, A. M. M. Cartney, S. Meiri, J. Melo-Ferreira,
X. Mengual, M. T. Monaghan, M. Montagna, R. W. Mysłajek, M. T. Neiber, V. Nicolas, M. Novo, P. Ozretić, F.
Palero, L. Pârvulescu, M. Pascual, O. S. Paulo, M. Pavlek, C. Pegueroles, L. Pellissier, G. Pesole, C. R.
Primmer, A. Riesgo, L. Rüber, D. Rubolini, D. Salvi, O. Seehausen, M. Seidel, B. Studer, S. Theodoridis, M.
Thines, L. Urban, A. Vasemägi, A. Vella, N. Vella, S. C. Vernes, C. Vernesi, D. R. Vieites, C. W. Wheat, G.
Wörheide, Y. Wurm, G. Zammit, How genomics can help biodiversity conservation. *Trends in Genetics* **39**,
545–559 (2023).
24. D. Díez-del-Molino, F. Sánchez-Barreiro, I. Barnes, M. T. P. Gilbert, L. Dalén, Quantifying Temporal Genomic
Erosion in Endangered Species. *Trends in Ecology & Evolution* **33**, 176–185 (2018).
25. R. Gargiulo, K. B. Budde, M. Heuertz, Mind the lag: understanding genetic extinction debt for conservation.
Trends in Ecology & Evolution **40**, 228–237 (2025).
26. C. W. Epps, N. Keyghobadi, Landscape genetics in a changing world: disentangling historical and
contemporary influences and inferring change. *Molecular Ecology* **24**, 6021–6040 (2015).
27. X. Liu, E. Milesi, C. Fontsere, H. L. Owens, R. Heinsohn, M. T. P. Gilbert, R. Crates, D. Nogués-Bravo, H. E.
Morales, Time-lagged genomic erosion and future environmental risks in a bird on the brink of extinction.
Proceedings of the Royal Society B: Biological Sciences **292**, 20242480 (2025).
28. A. V. Pinto, B. Hansson, I. Patramanis, H. E. Morales, C. van Oosterhout, The impact of habitat loss and
population fragmentation on genomic erosion. *Conserv Genet* **25**, 49–57 (2024).
29. C. Fontsere, S. A. Speak, A. J. Caven, J. A. Rodríguez, X. Wang, C. Pacheco, M. Cassatt-Johnstone, G.
Femerling, B. Maloney, J. Balacco, J. Collins, Y. Sims, L. Abueg, O. Fedrigo, E. D. Jarvis, B. K. Hartup, B.
Shapiro, M. T. P. Gilbert, C. van Oosterhout, H. E. Morales, Persistent Genomic Erosion in Whooping Cranes
Despite Demographic Recovery. *Molecular Ecology*, e70088 (2025).

30.   K. S. Mualim, J. P. Spence, C. Weiß, O. Selmoni, M. Lin, M. Exposito-Alonso, Genetic diversity loss in the
666   Anthropocene will continue long after habitat destruction ends. bioRxiv [Preprint] (2024).
667   https://doi.org/10.1101/2024.10.21.619096.
668   31.   H. Ellegren, N. Galtier, Determinants of genetic diversity. *Nat Rev Genet* **17**, 422–433 (2016).
669   32.   J. Romiguier, P. Gayral, M. Ballenghien, A. Bernard, V. Cahais, A. Chenuil, Y. Chiari, R. Dernat, L. Duret, N.
670   Faivre, E. Loire, J. M. Lourenco, B. Nabholz, C. Roux, G. Tsagkogeorga, A. a.-T. Weber, L. A. Weinert, K.
671   Belkhir, N. Bierne, S. Glémin, N. Galtier, Comparative population genomics in animals uncovers the
672   determinants of genetic diversity. *Nature* **515**, 261–263 (2014).
673   33.   A. Mackintosh, D. R. Laetsch, A. Hayward, B. Charlesworth, M. Waterfall, R. Vila, K. Lohse, The determinants
674   of genetic diversity in butterflies. *Nat Commun* **10**, 3466 (2019).
675   34.   G. Hewitt, The genetic legacy of the Quaternary ice ages. *Nature* **405**, 907–913 (2000).
676   35.   M. K. Schwartz, G. Luikart, R. S. Waples, Genetic monitoring as a promising tool for conservation and
677   management. *Trends in Ecology & Evolution* **22**, 25–33 (2007).
678   36.   M. T. Webster, A. Beaurepaire, P. Neumann, E. Stolle, Population Genomics for Insect Conservation. *Annual
679   Review of Animal Biosciences* **11**, 115–140 (2023).
680   37.   M. Ihse, Swedish agricultural landscapes — patterns and changes during the last 50 years, studied by aerial
681   photos. *Landscape and Urban Planning* **31**, 21–37 (1995).
682   38.   R. Lindborg, O. Eriksson, Historical Landscape Connectivity Affects Present Plant Species Diversity. *Ecology*
683   **85**, 1840–1845 (2004).
684   39.   SLU Artdatabanken, *Rödlistade arter i Sverige 2020* (SLU Artdatabanken, Uppsala, 2020).
685   40.   D. L. Wagner, R. Fox, D. M. Salcido, L. A. Dyer, A window to the world of global insect declines: Moth
686   biodiversity trends are complex and heterogeneous. *Proceedings of the National Academy of Sciences* **118**,
687   e2002549117 (2021).
688   41.   M. S. Warren, D. Maes, C. A. M. van Swaay, P. Goffart, H. Van Dyck, N. A. D. Bourn, I. Wynhoff, D. Hoare,
689   S. Ellis, The decline of butterflies in Europe: Problems, significance, and possible solutions. *Proceedings of
690   the National Academy of Sciences* **118**, e2002551117 (2021).
691   42.   J. C. Habel, A. Segerer, W. Ulrich, O. Torchyk, W. W. Weisser, T. Schmitt, Butterfly community shifts over
692   two centuries. *Conservation Biology* **30**, 754–762 (2016).
693   43.   C. Fägerström, Lund University Biological Museum - Insect collections Inventory. doi: 10.15468/dahk2a
694   (2024).
695   44.   Z. J. Nolen, M. Rundlöf, A. Runemark, Species-specific erosion of genetic diversity in grassland butterflies
696   depends on landscape land cover. *Biological Conservation* **296**, 110694 (2024).
697   45.   M. T. P. Gilbert, J. Binladen, W. Miller, C. Wiuf, E. Willerslev, H. Poinar, J. E. Carlson, J. H. Leebens-Mack, S.
698   C. Schuster, Recharacterization of ancient DNA miscoding lesions: insights in the era of sequencing-by-
699   synthesis. *Nucleic Acids Research* **35**, 1–10 (2007).
700   46.   M. Stiller, R. E. Green, M. Ronan, J. F. Simons, L. Du, W. He, M. Egholm, J. M. Rothberg, S. G. Keates, N. D.
701   Ovodov, E. E. Antipina, G. F. Baryshnikov, Y. V. Kuzmin, A. A. Vasilevski, G. E. Wuenschell, J. Termini, M.
702   Hofreiter, V. Jaenicke-Després, S. Pääbo, Patterns of nucleotide misincorporations during enzymatic
703   amplification and direct large-scale sequencing of ancient DNA. *Proceedings of the National Academy of
704   Sciences* **103**, 13578–13584 (2006).
705   47.   R. McQuillan, A.-L. Leutenegger, R. Abdel-Rahman, C. S. Franklin, M. Pericic, L. Barac-Lauc, N. Smolej-
706   Narancic, B. Janicijevic, O. Polasek, A. Tenesa, A. K. MacLeod, S. M. Farrington, P. Rudan, C. Hayward, V.
707   Vitart, I. Rudan, S. H. Wild, M. G. Dunlop, A. F. Wright, H. Campbell, J. F. Wilson, Runs of Homozygosity in
708   European Populations. *The American Journal of Human Genetics* **83**, 359–372 (2008).
709   48.   E. A. Thompson, Identity by Descent: Variation in Meiosis, Across Genomes, and in Populations. *Genetics*
710   **194**, 301–326 (2013).
711   49.   S. Glémin, HOW ARE DELETERIOUS MUTATIONS PURGED? DRIFT VERSUS NONRANDOM MATING.
712   *Evolution* **57**, 2678–2687 (2003).
713   50.   M. Kirkpatrick, P. Jarne, The Effects of a Bottleneck on Inbreeding Depression and the Genetic Load. *The
714   American Naturalist* **155**, 154–167 (2000).
715   51.   M. Lynch, W. Gabriel, Mutation Load and the Survival of Small Populations. *Evolution* **44**, 1725–1737 (1990).
716   52.   B. Charlesworth, Mutational load, inbreeding depression and heterosis in subdivided populations. *Molecular
717   Ecology* **27**, 4991–5003 (2018).
718   53.   M. Dehasque, H. E. Morales, D. Díez-del-Molino, P. Pečnerová, J. C. Chacón-Duque, F. Kanellidou, H.
719   Muller, V. Plotnikov, A. Protopopov, A. Tikhonov, P. Nikolskiy, G. K. Danilov, M. Giannì, L. van der Sluis, T.
720   Higham, P. D. Heintzman, N. Oskolkov, M. T. P. Gilbert, A. Götherström, T. van der Valk, S. Vartanyan, L.
721   Dalén, Temporal dynamics of woolly mammoth genome erosion prior to extinction. *Cell* **187**, 3531-3540.e13
722   (2024).
723   54.   T. van der Valk, D. Díez-del-Molino, T. Marques-Bonet, K. Guschanski, L. Dalén, Historical Genomes Reveal
724   the Genomic Consequences of Recent Population Decline in Eastern Gorillas. *Current Biology* **29**, 165-
725   170.e6 (2019).

55. G. Femerling, C. van Oosterhout, S. Feng, R. M. Bristol, G. Zhang, J. Groombridge, M. T. P. Gilbert, H. E. Morales, Genetic Load and Adaptive Potential of a Recovered Avian Species that Narrowly Avoided Extinction. *Molecular Biology and Evolution* **40**, msad256 (2023).

56. D. Kleinman-Ruiz, M. Lucena-Perez, B. Villanueva, J. Fernández, A. P. Saveljev, M. Ratkiewicz, K. Schmidt, N. Galtier, A. García-Dorado, J. A. Godoy, Purging of deleterious burden in the endangered Iberian lynx. *Proceedings of the National Academy of Sciences* **119**, e2110614119 (2022).

57. E. Humble, M. A. Stoffel, K. Dicks, A. D. Ball, R. M. Gooley, J. Chuven, R. Pusey, M. A. Remeithi, K.-P. Koepfli, B. Pukazhenthi, H. Senn, R. Ogden, Conservation management strategy impacts inbreeding and mutation load in scimitar-horned oryx. *Proceedings of the National Academy of Sciences* **120**, e2210756120 (2023).

58. K. Johannesson, L. Laikre, Monitoring of genetic diversity in environmental monitoring. *Report to the Swedish Agency for Marine and Water Management (dnr. HaV 3642-2018, 3643-2018)* (2020).

59. SLU Artdatabanken, Artfakta: kronärtsblåvinge (Plebejus argyrognomon. https://artfakta.se/taxa/101598.

60. R. Lande, Neutral Theory of Quantitative Genetic Variance in an Island Model with Local Extinction and Colonization. *Evolution* **46**, 381–389 (1992).

61. M. Nei, N. Takahata, Effective population size, genetic diversity, and coalescence time in subdivided populations. *J Mol Evol* **37**, 240–244 (1993).

62. Z. Gompert, A. Springer, M. Brady, S. Chaturvedi, L. K. Lucas, Genomic time-series data show that gene flow maintains high genetic diversity despite substantial genetic drift in a butterfly species. *Molecular Ecology* **30**, 4991–5008 (2021).

63. P. W. Hedrick, J. R. Adams, J. A. Vucetich, Reevaluating and Broadening the Definition of Genetic Rescue. *Conservation Biology* **25**, 1069–1070 (2011).

64. P. W. Hedrick, A. Garcia-Dorado, Understanding Inbreeding Depression, Purging, and Genetic Rescue. *Trends in Ecology & Evolution* **31**, 940–952 (2016).

65. D. A. Bell, Z. L. Robinson, W. C. Funk, S. W. Fitzpatrick, F. W. Allendorf, D. A. Tallmon, A. R. Whiteley, The Exciting Potential and Remaining Uncertainties of Genetic Rescue. *Trends in Ecology & Evolution* **34**, 1070–1079 (2019).

66. E. L. Landguth, S. A. Cushman, M. K. Schwartz, K. S. McKELVEY, M. Murphy, G. Luikart, Quantifying the lag time to detect barriers in landscape genetics. *Molecular Ecology* **19**, 4179–4191 (2010).

67. Y. B. Simons, G. Sella, The impact of recent population history on the deleterious mutation load in humans and close evolutionary relatives. *Current Opinion in Genetics & Development* **41**, 150–158 (2016).

68. G. M. Cooper, E. A. Stone, G. Asimenos, E. D. Green, S. Batzoglou, A. Sidow, Distribution and intensity of constraint in mammalian genomic sequence. *Genome Res.* **15**, 901–913 (2005).

69. C. D. Huber, B. Y. Kim, K. E. Lohmueller, Population genetic models of GERP scores suggest pervasive turnover of constrained sites across mammalian evolution. *PLOS Genetics* **16**, e1008827 (2020).

70. C. C. Kyriazis, J. A. Robinson, K. E. Lohmueller, Using Computational Simulations to Model Deleterious Variation and Genetic Load in Natural Populations. *The American Naturalist* **202**, 737–752 (2023).

71. J. von Seth, N. Dussex, D. Díez-del-Molino, T. van der Valk, V. E. Kutschera, M. Kierczak, C. C. Steiner, S. Liu, M. T. P. Gilbert, M.-H. S. Sinding, S. Prost, K. Guschanski, S. K. S. S. Nathan, S. Brace, Y. L. Chan, C. W. Wheat, P. Skoglund, O. A. Ryder, B. Goossens, A. Götherström, L. Dalén, Genomic insights into the conservation status of the world's last remaining Sumatran rhinoceros populations. *Nat Commun* **12**, 2393 (2021).

72. N. Dussex, T. van der Valk, H. E. Morales, C. W. Wheat, D. Díez-del-Molino, J. von Seth, Y. Foster, V. E. Kutschera, K. Guschanski, A. Rhie, A. M. Phillippy, J. Korlach, K. Howe, W. Chow, S. Pelan, J. D. Mendes Damas, H. A. Lewin, A. R. Hastie, G. Formenti, O. Fedrigo, J. Guhlin, T. W. R. Harrop, M. F. Le Lec, P. K. Dearden, L. Haggerty, F. J. Martin, V. Kodali, F. Thibaud-Nissen, D. Iorns, M. Knapp, N. J. Gemmell, F. Robertson, R. Moorhouse, A. Digby, D. Eason, D. Vercoe, J. Howard, E. D. Jarvis, B. C. Robertson, L. Dalén, Population genomics of the critically endangered kākāpō. *Cell Genomics* **1**, 100002 (2021).

73. C. Bortoluzzi, C. J. Wright, S. Lee, T. Cousins, T. A. L. Genez, D. Thybert, F. J. Martin, L. Haggerty, T. D. T. of L. P. Consortium, M. Blaxter, R. Durbin, Lepidoptera genomics based on 88 chromosomal reference sequences informs population genetic parameters for conservation. bioRxiv [Preprint] (2023). https://doi.org/10.1101/2023.04.14.536868.

74. H. A. Jackson, L. Percival-Alwyn, C. Ryan, M. F. Albeshr, L. Venturi, H. E. Morales, T. C. Mathers, J. Cocker, S. A. Speak, G. G. Accinelli, T. Barker, D. Heavens, F. Willman, D. Dawson, L. Ward, V. Tatayah, N. Zuël, R. Young, L. Concannon, H. Whitford, B. Clavijo, N. Bunbury, K. M. Tyler, K. Ruhomaun, M. K. Grace, M. W. Bruford, C. G. Jones, S. Tollington, D. J. Bell, J. J. Groombridge, M. Clark, C. Van Oosterhout, Genomic erosion in a demographically recovered bird species during conservation rescue. *Conservation Biology* **36**, e13918 (2022).

75. D. L. Gilroy, K. P. Phillips, D. S. Richardson, C. van Oosterhout, Toll-like receptor variation in the bottlenecked population of the Seychelles warbler: computer simulations see the "ghost of selection past" and quantify the "drift debt." *J Evol Biol* **30**, 1276–1287 (2017).

76. C. Grossen, F. Guillaume, L. F. Keller, D. Croll, Purging of highly deleterious mutations through severe
bottlenecks in Alpine ibex. *Nat Commun* **11**, 1001 (2020).
77. A. García-Dorado, A. Caballero, Neutral genetic diversity as a useful tool for conservation biology. *Conserv*
Genet **22**, 541–545 (2021).
78. L. B. Pettersson, H. Arnberg, *Svensk Dagfjärilsövervakning, Årsrapport För 2023* (Biologiska institutionen,
Lunds universitet, Lund, 2024;
https://www.dagfjarilar.lu.se/sites/default/files/public/pdf/sebms_2023_korr.pdf).
79. V. G. Twort, J. Minet, C. W. Wheat, N. Wahlberg, Museomics of a rare taxon: placing Whalleyanidae in the
Lepidoptera Tree of Life. *Systematic Entomology* **46**, 926–937 (2021).
80. F. Mölder, K. Jablonski, B. Letcher, M. Hall, C. Tomkins-Tinch, V. Sochat, J. Forster, S. Lee, S. Twardziok, A.
Kanitz, A. Wilm, M. Holtgrewe, S. Rahmann, S. Nahnsen, J. Köster, Sustainable data analysis with
Snakemake [version 2; peer review: 2 approved]. *F1000Research* **10** (2021).
81. Z. J. Nolen, PopGLen—a Snakemake pipeline for performing population genomic analyses using genotype
likelihood-based methods. *Bioinformatics* **41**, btaf105 (2025).
82. S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**,
i884–i890 (2018).
83. H. Li, R. Durbin, Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**,
1754–1760 (2009).
84. R. Martiniano, E. Garrison, E. R. Jones, A. Manica, R. Durbin, Removing reference bias and improving indel
calling in ancient DNA data analysis by mapping to a sequence variation graph. *Genome Biol* **21**, 250 (2020).
85. K. Lohse, Wellcome Sanger Institute Tree of Life programme, Wellcome Sanger Institute Scientific
Operations: DNA Pipelines collective, Tree of Life Core Informatics collective, R. Vila, Darwin Tree of Life
Consortium, The genome sequence of the Common Blue, *Polyommatus icarus* (Rottemburg, 1775) [version
1; peer review: 1 approved]. *Wellcome Open Research* **8** (2023).
86. A. Hayward, K. Lohse, D. Laetsch, R. Vila, E. Taluy, The genome sequence of the silver-studded blue,
Plebejus argus (Linnaeus, 1758) [version 1; peer review: 2 approved]. *Wellcome Open Research* **7** (2022).
87. K. Lohse, A. Hayward, D. Laetsch, V. Marques, R. Vila, C. Tyler-Smith, Wellcome Sanger Institute Tree of
Life programme, Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective, Tree of Life Core
Informatics collective, Darwin Tree of Life Consortium, The genome sequence of the Mazarine Blue, *Cyaniris*
semiargus (Rottemburg, 1775) [version 1; peer review: 2 approved]. *Wellcome Open Research* **8** (2023).
88. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv
arXiv:1303.3997 [Preprint] (2013). https://doi.org/10.48550/arXiv.1303.3997.
89. Picard toolkit. *Broad Institute, GitHub Repository* (2019).
90. G. A. Van der Auwera, B. D. O'Connor, *Genomics in the Cloud: Using Docker, GATK, and WDL in Terra*
(O'Reilly Media, Inc., 2020).
91. P. Danecek, J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard, A. Whitwham, T. Keane, S. A.
McCarthy, R. M. Davies, H. Li, Twelve years of SAMtools and BCFtools. *GigaScience* **10** (2021).
92. E. Garrison, J. Sirén, A. M. Novak, G. Hickey, J. M. Eizenga, E. T. Dawson, W. Jones, S. Garg, C. Markello,
M. F. Lin, B. Paten, R. Durbin, Variation graph toolkit improves read mapping by representing genetic
variation in the reference. *Nat Biotechnol* **36**, 875–879 (2018).
93. A. Peltzer, G. Jäger, A. Herbig, A. Seitz, C. Kniep, J. Krause, K. Nieselt, EAGER: efficient ancient genome
reconstruction. *Genome Biol* **17**, 1–14 (2016).
94. G. Jun, M. K. Wing, G. R. Abecasis, H. M. Kang, An efficient and scalable analysis framework for variant
extraction and refinement from population-scale DNA sequence data. *Genome Res.* **25**, 918–925 (2015).
95. J. Neukamm, A. Peltzer, K. Nieselt, DamageProfiler: fast damage pattern calculation for ancient DNA.
Bioinformatics **37**, 3652–3653 (2021).
96. K. L. Korunes, K. Samuk, pixy: Unbiased estimation of nucleotide diversity and divergence in the presence of
missing data. *Molecular Ecology Resources* **21**, 1359–1368 (2021).
97. R. R. Hudson, M. Slatkin, W. P. Maddison, Estimation of levels of gene flow from DNA sequence data.
Genetics **132**, 583–589 (1992).
98. G. Bhatia, N. Patterson, S. Sankararaman, A. L. Price, Estimating and interpreting FST: The impact of rare
variants. *Genome Res.* **23**, 1514–1521 (2013).
99. R. K. Waples, A. Albrechtsen, I. Moltke, Allele frequency-free inference of close familial relationships from
genotypes or low-depth sequencing data. *Molecular Ecology* **28**, 35–48 (2019).
100. K. Hanghøj, I. Moltke, P. A. Andersen, A. Manica, T. S. Korneliussen, Fast and accurate relatedness
estimation from high-throughput sequencing data in the presence of inbreeding. *GigaScience* **8**, giz034
(2019).
101. A. Manichaikul, J. C. Mychaleckyj, S. S. Rich, K. Daly, M. Sale, W.-M. Chen, Robust relationship inference in
genome-wide association studies. *Bioinformatics* **26**, 2867–2873 (2010).
102. T. S. Korneliussen, A. Albrechtsen, R. Nielsen, ANGSD: Analysis of Next Generation Sequencing Data. *BMC*
Bioinformatics **15**, 1–13 (2014).

103. J. Meisner, A. Albrechtsen, Inferring Population Structure and Admixture Proportions in Low-Depth NGS
Data. *Genetics* **210**, 719–731 (2018).
104. E. A. Fox, A. E. Wright, M. Fumagalli, F. G. Vieira, ngsLD: evaluating linkage disequilibrium using genotype
likelihoods. *Bioinformatics* **35**, 3855–3856 (2019).
105. L. Skotte, T. S. Korneliussen, A. Albrechtsen, Estimating Individual Admixture Proportions from Next
Generation Sequencing Data. *Genetics* **195**, 693–702 (2013).
106. P. Pečnerová, G. Garcia-Erill, X. Liu, C. Nursyifa, R. K. Waples, C. G. Santander, L. Quinn, P. Frandsen, J.
Meisner, F. F. Stæger, M. S. Rasmussen, A. Brüniche-Olsen, C. Hviid Friis Jørgensen, R. R. da Fonseca, H.
R. Siegismund, A. Albrechtsen, R. Heller, I. Moltke, K. Hanghøj, High genetic diversity and low differentiation
reflect the ecological versatility of the African leopard. *Current Biology* **31**, 1862-1871.e5 (2021).
107. G. Garcia-Erill, A. Albrechtsen, Evaluation of model fit of inferred admixture proportions. *Molecular Ecology
Resources* **20**, 936–949 (2020).
108. V. Narasimhan, P. Danecek, A. Scally, Y. Xue, C. Tyler-Smith, R. Durbin, BCFtools/RoH: a hidden Markov
model approach for detecting autozygosity from next-generation sequencing data. *Bioinformatics* **32**, 1749–
1751 (2016).
109. E. Santiago, I. Novo, A. F. Pardiñas, M. Saura, J. Wang, A. Caballero, Recent Demographic History Inferred
by High-Resolution Analysis of Linkage Disequilibrium. *Molecular Biology and Evolution* **37**, 3642–3653
(2020).
110. I. Novo, P. Ordás, N. Moraga, E. Santiago, H. Quesada, A. Caballero, Impact of population structure in the
estimation of recent historical effective population size by the software GONE. *Genet Sel Evol* **55**, 86 (2023).
111. M. Petr, B. C. Haller, P. L. Ralph, F. Racimo, *slendr*: a framework for spatio-temporal population genomic
simulations on geographic landscapes. *Peer Community Journal* **3** (2023).
112. F. Baumdicker, G. Bisschop, D. Goldstein, G. Gower, A. P. Ragsdale, G. Tsambos, S. Zhu, B. Eldon, E. C.
Ellerman, J. G. Galloway, A. L. Gladstein, G. Gorjanc, B. Guo, B. Jeffery, W. W. Kretzschumar, K. Lohse, M.
Matschiner, D. Nelson, N. S. Pope, C. D. Quinto-Cortés, M. F. Rodrigues, K. Saunack, T. Sellinger, K.
Thornton, H. van Kemenade, A. W. Wohns, Y. Wong, S. Gravel, A. D. Kern, J. Koskela, P. L. Ralph, J.
Kelleher, Efficient ancestry and mutation simulation with msprime 1.0. *Genetics* **220**, iyab229 (2022).
113. P. D. Keightley, A. Pinharanda, R. W. Ness, F. Simpson, K. K. Dasmahapatra, J. Mallet, J. W. Davey, C. D.
Jiggins, Estimation of the Spontaneous Mutation Rate in Heliconius melpomene. *Molecular Biology and
Evolution* **32**, 239–243 (2015).
114. W. McLaren, L. Gil, S. E. Hunt, H. S. Riat, G. R. S. Ritchie, A. Thormann, P. Flicek, F. Cunningham, The
Ensembl Variant Effect Predictor. *Genome Biol* **17**, 122 (2016).
115. E. V. Davydov, D. L. Goode, M. Sirota, G. M. Cooper, A. Sidow, S. Batzoglou, Identifying a High Fraction of
the Human Genome to be under Selective Constraint Using GERP++. *PLOS Computational Biology* **6**,
e1001025 (2010).
116. V. E. Kutschera, M. Kierczak, T. van der Valk, J. von Seth, N. Dussex, E. Lord, M. Dehasque, D. W. G.
Stanton, P. E. Khoonsari, B. Nystedt, L. Dalén, D. Díez-del-Molino, GenErode: a bioinformatics pipeline to
investigate genome erosion in endangered and extinct species. *BMC Bioinformatics* **23**, 1–17 (2022).
117. K. Lohse, D. Laetsch, R. Vila, Darwin Tree of Life Barcoding collective, Wellcome Sanger Institute Tree of Life
programme, Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective, Tree of Life Core
Informatics collective, Darwin Tree of Life Consortium, The genome sequence of the small copper, *Lycaena
phlaeas* (Linnaeus, 1760) [version 1; peer review: 2 approved with reservations]. *Wellcome Open Research* **6**
(2021).
118. Darwin Tree of Life Consortium, A. Hayward, C. Wright, Darwin Tree of Life Barcoding collective, Wellcome
Sanger Institute Tree of Life programme, Wellcome Sanger Institute Scientific Operations: DNA Pipelines
collective, Tree of Life Core Informatics collective, The genome sequence of the holly blue, *Celastrina
argiolus* (Linnaeus, 1758) [version 1; peer review: 2 approved]. *Wellcome Open Research* **6** (2021).
119. Darwin Tree of Life Consortium, J. Hinojosa Galisteo, R. Vila, Darwin Tree of Life Barcoding collective,
Wellcome Sanger Institute Tree of Life programme, Wellcome Sanger Institute Scientific Operations: DNA
Pipelines collective, Tree of Life Core Informatics collective, The genome sequence of the green-underside
blue, *Glaucopsyche alexis* (Poda, 1761) [version 1; peer review: 2 approved]. *Wellcome Open Research* **6**
(2021).
120. Darwin Tree of Life Consortium, K. Lohse, A. Hayward, R. Vila, C. Howe, Wellcome Sanger Institute Tree of
Life programme, Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective, Tree of Life Core
Informatics collective, The genome sequence of the Adonis blue, *Lysandra bellargus* (Rottemburg, 1775)
[version 1; peer review: 2 approved, 2 approved with reservations]. *Wellcome Open Research* **7** (2022).
121. S. Kumar, M. Suleski, J. M. Craig, A. E. Kasprowicz, M. Sanderford, M. Li, G. Stecher, S. B. Hedges,
TimeTree 5: An Expanded Resource for Species Divergence Times. *Molecular Biology and Evolution* **39**,
msac174 (2022).

Supplementary Material References

122. The Darwin Tree of Life Project Consortium, Sequence locally, think globally: The Darwin Tree of Life Project.
Proceedings of the National Academy of Sciences **119**, e2115642118 (2022).
123. J. M. Flynn, R. Hubley, C. Goubert, J. Rosen, A. G. Clark, C. Feschotte, A. F. Smit, RepeatModeler2 for
automated genomic discovery of transposable element families. *Proceedings of the National Academy of*
Sciences **117**, 9451–9457 (2020).
124. A. Smit, R. Hubley, P. Green, RepeatMasker Open-4.0, version 4.1.2 (2013); http://www.repeatmasker.org.
125. S. Dolenz, T. van der Valk, C. Jin, J. Oppenheimer, M. B. Sharif, L. Orlando, B. Shapiro, L. Dalén, P. D.
Heintzman, Unravelling reference bias in ancient DNA datasets. *Bioinformatics* **40**, btae436 (2024).
126. M. Schubert, A. Ginolhac, S. Lindgreen, J. F. Thompson, K. A. AL-Rasheid, E. Willerslev, A. Krogh, L.
Orlando, Improving ancient DNA read mapping against modern reference genomes. *BMC Genomics* **13**, 178
(2012).
127. M. Kircher, "Analysis of High-Throughput Ancient DNA Sequencing Data" in *Ancient DNA: Methods and*
Protocols, B. Shapiro, M. Hofreiter, Eds. (Humana Press, Totowa, NJ, 2012; https://doi.org/10.1007/978-1-
61779-516-9_23), pp. 197–228.
128. R. N. Lou, A. Jacobs, A. P. Wilder, N. O. Therkildsen, A beginner's guide to low-coverage whole genome
sequencing for population genomics. *Molecular Ecology* **30**, 5966–5993 (2021).

Acknowledgements

We would like to thank Melina Eberhagen, Amanda Görnerup, Kajsa Svensson, Aske Brydegaard

Runemark, and Mikkel Brydegaard for assistance with collecting butterflies, Håkan Elmqvist for providing

pinned *Sc. orion* specimen, Göran Engqvist for providing tissue from the pinned specimen of *Pl.*

argyrognomon, Emma Kärrnäs and Isolde van Riemsdijk for assistance with DNA extractions, Jadranka

Rota for assistance with the entomological collections of the Biological Museum, Lund University

(https://doi.org/10.15468/dahk2a), Verena Kutschera and the Swedish Bioinformatics Advisory Program

for advice regarding the bioinformatic analyses, Anna Penna and Danielle Clake for comments on the

manuscript, and Isolde van Riemsdijk, Rachel Steward, Kalle Tunström, Simon Jacobsen Ellerstrand,

Colin Olito, Magne Friberg, Mikkel Skovrind, and Carolina Pacheco for helpful discussions regarding the

study species, methods and interpretations. We would like to thank Project Psyche and the Darwin Tree

of Life project for generating and providing the reference genomes that made these analyses possible.

We acknowledge support from the National Genomics Infrastructure in Genomics Production Stockholm

funded by Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish

Research Council. Computational resources were provided by the National Academic Infrastructure for

Supercomputing in Sweden (NAISS) and the Swedish National Infrastructure for Computing (SNIC) at

UPPMAX partially funded by the Swedish Research Council through grant agreements no. 2022-06725

and no. 2018-05973. This work was supported by grants from the strategic research area "Biodiversity

and Ecosystem services in a Changing Climate" (BECC), the Foundation Oscar och Lili Lamms Minne,

and the Swedish Research Council for Sustainable Development (grant 2021-01163) to AR and the

Jörgen Lindström's Foundation to ZJN.

Author Contributions

Z.J.N. - Conceptualization, Methodology, Formal analysis, Investigation (Lead), Data curation, Writing – original draft, Visualization, Funding acquisition. **P.J.** - Investigation (Supporting), Writing – review & editing. **A.S.T.L.** - Investigation (Supporting), Writing - review & editing. **N.W.** - Methodology, Writing – review & editing, Resources, Supervision. **A.R.** - Conceptualization, Methodology, Writing – original draft, Supervision, Funding acquisition.

Competing Interests

The authors declare that they have no competing interests.

Data Availability Statement

The genetic data that support these findings will be made available upon publication at NCBI in BioProjects PRJNA1068054 (focal species, modern) & PRJNA1168917 (focal species, historical; additional species). Bioinformatic workflow configurations, extensions, and resources specific to this study are available at https://github.com/zjnolen/polyommatini-temporal-genomics and archived at https://doi.org/10.5281/zenodo.13902816.