Species-specific loss of genetic diversity and exposure of deleterious mutations following agricultural intensification

Zachary J. Nolen[1*], Patrycja Jamelska[1,2], Ana Sofia Torres Lara[1,3], Niklas Wahlberg[1], Anna Runemark[1]

[1] Department of Biology, Lund University, Lund, Sweden 22362
[2] Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland 2000
[3] Faculty 2 Biology/Chemistry, University of Bremen, Bremen, Germany 28359

*Corresponding author: Zachary J. Nolen

Email: zachary.nolen@biol.lu.se

Author Contributions: Zachary J. Nolen: Conceptualization, Methodology, Formal analysis, Investigation (Lead), Data curation, Writing – original draft, Visualization, Funding acquisition. **Patrycja Jamelska**: Investigation (Supporting). **Ana Sofía Torres Lara**: Investigation (Supporting), Writing - review & editing. **Niklas Wahlberg:** Methodology, Writing – review & editing, Supervision. **Anna Runemark:** Conceptualization, Methodology, Writing – review & editing, Supervision, Funding acquisition.

Keywords: Conservation, agricultural intensification, inbreeding, genetic load, genomic erosion

Abstract

Agriculture driven land-use change is a major contributor to insect declines globally. As a result of population decline and subdivision, species lose genetic diversity, potentially reducing adaptive potential and increasing genetic load, reinforcing decline. We examine how genetic diversity has changed over the past century in three Polyommatinae butterfly species in Sweden, leveraging genome sequencing of specimens from museum collections. Based on 63 historical and 92 contemporary genomes, we document a general decline in genetic diversity and reveal that the extent of decline depends on habitat specialization. The habitat generalist has retained most of its genetic diversity and connectivity. In contrast, the two specialists have declined sufficiently that even common variation is being eroded by drift, with up to a 3.3% reduction in heterozygosity per decade in the strongly grassland associated *Cyaniris semiargus*. While this sharp decline has reduced the burden of strongly deleterious mutations in the specialist species, increased population isolation has led to inbreeding in *Cy. semiargus*, exposing a greater proportion of weakly deleterious mutations in homozygous state. This realized burden likely reduces mean fitness and reinforces population decline, highlighting the need for well-connected grassland landscapes to maintain grassland specialist functional connectivity and genetic diversity. We complement these analyses with seven additional Polyommatinae species, uncovering that declines are common across the group, with alarming rates of genetic decline in species listed as viable. This study highlights the urgency of incorporating genetic data into conservation planning, as threats from genetic erosion are not captured by current conservation assessment.

Introduction

Insect decline has received increased attention in recent years as reports uncover alarming declines in insect assemblages and biomass (1). Efforts to quantify this decline estimate that terrestrial insects are declining at an average rate of 11% per decade globally, with this trend being primarily driven by even stronger regional declines in Europe and North America (2). This stark decline is the result of a variety of anthropogenic pressures, primarily climate change and land use changes such as agricultural intensification and urbanization (1, 3–7). Intensive agricultural landscapes place a particularly strong pressure on grassland insect species (6), especially in Europe where formerly biodiverse semi-natural grasslands have been abandoned or converted into intensively cultivated land (8).

While these declines already have alarming implications for natural communities and ecosystem services in which insects play key roles, they can also leave insect populations with genetic challenges to long term persistence. Smaller populations are expected to lose genetic diversity faster than larger populations due to the stronger relative force of drift (9). This reduces their ability to adapt to changing environments (10, 11) and may allow partially recessive, weakly deleterious mutations to accumulate and become unmasked through inbreeding and weakened selection (12, 13). Such accumulation of weakly deleterious mutations increases the genetic load, classically defined as the reduction in population fitness from an optimum genotype (14). This, in turn, may reinforce demographic decline in a process referred to as mutational meltdown (15). An increasing number of studies even document that decline of genetic diversity and accumulation of deleterious mutations may continue after demographic recovery due to drift debt (16–19).

How to assess the threat genetic factors pose to natural populations is still an active area of research, with several genetic monitoring approaches being suggested and implemented (20–23). Interpretation of genetic diversity is complex, as it is influenced by a multitude of factors (24), such as ecology and life history (25), genomic architecture (26), and demographic trends over thousands to millions of years (27). Some empirical studies

have shown that populations may persist despite low levels of genetic diversity (28, 29)

and that relationships between genetic diversity and extinction risk are context dependent

(30). Therefore the value of neutral genetic diversity in conservation management has been

subject to some debate (31–33).

As a means to control for factors that may make interpreting absolute estimates of genetic

diversity challenging, measures that estimate the rate of genetic erosion have seen

increasing focus in genetic monitoring. Programs in several countries use genetic metrics

measured at multiple time points or that directly estimate ongoing genetic decline (21, 34,

35). Examining decline directly rather than setting thresholds for absolute levels of genetic

diversity enables identifying populations for which 'low' genetic diversity is likely to

increase extinction risk. For instance, larger populations are expected to have higher

numbers of segregating deleterious mutations (36), making them more prone to inbreeding

depression following population decline (12, 37, 38).

Temporal sampling is, however, limited to estimating genetic decline from the first

sampled time point, with considerable erosion likely to have occurred several decades

before. To overcome this, genomic resources from historic samples preserved in natural

history museums can extend monitoring further back in time. The direct measures of pre-

and early bottleneck genetic parameters museum samples provide are crucial to fully

understand the extent and character of genetic decline (23). Comparisons with historical

samples may even allow approximating changes in genetic load across bottlenecks by

comparing counts of putatively deleterious mutations (12, 19, 23, 39–41). Such studies

have the potential to identify populations that face reduced fitness from decline.

Here, we use museomics to address to what extent genetic diversity has been lost and

functional connectivity reduced in grassland associated butterflies during the second

wave of agricultural modernization in Sweden starting in the 1940s (42, 43). During this

modernization, small farms were abandoned, and their associated semi-natural

grasslands began to decline considerably (43). We examine three focal butterfly species in

this landscape that vary in habitat specialization, ranging from grassland generalist to

specialist, enabling us to examine if patterns of genetic decline are stronger in more

specialized species, in line with expectations from greater demographic declines in

specialist insect abundance compared to generalists (44–46). By comparing pairs of

historical population samples collected in southern Sweden in the early 20th century with

their contemporary counterparts, we document species-specific losses of genetic

diversity and functional connectivity, and increased homozygosity of putatively deleterious

mutations that is expected to have increased genetic load in the most specialized species.

We further examine if patterns of loss of genetic diversity are general, and if loss of genetic

diversity is reflected in current conservation status, by extending the dataset to include

historical and modern individuals of seven additional related butterfly species with

Swedish Red List (47) conservation statuses ranging from least concern to critically

endangered.

Results

We analyzed changes in genetic diversity, differentiation, and deleterious mutation burden

in three focal Polyommatinae (Lepidoptera: Lycaenidae) butterfly species: *Polyommatus*

icarus, a grassland generalist, *Plebejus argus*, a heathland specialist, and *Cyaniris*

semiargus, a grassland specialist (Figure 1A). We generated whole genome resequencing

data from pinned museum specimens from the entomological collections of the Biological

Museum at Lund University (48) collected from 1917-1956 (Table S1; *N* = 28 - *Po. icarus*; 12

- *Pl. argus*; 14 - *Cy. semiargus*) in southern Sweden. These individuals were sampled in

populations generally collected at the same locality during the same year, and compared

with WGS data from samples collected at corresponding localities in 2020-2023 by Nolen

et al (49) (Figure 1B; Table S1; *N* = 27 - *Po. icarus*; 29 - *Pl. argus*; 29 - *Cy. semiargus*). As

DNA extracted from historical specimens is fragmented and subject to post-mortem DNA

damage (50, 51) (Figure S1), we confirmed shorter fragments did not induce a reference

bias in the historical samples (Figure S2). To address DNA damage, we tested post-

mortem DNA damage correction I) using MapDamage2 (52) and II) removing all transition

sites, which are the most common changes caused by DNA damage (50, 51). Rescaling

base quality scores with MapDamage2 biased historical samples towards the reference in

two of the three focal species, whereas transition removal did not (Figure S2). All

subsequent results were therefore generated without MapDamage2 correction and with

transitions removed. To account for lower sequencing depth in historical samples, we

performed analyses both at full depth and with samples subsampled to equivalent mean

sequencing depths, presenting the subsampled results in the main text, with full depth

results included in the Appendix.

General reductions in genetic diversity

To assess if agricultural intensification and the associated reduction in semi-natural

grassland habitat has resulted in reduced genetic diversity, we compared individual

genome-wide heterozygosity of historical and modern samples within each species. In all

three species, heterozygosity is lower in modern individuals, but considerably more so in

the two specialist species compared to the generalist (Welch two-sample t-test: P = 0.006

in *Po. icarus*, < 0.001 in *Pl. argus*, and < 0.001 in *Cy. semiargus;* Figure 1C; Tables S2-3). In

the specialist species, *Pl. argus* and *Cy. semiargus*, population-specific estimates of

genetic diversity are lower in all modern populations compared to their historical

counterparts, both for nucleotide diversity (π; Wilcoxon signed-rank test: median effect

size range [-0.52, -0.40] in *Pl. argus* and [-0.76, -0.51] in *Cy. semiargus*) and Watterson's θ

(median effect size range [-0.77, -0.68] in *Pl. argus* and [-0.86, -0.70] in *Cy. semiargus*),

even after subsampling to account for variation in sample size (Figure S3; Tables S4-5). In

the generalist, *Po. icarus*, only Watterson's θ shows declines robust to variation in sample

size, and only in two of the three populations (median effect size range [-0.29, -0.24];

Figure S3; Table S4-5). Tajima's D is higher in modern populations of all three species

compared to their historical counterparts, consistent with population bottlenecks across

the past century (median effect size range [0.63, 0.68] in *Po. icarus*, [0.82, 0.84] in *Pl.*

argus, and [0.73, 0.85] in *Cy. semiargus*; Figure S3; Table S4-5).

Reduced connectivity may accelerate loss of genetic diversity

While genetic diversity decline is an expected product of demographic decline, it can

additionally be accelerated by reductions in gene flow (53). We reveal a loss of functional

connectivity among modern populations, reflected in the stronger contemporary

population structure uncovered by both genetic differentiation and clustering approaches,

likely contributing to the loss of genetic diversity. *Po. icarus* and *Cy. semiargus* have

stronger genetic differentiation (F_{ST}) between modern than historical population pairs

(Paired t-test: ΔF_{ST} = 0.014, *P* = 0.007 in *Po. icarus* and ΔF_{ST} = 0.053, *P* = 0.036 in *Cy.*

semiargus; Figure 1F; Table 6-7). F_{ST} increased in *Pl. argus* in the single population pair

present in both time periods (ΔF_{ST} = 0.021; Table S7). Including all modern population pairs

in the analysis estimates mean F_{ST} increases of 0.027 and 0.071 in *Pl. argus* and *Cy.*

semiargus, respectively (Welch two-sample t-test: *P* = 0.001 in *Cy. semiargus*; Table S7).

Principal component analyses (PCAs) and admixture analyses also reveal generally

stronger population structure in modern samples than historical, and strongest in the most

specialized *Cy. semiargus* (Figure 2A). Along PC1, the greatest axis of variation, both *Po.*

icarus and *Pl. argus* are clustered primarily into historical and modern samples, consistent

with the best fit admixture analyses for these species (K=2; Figure 2B; Table S8). In these

two species, modern samples show greater among population differences than historical

along PC2, however, consistent with increases in genetic differentiation between

localities. However, more than two clusters are visible on the PCA for these two species,

supported by the correlation of residuals for the admixture analyses (Figure S4). This

highlights that greater genetic structure likely exists in these two species, but described by

too little variation to reliably cluster samples by at these sample sizes. Each modern

population forms a distinct cluster in the specialized species, *Cy. semiargus*, while

historical populations cluster together in a single separate cluster (Figure 2B). The stronger

contemporary population structure is especially driven by isolation of populations in the

agricultural landscapes within Sweden's southernmost province, Skåne (Figure 1D;

Figure 2B; Figure S5).

Species-specific increases in inbreeding

The pronounced losses of functional connectivity and genetic diversity have resulted in

inbreeding within the specialist *Cy. semiargus*. This contemporary inbreeding is reflected

by an increase in the inbreeding coefficient F_{RoH} from historical to modern *Cy. semiargus*

samples (Welch two-sample t-test: ΔF_{RoH} = 0.065, P < 0.001; Figure 1D; Table S3&9). F_{RoH}

measures the proportion of the autosomal genome in runs of homozygosity greater than a

certain length (54), which we set to 100kb, roughly corresponding to identical by descent

tracts with shared ancestry in the past 200 generations (55). Across species, no historical

individual has a coefficient greater than 0.04, and historical species averages are all below

0.01. In *Cy. semiargus*, modern samples have a maximum coefficient of 0.33 and an

average coefficient of 0.072, an order of magnitude higher than the mean for historical

samples. In modern *Cy. semiargus*, individuals with higher inbreeding coefficients have

fewer runs of homozygosity relative to their cumulative length, which may indicate they

stem from close-kin matings (Figure 3A, Figure S6). For other individuals, there is a largely

linear relationship between the number of runs of homozygosity and their cumulative

length, suggesting that close-kin mating is not common overall and inbreeding is primarily

due to increases in background relatedness, consistent with patterns expected of a

population bottleneck (55).

Purging of deleterious mutation burden

Population decline and subdivision can increase the exposure of partially recessive,

deleterious mutations to selection due to increased homozygosity, facilitating purging of

strongly deleterious mutations (56–58). However, as populations decline and subdivide,

selection weakens relative to drift, allowing partially recessive, weakly deleterious

mutations to drift to higher frequencies and fix, increasing genetic load (13, 15, 59). Here,

we estimated deleterious mutation burden using relative counts of putatively deleterious

mutations per individual as a proxy for genetic load (as in e.g. 19, 39, 41, 60, 61). We

separate putatively deleterious mutations into three groups. Weakly deleterious include

missense mutations, strongly deleterious consist of e.g. loss of function mutations, and

highly conserved include derived mutations at positions with high sequence conservation

in Lepidoptera, presumed to be strongly deleterious. We find evidence consistent with

purging of deleterious mutations in both specialist species *Pl. argus* and *Cy. semiargus*,

with the relative count of strongly deleterious mutations being lower in modern compared

to historical individuals in both species (Welch two-sample t-test: P = 0.018 in *Pl. argus*

and 0.031 in *Cy. semiargus*; Figure 3B; Table S10). Burden from weakly deleterious

mutations has also decreased somewhat in *Pl. argus* (P < 0.001; Figure 3B; Table S10). In

generalist *Po. icarus*, burden from weakly and strongly deleterious mutations is similar in

modern and historical samples. Burden from derived alleles at highly conserved positions

positions has remained stable in all three species (Figure 3B; Table S10).

Increased homozygosity of weakly deleterious burden in *Cy. semiargus*

Weakly deleterious mutations are exposed in a homozygous state more often in modern

than historical *Cy. semiargus* individuals (P < 0.001; Figure 3C; Table S10). In contrast,

strongly deleterious mutations are homozygous at equal rates over time. In *Po. icarus* and

Pl. argus, we find no evidence for increases in homozygosity of weakly or strongly

deleterious mutations (Figure 3C; Table S10). However, derived variants at conserved

regions are homozygous to a greater extent in modern samples of both *Pl. argus* and *Cy.*

semiargus (P < 0.001 in *Pl. argus* and = 0.009 in *Cy. semiargus*; Figure 3C; Table S10). In *Cy.*

semiargus, individuals with higher inbreeding coefficients had higher homozygous, but not

total, deleterious burden across all categories of deleterious mutations (linear regression:

R^2 = 0.79, 0.18, 0.35 for weakly, strongly, and conserved deleterious variants, respectively;

Figure S7). This suggests that increased rates of inbreeding have increased homozygosity

of deleterious mutations, and that this effect is strongest for weakly deleterious mutations.

Genetic decline not reflected in current conservation status

We utilized the genetic decline indicator thresholds outlined by Andersson et al. (20) to

determine if current conservation status relates to the rate of genetic decline. These

thresholds suggest categorizations of 'Acceptable', 'Warning', and 'Alarm' rates of genetic

decline, reflecting >=95%, 75-94%, and <75% retention of genetic diversity over 100 years,

respectively. Based on changes in heterozygosity, nucleotide diversity, and Watterson's θ,

Po. icarus largely exhibits 'Acceptable' or 'Warning' declines, while *Pl. argus* and *Cy.*

semiargus exhibit 'Warning' or 'Alarm' declines, depending on the metric of genetic

diversity used (Figure 4&S8; Tables S5&11), which largely are in line with the concerns

raised by our deleterious mutation burden analyses. This is in stark contrast with the 2020

Swedish Red List of species (47), where all three focal species share a conservation status

of 'Least Concern'.

To assess if the loss of genetic diversity in the three focal species is representative of other

grassland butterfly species, we supplemented the dataset by generating whole genome

resequencing data for historical-modern pairs of seven additional species of

Polyommatinae butterfly (Table S1). We assessed if the conservation status of these

species, ranging from of 'Least Concern' to 'Critically Endangered' (47) (Table S12) capture

losses of genetic diversity uncovered in this data set. Comparing individual heterozygosity,

we find that current conservation status generally fails to capture the levels of genetic

erosion in these species (Figure 4A&S8; Table S11). Utilizing estimates at full individual

sequencing depth, least concern species predominantly fell into the warning category,

with only *Celastrina argiolus* and Öland *Po. icarus* falling in the acceptable category. Of

species classified as least concern, only *Cy. semiargus* fell into the alarm category. Two

endangered species, *Phengaris alcon* and *Scolitantides orion,* fell into the acceptable and

alarm categories, respectively, with *Ph. alcon* even increasing slightly in genetic diversity

over the time frame. *Plebejus argyrognomon*, a critically endangered species that is now

presumed extinct in Sweden (62), had already experienced genetic decline categorized as

'alarm' by the early 2000s, when our modern sample for this species was collected.

Results from our focal species illustrate that categorization can depend considerably on

the individuals/populations sampled (Figure 4). Categorization is largely consistent for

each species both at full sequencing depth and when subsampled to 2-3X sequencing

depth, but some species/populations bordering adjacent categories shift between them

upon subsampling (Table S11).

Discussion

Our analyses of 63 historical and 92 modern genomes uncover a decline in butterfly

genetic diversity over the past century. This decline is greater in the specialist species,

especially *Cy. semiargus*, where the once highly connected populations have become

isolated into distinct genetic clusters. This isolation and resulting decline in genetic

diversity has resulted in purging of highly deleterious burden in the two specialist species,

and even of weakly deleterious burden in *Pl. argus*, which showed intermediate decline.

However, in *Cy. semiargus*, where decline was greatest, homozygosity of weakly

deleterious burden has increased following inbreeding, most likely resulting in reduced

fitness for more inbred populations. The rates of decline uncovered in the three focal

species are also common in seven additional species of blue butterflies, despite most of

these species having conservation statuses of 'Least Concern'. Our findings reveal that

recent genetic erosion is likely to be ubiquitous across grassland butterflies in southern

Sweden. These results highlight the need for incorporation of genetic data into

conservation assessments of insect threat status, as census-based estimates fail to

detect threats posed by genetic decline.

Our findings of reduced genetic diversity across all three focal species are consistent with

widespread demographic decline in arthropod species in Europe over the past century,

with the modernization of agriculture as a major driver (5, 63). The highest average decline

in genetic diversity we document, an up to 3.3% reduction in heterozygosity per decade in

Cy. semiargus, appears to be accelerated by strengthened isolation between populations.

However, both *Po. icarus* and *Pl. argus*, where increases in population subdivision are

more moderate, have experienced declines of up to 0.7% and 2.7% per decade,

respectively. Population structure and genetic differentiation analyses suggest modest

increases in population isolation in these species, potentially reflecting that declines in

genetic diversity predate genomic signals of population subdivision. Population

differentiation accumulates over time, lagging behind barriers to gene flow (64, 65), with

the rate at which genetic differentiation accumulates after subdivision depending upon the

underlying effective population sizes of the populations (66).

Importantly, the finding that these three species have historically had low levels of genetic

differentiation across southern Sweden highlights that currently isolated populations may

be made up of partitions of historically well-connected metapopulations. Hence, concerns

about genetic rescue (67) wiping out locally adapted ancestry (68, 69) are not likely to be

relevant for these and other grassland associated insects in regions with historically

abundant habitat. Instead, restoring functional connectivity to reduce inbreeding and

associated accumulation of deleterious mutations (15, 70) is likely key to improve

prospects of long-term persistence. Previous work on *Cy. semiargus* has shown that the

inbreeding in these modern populations is related to low grassland land cover in the

surrounding landscape (49). Expanding green infrastructure and restoration of semi-

natural grassland habitat, enabling dispersal and buffering genetic decline, is likely needed

to remedy future isolation and inbreeding in grassland associated butterflies.

Understanding to what extent genetic diversity decline, resulting from habitat loss and

isolation, reduces the fitness of individuals and populations is crucial for determining

conservation priorities. To approximate such genetic load without direct fitness estimates,

we quantified deleterious mutation burden (71). Strongly deleterious mutation burden is

lower in modern individuals compared to historical for both specialist species. This is in

line with theoretical predictions (56–58) and empirical evidence (19, 60, 72–75) suggesting

that population decline can increase exposure of partially recessive, strongly deleterious

mutations to selection, reducing their frequency. Interestingly, weakly deleterious

mutation burden appears to have also decreased in *Pl. argus*, though only mildly (5%

reduction compared to 15.7% for strongly deleterious). Weakly deleterious mutations are

less likely to be efficiently purged in small populations than strongly deleterious ones (56,

58), often requiring low dominance coefficients for purging to be effective (56). Dominance

coefficients are generally lower for more deleterious segregating mutations (76), and

estimates for the mean dominance coefficients for weakly deleterious mutations are

around 0.2-0.25 (77, 78). Despite this, purging of the portion of weakly deleterious burden

with high recessivity has been described in the Iberian Lynx, largely occurring at

intermediate points of decline before populations were very small (60). Our findings of

reduced weakly deleterious burden in *Pl. argus*, which has declined intermediately to the

other two species, could partially be explained by purging of only the highly recessive

portion of burden, as predicted from models of deleterious mutation count during a

bottleneck (60).

Considering the partial recessivity of segregating deleterious mutations (77–79) and the

frequency of inbreeding depression in outcrossing organisms (59, 68, 80, 81), negative

fitness effects for deleterious mutations should be primarily realized when exposed in

homozygous state. Genetic load, in turn, is likely more closely approximated by the

homozygous rather than total deleterious burden. In the grassland specialist *Cy.*

semiargus, burden has increased in homozygosity at both conserved positions and for

weakly deleterious mutations. As weakly deleterious burden is strongly associated with

inbreeding coefficient, this can be interpreted as unmasking of the inbreeding load present

in the historical populations (12, 68, 82). Contemporary heterozygosity in *Cy. semiargus*

has a strong negative relationship to rates of inbreeding (49). A recent study associating

extinction risk with heterozygosity in the Glanville fritillary butterfly (*Melitaea cinxia*) found

that for small, isolated, recently declined populations, heterozygosity was negatively

associated with extinction and mortality (30). Increased homozygous deleterious mutation

burden as we observe in small, isolated, recently declined populations of *Cy. semiargus* is

likely what causes the fitness consequences observed in isolated *M. cinxia* populations

(30, 83, 84). Joint estimation of fitness and deleterious mutation burden is key to fully

understanding how well different measures of mutation burden approximate genetic load.

This information is necessary for using genomics based approximations of load to assess

conservation status of wild populations.

Furthermore, understanding the link between measures of deleterious mutation burden

and fitness-based estimates of load is important for assessing the validity of assumptions

underlying estimates of burden. For instance, the functional effect based estimates used

to classify weakly and strongly deleterious mutations are reliant upon the accuracy of

underlying gene models and, in our case, assume the reference allele is less deleterious

than the alternate. Estimates of burden using sequence conservation scores also assume

more conserved regions are more functionally important (12, 85). Our findings for

conserved sites more closely resemble those of weakly deleterious mutations, potentially

reflecting the limited ability of sequence conservation scores to distinguish between

weakly and strongly deleterious mutations (86, 87). In addition, GERP scores depend on

estimates of the rate of neutral evolution, approximating the rejected substitution rate. Our

GERP scores are based on a time tree (similar to 39, 40, 74, 86) and hence differ from

scores estimated based on neutral rates (e.g. 88). However, our identification of conserved

sites should be robust for within-species comparisons of deleterious burden, as they

represent the upper extreme of the distribution in our data. Despite these caveats, our

estimates of deleterious burden are consistent with expectations under the species-

specific patterns of genetic decline, underscoring their relevance to conservation. The

strong general pattern of increased deleterious burden expressed in a homozygous state

uncovers a potentially serious threat to *Cy. semiargus*. While we uncover fixation of

deleterious mutations within populations, almost none are fixed across all populations.

This calls for assisted migration and establishment of migration corridors to avoid local

mutational meltdown and extirpation in *Cy. semiargus*.

None of the three focal species are currently demographically declining at a rate to qualify

for the Swedish Red List based on demographic data. Systematic monitoring in Sweden

from 2010-2023 reveal declines in both *Po. icarus* and *Cy. semiargus*, while *Pl. argus*

appears to have stabilized (89). Demographic stability in *Pl. argus* does not necessarily

mean that genetic diversity has stabilized in the species, as populations may continue to

lose genetic diversity even after demographic recovery, a phenomenon termed drift debt

(16). In *Cy. semiargus*, genetic factors are likely already contributing to its demographic

decline. Our findings illustrate the importance of incorporating genetic monitoring into

conservation assessment (90, 91) to identify genetic decline before it results in expressed

genetic load. Temporal monitoring with historical samples provides an opportunity to

detect species facing recent declines in genetic diversity, separating them from those with

longer term low genetic diversity (23). Using this approach on several species of

Polyommatinae blues in southern Sweden, we document a general loss of genetic diversity

in grassland butterflies and uncover that additional species currently not on the Red List

likely face threats from reduced genetic diversity. Estimations of genetic decline along with

complementary estimates of deleterious burden provide a promising way to assess

genetic conservation concerns in the wild (19), but these analyses are limited by the

availability of high quality reference genomes. This underscores the importance of efforts

to generate high quality references for wild species to improve our ability to accurately

assess these metrics (92).

In conclusion, we document a general decline in genetic diversity in grassland butterflies in

southern Sweden, which is likely mirrored in other regions that have lost grasslands to

intensive agriculture and forestry. While demographic declines largely inform current

conservation status of these species, we show that this is insufficient to capture declines

in genetic diversity and resulting inbreeding, highlighting the acute need for genetic

monitoring of wild insects to prevent extirpation and extinction. *Cyaniris semiargus*, a once

widespread species in Sweden, provides a strong example of this, now persisting in

segmented populations with increased rates of inbreeding and expression of deleterious

burden, requiring prompt conservation action to reduce the risk of local extinction.

Methods

Study System and Sampling

To assess changes in genetic diversity, differentiation, inbreeding, and deleterious burden,

we selected three blue butterfly species (Polyommatinae) as focal species, *Polyommatus*

icarus, *Plebejus argus*, and *Cyaniris semiargus* (Figure 1A). These three species were

selected as they vary in habitat specialization, being grassland generalist, heathland

specialist, and grassland specialist, respectively, in order to assess how specialization

may impact genetic decline. For these three species, we used population-level genetic

data we previously generated (49) as our modern dataset. We selected pinned specimens

from the entomological collections of the Biological Museum, Lund University (48) that

corresponded to roughly the same geographic area as the modern samples for each

species. These historical specimens were sampled between 1917-1956, during the early

stages of the second wave of agricultural modernization in Sweden (42). For each historical

locality, we selected 4-11 specimens sampled in the same year to estimate population-

level metrics. The historical and modern population-level pairs were sampled in four broad

geographic areas: I) Western and II) Eastern Skåne, III) Småland, and IV) Öland (Figure 1B),

with historical localities ranging approximately 12-40km from their modern counterparts.

This resulted in historical-modern population-level pairs for three regions in *Po. icarus*

(Western and Eastern Skåne and Öland), two regions in *Pl. argus* (Småland and Öland), and

three regions in *Cy. semiargus* (Western and Eastern Skåne and Småland). While most

regions consist of only one historical-modern population pair, we additionally compared a

modern *Cy. semiargus* population in Southeastern Skåne to the historical Eastern Skåne

population. We were unable to sample population-level historical data for *Pl. argus* in

Skåne, however we did sample one historical specimen each from Western and Eastern

Skåne, which we include in individual level analyses, pairing them with the modern

samples from Southwestern and Eastern Skåne, respectively.

To more broadly assess the variation in rates of genetic decline in Polyommatinae of

differing conservation statuses, we sampled historical and modern individuals of seven

additional species: *Agriades optilete, Aricia agestis, Celastrina argiolus, Polyommatus*

amandus, Phengaris alcon, Scolitantides orion, and *Plebejus argyrognomon*. Modern

specimens of *Ag. optilete, Ar. agestis, Ce. argiolus,* and *Po. amandus* were collected

across the same geographic regions as the focal species in 2021 (Table S1). Pinned

specimens to be used as modern samples of *Ph. alcon* and *S. orion,* collected in 2003 in

Western Skåne and 2016 in Södermanland, respectively, were donated to this study by

Håkan Elmqvist. Legs from a pinned speciemen of *Pl. argyrognomon*, collected in 2003 in

Western Skåne, were donated by Göran Engqvist and used as the modern sample for this

species. The historical samples for these eight species all were selected to be proximal to

the modern specimen sampling locality with collection dates ranging from 1932-1967, all

provided as pinned specimens from the entomological collections of the Biological

Museum, Lund University (48).

DNA Extraction and Sequencing

DNA from fresh specimens was extracted and sequenced following the methods in Nolen

et al (49). We extracted DNA and prepared sequencing libraries from pinned museum

specimens in a facility dedicated to historical DNA processing at Lund University, following

the protocol described by Twort et al (93). Libraries prepared for pinned specimens were

pooled into two equimolar pools containing 37-38 libraries each, sequencing each pool on

a single NovaSeq 6000 S4 lane. DNA was extracted from the abdomens or legs of male

specimens without homogenization to preserve anatomical structures. Abdomens were

then stored in glycerol together with the original specimen.

Genomic Analyses

We performed the majority of our analyses using the PopGLen pipeline v0.3.0 (94), a

Snakemake (95) pipeline we have built that combines several genotype likelihood based

population genomic analyses. Genotype likelihood based analyses use a probabilistic

framework to account for the uncertainty that comes with low coverage sequencing data,

reducing biases related to lower sequencing depth in our historical samples. We utilized

genotype likelihood based analyses for all results, except for the estimation of inbreeding

coefficients and genetic load, where the chosen methods required called genotypes.

Sequence Processing

We trimmed both modern and historical raw sequencing data using fastp v0.23.4 (96) with

default settings, additionally collapsing overlapping read pairs in historical samples,

discarding those with less than 30bp overlap. This processing of historical samples

discards short (<30bp) fragments that contain little data usable for unique mapping, as

well as long fragments (>270bp) that are more likely to be modern contaminants.

We mapped the trimmed, paired modern reads separately per sequencing run to the

respective species-specific reference genomes, or to the closest relative with an available

chromosome-level assembly (Table S12), using BWA mem v0.7.17 (97). We mapped the

collapsed historical reads to the same reference genomes using BWA aln, which is the

recommended algorithm for reads shorter than 70bp (98) using settings optimized for

ancient/historical DNA (`-1 16500 -n 0.01 -o 2`) (99).

We merged mapped sequencing runs per sample using Samtools v1.17 (100) and removed

PCR and optical duplicates using Picard v2.27 (101) with default settings except for an

optical duplicate pixel distance of 2500, as recommended for patterned flow cell data. We

removed PCR duplicates for our historical samples using dedup v0.12.8 (102), which

identifies duplicates using both ends of the collapsed sequencing read.

We realigned around indels using the GATK IndelRealigner (103). As the primary tool we are

using, ANGSD (104), counts overlapping reads twice (105), we additionally clipped

overlapping read pairs in the modern samples using BamUtil v1.0.15 (106). We estimated

post-mortem DNA damage in historical samples using DamageProfiler v1.1 (107).

Site and read filtering

We utilized a site based filtering scheme similar to that implemented by Pečnerová et al

(108) in which we developed multiple independent sets of filtered sites, which we

intersected to create a global list of filtered sites for all analyses. We performed this

filtering for each species dataset separately.

We excluded sex-linked chromosomes, mitochondria, and scaffolds less than 1 Mb in

length. We excluded repeats identified by RepeatMasker v4.1.2 (109), using the reference

specific repeat libraries provided by the Darwin Tree of Life Project

(https://projects.ensembl.org/darwin-tree-of-life/). We estimated regions in the lower 1%

and upper 99% of global sequencing depth distribution separately across all samples,

historical samples, and modern samples, excluding sites that appeared in any of the three

sets. The number and proportion of sites passing each of these filters, as well as passing

the final intersected filter set, are reported for each species in Table S13. We estimated

mean sequencing depth without filtering, as well as within the filtered site set, for each

sample using ANGSD (Table S1). All subsequent analyses were performed on the species-

specific filtered site sets.

Reference bias and post-mortem DNA damage

We investigated two potential methods to account for post-mortem DNA damage: I)

recalibration of putatively damaged bases using MapDamage2 (52), and II) removal of

transitions, where most DNA damage is expected to occur (50, 51). We assessed their

performance by comparing identity by state (IBS) similarity to the reference at filtered sites

between the two correction methods and the uncorrected data (Figure S2). We estimated

IBS similarity to the reference for each sample using ANGSD, sampling a single base at

each position (`-doIBS 1`), counting reference bases as 1 and non-reference bases as 0 (-

`doMajorMinor 4 -output01 1`). We calculated the average IBS similarity per individual

from these values, with 1 indicating complete similarity to the reference and 0 indicating

complete dissimilarity.

Historical samples without correction had less similarity to the reference than modern

(Figure S2), as expected for historical samples without substantial mapping bias due to

drift from the sampling time of the historical samples and the modern reference as well as

differences introduced by post-mortem DNA damage (41, 110). After MapDamage2

rescaling, historical samples showed higher similarity to the reference than modern

samples in *Po. icarus* and *Pl. argus*. As transition removal did not introduce this reference

bias, we proceeded using transition removal in all analyses.

Genotype Likelihood Calculations

We estimated genotype likelihoods in ANGSD v0.940 (104) using the GATK model (`-GL 2`),

using reads with a mapping quality greater than 30 (`-mapMapQ 30`) and base quality greater

than 20 (`-minQ 20`). We filtered our analyses to sites present in the species-specific filtered

sites list using the `-sites` option.

Heterozygosity and genetic diversity metrics

We estimated individual genome-wide heterozygosity for all samples using the single

sample site frequency spectrum (SFS) produced by realSFS, included with ANGSD. First,

we generated a sample allele frequency (SAF) file (`-doSaf 1`) for each sample, setting the

ancestral state (`-anc`) to the reference sequence. We generated a folded (`-fold 1`) SFS in

realSFS using the SAF file and calculated heterozygosity as the second value (count of

heterozygous sites) divided by the total count of analyzed sites. To test for differences in

mean individual heterozygosity across time periods, we performed an unequal variances *t*-

test comparing all samples within each time category for each of the focal species.

For the three focal species with population-level samples, we estimated nucleotide

diversity, Watterson's θ, and Tajima's D per sampled population using thetaStat, included

with ANGSD. As the estimates of these metrics, especially Watterson's θ and Tajima's D,

can depend on sample size, we made our estimates using 100 subsamples of each

population to four individuals. For each subsample, we estimated a SAF file in the same

fashion as the heterozygosity estimates, additionally limited to sites for which three of the

four subsampled individuals had data (`-minInd 3`). We then used realSFS to generate a

folded SFS and thetaStat to estimate these three diversity metrics from the SAF file and

SFS in 10kb sliding windows with a step size of 5kb, excluding windows with less than 1000

usable sites. The per site estimate of these three metrics was obtained for each window by

dividing the window estimate by the number of usable sites (`nsites`). We report the median

estimate across subsamples as the population estimate of each metric. To statistically

assess differences between historical and modern populations, we performed a paired

difference Wilcoxon signed-rank test on each of the corresponding historical and modern

subsamples, pairing estimates by genomic window. We estimated the effect size using the

following equation:

531
$$\frac{Z}{\sqrt{N_{win}}}$$

where Z is the Z-score from the Wilcoxon signed-rank test and N_{win} is the number of paired

windows. We considered modern populations to have significantly different estimates of

nucleotide diversity, Watterson's θ, or Tajima's D if the distribution of effect sizes across

all 100 subsample tests did not overlap 0.

Genetic Differentiation

We estimated genetic differentiation using pairwise population F_{ST} using realSFS. We

generated population SAF files using the same parameters as for the genetic diversity

metrics and generated a two dimensional folded SFS for each population pair using

realSFS. We then used this SFS and the population SAF files to estimate a global estimate

of F_{ST} using the Bhatia-Hudson estimator (`-whichFst 1`), which is less sensitive to small or

uneven sample sizes (111). Where reported, F_{ST} estimates refer to the weighted F_{ST}. To

statistically assess differences in F_{ST} between modern and historical populations, we

performed two tests: I) a paired *t*-test pairing corresponding population pairs in historical

and modern population sets and II) an unequal variances unpaired *t*-test incorporating all

possible population pairs in each set, comparing the historical to the modern population

set.

Relatedness

As the analyses we utilize to assess population structure can be sensitive to the presence

of close relatives in the dataset, we estimated pairwise relatedness between all samples

within each species dataset. We utilized the IBSrelate SFS-based method (112), as

implemented in NGSrelate (113), to estimate R0, R1, and KING-robust kinship coefficients

(114) for all pairs of individuals in the dataset. For input, we generated a Beagle file using

ANGSD for each focal species dataset (`-doGlf 2`), utilizing only positions where greater

than 80% of the individuals had data, calling single nucleotide polymorphisms (SNPs) (-

`SNP_pval 1e-6`) and assuming known major and minor alleles inferred from the genotype

likelihoods (`-doMajorMinor 1 -doMaf 1`) (115, 116).

Population Structure

We explored population structure using principal component analysis (PCA) as

implemented in PCAngsd v1.10 (117). As the population structure methods we use

assume independence of sites, we first estimated linkage disequilibrium for SNPs less

than 50kb apart in the species dataset Beagle files using ngsLD v1.2.0 (118). We pruned

SNPs using the recommended prune_graph tool (https://github.com/fgvieira/prune_graph;

v0.3.2), assuming independence of SNPs >50kb apart and with a coefficient of correlation

(r^2) < 0.1 and subset the Beagle file to include only pruned sites. We then removed one

individual from each pair of third-degree or closer relatives in the dataset (expected KING-

kinship coefficient >= 0.0625; Table S14) before running the analysis.

We formally tested for the presence of population structure using an admixture analysis for

1-10 ancestral clusters (K) using the pruned, relative removed Beagle file in NGSadmix v33

(119). We performed at least 20 replicate runs for each value of K, up to 100 total runs,

stopping if the three highest likelihood replicates converged within 2 log-likelihood units of

each other. The 'best-fit' value for K was then determined to be the highest K for which

replicates met this convergence criteria, following Pečnerová et al (108). We further

assessed the quality of the model fit for each value of K using evalAdmix v0.961 (120).

Genotype Calling

To estimate runs of homozygosity and deleterious burden in our three focal species, we

utilized methods that require genotype calls. As deleterious burden estimates can be

biased by variation in sequencing depth (19), we subsampled all samples to a mean 5X

filtered sequencing depth using Samtools, excluding samples with a mean filtered

sequencing depth below 5X from genotype call based analyses. We called genotypes per

sample using BCFtools multi-allelic caller v1.19 (100). We called genotypes only for

regions in the species-specific filtered sites files, using reads with a minimum mapping

quality of 30 and bases with a minimum quality of 20. We set genotypes with a read depth

<4 to missing and heterozygous genotypes where one allele was supported by <20% of the

reads to missing. We then merged both historical and modern samples into single species-

specific BCFs, removing SNPs within 5bp of indels, indels, and including only biallelic

SNPs which segregated in the combined historical and modern sample dataset. Finally, we

only included sites with called genotypes in 80% of samples and excluded all transition

variants.

Runs of Homozygosity

We assessed rates of recent and background inbreeding by estimating the inbreeding

coefficient F_{RoH}, the proportion of the autosomal genome existing in runs of homozygosity

over a specified length, for each individual from the genotype calls. We incorporated runs

of homozygosity >100kb into our estimates of F_{RoH}. We estimated runs of homozygosity

using BCFtools/RoH v1.19 (121), setting a constant species-specific recombination rate (-

M; estimated assuming 50 centimorgans per autosomal chromosome as in Mackintosh et

al (26), estimates in Table S12) and running the analysis on genotype calls only, with the PL

of unseen genotypes set to 30 (`-G30`). As we did not have sufficient samples to confidently

estimate population allele frequencies, we used a default minor allele frequency of 0.4 (`--`

`AF-dflt 0.4`). We additionally ignored homozygous reference positions (`--ignore-`

`homref`), as including these can inflate estimates of F_{RoH} when using a default allele

frequency based on our testing (Figure S9). We calculated F_{RoH} for each individual by

dividing the total sum of runs of homozygosity greater than 100kb in length by the total

length of the autosomes.

Deleterious mutation burden

We estimated individual levels of deleterious mutation burden from the genotype calls

using two complementary methods, one targeting conserved regions of the genome (122)

and the other utilizing gene annotations to infer functional effects (123). For these

analyses, we utilized the species-specific BCFs, additionally filtering to include only

positions with a genotyping rate of 80% in both the historical and the modern sample

subsets. As variants were called across the combined historical and modern sample set

for each species, alleles that are fixed within one time category, but variable or fixed for the

opposing allele in the other, are included.

To estimate to which extent the focal species had accumulated putatively deleterious

variants in conserved regions of the genome, we used the GenErode pipeline v0.6.0 (124)

to estimate Genomic Evolutionary Rate Profiling (GERP) scores (85) across each of the

three focal species genomes. The method implemented in this pipeline does not require a

multi-species alignment, instead fragmenting the outgroup genomes and mapping the

fragments to the target to generate alignments which are then used to estimate scores. We

used a dataset of 106 Lepidoptera chromosome-level outgroup genomes (Table S15) to

estimate the scores and ancestral states, using divergence times estimated with TimeTree

5 (125). We selected the top 1% of GERP scores as the most conserved positions using the

`get_gerp_score_percentile.py` script included with GenErode. For each individual, we

estimated deleterious burden at conserved positions using a GERP-scaled relative count

of derived alleles at the top conserved positions, considering the derived allele to be

putatively deleterious. Total deleterious burden at conserved positions for each individual

was calculated by summing the multiple of the derived allele count and GERP score at

each conserved position, scaled per 1000 derived alleles. We estimated homozygous

deleterious burden by calculating the same relative count, but only including homozygous

derived genotypes.

For the functional annotation based method, we used the genome annotations provided by

the Darwin Tree of Life Project (https://projects.ensembl.org/darwin-tree-of-life/). We

assigned impacts to each variant using Ensembl's Variant Effect Predictor (VEP) v112.0

(123). VEP categorizes variants into four classes: 'High', 'Moderate', 'Low', or 'Modifier',

which largely correspond to loss of function, missense, synonymous, and intergenic

variants. As variants often overlap multiple transcripts, multiple impacts will be assigned,

so we required VEP to pick one effect per variant (`--flag_pick`), using the default selection

order (123). For each individual, we estimated deleterious burden from strongly (including

variants classified as 'High' impact, e.g. loss of function) and weakly (including variants

classified as 'Moderate' impact, i.e. missense) deleterious variants using a relative count

of alternate alleles at these positions, which we treated as the putatively deleterious

alleles. Total deleterious burden for each individual is the count of alternate alleles for the

variant category of interest per 1000 alternate alleles. Homozygous deleterious burden is

the count of alternate alleles at homozygous positions for the variant category of interest

per 1000 alternate alleles.

Categorizing rates of genetic decline

To assess the changes in genetic diversity in a framework developed for conservation

focused genetic monitoring, we used the indicator thresholds suggested by Andersson et

al (20), which are based upon indicators applied for national use in Sweden (126). Using

these thresholds, genetic decline within populations is measured as the amount of genetic

diversity retained over a period of 100 years, assuming a constant rate of change. Three

threat categories are suggested: retention >=95% is considered 'Acceptable', 75-94% as

'Warning', and <75% as 'Alarm'. As different population genetic estimates of genetic

diversity are influenced by different components of genetic diversity, such as rare and

common variants, this metric can be assessed for each estimate separately.

We estimated the predicted retention of genetic diversity over 100 years for each

population using individual heterozygosity, nucleotide diversity, and Watterson's θ.

Genetic diversity retention per year was estimated using the following equation:

$$e^{\frac{ln(H_{mod}/H_{hist})}{Y_{mod}-Y_{hist}}}$$

where H is the estimate of genetic diversity for a given sample/population and Y is the

sampling year for that sample/population. As different populations had different sample

sizes, we used repeated subsamples to generate estimates of decline. For individual

heterozygosity, we performed all pairwise comparisons of historical and modern individual

estimates of heterozygosity within corresponding historical-modern sample pairs,

generating a single estimate for species with only one specimen per time period and a

range for ones with more (see Figure 4). For nucleotide diversity and Watterson's θ, we

utilized the previously generated 100 subsamples (N=4) for each population, estimating

H_{mod}/H_{hist} for paired historical and modern populations for each subsample, utilizing the

median value of H_{mod}/H_{hist} across subsamples to calculate the predicted retention.

Acknowledgements

We would like to thank Melina Eberhagen, Amanda Görnerup, Kajsa Svensson, Aske

Brydegaard Runemark, and Mikkel Brydegaard for assistance with collecting butterflies,

Håkan Elmqvist for providing pinned specimens of *Ph. alcon* and *S. orion*, Göran Engqvist

for providing tissue from the pinned specimen of *Pl. argyrognomon*, Emma Kärrnäs and

Isolde van Riemsdijk for assistance with DNA extractions, Jadranka Rota for assistance

with the entomological collections of the Biological Museum, Lund University

(https://doi.org/10.15468/dahk2a), Verena Kutschera and the Swedish Bioinformatics

Advisory Program for advice regarding the bioinformatic analyses, Anna Penna for

comments on the manuscript, and Isolde van Riemsdijk, Rachel Steward, Kalle Tunström,

Simon Jacobsen Ellestrand, Colin Olito, and Magne Friberg for helpful discussions

regarding the study species, methods and interpretations. We acknowledge support from

the National Genomics Infrastructure in Genomics Production Stockholm funded by

Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish

Research Council. The computations were enabled by resources provided by the National

Academic Infrastructure for Supercomputing in Sweden (NAISS) and the Swedish National

Infrastructure for Computing (SNIC) at UPPMAX partially funded by the Swedish Research

Council through grant agreements no. 2022-06725 and no. 2018-05973. This work was

supported by grants from the Strategic Research Area "Biodiversity and Ecosystem

services in a Changing Climate" (BECC), funded by the Swedish government, the

Foundation Oscar och Lili Lamms Minne, the Jörgen Lindström's Foundation, and the

Swedish Research Council for Sustainable Development (grant 2021-01163).

Data Availability Statement

693   The genetic data that support these findings will be made available upon publication at

694   NCBI in BioProjects PRJNA1068054 (focal species, modern) & PRJNA1168917 (focal

695   species, historical; additional species). The main population genomic pipeline is deposited

696   at https://doi.org/10.5281/zenodo.13785563. Workflow configurations, extensions, and

697   resources specific to this study are archived at https://doi.org/10.5281/zenodo.13902816.

References

1.	D. L. Wagner, E. M. Grames, M. L. Forister, M. R. Berenbaum, D. Stopak, Insect decline in the anthropocene: Death by a thousand cuts. *Proceedings of the National Academy of Sciences* **118**, e2023989118 (2021).

2.	R. van Klink, *et al.*, Meta-analysis reveals declines in terrestrial but increases in freshwater insect abundances. *Science* **368**, 417–420 (2020).

3.	R. Dirzo, *et al.*, Defaunation in the Anthropocene. *Science* **345**, 401–406 (2014).

4.	C. A. Halsch, *et al.*, Insects and recent climate change. *Proceedings of the National Academy of Sciences* **118**, e2002543117 (2021).

5.	P. H. Raven, D. L. Wagner, Agricultural intensification and climate change are rapidly decreasing insect biodiversity. *Proceedings of the National Academy of Sciences* **118**, e2002548117 (2021).

6.	S. Seibold, *et al.*, Arthropod decline in grasslands and forests is associated with landscape-level drivers. *Nature* **574**, 671–674 (2019).

7.	D. H. Boyes, D. M. Evans, R. Fox, M. S. Parsons, M. J. O. Pocock, Is light pollution driving moth population declines? A review of causal mechanisms across the life cycle. *Insect Conservation and Diversity* **14**, 167–187 (2021).

8.	J. R. Shipley, *et al.*, Agricultural practices and biodiversity: Conservation policies for semi-natural grasslands in europe. *Current Biology* **34**, R753–R761 (2024).

9.	R. C. Lacy, Loss of Genetic Diversity from Managed Populations: Interacting Effects of Drift, Mutation, Immigration, Selection, and Population Subdivision. *Conservation Biology* **1**, 143–158 (1987).

10.	R. Lande, Mutation and Conservation. *Conservation Biology* **9**, 782–791 (1995).

11.	R. Lande, S. Shannon, The Role of Genetic Variation in Adaptation and Population Persistence in a Changing Environment. *Evolution* **50**, 434–437 (1996).

12.	G. Bertorelle, *et al.*, Genetic load: genomic estimates and applications in non-model animals. *Nature Reviews Genetics* **23**, 492–503 (2022).

13.	M. Lynch, J. Conery, R. Burger, Mutation accumulation and the extinction of small populations. *The American Naturalist* **146**, 489–518 (1995).

14.	J. F. Crow, Some possibilities for measuring selection intensities in man. *Human Biology* **30**, 1–13 (1958).

15.	M. Lynch, W. Gabriel, Mutation Load and the Survival of Small Populations. *Evolution* **44**, 1725–1737 (1990).

16. D. L. Gilroy, K. P. Phillips, D. S. Richardson, C. van Oosterhout, Toll-like receptor variation in the bottlenecked population of the Seychelles warbler: computer simulations see the 'ghost of selection past' and quantify the 'drift debt'. *Journal of Evolutionary Biology* **30**, 1276–1287 (2017).

17. A. V. Pinto, B. Hansson, I. Patramanis, H. E. Morales, C. van Oosterhout, The impact of habitat loss and population fragmentation on genomic erosion. *Conservation Genetics* **25**, 49–57 (2024).

18. H. A. Jackson, *et al.*, Genomic erosion in a demographically recovered bird species during conservation rescue. *Conservation Biology* **36**, e13918 (2022).

19. M. Dehasque, *et al.*, Temporal dynamics of woolly mammoth genome erosion prior to extinction. *Cell* **187**, 3531–3540.e13 (2024).

20. A. Andersson, S. Karlsson, N. Ryman, L. Laikre, Monitoring genetic diversity with new indicators applied to an alpine freshwater top predator. *Molecular Ecology* **31**, 6422–6439 (2022).

21. D. O'Brien, *et al.*, Bringing together approaches to reporting on within species genetic diversity. *Journal of Applied Ecology* **59**, 2227–2233 (2022).

22. K. Theissinger, *et al.*, How genomics can help biodiversity conservation. *Trends in Genetics* **39**, 545–559 (2023).

23. D. Díez-del-Molino, F. Sánchez-Barreiro, I. Barnes, M. T. P. Gilbert, L. Dalén, Quantifying temporal genomic erosion in endangered species. *Trends in Ecology & Evolution* **33**, 176–185 (2018).

24. H. Ellegren, N. Galtier, Determinants of genetic diversity. *Nature Reviews Genetics* **17**, 422–433 (2016).

25. J. Romiguier, *et al.*, Comparative population genomics in animals uncovers the determinants of genetic diversity. *Nature* **515**, 261–263 (2014).

26. A. Mackintosh, *et al.*, The determinants of genetic diversity in butterflies. *Nature Communications* **10**, 3466 (2019).

27. G. Hewitt, The genetic legacy of the Quaternary ice ages. *Nature* **405**, 907–913 (2000).

28. J. A. Robinson, *et al.*, Genomic flatlining in the endangered island fox. *Current Biology* **26**, 1183–1189 (2016).

29. J. A. Robinson, *et al.*, The critically endangered vaquita is not doomed to extinction by inbreeding depression. *Science* **376**, 635–639 (2022).

30. M. F. DiLeo, A. Nair, M. Kardos, A. Husby, M. Saastamoinen, Demography and
environment modulate the effects of genetic diversity on extinction risk in a butterfly
metapopulation. *Proceedings of the National Academy of Sciences* **121**, e2309455121
(2024).

31. J. C. Teixeira, C. D. Huber, The inflated significance of neutral genetic diversity in
conservation genetics. *Proceedings of the National Academy of Sciences* **118**,
e2015096118 (2021).

32. A. García-Dorado, A. Caballero, Neutral genetic diversity as a useful tool for
conservation biology. *Conservation Genetics* **22**, 541–545 (2021).

33. M. Kardos, H. R. Taylor, H. Ellegren, G. Luikart, F. W. Allendorf, Genomics advances
the study of inbreeding depression in the wild. *Evolutionary Applications* **9**, 1205–1218
(2016).

34. M. K. Schwartz, G. Luikart, R. S. Waples, Genetic monitoring as a promising tool for
conservation and management. *Trends in Ecology & Evolution* **22**, 25–33 (2007).

35. S. Hoban, *et al.*, Effective population size remains a suitable, pragmatic indicator of
genetic diversity for all species, including forest trees. *Biological Conservation* **253**, 108906
(2021).

36. M. Kardos, *et al.*, The crucial role of genome-wide genetic variation in conservation.
Proceedings of the National Academy of Sciences **118**, e2104642118 (2021).

37. N. Dussex, H. E. Morales, C. Grossen, L. Dalén, C. van Oosterhout, Purging and
accumulation of genetic load in conservation. *Trends in Ecology & Evolution* **38**, 961–969
(2023).

38. C. van Oosterhout, Mutation load is the spectre of species conservation. *Nature
Ecology & Evolution* **4**, 1004–1006 (2020).

39. T. van der Valk, D. Díez-del-Molino, T. Marques-Bonet, K. Guschanski, L. Dalén,
Historical Genomes Reveal the Genomic Consequences of Recent Population Decline in
Eastern Gorillas. *Current Biology* **29**, 165–170.e6 (2019).

40. J. von Seth, *et al.*, Genomic insights into the conservation status of the world's last
remaining Sumatran rhinoceros populations. *Nature Communications* **12**, 2393 (2021).

41. G. Femerling, *et al.*, Genetic Load and Adaptive Potential of a Recovered Avian
Species that Narrowly Avoided Extinction. *Molecular Biology and Evolution* **40**, msad256
(2023).

42. M. Ihse, Swedish agricultural landscapes patterns and changes during the last 50
years, studied by aerial photos. *Landscape and Urban Planning* **31**, 21–37 (1995).

43.   R. Lindborg, O. Eriksson, Historical Landscape Connectivity Affects Present Plant
Species Diversity. *Ecology* **85**, 1840–1845 (2004).

44.   D. L. Wagner, R. Fox, D. M. Salcido, L. A. Dyer, A window to the world of global
insect declines: Moth biodiversity trends are complex and heterogeneous. *Proceedings of*
the National Academy of Sciences **118**, e2002549117 (2021).

45.   M. S. Warren, *et al.*, The decline of butterflies in europe: Problems, significance, and
possible solutions. *Proceedings of the National Academy of Sciences* **118**, e2002551117
(2021).

46.   J. C. Habel, *et al.*, Butterfly community shifts over two centuries. *Conservation*
Biology **30**, 754–762 (2016).

47.   SLU Artdatabanken, *Rödlistade arter i Sverige 2020* (SLU, 2020).

48.   C. Fägerström, Lund University Biological Museum - Insect collections Inventory.
(2024). https://doi.org/10.15468/dahk2a.

49.   Z. J. Nolen, M. Rundlöf, A. Runemark, Species-specific erosion of genetic diversity in
grassland butterflies depends on landscape land cover. *Biological Conservation* **296**,
110694 (2024).

50.   M. T. P. Gilbert, *et al.*, Recharacterization of ancient DNA miscoding lesions:
Insights in the era of sequencing-by-synthesis. *Nucleic Acids Research* **35**, 1–10 (2007).

51.   M. Stiller, *et al.*, Patterns of nucleotide misincorporations during enzymatic
amplification and direct large-scale sequencing of ancient DNA. *Proceedings of the*
National Academy of Sciences **103**, 13578–13584 (2006).

52.   H. Jónsson, A. Ginolhac, M. Schubert, P. L. F. Johnson, L. Orlando, mapDamage2.0:
Fast approximate bayesian estimates of ancient DNA damage parameters. *Bioinformatics*
29, 1682–1684 (2013).

53.   P. W. Hedrick, *Genetics of Populations* (Jones & Bartlett Publishers, 2009).

54.   R. McQuillan, *et al.*, Runs of homozygosity in european populations. *The American*
Journal of Human Genetics **83**, 359–372 (2008).

55.   E. A. Thompson, Identity by descent: Variation in meiosis, across genomes, and in
populations. *Genetics* **194**, 301–326 (2013).

56.   S. Glémin, How are deleterious mutations purged? Drift versus nonrandom mating.
Evolution **57**, 2678–2687 (2003).

57.   M. Kirkpatrick, P. Jarne, The effects of a bottleneck on inbreeding depression and
the genetic load. *The American Naturalist* **155**, 154–167 (2000).

58. J. Wang, W. G. Hill, D. Charlesworth, B. Charlesworth, Dynamics of inbreeding
depression due to deleterious mutations in small populations: mutation parameters and
inbreeding rate. *Genetics Research* **74**, 165–178 (1999).

59. B. Charlesworth, Mutational load, inbreeding depression and heterosis in
subdivided populations. *Molecular Ecology* **27**, 4991–5003 (2018).

60. D. Kleinman-Ruiz, *et al.*, Purging of deleterious burden in the endangered iberian
lynx. *Proceedings of the National Academy of Sciences* **119**, e2110614119 (2022).

61. E. Humble, *et al.*, Conservation management strategy impacts inbreeding and
mutation load in scimitar-horned oryx. *Proceedings of the National Academy of Sciences*
120, e2210756120 (2023).

62. SLU Artdatabanken, Artfakta: kronärtsblåvinge (Plebejus argyrognomon).

63. J. C. Habel, M. J. Samways, T. Schmitt, Mitigating the precipitous decline of
terrestrial European insects: Requirements for a new strategy. *Biodiversity and*
Conservation **28**, 1343–1360 (2019).

64. E. L. Landguth, *et al.*, Quantifying the lag time to detect barriers in landscape
genetics. *Molecular Ecology* **19**, 4179–4191 (2010).

65. C. W. Epps, N. Keyghobadi, Landscape genetics in a changing world: disentangling
historical and contemporary influences and inferring change. *Molecular Ecology* **24**, 6021–
6040 (2015).

66. W. H. Lowe, F. W. Allendorf, What can genetics tell us about population
connectivity? *Molecular Ecology* **19**, 3038–3051 (2010).

67. P. W. Hedrick, J. R. Adams, J. A. Vucetich, Reevaluating and Broadening the
Definition of Genetic Rescue. *Conservation Biology* **25**, 1069–1070 (2011).

68. P. W. Hedrick, A. Garcia-Dorado, Understanding inbreeding depression, purging,
and genetic rescue. *Trends in Ecology & Evolution* **31**, 940–952 (2016).

69. D. A. Bell, *et al.*, The Exciting Potential and Remaining Uncertainties of Genetic
Rescue. *Trends in Ecology & Evolution* **34**, 1070–1079 (2019).

70. R. Lande, Risk of population extinction from fixation of new deleterious mutations.
Evolution **48**, 1460–1469 (1994).

71. Y. B. Simons, G. Sella, The impact of recent population history on the deleterious
mutation load in humans and close evolutionary relatives. *Current Opinion in Genetics &*
Development **41**, 150–158 (2016).

72. Y. Xue, *et al.*, Mountain gorilla genomes reveal the impact of long-term population
decline and inbreeding. *Science* **348**, 242–245 (2015).

73.   C. Grossen, F. Guillaume, L. F. Keller, D. Croll, Purging of highly deleterious
mutations through severe bottlenecks in Alpine ibex. *Nature Communications* **11**, 1001
(2020).

74.   N. Dussex, *et al.*, Population genomics of the critically endangered kākāpō. *Cell*
Genomics **1**, 100002 (2021).

75.   A. Khan, *et al.*, Genomic evidence for inbreeding depression and purging of
deleterious genetic variation in indian tigers. *Proceedings of the National Academy of*
Sciences **118**, e2023018118 (2021).

76.   B. Charlesworth, Evidence against Fisher's theory of dominance. *Nature* **278**, 848–
849 (1979).

77.   F. Manna, G. Martin, T. Lenormand, Fitness landscapes: An alternative theory for
the dominance of mutation. *Genetics* **189**, 923–937 (2011).

78.   A. García-Dorado, A. Caballero, On the average coefficient of dominance of
deleterious spontaneous mutations. *Genetics* **155**, 1991–2001 (2000).

79.   C. D. Huber, A. Durvasula, A. M. Hancock, K. E. Lohmueller, Gene expression drives
the evolution of dominance. *Nature Communications* **9**, 2750 (2018).

80.   B. Charlesworth, D. Charlesworth, *Elements of evolutionary genetics* (Roberts;
Company, 2010).

81.   L. F. Keller, D. M. Waller, Inbreeding effects in wild populations. *Trends in Ecology &*
Evolution **17**, 230–241 (2002).

82.   D. Charlesworth, J. H. Willis, The genetics of inbreeding depression. *Nature Reviews*
Genetics **10**, 783–796 (2009).

83.   A. L. K. Mattila, *et al.*, High genetic load in an old isolated butterfly population.
Proceedings of the National Academy of Sciences **109**, E2496–E2505 (2012).

84.   I. Saccheri, *et al.*, Inbreeding and extinction in a butterfly metapopulation. *Nature*
392, 491–494 (1998).

85.   G. M. Cooper, *et al.*, Distribution and intensity of constraint in mammalian genomic
sequence. *Genome Research* **15**, 901–913 (2005).

86.   C. D. Huber, B. Y. Kim, K. E. Lohmueller, Population genetic models of GERP scores
suggest pervasive turnover of constrained sites across mammalian evolution. *PLOS*
Genetics **16**, e1008827 (2020).

87.   C. C. Kyriazis, J. A. Robinson, K. E. Lohmueller, Using computational simulations to
model deleterious variation and genetic load in natural populations. *The American*
Naturalist **202**, 737–752 (2023).

88.   C. Bortoluzzi, *et al.*, Lepidoptera genomics based on 88 chromosomal reference
sequences informs population genetic parameters for conservation. *bioRxiv* (2023).
https://doi.org/10.1101/2023.04.14.536868.

89.   L. B. Pettersson, H. Arnberg, *Svensk dagfjärilsövervakning, årsrapport för 2023*
(Biologiska institutionen, Lunds universitet, 2024).

90.   S. Hoban, *et al.*, Genetic diversity targets and indicators in the CBD post-2020
Global Biodiversity Framework must be improved. *Biological Conservation* **248**, 108654
(2020).

91.   L. Laikre, *et al.*, Post-2020 goals overlook genetic diversity. *Science* **367**, 1083–1085
(2020).

92.   G. Formenti, *et al.*, The era of reference genomes in conservation genomics. *Trends*
in Ecology & Evolution **37**, 197–202 (2022).

93.   V. G. Twort, J. Minet, C. W. Wheat, N. Wahlberg, Museomics of a rare taxon: placing
Whalleyanidae in the Lepidoptera Tree of Life. *Systematic Entomology* **46**, 926–937 (2021).

94.   Z. J. Nolen, Zjnolen/PopGLen: V0.3.0. (2024).
https://doi.org/10.5281/zenodo.13785563. Deposited September 2024.

95.   F. Mölder, *et al.*, Sustainable data analysis with snakemake [version 2; peer review:
2 approved]. *F1000Research* **10** (2021).

96.   S. Chen, Y. Zhou, Y. Chen, J. Gu, Fastp: An ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

97.   H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. (2013). https://doi.org/10.48550/arXiv.1303.3997.

98.   H. Li, R. Durbin, Fast and accurate long-read alignment with BurrowsWheeler
transform. *Bioinformatics* **26**, 589–595 (2010).

99.   E. Palkopoulou, *et al.*, Complete genomes reveal signatures of demographic and
genetic declines in the woolly mammoth. *Current biology: CB* **25**, 1395–1400 (2015).

100.   P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *GigaScience* **10** (2021).

101.   Picard toolkit. *Broad Institute, GitHub Repository* (2019). Deposited 2019.

102.   A. Peltzer, *et al.*, EAGER: efficient ancient genome reconstruction. *Genome Biology*
17, 1–14 (2016).

103.   G. A. van der Auwera, B. D. O'Connor, *Genomics in the Cloud: Using Docker, GATK,*
and WDL in Terra (O'Reilly Media, Inc., 2020).

104.   T. S. Korneliussen, A. Albrechtsen, R. Nielsen, ANGSD: Analysis of Next Generation
Sequencing Data. *BMC Bioinformatics* **15**, 1–13 (2014).

105.   R. N. Lou, A. Jacobs, A. P. Wilder, N. O. Therkildsen, A beginner's guide to low-
coverage whole genome sequencing for population genomics. *Molecular Ecology* **30**,
5966–5993 (2021).

106.   G. Jun, M. K. Wing, G. R. Abecasis, H. M. Kang, An efficient and scalable analysis
framework for variant extraction and refinement from population-scale DNA sequence
data. *Genome Research* **25**, 918–925 (2015).

107.   J. Neukamm, A. Peltzer, K. Nieselt, DamageProfiler: Fast damage pattern
calculation for ancient DNA. *Bioinformatics* **37**, 3652–3653 (2021).

108.   P. Pečnerová, *et al.*, High genetic diversity and low differentiation reflect the
ecological versatility of the African leopard. *Current Biology* **31**, 1862–1871.e5 (2021).

109.   A. F. A. Smit, R. Hubley, P. Green, RepeatMasker open-4.0. (2013). Deposited 2013.

110.   S. Gopalakrishnan, *et al.*, The population genomic legacy of the second plague
pandemic. *Current Biology* **32**, 4743–4751.e6 (2022).

111.   G. Bhatia, N. Patterson, S. Sankararaman, A. L. Price, Estimating and interpreting
FST: The impact of rare variants. *Genome Research* **23**, 1514–1521 (2013).

112.   R. K. Waples, A. Albrechtsen, I. Moltke, Allele frequency-free inference of close
familial relationships from genotypes or low-depth sequencing data. *Molecular Ecology* **28**,
35–48 (2019).

113.   K. Hanghøj, I. Moltke, P. A. Andersen, A. Manica, T. S. Korneliussen, Fast and
accurate relatedness estimation from high-throughput sequencing data in the presence of
inbreeding. *GigaScience* **8**, giz034 (2019).

114.   A. Manichaikul, *et al.*, Robust relationship inference in genome-wide association
studies. *Bioinformatics* **26**, 2867–2873 (2010).

115.   L. Skotte, T. S. Korneliussen, A. Albrechtsen, Association testing for next-generation
sequencing data using score statistics. *Genetic Epidemiology* **36**, 430437 (2012).

116.   S. Y. Kim, *et al.*, Estimation of allele frequency and association mapping using next-
generation sequencing data. *BMC Bioinformatics* **12**, 231 (2011).

117.   J. Meisner, A. Albrechtsen, Inferring population structure and admixture proportions
in low-depth NGS data. *Genetics* **210**, 719–731 (2018).

118.   E. A. Fox, A. E. Wright, M. Fumagalli, F. G. Vieira, ngsLD: Evaluating linkage
disequilibrium using genotype likelihoods. *Bioinformatics* **35**, 3855–3856 (2019).

119.   L. Skotte, T. S. Korneliussen, A. Albrechtsen, Estimating individual admixture
proportions from next generation sequencing data. *Genetics* **195**, 693–702 (2013).

120.   G. Garcia-Erill, A. Albrechtsen, Evaluation of model fit of inferred admixture
proportions. *Molecular Ecology Resources* **20**, 936–949 (2020).

121.   V. Narasimhan, *et al.*, BCFtools/RoH: A hidden markov model approach for
detecting autozygosity from next-generation sequencing data. *Bioinformatics* **32**, 1749–
1751 (2016).

122.   E. V. Davydov, *et al.*, Identifying a High Fraction of the Human Genome to be under
Selective Constraint Using GERP++. *PLOS Computational Biology* **6**, e1001025 (2010).

123.   W. McLaren, *et al.*, The Ensembl Variant Effect Predictor. *Genome Biology* **17**, 122
(2016).

124.   V. E. Kutschera, *et al.*, GenErode: a bioinformatics pipeline to investigate genome
erosion in endangered and extinct species. *BMC Bioinformatics* **23**, 1–17 (2022).

125.   S. Kumar, *et al.*, TimeTree 5: An expanded resource for species divergence times.
Molecular Biology and Evolution **39**, msac174 (2022).

126.   K. Johannesson, L. Laikre, Monitoring of genetic diversity in environmental
monitoring [in Swedish]. *Report to the Swedish Agency for Marine and Water Management*
(dnr. HaV 3642-2018, 3643-2018) (2020).

*Figure 1: **Sampling design and changes in genetic diversity, differentiation, and inbreeding in the focal species.** (A) Images of the three focal species. (B) Map of the sampling area in southern Sweden, sampling locations are approximate. Circles represent areas with modern samples in at least one of the species and triangles represent areas with historical samples, see Table S1 for complete sampling information. (C) Change in individual heterozygosity over time, measured as heterozygous sites per 1000bp, excluding transitions. Each point represents an individual sample, grouped into historical and modern samples per species. Coloring refers to sampling region and corresponds to the map in panel B. (D) Change in inbreeding coefficients over time, estimated using runs of homozygosity larger than 100kb (F_{RoH}). (E) Change in pairwise population genetic differentiation, F_{ST} over time between equivalent comparisons in each time period. Color of the points and line refer to the geographic distance between populations. Populations with no connecting line refer to pairs of modern sample populations that have no comparable historic pair. In C & D, results of unequal variance t-tests between historical and modern samples for each species is shown using the following scale, where panels without * are not significant (P > 0.05): * P <= 0.05, ** P <= 0.01, *** P <= 0.001, **** P <= 0.0001. Statistical assessment of genetic differentiation is not shown, as it could not be assessed for all three species (Table S7).*

Figure 2: **Population structure of the focal species.** *(A) Principal Component Analysis (PCA) of the three focal species. Shapes refer to sampling period, with each circle representing a modern and each triangle a historical individual, with colors corresponding to the sampling locations in Figure 1B. Percentages on each axis refer to the proportion of total variance explained by the component. (B) Best fit admixture analyses for the three focal species. Each bar represents an individual and each color one of K genetic clusters. Each bar is colored in proportion to the probability of the individual belonging to each cluster.*

*Figure 3: **Temporal changes in deleterious burden in the focal species** (A) Individual inbreeding coefficients (F_{RoH}) partitioned by RoH size class. Longer RoH indicate more recent common ancestry of the identical haplotypes. (B) Estimates of total deleterious mutation burden for each individual of the three focal species across three categories of putatively deleterious mutations: I) missense (i.e. weakly deleterious; upper), II) loss of function (i.e. strongly deleterious; middle), and III) derived mutations in highly conserved regions (lower). Total burden for missense and LoF mutations is the total count of alternate alleles for the given mutation category per 1000 alternate alleles. Total burden for derived mutations at conserved positions is the total count of derived alleles at the top 1% most genomically conserved positions multiplied by their GERP scores, per 1000 derived alleles.*

C) Homozygous deleterious mutation burden for the same mutation categories as in B, but only calculated using positions homozygous for the putatively deleterious alleles. In B & C, results of unequal variance t-tests comparing deleterious burden between historical and modern samples are shown in each panel, where panels without * are not significant (P > 0.05): * P <= 0.05, ** P <= 0.01, *** P <= 0.001, **** P <= 0.0001.

Figure 4: Changes in heterozygosity predicted over 100 years based on the rate of change estimated from modern and historical samples for each of 11 species. Species are grouped by conservation status, which follows the species name on the x-axis. Background coloration refers to the three genetic decline thresholds outlined by Andersson et al (20), where a retention of >=95% over 100 year is considered 'acceptable' (green), a retention of 75-94% as 'warning' (yellow), and a retention of <75% as 'alarm' (red). Points represent mean estimates for change in heterozygosity across all sample pairs for a species. Species with bars have more than one possible sample pair, with error bars illustrating the spread of estimates across all sample pairs. Species with a * next to their species name on the x-axis have been mapped to a related species' reference genome as they lack a species-specific reference (Table S12).