1   **Modular cloning of multigene vectors for the baculovirus system and yeast**

2   **Zhihao Lai[1], Sarena F. Flanigan[1], Marion Boudes[1,3], Chen Davidovich[1,2,3]**

[1]Department of Biochemistry and Molecular Biology, Biomedicine Discovery Institute, Faculty

of Medicine, Nursing and Health Sciences, Monash University, Clayton, Victoria, Australia.

[2]EMBL-Australia, Clayton, Victoria, Australia

[3]Corresponding authors: Marion Boudes (marion.boudes@monash.edu) and Chen

Davidovich (chen.davidovich@monash.edu)

Abstract

Recombinant macromolecular complexes are often produced by the baculovirus system,

using multigene expression vectors. Yet, the construction of baculovirus-compatible

multigene expression vectors is complicated and time-consuming. Furthermore, while the

baculovirus and yeast are popular protein expression systems, no single method for

multigene vector construction is compatible with both. Here we present the modular cloning

(MoClo) Baculo toolkit for constructing multigene expression vectors for the baculovirus

system and, through compatibility with the MoClo Yeast toolkit, also for yeast. Vector

construction by MoClo Baculo does not require PCR, primers or the sequencing of

intermediate products. As a proof of principle, MoClo Baculo was used to construct

baculovirus and yeast multigene vectors expressing the four- and five-subunit human

Polycomb Repressive Complex 2. We show that MoClo Baculo simplifies and expedites the

construction of multigene expression vectors for the baculovirus system and provides

compatibility with yeast as an alternative expression system.

Introduction

An increasing number of large multi-subunit macromolecular complexes are subjected to

structure determination using cryo-electron microscopy [1]. Such structural and biophysical

studies often require the expression and purification of recombinant multisubunit protein

complexes. When the macromolecule complex of interest is of a eukaryotic origin, its

subunits are often required to be co-expressed in a eukaryotic host [2,3]. Among the most

commonly used eukaryotic systems for recombinant protein expression are the baculovirus

system [4] and yeast [5], each offering different advantages.

There are several widely used baculovirus expression systems. The Bac-to-Bac system is

one of the most commonly used systems for the expression of protein complexes with a few

protein subunits [6]. The Bac-to-Bac system allows the cloning of up to two genes per

expression vector, which is then integrated into the baculovirus DNA and packed into

baculovirus particles [6]. Complexes with more than two protein subunits can be expressed

by the Bac-to-Bac system. This is commonly done by the co-infection of insect cells using

multiple different baculoviruses, each carrying a different gene. Accordingly, there are many

examples of protein complexes that were expressed using co-infection with 3-5

baculoviruses (e.g. [7,8,9,10,11]). Yet, beyond that, the efficiency and reproducibility of co-

infections is reduced and, therefore, the expression of complexes with more than a few

subunits commonly requires the usage of multi-gene vectors.

There are several multigene vector systems for protein expression using the baculovirus

system. MultiBac recombines multiple genes into one vector using the Cre-lox technology

[12]. BiGBac enables the construction of similar multigenic vectors using a Gibson Assembly

[13], allowing a maximum of 25 genes in one plasmid [14]. GoldenBac utilises Golden Gate

assembly that simplifies vector construction, where individual genes are PCR amplified only

once, at the beginning of the process [15].

All current multigene baculovirus expression systems [6,11,12,14,15] require PCR or Gibson

Assembly [13], and therefore oligonucleotide synthesis. PCR-based approaches may

complicate the usage of GC-rich or repetitive templates for the construction of some

multigene vectors. Furthermore, most multigene baculovirus vectors do not have a protein

tag pre-incorporated in the vector. Instead, large protein tags are often added through a prior

cloning step while small tags are included in the cloning primers. These factors increase the

reliance on oligonucleotide synthesis and the subsequent sequencing of intermediate

products, which increase the time, cost and complexity of the project.

The problem of multigene vector construction has already been solved in various organisms

using modular cloning (MoClo) [16]. In MoClo, different 'parts' are functional DNA sequence

elements that are assembled together to form 'devices', which are more complex plasmids.

Each 'part' is a plasmid that includes one functional element, such as a protein-coding

sequence, tag, promoter, or backbone of an expression vector. Each part is generated only

once, usually with the aid of gene synthesis or gBlocks, and can then be used repeatedly in

different projects or shared between labs. Multiple plasmids, each holding a single part, are

digested together by type IIS restriction enzymes to release DNA fragments with cohesive

ends that allow multiple parts to assemble together through Golden Gate assembly [17]. This

process can be carried out iteratively, where plasmids are ligated, digested and ligated again

to form increasingly complex plasmids. MoClo-constructed plasmids may include multiple

genes and all the elements that are required for their delivery, integration into the host,

selection, protein expression and protein purification. MoClo is fast, efficient, affordable, and

does not require PCR, Gibson Assembly or custom primer design.

The MoClo Yeast Toolkit [18] was designed for modular cloning of multigene expression

cassettes, aiming for synthetic biology applications in the yeast *S. cerevisiae*. Although the

MoClo Yeast Toolkit was not designed primarily for the expression and purification of

recombinant proteins, we reasoned that it would be feasible to modify it for that purpose.

Based on a cloning standard inspired by the MoClo Yeast Toolkit, we set out to develop a

method for cross-platform multi-gene vector building for protein expression and purification.

We focused on constructing parts for modular cloning of expression vectors dedicated to the

baculovirus system and yeast, given their broad applicability.

Here we present the MoClo Baculo, a multi-gene modular cloning (MoClo) method designed

for the expression of proteins using the baculovirus system and in yeast. MoClo Baculo

utilises Golden Gate as the only cloning method. The mutation rate is extremely low, given

that MoClo does not involve the usage of PCR, Gibson Assembly, in vitro polymerases or

primers. Therefore, the sequencing of intermediate product plasmids is not required, which

dramatically reduces the time, cost and complexity of building multigene constructs. Modular

cloning allows for flexible replacement or removal of N- or C-terminal affinity tags when

developing a new purification scheme. MoClo Baculo supports the expression of up to 6

proteins from a single multigene construct, however this number can theoretically be

expanded by adding adapters. The number of co-expressed subunits can further be

expanded through the delivery of multiple multigene constructs. As a proof of concept, we

used MoClo Baculo to build multi-gene expression vectors of the 4- and 5-subunits

Polycomb repressive complex 2 (PRC2). We demonstrate the suitability of such MoClo

Baculo expression vectors to express catalytically active PRC2 complexes using the

baculovirus system and yeast.

Results

Modification of the MoClo Yeast toolkit for better compatibility with applications of recombinant protein expression and purification.

We set out to generate a system that would utilise the same set of open reading frames (ORFs) and tag constructs for building different multi-protein expression vectors. Furthermore, we wished to have compatibility with protein expression using the baculovirus system and yeast. We prioritised methods that do not rely on PCR, in vitro polymerase reactions or primers, aiming to avoid the sequencing of intermediate products. We reasoned that the MoClo Yeast Toolkit [18], originally developed for synthetic biology applications, would be an ideal starting point.

The MoClo Baculo toolkit design is inherited from the MoClo Yeast Toolkit and therefore includes the following part types that are assembled together to construct expression vectors (Supplementary Fig. 1 and Supplementary Table 1): Left Connector (ConL; part 1) that connects functional elements to the plasmid backbone; Promoter, designed for protein expression with (part 2a) or without (part 2) N-terminal tag; an optional N-terminal tag (part 2b); a protein Coding Sequence (CDS; part 3); an optional C-terminal tag (part 4a); a transcription Terminator sequence that is designed to be used with or without C-terminal tag (part 4b or 4, respectively); a right Connector (ConR; part 5) and a backbone (part 6) that can be divided into up to 4 different parts (parts 6a-d) with different functionalities (more below).

One important difference between the MoClo Yeast Toolkit and the MoClo Baculo designs is that in the latter one can use the same CDS part with or without tags(Supplementary Fig. 1 and Supplementary Table 1). Specifically, in the MoClo Yeast toolkit, the construction of vectors for expressing untagged and N-terminally tagged proteins is done using two different CDS parts (Supplementary Fig. 1). This design is undesirable for structural biologists, who would benefit from the frequent removal or inclusion of N-terminal tags without having to source two separate CDS parts. Therefore, to allow for removing the N-terminal tag without changing the CDS part, we modified the MoCo Yeast general design of promoters and N-terminal parts (Supplementary Fig. 1A).

To enable the modular cloning using Golden Gate assembly, each part is domesticated into a chloramphenicol resistance plasmid (pYTK001) that originated from the MoClo Yeast toolkit (Supplementary Fig. 1D). The resulting plasmid holds a single part and is referred to as "Level 0" plasmid [18] (see methods section and Supplementary Fig. 1C,D for the flanking

sequences that are added to each part, to allow for its domestication and subsequent

usage).

The newly modified design of MoClo Baculo still allows the replacement of promoters

through modular cloning, although it is not fully compatible with the promoter design of the

MoClo Yeast toolkit [18]. Therefore, we generated several useful MoClo Baculo-compatible

S. cerevisiae promoters, based on promoter sequences from the MoClo Yeast toolkit

(Supplementary Table 1). More *S. cerevisiae* promoters can be made as needed. We also

designed and generated parts carrying N- and C-terminal tags that are useful for protein

expression and purification. Such tags for protein purification include a PreScission (3C)

protease cleavable maltose binding protein (MBP) tag, a dual protein A tag and a Strep-tag

(Supplementary Table 1). Additional tags can readily be added as individual parts, without

bringing the coding sequence into consideration and as long as the correct flanking

sequences are added to them (Supplementary Fig. 1C,D).

Adding compatibility with the baculovirus system for protein expression

We next sought to enable compatibility between the MoClo Yeast toolkit to the baculovirus

system. To this end, we modified parts of an expression vector used in the Bac-to-Bac

expression system (pFastBac1 [6]), aiming to enable modular cloning. For that, we first

aligned backbone sequences from different existing baculovirus expression systems that

were previously modified from the Bac-to-Bac systems. This was done to identify commonly

used modules (Supplementary Fig. 1B) and regions where the pFastBac1 vector can be

modified without negatively impacting performances. Based on this sequence analysis we

designed two new parts that originated from the pFastBac1 vector and are termed here BV

'Backbone' (part 6b-d in Supplementary Table 1 and plasmid pMB.001 in Supplementary

Table 2) and 'Recombination into BV DNA' (part 6a in Supplementary Table 1 and Fig. 1A,

top right, and plasmid pF003 in Supplementary Table 2). The BV Backbone (part 6b-d;

plasmid pMB.001) includes bacterial replication origin, ampicillin and gentamicin resistance

and the right transposition site (Tn7R), which is used for recombination into the baculovirus

DNA. The left transposition site (Tn7L) is included within the 'Recombination into BV DNA'

part (part 6a; plasmid pF003). Together, the 'Backbone' part and the 'Recombination into BV

DNA' part include all the functional elements required to connect to each other and to all the

other parts that are collectively required to express a single open reading frame, either

tagged or not. We envision that the usage of two parts for the backbone would facilitate

further backbone modifications in future works. For compatibility with the MoClo Yeast

cloning standard, we used codon optimization to remove BsaI and BsmBI sites from the

sequences that were used to construct all these parts. In the case of parts that are specific

for yeast, or may be used for both yeast and baculovirus expression, we also removed NotI

sites, as NotI is used to linearise vectors for yeast genome integration. If one wishes to use

the optional compatibility between MoClo Baculo to the biGBac system (more below), then

PmeI sites should be removed too.

For compatibility with the baculovirus expression system, we designed additional new parts

using the MoClo cloning standard: polyhedrin promoter (parts 2 and 2a) and SV40

terminator (parts 4 and 4b). All these parts were domesticated into Level 0 plasmids and are

compatible with Golden Gate assembly using the type IIS restriction enzyme BsaI. At this

point, the system included all the elements required to assemble expression vectors that are

compatible with the Bac-to-Bac system. Each of the resulting expression vectors are termed

"Level 1" plasmids (Fig. 1, top and Supplementary Fig. 1A). Level 1 plasmids can express a

single protein from an element that is termed a "Transcription Unit": a functional element

required to transcribe an mRNA that is coding for the expression of a desired protein.

To enable the assembly of multiple Transcription Units (from multiple Level 1 plasmids) into

a single multi-gene baculovirus expression vector (termed "Level 2" plasmid), we required a

new backbone that confers additional antibiotic resistance. For that, we added

spectinomycin resistance into the 'Recombination into BV DNA' part (part 6a). The new part

6a ('Recombination into BV DNA + SpecR') can now be assembled together with the

Backbone (part 6b-d), a Spacer (part 2-4) and Connectors (parts 1 and 5) through a Golden

Gate reaction using BsaI. The result is a Level 1 plasmid that is termed here 'Level 2

Backbone'. The Level 2 Backbone plasmid does not carry a Transcription Unit but, instead,

contributes the backbone during the assembly of Level 2 plasmids. Users can assemble

their own 'Level 2 Backbone' plasmids using parts from the MoClo Baculo toolkit, together

with Connector parts from the MoClo Yeast toolkit. However, for the ease of use, we

included in the MoClo Baculo toolkit a Level 2 Backbone plasmid (pMB.002) that includes

the ConLS' and ConRE' connectors. The pMB.002 plasmid allows the integration of up to 6

transcription units into a Level 2 plasmid (Fig. 1B, right, and Supplementary Table 2).

Overall, this design ensures that both single-gene (Level 1) and multi-gene (Level 2)

expression vectors are suitable for protein expression using the Bac-to-Bac system and,

therefore, confer resistance to ampicillin and gentamicin. In addition to these, multi-gene

(Level 2) baculovirus expression vectors also confer resistance to spectinomycin.

Adding compatibility with the biGBac system

We envisioned that some users may benefit from compatibility between MoClo Baculo and

the biGBac system [14]. This is because the biGBac system allows to connect together 5

multi-gene vectors using Gibson Assembly. Therefore, we decided to generate MoClo
Baculo parts that would include the Gibson Assembly linkers of the biGBac system. For that,
we modified the pMB.002 plasmid by inserting linker sequences ("A-F") from pBIG1a-e [14]
and PemI restriction enzyme sites (Supplementary figure 4A). We named the resulting
plasmids pMB.BIG1a to pMB.BIG1e. These plasmids are functionally identical to the biGBac
system plasmids pBIG1a to pBIG1e plasmids, respectively. The compatibility between
MoClo Baculo to the biGBac system may allow the optional connection of up to five Level 2
MoClo Baculo plasmids, each with up to six transcription units. The resulting biGBac
expression vectors are referred to as Level 3 plasmids here and contain up to 30
transcription units.

We believe that most users will not need the compatibility with the biGBac system, and
therefore it is marked here as optional. This is because compatibility with biGBac is
beneficial only if one wishes to increase the maximal number of transcriptional units from six
(Level 2 plasmids) to 30 (optional Level 3; Supplementary Fig. 4B) without adding new
MoClo Baculo adapters. However, the added compatibility between the MoClo Baculo to the
biGBac system may allow regular users of the biGBac system to construct their expression
vectors with the aid of Modular Cloning. This may provide biGBac users with the simplicity of
removing or replacing tags without PCR or gibson assembly.

As a proof of principle, we assembled a baculovirus vector aiming for the expression of an
eight subunit complex (Supplementary Fig. 5). This was done first with the MoClo Baculo
system, using Golden Gate assembly, to construct the three Level 2 vectors (Supplementary
Fig. 5B): pMB.BIG1a.PRC2 (expresses four PRC2 subunits), pMB.BIG1b.MTF2-PALI1 (for
the expression of two accessory subunits of PRC2: MTF2 and PALI1), and pMB.BIG1c.G9A-
GLP (for the expression of the H3K9me3 methyltransferase G9A and GLP). The resulted
three Level 2 vectors are compatible with the biGBac system, which was used to combine
them using Gibson Assembly into one Level 3 vector of 32 kb (pBIG2abc.PRC2.1-EHMT),
aiming for the co-expression of all eight subunits (Supplementary Fig. 5C). The efficiency of
constructing the Level 3 vector was 33% (2 out of 6 clones were correct; Supplementary Fig.
5A). The produced Level 3 vectors were verified using whole plasmid sequencing, with the
exception of sporadic mutations in the homology regions used for Gibson Assembly.

Modular cloning of baculovirus expression vectors

From this point, modular cloning is carried out using the same workflow that has been
described for the MoClo Yeast toolkit [18]. As the process is described below in brief, it is
highly recommended to read the description of modular cloning by Weber et al [16] and Lee

et al [18]. Briefly, all the Level 0 (chloramphenicol resistant) vectors that carry the parts for a
desired Transcription Unit are combined in a single pot Golden Gate assembly, using the
type IIS restriction enzyme BsaI (Fig 1a). The produced Level 1 plasmid has lost its
chloramphenicol resistance and now confers resistance to ampicillin in *E. coli*. The Level 1
baculovirus expression vectors also confer resistance to gentamicin in *E. coli*. This transition
between antibiotic selection markers is inherited from the modular cloning method and
allows selection against precursor plasmids while progressing up to the next plasmid Level
through iterative cloning. For the production of baculovirus expression vectors, we use
chloramphenicol to select Level 0 plasmids, ampicillin or gentamicin to select Level 1
plasmids and spectinomycin to select Level 2 plasmids. Each Level 1 plasmid can now be
used to express a single protein, depending on the transcription unit and the backbone.
Multiple Level 1 plasmids can be used to assemble one Level 2 plasmid—a multi-gene
expression vector—using the Golden Gate assembly region with BsmBI digestion (Fig. 1B,
bottom). Yet, the assembly of Level 2 vectors requires some planning, because the order of
the transcription units in a Level 2 vector is defined based on the Connectors (parts 1 and 5)
that are included in each of the Level 1 vectors that are combined to assemble a multi-gene
expression vector (Level 2).

Lee et al. [18] and Weber et al. [16] are highly recommended readings on the topic of
connector design, as the MoClo Baculo toolkit uses the connector parts of the MoClo Yeast
toolkit [18]. In brief, connectors are designed such that a right Connector (ConR, part 5) of a
given Transcription Unit can be connected only to a complementary left Connector (ConL,
part 1) of another Transcription unit. Exceptions are the ConLS' and ConRE' connectors that
are reserved for the backbone. For instance, ConR1 is connected to ConL1 of another
Transcription Unit, ConR2 is connected to ConL2 in another Transcription unit, and so forth
(see Supplementary Table 1 for the list of connectors). Practically, this means that the order
of the transcription units in a given Level 2 vector usually need to be defined at the time
when one designs the relevant set of Level 1 vectors that are intended to be used together in
a subsequent Golden Gate assembly.

Even when Level 2 expression vector is intended for the baculovirus system, we find it
convenient to use the same connector parts that were designed by Lee et al for the MoClo
Yeast toolkit [18]. The MoClo Yeast toolkit provides connectors for combining up to 6
transcription units together. If less than 6 subunits are assembled, a linker plasmid can be
used to connect the last transcription unit to the backbone (an example for that is described
in the next section). It is anticipated that more connectors can be added to increase the

number of transcription units in a given multi-gene expression vector, as long as the total
length of the construct permits Golden Gate assembly.

We find that the sequencing of Level 1 plasmids is not required if they are only used as
intermediate products for assembling a Level 2 vector. This is because mutations are
uncommon, since no PCR or oligonucleotides are used. Therefore, when Level 1 plasmids
are only used as intermediate products, we commonly validate the correct Level 1 plasmid
assembly by restriction digestion before proceeding to Level 2 assembly. At the end of the
process, the correct assembly of the Level 2 plasmid is validated by restriction digestion
followed by whole plasmid sequencing. The obtained Level 2 plasmids are multi-gene
expression vectors compatible with the bac-to-bac baculovirus expression system (Fig. 1C).

Modular cloning of yeast expression vectors

The construction of yeast expression vectors is carried out using the same CDS and tag
parts that are used for baculovirus vector building. The process of vector building for yeast
expression is nearly identical to the construction of baculovirus vectors, as described above.
An exception is that for the construction of yeast expression vectors we normally use
chloramphenicol to select Level 0 plasmids, ampicillin to select Level 1 plasmids and
kanamycin to select Level 2 plasmids, as described by Lee et al [18]. Most of the parts that
are used for building Level 1 and 2 plasmids for yeast expression are taken from the MoClo
Yeast toolkit [18] (Fig. 1A; boxes with black lines). Exceptions are the following parts that
were generated for the MoClo Baculo toolkit (Fig. 1A; boxes with a blue line or dashed blue
line): yeast promoter parts dedicated for using together with N-terminal tags according to the
MoClo Baculo cloning standard and the same N-terminal tag parts, CDS parts and C-
terminal tag parts that are used for baculovirus expression.

Proof of principle: multi-gene vector construction

As MoClo Baculo adapted the Modular Cloning strategy of the MoClo Yeast toolkit [18], the
efficiency of cloning Level 0 and Level 1 vectors is close to 100 %. As a demonstration of
Level 1 cloning, we constructed six different Level 1 plasmids (Fig. 2A). These included a
backbone plasmid for baculovirus expression (pMB.002) and five Transcription Unites for the
expression of each of the five subunits of the PRC2-AEBP2 complex (Fig. 2A,B). These five
Transcription Units designed for the expression of the PRC2 protein subunits MBP-EZH2,
MBP-SUZ12, MBP-EED MBP-RBBP4 and AEBP2, where MBP is a PreScission-cleavable
maltose binding protein tag. After the Golden Gate assembly reaction, two clones were

picked per each of these Level 1 plasmids. All these clones were then assayed using

restriction digestion, confirming a success rate of 100% (12/12; Fig 2A).

As with other systems of Modular Cloning [16,18], the efficiency of assembling Level 2

vectors may be reduced with the size and the number of parts. Here we demonstrate the

construction of two Level 2 vectors, for the baculovirus expression of the PRC2 core

complex in the presence or absence of its accessory subunit AEBP2 (Fig. 2C,D).

Specifically, a 19 kbp PRC2 expression vector was designed for the expression of a 4-

protein complex (MBP-EZH2, MBP-SUZ12, MBP-EED and MBP-RBBP4) and assembled

with 80 % efficiency (4/5 colonies; Fig. 2C). A 21 kbp PRC2-AEBP2 expression vector

designed to express five protein subunits (MBP-EZH2, MBP-SUZ12, MBP-EED, MBP-

RBBP4 and AEBP2) and was assembled with an efficiency of 50% (2/4 colonies; Fig. 2C).

Each of the Golden Gate assemblies that were carried out to generate the expression

vectors for the 4- and 5-subunit PRC2 complexes included a linker plasmid (L6 and L56 in

Fig 2B). Such a linker plasmid is required when less than 6 Transcription Units are

incorporated and aims to connect the last Transcription Unit to the backbone. The linker

plasmid is assembled in a Golden Gate reaction using BsaI, by combining an ampicillin

resistant *E. coli* Backbone plasmid (plasmid pYTK095 from the MoClo Yeast Toolkit), a

Spacer (parts 2-4, plasmid pYTK048 from the MoClo Yeast toolkit) and the appropriate

Connector parts (parts 1 and 5; Supplementary Table 1) that are required to connect the last

Transcription Unit to the Level 2 Backbone plasmid (pMB.002; Supplementary Table 2).

The sizes of the two Level 2 plasmids that we generated—19 kbp and 21 kbp—approach the

upper size of plasmids that can be efficiently assembled using Golden Gate assembly

(commonly reported as 20-30 kbp [16,18,19]). The expression of all subunits is driven by the

same polyhedrin promoter and SV40 terminator, similar to biGBac [14] and Multibac systems

[12]. Selected Level 2 clones were subjected to whole plasmid sequencing before

downstream protein expression and purification.

Proof of principle: protein expression and purification

As proof of principle, we used the PRC2- and PRC2-AEBP2 multigene baculovirus

expression vectors that were constructed above (pMB.PRC2 and pMB.PRC2.AEBP2,

respectively; Fig. 2D) for protein expression and purification. The obtained macromolecular

complexes were pure, assembled and monodispersed (Fig. 3A-D,G). We obtained about 4

mg and 1.5 mg of PRC2 and PRC2-AEBP2, respectively, per litre of Hi5 insect cell culture.

We also demonstrated PRC2 protein expression from the biGBac-compatible vector

pMB.BIG1a.PRC2 (Fig. 3H), which was assembled solely by Golden Gate assembly using

MoClo Baculo (Supplementary Figure. 5B).

As a proof-of-principle of protein expression using the yeast *S. cerevisiae*, we generated an

expression vector for PRC2 (pMY.PRC2; Supplementary Fig. 2). For vector building, we

used the same CDS (part 3) and Connectors (parts 1 and 5) parts that were used to build

the baculovirus expression vectors above. To allow for yeast expression, we replaced the

promoters (part 2/2a) and terminators (part 4) with *S. cerevisiae* promoters and terminators.

For the same reason, we replaced the parts responsible for integration into the host DNA

(parts 4a-d) with parts that were designed for integration into *S. cerevisiae* chromosome in

yeast (Fig. 1A and Supplementary Fig. 2). We also replaced the MBP tags by a dual Protein

A tag (Protein A) that we find more suitable in the yeast system, possibly given the lower

expression level of human PRC2 in S. *cerevisiae* (more below). We then integrated the

multi-gene PRC2 expression vector into *S. cerevisiae* chromosome through homologous

recombination and used the obtained strain for the expression of the human PRC2. The

obtained PRC2 was then purified as an assembled complex (Fig. 3E,F).

PRC2 is a histone methyltransferase enzyme. Therefore, we wished to determine that the

recombinant PRC2 protein complex that we obtained is enzymatically active *in vitro*. To this

end, we carried out an *in vitro* histone methyltransferase assay, to compare the enzymatic

activities of the recombinant 4-subunits human PRC2 complexes that we purified using three

different approaches: (i) Baculovirus expression, through the co-infection of four baculovirus

stocks that were made using standard pFastBac1 vectors, each designed to express a

single PRC2 subunit (Fig. 4, "pFB1 PRC2"), including the protein subunits MBP-EZH2, MBP-

SUZ12, MBP-RBBP4 and MBP-EED. (ii) Baculovirus expression using a single MoClo

Baculo-derived multi-gene vector carries Transcription Units for the same four PRC2

subunits with the same MBP tags (Fig. 4, "MoClo Baculo PRC2, insect cells"). (iii) A yeast

expression of PRC2 using a single multi-gene expression vector including four Transcription

Units of the same PRC2 subunits as above, with the exception that EZH2 carries a 2xProtein

A tag and the rest of the subunits are untagged (Fig. 4, "MoClo Baculo PRC2, yeast"). In all

cases, all tags were removed during the protein purification process and were not present in

the enzymatic assay.

All the PRC2 samples exhibited a substantial enzymatic activity only in the presence of an

allosteric effector peptide H3K27me3 (Fig. 4B, blue bars), in agreement with PRC2 being an

allosteric enzyme [20]. We also observed the automethylation of the catalytic subunit EZH2

(Supplementary Fig. 4A,B), which has to be methylated for PRC2 to become active [21,22].

We observed some variations in the enzymatic activity of the different recombinant human

PRC2 preparations. These variations in activity were rather small, within <2-fold of the

apparent reaction rate (Fig. 4B). These variations in activity might be attributed to the

different expression systems. We also cannot exclude slight variations in the quantification of

the protein concentration or subunit stoichiometry between samples, although SDS-PAGE

did not identify large differences between them (Fig 4C). Overall, we demonstrated the

construction of multi-gene expression vectors using MoClo Baculo and their subsequent

application of protein expression and purification using the baculovirus system and yeast.

Discussion

The MoClo Baculo system allows a high flexibility in building expression vectors while

altering tags and expression systems as needed. According to data herein (Table 1) and

elsewhere [23, 24], the baculovirus system provides a higher protein yield per litre than

yeast. On the other hand, from the point plasmids are constructed, protein expression using

the baculovirus system is a slow process (20-30 days under ideal conditions), while the

yeast expression system provides a faster progress (6-7 days; Table 1 and [24]). Another

advantage of the yeast expression system is the lower cost of the media: roughly 40-20% of

the per-litre cost compared to the baculovirus system (Table 1). We propose that one could

use MoClo Baculo in order to construct expression vectors for fast and cost-effective protein

expression trials in yeast. Subsequently, the best-performing set of ORFs and tags can be

assembled into baculovirus expression vectors, aiming for a high-yield protein production in

insect cells.

A benefit of MoClo Baculo, with respect to most other baculovirus systems (Table 2), is that

the entire vector construction process can be automated, allowing high throughput cloning.

Specifically, MoClo Baculo is PCR free, which reduces the likelihood of mutations and

eliminates the need for primer design and PCR optimization. Open reading frames can be

synthesised as plasmids or gene fragments (e.g. gBlocks), such that they can be directly

domesticated into Level 0 vectors by Golden Gate assembly. For the same reason, MoClo

Baculo would be beneficial also by reducing the time and training required to design and

construct complex expression vectors in any lab.

As for other MoClo cloning standards, MoClo Baculo would allow labs and institutes to

manufacture, store and share collections of ORFs and tags ready for their assembly into

multi-gene expression vectors for baculovirus and yeast. MoClo Baculo was designed as a

general cloning standard, with the end goal of expression and purification of multi-subunit
complexes for applications of structural biology, biophysics and biochemistry. As such,
MoClo Baculo could potentially be expanded to other protein expression systems, as done
with other MoClo tools for various applications [25,26,27,28,29]. Overall, MoClo Baculo is a
cost-effective, fast and robust system for constructing multi-gene expression vectors,
demonstrated here using the baculovirus system and yeast.

Acknowledgements

406 We thank the support of the Micromon Genomics facility. This work was supported by Sylvia
407 and Charles Viertel Senior Medical Research Fellowship (C.D.) and the National Health and
408 Medical Research Council (NHMRC) grant numbers APP1162921, APP1184637,
409 APP2011767 and APP2020900 (C.D.).

Competing interests

411 The authors declare no competing interests.

Author contributions

413 Z.L., M.B. and C.D. designed the project, Z.L., S.F.F. and M.B. carried out experiments while
414 Z.L. led aspects related to the baculovirus system and M.B. led aspects related to yeast, Z.L.
415 and M.B. analyzed data, Z.L., M.B. and C.D. wrote the manuscript and C.D. supervised the
416 project.

Methods

Plasmid sequences

Part plasmids (Level 0) and expression vectors (Level 1 and 2) that were generated in this
study are listed in Supplementary Table 2 and their annotated sequences are provided in
GenBank file format in Supplementary Data 1. Plasmid sequence annotation was carried out
with the aid of pLannotate [30]. Detailed information about plasmid construction is provided
below. Even if one wishes to use MoClo Baculo exclusively for protein expression using the
baculovirus system they would still need some plasmids from the MoClo Yeast toolkit, such
as the Level 0 backbone (plasmid pYTK001), Connector part plasmids, Spacer part (plasmid
pYTK048) and an *E. coli* backbone (plasmid pYTK095). The MoClo Yeast toolkit plasmids
are available on AddGene (Kit #1000000061) and their sequences are provided in [18].

Design of coding sequences (CDS) for MoClo Baculo

All CDS sequences were designed for compatibility with the MoClo Yeast system [18]. This includes the addition of overhangs that are reserved for part 3 (CDS; Supplementary Fig. 2C). Flanking BsmBI and BsaI sites were added to the CDS sequence, designed to generate the desired overhands for Golden Gate assembly into Level 0 and Level 1 plasmids, respectively (See Supplementary Fig. 1D).

In the MoClo Baclo system, part 3 allows for protein expression either with or without an N-terminal tag (Supplementary Fig. 2A), so only one CDS part is needed per project. As with the MoClo Yeast toolkit, the design of the CDS involved the introduction of synonymous mutations to remove the restriction sites for NotI, BsmBI, and BsaI from within the protein open reading frame (i.e., with the exception of the flanking BsmBI and BsaI sites). If one wishes to use MoClo Baculo to construct biGBac plasmids, they will also have to ensure that CDS and tag parts do not include PemI sites, although this would rarely be a problem as PemI sites are extremely uncommon. If one wishes only to construct expression vectors for the baculovirus system, not for yeast, then it is not required to remove NotI sites from CDS parts. This is because NotI is only used to linearise plasmids before their homologous recombination into a yeast chromosome. Finally, CDS are then domesticated into Level 0 plasmids, as described in the next section.

Generation of new part (Level 0) plasmids

This process does not need to be repeated in cases where part plasmids already exist. It would typically be carried out once when a new part (e.g. a new tag) is introduced for the first time. The appropriate part-specific flanking sequences (Supplementary Fig. 1C) and universal flanking sequences (Supplementary Fig. 1D) were added to part sequences, either through gene synthesis or PCR primers, to enable domestication into the Level 0 vector and the subsequent modular cloning. DNA with the sequence of the promoter, terminator and tag parts were obtained either as ampicillin-resistant plasmids from gene synthesis (GenScript), gene fragments (IDT gBlocks), or PCR amplicons (Phusion® High-Fidelity DNA Polymerase, NEB) that were generated from plasmids with a pFastBac1.HMBP backbone. The parts were then cloned using Golden Gate Assembly (BsmBI-v2, NEB #R0739) into the Level 0 backbone plasmid (pYTK001) of the MoClo yeast toolkit. The Backbone part (Part 6b-c for the baculovirus system; Supplementary Table 1; plasmid pMB.001) for Level 1 MoClo Baculo baculovirus expression plasmids, was synthesised by Genscript to remove the BsaI and BsmBI sites and insertion of GFP-dropout.

Generation of level 2 backbone plasmids

pMB.002 was generated by the golden gate assembly with BsaI enzyme using pMB.001 and pMB.004 from the MoClo Baculo toolkit, together with pYTK048, pYTK008, pYTK073 from yeast MoClo toolkit [18]. To construct the biGBac compatible backbones pMB.BIG1a to pMB.BIG1e (Supplementary Fig. 4A), the same reaction was performed using modified pYTK008 plasmids (ConLS'A to ConLS'E, respectively) and modified pYTK073 plasmids (ConLE'B to ConLE'F, respectively), where the biGBac compatible homology regions for Gibson Assembly (sequences A to F [14]; Supplementary Table 3) were inserted by BsmBI linearisation and HiFi DNA assembly (NEB, #E2621L) using ssDNA templates.

Golden Gate assembly using BsaI and BsmBI

For Golden Gate assembly using BsaI, 25 fmol of each plasmid was added per reaction and assembled with NEBridge® Ligase Master Mix (NEB #M1100) and BasI-HF-V2 (NEB #R3733) according to protocol provided by the manufacturer. For Golden Gate assembly using BsmBI, 25 fmol of each plasmid was added per reaction with 1 µLBsmBI-v2 (NEB #R0739), 1 µL HiT4 ligase (NEB #R2622), 1.5 µL 10x T4 ligase buffer (NEB #B0202), 1 µL 500 mM NaCl making up a 15 µL reaction. The mixture was cycled 60 times between 16 °C and 50 °C for 5 min each, followed by 30 min final ligation at 16 °C and digestion at 50 °C for 15 min. The reaction was held at 10 °C and stored at -20 °C until use.

XL1-Blue (Agilent) was used as the only bacterial strain for all the cloning. Transformation and culturing of XL1-Blue is carried out according to the manufacturer's protocol. For the propagation of pMB.001, the transformed cells were selected on an ampicillin plate. For the selection of Level 1 clones, we find that it is easier to identify GFP positive colonies (i.e. where Golden Gate assembly failed) on ampicillin plates.

Construction of Level 3 plasmids using the biGBac system (optional)

pBIG2 plasmids were linearised in PmeI (NEB #R0560) restriction digest reaction at a final concentration of 10 fmol/µL. All multigene-carrying pMB.BIG1a-e plasmids were miniprep (Qiagen #27106) with an extra PE buffer wash and eluted in 40 µL 60 °C Milli-Q water twice. Then, 500 fmol of each plasmids were mixed in a total of a 10 µL reaction of 1x Cutsmart (NEB) and with 1 µL PmeI (NEB #R0560). If plasmid concentration was not sufficient, all plasmids were mixed and dried in speedvac and re-dissolved in 9 µL 1x Cutsmart (NEB). All digestions were carried out at 37 °C for 90 minutes. Then, 1 µL of linearised backbone and 1 µL digested pMB.BIG1a-e mix were used to set up a 20 µL Gibson Assembly (NEB #E2611) reaction. The ligation mix were used to transform chemical-competent NEB10-beta (NEB

495 #C3019) or electro-competent XL1-Blue, depending on the size of the final product.

496 Generally, for plasmid between 20-30 kbp, electroporation is beneficial but not essential. For

497 plasmid >30 kbp, electroporation is required.

Expression and purification of PRC2 using the baculovirus system

Baculovirus was generated using Sf9 cells according to the instructions by the manufacturer

(ThermoFisher). The virus titer was determined by the viability of infected SF9 cells using the

MTS kit (Promega #G3580).

Protein expression was carried out according to the method previously described by Zhang

et al [11]. Specifically, High Five cells were grown in Insect-XPRESS (Lonza #12-730Q)

media and were infected with baculovirus at a cell density of 1.8-2.0*10^6 cells/ml. 70 hours

post-infection, cell pellets were harvested at 1500 g and lysed in lysis buffer (50 mM Tris

pH8 at 25 °C, 300 mM NaCl, 15 % glycerol, 0.5 % NP-40, 1 mM TCEP, cOmplete protease

inhibitor (11836170001)) for 30 min at 4 °C. The lysate was clarified by centrifugation at

30,000 g for 30 min and the supernatant was incubated with amylose resin (NEB #E8021)

for 60 min with rotation. Then, the resin was washed with 5 column volume of lysis buffer,

10-column volumes of HSW buffer (50 mM Tris pH8 at 25 °C, 500 mM NaCl) and 10 c.v. of

LSW buffer (50 mM Tris pH8 at 25 °C, 150 mM NaCl). The protein was eluted from the resin

with LSW buffer supplemented with 10 mM maltose and 1 mM TCEP. Then the protein was

digested overnight with PreScission protease and loaded to a 5 ml prepacked Heparin

column (Cytiva #17040701) and eluted with a linear gradient over 10 c.v., starting from

Buffer A (150 mM NaCl, 20 mM Tris pH 8.0 at 25 °C) to 50% Buffer B (2000 mM NaCl, 20

mM Tris pH 8.0 at 25°C). Finally, the eluate was diluted 1:1 with 20 mM Tris pH 8.0, 1 mM

TCEP, concentrated using 30 kDa Amicon Ultra centrifugal filter and injected to a Superose

6 Increase 10/300 column (Cytiva, #29091596) in HEPES pH7.5 at 25 °C and 150 mM NaCl.

Fractions that contain protein of interest were supplemented with 1 mM TCEP, concentrated

and frozen in liquid nitrogen. The entire purification was carried out at 4-8 °C.

Expression and purification of PRC2 using S. cerevisiae

For the purification of the 4-subunits PRC2 from yeast, cells were picked from a single

colony and grown overnight in -Ura synthetic minimal media (2 g/L Yeast Synthetic Drop-out

Medium Supplement without uracil (Merck #1501), 67 g/L Yeast Nitrogen Base Without

Amino Acids (Merck #Y0626)) supplemented with 3% Glucose. The following day, cells were

washed twice and used to inoculate 12 L of YP-lactate (1.5% Yeast extract (Merck #Y1625),

3% Meat Peptone (ThermoScientific H26694.36), 2% Lactate (Chem Supply #LL008)) in 6

flasks. Cultures were grown at 30 °C and 180 rpm for 31h. EZH2 expression was induced by

the addition of 3% galactose and the cultures were grown for 19h. Cells were harvested,

washed in water, pelleted (335 g), flash-frozen and stored at -80 °C until the protein

purification was carried out. For protein purification, the cell pellet was thawed in an equal

volume of 2X lysis buffer (400 mM NaCl, 50 mM Tris pH 7.5, 0.5 mM DTT, 17 mg/mL

Phenylmethanesulfonyl Fluoride (Merck #P7626), 25 µg/mL Leupeptin (Merck #2884), 125

µg/mL Pepstatin A (Merck #516481), 33 mg/mL Benzamidine (Merck #B6506)), and lysed

using a DynoMill. Lysate was clarified by centrifugation at 30,000 g, followed by

ultracentrifugation at 45,000 rpm. The supernatant was bound overnight to 20 mL of IgG

resin. Beads were washed with a high salt buffer (500 mM NaCl, 20 mM Tris pH 7.5, 0.5 mM

DTT), followed by a low salt buffer (200 mM NaCl, 20 mM Tris pH 7.5, 0.5 mM DTT). PRC2

complex was eluted using an overnight cleavage by TEV protease (100 µg) and subjected to

heparin purification. The heparin peak was concentrated and injected into a Superose 6

10/300 column (Cytiva, #29091596). The protein in the peak corresponding to PRC2 was

concentrated and flash-frozen.

SDS-PAGE

To resolve all subunits using SDS-PAGE, 3-8% Tris-Acetate gel (ThermoFisher

#EA0375BOX) and 1X MES SDS Running buffer (ThermoFisher #NP0002) were used. 3 µg

or 1 µg PRC2 complexes were supplemented to a final concentration of 1X LDS sample

buffer (ThermoFisher #NP0007) with 1% 2-Mercaptoethanol (Sigma-Aldrich #M3148) and

heated at 95 °C for 5 min before loading onto a Tris-Acetate gel. The gel was run for 60 min

at 160 V before staining with InstantBlue Coomassie protein stain (Expedeon, #ISB1L).

Nucleosome reconstitution

Histones, octamers, and polynucleosome reconstitution were carried out as previously

described [11]. In brief, recombinant histones were purified from inclusion bodies and

reconstituted into histone octamers. 182 bp DNA was amplified using Pfu DNA polymerase

and purified by ion exchange chromatography and isopropanol precipitation. Chromatin was

assembled using gradient salt dialysis at 4°C. Chromatin was reconstituted by initially

titrating across a range of octamer ratios (from 1:0.9 to 1:1.1 DNA:octamer molar ratio) in a

20 µL mixture of 3 µM DNA, 20 mM Tris pH 7.5 at 25 °C, 2 M NaCl, 1 mM EDTA and 1 mM

DTT. For each of these samples, gradient salt dialysis was used at 4°C in a dialysis device

(ThermoFisher #69572), starting from refolding buffer (20 mM Tris pH 7.5 at 25 °C, 2 M KCl,

1 mM EDTA and 1 mM DTT) to a medium salt buffer containing 20 mM Tris pH 7.5 at 25 °C,

250 mM KCl, 1 mM EDTA and 1 mM DTT over 18 h, and the final step dialysis was carried

out using a low salt buffer containing 20 mM Tris pH 7.5 at 25 °C, 2.5 mM KCl, 1 mM DTT.

The quality of mononucleosomes was assessed by 0.8 % agarose TBE and 6 %

polyacrylamide TBE gel electrophoresis, and the most appropriate molar ratio of

DNA:octamer was selected for a large-scale batch. Large-scale reconstitution was

conducted as above, except in a volume of 0.5 mL using dialysis tubing (Spectrum #888-

11527). The folded mononucleosome was concentrated with 500 µL 100 kDa MWCO

concentrator (Cytiva, #28932237). The concentration of the mononucleosome was

measured using a Qubit protein assay kit (ThermoFisher #Q3321). Mononucleosomes were

stored at 4 °C until use.

In vitro **HMTase activity assays using radiolabeled S-adenosyl-L-methionine**

For the HMTase reactions, each 10 µL reaction contained 0.6 µM PRC2 produced from

either yeast or insect cells, 0.6 µM chromatinised gene, and 5 µM S-[methyl-14C]-adenosyl-

L-methionine (PerkinElmer, #NEC363050UC). Reactions were incubated in reaction buffer

(50 mM Tris-HCl pH 8.0 at 30°C, 100 mM KCl, 2.5 mM $MgCl_2$, 0.1 mM $ZnCl_2$, 2 mM 2-

mercaptoethanol, 0.1 mg/mL BSA, and 5% v/v glycerol) for 30, 60, or 120 min at 30°C.

Reactions were stopped by adding 4X LDS loading dye (Thermo Fisher Scientific, #NP0007)

supplemented with 4% 2-Mercaptoethanol (Sigma-Aldrich #M3148) to a final concentration

of 1X LDS and heating at 95°C for 5 min. The reactions were then loaded onto 16.5 % SDS-

PAGE gels and ran on ice for 120 min at 160 V in 1X Tris-glycine buffer. Gels were stained

with InstantBlue Coomassie protein stain (Expedeon, #ISB1L) before vacuum-drying for 1 h

at 60°C. Dried gels were then exposed to a storage phosphor screen for six days before

acquiring radiograms using a Typhoon 5 Imager (GE Healthcare). All experiments were

performed in three independent replicates that were carried out on three separate days.

Densitometry was carried out using ImageLab software (Bio-Rad). Values were interpolated

from a standard curve using GraphPad Prism 10.2.1 and the background was subtracted,

then values plotted with SD displayed.

Figures and Tables

Figure 1. Design of the MoClo Baculo system. (A) Level 0 plasmids carrying individual
parts, each with a different function as indicated. Connectors: designed to connect the left
(L) and right (R) parts of elements from multiple Level 1 plasmids. Promoter: a promoter
compatible with the intended host organism. N' Tag: optional N-terminal tag. CDS: coding
sequence of the protein of interest. C-tag: optional C-terminal tag. Terminator: transcription
termination sequence. Promoter and terminator parts are marked with an asterisk if they
were designed to connect to an optional N-terminal or C-terminal tag, respectively. 3'Hom
and 5'Hom are the 3' and 5' homologous arms aiming for chromosomal integration in *S.
cerevisiae*. Origin is from an *S. cerevisiae* plasmid (e.g. 2 micron; in case the user does not
wish to integrate the expression cassette into a chromosome). Backbone: includes elements
critical for the propagation of the plasmid in *E. coli*, such as antibiotic resistance and *E. coli*
origin of replication. Parts specific to the baculovirus system (BV) or yeast are indicated in
blue and red font, respectively. Parts that were constructed for MoClo Baculo are

represented by boxes with a blue border. Parts that originated from the MoClo Yeast Toolkit

are represented by boxes of black border. Dashed blue boxes represent the ORF part, which

was only partially inherited from the MoClo Yeast toolkit (see Supplementary Fig. 1A), and

the C-terminal tag part that follows the same design as the MoClo Yeast toolkit but now

includes a broader selection of tags for protein purification. The ampersand (&) operator

represents an obligatory selection of parts that were designed to be assembled directly.

When parts or elements within brackets are stacked vertically, only one of them should be

selected. Bottom: For the assembly of Level 1 plasmids, parts are subjected to Golden Gate

assembly using the BsaI restriction enzyme. (B) Level 1 plasmids are transcription units,

each capable of expressing a single protein. An exception is where a simple backbone

(bottom right in (A)) is used for the assembly of the transcription unit, which leads to a

transcription unit that can be assembled into a functional Level 2 expression vector, although

the Level 1 plasmid is not designed to function as an expression vector by itself. Multiple

Level 1 vectors, including multiple transcription units and one backbone, are subjected to

Golden Gate assembly using the BsmBI enzyme to form one Level 2 vector. In some cases,

a 'Linker' plasmid can replace a transcription unit in order to link between another

transcription unit to the backbone (bottom in (B)). (C) Level 2 plasmids are multi-gene

expression vectors, designed to express a multi-subunit macromolecular complex. The

compatible expression system is determined based on the parts selected in (A). Yeast

expression vectors were designed to express proteins either after homologous

recombination into a chromosome or using a yeast plasmid. The order of the transcription

units is determined based on the connectors that were selected in (A).

Figure 2. Proof-of-principle: construction of multigene baculovirus expression vectors for the expression of 4- and 5-subunit PRC2 complexes. (A) Golden Gate assembly of Level 1 plasmids was carried out using BsaI and two of the resulting clones were proceeded

for restriction digestion using BsmBI. The plasmids pMB.M.E2, pMB.M.S12, pMB.M.ED,

pMB.M.R4 and pMB.AI3 are Level 1 baculovirus expression vectors for the expression of

EZH2, SUZ12, EED, RBBP4 and AEBP2 (isoform 3). The plasmid pMB.002 is a Level 1

plasmid that was assembled to contribute to the backbone during subsequent assemblies of

Level 2 plasmids. (B) Schematic representation of the plasmids resulted from the Golden

Gate assembly reaction that was analysed in (A). Presented are also two 'linker plasmids',

L6 and L56, that are Level 1 plasmids that include a Spacer and Connector parts from the

MoClo Yeast toolkit [18], designed to connect the last Transcription Unit to the backbone

during the assembly of a Level 2 plasmid. (C) Golden Gate assembly using the Level 1

plasmids from (B) was carried out using BsmBI to construct Level 2 plasmids for a

baculovirus expression of the 4-subunit PRC2 (left) or the 5-subunit PRC2-AEBP2 (right).

Left: pMB.M.E2, pMB.M.S12, pMB.M.ED, pMB.M.R4, pMB.002 and L56 were assembled to

form an expression vector coding for PRC2. In this context, the linker L56 was designed to

connect the 4th transcription unit to the backbone. Right: PRC2-AEBP2 expression vector

was assembled from the same Level 1 plasmids, with the exception that the Level 1 plasmid

pMB.AI3 was added to include also the expression of AEBP2 and the L56 linker plasmid was

replaced by the L6 linker plasmid, to connect the 5th transcription unit to the backbone. Four

clones were selected from each Golden Gate reaction and were subjected to restriction

digestion using MfeI and SalI. (D) Schematics of the obtained Level 2 multigene expression

vectors, as indicated. In (B) and (D), the colour key is as in Fig. 1. PH: polyhedrin promoter;

SV40: SV40 polyA terminator; MBP: maltose-binding protein with PreScission cleavage site;

Spacer: DNA sequence that was designed to fill up a gap during vector building (originated

from the MoClo Yeast toolkit [18]); Backbone: vector backbone; BV rec and BV rec + SecR:

Tn7 recombination sites compatible with the Bac-to-Bac baculovirus expression system, in

the presence and absence of Spectinomycin resistance, respectively. The Level 1 plasmids

presented in (B) confer Ampicillin and Gentamicin resistance in *E. coli*. The Level 2 plasmids

presented in (D) confer Ampicillin, Gentamicin and Spectinomycin in *E. coli*. In (A) and (B),

fragments were analysed on 0.7% agarose TAE gel.

Figure 3. MoClo Baculo-expressed 4- and 5-subunit PRC2 complexes are pure and assembled. (A) PRC2 was expressed from baculovirus (BV) using the pMB.PRC2 multi-gene expression vector (Fig. 2D, left) and was fractionated over a Superose 6 Increase 10/300 column. UV absorbance at 280 nm was plotted to trace protein elution. The void volume is marked as V_0. The black horizontal line above the peak indicates fractions that were taken for SDS-PAGE analysis in (B). (B) PRC2 gel filtration fractions from (A) were analysed by SDS-PAGE (Tris-Glycine, 12% acrylamide) and stained by coomassie. (C) PRC2-AEBP2 expressed using pMB.PRC2.AEBP2 construct (Fig. 2D, right) was analysed by Superose 6 Increase 10/300 column. (D) Fractions marked by the black line in (C) were analysed by SDS-PAGE (Tris-Glycine, 12% acrylamide). (E) A PRC2 yeast expression construct (pMY.PRC2; see supplementary Fig. 2 for vector construction) was integrated into the genome of *S. cerevisiae* using homologous recombination for protein expression and then PRC2 that was purified and fractionated over a Superose 6 Increase 10/300 column. (F) Fractions that are marked by the horizontal black line in (E) were analysed on an SDS-PAGE (Tris-Glycine, 12% acrylamide). "Inp" indicates a sample before the size exclusion chromatography. (G) Purified PRC2 and PRC2-AEBP2 from (A) and (C), respectively, were analysed on SDS-PAGE (Bis-Tris, 4%-12% acrylamide) after fractions were pooled and proteins were concentrated. (H) Amylose affinity purification of PRC2 from lysate of SF9

cells that were infected with baculovirus carry the pMB.BIG1a.PRC2 vector (Supplementary

Fig. 5B), analysed on SDS-PAGE (Bis-Tris, 8%-16% acrylamide).

Figure 4. MoClo Baculo-expression vectors designed either for the baculovirus

system or for yeast produced recombinant human PRC2 complexes that are

enzymatically active. A. HMTase assay performed with 600 nM PRC2, 600 nM

mononucleosomes, 5 μM S-Adenosyl-L-methionine [methyl-^{14}C] and in the presence or

absence of 50 μM H3K27me3 peptide, as indicated. The reaction mixture was separated

with SDS-PAGE, stained with coomassie and exposed to a phosphor screen to produce a

radiogram. (B) The bar plot (bottom) represents the relative HMTase activities of PRC2 in

the absence (grey) or presence of H3K27me3 (blue) peptides, as indicated. Individual

observations are presented, mean represents the normalised densitometry and error bars

represent standard deviation of three independent experiments that were carried out on

different days. (C) Equal amounts (1 or 3 μg) of the PRC2 complexes that were used for the

HMTase assays were analysed on an SDS-PAGE (Tris-Acetate, 3%-8% acrylamide). In all

panels: "insect" and "yeast" referred to protein that was expressed using the baculovirus

expression system or *S. cerevisiae*, respectively. "pFB1" referred to a PRC2 complex that

was expressed by the co-infection of 4 baculovirus stocks, each made of one pFastBac1

plasmid coding for one protein. "MoClo Baculo" referred to PRC2 complexes that were

expressed from either baculovirus or yeast expression vectors, as indicated, constructed

using parts of the MoClo Baculo toolkit.

	Baculovirus	Yeast
Time required to build a multi-gene (Level 2) plasmid	5 days	5 days
Protein expression [a]	20-30 days	6-7 days
Yield per litre (PRC2)	3-5 mg	0.03-0.05 mg
Cost of protein expression [a]	$ 17/mg	$ 500-830/mg
Expression cost per batch [a]	$ 270 per 4 L batch	$ 300 per 12 L batch [b]
Estimated labour per batch [c]	4.6 h per 4 L batch	4.8 h per 12 L batch

Table 1. Comparison between the Baculovirus system to yeast in terms of protein

expression. Values represent the best estimate based on the expression of the 4-subunit

PRC2 complex from multigene expression vectors generated using MoClo Baculo.

[a] 'Protein expression' and 'Expression cost' are defined here as the process from the stage

the plasmid is made until the cells that expressed the protein are pelleted and collected, but

before the protein purification process has started. For the baculovirus system, this process

includes the preparation, amplification and quantification of the baculovirus particles, insect
cell culture buildup, infection, incubation during the protein expression period and cell
harvest. For the yeast system, this process includes transformation and colony growth,
media prep, culture starter and inoculation, induction, growth and cell harvest. Expression
trial in yeast was carried out using an *S. cerevisiae* line that was generated through the
homologous recombination of the multi-gene expression vector into a chromosome.
Estimated costs are provided in US dollars.
[b] In this work, protein in *S. cerevisiae* was carried out using a galactose-inducible promoter
on one of the PRC2 subunits (EZH2). This strategy added a substantial cost because of the
usage of lactate (expression suppressor) and galactose (expression inducer). A usage of
constitutive promoter, when feasible, would avoid galactose and lactate and therefore
reduce the expression cost by about 50%.
[c] Labour was estimated in the form of cumulative working hours that were dedicated to
the process of protein expression, as defined in [a].

	MoClo Baculo	Multibac [12]	Bac-to-Bac [6]	BiGBac [14]	GoldenBac [15]
Expression system	BV and Yeast (*S. cerevisiae*)	BV	BV	BV	BV
Modular Cloning	Yes	No	No	Modular for the multigene level, but not for building the transcription units.	Modular for the multigene level, but not for building the transcription units.
PCR or Gibson Assembly required	N*	Y	Y	Y	Y
Maximal number of ORFs per expression vector	6 (or 30 through compatibility with biGBac)**	Theoretically unlimited	2	25	15

Table 2. Comparison between MoClo Baculo to other baculovirus expression systems.

*The construction of up to Level 2 plasmids with up to 6 transcription units is done using Golden Gate assembly, even if the Level 2 plasmids are compatible with the biGBac system. Yet, the optional connection of multiple biGBac-compatible Level 2 plasmids with up to 6 transcription units each into a single Level 3 plasmids of up to 30 transcription units is done using the biGBac system and required Gibson Assembly.

**Pre-build parts support up to 6 ORFs per MoClo Baculo vector (Level 2) and up to 30 ORFs if biGBac compatible parts are used to connect multiple Level 2 vectors into biGBac vectors (Level 3 here). Yet, the number of ORFs per MoClo Baculo vectors (Level 2), and therefore the number of ORFs in Level 3 vectors, can be expended if more MoClo Baculo Connector parts are designed, as in the case of the MoClo Yeast Toolkit [18]. However, we

733   find that the Golden Gate assembly of Level 2 plasmids becomes very inefficient when the

734   final product size approaches 30 kbp.

Supplementary Figure 1. Level 1 plasmid architecture of the MoClo toolkit. (A) The

MoClo Baculo Level 1 plasmid design is illustrated, including the left connector (ConL),

promoter (Pro), N-terminal tag (N'tag), coding sequence (CDS), C-terminal tag (C'Tag),

terminator (Term) and the right connector (ConR). Changes that were made to the original

MoClo Yeast toolkit design [18] in order to enable the usage of the same CDS part either in

the presence or absence of an N-terminal tag are indicated above (highlighted in blue

boxes). Presented are also the four base pair overhangs that are generated in each part

after BsaI digestion. The parts are numbered according to the MoClo Yeast numbering

system, with the exception of parts that were modified (in the blue box) and the backbone

parts that uses the same overhangs as in the MoClo Yeast toolkit but are all collectively

numbered as part 6 for simplicity (specifically, the MoClo Baculo parts 6a, 6b, 6c and 6d

were originally termed parts 6, 7, 8a and 8b, respectively, in the Moclo Yeast toolkit). Parts

2-4 are related to the expressed gene, part 6 allows recombination into baculovirus DNA or

into the yeast genome, and parts 1 and 5 are for the integration into multi-gene vectors

(Level 2 plasmids). (B) A comparison between the organisation of different functional

modules of various baculovirus expression vectors that are compatible with the Bac-to-Bac

expression system for the expression of single (top) or multiple (bottom) genes and the

single- (Level 1; "Lv1") and multi-gene (Level 2; "Lv2") MoClo Baculo expression vectors.

Presented are the Tn7 left and right recombination sites (Tn7R and Tn7L), gentamicin,

kanamycin, spectinomycin, ampicillin and chloramphenicol resistance (GenR, KanR, SpecR,

AmpR and CamR), LoxP recombination site (LoxP), left and right connectors for Golden

Gate assembly (ConL and ConR), a region that hosts multiple genes or transcription units

(TU integration), Simian virus polyadenylation signal (SV40), Herpes Simplex Virus

thymidine kinase polyadenylation signal (HSV), coding sequence (CDS), baculovirus p10

and polyhedrin promoters (p10 and polH), f1 origin (F1 ori), and pUC origin (ori). (C)

Flanking sequence of promoter (Pro), N-terminal tag (N Tag), coding sequence (CDS), C-

terminal tag (C'Tag), terminator (Term) within the context of Level 0 vectors. BsaI restriction

site is highlighted in orange and the overhang in yellow. Overhang sequences are inherited

from the MoClo yeast toolkit, with the exception of parts 2a, 2b and 3, which were

redesigned. (D) In order to generate Level 0 plasmids that holds a given part (presented in

panel (C)), the part can be obtained as a double strand DNA (e.g. as a gene fragment or a

plasmid from gene synthesis) while including flanking sequences suitable for its Golden Gate

Assembly into the pYTK001 plasmid (originated from the MoClo Yeast Tool Kit [18]) using

BsmBI. As an example, presented are the complete flanking sequences that are added to a

CDS part ("3 CDS" representing an open reading frame sequence) that is ordered from gene

synthesis. BsmBI restriction sites are highlighted in grey, the overhang for assembly with

pYTK001 is underlined and the rest of the colours are as in panel (C). The black crosses

represent the site of ligation during BsmBI golden gate reaction.

A

Supplementary Figure 2. A construct design of a yeast expression plasmid. Schematic of the Level 1 and Level 2 constructs that were made using parts of the MoClo Baculo and MoClo Yeast toolkits [18] for the expression of the human PRC2 in *S. cerevisiae*. Parts from the MoClo Yeast toolkit are in black boxes and parts of the MoClo Baculo (parts 2b and 2a; see in Supplementary Fig. 1A) are in blue boxes. Dashed blue boxes represent the same CDS parts that were also used to construct baculovirus expression vectors in Fig. 2. "BB-URA3" mark a backbone plasmid that originated from the MoClo Yeast toolkit [18] (pYTK096), aiming to enable genomic integration into the *URA3* locus of the *S. cerevisiae* chromosome, using selection by uracil prototrophy.

Supplementary Figure 3. Human PRC2 complexes expressed using MoClo Baculo

vectors in insect cells and in yeast are enzymatically active. (A) Uncropped radiogram

and coomassie stained gels from three independent replicates of HMTase activity assay

that were carried out on three different days (the cropped gels are in Fig. 4A). The ^{14}C-

labelled H3 standard that is used for normalisation between different replicates is shown on

the right side of the gel. B. The EZH2 automethylation signal was quantified using ^{14}C-

labelled H3 standards and the values were plotted as a bar chart. C. H3 standard curves that

were used to normalise data from independent replicates of HMTase activity assays. The

standard curves of the different replicates are presented, with colour key as indicated.

Supplementary Figure 4. MoClo Baculo level 2 plasmids can be connected into larger

Level 3 plasmids through compatibility with the biGBac system. (A) pMB.BIG1a to

pMB.BIG1e plasmids (bottom) were generated by inserting biGBac linker sequence (box

marked with A to F) and PmeI site between the pMB.002 plasmid backbone and BsmBI site.

(B) The workflow of constructing biGBac compatible level 2 vectors is essentially the same

as other MoClo Baculo vectors, except that the pMB.BIG1a to pMB.BIG1e plasmids are

used as Level 2 backbone instead of the pMB.002 plasmid. Colour key is as in Fig. 1 with

addition of biGBac linker sequences A to F in assorted colours. TU: transcriptional unit; TUs:

BsmBI golden gate ligated transcriptional units; SpecR: spectinomycin resistance; CamR:

chloramphenicol resistance.

Supplementary Figure 5. Proof-of-principle: Construction of a Level 3 vector through compatibility between the MoClo Baculo to the biGBac system. (A) Gibson assembly of

pMB.BIG1a.PRC2, pMB.BIG1b.MTF2-PALI1 and pMB.BIG1c.G9A-GLP plasmids were

carried out using pBIG2abc as backbone. The final plasmid, pBIG2abc.PRC2.1-EHMT, is

validated by restriction digestion using MfeI and PmeI. Chemical- and electroporation

transformation were benchmarked here. Clone 1 are correct clones, while 2 and 3 are empty

pBIG2abc. (B) Schematic representation of the plasmids used for the Gibson assembly

reaction that was analysed in (A). (C) Schematic representation of the plasmids produced by

the Gibson assembly reaction was analysed in (A).

Legends for Supplementary Tables and Data

Supplementary Table 1. Compatibility between parts for the construction of expression vectors for baculovirus and yeast.

Tabulated are relevant parts (level 0 plasmids) from the MoClo Baculo and MoClo yeast toolkits, where the latter are indicated by an asterisk. Part numbers and sub-part numbers are as in Supplementary Fig 1A. Expression system-specific parts, either for the baculovirus (BV) system or yeast, are clustered together (see left column). All the tags, coding sequences (CDS) and connector parts of the MoClo Baculo system are compatible with vector building for both yeast and baculovirus expression. Promoter, terminator and backbone parts are system-specific.

Supplementary Table 2. Plasmids that were generated during this study. Indicated are plasmids that were constructed during this study, including new parts (Level 0 plasmids) and expression vectors (Level 1 and 2 plasmids). Excluded are vectors that were constructed as intermediate products of multi-gene vector buildings for the yeast expression system, as they were not designed to express proteins.

Supplementary Table 3. Sequences of ssDNAs that were used to generate linker sequences for compatibility with the biGBac system. Indicated are the sequences that were added to generate the modified ConLS'A to ConLS'E and ConRE'B to ConRE'F parts that were used as precursors for generating the biGBac compatible plasmids pMB.BIG1a to pMB.BIG1e, respectively.

Supplementary Data 1. Plasmid sequences. Sequences of baculovirus-specific Level 0 plasmids that were generated during this study are included within a zipped directory as GenBank files.

Supplementary Data 2. Plasmid sequences. Sequences of yeast-specific Level 0 plasmids that were generated during this study are included within a zipped directory as GenBank files.

Supplementary Data 3. Plasmid sequences. Sequences of Level 0 plasmids that are used for both yeast and the baculovirus system and that were generated during this study are included within a zipped directory as GenBank files.

Supplementary Data 4. Plasmid sequences. Sequences of ORFs as Level 0 plasmids that were synthesised during this study are included within a zipped directory as GenBank files.

Supplementary Data 5. Plasmid sequences. Sequences of single-gene and multi-gene

baculovirus expression vectors (Level 1 and Level 2) plasmids, respectively, that were

generated during this study are included within a zipped directory as GenBank files.

Supplementary Data 6. Plasmid sequences. The sequence of a yeast expression

multigene plasmid (Level 2) that was generated during this study is included within a zipped

directory as GenBank files.

Reference

1. Saibil HR. Cryo-EM in molecular and cellular biology. Mol Cell. 2022 Jan
20;82(2):274-284. doi: 10.1016/j.molcel.2021.12.016. PMID: 35063096.
2. Baser B, van den Heuvel J. Assembling Multi-subunit Complexes Using Mammalian
Expression. Adv Exp Med Biol. 2016;896:225-38. doi: 10.1007/978-3-319-27216-
0_15. PMID: 27165329.
3. Assenberg R, Wan PT, Geisse S, Mayr LM. Advances in recombinant protein
expression for use in pharmaceutical research. Curr Opin Struct Biol. 2013
Jun;23(3):393-402. doi: 10.1016/j.sbi.2013.03.008. Epub 2013 Jun 1.
PMID:23731801.
4. Kost TA, Condreay JP, Jarvis DL. Baculovirus as versatile vectors for protein
expression in insect and mammalian cells. Nat Biotechnol. 2005 May;23(5):567-75.
doi: 10.1038/nbt1095. PMID: 15877075; PMCID: PMC3610534.
5. Baghban R, Farajnia S, Rajabibazl M, Ghasemi Y, Mafi A, Hoseinpoor R, Rahbarnia
L, Aria M. Yeast Expression Systems: Overview and Recent Advances. Mol
Biotechnol. 2019 May;61(5):365-384. doi: 10.1007/s12033-019-00164-8. PMID:
30805909.
6. Anderson D, Harris R, Polayes D. Rapid generation of recombinant baculoviruses
and expression of foreign genes using the Bac-to-Bac baculovirus expression
system. 1996; Focus 17:53–58
7. Sokolenko S, George S, Wagner A, Tuladhar A, Andrich JM, Aucoin MG. Co-
expression vs. co-infection using baculovirus expression vectors in insect cell culture:
Benefits and drawbacks. Biotechnol Adv. 2012 May-Jun;30(3):766-81. doi:
10.1016/j.biotechadv.2012.01.009. Epub 2012 Jan 28. PMID: 22297133; PMCID:
PMC7132753.
8. Aucoin MG, Perrier M, Kamen AA. Improving AAV vector yield in insect cells by
modulating the temperature after infection. Biotechnol Bioeng. 2007 Aug
15;97(6):1501-9. doi: 10.1002/bit.21364. PMID: 17274066.
9. Aucoin MG, Perrier M, Kamen AA. Production of adeno-associated viral vectors in
insect cells using triple infection: optimization of baculovirus concentration ratios.
Biotechnol Bioeng. 2006 Dec 20;95(6):1081-92. doi: 10.1002/bit.21069. PMID:
16952153.
10. Meghrous J, Aucoin MG, Jacob D, Chahal PS, Arcand N, Kamen AA. Production of
recombinant adeno-associated viral vectors using a baculovirus/insect cell
suspension culture system: from shake flasks to a 20-L bioreactor. Biotechnol Prog.
2005 Jan-Feb;21(1):154-60. doi: 10.1021/bp049802e. PMID: 15903253.

11. Zhang Q, McKenzie NJ, Warneford-Thomson R, Gail EH, Flanigan SF, Owen BM,
Lauman R, Levina V, Garcia BA, Schittenhelm RB, Bonasio R, Davidovich C. RNA
exploits an exposed regulatory site to inhibit the enzymatic activity of PRC2. Nat
Struct Mol Biol. 2019 Mar;26(3):237-247. doi: 10.1038/s41594-019-0197-y. Epub
2019 Mar 4. PMID: 30833789; PMCID: PMC6736635.

12. Pelosse M, Crocker H, Gorda B, Lemaire P, Rauch J, Berger I. MultiBac: from protein
complex structures to synthetic viral nanosystems. BMC Biol. 2017 Oct 30;15(1):99.
doi: 10.1186/s12915-017-0447-6. PMID: 29084535; PMCID: PMC5661938.

13. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA 3rd, Smith HO.
Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat
Methods. 2009 May;6(5):343-5. doi: 10.1038/nmeth.1318. Epub 2009 Apr 12. PMID:
19363495.

14. Weissmann F, Petzold G, VanderLinden R, Huis In 't Veld PJ, Brown NG, Lampert F,
Westermann S, Stark H, Schulman BA, Peters JM. biGBac enables rapid gene
assembly for the expression of large multisubunit protein complexes. Proc Natl Acad
Sci U S A. 2016 May 10;113(19):E2564-9. doi: 10.1073/pnas.1604935113. Epub
2016 Apr 25. PMID: 27114506; PMCID: PMC4868461.

15. Neuhold J, Radakovics K, Lehner A, Weissmann F, Garcia MQ, Romero MC, Berrow
NS, Stolt-Bergner P. GoldenBac: a simple, highly efficient, and widely applicable
system for construction of multi-gene expression vectors for use with the baculovirus
expression vector system. BMC Biotechnol. 2020 May 12;20(1):26. doi:
10.1186/s12896-020-00616-z. PMID: 32398045; PMCID: PMC7216392.

16. Weber E, Engler C, Gruetzner R, Werner S, Marillonnet S. A modular cloning system
for standardized assembly of multigene constructs. PLoS One. 2011 Feb
18;6(2):e16765. doi: 10.1371/journal.pone.0016765. PMID: 21364738; PMCID:
PMC3041749.

17. Engler C, Gruetzner R, Kandzia R, Marillonnet S. Golden gate shuffling: a one-pot
DNA shuffling method based on type IIs restriction enzymes. PLoS One.
2009;4(5):e5553. doi: 10.1371/journal.pone.0005553. Epub 2009 May 14. PMID:
19436741; PMCID: PMC2677662.

18. Lee ME, DeLoache WC, Cervantes B, Dueber JE. A Highly Characterized Yeast
Toolkit for Modular, Multipart Assembly. ACS Synth Biol. 2015 Sep 18;4(9):975-86.
doi: 10.1021/sb500366v. Epub 2015 May 1. PMID: 25871405.

19. Enghiad B, Xue P, Singh N, Boob AG, Shi C, Petrov VA, Liu R, Peri SS, Lane ST,
Gaither ED, Zhao H. PlasmidMaker is a versatile, automated, and high throughput
end-to-end platform for plasmid construction. Nat Commun. 2022 May 16;13(1):2697.
doi: 10.1038/s41467-022-30355-y. PMID: 35577775; PMCID: PMC9110713.

20. Margueron R, Justin N, Ohno K, Sharpe ML, Son J, Drury WJ 3rd, Voigt P, Martin
SR, Taylor WR, De Marco V, Pirrotta V, Reinberg D, Gamblin SJ. Role of the
polycomb protein EED in the propagation of repressive histone marks. Nature. 2009
Oct 8;461(7265):762-7. doi: 10.1038/nature08398. Epub 2009 Sep 20. PMID:
19767730; PMCID: PMC3772642.

21. Lee CH, Yu JR, Granat J, Saldaña-Meyer R, Andrade J, LeRoy G, Jin Y, Lund P,
Stafford JM, Garcia BA, Ueberheide B, Reinberg D. Automethylation of PRC2
promotes H3K27 methylation and is impaired in H3K27M pediatric glioma. Genes
Dev. 2019 Oct 1;33(19-20):1428-1440. doi: 10.1101/gad.328773.119. Epub 2019
Sep 5. PMID: 31488577; PMCID: PMC6771381.

22. Wang X, Long Y, Paucek RD, Gooding AR, Lee T, Burdorf RM, Cech TR. Regulation
of histone methylation by automethylation of PRC2. Genes Dev. 2019 Oct 1;33(19-
20):1416-1427. doi: 10.1101/gad.328849.119. Epub 2019 Sep 5. PMID: 31488576;
PMCID: PMC6771386.

23. Lowin T, Raab U, Schroeder J, Franssila R, Modrow S. Parvovirus B19 VP2-proteins
produced in Saccharomyces cerevisiae: comparison with VP2-particles produced by
baculovirus-derived vectors. J Vet Med B Infect Dis Vet Public Health. 2005 Sep-
Oct;52(7-8):348-52. doi: 10.1111/j.1439-0450.2005.00871.x. PMID: 16316399.

24. Morton CL, Potter PM. Comparison of Escherichia coli, Saccharomyces cerevisiae,
Pichia pastoris, Spodoptera frugiperda, and COS7 cells for recombinant gene
expression. Application to a rabbit liver carboxylesterase. Mol Biotechnol. 2000
Nov;16(3):193-202. doi: 10.1385/MB:16:3:193. PMID: 11252804.

25. Moore SJ, Lai HE, Kelwick RJ, Chee SM, Bell DJ, Polizzi KM, Freemont PS.
EcoFlex: A Multifunctional MoClo Kit for E. coli Synthetic Biology. ACS Synth Biol.
2016 Oct 21;5(10):1059-1069. doi: 10.1021/acssynbio.6b00031. Epub 2016 May 2.
Erratum in: ACS Synth Biol. 2020 May 15;9(5):1225. doi:
10.1021/acssynbio.0c00177. PMID: 27096716.

26. Otto M, Skrekas C, Gossing M, Gustafsson J, Siewers V, David F. Expansion of the
Yeast Modular Cloning Toolkit for CRISPR-Based Applications, Genomic Integrations
and Combinatorial Libraries. ACS Synth Biol. 2021 Dec 17;10(12):3461-3474. doi:
10.1021/acssynbio.1c00408. Epub 2021 Dec 3. PMID: 34860007; PMCID:
PMC8689691.

27. Iverson SV, Haddock TL, Beal J, Densmore DM. CIDAR MoClo: Improved MoClo
Assembly Standard and New E. coli Part Library Enable Rapid Combinatorial Design
for Synthetic and Traditional Biology. ACS Synth Biol. 2016 Jan 15;5(1):99-103. doi:
10.1021/acssynbio.5b00124. Epub 2015 Nov 4. PMID: 26479688.

28. Obst U, Lu TK, Sieber V. A Modular Toolkit for Generating Pichia pastoris Secretion
Libraries. ACS Synth Biol. 2017 Jun 16;6(6):1016-1025. doi:
10.1021/acssynbio.6b00337. Epub 2017 Mar 15. PMID: 28252957.
29. Fonseca JP, Bonny AR, Kumar GR, Ng AH, Town J, Wu QC, Aslankoohi E, Chen
SY, Dods G, Harrigan P, Osimiri LC, Kistler AL, El-Samad H. A Toolkit for Rapid
Modular Construction of Biological Circuits in Mammalian Cells. ACS Synth Biol.
2019 Nov 15;8(11):2593-2606. doi: 10.1021/acssynbio.9b00322. Epub 2019 Nov 5.
PMID: 31686495.
30. McGuffie MJ, Barrick JE. pLannotate: engineered plasmid annotation. Nucleic Acids
Res. 2021 Jul 2;49(W1):W516-W522. doi: 10.1093/nar/gkab374. PMID: 34019636;
PMCID: PMC8262757.