1

Effective Polarization in Potassium Channel Simulations: Ion Conductance, Occupancy, Voltage Response, and Selectivity

Chenggong Hui[1], Reinier de Vries[1], Wojciech Kopec[1,2]*, Bert L. de Groot[1]*

[1]Computational Biomolecular Dynamics Group, Max Planck Institute for Multidisciplinary Sciences, Am Fassberg 11, 37077 Göttingen, Germany

[2]Department of Chemistry, Queen Mary University of London, 327 Mile End Road, London E1 4NS, United Kingdom

*corresponding authors: w.kopec@qmul.ac.uk, bgroot@gwdg.de

Classification : Biological Science, Biophysics and Computational Biology

Keywords : Potassium Channel, MD simulation, Force Field, Charge Scaling, Selectivity

Significance Statement

Potassium (K^+) channels are essential membrane proteins that facilitate the selective conduction of K^+ ions while excluding Na^+ ions. Incorporation of electronic polarization effects through the Electronic Continuum Correction (ECC) method enables molecular dynamics simulations to reproduce rapid K^+ permeation under physiological voltage conditions. In this study, we demonstrate that employing ECC approximated polarization in molecular dynamics simulations allows for the prediction of multiple channel properties: conductance, ion occupancy, voltage response, and selectivity, with unprecedented accuracy. Furthermore, our simulations provide atomistic insights into the asymmetrical current-voltage (I-V) relationship observed in the MthK channel.

Abstract

Potassium (K^+) channels are widely distributed in many types of organisms. They combine high efficiency (~100 pS) and K^+/Na^+ selectivity by a conserved selectivity filter (SF). Molecular Dynamics (MD) simulations can provide detailed, atomistic mechanisms of this sophisticated ion permeation. However, currently there are clear inconsistencies between computational predictions and experimental results. Firstly, the ion occupancy of the SF in simulations is lower than expected (~2.5 in MD compared to ~4 in X-ray crystallography). Secondly, in many reported MD simulations of K^+ channels, K^+ conductance is typically an order of magnitude lower than experimental values. This discrepancy is in part because the force fields used in MD simulations of potassium channels do not account for polarization. One of the proposed solutions is the Electronic Continuum Correction (ECC), a force field modification that scales down formal charges, to introduce the polarization in a mean-field way. When the ECC is used in conjunction with the Charmm36m force field, the simulated K^+ conductance increases 13-fold. Following the analysis of ion occupancy states using Hamiltonian Replica Exchange (HRE) simulations, we propose a new parameter set for Amber14sb, that also leads to a similar increase in conductance. These two force fields are then used to compute, for the first time, the full current-voltage (I-V) curves from MD simulations, approaching quantitative agreement with experiments at all voltages.

48   In general, the ECC-enabled simulations are in excellent agreement with experiment, in terms of

49   ion occupancy, conductance, current-voltage response, and K^+/Na^+ selectivity.

1.   Introduction

Potassium channels play an important role in many cell types. They set the resting potential in

non-excitable cells or reset the potential in depolarization in excitable cells. Potassium channels

are known for their efficient and selective permeation of potassium ions. For instance, the NaK2K

channel conducts Na^+ ions at ~10 pS in NaCl solution while conducting K^+ ions at 120 pS in KCl

solution.[1] All potassium channels share a conserved selectivity filter (SF), which forms 4 adjacent

ion binding sites made of backbone carbonyl oxygens (sites S1-S3), plus typically 4 threonine

hydroxyl oxygens at the S4 site. When the high-resolution crystal structure of KcsA was first

published[2], the electron density in the 4 binding sites was initially interpreted as two K^+ ions

occupying two non-adjacent binding sites and two water molecules placed in between the ions.

This interpretation suggests that the four peaks in the electron density represent an overlap of two

distinct states: K^+-water-K^+-water and water-K^+-water-K^+.[3] Consequently, the hypothesis of a

water-mediated "soft knock-on" permeation mechanism was coined. More recent anomalous X-

ray diffraction studies in NaK2K and KcsA channels suggested that all of the 4 sites are fully

occupied by K^+ ions,[4–6] which supports an alternative hypothesis of direct knock-on mechanism,

observed in numerous Molecular Dynamics (MD) studies,[7–10] that does not include any co-

permeation of water. A more detailed discussion can be found in our recent review.[11]

Despite the debated permeation mechanism, low conductance and low K^+ occupancy of the

binding sites were consistently observed in MD simulations of potassium channels displaying both

soft and direct knock-on mechanisms, although direct knock-on typically results in higher

conductance[7,12]. In our previous work, direct knock-on was observed during continuous

permeation under membrane voltage, with the SF typically containing 2 or 3 ions.[7] Specifically,

spontaneous permeation under voltage has been observed on MthK, KcsA, NaK2K, BK, Kv1.2,

TREK2, and TRAAK,[8,13,14] but the conductance was generally one order of magnitude lower than

the experimental values. Ion occupancy in the SF, permeation mechanism, and conductance in MD

simulations all eventually result from a subtle balance of various interactions between ions, protein,

and water. Even tiny changes in the interaction parameters between them might change the kinetics,

thermodynamics, and mechanism of ion permeation.

One often discussed cause for the observed shortcomings in current simulations of K^+ channels is

the lack of explicit electronic polarization in classical force fields.[15] In a classical, fixed-charged

force field (e.g. Amber, Charmm, OPLS-AA), atoms are modeled as spheres (particles) with

assigned partial charges, as an approximation of positively charged nuclei with clouds of electrons

around them. In MD simulations, electrostatic interactions are computed between these particles

by applying the Coulomb law and using the dielectric constant of vacuum. In reality however, the

electrostatic interactions between the nuclei will be (differently) screened by the instantaneous

distribution of the electrons, that will be in turn affected by the position of the nuclei (i.e. electronic

polarization), resulting in the relative permittivity >1. Therefore, for the more accurate modeling

of electrostatic interactions, two general solutions have been proposed. In the first one, force fields

with explicit electronic polarization are developed, whereas in the second one, the so-called

Electronic Continuum Correction (ECC) is applied to existing, classical force fields. Ren and

coworkers have tested both solutions in MD simulations of a K^+ channel[16], focusing on the

different K^+ occupancy states of the SF. They found that when the ECC is applied, Charmm36m

behaves similarly to the AMOEBA polarizable force field, and both show the SF being fully

occupied with four K^+ ions to be energetically the most stable configuration. In practice, the ECC

is achieved by rescaling the formal partial charges of ionized groups (such as side chains of

arginine, lysine, glutamine, and aspartate amino acids) and ions, which is mathematically the same

as changing the dielectric constant in computing electrostatic interactions via Coulomb law.[17,18]

This approach has been recently applied to improve force fields for ions, proteins, and lipids.[19–23]

In general, the argument is made that the rescaling factor should be ~sqrt(1/2) ≈ 0.7, because the

electronic screening factor (high-frequency dielectric constant) is about 2 for most organic

media.[24] However, there are at least two reasons why directly applying this number might not be

optimal. First, the exact dielectric constant of the approximated medium between MD particles is

not known and might be system-dependent. Our main interest here is the magnitude of channel

conductance and corresponding free energy difference between relevant SF occupancies. Second,

most of the force field parameters were developed without ECC in mind, and might compensate

for the mean polarization to some extent. Therefore, instead of focusing on a single scaling factor, in this work, we combine the ECC approach with two common fixed-charge force fields: Charmm36m and Amber14sb, at various scaling factors (1.0-0.65), to study ion permeation in three different potassium channels. We carefully developed new Lennard-Jones (LJ) parameters, describing van der Waals (vdW) interactions, at each scaling factor, so that the ion radius and solvation free energy are consistent with the original force field. We observe that the ECC dramatically increases the simulated conductance in Charmm36m, to a level that quantitatively agrees with the experimental single-channel conductance. In contrast, the ECC has only a minor effect with the Amber14sb force field. These discrepancies are rationalized using Hamiltonian Replica Exchange (HRE) simulations, providing insights into the effect of the ECC on the SF ion occupancy, and resulting in adjusted SF partial charges for Amber14sb, which consequently predicts conductance close to the experiments as well. We further apply these force fields in simulations under physiological voltage <-150, +150 mV>, recording, for the first time, full I-V curves, compatible with experiments. The K^+/Na^+ selectivity is preserved, indicating the necessity of adjusting LJ parameters of ions in ECC. Finally, applying the recently introduced permeation cycle analysis, we provide cues into the physical basis of inward rectification in the MthK channel.

2.   Results and Discussion

2.1.   Low conductance and SF occupancy in MD with Charmm36m and Amber14sb force fields

To obtain a comprehensive view on ion permeation in K^+ channels, we simulated three different channels, namely NaK2K, TRAAK, and MthK. Whereas the archaeal MthK channel shares the SF sequence (TVGYGD) with the NaK-derived NaK2K channel, the SF sequence of TRAAK is more varied: being a dimer, TRAAK has two SF strands (called SF1) with a TIGYGN sequence, and two with TVGFGD (SF2). The inclusion of TRAAK simulations allows us to generalize our findings beyond the most common SF sequence. The channels were simulated using both the Charmm36m and Amber14sb force fields. When compared with experiments, a clear discrepancy in quantities describing ion permeation, such as conductance and SF occupancy is seen, as previously reported by us and others.[7–9,13] Generally speaking, single channel conductance (Fig.

1a) is underestimated by an order of magnitude in all channels and force fields tested. The

instantaneous occupancy of the SF binding sites during permeation (Fig. 1b) predominantly shows

(despite differences between force fields, Table S1) an alternation between 3-ion and 2-ion

(counting ions in S1 to S4) SF occupancy states. Although this quantity is not readily available

from experiments, and the SF occupancy during permeation is debated (see Introduction), we note

here that the 3-ion and 2-ion states underestimate the possible number of ions in the SF, as

compared to anomalous diffraction data (full SF occupancy in S1-S4 at 100 K and 3.44 total

occupancy at 292 K).[4,25] These results clearly show the need for improved force fields for MD

simulations of K^+ channels. Efforts have been made in this direction. For example a special

exception interaction (NBFIX) was developed to weaken the interaction between carbonyl

oxygens and K^+ ions,[26] but reproducing the conductance was found to be challenging.[27] We reason

that single channel potassium conductance is the central quantity describing ion permeation, which

can be directly compared between simulations and experiments. Therefore, an improved force field

should predict the single channel conductance close to the experimental value, across the same

voltage range as used in electrophysiology (-150 ~ 150 mV), and ideally also in the physiological

range (resting potential: -10 ~ -100 mV, action potential: ~+100 mV).[28,29] We further reason that

such a force field would likely better describe any secondary observables characterizing ion

permeation (i.e. SF occupancy states, rate limit steps, SF geometries etc.) as well.

In this contribution, we show how the ECC-introduced polarization in Charmm36m and

Amber14sb force fields improves simulated single channel conductance in all channels tested.

Interestingly, the application of charge scaling simultaneously results in SFs occupied by more K^+

ions (3-ion and 4-ion states) than in non-scaled force fields, agreeing well with experimental data,

and further highlighting the role of direct ion-ion contact in rapid K^+ permeation through K^+

channels.

2.2.   ECC-Charmm36m improves the conductance, but ECC-Amber does not.

We performed simulations with different scaling factors, starting at the nominal charge of +1.0,

with a step of 0.05, down to +0.65, in Charmm36m and Amber14sb. Similarly to simulations with

unscaled charges, we initially used an applied voltage of $+150$ mV to drive ion permeation. The
conductance at different levels of ECC (scaling factors) in the NaK2K channel is shown in Fig.
2a-c (see Fig. S4 and S5 for TRAAK and MthK channels, respectively). We first focus on
simulations using Charmm36m (Fig. 2a, d, g). Compared to the standard parameters (scaling factor
of 1.00), the conductance decreases at $+0.9$ charge, but then significantly increases around $\sim +0.75$.
At the scaling factor of 0.75, the conductance is 122.4 ± 10.1 pS, which is a 13-fold increase
compared to standard Charmm36m (9.2 ± 1.7 pS), and in apparent excellent agreement with the
experimental single channel conductance of ~ 120 pS.[1] The equilibrated populations of the SF
states show major changes with the applied scaling factor. Compared to standard Charmm36m, 4-
ion states become gradually more populated with stronger scaling (Fig. 2d, g): the WKKKK0 state
is dominant (79%) with 0.7 scaling, in stark contrast to unscaled Charmm36m (WKK0K0 (37%),
WK0KK0 (20%), WK0K0K (20%)). The increase of ion occupancy is consistent with the
previously reported ECC-Charmm36m simulation in KcsA.[16] A similar effect of increased ion
occupancy with charge scaling is observed in Amber14sb (Fig. 2e,h), however there is no effect
on conductance, as it remains low (~ 20 pS) at all scaling factors tested, also in the other two
channels (Fig. S4 and S5).

2.3.   Redistributing the charge in the SF in Amber14sb increases the conductance

Prompted by the conductance insensitivity to charge scaling in Amber14sb, we decided to
investigate this issue in detail. By comparing simulations carried out with Charmm36m and
Amber14sb we noticed two important (and possibly related) differences: 1. In Charmm36m, the
carbonyl oxygen has a partial charge of -0.51, whereas it is -0.5679 in Amber14sb, potentially
resulting in stronger binding of a K^+ ion to the SF, and 2. For the <0.85, 0.75> range of charge
scaling in Amber14sb, the K0KKK0 state is the dominant one, and, for scalings < 0.75, WKKKK0
dominates (Fig. 2e). These observations suggest that K^+ ions in Amber14sb are potentially too
strongly bound to the SF, especially to its lower part, which might slow down permeation. To test
this idea, we reduced the negative charge on the carbonyl oxygen atoms at S3 to different values
(-0.55, -0.53, -0.51), and evenly distributed the remaining charge on the other atoms in the SF (see
Fig S1 for the illustration of oxygen charge in the SF). -0.53 gave the SF state population similar

to Charmm36m (see Fig. 2 and Fig. S2 for SF state population). We further tested the -0.53 charge modification and call it Amber14sb-S3 from now on.

With this modified force field at hand, we repeated all the simulations for all three channels (Fig. 2c, f, i, Fig. S4, Fig. S5). Interestingly, the state populations (Fig. 2f and i) are now more similar to ECC-Charmm36m. When compared to the original Amber14sb, the major difference is that the population of K0KKK0 which dominated the distribution at 0.8 (Fig. 2e) is now significantly reduced (Fig. 2f). Importantly, these effects resulted in a major increase in conductance, akin to ECC-Charmm36m (Fig. 2c, maximal conductance at the scaling factor of 0.75: 88.0 ± 3.2 pS). We observe essentially the same behavior in simulations of TRAAK and MthK channels (Fig. S4 and S5).

To obtain clearer mechanistic insights into the factors governing rapid ion permeation at around 0.80-0.75 scaling in both Charmm36m and Amber14sb-S3, we looked again at the populations of SF states, but this time combining states together with the same number of K^+ ions bound to the SF (Fig. 2g-i, Fig. S4, Fig. S5). We find that in high conductance regions, there is a similar population for 3-ion and 4-ion population states (crossing of blue and orange lines), suggesting a smooth (barrierless) free energy landscape between microscopic states (SF occupancies). We hence hypothesize that a balanced (low barrier) free energy surface between multiple 3- and 4-ion states is a prerequisite for efficient, rapid ion permeation through the SF of K^+ channels. Accordingly, if certain states dominate the SF state distribution the permeation cycle would be slowed down by the barrier to leaving this state. For example, in Charmm36m at 0.9, the SF is mostly populated by a single WKK0K0 state, whereas in Amber14sb at 0.8, the SF is mostly populated by K0KKK0. Those extra stable states might contribute to the underestimation of conductance in MD simulations.

Our proof-of-concept Amber14sb-S3 modification opens up a new degree of freedom for force field development in K^+ channel simulations. The default backbone parameters for the SF in Amber14sb do not yield reasonable conductance, instead they result in an SF being stuck at a certain SF state (e.g. K0KKK0 at 0.8 scale). By a slight modification of a partial charge on the

backbone carbonyl oxygen (from -0.5679 e to -0.53 e) at the S3 binding site, we adjusted the

relative free energies of SF states resulting in an efficient ion permeation. We do not suggest our

applied parameter changes as a general force field patch, but a special parameter set for K^+ channel

SF.

Comparing the conductance-scaling relationship for all three channels (Fig. 2a-c, Fig. S4a-c, Fig.

S5a-c), the maximal conductance is observed at remarkably similar scaling factors. In ECC-

Charmm36, the conductance for TRAAK and NaK2K reached their maximum at 0.78 and 0.77,

whereas for MthK at 0.75. In ECC-Amber14sb-S3, 0.78 for TRAAK, 0.75 for NaK2K and MthK.

Similarly, the crossing point of 3-ion and 4-ion state populations appears in high conductance

regions for all three channels, following the same order: TRAAK 0.82, NaK2K 0.78, MthK 0.77

(Charmm36m, Fig. 2g, Fig. S4, Fig. S5), TRAAK 0.79, NaK2K 0.77, MthK 0.77 (Amber14sb-S3,

Fig. 2i, Fig. S4, Fig. S5). These results agree well with our hypothesis that the presence of multiple

3- and 4-ion states at similar (balanced) free energy levels is a prerequisite for high conductance

in K^+ channels. And also agree with the room temperature anomalous scattering occupancy

result.[25]

2.4. The ECC force fields predict I-V curves in agreement with the experiment

Encouraged by the unprecedented increase of the simulated single channel conductance at a single

voltage of +150 mV when using ECC-Charmm36 and ECC-Amber14sb-S3, we decided to test

these force fields in the whole experimental voltage range. As a result, we obtained, for the first

time, the full current-voltage (I-V) curves of K^+ channels derived purely from MD simulations, for

experimentally and physiologically relevant voltage ranges (Fig. 3). The ECC correction improves

conductance at all voltages tested, bringing simulated currents with the ~0.80-0.75 scaling factors

very close to the experimental currents (black traces) for all three channels, and in both force fields.

This effect is particularly pronounced for the experimentally relevant voltage range (<-150 mV,

+150 mV>), where both outward and inward currents with unscaled charges are very low (blue

traces). Notably, in all conditions tested, the rapid ion permeation occurs through some variations

of the direct knock-on mechanism, that is without any co-permeating water molecules. Generally,

the probability of finding water in SF is very low. (K^+/Water occupancy in SF is listed in Table

S3). Similar to our findings in the previous section, the highest conductance at all voltages is

enabled by the co-existence of several 3-ion and 4-ion states in the SF with similar populations.

In some cases, the application of the ECC correction leads to simulated currents that are higher

than their experimental counterparts (e.g. MthK in ECC-Charmm36m 0.75 scaling at positive

voltages, or TRAAK ECC-Amber14sb-S3 0.80 scaling at all voltages), which, to the best of our

knowledge, has never been reported before for any MD simulation. This observation poses a

challenge in deciding which scaling factor is the optimal one (for a given force field). In this

context, it is worth recalling that the shapes of experimental I-V curves might substantially vary

between K^+ channels. For the channels tested here, both NaK2K and TRAAK display linear I-V

curves[1,30], whereas MthK shows inward rectification[10]. Taking this into account, we notice that

ECC-Charmm36m with the 0.78 scaling factor (Fig. 3 (a-c), green traces) is in best agreement with

the experimental data, reproducing well shapes of experimental I-V curves, including the inward

rectification phenomenon in MthK. A more complex situation is seen in ECC-Amber14sb-S3.

Here, the 0.78 scaling factor (Fig. 3 (d-f), green traces) is in almost perfect agreement with

experimental data for MthK, and it also predicts the linear I-V curves for NaK2K and TRAAK,

but underestimates the current for Nak2K and TRAAK. These examples show that none of the

ECC scaling tested here is ideal, and further reparameterization work is needed to obtain a robust

set of parameters applicable to MD of any K^+ channel. The observed differences between studied

channels (and also scaling factors) might be rationalized by sequence variations in the selectivity

filter and the second layer of residues surrounding the selectivity filter. Indeed, while the SF

sequence is highly conserved, the residues around it show a higher degree of variability, likely

leading to ion occupancy and conductance changes, due to subtle differences in dynamics and

intermolecular interactions. However, these observations remain secondary to the primary effect

of the ECC correction, that is the substantial (order of magnitude) increase of the single channel

current (conductance), observed in the 0.80-0.75 scaling range, to the level comparable with

experimental estimates. Thus our work opens the possibility to use, for the first time, MD

simulations of K^+ channels in a quantitative manner to study a wide range of phenomena, using

the same observable (the current-voltage relationship) as in electrophysiological experiments.

285 Our ECC simulations carried out at voltages outside of the experimental range of <-150 mV, +150

286 mV> provide additional insights. For example, the scaling factor showing the maximal

287 conductance in ECC-Charmm36m for NaK2K at 200 mV appears to be 0.75, and, in the absence

288 of experimental data at this voltage, it could have been deemed optimal. However, simulations at

289 lower positive voltages show that this is not the case. Therefore, we conclude that in the future

290 MD simulations of ion channels should be performed, whenever possible, at experimentally and/or

291 physiologically relevant voltages. High voltages have been traditionally used to speed up ion

292 permeation in MD simulations of K^+ channels,[7,10,13,27] occasionally reaching 500 mV or even 750

293 mV,[27,31,32] although the effects of such high voltages on the protein structure and the permeation

294 mechanism itself are not clear.[7] With the ECC approach, there does not seem to be a need to use

295 such high voltages anymore, since robust currents are achievable at experimental voltages as well.

297 Finally, we tested two more channels, Kv1.2-2.1 and KcsA E71A with the optimal parameters for

298 ECC-Charmm36m at 0.78 scaling factor. Good agreement with experimental currents at both +150

299 mV and -150 mV are observed. (Fig. S7)

301 2.5. K^+/Na^+ selectivity is sensitive to both the ion partial charge and LJ parameters

302 All K^+ channels show exquisite K^+/Na^+ selectivity (>10:1 permeability ratio, i.e. the ratio of

303 conductances in pure K^+ and Na^+ solutions),[1,33] therefore this important characteristic should be

304 captured in simulations as well. In this section, we first present thermodynamic properties of our

305 newly developed radius-corrected LJ parameters for ECC ions (All of the simulation results above

306 were using these radius-corrected LJ parameters). Secondly, we compare ECC K^+ channel

307 simulations with these new parameters versus unmodified LJ parameters, highlighting the

308 importance of compatible ion LJ parameters in ECC force fields.

310 The straightforward application of the charge scaling procedure in the ECC has a big effect on ion

311 properties in water. Particularly relevant for ion permeation is the radius of an ion, which is

typically carefully checked versus experimental data during the development of ion parameters.

When scaling of the charge is applied, cations will have effectively larger radii, due to the reduced

interactions of a scaled charge on an ion with the surrounding water molecules (Fig. 4a),

thermodynamically leading to an underestimation of the solvation free energies (Fig. 4b, see

Methods for the derivation of the solvation free energy scaling in the ECC).

To amend this situation, we developed new radius-corrected LJ parameters for ECC ions. This was

done in a way that at any given scaling factor, the radius of the first solvation shell reproduces the

radius in the original (not scaled) force field. We decided to choose the original force field as the

fitting target rather than the experimental value to: 1. facilitate the comparison with the original

force field, and 2. reduce the perturbation to the original force field. As shown in Fig. 4a, our new

parameters for K^+ and Na^+ consistently reproduce the originally parametrized radius. Importantly,

also solvation free energies of our radius-corrected ECC ions closely follow experimental values

(Fig. 4b).

To test the radius-corrected LJ parameters, we performed MD simulations of the same K^+ channels

with Na^+ ions at +150 mV at different scaling factors, and compared simulated outward

conductance with K^+ conductance and experimental values (Fig. 5, Fig. S14). The unmodified

vdW is only used in Fig. 5 (d-e) and Fig. S14 (d-e) for comparison, it is not recommended to use

the unmodified vdW in charge scaling simulations.

With our radius-corrected LJ parameters (Fig. 5 a-c, Fig. S14, Fig. 2, Fig. 3), we observe

consistently larger K^+ conductances than Na^+ conductances at all scaling factors larger than ~0.7.

Specifically, combined with our I-V curves, we deem the scaling factor of 0.78 optimal for current

simulations, i.e. it produces I-V curves that qualitatively match experimental ones, and

simultaneously provides robust K^+/Na^+ selectivity for all three tested channels (Table 1). In

contrast, for a smaller scaling factor (<0.7), all three channels conduct more Na^+ than K^+ ions,

which indicates incorrect selectivity. This observation highlights the balance of strong interactions

governing ion permeation and selectivity in K^+ channels, leading to the necessity of careful

validation of both the scaling factor and LJ parameters of their accurate simulations. We

acknowledge that we did not achieve a perfect match for the conductance ratio across all channels
(Table 1), but accurately obtaining energy barrier differences to predict the exact ratio may exceed
the precision limits of the classical force field. According to the Eyring equation,[34] a 10:1 ratio in
permeation rates at 310K corresponds to approximately the 1.4 kcal/mol difference in the barrier
height, and the 2.5:1 ratio corresponds to the 0.6 kcal/mol difference.

In contrast, when the original (unmodified) LJ parameters are used, K^+/Na^+ selectivity is lost in
almost all conditions tested (Fig. 5 d-e, Fig. S14). Our results therefore suggest that the adjustment
of LJ parameters, and, consequently, maintaining the correct radii of ions, is necessary to capture
their proper permeation and selectivity behavior in K^+ channels with the ECC correction, and will
be likely relevant in other systems showing significant ion selectivity.

2.6.   Mechanistic Basis of Inward Rectification

As a final insight from our ECC simulations, we decided to take a closer look at simulations
displaying inward rectification in the MthK channel (ECC-Charmm36m at 0.78 scaling, Fig. 3c,
green curve), that is the larger current at negative voltages than at positive ones. This inward
rectification is in excellent agreement with experiment (Fig. 3c, black curve). Importantly, this
inward rectification is observed, both experimentally and computationally, for the pore domain
only (i.e. without the additional intracellular calcium-binding domains), indicating that it is an
inherent property of the K^+ permeation mechanism in this channel. To study it in more detail, we
applied a recently proposed analysis of ion permeation mechanisms using permeation cycles.[9] We
simplified the cycle by lumping together several states when their exchange rate was above a
certain cutoff. The comparison of permeation cycles at +150 mV and -150 mV (Fig. 4) reveals that
the rate-limiting step at both voltages is the transition between WKKK0K and WKK0KK states
(i.e. a K^+ ion transitioning from S4 to S3 sites during outward permeation, and in the opposite
direction during inward permeation; while both S1 and S2 remain occupied by K^+ ions). However,
the rate (inverse of the mean first passage time) of this transition differs by 3-fold for these two
voltages (0.318 ns^{-1} for +150 mV and 0.912 ns^{-1} for -150 mV). Our approach allows finding such
rate limiting states in MD simulations of K^+ channels, and thus deciphering the molecular

determinants governing the existence of such states, and rectification properties in different K^+

channels in general.

3. Conclusion

In this work, we introduced and tested a range of parameters for K^+ and Na^+ ions consistent with

the ECC framework, i.e. including effective electronic polarization, in MD simulations of

potassium channels. We applied the same methodology to two fundamentally different force fields

(Charmm36m and Amber14sb), ran simulations of three different K^+ channels over the whole

range of physiologically and experimentally relevant voltages. Despite minor differences between

specific cases, our findings can be broadly summarized in three major points: 1. The introduction

of the electronic polarization of cations (here through the ECC approach) is crucial to accurately

model K^+ ion permeation through potassium channels, allowing to simulate, for the first time, full

I-V curves from atomistic MD simulations that are in very good agreement with experimental data;

2. This efficient K^+ permeation occurs in all cases through the direct knock-on mechanism, with

an even higher number of ions occupying the SF than seen in previous simulations, aided by

reduced ion-ion repulsion due to electronic polarization (Fig. 7). Specifically, a highly conductive

K^+ channel can accommodate 3 or 4 ions in the SF. Furthermore, achieving balanced free energy

between the 3-ion and 4-ion states is crucial for facilitating low barrier permeation (Fig. 7).

Consequently, throughout all simulations conducted in this study, virtually no water molecules

cross any potassium channels; 3. A careful adjustment of ion LJ parameters is necessary to recover

correct ion radii and solvation free energies, which, as we showed, is critical for correct K^+/Na.

selectivity in K. channels. We therefore postulate, based on our and other works[4,16], that the

conductive state of the SF is characterized by near-full occupancy of K^+ in many conditions.

In our simulations, the best overall performing force field is CHARMM36m at a scaling factor of

0.78 with radius-corrected LJ parameters for ECC ions. This force field correctly reproduces I-V

curves in all three channels, as well as their K^+/Na^+ selectivity and inward rectification in the MthK

channel. We would therefore recommend it for simulations of other potassium channels, however

slight adjustments of the scaling factor might result in a better performance for a specific channel.

Our newly developed parameters allow for MD simulations of K$^+$ channels at an unprecedented
level of accuracy. As almost any process involving K. channels is related to the SF behavior and
its occupancy by K$^+$ ions, we envision future simulation studies to benefit from these findings.
Computational insights into the mechanisms of, for example, current regulation, selectivity filter
gating or channel modulation by small molecules are now accessible at a much higher accuracy,
especially as current readout is now readily accessible at unprecedented accuracy.

408 4. Methods

409 4.1. System preparation

410 4.1.1. NaK2K

The crystal structure 3ouf[1] was used and oriented in the membrane using OPM.[35] 8 water
molecules present in the crystal structure, behind the SF of NaK2K, were preserved. The current-
increasing F92A mutation[1,36], ACE and NME termini patching, membrane and solvation box
building were performed using Charmm-GUI,[37,38] webserver. The final system consisted of one
tetrameric channel, 168 POPC molecules, 11,134 water molecules, 160 K$^+$ ions, 152 Cl$^-$ ions,
corresponding to 0.8 m/L salt concentration.

417 4.1.2. TRAAK

The crystal structure 4wfe[39] from OPM was used for TRAAK. 6 water behind the SF was preserved.
The I159C and R284C mutations were introduced based on experimental results indicating that
this disulfide bond can prevent the channel from closing.[39] In the previously reported WT
simulation, a lipid can enter the pore from this fenestration site and possibly close the channel.[8]
246 POPC molecules were added using Charmm-GUI.[37,38] ACE and NME were added to cap the
termini. The final system consisted of one channel, 246 POPC molecules, 27,973 water molecules,
403 K$^+$ ions, 389 Cl$^-$ ions, corresponding to 0.8 m/L salt concentration.

4.1.3. MthK

For MthK the 3ldc[33] crystal structure was used. This structure was modified using the PyMOL mutagenesis tool to perform H68S and C77V mutations to match the WT amino acid sequence.[40] This structure was aligned using OPM. The Charmm-GUI Bilayer Builder was used to add ACE and NME capping groups and to create the simulation box. Crystal water and potassium ions were kept during this procedure. The final system contained 169 POPC molecules and 244 K^+ ions, 232 Cl^- and 16,900 water molecules, corresponding to 0.8 m/L salt concentration.

4.1.4. Kv1.2-2.1

The crystal structure of the Kv1.2-2.1 paddle chimera (2r9r[41]) from OPM was used. 8 water behind SF was preserved. The pore and voltage-sensing domains (E144 to T417) were embedded in a POPC membrane using Charmm-GUI. The functionally nonessential[42] tetramerization domain (T1) and the regulatory β subunit were omitted. ACE and NME capping were added in Charmm-GUI. The final system contained 313 POPC molecules and 460 K^+ ions, 452 Cl^- and 31,908 water molecules, corresponding to 0.8 m/L salt concentration.

4.1.5. KcsA

A pre-equilibrated structure of the E71A mutant (that removes C-type inactivation), including 17 water molecules behind and on top of the SF, based on the 5vk6[43] crystal structure as used in previous studies[44,45] was embedded in a POPC membrane using Charmm-GUI. ACE and NME capping groups were used for the N- and C-terminus respectively. GLU118 and GLU120 were protonated to mimic the state at pH 4. The final system contained 153 POPC molecules, 234 K^+ ions, 250 Cl^- and 16,196 water molecules, corresponding to 0.8 m/L salt concentration.

4.2.   Force field-specific parameters

4.2.1.   Charmm36m

For the Charmm36m[46] force field, Charmm lipid[47], Charmm TIP3P[48], and Charmm ions[49] were
used. A cutoff of 1.2 nm was used for van der Waals (vdW) interactions, and the forces were
switched smoothly to zero between 1.0 to 1.2 nm. The particle mesh Ewald (PME) method with a
1.2 nm cutoff was used for long-range electrostatic interactions.

4.2.2.   Amber14sb

For the Amber14sb[50] force field, lipid21[51], TIP3P water[52], and Joung and Cheatham ion
parameters[53] were used. A cutoff of 1.0 nm was used for van der Waals (vdW) interactions, and a
long-range analytical dispersion correction was applied to the energy and pressure. The PME
method with a 1.0 nm cutoff was used for long-range electrostatic interactions.

4.2.3.   Rescaled force fields

In one simulation box, the partial charges for all the charged residues including ions and charged
amino acids were adjusted with the same scaling factor, resulting in overall neutral simulation
systems. The partial charges for the charged amino acids were adjusted as follows. In Charmm36m
the partial charges of the charge group contributing to the net charge were multiplied by the
selected scaling factor. For example, in ASP the -CH_2-CO_2 group has a charge of -1. When charge
scaling is applied to ASP, the charge of the -CH_2-CO_2 group is scaled with the scaling factor (Fig.
S21). This approach could not be used in Amber, because the partial charges are also distributed
over the backbone atoms. Therefore, the scaled charges were obtained by taking a weighted
average (scaling factor as the weight) over the protonated and deprotonated charges (Fig. S21).
The deprotonated arginine, that was not available in the original amber14sb distribution, was
optimized at MP2/6-311G(d) using Gaussian 09 and RESP charge was fitted to HF/6-31G(d) with
antechamber.[54,55] All the partial charges used in this work at every scaling factors are provided in
the github repository:

https://github.com/deGrootLab/Charge_Scaling_in_Potassium_Channel_Simulations_paper/

4.2.4. Radius corrected vdW for charge scaled ion

The radius corrected vdW for the charge scaled ion was obtained and validated in the following 3 steps.

Step 1: Sigma optimization with grid search.

At each 0.05 charge from 1.00 (1.00, 0.95, 0.90 …) the sigma parameter of the ion decreased to the point when the radius of the first solvation shell reproduced the original radius in 1.00 charge. This sigma optimization was done by a grid search (a grid with (sigma, charge) pair). The setting and results of this grid search can be found in Fig. S8, and Fig. S9. The first peak in the ion-water oxygen RDF was used as the radius of the first solvation shell. Epsilon was kept the same as the original paper in the respective force field.[49,53] The sigma parameter between 0.05 charge (0.99, 0.98, 0.97, 0.96, 0.94 …) was linearly interpolated. The optimized vdW for K^+, Na^+, Cl^- is listed in Table S5.

In this step a simulation box with 2163 water molecules and 1 ion was used. Each (sigma, charge) pair on the grid was simulated for 40 ns, and trajectory was saved every 0.2 ps. Each grid (K^+, Na^+, Cl^-) was run in a Hamiltonian replica exchange simulation, and an exchange is attempted every 10 ps. RDF between ion and water oxygen is computed with "gmx rdf", using a bin width of 0.002 nm. The highest 7 points in the RDF were fitted with a Gaussian function (illustrated in Fig. S11). The same non-bonded settings for the corresponding force fields mentioned in Sections 4.2.1 and 4.2.2 were used.

Step 2: Solvation free energy validation

The ion solvation free energy calculation was performed using Gromacs 2022.5. The non-bonded parameters were set the same as previously mentioned in 4.2.1 and 4.2.2. Temperature was controlled by langevin integrator at 298 K and pressure was set at 1 bar using Parrinello-Rahman. The Coulomb interaction was gradually turned off in 10 lambda windows (0.0, 0.1, 0.2, 0.32, 0.44,

0.56, 0.68, 0.8, 0.9, 1.0, window 1-10), and vdw interaction was then turned off in the next 11

windows (0.0, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, window 10-20). HRE was used to speed

up the sampling with an exchange attempt every 100 steps (0.2 ps). Each lambda window was

simulated for 3 ns. The $\partial H/\partial \lambda$ and energy were saved every 2 ps. At least 1000 uncorrelated

samples were recorded in each lambda window. "Alchemical-analysis" tool[56] was used and the

reported value was estimated by the Multistate Bennett Acceptance Ratio (MBAR) method. A

soft-core potential[57] was employed for the van der Waals interactions. The solvation free energy

is shown in Fig. 4 and Fig. S10.

Step 3: RDF validation in dilute solution and 4M solution.

The RDF was validated under two conditions: (1) an infinitely diluted system, where the same

simulation box as in step 1 was used, and (2) a 4M KCl solution, where the concentration matched

that of the experiment.[58]

In the infinitely diluted system, each condition (charge with optimized sigma) was simulated for

100 ns and the trajectory is saved every 2 ps. The highest 7 points in the RDF were fitted with a

Gaussian function (Fig. S11, Fig. S12). The radius is shown in Fig. 4 and Fig. S10.

A 4M system was constructed to compare with the neutron scattering experiment. The simulation

box consisted of 6661 water, 120 K^+ ions, and 120 Cl^- ions. Each condition (charge with optimized

sigma) was simulated for 20 ns, and the trajectory was saved every 1 ps. The experimental RDF

was provided by the authors of the publication.[58] The RDF comparison is shown in Fig. S13.

All the initial structures, topology, and MD parameter files are provided in the github repository:

https://github.com/deGrootLab/Charge_Scaling_in_Potassium_Channel_Simulations_paper/

4.3. MD simulation

4.3.1. Equilibration and general setting

Gromacs 2022.5 was used except for the charge-scaled HRE.[59,60] The initial structure was minimized using the steepest descent until the maximum force was lower than 1000 kJ/(mol*nm). 100 ps of NVT with a harmonic restraint of 1000 kJ/(mol*nm^2) on all heavy atoms of protein and POPC. The temperature in this equilibration was controlled by either the Berendsen (Charmm36m) or v-rescale (Amber14sb) thermostat at 310 K. An additional 110 ns NPT equilibration was performed to relax the membrane with harmonic restraints of 1000 kJ/(mol*nm^2) on all heavy atoms of the protein. The temperature was controlled by either the Berendsen (Charmm36m) or V-rescale (Amber14sb) thermostat at 310 K. The semi-isotropic pressure coupling was controlled by the Berendsen barostat at 1 bar. The averaged box z dimension in the last 100 ns was used to set the electric field. In the later production run, we switched to the Nose-Hoover (for Chamm36m) or V-rescale (for Amber14sb) thermostat and the Parrinello-Rahman (for both Charmm36m and Amber14sb) barostat. A time step of 2 fs was used, and bonds with Hydrogen atoms were constrained by LINCS.[61]

4.3.2. Non-equilibrium simulation under voltage

To simulate the channel under voltage, an electric field is applied.[62] In Charmm36m and Amber14sb, the simulation was started after the equilibration, while in Amber14sb-S3, they were started from the ensemble generated by HRE with randomly generated velocity at 310 K. For every condition (including channel, force field, scaling factors, and voltage), 10 replicas of 500 ns were performed.

When counting the K$^+$ current crossing the channel, we assume that 1 K$^+$ ion contributes to 1 elementary charge even if the partial charge is less than 1. The scaling of the partial charge is a mean field treatment to approximate the polarization of the electron cloud between the nuclei, thus the electric field (voltage) in the simulation is set the same for the 1.00 charge force field and a charge scaled force field, and transporting 1 K$^+$ ion would generate the current of 1 elementary charge regardless of the parameter.

4.3.3. Hamiltonian replica exchange (HRE)

The HRE simulations were performed on Gromacs 2021.7 with plumed 2.8.2.[63–65] Although Gromacs has its own HRE implementation, it only works with the free energy code and can only interpolate states using the lambda definition. The actual interpolation can be found in Gromacs documentation (https://manual.gromacs.org/2022.5/reference-manual/functions/free-energy-interactions.html).[59,60] In the Plumed Hamiltonian Replica Exchange (HRE) setup, each replica's Hamiltonian can be defined independently using separate topology files. We opted for Plumed HRE because this implementation allows every replica to have physically meaningful parameters, enabling us to assign distinct charges to each replica. Replica exchange is meaningful only in equilibrium conditions, so no voltage is applied. 36 replicas were set up with charge scaling factors decreasing from 1.00 to 0.65 in steps of 0.01, using the same simulation boxes as mentioned in the system preparation section. Every 100 MD steps (0.2 ps), an exchange attempt was tested. Each replica was simulated for 80 ns and the trajectory was saved every 20 ps. In our simulation practice, we found the HRE approach to be an efficient method to probe the scaling factor space. The conductance-scaling bar plot in Fig. 2(a) was obtained from 40 μs simulations with a scaling factor from 1.00 to 0.65 and a step of 0.05 (8 scaling factors, 10 replicas per scale, 500 ns every replica), while the SF state population plot in Fig. 2(d) was obtained from 2.88 μs simulation on the same simulation box with a scaling factor from 1.00 to 0.65 and a step of 0.01 (36 replicas, 80 ns each). Using HRE, with a fraction of the computation power, the high conductance parameters can be quickly located with fine resolution.

4.3.4. Comparison of solvation free energy with experiment

The solvation free energy calculated from the scaled ion cannot be directly compared with the experiment, because part of the energy from electric polarization (E_{ele}) is removed from the force field (E_{ff}) and treated in the mean field approximation. (Eq. 1) The actual solvation free energy can be splitted into 2 terms, ΔG_{nuc} which can be computed from MD solvation free energy calculation, and ΔG_{ele} which comes from the mean field approximation.

$$E_{total} = E_{ff} + E_{ele} \qquad \text{(Eq. 1)}$$

$$\Delta G_{Exp} = \Delta G_{nuc} + \Delta G_{ele} \qquad \text{(Eq. 2)}$$

591 Assuming the approximated medium between MD particles has a relative dielectric constant of ε_r,

592 the ΔG_{ele} can be estimated with Born model (Eq. 3)

$$\Delta G = -\frac{N_A z^2 e^2}{8\pi\varepsilon_0 r_0}\left(1 - \frac{1}{\varepsilon_r}\right) \qquad \text{(Eq. 3)}$$

N_A	Avogadro constant
z	charge of ion
e	elementary charge
ε_0	dielectric constant of vacuum
r_0	effective radius of ion in Born model

595 If we take scaling factor q as $1/\sqrt{\varepsilon_r}$, Eq. 3 becomes Eq. 4.

$$\Delta G_{ele} = -\frac{N_a z^2 e^2}{8\pi\varepsilon_0 r_0}(1 - q^2) \qquad \text{(Eq. 4)}$$

596 The relative dielectric constant of water is significantly greater than 1 (ε_r for water = 78.4). We

597 approximate the experimental solvation free energy using the Born model, as shown in Eq. 5.

$$\Delta G_{Exp} = -\frac{N_a z^2 e^2}{8\pi\varepsilon_0 r_0}\left(1 - \frac{1}{\varepsilon_{r\_water}}\right) \approx -\frac{N_a z^2 e^2}{8\pi\varepsilon_0 r_0} \qquad \text{(Eq. 5)}$$

599 So that the MD simulated solvation free energy should target the rest of the solvation free energy,

600 which is Eq. 6.

$$\Delta G_{MD} = \Delta G_{Exp} - \Delta G_{ele} \qquad \text{(Eq. 6)}$$

$$\approx -\frac{N_a z^2 e^2}{8\pi\varepsilon_0 r_0} + \frac{N_a z^2 e^2}{8\pi\varepsilon_0 r_0}(1 - q^2)$$

$$= \Delta G_{Exp} * q^2$$

The comparing of experimental solvation free energy with ECC ion was discussed in several other

references.[24,66]

4.4. Mechanism Graphs

The simulation was saved every 0.2 ps. The trajectory was discretized by the SF state. The rate

between two states was defined as the inverse mean first passage time. All of the rates were first

listed (every state pair, forward and backward), and the mechanism graph was progressively

lumped from the state pair which had the fastest transition. If at least one of the forward or

backward rates between a pair of states exceeded the cutoff rate, this pair of states would be lumped

together. The cutoff rate when the lumping stopped is listed under each graph.

5. Acknowledgments

We would like to thank Youxing Jiang, Thomas Boulin, Dawon Kang, and Brad Rothberg for

kindly sharing the experimental IV curves of NaK2K[1], TRAAK[30], and MthK[10]. We also thank

Carter J. Wilson for providing the RESP charge of deprotonated arginine and Andrei Mironenko

for providing the KcsA structure. R.d.V., W.K. and B.L.d.G. acknowledge funding from the

Leibniz Collaborative Excellence Project "Ion Selectivity and Conduction Mechanism of Cation

Channels" K305/2020. W.K and B.L.d.G. acknowledge funding from the German Research
Foundation (DFG) through FOR2518 'Dynion', Project P5.

6. Author contributions

622 W.K. and B.L.d.G. conceived and supervised the project. C.H. and R.d.V. performed all
623 computational work and analyzed the data. C.H., R.d.V., W.K. and B.L.d.G. wrote the manuscript.

7. References

(1) Derebe, M. G.; Sauer, D. B.; Zeng, W.; Alam, A.; Shi, N.; Jiang, Y. Tuning the Ion Selectivity
of Tetrameric Cation Channels by Changing the Number of Ion Binding Sites. *Proc. Natl.*
Acad. Sci. **2011**, *108* (2), 598–602. https://doi.org/10.1073/pnas.1013636108.
(2) Zhou, Y.; Morais-Cabral, J. H.; Kaufman, A.; MacKinnon, R. Chemistry of Ion Coordination
and Hydration Revealed by a K+ Channel–Fab Complex at 2.0 Å Resolution. *Nature* **2001**,
414 (6859), 43–48. https://doi.org/10.1038/35102009.
(3) Morais-Cabral, J. H.; Zhou, Y.; MacKinnon, R. Energetic Optimization of Ion Conduction
Rate by the K+ Selectivity Filter. *Nature* **2001**, *414* (6859), 37–42.
https://doi.org/10.1038/35102000.
(4) Langan, P. S.; Vandavasi, V. G.; Weiss, K. L.; Afonine, P. V.; el Omari, K.; Duman, R.;
Wagner, A.; Coates, L. Anomalous X-Ray Diffraction Studies of Ion Transport in K+
Channels. *Nat. Commun.* **2018**, *9* (1), 4540. https://doi.org/10.1038/s41467-018-06957-w.
(5) Boiteux, C.; Posson, D. J.; Allen, T. W.; Nimigean, C. M. Selectivity Filter Ion Binding
Affinity Determines Inactivation in a Potassium Channel. *Proc. Natl. Acad. Sci.* **2020**, *117*
(47), 29968–29978. https://doi.org/doi:10.1073/pnas.2009624117.
(6) Köpfer, D. A.; Song, C.; Gruene, T.; Sheldrick, G. M.; Zachariae, U.; De Groot, B. L. Ion
Permeation in K $^+$ Channels Occurs by Direct Coulomb Knock-On. *Science* **2014**, *346* (6207),
352–355. https://doi.org/10.1126/science.1254840.
(7) Kopec, W.; Köpfer, D. A.; Vickery, O. N.; Bondarenko, A. S.; Jansen, T. L. C.; de Groot, B.
L.; Zachariae, U. Direct Knock-on of Desolvated Ions Governs Strict Ion Selectivity in K+
Channels. *Nat. Chem.* **2018**, *10* (8), 813–820. https://doi.org/10.1038/s41557-018-0105-9.
(8) Ocello, R.; Furini, S.; Lugli, F.; Recanatini, M.; Domene, C.; Masetti, M. Conduction and
Gating Properties of the TRAAK Channel from Molecular Dynamics Simulations with
Different Force Fields. *J. Chem. Inf. Model.* **2020**, *60* (12), 6532–6543.
https://doi.org/10.1021/acs.jcim.0c01179.
(9) Lam, C. K.; De Groot, B. L. Ion Conduction Mechanisms in Potassium Channels Revealed
by Permeation Cycles. *J. Chem. Theory Comput.* **2023**, *19* (9), 2574–2589.
https://doi.org/10.1021/acs.jctc.3c00061.

(10) Kopec, W.; Thomson, A. S.; De Groot, B. L.; Rothberg, B. S. Interactions between Selectivity Filter and Pore Helix Control Filter Gating in the MthK Channel. *J. Gen. Physiol.* **2023**, *155* (8), e202213166. https://doi.org/10.1085/jgp.202213166.

(11) Mironenko, A.; Zachariae, U.; de Groot, B. L.; Kopec, W. The Persistent Question of Potassium Channel Permeation Mechanisms. *Ion Channels Intersect. Struct. Funct. Pharmacol.* **2021**, *433* (17), 167002. https://doi.org/10.1016/j.jmb.2021.167002.

(12) Jensen, M. Ø.; Borhani, D. W.; Lindorff-Larsen, K.; Maragakis, P.; Jogini, V.; Eastwood, M. P.; Dror, R. O.; Shaw, D. E. Principles of Conduction and Hydrophobic Gating in K$^+$ Channels. *Proc. Natl. Acad. Sci.* **2010**, *107* (13), 5833–5838. https://doi.org/10.1073/pnas.0911691107.

(13) Domene, C.; Ocello, R.; Masetti, M.; Furini, S. Ion Conduction Mechanism as a Fingerprint of Potassium Channels. *J. Am. Chem. Soc.* **2021**, *143* (31), 12181–12193. https://doi.org/10.1021/jacs.1c04802.

(14) Brennecke, J. T.; de Groot, B. L. Mechanism of Mechanosensitive Gating of the TREK-2 Potassium Channel. *Biophys. J.* **2018**, *114* (6), 1336–1343. https://doi.org/10.1016/j.bpj.2018.01.030.

(15) Guidoni, L.; Carloni, P. Potassium Permeation through the KcsA Channel: A Density Functional Study. *Biochim. Biophys. Acta BBA - Biomembr.* **2002**, *1563* (1–2), 1–6. https://doi.org/10.1016/S0005-2736(02)00349-8.

(16) Jing, Z.; Rackers, J. A.; Pratt, L. R.; Liu, C.; Rempe, S. B.; Ren, P. Thermodynamics of Ion Binding and Occupancy in Potassium Channels. *Chem. Sci.* **2021**, *12* (25), 8920–8930. https://doi.org/10.1039/D1SC01887F.

(17) Kirby, B. J.; Jungwirth, P. Charge Scaling Manifesto: A Way of Reconciling the Inherently Macroscopic and Microscopic Natures of Molecular Simulations. *J. Phys. Chem. Lett.* **2019**, *10* (23), 7531–7536. https://doi.org/10.1021/acs.jpclett.9b02652.

(18) E. Duboué-Dijon; M. Javanainen; P. Delcroix; P. Jungwirth; H. Martinez-Seara. A Practical Guide to Biologically Relevant Molecular Simulations with Charge Scaling for Electronic Polarization. *J. Chem. Phys.* **2020**, *153* (5), 050901. https://doi.org/10.1063/5.0017775.

(19) Melcr, J.; Ferreira, T. M.; Jungwirth, P.; Ollila, O. H. S. Improved Cation Binding to Lipid Bilayers with Negatively Charged POPS by Effective Inclusion of Electronic Polarization. *J. Chem. Theory Comput.* **2020**, *16* (1), 738–748. https://doi.org/10.1021/acs.jctc.9b00824.

(20) Melcr, J.; Martinez-Seara, H.; Nencini, R.; Kolafa, J.; Jungwirth, P.; Ollila, O. H. S. Accurate Binding of Sodium and Calcium to a POPC Bilayer by Effective Inclusion of Electronic Polarization. *J. Phys. Chem. B* **2018**, *122* (16), 4546–4557. https://doi.org/10.1021/acs.jpcb.7b12510.

(21) Nencini, R.; Tempra, C.; Biriukov, D.; Riopedre-Fernandez, M.; Cruces Chamorro, V.; Polák, J.; Mason, P. E.; Ondo, D.; Heyda, J.; Ollila, O. H. S.; Jungwirth, P.; Javanainen, M.; Martinez-Seara, H. Effective Inclusion of Electronic Polarization Improves the Description of Electrostatic Interactions: The prosECCo75 Biomolecular Force Field. *J. Chem. Theory Comput.* **2024**, *20* (17), 7546–7559. https://doi.org/10.1021/acs.jctc.4c00743.

(22) Laage, D.; Stirnemann, G. Effect of Ions on Water Dynamics in Dilute and Concentrated Aqueous Salt Solutions. *J. Phys. Chem. B* **2019**, *123* (15), 3312–3324. https://doi.org/10.1021/acs.jpcb.9b01053.

(23) Wilson, C. J.; Gapsys, V.; De Groot, B. L. Improving pKa Prediction Accuracy with Reparametrized Force Fields and Free Energy Calculations. January 9, 2025. https://doi.org/10.26434/chemrxiv-2024-btnq5-v2.

(24) Leontyev, I. V.; Stuchebrukhov, A. A. Electronic Continuum Model for Molecular Dynamics Simulations of Biological Molecules. *J. Chem. Theory Comput.* **2010**, *6* (5), 1498–1508. https://doi.org/10.1021/ct9005807.

(25) Lee, B.; White, K. I.; Socolich, M.; Klureza, M. A.; Henning, R.; Srajer, V.; Ranganathan, R.; Hekstra, D. R. Direct Visualization of Electric-Field-Stimulated Ion Conduction in a Potassium Channel. *Cell* **2025**, *188* (1), 77-88.e15. https://doi.org/10.1016/j.cell.2024.12.006.

(26) Bernèche, S.; Roux, B. Energetics of Ion Conduction through the K+ Channel. *Nature* **2001**, *414* (6859), 73–77. https://doi.org/10.1038/35102067.

(27) Jensen, M. Ø.; Jogini, V.; Eastwood, M. P.; Shaw, D. E. Atomic-Level Simulation of Current–Voltage Relationships in Single-File Ion Channels. *J. Gen. Physiol.* **2013**, *141* (5), 619–632. https://doi.org/10.1085/jgp.201210820.

(28) Abdul Kadir, L.; Stacey, M.; Barrett-Jolley, R. Emerging Roles of the Membrane Potential: Action Beyond the Action Potential. *Front. Physiol.* **2018**, *9*, 1661. https://doi.org/10.3389/fphys.2018.01661.

(29) Lazzari-Dean, J. R.; Gest, A. M. M.; Miller, E. W. Measuring Absolute Membrane Potential Across Space and Time. *Annu. Rev. Biophys.* **2021**, *50* (1), 447–468. https://doi.org/10.1146/annurev-biophys-062920-063555.

(30) Ben Soussia, I.; El Mouridi, S.; Kang, D.; Leclercq-Blondel, A.; Khoubza, L.; Tardy, P.; Zariohi, N.; Gendrel, M.; Lesage, F.; Kim, E.-J.; Bichet, D.; Andrini, O.; Boulin, T. Mutation of a Single Residue Promotes Gating of Vertebrate and Invertebrate Two-Pore Domain Potassium Channels. *Nat. Commun.* **2019**, *10* (1), 787. https://doi.org/10.1038/s41467-019-08710-3.

(31) Miranda, W. E.; DeMarco, K. R.; Guo, J.; Duff, H. J.; Vorobyov, I.; Clancy, C. E.; Noskov, S. Yu. Selectivity Filter Modalities and Rapid Inactivation of the hERG1 Channel. *Proc. Natl. Acad. Sci.* **2020**, *117* (6), 2795–2804. https://doi.org/10.1073/pnas.1909196117.

(32) Sumikama, T.; Oiki, S. Digitalized K^+ Occupancy in the Nanocavity Holds and Releases Queues of K^+ in a Channel. *J. Am. Chem. Soc.* **2016**, *138* (32), 10284–10292. https://doi.org/10.1021/jacs.6b05270.

(33) Ye, S.; Li, Y.; Jiang, Y. Novel Insights into K+ Selectivity from High-Resolution Structures of an Open K+ Channel Pore. *Nat. Struct. Mol. Biol.* **2010**, *17* (8), 1019–1023. https://doi.org/10.1038/nsmb.1865.

(34) Atkins, P.; De Paula, J.; Keeler, J. *Atkins' Physical Chemistry*, 11th ed.; Oxford University Press: London, England, 2017.

(35) Lomize, M. A.; Pogozheva, I. D.; Joo, H.; Mosberg, H. I.; Lomize, A. L. OPM Database and PPM Web Server: Resources for Positioning of Proteins in Membranes. *Nucleic Acids Res.* **2012**, *40* (D1), D370–D376. https://doi.org/10.1093/nar/gkr703.

(36) Sauer, D. B.; Zeng, W.; Canty, J.; Lam, Y.; Jiang, Y. Sodium and Potassium Competition in Potassium-Selective and Non-Selective Channels. *Nat. Commun.* **2013**, *4* (1), 2721. https://doi.org/10.1038/ncomms3721.

(37) Jo, S.; Lim, J. B.; Klauda, J. B.; Im, W. CHARMM-GUI Membrane Builder for Mixed Bilayers and Its Application to Yeast Membranes. *Biophys. J.* **2009**, *97* (1), 50–58. https://doi.org/10.1016/j.bpj.2009.04.013.

(38) Jo, S.; Kim, T.; Iyer, V. G.; Im, W. CHARMM-GUI: A Web-based Graphical User Interface for CHARMM. *J. Comput. Chem.* **2008**, *29* (11), 1859–1865. https://doi.org/10.1002/jcc.20945.

(39) Brohawn, S. G.; Campbell, E. B.; MacKinnon, R. Physical Mechanism for Gating and
Mechanosensitivity of the Human TRAAK K+ Channel. *Nature* **2014**, *516* (7529), 126–130.
https://doi.org/10.1038/nature14013.
(40) The PyMOL Molecular Graphics System, Version 2.5.4 Schrödinger, LLC.
(41) Long, S. B.; Tao, X.; Campbell, E. B.; MacKinnon, R. Atomic Structure of a Voltage-
Dependent K+ Channel in a Lipid Membrane-like Environment. *Nature* **2007**, *450* (7168),
376–382. https://doi.org/10.1038/nature06265.
(42) Miller, C.; Kobertz, W. R. K+ Channels Lacking the "tetramerization" Domain: Implications
for Pore Structure. *Nat. Struct. Biol.* **1999**, *6* (12), 1122–1125. https://doi.org/10.1038/70061.
(43) Cuello, L. G.; Cortes, D. M.; Perozo, E. The Gating Cycle of a K+ Channel at Atomic
Resolution. *eLife* **2017**, *6*, e28032. https://doi.org/10.7554/eLife.28032.
(44) Mironenko, A.; De Groot, B. L.; Kopec, W. Selectivity Filter Mutations Shift Ion Permeation
Mechanism in Potassium Channels. *PNAS Nexus* **2024**, *3* (7), pgae272.
https://doi.org/10.1093/pnasnexus/pgae272.
(45) Pérez-Conesa, S.; Delemotte, L. Free Energy Landscapes of KcsA Inactivation. July 4, 2023.
https://doi.org/10.7554/eLife.88403.1.
(46) Huang, J.; Rauscher, S.; Nawrocki, G.; Ran, T.; Feig, M.; de Groot, B. L.; Grubmüller, H.;
MacKerell, A. D. CHARMM36m: An Improved Force Field for Folded and Intrinsically
Disordered Proteins. *Nat. Methods* **2017**, *14* (1), 71–73. https://doi.org/10.1038/nmeth.4067.
(47) Klauda, J. B.; Venable, R. M.; Freites, J. A.; O'Connor, J. W.; Tobias, D. J.; Mondragon-
Ramirez, C.; Vorobyov, I.; MacKerell, A. D.; Pastor, R. W. Update of the CHARMM All-
Atom Additive Force Field for Lipids: Validation on Six Lipid Types. *J. Phys. Chem. B* **2010**,
114 (23), 7830–7843. https://doi.org/10.1021/jp101759q.
(48) MacKerell, A. D.; Bashford, D.; Bellott, M.; Dunbrack, R. L.; Evanseck, J. D.; Field, M. J.;
Fischer, S.; Gao, J.; Guo, H.; Ha, S.; Joseph-McCarthy, D.; Kuchnir, L.; Kuczera, K.; Lau, F.
T. K.; Mattos, C.; Michnick, S.; Ngo, T.; Nguyen, D. T.; Prodhom, B.; Reiher, W. E.; Roux,
B.; Schlenkrich, M.; Smith, J. C.; Stote, R.; Straub, J.; Watanabe, M.; Wiórkiewicz-Kuczera,
J.; Yin, D.; Karplus, M. All-Atom Empirical Potential for Molecular Modeling and Dynamics
Studies of Proteins. *J. Phys. Chem. B* **1998**, *102* (18), 3586–3616.
https://doi.org/10.1021/jp973084f.
(49) Beglov, D.; Roux, B. Finite Representation of an Infinite Bulk System: Solvent Boundary
Potential for Computer Simulations. *J. Chem. Phys.* **1994**, *100* (12), 9050–9063.
https://doi.org/10.1063/1.466711.
(50) Maier, J. A.; Martinez, C.; Kasavajhala, K.; Wickstrom, L.; Hauser, K. E.; Simmerling, C.
ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from
ff99SB. *J. Chem. Theory Comput.* **2015**, *11* (8), 3696–3713.
https://doi.org/10.1021/acs.jctc.5b00255.
(51) Dickson, C. J.; Walker, R. C.; Gould, I. R. Lipid21: Complex Lipid Membrane Simulations
with AMBER. *J. Chem. Theory Comput.* **2022**, *18* (3), 1726–1736.
https://doi.org/10.1021/acs.jctc.1c01217.
(52) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison
of Simple Potential Functions for Simulating Liquid Water. *J. Chem. Phys.* **1983**, *79* (2),
926–935. https://doi.org/10.1063/1.445869.
(53) Joung, I. S.; Cheatham, T. E. Determination of Alkali and Halide Monovalent Ion Parameters
for Use in Explicitly Solvated Biomolecular Simulations. *J. Phys. Chem. B* **2008**, *112* (30),
9020–9041. https://doi.org/10.1021/jp8001614.

(54) Wang, J.; Wang, W.; Kollman, P. A.; Case, D. A. Automatic Atom Type and Bond Type
Perception in Molecular Mechanical Calculations. *J. Mol. Graph. Model.* **2006**, *25* (2), 247–
260. https://doi.org/10.1016/j.jmgm.2005.12.005.
(55) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J.
R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.;
Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.;
Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao,
O.; Nakai, H.; Vreven, T.; Montgomery, J. A.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd,
J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.;
Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.;
Millam, M. J.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.;
Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski,
J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.;
Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.;
Cioslowski, J.; Fox, D. J. Gaussian 09 Revision D.01.
(56) Klimovich, P. V.; Shirts, M. R.; Mobley, D. L. Guidelines for the Analysis of Free Energy
Calculations. *J. Comput. Aided Mol. Des.* **2015**, *29* (5), 397–411.
https://doi.org/10.1007/s10822-015-9840-9.
(57) Beutler, T. C.; Mark, A. E.; Van Schaik, R. C.; Gerber, P. R.; Van Gunsteren, W. F. Avoiding
Singularities and Numerical Instabilities in Free Energy Calculations Based on Molecular
Simulations. *Chem. Phys. Lett.* **1994**, *222* (6), 529–539. https://doi.org/10.1016/0009-
2614(94)00397-1.
(58) Mason, P. E.; Tavagnacco, L.; Saboungi, M.-L.; Hansen, T.; Fischer, H. E.; Neilson, G. W.;
Ichiye, T.; Brady, J. W. Molecular Dynamics and Neutron Scattering Studies of Potassium
Chloride in Aqueous Solution. *J. Phys. Chem. B* **2019**, *123* (50), 10807–10813.
https://doi.org/10.1021/acs.jpcb.9b08422.
(59) Bauer, P.; Hess, B.; Lindahl, E. GROMACS 2022.5 Manual. **2023**.
https://doi.org/10.5281/ZENODO.7586765.
(60) Abraham, M. J.; Murtola, T.; Schulz, R.; Páll, S.; Smith, J. C.; Hess, B.; Lindahl, E.
GROMACS: High Performance Molecular Simulations through Multi-Level Parallelism
from Laptops to Supercomputers. *SoftwareX* **2015**, *1–2*, 19–25.
https://doi.org/10.1016/j.softx.2015.06.001.
(61) Hess, B.; Bekker, H.; Berendsen, H. J. C.; Fraaije, J. G. E. M. LINCS: A Linear Constraint
Solver for Molecular Simulations. *J. Comput. Chem.* **1997**, *18* (12), 1463–1472.
https://doi.org/10.1002/(SICI)1096-987X(199709)18:12<1463::AID-JCC4>3.0.CO;2-H.
(62) Roux, B. The Membrane Potential and Its Representation by a Constant Electric Field in
Computer Simulations. *Biophys. J.* **2008**, *95* (9), 4205–4216.
https://doi.org/10.1529/biophysj.108.136499.
(63) The PLUMED consortium. Promoting Transparency and Reproducibility in Enhanced
Molecular Simulations. *Nat. Methods* **2019**, *16* (8), 670–673. https://doi.org/10.1038/s41592-
019-0506-8.
(64) Tribello, G. A.; Bonomi, M.; Branduardi, D.; Camilloni, C.; Bussi, G. PLUMED 2: New
Feathers for an Old Bird. *Comput. Phys. Commun.* **2014**, *185* (2), 604–613.
https://doi.org/10.1016/j.cpc.2013.09.018.
(65) Bonomi, M.; Branduardi, D.; Bussi, G.; Camilloni, C.; Provasi, D.; Raiteri, P.; Donadio, D.;
Marinelli, F.; Pietrucci, F.; Broglia, R. A.; Parrinello, M. PLUMED: A Portable Plugin for

Free-Energy Calculations with Molecular Dynamics. *Comput. Phys. Commun.* **2009**, *180* (10), 1961–1972. https://doi.org/10.1016/j.cpc.2009.05.011.

(66) Leontyev, I.; Stuchebrukhov, A. Accounting for Electronic Polarization in Non-Polarizable Force Fields. *Phys. Chem. Chem. Phys.* **2011**, *13* (7), 2613–2626. https://doi.org/10.1039/C0CP01971B.

(67) Marcus, Y. Ionic Radii in Aqueous Solutions. *Chem. Rev.* **1988**, *88* (8), 1475–1498. https://doi.org/10.1021/cr00090a003.

(68) Schmid, R.; Miah, A. M.; Sapunov, V. N. A New Table of the Thermodynamic Quantities of Ionic Hydration: Values and Some Applications (Enthalpy–Entropy Compensation and Born Radii). *Phys. Chem. Chem. Phys.* **2000**, *2* (1), 97–102. https://doi.org/10.1039/A907160A.

Figures

Figure 1. Main features of K[+] permeation through K[+] channels in MD simulations with traditional force fields. (a) Comparison of experimental conductances[1,10,30] and MD simulated conductances in 3

channels (NaK2K, TRAAK, MthK), with 2 force fields (Charmm36m and Amber14sb), at +100 mV. **(b)** A schematic mechanism of ion permeation in NaK2K in MD simulations using Charmm36m at +100 mV. Trajectories were discretized based on the number of K^+ ions in positions S0-SCav. The SF occupancy is represented using a 6-letter code showing the occupancy at binding sites S0 to SCav: 'K' for a K^+ ion, 'W' for water, and '0' for an unoccupied site. Populations of SF states are expressed as percentages, with WKK0K0, WK0KK0, KWK0K0, WKWK0K, and WK0K0K being the only states exceeding a 5% population.

Figure 2. Charge scaling has a major impact on ion permeation characteristics in the NaK2K channel.
(**a-c**) Average conductances at different scaling factors in MD simulations using Charmm36m, Amber14sb, and Amber14sb-S3 (see text for explanation) force fields under +150 mV in NaK2K. The error bar is the standard error of the mean (N=10). (**d-f**) Equilibrated populations of the SF states recorded in HRE in NaK2K, using a 6-letter code showing the occupancy at binding sites S0 to Scav: 'K' for a K^+ ion, 'W' for water, and '0' for an unoccupied site. For clarity, only the even replicas are plotted (replica 0, 2, 4 etc at scale 1.00, 0.98, 0.96 etc). The odd replicas are not shown (replica 1, 3, 5 etc at scale 0.99, 0.97, 0.95 etc). (**g-i**) Ion occupation state distribution from HRE simulation, the states with the same number of ions in the SF are summed together. In (d-i) The error bar is the 95% confidence interval when bootstrapping frames.

Figure 3. Simulated full current-voltage (I-V) curves for NaK2K, TRAAK, and MthK channels at different scaling factors. (a-c) Simulated I-V curves with Charmm36m. **(d-f)** Simulated I-V curve with Amber14sb-S3. MD simulations were performed at ± 50 mV, ± 100 mV, ± 150 mV, ± 200 mV (if the experiment data covered ± 200 mV). The experimental data are shown in black and were provided by the authors of respective publications.[1,10,30] Simulated data are shown in colors for different scaling factors. The error bar is the standard error of the mean in all cases. The digital data can be found in the SI file.

Figure 4. Comparison of the radius corrected LJ parameters for ECC ion in this work (solid line) and the original vdW (dotted line) in Charmm36m. (a) The radius of the first solvation shell for K^+ and Na^+ with two sets of LJ parameters, compared to the experimental value.[67] (b) The solvation free energy for K^+ and Na^+ with two sets of vdW parameters. The experimental solvation free energy from Schimdt et al.[68] was scaled at each scaling factor, following the Born model (see Methods for details). The experiment value was from the respective references. The vdW parameters at 1.0 scale were taken from the original force field. The same plot for Amber14sb is shown in Fig. S10.

Figure 5. K⁺ and Na⁺ conductance with two treatments of LJ parameters in Charmm36m. (a-c)
Simulated conductances with our newly developed radius-corrected LJ parameters for ECC ions. **(d-f)**
Simulated conductance with unmodified LJ parameters in Charmm36m. All simulations were performed
under +150 mV voltage. The analogous plot for Amber14-S3 is in Fig S14. The experimental values were
taken from the respective references.[1,33]

Table 1. K⁺/ Na⁺ conductance ratio at 0.78 scale, at +150 mV

	NaK2K	TRAAK	MthK
Charmm36m	3.3	2.5	16.1
Amber14sb-S3	2.5	7.7	8.4

Figure 6. The mechanism graph from simulations of MthK in Charmm36m with 0.78 scaling under +150 mV or -150 mV. 515 (a) and 970 (b) permeation events were observed in 500 ns x 10 replica simulations at each voltage, respectively. The numbers shown by the edges are the net flux. The thickness of the edge (arrow) is proportional to the rate (defined as the inverse of the mean first passage time). The size of a node corresponds to the overall population of all states in this node. Nodes with 4 K^+ ions in SF are shown in orange, and nodes with 3 K^+ ions in SF are shown in blue. In panel (a), states A and B are lumped together when rate exceeds 125.0 ns^{-1}, similarly, in panel (b), they are lumped when this product exceeds 180.0 ns^{-1}. The rate-limiting step is marked in red. The exact flux and rate between all state pairs in the rate limiting step can be found in Table S4. The state pair WKKK0K-WKK0KK contributes to more than 85% of the net flux, and is identified as the rate-limiting step without lumping.

Figure 7. Schematic energy landscape between 3 and 4 K⁺ states with/without polarization. Only 1 SF state is drawn for 3 and 4 K⁺ states, and in reality, multiple 3 K⁺ and multiple 4 K⁺ states are visited. When the ECC polarization is introduced, the dominating effect is a decreased ion-ion repulsion, leading to more ion-ion contacts and fully ion filled filters, as well as a reduced barrier to form such contacts, and thus a more efficient knock-on permeation.