1 *Salmonella* **exploits host- and bacterial-derived β-alanine for**

2 **replication inside host macrophages**

Shuai Ma[1,†], Bin Yang[1,†], Yuyang Sun[1], Xinyue Wang[1], Houliang Guo[1],

Ruiying Liu[1], Ting Ye[1], Chenbo Kang[1], Jingnan Chen[1], Lingyan Jiang[1,*]

[1]*National Key Laboratory of Intelligent Tracking and Forecasting for Infectious*

Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai

University, Tianjin 300457, China.

[†] Shuai Ma and Bin Yang contributed equally to this work.

[*]Correspondence and requests for materials should be addressed to Lingyan Jiang

(email: jianglingyan@nankai.edu.cn).

Abstract

Salmonella is a major foodborne pathogen that can effectively replicate inside

host macrophages to establish life-threatening systemic infections. *Salmonella* must

utilize diverse nutrients for growth in nutrient-poor macrophages, but which nutrients

are required for intracellular *Salmonella* growth is largely unknown. Here, we found

that either acquisition from the host or *de novo* synthesis of a nonprotein amino

acid, β-alanine, is critical for *Salmonella* replication inside macrophages. The

concentration of β-alanine is decreased in *Salmonella*-infected macrophages, while

the addition of exogenous β-alanine enhances *Salmonella* replication in macrophages,

suggesting that *Salmonella* can uptake host-derived β-alanine for intracellular growth.

Moreover, the expression of *panD,* the rate-limiting gene required for β-alanine

synthesis in *Salmonella,* is upregulated when *Salmonella* enters macrophages.

Mutation of *panD* impaired *Salmonella* replication in macrophages and colonization

in the mouse liver and spleen, indicating that *de novo* synthesis of β-alanine is

essential for intracellular *Salmonella* growth and systemic infection. Additionally, we

revealed that β-alanine influences *Salmonella* intracellular replication and *in vivo*

virulence partially by increasing expression of the zinc transporter genes *znuABC*,

which in turn facilitates the uptake of the essential micronutrient zinc by *Salmonella*.

Taken together, these findings highlight the important role of β-alanine in the

intracellular replication and virulence of *Salmonella*, and *panD* is a promising target

for controlling systemic *Salmonella* infection.

Keywords: *Salmonella*, replication in macrophage, virulence, β-alanine, zinc

uptake

Introduction

Salmonella is a major foodborne pathogen worldwide that can cause self-limiting gastroenteritis or life-threatening systemic disease in a wide range of animals (Fàbrega & Vila, 2013; Ohl & Miller, 2001). *Salmonella* infection remains a significant global public health concern. An estimated 93.8 million cases of gastroenteritis and 27 million cases of systemic diseases caused by *Salmonella* species occur annually worldwide, with 355,000 deaths (Kim et al., 2019; Majowicz et al., 2010). The ability to survive and replicate in host macrophages is a key determinant for *Salmonella* to induce systemic infection (Fields, Swanson, Haidaris, & Heffron, 1986; LaRock, Chaudhary, & Miller, 2015; Leung & Finlay, 1991). After internalization by macrophages, *Salmonella* delivers a set of more than 30 effector proteins to the macrophage cytoplasm, mainly through a type III secretion system (T3SS) encoded by *Salmonella* pathogenicity island-2 (SPI-2) (Pillay et al., 2023). SPI-2 effectors manipulate diverse cellular processes to promote the formation of a membrane-bound compartment, termed the *Salmonella*-containing vacuole (SCV), a niche where *Salmonella* resides and grows (Castanheira & García-Del Portillo, 2017; Rosenberg, Riquelme, Prince, & Avraham, 2022; Steeb et al., 2013). SCV protects *Salmonella* from contact with antimicrobial agents in macrophages (Figueira & Holden, 2012; Li et al., 2023). Moreover, SPI-2 effectors induce the formation of specific tubular membrane compartments that extend from the SCV, known as *Salmonella*-induced filaments (SIFs). These filaments allow *Salmonella* to access various types of endocytosed nutrients, thereby facilitating efficient replication

within macrophages (Brumell, Tang, Mills, & Finlay, 2001; Liss et al., 2017; Rajashekar, Liebl, Seitz, & Hensel, 2008).

As the SCV of macrophages is a nutrient-poor environment (Fields et al., 1986; Kehl, Noster, & Hensel, 2020), to effectively replicate in the SCV, *Salmonella* needs to acquire a wide range of host nutrients or host-derived metabolites and synthesize metabolites *de novo* that cannot be sufficiently accessed from the host (Dandekar et al., 2014; Röder, Felgner, & Hensel, 2021; Tuli & Sharma, 2019). Nutrients/metabolites are used by intracellular *Salmonella* either as carbon sources to generate energy or for the synthesis of fatty acids and proteins (Dandekar et al., 2014; Steeb et al., 2013). Moreover, several metabolites were found to be employed by *Salmonella* as environmental cues to induce the expression of virulence genes (Jiang et al., 2021; X. Wang et al., 2023). In recent years, an increasing number of studies have focused on the intracellular nutrition of *Salmonella* (Bumann & Schothorst, 2017; Liss et al., 2017; Röder et al., 2021); however, the nutrients that are required for *Salmonella* replication in macrophages remain largely unknown.

β-Alanine, also known as 3-aminopropionic acid (3-AP), is the only naturally occurring β-type amino acid and is found in all living organisms (Song, Zhang, Wang, Xu, & Wei, 2023). β-Alanine can be synthesized *de novo* by bacteria, fungi, and plants, whereas animals need to obtain it from food or generate it via the catabolism of cytosine and uracil (Gojković, Sandrini, & Piskur, 2001; L. Wang, Mao, Wang, Ma, & Chen, 2021). In bacteria, β-alanine is synthesized *via* the decarboxylation of L-aspartate, a reaction catalyzed by L-aspartate decarboxylase (PanD) (Begley,

Kinsland, & Strauss, 2001; Schmitzberger, Smith, Abell, & Blundell, 2003; West,

Traut, Shanley, & O'Donovan, 1985). The *panD* gene is conserved among most

bacteria (Nozaki, Webb, & Niki, 2012). Although β-alanine is a nonprotein amino

acid that is not incorporated into proteins, it has important physiological functions in

the metabolism of organisms (L. Wang et al., 2021; Yuan et al., 2022). First, β-alanine

forms a part of pantothenate (vitamin B5), which is the key precursor for the

biosynthesis of coenzyme A (CoA) (Webb, Smith, & Abell, 2004; White, Gunyuzlu,

& Toyn, 2001). CoA is an essential cofactor involved in many metabolic pathways,

including the synthesis and degradation of fatty acids, pyruvate oxidation through the

tricarboxylic acid (TCA) cycle, and the production of secondary metabolites

(Davaapil, Tsuchiya, & Gout, 2014; Gout, 2019; Sibon & Strauss, 2016; Theodoulou,

Sibon, Jackowski, & Gout, 2014). Second, β-alanine is a limiting precursor of

carnosine, a nonenzymatic free radical scavenger and a natural antioxidant, with

anti-inflammatory and neuroprotective effects in animals (Boldyrev, Aldini, & Derave,

2013; Hoffman, Varanoske, & Stout, 2018). In the past 15 years, β-alanine has

become one of the most commonly used sports supplements worldwide (Bellinger,

2014; Hoffman et al., 2018; Huerta Ojeda, Tapia Cerda, Poblete Salvatierra,

Barahona-Fuentes, & Jorquera Aguilera, 2020). Although both *Salmonella* and host

cells are capable of producing β-alanine, whether β-alanine contributes to the

pathogenicity and intracellular growth of *Salmonella* remains unknown.

In this work, using targeted metabolic profiling, *in vitro* and *in vivo* infection

assays, and many other molecular techniques, we demonstrated that the utilization of

β-alanine is essential for *Salmonella* replication in host macrophages and virulence in

mice. *Salmonella* acquires β-alanine both via the uptake of β-alanine from host

macrophages and the *de novo* synthesis of β-alanine. Further investigation revealed

the molecular mechanism underlying the contribution of β-alanine to *Salmonella*

intracellular replication and pathogenicity, wherein β-alanine promotes the expression

of zinc transporter genes to facilitate the uptake of the essential micronutrient zinc by

intracellular *Salmonella,* therefore promoting *Salmonella* replication in macrophages

and subsequent systemic infection. Taken together, these findings demonstrate a

correlation between *Salmonella* β-alanine utilization and zinc uptake during

intracellular infection and provide new insights into the intracellular nutrition of

Salmonella. The rate-limiting gene (*panD*) in the β-alanine synthesis pathway of

Salmonella might be a future target for the prevention and treatment of this pathogen.

Results

Host-derived β-alanine promotes *Salmonella* replication inside macrophages

To explore changes in the levels of different amino acids inside macrophages upon *Salmonella* infection, we performed targeted metabolomics analysis of mouse RAW264.7 macrophages that were mock-infected or infected with wild-type *Salmonella* (*Salmonella enterica* serovar Typhimurium ATCC 14028s, STM) for 8 h using liquid chromatography-tandem mass spectrometry (LC□MS/MS) (Figure 1A). Principal component analysis (PCA) demonstrated a clear separation between the mock- and *Salmonella*-infected groups (Figure 1B). A total of 26 free amino acids were analyzed, and 8 showed significant differences in abundance between the two groups (VIP (Variable Importance in the Projection) > 1 and $P < 0.05$; FC (fold change) >1.5 or<0.667) (Figure 1C，D). Compared with those in the mock-infected group, the concentrations of 3 amino acids (L-hydroxyproline, L-citrulline and L-cysteine) were upregulated (Figure 1C), and 5 amino acids (L-asparagine, L-serine, L-aspartate, β-alanine and γ-aminobutyric acid) were downregulated (Figure 1D) in the *Salmonella*-infected group. Consistent with previous findings, intracellular serine concentrations were downregulated due to the reprogramming of macrophage glucose metabolism during *Salmonella* infection (Jiang et al., 2021). *Salmonella* can use host-derived aspartate and asparagine for growth in macrophages (Popp et al., 2015); therefore, the decrease in intracellular aspartate and asparagine upon *Salmonella* infection is likely due to their utilization by bacteria. Interestingly, β-alanine concentrations were also downregulated in the *Salmonella*-infected group (Figure 1D),

suggesting that intracellular *Salmonella* may use host-derived β-alanine for growth.

To investigate whether host-derived β-alanine can promote intracellular

Salmonella replication, we added an additional 0.5, 1, 2, 4 mM β-alanine (Schneider,

Krämer, & Burkovski, 2004) to the culture medium (RPMI) of RAW264.7 cells and

then infected them with *Salmonella* to test the influence of β-alanine addition on the

ability of *Salmonella* to replicate in macrophages. The results showed that the

replication of *Salmonella* in RAW264.7 cells significantly ($P < 0.001$) increased with

the addition of 1, 2, or 4 mM β-alanine (Figure 1E). Furthermore, β-alanine enhanced

Salmonella intracellular replication in a dose-dependent manner (Figure 1E). The

results suggest that host-derived β-alanine facilitates *Salmonella* replication inside

macrophages. We then investigated whether β-alanine-mediated *Salmonella* growth

promotion is due to the changes in antimicrobial activity of the macrophages. We

observed that the addition of 1 mM β-alanine did not influence the ROS (reactive

oxygen species) and RNS (reactive nitrogen species) levels in *Salmonella*-infected

RAW264.7 cells (Figure 1_figure Supplement 1). Flow cytometry analysis indicated

that the addition of 1 mM β-alanine did not affect the percentage of pro-inflammatory

M1 macrophages (CD86+) and anti-inflammatory M2 macrophages (CD163+) during

Salmonella infection (Figure 1_figure Supplement 2), implying that the addition of

β-alanine to macrophages does not change their immune response. Combining these

results, we can further infer that *Salmonella* use host-derived β-alanine for

intracellular growth.

Direct validation of *Salmonella* using host-derived β-alanine for intracellular

growth requires a mutant that has a defect in β-alanine uptake. *Escherichia coli*

uptakes β-alanine via the transporter protein CycA (Schneider et al., 2004). However,

the *Salmonella* Δ*cycA* mutant was able to use β-alanine as the sole carbon source for

growth in minimal medium (Figure 1_figure Supplement 3), indicating that CycA is

not a transporter for β-alanine in *Salmonella* and that other unknown β-alanine import

mechanisms are involved. Consistent with these results, mutation of *cycA* did not

influence the replication of *Salmonella* in RAW264.7 cells (Figure 1_figure

Supplement 4) or colonization in mouse systemic tissues (liver and spleen; Figure

1_figure Supplement 5).

***De novo* β-alanine synthesis is critical for *Salmonella* replication inside**

macrophages

Salmonella can *de novo* synthesize β-alanine *via* the decarboxylation

of L-aspartate, which is catalyzed by L-aspartate decarboxylase (PanD) (Figure 2A)

and is reportedly the rate-limiting step of β-alanine generation (Begley et al., 2001;

Schmitzberger et al., 2003; West et al., 1985). To further assess the role of β-alanine in

Salmonella intracellular replication, we analyzed the expression level of the

Salmonella panD gene in macrophages and the impact of *panD* mutation on the

ability of *Salmonella* to replicate in macrophages. Quantitative real-time PCR

(qRT□PCR) assays revealed that the expression level of *panD* was significantly ($P <$

0.01) greater in RAW264.7 cells than in RPMI-1640 medium (Figure 2B). Increased

expression of *panD* was also observed in N-minimal medium, a widely used medium

that mimics the conditions inside macrophages, as revealed by qRT□PCR and

bioluminescent reporter assays (Figure 2C, D). These results demonstrate that *panD*

expression is enhanced during *Salmonella* growth inside macrophages, suggesting a

relationship between *panD* expression and intracellular *Salmonella* growth.

We then constructed the *panD* mutant strain Δ*panD* and compared the replication

ability of the Δ*panD* strain and the *Salmonella* Typhimurium 14028s wild type (WT)

strain in RAW264.7 cells. Gentamicin protection assays showed that the replication of

Δ*panD* in RAW264.7 cells decreased 2.4-fold at 20 h postinfection compared with

that of the WT strain ($P < 0.001$), while complementation of Δ*panD* with the *panD*

gene restored the replication ability of the mutant strain in RAW264.7 cells (Figure

2E). Immunofluorescence analysis revealed that the number of Δ*panD* in each

infected RAW264.7 cell was comparable to that of the WT strain at the initial

infection stage (2 h), but at 20 h postinfection, the number of Δ*panD* in each infected

RAW264.7 cell was significantly ($P < 0.0001$) lower than that of the WT strain

(Figure 2F, G). These results indicate that *panD* contributes to *Salmonella* replication

in macrophages. The growth rates of Δ*panD* in LB medium and RPMI medium

resembled those of the WT (Figure 2_figure Supplement 1, Figure 2_figure

Supplement 2), indicating that the impaired intracellular replication ability of the

mutant was not due to a growth defect. Moreover, the replication defect of Δ*panD* in

RAW264.7 cells was relieved by the addition of 1 mM β-alanine to the RPMI medium

(Figure 2H). Furthermore, we examined the role of β-alanine synthesis in the

intracellular replication of *Salmonella* within another typical serovar, *Salmonella*

enterica serovar Typhi (*S.* Typhi), a serovar specific to humans and the causative

agent of typhoid fever (de Jong, Parry, van der Poll, & Wiersinga, 2012). We found

that the replication of *S.* Typhi Δ*panD* in human THP-1 macrophages was reduced by

2.6-fold compared to the the *S.* Typhi Ty2 WT strain ($P < 0.01$) (Figure 2_figure

Supplement 3), suggesting that *panD* also facilitates *S.* Typhi replication within

human macrophages.

These data collectively suggest that β-alanine synthesis is critical for *Salmonella*

replication inside macrophages.

***De novo* β-alanine synthesis is critical for systemic *Salmonella* infection in mice**

As replication in macrophages is a key determinant of systemic *Salmonella*

infection, we reasoned that β-alanine synthesis also influences *Salmonella* systemic

infection *in vivo*. To determine whether β-alanine influences systemic *Salmonella*

infection, we conducted mouse infection assays using intraperitoneal (i.p.) injection.

This method allows *Salmonella* to disseminate directly to systemic sites *via* the

lymphatic and bloodstream system, bypassing the need for intestinal invasion

(Pandeya et al., 2023; Silva et al., 2016). BALB/c mice were infected by i.p. injection

of 5,000 CFU of WT, Δ*panD*, or the complemented strain c*panD*. The survival rate,

body weight, bacterial burden in the liver and spleen, and liver histopathological

alterations of the infected mice were measured (Figure 3A). The WT-infected mice

exhibited high lethality and marked loss of body weight within 5 days, and all mice

died within 9 days of infection (Figure 3B, C). In contrast, the Δ*panD*-infected mice

displayed significantly improved survival rates and body weights, and no mice died

within the 10-day surveillance period (Figure 3B, C). Consistent with these results,

the bacterial burden in the liver and spleen of the Δ*panD*-infected mice was

significantly decreased, and the body weight was significantly increased compared

with that of the WT-infected mice on day 3 postinfection (Figure 3D).

Complementation of Δ*panD* with *panD* significantly decreased the survival rate and

body weight of infected mice but significantly increased the bacterial burden in the

liver and spleen of the infected mice (Figure 3B-D). Through immunofluorescence

staining, we examined the bacterial count in liver macrophages of mice infected with

WT, Δ*panD*, and complemented strain. The results showed that the bacterial count in

each macrophage from Δ*panD*-infected mice was significantly ($P < 0.0001$) lower

than that in WT-infected mice, on day 5 post-infection. Complementation of Δ*panD*

with *panD* restored the bacterial count in each macrophage to WT level (Figure 3E).

Furthermore, H&E staining revealed increased aggregation of inflammatory cells and

pyknosis in the livers of the WT-infected mice on day 5 postinfection, while these

histopathological alterations were obviously reduced in the livers of Δ*panD*-infected

mice (Figure 3F). Taken together, these results reveal that β-alanine synthesis is

critical for systemic *Salmonella* infection in mice.

β-Alanine is involved in the regulation of several metabolic pathways in

Salmonella

To explore the mechanism(s) associated with β-alanine-mediated promotion of

Salmonella replication in macrophages and *in vivo* virulence, we performed RNA

sequencing (RNA-seq) to reveal the differences in gene transcripts between

Salmonella WT and Δ*panD* strains grown in N-minimal medium. PCA plot of the

global transcriptomic profiles clearly demonstrated separation between the WT and

Δ*panD* strains (Figure 4A). Remarkable transcriptional changes were observed due to

the mutation of *panD*. Compared with those in the WT strain, 1379 genes were

differentially expressed in the Δ*panD* strain, with 561 upregulated genes and 618

downregulated genes (fold change ≥ 2 and *P* value < 0.05; Figure 4B). Gene Ontology

(GO) enrichment analysis revealed that the differentially expressed genes (DEGs)

were mainly involved in the metabolism and biosynthesis of several amino acids

(including arginine, leucine, histidine, and branched amino acids), carboxylic acid

metabolism, small molecule biosynthesis, and aerobic respiration (Figure 4C). Kyoto

Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis also

revealed a high frequency of terms related to metabolism, including amino acid

metabolism, lipid metabolism, carbohydrate metabolism, energy metabolism, and

nucleotide metabolism (Figure 4D). These data collectively indicate that β-alanine is

involved in the regulation of a series of metabolic pathways in *Salmonella*.

Further analysis of the downregulated DEGs (activated by PanD) revealed that

mutation of *panD* decreased the expression of genes involved in 7 pathways that are

associated with the virulence of *Salmonella* or other bacterial pathogens, including

methionine metabolism, fatty acid β-oxidation, histidine biosynthesis, and the

transport of zinc, galactose, potassium, and polyamine (Figure 4E). Zinc and

potassium uptake are associated with the virulence of *Salmonella* (zinc acquisition

promotes *Salmonella* Typhimurium virulence in mice, and potassium acquisition

promotes *Salmonella* Enteritidis virulence in chickens) (Ammendola et al., 2007;

Battistoni, Ammendola, Chiancone, & Ilari, 2017; Ilari et al., 2016; Liu et al., 2013),

while the remaining 5 pathways are involved in the pathogenicity of other pathogens

(Basavanna et al., 2013; de Paiva et al., 2016; Feldman et al., 2016; Lauriano et al.,

2004; Martínez-Guitián et al., 2019). In addition, the expression of the LysR-type

transcriptional regulator LeuO, which activates the expression of the *leuABCD*

leucine synthesis operon and numerous virulence genes in *Salmonella* Typhimurium

(Dillon et al., 2012; Guadarrama, Villaseñor, & Calva, 2014; Hernández-Lucas et al.,

2008), was also downregulated in the Δ*panD* strain (Figure 4E). In line with the

decreased expression of *leuO*, the expression of *leuABCD* was downregulated in the

Δ*panD* strain (Figure 4E).

We selected 16 downregulated DEGs (including the regulatory gene *leuO* and

genes from the above 7 pathways) for qRT□PCR analysis. The results showed that

the expression of all 16 genes significantly ($P < 0.05$) decreased in the Δ*panD* mutant

compared with the WT strain (Figure 4F), and complementation of Δ*panD* with *panD*

restored the gene expression to the WT level (Figure 4F), thus confirming the positive

regulation of these pathways and LeuO by β-alanine.

Salmonella pathogenesis largely depends on virulence genes encoded by

Salmonella pathogenicity islands (SPIs), with SPI-1 to SPI-5 being well-characterized

for their involvement in *Salmonella* virulence (Han et al., 2024). SPI-2 gene

expression is essential for *Salmonella* replication in macrophages and systemic

infection (Deng et al., 2017), yet the expression of SPI-2 genes was unaffected by

panD mutation (Figure 4_figure Supplement 1). Moreover, the gene expression of

four other virulence-associated SPIs, SPI-1, SPI-3, SPI-4, and SPI-5, is also

unaffected by *panD* mutation (Figure 4_figure Supplement 2, Figure 4_figure

Supplement 3).

Taken together, these data suggest that β-alanine might promote *Salmonella*

intracellular replication and virulence by activating virulence-associated pathway(s)

or activating the virulence-associated regulator LeuO, rather than by activating the

expression of virulence genes encoded within pathogenicity islands.

β-Alanine promotes *Salmonella* virulence *in vivo* partially by increasing the

expression of zinc transporter genes

Next, we inactivated the 7 downregulated pathways mentioned above, as well as

the regulatory gene *leuO* in *Salmonella,* to uncover the mechanism(s) by which

β-alanine promotes *Salmonella* virulence *in vivo*. Mouse infection assays revealed that

mutations in *fadAB*, *metR*, *hisABCDFGHL*, *kdpABC*, *mglABC*, and *potFGHI,* which

are associated with fatty acid β-oxidation, methionine metabolism, histidine

biosynthesis, potassium uptake, galactose uptake, and polyamine uptake, respectively,

did not influence *Salmonella* colonization in the mouse liver or spleen (Figure 5A, B)

or the body weight of infected mice (Figure 5C). Interestingly, although LeuO has

been reported to be associated with the regulation of a diverse set of virulence factors

(Dillon et al., 2012; Guadarrama et al., 2014), mutation of the regulatory gene *leuO*

did not influence *Salmonella* colonization in the mouse liver or spleen (Figure 5A, B)

or the body weight of infected mice (Figure 5C). In contrast, mutation of the zinc

transporter gene *znuA* significantly decreased *Salmonella* colonization in the mouse

liver and spleen (Figure 5D, left and middle panels); this result is consistent with

previous studies (Ammendola et al., 2007; Battistoni et al., 2017; Ilari et al., 2016).

Accordingly, the body weight of Δ*znuA*-infected mice was significantly (P < 0.001)

greater than that of the WT-infected mice (Figure 5D, right panel). These results

indicate that β-alanine might promote *Salmonella* virulence *in vivo* by promoting zinc

uptake.

To test this hypothesis, we initially evaluated the zinc content in the livers of

WT- and Δ*panD*-infected mice, on day 3 post-infection. We observed that the zinc

concentration in the macrophages of Δ*panD*-infected mouse livers was 3.2-fold higher

than in those of WT-infected mice (*P* < 0.0001; Figure 5E), suggesting that the *panD*

gene and β-alanine are crucial for *Salmonella* to obtain zinc from host cells.

Next, we constructed a double mutant, Δ*panD*Δ*znuA*, and compared colonization

of the mouse liver and spleen of the double mutant to that of the single mutant, Δ*znuA*.

The results showed that colonization of the liver and spleen of infected mice by

Δ*panD*Δ*znuA* was significantly lower than that of infected mice colonized by Δ*znuA*

(Figure 5D, left and middle panels). In agreement with these results, the body weight

of Δ*panD*Δ*znuA*-infected mice was greater than that of Δ*znuA*-infected mice (Figure

5D, right panel), suggesting that the contribution of *panD* to the virulence of

Salmonella is partially dependent on *znuA*.

Collectively, these data indicate that β-alanine promotes *in vivo* virulence of

Salmonella partially by increasing the expression of zinc transporter genes.

β-Alanine promotes *Salmonella* replication within macrophages partially by

increasing the expression of zinc transporter genes

To determine whether β-alanine influences *Salmonella* intracellular replication

by acting on zinc transporters, the zinc content in RAW 264.7 macrophages infected

with WT and Δ*panD* was also examined. We observed that the zinc concentration in

Δ*panD*-infected RAW 264.7 cells increased by 1.8-fold compared to WT-infected

cells ($P < 0.0001$; Figure 6A), further indicating the *panD* gene and β-alanine are

crucial for *Salmonella* to absorb zinc from macrophages.

We then analyzed the ability of Δ*znuA* and Δ*panD*Δ*znuA* to replicate in

RAW264.7 macrophages *via* gentamicin protection assays. The results showed that

the replication of Δ*panD*Δ*znuA* in RAW264.7 cells was significantly reduced

compared with that of the single mutant Δ*znuA* (Figure 6B), implying that the

contribution of *panD* to the intracellular replication of *Salmonella* is partially

dependent on *znuA*. The addition of 100 μM zinc to RPMI medium increased the

replication of Δ*panD* in RAW264.7 cells (Figure 6C), while the addition of 1 mM

β-alanine to RPMI medium did not increase the replication of Δ*znuA* (Figure 6D),

suggesting that the impaired replication due to the decrease in β-alanine can be

relieved by zinc supplementation.

Taken together, these data indicate that β-alanine promotes *Salmonella*

replication within macrophages by increasing the expression of zinc transporter genes.

Discussion

Replication within host macrophages is a crucial step for *Salmonella* to cause

life-threatening systemic infection in the host (Bomjan, Zhang, & Zhou, 2019;

Lathrop et al., 2015), while the crosstalk between *Salmonella* and macrophages at the

metabolic interface is critical for intracellular *Salmonella* replication (Dandekar et al.,

2014; Herrero-Fresno & Olsen, 2018; Lynch & Lesser, 2021; Rosenberg et al., 2021;

Thompson, Fulde, & Tedin, 2018). Emerging evidence suggests that several

metabolites affect the replication of *Salmonella* in macrophages. The promotion of

intracellular replication by metabolites is possibly achieved in three ways: i)

metabolites are utilized by *Salmonella* as nutrients for intracellular growth (Bowden,

Rowley, Hinton, & Thompson, 2009; Eisenreich, Dandekar, Heesemann, & Goebel,

2010; J. Wang et al., 2021); ii) *Salmonella* senses metabolites as environmental cues

to activate the expression of virulence genes (Jiang et al., 2021; X. Wang et al., 2023);

and iii) metabolites can regulate the immune responses of macrophages (Michelucci

et al., 2013; Peace & O'Neill, 2022; Yang & Cong, 2021). In this study, we

demonstrated that *Salmonella* promotes its replication inside macrophages by utilizing

both host- and bacterial-derived β-alanine (Figure 6E). We showed that β-alanine

promotes *Salmonella* intracellular replication and systemic infection partially by

increasing the expression of zinc transporter genes and therefore the uptake of zinc by

intracellular *Salmonella* (Figure 6E). Therefore, this work identified another

metabolite that can influence the replication of *Salmonella* in macrophages and

illustrated the mechanism by which β-alanine promotes intracellular *Salmonella*

replication.

We observed that *Salmonella*-infected macrophages contained lower β-alanine

levels than mock-infected macrophages, while β-alanine supplementation in the cell

medium increased the replication of *Salmonella* in macrophages, revealing that

Salmonella uptakes host-derived β-alanine to promote intracellular replication. In

addition, a deficiency in the biosynthesis of β-alanine (*via* mutation of the

rate-limiting gene *panD*) reduced *Salmonella* replication in macrophages and

systemic infection in mice, suggesting that *Salmonella* also utilizes bacterial-derived

β-alanine to promote intracellular replication and pathogenicity. It is known that

bacteria are quite stringent with their energy resources (Bergkessel, 2020; Bosdriesz,

Molenaar, Teusink, & Bruggeman, 2015), while the results of this work indicate that

either acquisition from the host or *de novo* synthesis of β-alanine is critical for

Salmonella replication inside macrophages. We speculate that *Salmonella* relies on a

large amount of β-alanine to efficiently replicate in macrophages, thereby highlighting

the importance of β-alanine for *Salmonella* intracellular growth. Nevertheless, unlike

the closely related species *E. coli*, which takes up β-alanine via the transporter protein

CycA (Schneider et al., 2004), *Salmonel la* does not use CycA to uptake β-alanine,

implying the presence of other yet unknown β-alanine import mechanisms in

Salmonella.

Several amino acids, including lysine, proline, arginine, aspartate and asparagine,

have previously been reported to be associated with the pathogenicity of *Salmonella*

(Popp et al., 2015; Steeb et al., 2013). These amino acids are involved in the synthesis

of proteins in *Salmonella*. In contrast, β-alanine is not incorporated into proteins but

can participate in the regulation of bacterial activity through the synthesis of

pantothenate and CoA (Webb et al., 2004; White et al., 2001; Yuan et al., 2022).

Accordingly, our transcriptome data showed that a deficiency in β-alanine

biosynthesis affected the expression of *Salmonella* genes involved in a series of

important metabolic pathways. Importantly, although β-alanine does not influence the

gene expression of SPIs, it activates methionine metabolism; fatty acid β-oxidation;

histidine biosynthesis; and the transport of zinc, galactose, potassium, and polyamine,

which have been previously known to be associated with the virulence of *Salmonella*

and other bacterial pathogens. Further analysis revealed that β-alanine promotes

Salmonella intracellular replication and systemic infection partially by promoting the

uptake of zinc. As for the other virulence-related metabolic pathways activated by

β-alanine, methionine metabolism, fatty acid β-oxidation, and histidine biosynthesis

contribute to the virulence of *Streptococcus pneumoniae* (Basavanna et al., 2013),

Yersinia pestis (Feldman et al., 2016), and *Acinetobacter baumannii*

(Martínez-Guitián et al., 2019), respectively; the uptake of galactose and polyamine

influences the pathogenicity of *Francisella tularensis* (Lauriano et al., 2004) and the

avian pathogenic *Escherichia coli* (de Paiva et al., 2016), respectively; and the uptake

of potassium is associated with the virulence of *Salmonella* Enteritidis in chickens

(Liu et al., 2013). However, blocking these pathways did not influence the systemic

Salmonella Typhimurium infection in mice, implying that different bacterial

pathogens adopt different virulence strategies to establish infection. Determining

other mechanism(s) by which β-alanine promotes the intracellular replication and

systemic infection of *Salmonella* require further investigation.

We observed that β-alanine also activates the expression of the LysR-type

transcriptional regulator LeuO, which is known to regulate the expression of a wide

variety of *Salmonella* genes that impact the stress response and virulence (Dillon et al.,

2012; Guadarrama et al., 2014; Hernández-Lucas et al., 2008). Typically, LeuO

activates the synthesis of the quiescent porins OmpS1 and OmpS2, which are required

for *Salmonella* virulence in mice (De la Cruz et al., 2007; Fernández-Mora, Puente, &

Calva, 2004; Rodríguez-Morales et al., 2006). Consistent with the positive regulation

of OmpS1 and OmpS2 by LeuO, lack of *leuO* in *Salmonella* also attenuated virulence

in a mouse model (Rodríguez-Morales et al., 2006). However, the attenuated

phenotypes of the *leuO* mutant in mice were not evident after i.p. injection relative to

oral infection, as a previous report showed that the competitive index for the *leuO*

mutant indicated approximately 1,000-fold reduced colonization in mouse systemic

tissues after oral infection but much less reduced colonization after i.p. injection (less

than 10-fold) (Rodríguez-Morales et al., 2006). These results imply that LeuO might

be predominantly associated with the invasion and intestinal infection of *Salmonella*

but is weakly implicated in *Salmonella* intracellular replication and systemic infection.

Therefore, it is not surprising that lack of *leuO* did not significantly affect *Salmonella*

colonization in the systemic tissues of mice after i.p. injection, as revealed by our

results.

Zinc is an essential micronutrient for all living organisms and is used as a

cofactor for various enzymes and proteins (Bock, Müller, & Blankenfeldt, 2016; Ilari

et al., 2016; Tan, Bramlett, & Lindahl, 2004). In bacteria, zinc-binding proteins

account for approximately 5% of the bacterial proteome and play crucial roles in

bacterial metabolism and virulence (Andreini, Banci, Bertini, & Rosato, 2006).

Knockout of the zinc transporter ZnuABC reduces the virulence of *Salmonella*,

Campylobacter jejuni, *Haemophilus ducreyi*, *Moraxella*, and urinary tract pathogenic

Escherichia coli (UPEC) in the host (Ilari et al., 2016). Moreover, zinc is also utilized

by *Salmonella* to subvert the antimicrobial host defense of macrophages by inhibiting

NF-κB activation and impairing NF-κB-dependent bacterial clearance (Jennings,

Esposito, Rittinger, & Thurston, 2018; Wu et al., 2017). Therefore, the efficient

acquisition of zinc may be crucial for *Salmonella*'s survival and replication within

macrophages, where zinc availability is limited (*Infect Immun.* 2007, 75(12):5867-76;

Biochim Biophys Acta. 2016, 1860(3):534-41). It has been reported that *Salmonella*

utilizes the high-affinity ZnuABC zinc transporter to optimize zinc availability in host

cells (*Infect Immun.* 2007, 75(12):5867-76). In this study, we found that β-alanine can

increase the expression of the zinc transporter genes *znuABC*, which could represent

an additional mechanism for the efficient zinc uptake of *Salmonella* in macrophages.

The results demonstrate a correlation between *Salmonella* β-alanine utilization and

zinc uptake during intracellular infection and provide evidence that β-alanine can

influence the macrophage immune response by acting on zinc uptake.

Overall, our findings suggest a model in which *Salmonella* exploits host- and

bacterial-derived β-alanine to efficiently replicate in host macrophages and cause

systemic disease. We propose that *Salmonella* requires a large amount of β-alanine

during intracellular infection. The utilization of β-alanine promotes *Salmonella* uptake

of the essential micronutrient zinc, which was previously shown to be required for the

metabolic needs of intracellular *Salmonella* and to subvert the antimicrobial defense

of macrophages by *Salmonella*. These observations provide new insight into

Salmonella pathogenesis and the crosstalk between *Salmonella* and macrophages

during intracellular infection. Considering that the *panD* gene is present in the

genomes of all *Salmonella* strains and that mutation of *panD* markedly reduced

Salmonella replication ability in macrophages, as well as virulence in the mouse

model, this gene may be used as a potential target to control systemic *Salmonella*

infection.

Methods

Ethics statement

Six-week-old female BALB/c mice were obtained from Beijing Vital River

Laboratory Animal Technology (Beijing, China). Mice were housed in barrier

facilities under specific pathogen-free conditions with a 12 h light/dark cycle at a

temperature of 24 ± 2 °C and a relative humidity of 50 ± 5%. Mice were fed a

standard mouse chow diet, and they consumed food and water *ad libitum* throughout

the experiment. All animal experiments were conducted in accordance with the

policies of the Institutional Animal Care Committee of Nankai University (Tianjin,

China) and performed under protocol no. 2021-SYDWLL-000029.

Cell culture

The RAW264.7 mouse macrophage-like cell line (ATCC TIB-71) was obtained

from the Shanghai Institute of Biochemistry and Cell Biology of the Chinese

Academy of Sciences (Shanghai, China). Cells were cultured in RPMI-1640 medium

(Gibco #11879020) supplemented with 10% (v/v) fetal bovine serum (FBS, Gibco

#10100147) at 37 □ with 5% CO_2. Cells were seeded in 24-well tissue culture plates

at 1×10^5 cells per well 24 h before infection.

Bacterial strains, plasmids, and growth conditions

The bacterial strains and plasmids used in this study are listed in Supplementary

Table 1. The *Salmonella enterica* serovar Typhimurium (STM) strain ATCC 14028s

was used as the WT strain throughout this study and for the construction of the

mutants. Mutant strains were generated using the λ Red recombination system with

the plasmid pSIM17 (Jiang et al., 2017). To construct the complemented strain of

Δ*panD*, the amplified DNA fragments of the *panD* ORF and its upstream promoter

were digested and inserted into the BamHI–EcoRI site of the low-copy-number

plasmid pBR322. To generate the *panD-lux* reporter fusion, the amplification

products of the *panD* promoter region were digested and cloned into the XhoI–BamHI

site of the plasmid pMS402, which carries a promoter-less *lux*CDABE reporter gene

cluster (Liang, Li, Dong, Surette, & Duan, 2008). The sequences of primers used for

the construction of the strains are listed in Supplementary Table 2. All the strains were

verified by PCR amplification and sequencing.

Bacterial strains were conventionally grown overnight in Luria–Bertani (LB)

medium (10 g/L tryptone, 5 g/L yeast extract, and 10 g/L NaCl) or in N-minimal

medium (10 μM $MgCl_2$, 110 μM KH_2PO_4, 7.5 mM $(NH_4)_2SO_4$, 0.5 mM K_2SO_4, 5 mM

KCl, 38 mM glycerol, and 0.1% [w/v] casamino acids) supplemented with appropriate

antibiotics at 37 °C with shaking at 180 rpm or on LB agar plates. All antibiotics were

used at their standard concentrations (chloramphenicol, 25 μg/mL; kanamycin, 50

μg/mL; ampicillin, 100 μg/mL; gentamicin, 10 or 100 μg/mL) unless otherwise

mentioned.

Growth curve

Bacterial strains were conventionally grown overnight in LB medium. The next

day, they were subcultured (1:100) in new LB medium and RPMI-1640 medium or

subcultured in N-minimal medium supplemented with glycerol or β-alanine as the

sole carbon source at 37 °C with shaking at 180 rpm. To measure the growth of

bacteria, 200 μL of the bacterial cultures were transferred to the microplate wells. The

absorbance (OD_{600}) of the bacteria was measured every half hour for 12 h with a

Spark multimode microplate reader (Tecan).

Bioluminescent reporter assays

STM WT carrying the *panD-luxCDABE* fusion plasmid was conventionally

grown overnight in LB medium, and the next day, the cells were subcultured (1:100)

in new LB medium or N-minimal medium for 8 h. The luminescence of the cultured

bacteria (200 μl) was measured with a Spark multimode microplate reader (Tecan).

Moreover, the cultured bacteria (100 μl) were serially diluted and plated on LB agar

plates to estimate bacterial CFUs. Bacterial CFUs were used to normalize

luminescence values.

Salmonella **infection of macrophages**

Bacterial strains were conventionally grown overnight in LB medium to the late

stationary phase, and the next day, the bacteria were diluted to 2×10^6 CFUs/mL and

opsonized in RPMI-1640 medium supplemented with 10% FBS for 15 min. The

macrophage monolayers were infected with the opsonized bacteria culture (0.5

mL/well, multiplicity of infection (MOI) = 10) and centrifuged at $800 \times g$ for 5 min to

synchronize infection. The infected cells were incubated for 30 min at 37 °C in 5%

CO_2 and then washed three times with $1\times$ PBS. Fresh RPMI-1640 medium containing

100 μg/mL gentamicin was added to the infected cells to kill extracellular bacteria.

After 1 h, fresh RPMI-1640 medium containing 10 μg/mL gentamicin was added to

the infected cells for the remainder of the experiment. To assess the intarcelluar

growth of *Salmonella*, the infected cells were lysed with 1% Triton X-100 at 2 hpi and

20 hpi, and the abundance of the intracellular bacteria CFUs was estimated on LB

agar plates. The relative fold replication of intracellular bacterial strains was denoted

as the CFUs recovered at 20 hpi relative to those at 2 hpi. The relative fold change in

replication was normalized to the number of RAW264.7 cells. When indicated, 1 mM

β-alanine or 100 μM $ZnSO_4$ were added after 1 h of gentamicin treatment.

Targeted metabolomics analysis of amino acids in macrophages

Targeted metabolomics was conducted as previously described (Jiang et al., 2021), with the combined metabolites of infected cells and intracellular bacteria were extracted for analysis. The metabolites have been confirmed to be dominated by the host metabolites (Jiang et al., 2021). RAW264.7 cells were mock-infected or infected with STM WT for 8 h. The cells were harvested and washed with precooled PBS solution to remove the medium. Cellular metabolites were extracted using ice-cold extraction solvent (40:40:20 vol/vol/vol acetonitrile:methanol:water, 0.1 M formic acid), incubated at -20 □ for 20 min, and then centrifuged for 10 min at 12,000×g and 4 □ to obtain the supernatant. Subsequently, the supernatant was transferred to an LC−MS vial and analyzed using ultrahigh-performance liquid chromatography (Acquity; Waters, Milford, MA, USA) coupled with mass spectrometry (Q Exactive Hybrid Quadrupole-Orbitrap; Thermo Fisher Scientific, Waltham, MA, USA). Metabolites were separated with a Luna NH$_2$ column (2 mm × 100 mm, 3 µm particle size; Phenomenex). Mobile phase A was 20 mM ammonium acetate (pH 9.0), and mobile phase B was acetonitrile containing 0.1% formic acid. The flow rate was 0.4 ml/min. Xcalibur 4.0 software (Thermo Fisher) was used for data acquisition and processing. Metabolite identification was achieved by high-resolution mass and retention time matching to authentic standards. The absolute quantification of amino acids was performed using the standard curve method, and the values were normalized to the cell number. Four biological replicates of each sample were analyzed.

Immunofluorescence staining

RAW264.7 cells were infected with STM WT or Δ*panD* and the complemented strain c*panD* as described above. After 2 and 20 h of cultivation, the infected cells were fixed for 15 min in 4% paraformaldehyde, washed in PBS, and permeabilized with 0.1% Triton X-100 in PBS for 15 min. The fixed samples were blocked in 5% bovine serum albumin for 30 min, followed by staining with a FITC-conjugated anti-*Salmonella* antibody (1:100 dilution, Abcam #ab20320) for 1 h at room temperature in the dark. The nuclei were then stained with DAPI (Invitrogen #D21490) for 2 min. A confocal laser scanning microscope (Zeiss LSM800) and ZEN 2.3 software (blue edition) were used to acquire and analyze the cell images (Objective lense: 40×; The number of intracellular bacteria per infected cell was estimated in random fields by Fiji-ImageJ).

RNA isolation

RNA was extracted from *Salmonella* strains cultured in N-minimal medium or LB medium. To investigate the expression of the *Salmonella panD* gene inside macrophages, we obtained RNA from intracellular bacteria in RAW264.7 cells at 8 h postinfection and from bacteria in RPMI medium. RNA was extracted using an EASYspinPlus bacterial RNA rapid extraction kit (Aidlab #RN0802) according to the manufacturer's protocol. RNA quantity and purity were determined using a NanoDrop 2000 spectrophotometer (NanoDrop Technologies). RNA samples were stored at −80 °C before use.

Quantitative real-time PCR (qRT□PCR)

According to the manufacturer's protocols, qRT□PCR was performed using 2× RealStar Power SYBR qPCR Mix (Genstar #A304) in a QuantStudio 5 Real Time PCR system (Applied Biosystems). cDNA was synthesized using a StarScript III RT Kit (Genstar #A232). Each sample was subjected to qRT□PCR in triplicate. The expression level of the 16S *rRNA* gene was used to normalize that of the target genes. We estimated the expression of each target gene using the $2^{-\Delta\Delta Ct}$ method.

RNA sequencing and analyses

The STM WT and Δ*panD* strains were conventionally grown overnight in LB medium, subcultured (1:100) in N-minimal medium for 8 h and then collected by centrifugation for RNA extraction. Sequencing libraries were generated using the NEBNext Ultra RNA Library Prep Kit for Illumina (New England Biolabs) according to the manufacturer's instructions, and sequencing was conducted using the Illumina HiSeq 2000 platform at Shanghai Majorbio Bio-Pharm Technology Co., Ltd. (Shanghai, China). The sequencing data were deposited in the NCBI Sequence Read Archive under accession number (SRA, PRJNA1124637). The clean reads were mapped to the STM ATCC 14028 reference genome (CP001363 and CP001362) by using the short-sequence alignment software Bowtie 2. Gene expression was evaluated using the fragments per kilobase of transcript per million mapped reads (FPKM) method. DEGs in the *panD* mutant relative to the WT were determined using the R statistical package software EdgeR. The thresholds for statistically significant differences were set to a fold change□≥□2 or□≤□0.5 and a false discovery rate

(FDR)≤0.05. P values were adjusted using the Benjamini–Hochberg procedure for

controlling the FDR. Enrichment analysis of DEGs was conducted using GO and

KEGG enrichment analyses.

Mouse infection

Salmonella strains were conventionally cultured overnight in LB medium, and

the next day, they were subcultured (1:100) in new LB medium and grown at 37 °C

with shaking at 200 rpm to stationary phase (OD_{600}~2). The bacteria were diluted to 5

$\times 10^4$ CFUs/mL in 0.9% NaCl. Groups of BALB/c mice were infected i.p. with 0.1

mL of the NaCl suspension. For survival assays, we recorded and monitored the

mortality and body weight of the infected mice daily for 10 days. To analyze the

bacterial burden of the mouse liver and spleen, we weighed the infected mice first on

day 3 postinfection and then harvested the liver and spleen. The liver and spleen of

infected mice were homogenized in ice-cold PBS, serially diluted, and plated on LB

plates containing the appropriate antibiotics to determine bacterial CFUs. H&E

staining of the mouse liver was performed to investigate the histopathological

alterations in the liver of infected mice. To evaluate the histopathological alterations

in the mouse liver, we harvested the liver of the infected mice on day 5 postinfection.

The mouse liver was washed with 0.9% NaCl, fixed with 10% neutral formalin for 48

h and subsequently processed for routine paraffin embedding. Paraffin-embedded

tissues were sectioned at a thickness of 5 μm and then stained with hematoxylin

(Sangon Biotech #E607318) and eosin (Sangon Biotech #E607318) for

histopathological examination. The stained sections were then examined by light

microscopy (Leica DM2500 LED).

Flow cytometry

RAW264.7 cells were infected with *Salmonella* WT for 8 h, in the absence or

presence of 1 mM β-alanine, which was added to the infected cells at 1 h

post-infection. The infected cells were fixed for 15 min in 4% paraformaldehyde,

washed in PBS, and permeabilized with 0.1% Triton X-100 in PBS for 15 min. The

fixed samples were blocked in 5% bovine serum albumin for 30 min, followed by

staining with an anti-CD86 antibody (Abcam, ab119857), an anti-CD163 antibody

(Abcam, ab182422) for 30 min, and a goat anti-rat IgG H&L (Alexa Fluor® 488)

(Abcam, ab150165), a donkey anti-rabbit IgG H&L (Alexa Fluor® 647) (Abcam,

ab150075) for 30 min in the dark. The infected cells were analyzed using a BD

FACSAria Flow Cytometer (BD Biosciences).

Assessment of intracellular ROS/ RNS levels

RAW264.7 cells were infected with *Salmonella* WT for 8 hours, either in the

absence or presence of 1 mM β-alanine. The β-alanine was added to the infected cells

at 1 h post-infection. Following the infection, the cells were washed three times with

PBS, and the levels of ROS and RNS were measured using a ROS fluorescence probe

(BBoxiProbe® O06, Bestbio, BB-46051) and an RNS fluorescence probe

(BBoxiProbe® O52, Bestbio, BB-470567), respectively, in accordance with the

manufacturer's protocol. The ROS/ RNS levels were normalized to the cell count.

Assessment of zinc levels in mouse liver and RAW 264.7 macrophages

To assess the zinc level in mouse liver, mice were i.p. infected with

approximately 5000 CFUs of *Salmonella* WT or Δ*panD* for 3 days, with three mice in

each injection group. Following infection, mouse livers were collected and then

dissociated into single cells using the gentleMACS/Mouse Liver Dissociation Kit

(Miltenyi Biotec, 130-105-807) in accordance with the manufacturer's protocol. The

zinc concentration in mouse liver was determined using a zinc fluorescence probe

(Zinquin ethyl ester, MKBio, MX4516) following the manufacturer's instructions and

normalized by the cell count.

To assess the zinc level in RAW 264.7 macrophages, RAW 264.7 cells were

infected with *Salmonella* WT or Δ*panD* for 8 hours. The infected cells were washed

three times with PBS, and zinc concentration was measured using a zinc fluorescence

probe (Zinquin ethyl ester, MKBio, MX4516) following the manufacturer's protocol.

The zinc concentration was normalized by the cell count.

Statistical analysis

The data are presented as the mean ± SD. All in vitro experiments were

conducted in duplicate and repeated at least three times (n ≥ 3). Mouse assays were

performed twice, with at least two mouse (n ≥ 2) in each injection group, and the

combined data from the two experiments was used for statistical analysis. Statistical

analyses were performed using GraphPad InStat software (version GraphPad Prism

9.5.1, San Diego, CA, USA) with two-sided Student's *t* tests, one-way ANOVA,

two-way ANOVA, log-rank Mantel–Cox tests, or Mann□Whitney U tests according

684 to the test requirements (as stated in the figure legends). A P value < 0.05 indicated a

685 statistically significant difference. ns represents no statistical significance.

686 **Data availability statement**

687 The RNA-seq data generated in this study have been deposited in the NCBI

688 Sequence Read Archive (SRA) database under the accession number PRJNA1124637.

689 All other data associated with this study are available in the main text and

690 supplementary materials. Source data are provided in Supplementary Data 1.

691 **Acknowledgements**

692 This work was supported by the National Natural Science Foundation of China

693 (NSFC) Program (grant no. 32170110, 32270191), the Natural Science Foundation of

694 Tianjin (grant no. 22JCYBJC01060), and the Fundamental Research Funds for the

695 Central Universities, Nankai University (grant no. 63241588, 63243161).

696 **Author contributions**

697 LJ designed the research; SM, YS, XW, HG, RL, TY, CK, and JC performed the

698 research; BY provided technical support and insights; SM and BY analyzed the data;

699 and LJ and SM wrote the manuscript.

700 **Disclosure statement**

701 The authors report there are no competing interests to declare.

702 **ORCID**

703 Lingyan Jiang http://orcid.org/0000-0001-8580-606X

References

Ammendola, S., Pasquali, P., Pistoia, C., Petrucci, P., Petrarca, P., Rotilio, G., &

Battistoni, A. (2007). High-affinity Zn2+ uptake system ZnuABC is required for

bacterial zinc homeostasis in intracellular environments and contributes to the

virulence of *Salmonella enterica*. *Infect Immun, 75*(12), 5867-5876.

doi:10.1128/iai.00559-07

Andreini, C., Banci, L., Bertini, I., & Rosato, A. (2006). Zinc through the three

domains of life. *J Proteome Res, 5*(11), 3173-3178. doi:10.1021/pr0603699

Basavanna, S., Chimalapati, S., Maqbool, A., Rubbo, B., Yuste, J., Wilson, R. J., . . .

Brown, J. S. (2013). The effects of methionine acquisition and synthesis on

Streptococcus pneumoniae growth and virulence. *PLoS One, 8*(1), e49638.

doi:10.1371/journal.pone.0049638

Battistoni, A., Ammendola, S., Chiancone, E., & Ilari, A. (2017). A novel

antimicrobial approach based on the inhibition of zinc uptake in *Salmonella*

enterica. *Future Med Chem, 9*(9), 899-910. doi:10.4155/fmc-2017-0042

Begley, T. P., Kinsland, C., & Strauss, E. (2001). The biosynthesis of coenzyme A in

bacteria. *Vitam Horm, 61*, 157-171. doi:10.1016/s0083-6729(01)61005-7

Bellinger, P. M. (2014). β-Alanine supplementation for athletic performance: an

update. *J Strength Cond Res, 28*(6), 1751-1770.

doi:10.1519/jsc.0000000000000327

Bergkessel, M. (2020). Regulation of protein biosynthetic activity during growth

arrest. *Curr Opin Microbiol, 57*, 62-69. doi:10.1016/j.mib.2020.07.010

Bock, T., Müller, R., & Blankenfeldt, W. (2016). Crystal structure of AibC, a reductase involved in alternative de novo isovaleryl coenzyme A biosynthesis in *Myxococcus xanthus*. *Acta Crystallogr F Struct Biol Commun, 72*(Pt 8), 652-658. doi:10.1107/s2053230x16011146

Boldyrev, A. A., Aldini, G., & Derave, W. (2013). Physiology and pathophysiology of carnosine. *Physiol Rev, 93*(4), 1803-1845. doi:10.1152/physrev.00039.2012

Bomjan, R., Zhang, M., & Zhou, D. (2019). YshB promotes intracellular replication and is required for *salmonella* virulence. *J Bacteriol, 201*(17), e00314-19. doi:10.1128/jb.00314-19

Bosdriesz, E., Molenaar, D., Teusink, B., & Bruggeman, F. J. (2015). How fast-growing bacteria robustly tune their ribosome concentration to approximate growth-rate maximization. *Febs j, 282*(10), 2029-2044. doi:10.1111/febs.13258

Bowden, S. D., Rowley, G., Hinton, J. C., & Thompson, A. (2009). Glucose and glycolysis are required for the successful infection of macrophages and mice by *Salmonella enterica* serovar Typhimurium. *Infect Immun, 77*(7), 3117-3126. doi:10.1128/iai.00093-09

Brumell, J. H., Tang, P., Mills, S. D., & Finlay, B. B. (2001). Characterization of *Salmonella*-induced filaments (Sifs) reveals a delayed interaction between *Salmonella*-containing vacuoles and late endocytic compartments. *Traffic, 2*(9), 643-653. doi:10.1034/j.1600-0854.2001.20907.x

Bumann, D., & Schothorst, J. (2017). Intracellular *Salmonella* metabolism. *Cell*

Microbiol, 19(10). doi:10.1111/cmi.12766

Castanheira, S., & García-Del Portillo, F. (2017). *Salmonella* populations inside host

cells. *Front Cell Infect Microbiol, 7*, 432. doi:10.3389/fcimb.2017.00432

Dandekar, T., Fieselmann, A., Fischer, E., Popp, J., Hensel, M., & Noster, J. (2014).

Salmonella-how a metabolic generalist adopts an intracellular lifestyle during

infection. *Front Cell Infect Microbiol, 4*, 191. doi:10.3389/fcimb.2014.00191

Davaapil, H., Tsuchiya, Y., & Gout, I. (2014). Signalling functions of coenzyme A and

its derivatives in mammalian cells. *Biochem Soc Trans, 42*(4), 1056-1062.

doi:10.1042/bst20140146

de Jong, H. K., Parry, C. M., van der Poll, T., & Wiersinga, W. J. (2012).

Host-pathogen interaction in invasive Salmonellosis. *PLoS Pathog, 8*(10),

e1002933. doi:10.1371/journal.ppat.1002933

De la Cruz, M. A., Fernández-Mora, M., Guadarrama, C., Flores-Valdez, M. A.,

Bustamante, V. H., Vázquez, A., & Calva, E. (2007). LeuO antagonizes H-NS

and StpA-dependent repression in *Salmonella enterica* ompS1. *Mol Microbiol,*

66(3), 727-743. doi:10.1111/j.1365-2958.2007.05958.x

de Paiva, J. B., da Silva, L. P., Casas, M. R., Conceição, R. A., Nakazato, G., de Pace,

F., . . . da Silveira, W. D. (2016). *In vivo* influence of in vitro up-regulated genes

in the virulence of an APEC strain associated with swollen head syndrome. *Avian*

Pathol, 45(1), 94-105. doi:10.1080/03079457.2015.1125995

Deng, W., Marshall, N. C., Rowland, J. L., McCoy, J. M., Worrall, L. J., Santos, A.

S., . . . Finlay, B. B. (2017). Assembly, structure, function and regulation of type

III   secretion   systems.   *Nat   Rev   Microbiol,   15*(6),   323-337.

doi:10.1038/nrmicro.2017.20

Dillon, S. C., Espinosa, E., Hokamp, K., Ussery, D. W., Casadesús, J., & Dorman, C. J.

(2012). LeuO is a global regulator of gene expression in *Salmonella enterica*

serovar   Typhimurium.   *Mol   Microbiol,   85*(6),   1072-1089.

doi:10.1111/j.1365-2958.2012.08162.x

Eisenreich, W., Dandekar, T., Heesemann, J., & Goebel, W. (2010). Carbon

metabolism of intracellular bacterial pathogens and possible links to virulence.

Nat Rev Microbiol, 8(6), 401-412. doi:10.1038/nrmicro2351

Fàbrega, A., & Vila, J. (2013). *Salmonella enterica* serovar Typhimurium skills to

succeed in the host: virulence and regulation. *Clin Microbiol Rev, 26*(2), 308-341.

doi:10.1128/cmr.00066-12

Feldman, M., Harbeck, M., Keller, M., Spyrou, M. A., Rott, A., Trautmann, B., . . .

Krause, J. (2016). A High-Coverage *Yersinia pestis* Genome from a

Sixth-Century Justinianic Plague Victim. *Mol Biol Evol, 33*(11), 2911-2923.

doi:10.1093/molbev/msw170

Fernández-Mora, M., Puente, J. L., & Calva, E. (2004). OmpR and LeuO positively

regulate the *Salmonella enterica* serovar Typhi ompS2 porin gene. *J Bacteriol,*

186(10), 2909-2920. doi:10.1128/jb.186.10.2909-2920.2004

Fields, P. I., Swanson, R. V., Haidaris, C. G., & Heffron, F. (1986). Mutants of

Salmonella Typhimurium that cannot survive within the macrophage are

avirulent.   *Proc   Natl   Acad   Sci   U   S   A,   83*(14),   5189-5193.

791 doi:10.1073/pnas.83.14.5189

792 Figueira, R., & Holden, D. W. (2012). Functions of the *Salmonella* pathogenicity

793 island 2 (SPI-2) type III secretion system effectors. *Microbiology (Reading),*

794 *158*(Pt 5), 1147-1161. doi:10.1099/mic.0.058115-0

795 Gojković, Z., Sandrini, M. P., & Piskur, J. (2001). Eukaryotic beta-alanine synthases

796 are functionally related but have a high degree of structural diversity. *Genetics,*

797 *158*(3), 999-1011. doi:10.1093/genetics/158.3.999

798 Gout, I. (2019). Coenzyme A: a protective thiol in bacterial antioxidant defence.

799 *Biochem Soc Trans, 47*(1), 469-476. doi:10.1042/bst20180415

800 Guadarrama, C., Villaseñor, T., & Calva, E. (2014). The subtleties and contrasts of the

801 LeuO regulator in *Salmonella* Typhi: implications in the immune response. *Front*

802 *Immunol, 5*, 581. doi:10.3389/fimmu.2014.00581

803 Han, J., Aljahdali, N., Zhao, S., Tang, H., Harbottle, H., Hoffmann, M., . . . Foley, S. L.

804 (2024). Infection biology of *Salmonella* enterica. *EcoSal Plus, 12*(1),

805 eesp00012023. doi:10.1128/ecosalplus.esp-0001-2023

806 Hernández-Lucas, I., Gallego-Hernández, A. L., Encarnación, S., Fernández-Mora, M.,

807 Martínez-Batallar, A. G., Salgado, H., . . . Calva, E. (2008). The LysR-type

808 transcriptional regulator LeuO controls expression of several genes in

809 *Salmonella enterica* serovar Typhi. *J Bacteriol, 190*(5), 1658-1670.

810 doi:10.1128/jb.01649-07

811 Herrero-Fresno, A., & Olsen, J. E. (2018). *Salmonella* Typhimurium metabolism

812 affects virulence in the host - A mini-review. *Food Microbiol, 71*, 98-110.

doi:10.1016/j.fm.2017.04.016

Hoffman, J. R., Varanoske, A., & Stout, J. R. (2018). Effects of β-Alanine

supplementation on carnosine elevation and physiological performance. *Adv*

Food Nutr Res, 84, 183-206. doi:10.1016/bs.afnr.2017.12.003

Huerta Ojeda, Á., Tapia Cerda, C., Poblete Salvatierra, M. F., Barahona-Fuentes, G.,

& Jorquera Aguilera, C. (2020). Effects of beta-Alanine supplementation on

physical performance in aerobic-anaerobic transition zones: a systematic review

and meta-analysis. *Nutrients, 12*(9), 2490. doi:10.3390/nu12092490

Ilari, A., Pescatori, L., Di Santo, R., Battistoni, A., Ammendola, S., Falconi, M., . . .

Chiancone, E. (2016). *Salmonella enterica* serovar Typhimurium growth is

inhibited by the concomitant binding of Zn(II) and a pyrrolyl-hydroxamate to

ZnuA, the soluble component of the ZnuABC transporter. *Biochim Biophys Acta,*

1860(3), 534-541. doi:10.1016/j.bbagen.2015.12.006

Jennings, E., Esposito, D., Rittinger, K., & Thurston, T. L. M. (2018).

Structure-function analyses of the bacterial zinc metalloprotease effector protein

GtgA uncover key residues required for deactivating NF-κB. *J Biol Chem,*

293(39), 15316-15329. doi:10.1074/jbc.RA118.004255

Jiang, L., Feng, L., Yang, B., Zhang, W., Wang, P., Jiang, X., & Wang, L. (2017).

Signal transduction pathway mediated by the novel regulator LoiA for low

oxygen tension induced *Salmonella* Typhimurium invasion. *PLoS Pathog, 13*(6),

e1006429. doi:10.1371/journal.ppat.1006429

Jiang, L., Wang, P., Song, X., Zhang, H., Ma, S., Wang, J., . . . Wang, L. (2021).

Salmonella Typhimurium reprograms macrophage metabolism via T3SS effector

SopE2 to promote intracellular replication and virulence. *Nat Commun, 12*(1),

879. doi:10.1038/s41467-021-21186-4

Kehl, A., Noster, J., & Hensel, M. (2020). Eat in or Take out? Metabolism of

Intracellular Salmonella enterica. *Trends Microbiol, 28*(8), 644-654.

doi:10.1016/j.tim.2020.03.005

Kim, S., Lee, K. S., Pak, G. D., Excler, J. L., Sahastrabuddhe, S., Marks, F., . . .

Mogasale, V. (2019). Spatial and temporal patterns of typhoid and paratyphoid

fever outbreaks: a worldwide review, 1990-2018. *Clin Infect Dis, 69*(Suppl 6),

S499-s509. doi:10.1093/cid/ciz705

LaRock, D. L., Chaudhary, A., & Miller, S. I. (2015). Salmonellae interactions with

host processes. *Nat Rev Microbiol, 13*(4), 191-205. doi:10.1038/nrmicro3420

Lathrop, S. K., Binder, K. A., Starr, T., Cooper, K. G., Chong, A., Carmody, A. B., &

Steele-Mortimer, O. (2015). Replication of *Salmonella enterica* serovar

Typhimurium in human monocyte-derived macrophages. *Infect Immun, 83*(7),

2661-2671. doi:10.1128/iai.00033-15

Lauriano, C. M., Barker, J. R., Yoon, S. S., Nano, F. E., Arulanandam, B. P., Hassett,

D. J., & Klose, K. E. (2004). MglA regulates transcription of virulence factors

necessary for *Francisella tularensis* intraamoebae and intramacrophage survival.

Proc Natl Acad Sci U S A, 101(12), 4246-4249. doi:10.1073/pnas.0307690101

Leung, K. Y., & Finlay, B. B. (1991). Intracellular replication is essential for the

virulence of *Salmonella* Typhimurium. *Proc Natl Acad Sci U S A, 88*(24),

11470-11474. doi:10.1073/pnas.88.24.11470

Li, W., Ren, Q., Ni, T., Zhao, Y., Sang, Z., Luo, R., . . . Li, S. (2023). Strategies

adopted by *Salmonella* to survive in host: a review. *Arch Microbiol, 205*(12), 362.

doi:10.1007/s00203-023-03702-w

Liang, H., Li, L., Dong, Z., Surette, M. G., & Duan, K. (2008). The YebC family

protein PA0964 negatively regulates the Pseudomonas aeruginosa quinolone

signal system and pyocyanin production. *J Bacteriol, 190*(18), 6217-6227.

doi:10.1128/jb.00428-08

Liss, V., Swart, A. L., Kehl, A., Hermanns, N., Zhang, Y., Chikkaballi, D., . . . Hensel,

M. (2017). *Salmonella enterica* Remodels the Host Cell Endosomal System for

Efficient Intravacuolar Nutrition. *Cell Host Microbe, 21*(3), 390-402.

doi:10.1016/j.chom.2017.02.005

Liu, Y., Ho, K. K., Su, J., Gong, H., Chang, A. C., & Lu, S. (2013). Potassium

transport of *Salmonella* is important for type III secretion and pathogenesis.

Microbiology (Reading), 159(Pt 8), 1705-1719. doi:10.1099/mic.0.068700-0

Lynch, J. P., & Lesser, C. F. (2021). A host metabolite promotes *Salmonella* survival.

Science, 371(6527), 344-345. doi:10.1126/science.abf8414

Majowicz, S. E., Musto, J., Scallan, E., Angulo, F. J., Kirk, M., O'Brien, S. J., . . .

Hoekstra, R. M. (2010). The global burden of nontyphoidal *Salmonella*

gastroenteritis. *Clin Infect Dis, 50*(6), 882-889. doi:10.1086/650733

Martínez-Guitián, M., Vázquez-Ucha, J. C., Álvarez-Fraga, L., Conde-Pérez, K.,

Lasarte-Monterrubio, C., Vallejo, J. A., . . . Beceiro, A. (2019). Involvement of

HisF in the persistence of *Acinetobacter baumannii* during a pneumonia

infection. *Front Cell Infect Microbiol, 9*, 310. doi:10.3389/fcimb.2019.00310

Michelucci, A., Cordes, T., Ghelfi, J., Pailot, A., Reiling, N., Goldmann, O., . . . Hiller,

K. (2013). Immune-responsive gene 1 protein links metabolism to immunity by

catalyzing itaconic acid production. *Proc Natl Acad Sci U S A, 110*(19),

7820-7825. doi:10.1073/pnas.1218599110

Nozaki, S., Webb, M. E., & Niki, H. (2012). An activator for pyruvoyl-dependent

l-aspartate α-decarboxylase is conserved in a small group of the γ-proteobacteria

including Escherichia coli. *Microbiologyopen, 1*(3), 298-310.

doi:10.1002/mbo3.34

Ohl, M. E., & Miller, S. I. (2001). *Salmonella*: a model for bacterial pathogenesis.

Annu Rev Med, 52, 259-274. doi:10.1146/annurev.med.52.1.259

Pandeya, A., Zhang, Y., Cui, J., Yang, L., Li, J., Zhang, G., . . . Wei, Y. (2023).

Inflammasome activation and pyroptosis mediate coagulopathy and

inflammation in *Salmonella* systemic infection. *Microbiol Res, 275*, 127460.

doi:10.1016/j.micres.2023.127460

Peace, C. G., & O'Neill, L. A. (2022). The role of itaconate in host defense and

inflammation. *J Clin Invest, 132*(2), e148548. doi:10.1172/jci148548

Pillay, T. D., Hettiarachchi, S. U., Gan, J., Diaz-Del-Olmo, I., Yu, X. J., Muench, J.

H., . . . Pearson, J. S. (2023). Speaking the host language: how *Salmonella*

effector proteins manipulate the host. *Microbiology (Reading), 169*(6), 001342.

doi:10.1099/mic.0.001342

Popp, J., Noster, J., Busch, K., Kehl, A., Zur Hellen, G., & Hensel, M. (2015). Role of host cell-derived amino acids in nutrition of intracellular *Salmonella enterica*. *Infect Immun, 83*(12), 4466-4475. doi:10.1128/iai.00624-15

Rajashekar, R., Liebl, D., Seitz, A., & Hensel, M. (2008). Dynamic remodeling of the endosomal system during formation of *Salmonella*-induced filaments by intracellular *Salmonella* enterica. *Traffic, 9*(12), 2100-2116. doi:10.1111/j.1600-0854.2008.00821.x

Röder, J., Felgner, P., & Hensel, M. (2021). Comprehensive Single Cell Analyses of the Nutritional Environment of Intracellular *Salmonella enterica*. *Front Cell Infect Microbiol, 11*, 624650. doi:10.3389/fcimb.2021.624650

Rodríguez-Morales, O., Fernández-Mora, M., Hernández-Lucas, I., Vázquez, A., Puente, J. L., & Calva, E. (2006). *Salmonella enterica* serovar Typhimurium ompS1 and ompS2 mutants are attenuated for virulence in mice. *Infect Immun, 74*(2), 1398-1402. doi:10.1128/iai.74.2.1398-1402.2006

Rosenberg, G., Riquelme, S., Prince, A., & Avraham, R. (2022). Immunometabolic crosstalk during bacterial infection. *Nat Microbiol, 7*(4), 497-507. doi:10.1038/s41564-022-01080-5

Rosenberg, G., Yehezkel, D., Hoffman, D., Mattioli, C. C., Fremder, M., Ben-Arosh, H., . . . Avraham, R. (2021). Host succinate is an activation signal for *Salmonella* virulence during intracellular infection. *Science, 371*(6527), 400-405. doi:10.1126/science.aba8026

Schmitzberger, F., Smith, A. G., Abell, C., & Blundell, T. L. (2003). Comparative

analysis of the *Escherichia coli* ketopantoate hydroxymethyltransferase crystal structure confirms that it is a member of the (betaalpha)8 phosphoenolpyruvate/pyruvate superfamily. *J Bacteriol, 185*(14), 4163-4171. doi:10.1128/jb.185.14.4163-4171.2003

Schneider, F., Krämer, R., & Burkovski, A. (2004). Identification and characterization of the main beta-alanine uptake system in *Escherichia coli*. *Appl Microbiol Biotechnol, 65*(5), 576-582. doi:10.1007/s00253-004-1636-0

Sibon, O. C., & Strauss, E. (2016). Coenzyme A: to make it or uptake it? *Nat Rev Mol Cell Biol, 17*(10), 605-606. doi:10.1038/nrm.2016.110

Silva, A. F., Matos, M. P., Ralph, M. T., Silva, D. L., de Alencar, N. M., Ramos, M. V., & Lima-Filho, J. V. (2016). Comparison of immunomodulatory properties of mannose-binding lectins from Canavalia brasiliensis and Cratylia argentea in a mice model of *Salmonella* infection. *Int Immunopharmacol, 31*, 233-238. doi:10.1016/j.intimp.2015.12.036

Song, P., Zhang, X., Wang, S., Xu, W., & Wei, F. (2023). Advances in the synthesis of β-alanine. *Front Bioeng Biotechnol, 11*, 1283129. doi:10.3389/fbioe.2023.1283129

Steeb, B., Claudi, B., Burton, N. A., Tienz, P., Schmidt, A., Farhan, H., . . . Bumann, D. (2013). Parallel exploitation of diverse host nutrients enhances *Salmonella* virulence. *PLoS Pathog, 9*(4), e1003301. doi:10.1371/journal.ppat.1003301

Tan, X., Bramlett, M. R., & Lindahl, P. A. (2004). Effect of Zn on acetyl coenzyme a synthase: evidence for a conformational change in the alpha subunit during

catalysis. *J Am Chem Soc, 126*(19), 5954-5955. doi:10.1021/ja039600z

Theodoulou, F. L., Sibon, O. C., Jackowski, S., & Gout, I. (2014). Coenzyme A and

its derivatives: renaissance of a textbook classic. *Biochem Soc Trans, 42*(4),

1025-1032. doi:10.1042/bst20140176

Thompson, A., Fulde, M., & Tedin, K. (2018). The metabolic pathways utilized by

Salmonella Typhimurium during infection of host cells. *Environ Microbiol Rep,*

10(2), 140-154. doi:10.1111/1758-2229.12628

Tuli, A., & Sharma, M. (2019). How to do business with lysosomes: *Salmonella* leads

the way. *Curr Opin Microbiol, 47*, 1-7. doi:10.1016/j.mib.2018.10.003

Wang, J., Ma, S., Li, W., Wang, X., Huang, D., Jiang, L., & Feng, L. (2021).

Salmonella enterica serovar Typhi induces host metabolic reprogramming to

increase glucose availability for intracellular replication. *Int J Mol Sci, 22*(18),

10003. doi:10.3390/ijms221810003

Wang, L., Mao, Y., Wang, Z., Ma, H., & Chen, T. (2021). Advances in

biotechnological production of β-alanine. *World J Microbiol Biotechnol, 37*(5),

79. doi:10.1007/s11274-021-03042-1

Wang, X., Yang, B., Ma, S., Yan, X., Ma, S., Sun, H., . . . Jiang, L. (2023). Lactate

promotes *Salmonella* intracellular replication and systemic infection via driving

macrophage M2 polarization. *Microbiol Spectr, 11*(6), e0225323.

doi:10.1128/spectrum.02253-23

Webb, M. E., Smith, A. G., & Abell, C. (2004). Biosynthesis of pantothenate. *Nat*

Prod Rep, 21(6), 695-721. doi:10.1039/b316419p

West, T. P., Traut, T. W., Shanley, M. S., & O'Donovan, G. A. (1985). A *Salmonella*

typhimurium strain defective in uracil catabolism and beta-alanine synthesis. *J*

Gen Microbiol, 131(5), 1083-1090. doi:10.1099/00221287-131-5-1083

White, W. H., Gunyuzlu, P. L., & Toyn, J. H. (2001). *Saccharomyces cerevisiae* is

capable of de *novo* pantothenic acid biosynthesis involving a novel pathway of

beta-alanine production from spermine. *J Biol Chem, 276*(14), 10794-10800.

doi:10.1074/jbc.M009804200

Wu, A., Tymoszuk, P., Haschka, D., Heeke, S., Dichtl, S., Petzer, V., . . . Weiss, G.

(2017). *Salmonella* utilizes zinc to subvert antimicrobial host defense of

macrophages via modulation of NF-κB signaling. *Infect Immun, 85*(12),

e00418-17. doi:10.1128/iai.00418-17

Yang, W., & Cong, Y. (2021). Gut microbiota-derived metabolites in the regulation of

host immune responses and immune-related inflammatory diseases. *Cell Mol*

Immunol, 18(4), 866-877. doi:10.1038/s41423-021-00661-4

Yuan, S. F., Nair, P. H., Borbon, D., Coleman, S. M., Fan, P. H., Lin, W. L., & Alper,

H. S. (2022). Metabolic engineering of E. coli for β-alanine production using a

multi-biosensor enabled approach. *Metab Eng, 74*, 24-35.

doi:10.1016/j.ymben.2022.08.012

 Figure legends

 Figure 1. Host-derived β-alanine promotes *Salmonella* replication inside

 macrophages. (A) Schematic workflow for targeted metabolomics investigation of

 mock- and *Salmonella*-infected (STM) mouse RAW264.7 macrophages. Picture

 materials were used from bioicons (https://bioicons.com/). (B) Principal component

 analysis (PCA) score plots of metabolic profiles in the mock- and *Salmonella*-infected

 (STM) groups (n = 4 biologically independent samples). (C) The concentrations of

 upregulated amino acids in the mock- and *Salmonella*-infected groups (n = 4

 biologically independent samples). (D) The concentrations of downregulated amino

 acids in the mock- and *Salmonella*-infected groups (n = 4 biologically independent

 samples). (E) Fold intracellular replication (20 h *vs.* 2 h) of *Salmonella* WT in

 RAW264.7 cells in the presence of 0.5, 1, 2, 4 mM β-alanine. The data are presented

 as the mean ± SD, n = 3 independent experiments (E). Statistical significance was

 assessed using two-sided Student's *t*-test (C, D) and one-way ANOVA   (E).

 Figure 2. *De novo* β-alanine synthesis is critical for *Salmonella* replication inside

 macrophages. (A) Scheme of β-alanine and the downstream CoA biosynthesis

 pathway in *Salmonella*. (B) qRT☐PCR analysis of the expression of the *Salmonella*

 panD gene in RAW264.7 cells (8 h postinfection) and RPMI-1640 medium. (C)

 qRT☐PCR analysis of the expression of the *Salmonella panD* gene in N-minimal

 medium and LB medium. (D) Expression of the *panD*-lux transcriptional fusion in

 N-minimal medium and LB medium. Luminescence values were normalized to 10^5

bacterial CFUs. (E) Fold intracellular replication (20 h *vs.* 2 h) of *Salmonella*

Typhimurium 14028s wild type (WT), the *panD* mutant (Δ*panD*) and the

complemented strain (c*panD*) in RAW264.7 cells. (F) Number of intracellular

Salmonella WT, Δ*panD*, and c*panD* strains per RAW264.7 cell at 2 and 20 h

postinfection. The number of intracellular bacteria per infected cell was estimated in

random fields, n = 80 cells per group from 3 independent experiments. (G)

Representative immunofluorescence images of *Salmonella* WT, Δ*panD*, and c*panD* in

RAW264.7 cells at 20 h postinfection (green, *Salmonella*; blue, nuclei; scale bars, 50

µm). Images are representative of three independent experiments. (H) Replication of

Salmonella WT and Δ*panD* in RAW264.7 cells in the presence or absence of 1 mM

β-alanine. The data are presented as the mean ± SD, n = 3 (B–E, H) independent

experiments. Statistical significance was assessed using two-sided Student's *t*-test

(B-D) or one-way ANOVA (E, F, H). ns, not Significant.

1021

Figure 3. ***De novo* β-alanine synthesis is critical for systemic *Salmonella* infection**

in mice. (A) Schematic illustration of the mouse infection assays. Picture materials

were used from bioicons (https://bioicons.com/). (B, C) Survival curves (B) and body

weight dynamics (C) of mice infected i.p. with *Salmonella* WT, Δ*panD*, or c*panD*. n =

10 mice per group. (D) Liver and spleen bacterial burdens and body weights of mice

infected with *Salmonella* WT, Δ*panD*, or c*panD* on day 3 postinfection. n = 7 mice

per group. (E) Representative immunofluorescence images and intracellular bacterial

counts of *Salmonella* WT, Δ*panD*, and c*panD* in mouse liver at 5 days post-infection

(green, *Salmonella*; blue, nuclei; scale bars, 50 µm). Images are representative of

three independent experiments. The number of intracellular bacteria per infected cell

was estimated in random fields, with n = 80 cells per group from 3 independent

experiments. (F) Representative H&E-stained liver sections from mice that were left

uninfected or infected with *Salmonella* WT, Δ*panD*, or c*panD* on day 5 postinfection.

Arrows indicate severe inflammatory cell infiltration in the mouse liver. Images are

representative of three independent experiments. The data are presented as the mean ±

SD (B-E). Statistical significance was assessed using the log-rank Mantel–Cox test

(B), two-sided Student's *t*-test (C), or one-way ANOVA (D, E).

1039

Figure 4. β-Alanine is involved in the regulation of several metabolic pathways in

Salmonella. (A) Principal component analysis (PCA) score plots of transcriptomic

profiles of *Salmonella* WT and Δ*panD* (n = 3 biologically independent samples). (B)

Volcano plot of the differentially expressed genes (DEGs) in *Salmonella* WT versus

Δ*panD*. The upper right section (red dots) indicates the upregulated DEGs, and the

upper left section (green dots) indicates the downregulated DEGs. (C) Gene Ontology

(GO) enrichment analysis of DEGs. Bubble chart showing the top 20 enriched GO

terms. (D) Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment

analysis of DEGs. (E) Expression of the downregulated pathways (activated by PanD)

is shown in the Z score-transformed heatmap, with red representing higher abundance

and blue representing lower abundance. (F) qRT☐PCR analysis of the mRNA levels

of 16 selected downregulated DEGs in *Salmonella* WT, Δ*panD*, and c*panD*. The data

are presented as the mean ± SD, n = 3 independent experiments. Statistical

significance was assessed using two-way ANOVA.

1054

Figure 5. β-Alanine promotes *Salmonella* virulence *in vivo* partially by increasing

the expression of zinc transporter genes. (A, B, C) Liver (A) and spleen (B)

bacterial burdens and body weight (C) of mice infected with *Salmonella* WT, Δ*fadAB*,

Δ*metR*, Δ*hisABCDFGHL*, Δ*kdpABC*, Δ*mglABC*, Δ*potFGHI*, or Δ*leuO* on day 3

postinfection. n = 5 mice per group. (D) Liver and spleen bacterial burdens and body

weights of mice infected with *Salmonella* WT, Δ*panD,* Δ*znuA* or Δ*panD*Δ*znuA* on day

3 postinfection. n = 5 mice per group. (E) The zinc levels in the livers of mice

infection with either *Salmonella* WT or Δ*panD* for 3 days, n = 5 mice per group. The

data are presented as the mean ± SD (A–E). Statistical significance was assessed

using one-way ANOVA (A–D), two-sided Student's *t*-test (E). ns, not Significant.

1065

Figure 6. β-Alanine promotes *Salmonella* replication within macrophages

partially **by increasing the expression of zinc transporter genes.** (A) The zinc levels

in RAW264.7 cells after infection with *Salmonella* WT or Δ*panD* for 8 h. (B)

Replication of *Salmonella* WT, Δ*panD,* Δ*znuA* and Δ*panD*Δ*znuA* in RAW264.7 cells.

(C) Replication of *Salmonella* WT and Δ*panD* in RAW264.7 cells in the presence or

absence of 100 μM $ZnSO_4$. (D) Replication of *Salmonella* WT and Δ*znuA* in

RAW264.7 cells in the presence or absence of 1 mM β-alanine. The data are presented

as the mean ± SD, n = 3 independent experiments (A–D). Statistical significance was

assessed using two-sided Student's *t*-test (A), one-way ANOVA (B–D). ns, not

Significant. (E) Schematic model of β-alanine-mediated *Salmonella* replication inside

macrophages. In macrophages, *Salmonella* acquires β-alanine both via the uptake of

β-alanine from host macrophages and the *de novo* synthesis of β-alanine. β-alanine

promotes the expression of zinc transporter genes *ZnuABC,* which facilitate the uptake

of zinc by intracellular *Salmonella,* therefore promote *Salmonella* replication in

macrophages and subsequent systemic infection.

1081

1082 **Figure 1_figure Supplement 1.** The levels of ROS and RNS in RAW264.7 cells after

1083 infection with *Salmonella* WT for 8 hours, in the presence or absence of 1 mM

1084 β-alanine. Data are presented as mean ± SD, n = 3 independent experiments.

1085 Statistical significance was assessed using two-sided Student's *t*-test. ns, not

1086 Significant.

1087

Figure 1_figure Supplement 2. Flow cytometry analysis was conducted to determine

the percentage of pro-inflammatory M1 macrophages (CD86+) and anti-inflammatory

M2 macrophages (CD163+). RAW264.7 cells were infected with *Salmonella* WT for

8 hours, in the presence or absence of 1 mM β-alanine. Subsequently, the infected

cells were collected for flow cytometry analysis. Representative dot plots and

quantification of M1 (CD86+) and M2 (CD163+) macrophages are displayed in the

left and right panels, respectively. Data are presented as mean ± SD, n = 3

independent experiments. Statistical significance was assessed using two-way

ANOVA. ns, not Significant.

1097

1098   **Figure 1_figure Supplement 3.** Growth curves of *Salmonella* WT and the *cycA*

1099   mutant (Δ*cycA*) in N-minimal medium supplemented with β-alanine (1 mM) as the

1100   sole carbon source. Data are presented as mean ± SD, n = 4 independent experiments.

1101

1102   **Figure 1_figure Supplement 4.** Fold intracellular replication (20 h *vs.* 2 h) of

1103   *Salmonella* wild-type and Δ*cycA* in RAW264.7 cells. Data are presented as mean ±

1104   SD, n = 4 independent experiments. Statistical significance was assessed using

1105   two-way ANOVA. ns, not Significant.

1106

1107   **Figure 1_figure Supplement 5.** Liver and spleen bacterial burdens, and body weight

1108   of mice infected with *Salmonella* wild-type (STM) and *cycA* mutant (Δ*cycA*), at day 3

1109   post-infection. n = 5 mice per group. Statistical significance was assessed using

1110   Mann-Whitney U test. ns, not Significant.

1111

1112   **Figure 2_figure Supplement 1.** Growth curves of *Salmonella* WT, *panD* mutant

1113   (Δ*panD*) and the complemented strain (c*panD*) in LB medium. Data are presented as

1114   mean ± SD, n = 4 independent experiments.

1115

1116   **Figure 2_figure Supplement 2.** Growth curves of *Salmonella* WT, *panD* mutant

1117   (Δ*panD*) and the complemented strain (c*panD*) in RPMI-1640 medium (B). Data are

1118 presented as mean ± SD, n = 4 independent experiments.

1119

1120 **Figure 2_figure Supplement 3.** Fold intracellular replication (20 h *vs*. 2 h) of

1121 *Salmonella enterica* serovar Typhi Ty2 wild type (WT), the *panD* mutant (Δ*panD*) in

1122 human THP-1 monocyte like cell line (ATCC TIB-22). Data are presented as mean ±

1123 SD, n = 4 independent experiments. Statistical significance was assessed using

1124 two-sided Student's *t*-test. ns, not Significant.

1125

1126 **Figure 4_figure Supplement 1.** Expression of SPI-2 is shown in the Z-score

1127 transformed heatmap, with orange representing higher and blue representing lower

1128 abundance.

1129

1130 **Figure 4_figure Supplement 2.** Expression of SPI-1 is shown in the Z-score

1131 transformed heatmap, with orange representing higher and blue representing lower

1132 abundance.

1133

1134 **Figure 4_figure Supplement 3.** Expression of SPI-3, SPI-4, and SPI-5 genes is

1135 shown in the Z-score transformed heatmap, with orange representing higher and blue

1136 representing lower abundance.

1137

1138 **Figure 1_source data 1.** The numerical source data corresponds to Figure 1.

1139

1140 **Figure 2_source data 1.** The numerical source data corresponds to Figure 2.

1141   **Figure 3_source data 1.** The numerical source data corresponds to Figure 3.

1142

1143   **Figure 4_source data 1.** The numerical source data corresponds to Figure 4.

1144

1145   **Figure 5_source data 1.** The numerical source data corresponds to Figure 5.

1146

1147   **Figure 6_source data 1.** The numerical source data corresponds to Figure 6.

1148

1149   **Figure 1_figure Supplement 1_source data 1.** The numerical source data

1150   corresponds to Figure 1_figure Supplement 1.

1151

1152   **Figure 1_figure Supplement 2_source data 1.** The numerical source data

1153   corresponds to Figure 1_figure Supplement 2.

1154

1155   **Figure 1_figure Supplement 3_source data 1.** The numerical source data

1156   corresponds to Figure 1_figure Supplement 3.

1157

1158   **Figure 1_figure Supplement 4_source data 1.** The numerical source data

1159   corresponds to Figure 1_figure Supplement 4.

1160

1161   **Figure 1_figure Supplement 5_source data 1.** The numerical source data

1162   corresponds to Figure 1_figure Supplement 5.

1163

1164 **Figure 2_figure Supplement 1_source data 1.** The numerical source data

1165 corresponds to Figure 2_figure Supplement 1.

1166

1167 **Figure 2_figure Supplement 2_source data 1.** The numerical source data

1168 corresponds to Figure 2_figure Supplement 2.

1169

1170 **Figure 2_figure Supplement 3_source data 1.** The numerical source data

1171 corresponds to Figure 2_figure Supplement 3.

1172

1173 **Figure 4_figure Supplement 1_source data 1.** The numerical source data

1174 corresponds to Figure 4_figure Supplement 1.

1175

1176 **Figure 4_figure Supplement 2_source data 1.** The numerical source data

1177 corresponds to Figure 4_figure Supplement 2.

1178

1179 **Figure 4_figure Supplement 3_source data 1.** The numerical source data

1180 corresponds to Figure 4_figure Supplement 3.

1181

1182

1183

1184

Supplementary Table 1 Strains and plasmids used in this study

Plasmid or strain	Genotype or description	Source
Plasmid		
pSim17	temperature-induced λ-Red recombinase system; BlaS^R	Laboratory collection
pKD3	template plasmid containing the Cm cassette for λ-Red recombination; Cm^R	Laboratory collection
pKD4	template plasmid containing the Km cassette for λ-Red recombination; Km^R	Laboratory collection
pBR322	low-copy-number expression vector; Ap^R	Laboratory collection
pMS402	Containing a promoterless *luxCDABE* reporter gene cluster; Km^R	Laboratory collection
pBR322-*panD*	pBR322 carrying the STM *panD* gene; Ap^R	This study
pMS402-*panD*	pMS402 carrying the STM *panD* promoter; Km^R	This study
strain		
wild-type (WT) STM	wild-type *Salmonella* enterica serovar Typhimurium strain ATCC 14028s	Laboratory collection
Δ*panD*	WT strain *panD*::Cm; Cm^R	This study
c*panD*	Δ*panD* containing plasmid pBR322-*panD*; Cm^R , Ap^R	This study
Δ*cycA*	WT strain cycA::Cm; Cm^R	This study
Δ*fadAB*	WT strain *fadAB*::Cm; Cm^R	This study
Δ*metR*	WT strain *metR*::Cm; Cm^R	This study
Δ*leuABCD*	WT strain *leuABCD*::Cm; Cm^R	This study
Δ*leuO*	WT strain *leuO*::Km; Km^R	This study
Δ*hisABCDFGHL*	WT strain *hisABCDFGHL*::Cm; Cm^R	This study
Δ*kdpABC*	WT strain *kdpABC*::Cm; Cm^R	This study
Δ*mglABC*	WT strain *mglABC*::Cm; Cm^R	This study
Δ*potFGHI*	WT strain *potFGHI*::Cm; Cm^R	This study
Δ*znuA*	WT strain *znuA*::Cm; Cm^R	This study

1186

1187

 Supplementary Table 2 Primers involved in this study

Targets		Primer Sequences (5′-3′)
Primers used for the construction and identification of mutant strains		
Δ*cycA*	construction	F: GTATCATAGACCAAAGGCCGTAGAGCCCGCACAACACAG ACAGGTACAGGAAGAAAAACGTGTAGGCTGGAGCTGCTTC R: CCTCCAAATCAACGTTACGCTGTAAGCCCGGTAAGCGCC AGCGCCACCGGGCAAAACAACATATGAATATCCTCCTTAG
	identification	F: TGTGAGCTAATTCGCATTATCAAAG R: ACGGCATTAATGAGATGATTGATGA
Δ*panD*	construction	F: CGGATTCGCTGGAAACCATGTCGCGGCTGATCAGCAACC AGCCGCAGGGATAAGGACTAGTGTAGGCTGGAGCTGCTT R: TTTCCAAAGCAGGCCACAAGCGCCTGCTTAGCCAAGGTA AACGACAGGGTAAAGAAGTTCATATGAATATCCTCCTTAG
	identification	F: CGGTTATAACGCGACCGCGATCAAA R: GCTGCTGGAGCTGACGGAAACCAGC
Δ*fadAB*	construction	F: GGATTCGCTCAGTTGCCGCTGCGCTGCAATGCGAGTTATT TAGGGGATATTATCTTTGAGTGTAGGCTGGAGCTGCTTC R: AATACACACTTCGCTTCATCTGGTACGACCAGATCACTTT GTGGATTCAGGAGACTGACCATATGAATATCCTCCTTAG
	identification	F: AAACGGCAACTAAACTGTTTCCCGT R: AGTGATTCCATTTTTTACCCTTCTG
Δ*metR*	construction	F: TAAAGCTTGGCCAGTGGGGCTGCTGATGCCAGATGGCAA GATTAACCAGCAGTCCTGCGGTGTAGGCTGGAGCTGCTTC R: GAAAGTCCTTCACTTCGCCATGAACAAATTGCGCTTGAG GAATATACAGTACCTTTACACATATGAATATCCTCCTTAG
	identification	F: CCATTGCAGCGTGGTTTCCGGCAGC R: TGCAAGCCGAACTATTTAGAGTTTT
Δ*leuABCD*	construction	F: GAATATAATGGCAACCGGGTGATATTGCATAACCTGTAG GCCCGGTAAGGCGTCAAGGGGTGTAGGCTGGAGCTGCTTC R: CCGTTGCGCGGGTTTTTTTATGCCTGACGCAAGGCGCCCC TGGAGACAAGGACCACATCCATATGAATATCCTCCTTAG
	identification	F: TTTTTTATTTCGGCTGTTTCTGGAT R: GTTGACATTAAACGGCATATCCAGT
Δ*leuO*	construction	F: CCAGAAAAAGGGAGTTAAGCGTGACAGTGGAGTTAAATGTG TAGGCTGGAGCTGCTTC R: ATAAACCAGAATTTGTTTCTGATTTATTCTGCCCGGTTCATATG AATATCCTCCTTAG
	identification	F: CATTATGAATCGCAATGGTGTGAC R: CTCCGTCTGAATCACACCTGGT

 Supplementary Table 2 Primers involved in this study (continue)

Targets		Primer Sequences (5′-3′)
		Primers used for the construction and identification of mutant strains
Δ*hisABCDFGHL*	construction	F: GGCGTAAAAGTGGTTTAGGTTAAAAGGTATCAAATGAATA AGCATTCATCGGAATTTTTGTGTAGGCTGGAGCTGCTTC
		R: GATGGCTGGCATCAGGCCATCGGTTTTTTCCCAGTCCAGC TCGCGGCGTTGTTGCTCTGCATATGAATATCCTCCTTAG
	identification	F: CCTAACCAACCTAAACCGACAATTG
		R: AAACGCTGTTTCGTGCGTGAGAAGA
Δ*kdpABC*	construction	F: GTGGGGCGGGGGCGTTTGTTCCAACAAGCGATCCGGATC AGGACGCAAGGGTTCGTCGTGTGTAGGCTGGAGCTGCTTC
		R:CTGGTTTTCTTACTTTTAGGTTATCTGGTCTATGCCCTGATTAAT GCGGAGGCGTTCTGCATATGAATATCCTCCTTAG
	identification	F: CAGAATGGTCAGCCCGTCAAG
		R: ATACTTTTTTTACACTCCGCCC
Δ*mglABC*	construction	F: CCGGCGCGCAGTTTCGCTGCGCCGGGTTATCTGATAAGC AAGGCAATAGGTCTGGATAAGTGTAGGCTGGAGCTGCTTC
		R: GTGCTGAACAGCCGGGCATTTTTTTACGCTATACCCTACA TAATAAAACCGGAGCTACCCATATGAATATCCTCCTTAG
	identification	F: AAAAAGAGCACGCGTTAACACTGTA
		R: AGGCTCTAAATACGCTTCGGC
Δ*potFGHI*	construction	F: ATGTTTTAAACCACGCCTAATGGGTTCATTTGTTAACGGAT TTCAGAAGGAAAGCGATGGTGTAGGCTGGAGCTGCTTC
		R: GAAAATGATGCCGCGACCCGCGCGGCAACGCATTGCCAT AGTGGAAGATTTTAGTGGCTCATATGAATATCCTCCTTAG
	identification	F: AAAGGCGTTTTTTAATCTGGGCTAT
		R: TATTCATAATCATCAGCACATCGAG
Δ*znuA*	construction	F: AGATTATTAAATGCCAGGGCGACAGAGCGGGCTATCTGTT GCACGTATTCACTTCCTCGGTGTAGGCTGGAGCTGCTTC
		R: AAGTGATAGAATGTTATAATATCACATTTCACACATTCATT ACGATGATTAGTCGCATTCATATGAATATCCTCCTTAG
	identification	F: AGGGAACGAATCTCGCTTTTCTC
		R: ATTCAAGCGACACGTCAGAGAGGAC
		Primers used for the construction of complementary strains
pBR322-*panD*		F: CCCTTTCGTCTTCAAGAATTCGCTGCTGGAGCTGACGGAAAC
		R: TGCGTCCGGCGTAGAGGATCCTCAGGCAACCTGTACCGGAAT
pBR322		F: GACACGGAAATGTTGAATACTCATA
		R: CAAGGAATGGTGCATGCAAG

 Supplementary Table 2 Primers involved in this study (continue)

Targets	Primer Sequences (5'-3')
Primers used for luciferase reporter system	
pMS402-*panD*	F: CGAGGCCCTTTCGTCTTCACCTCGAGTCAAGGAAATCCTGGTCAATG
	R: ATTTTGCGGCCGCAACTAGAGGATCCCGGGTAATCGTGAGCTGC
pMS402	F: CTGTCTCTTGATCAGATCTT
	R: TTATTTCCATTACAATCAAT
Primers used for RT-qPCR analysis	
16s	F:GAAAGCGTGGGAGCAAAC
	R:ACATGCTCCACCGCTTGTG
panD	F:AAGGTTCCTGTGCCATTGAC
	R:ACCGTTCACCGAGATGATTC
fadH	F:GAAACCCACATGCCGATAAC
	R:AAACTGACCGCCAATTTCAC
fadB	F:GTTCGCTGTCTGGAAGAAGG
	R: TACTGCTGCGCCATATCAAG
fadI	F:TCTGACGCTTTTTGACATGC
	R:TTCATGCAACGTTTGGGTTA
fadD	F:AAGATCTGGCTTTCCTGCAA
	R:CCCAGCTCGATAAAAAGCAG
fadE	F:AATTGGTCGTCGTCACTTCC
	R: GCGTCCTACCGACAAACATT
metR	F:TTTGGTGATGACGTCGGATA
	R:CGGGTAAATCAACAGCGTTT
hisG	F:GTGCGTGATGATGACATTCC
	R:GGAGGTGCGGATATGAGGTA
hisF	F:TGTATTGTCGTCGGGATTGA
	R:TTTTCAACTGCGTCAGATCG
leuD	F:CCGGAATTCGTGTTGAACTT
	R:GTAGAAGATGTCGGCGAAGC
leuO	F:AGATATGGGCAAACCACAGC
	R: CCTGACGGACTGAACCAAAT
mglA	F:AGCGATTAACCACGGTTTTG
	R:ATTGGGTGTCGCTTTTCATC
mglB	F:AGAATCCGGTGTGATTCAGG
	R:CAGCCAGGCATCCATCTTAT
kdpA	F:ACTAATCTGGCGCAGATGCT
	R:ACCCTCCATGTTGACGCTAC
kdpB	F:TCGGTAAGCAGATGCTGATG
	R:CCAGCGGGATCAGAAAAATA

1193

1194

1195

1196 **Supplementary Table 2 Primers involved in this study (continue)**

Targets	Primer Sequences (5′-3′)
Primers used for RT-qPCR analysis	
potG	F:ATTTACGAGCATCCGACGAC
	R:AGAGGCATCTGCATCGACTT
potF	F:TCTGATGGTCGTCTCTGTCG
	R:GCCTTCCAGCACTTCGTTAG
znuA	F:TTAAACCGCTTGGGTTCATC
	R:AACGGTTTTACATCGGCAAG

1197

A

Liver
Spleen
Bacterial CFU determination

0   3   5

Immunofluorescence staining of liver slice

H&E staining of liver slice

i.p. 5,000 CFU

Euthanize

0   5   10

Mice were observed for 10 days for survival and body weight was documented.

B

Survival (%)

WT
ΔpanD
cpanD

P < 0.0001
P < 0.0001

Days post infection

C

Body weight (g)

WT vs ΔpanD P < 0.0001

WT
ΔpanD
cpanD

Days post infection

D

$\log_{10}$(CFU/g liver)

P < 0.0001
P < 0.0001

WT   ΔpanD   cpanD

$\log_{10}$(CFU/g spleen)

P < 0.0001
P < 0.0001

WT   ΔpanD   cpanD

3dpi body weight (g)

P < 0.0001
P = 0.0001

WT   ΔpanD   cpanD

E

Salmonella   Nucleus   Merge

WT

ΔpanD

cpanD

Bacteria per Macrophage (liver)

P < 0.0001
P < 0.0001

WT   ΔpanD   cpanD

F

Uninfect   WT   ΔpanD   cpanD

A

Zinc level/ 10^5 macrophages (RAW264.7 cell)

P < 0.0001

WT $\Delta panD$

B

Fold intracellular replication (20 h $vs.$ 2 h)

P = 0.0002
P = 0.0004
P < 0.0001
P = 0.0422

WT $\Delta panD$ $\Delta znuA$ $\Delta panD\Delta znuA$

C

Fold intracellular replication (20 h $vs.$ 2 h)

P = 0.0020
ns

WT $\Delta panD$ $\Delta panD$+100μM ZnSO$_4$

D

Fold intracellular replication (20 h $vs.$ 2 h)

P = 0.0028
P = 0.0027

WT $\Delta znuA$ $\Delta znuA$+1mM β-alanine

E

Macrophage

Zn²⁺

SCV

β-alanine → Uptake by *Salmonella* → *Salmonella*

Promote *Salmonella* growth

SCV

Dissemination and systemic infection

L-aspartate
PanD
β-alanine → β-alanine → ZnuABC ↑
Zn²⁺ uptake

Macrophage
SCV (*Salmonella*-containing vacuole)
Salmonella
Zn²⁺
Activation
Upregulation