Salmonella **exploits host- and bacterial-derived β-alanine for**

replication inside host macrophages

Shuai Ma[1†], Bin Yang[1†], Yuyang Sun[1], Xinyue Wang[1], Houliang Guo[1],

Ruiying Liu[1], Ting Ye[1], Chenbo Kang[1], Jingnan Chen[1], Lingyan Jiang[1*]

[1]National Key Laboratory of Intelligent Tracking and Forecasting for Infectious

Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai

University, Tianjin 300457, China.

[†] Shuai Ma and Bin Yang contributed equally to this work.

[*]Correspondence and requests for materials should be addressed to Lingyan Jiang

(email: jianglingyan@nankai.edu.cn).

Abstract

Salmonella is a major foodborne pathogen that can effectively replicate inside host macrophages to establish life-threatening systemic infections. *Salmonella* must utilize diverse nutrients for growth in nutrient-poor macrophages, but which nutrients are required for intracellular *Salmonella* growth is largely unknown. Here, we found that either acquisition from the host or *de novo* synthesis of a nonprotein amino acid, β-alanine, is critical for *Salmonella* replication inside macrophages. The concentration of β-alanine is decreased in *Salmonella*-infected macrophages, while the addition of exogenous β-alanine enhances *Salmonella* replication in macrophages, suggesting that *Salmonella* can uptake host-derived β-alanine for intracellular growth. Moreover, the expression of *panD,* the rate-limiting gene required for β-alanine synthesis in *Salmonella,* is upregulated when *Salmonella* enters macrophages. Mutation of *panD* impaired *Salmonella* replication in macrophages and colonization in the mouse liver and spleen, indicating that *de novo* synthesis of β-alanine is essential for intracellular *Salmonella* growth and systemic infection. Additionally, we revealed that β-alanine influences *Salmonella* intracellular replication and *in vivo* virulence partially by increasing expression of the zinc transporter genes *znuABC*, which in turn facilitates the uptake of the essential micronutrient zinc by *Salmonella*. Taken together, these findings highlight the important role of β-alanine in the intracellular replication and virulence of *Salmonella*, and *panD* is a promising target for controlling systemic *Salmonella* infection.

Keywords: *Salmonella*, replication in macrophage, virulence, β-alanine, zinc

37   uptake

Introduction

Salmonella is a major foodborne pathogen worldwide that can cause self-limiting gastroenteritis or life-threatening systemic disease in a wide range of animals (Fàbrega & Vila, 2013; Ohl & Miller, 2001). *Salmonella* infection remains a significant global public health concern. An estimated 93.8 million cases of gastroenteritis and 27 million cases of systemic diseases caused by *Salmonella* species occur annually worldwide, with 355,000 deaths (Kim et al., 2019; Majowicz et al., 2010). The ability to survive and replicate in host macrophages is a key determinant for *Salmonella* to induce systemic infection (Fields, Swanson, Haidaris, & Heffron, 1986; LaRock, Chaudhary, & Miller, 2015; Leung & Finlay, 1991). After internalization by macrophages, *Salmonella* delivers a set of more than 30 effector proteins to the macrophage cytoplasm, mainly through a type III secretion system (T3SS) encoded by *Salmonella* pathogenicity island-2 (SPI-2) (Pillay et al., 2023). SPI-2 effectors manipulate diverse cellular processes to promote the formation of a membrane-bound compartment, termed the *Salmonella*-containing vacuole (SCV), a niche where *Salmonella* resides and grows (Castanheira & García-Del Portillo, 2017; Rosenberg, Riquelme, Prince, & Avraham, 2022; Steeb et al., 2013). SCV protects *Salmonella* from contact with antimicrobial agents in macrophages (Figueira & Holden, 2012; Li et al., 2023). Moreover, SPI-2 effectors induce the formation of specific tubular membrane compartments that extend from the SCV, known as *Salmonella*-induced filaments (SIFs). These filaments allow *Salmonella* to access various types of endocytosed nutrients, thereby facilitating efficient replication

within macrophages (Brumell, Tang, Mills, & Finlay, 2001; Liss et al., 2017;

Rajashekar, Liebl, Seitz, & Hensel, 2008).

As the SCV of macrophages is a nutrient-poor environment (Fields et al., 1986;

Kehl, Noster, & Hensel, 2020), to effectively replicate in the SCV, *Salmonella* needs

to acquire a wide range of host nutrients or host-derived metabolites and synthesize

metabolites *de novo* that cannot be sufficiently accessed from the host (Dandekar et

al., 2014; Röder, Felgner, & Hensel, 2021; Tuli & Sharma, 2019).

Nutrients/metabolites are used by intracellular *Salmonella* either as carbon sources to

generate energy or for the synthesis of fatty acids and proteins (Dandekar et al., 2014;

Steeb et al., 2013). Moreover, several metabolites were found to be employed by

Salmonella as environmental cues to induce the expression of virulence genes (Jiang

et al., 2021; X. Wang et al., 2023). In recent years, an increasing number of studies

have focused on the intracellular nutrition of *Salmonella* (Bumann & Schothorst, 2017;

Liss et al., 2017; Röder et al., 2021); however, the nutrients that are required for

Salmonella replication in macrophages remain largely unknown.

β-Alanine, also known as 3-aminopropionic acid (3-AP), is the only naturally

occurring β-type amino acid and is found in all living organisms (Song, Zhang, Wang,

Xu, & Wei, 2023). β-Alanine can be synthesized *de novo* by bacteria, fungi, and

plants, whereas animals need to obtain it from food or generate it via the catabolism

of cytosine and uracil (Gojković, Sandrini, & Piskur, 2001; L. Wang, Mao, Wang, Ma,

& Chen, 2021). In bacteria, β-alanine is synthesized *via* the decarboxylation

of L-aspartate, a reaction catalyzed by L-aspartate decarboxylase (PanD) (Begley,

Kinsland, & Strauss, 2001; Schmitzberger, Smith, Abell, & Blundell, 2003; West,

Traut, Shanley, & O'Donovan, 1985). The *panD* gene is conserved among most

bacteria (Nozaki, Webb, & Niki, 2012). Although β-alanine is a nonprotein amino

acid that is not incorporated into proteins, it has important physiological functions in

the metabolism of organisms (L. Wang et al., 2021; Yuan et al., 2022). First, β-alanine

forms a part of pantothenate (vitamin B5), which is the key precursor for the

biosynthesis of coenzyme A (CoA) (Webb, Smith, & Abell, 2004; White, Gunyuzlu,

& Toyn, 2001). CoA is an essential cofactor involved in many metabolic pathways,

including the synthesis and degradation of fatty acids, pyruvate oxidation through the

tricarboxylic acid (TCA) cycle, and the production of secondary metabolites

(Davaapil, Tsuchiya, & Gout, 2014; Gout, 2019; Sibon & Strauss, 2016; Theodoulou,

Sibon, Jackowski, & Gout, 2014). Second, β-alanine is a limiting precursor of

carnosine, a nonenzymatic free radical scavenger and a natural antioxidant, with

anti-inflammatory and neuroprotective effects in animals (Boldyrev, Aldini, & Derave,

2013; Hoffman, Varanoske, & Stout, 2018). In the past 15 years, β-alanine has

become one of the most commonly used sports supplements worldwide (Bellinger,

2014; Hoffman et al., 2018; Huerta Ojeda, Tapia Cerda, Poblete Salvatierra,

Barahona-Fuentes, & Jorquera Aguilera, 2020). Although both *Salmonella* and host

cells are capable of producing β-alanine, whether β-alanine contributes to the

pathogenicity and intracellular growth of *Salmonella* remains unknown.

In this work, using targeted metabolic profiling, *in vitro* and *in vivo* infection

assays, and many other molecular techniques, we demonstrated that the utilization of

β-alanine is essential for *Salmonella* replication in host macrophages and virulence in

mice. *Salmonella* acquires β-alanine both via the uptake of β-alanine from host

macrophages and the *de novo* synthesis of β-alanine. Further investigation revealed

the molecular mechanism underlying the contribution of β-alanine to *Salmonella*

intracellular replication and pathogenicity, wherein β-alanine promotes the expression

of zinc transporter genes to facilitate the uptake of the essential micronutrient zinc by

intracellular *Salmonella,* therefore promoting *Salmonella* replication in macrophages

and subsequent systemic infection. Taken together, these findings demonstrate a

correlation between *Salmonella* β-alanine utilization and zinc uptake during

intracellular infection and provide new insights into the intracellular nutrition of

Salmonella. The rate-limiting gene (*panD*) in the β-alanine synthesis pathway of

Salmonella might be a future target for the prevention and treatment of this pathogen.

Results

Host-derived β-alanine promotes *Salmonella* replication inside macrophages

To explore changes in the levels of different amino acids inside macrophages upon *Salmonella* infection, we performed targeted metabolomics analysis of mouse RAW264.7 macrophages that were mock-infected or infected with wild-type *Salmonella* (*Salmonella enterica* serovar Typhimurium ATCC 14028s, STM) for 8 h using liquid chromatography-tandem mass spectrometry (LC□MS/MS) (Figure 1A). Principal component analysis (PCA) demonstrated a clear separation between the mock- and *Salmonella*-infected groups (Figure 1B). A total of 26 free amino acids were analyzed, and 8 showed significant differences in abundance between the two groups (VIP (Variable Importance in the Projection) > 1 and P < 0.05; FC (fold change) >1.5 or<0.667) (Figure 1C，D). Compared with those in the mock-infected group, the concentrations of 3 amino acids (L-hydroxyproline, L-citrulline and L-cysteine) were upregulated (Figure 1C), and 5 amino acids (L-asparagine, L-serine, L-aspartate, β-alanine and γ-aminobutyric acid) were downregulated (Figure 1D) in the *Salmonella*-infected group. Consistent with previous findings, intracellular serine concentrations were downregulated due to the reprogramming of macrophage glucose metabolism during *Salmonella* infection (Jiang et al., 2021). *Salmonella* can use host-derived aspartate and asparagine for growth in macrophages (Popp et al., 2015); therefore, the decrease in intracellular aspartate and asparagine upon *Salmonella* infection is likely due to their utilization by bacteria. Interestingly, β-alanine concentrations were also downregulated in the *Salmonella*-infected group (Figure 1D),

suggesting that intracellular *Salmonella* may use host-derived β-alanine for growth.

To investigate whether host-derived β-alanine can promote intracellular

Salmonella replication, we added an additional 0.5, 1, 2, 4 mM β-alanine (Schneider,

Krämer, & Burkovski, 2004) to the culture medium (RPMI) of RAW264.7 cells and

then infected them with *Salmonella* to test the influence of β-alanine addition on the

ability of *Salmonella* to replicate in macrophages. The results showed that the

replication of *Salmonella* in RAW264.7 cells significantly ($P < 0.001$) increased with

the addition of 1, 2, or 4 mM β-alanine (Figure 1E). Furthermore, β-alanine enhanced

Salmonella intracellular replication in a dose-dependent manner (Figure 1E). The

results suggest that host-derived β-alanine facilitates *Salmonella* replication inside

macrophages. We then investigated whether β-alanine-mediated *Salmonella* growth

promotion is due to the changes in antimicrobial activity of the macrophages. We

observed that the addition of 1 mM β-alanine did not influence the ROS (reactive

oxygen species) and RNS (reactive nitrogen species) levels in *Salmonella*-infected

RAW264.7 cells (Figure 1_figure Supplement 1). Flow cytometry analysis indicated

that the addition of 1 mM β-alanine did not affect the percentage of pro-inflammatory

M1 macrophages (CD86+) and anti-inflammatory M2 macrophages (CD163+) during

Salmonella infection (Figure 1_figure Supplement 2), implying that the addition of

β-alanine to macrophages does not change their immune response. Combining these

results, we can further infer that *Salmonella* use host-derived β-alanine for

intracellular growth.

Direct validation of *Salmonella* using host-derived β-alanine for intracellular

growth requires a mutant that has a defect in β-alanine uptake. *Escherichia coli*

uptakes β-alanine via the transporter protein CycA (Schneider et al., 2004). However,

the *Salmonella* Δ*cycA* mutant was able to use β-alanine as the sole carbon source for

growth in minimal medium (Figure 1_figure Supplement 3, Figure 1_figure

Supplement 4), indicating that CycA is not a transporter for β-alanine in *Salmonella*.

Consistent with these results, mutation of *cycA* did not influence the replication of

Salmonella in RAW264.7 cells (Figure 1_figure Supplement 5) or colonization in

mouse systemic tissues (liver and spleen; Figure 1_figure Supplement 6). In *E. coli*,

GabP transports γ-aminobutyric acid (GABA), a structural analogue of β-alanine, and

may also transport β-alanine (Pavić et al., 2021). Nevertheless, *Salmonella* Δ*gabP*

mutant displayed no growth defect in minimal medium with β-alanine as the sole

carbon source (Figure 1_figure Supplement 7, Figure 1_figure Supplement 8),

indicating that GabP is not involved in β-alanine uptake in *Salmonella*. Strikingly, the

Δ*argT* mutant—defective in arginine uptake—showed markedly decreased growth in

the minimal medium with β-alanine as the sole carbon source (Figure 1F), suggesting

that ArgT also transports β-alanine in *Salmonella*. It has been reported that ArgT is

essential for *Salmonella* replication within macrophages and full virulence *in vivo*

(Das et al., 2010). Given that ArgT is involved in both arginine and β-alanine uptake

(as verified in this study), whether the attenuated virulence of the Δ*argT* mutant is due

to a deficiency in β-alanine or arginine requires further investigation.

180   ***De novo* β-alanine synthesis is critical for *Salmonella* replication inside**

181   **macrophages**

182   *Salmonella* can *de novo* synthesize β-alanine *via* the decarboxylation

183   of L-aspartate, which is catalyzed by L-aspartate decarboxylase (PanD) (Figure 2A)

184   and is reportedly the rate-limiting step of β-alanine generation (Begley et al., 2001;

185   Schmitzberger et al., 2003; West et al., 1985). To further assess the role of β-alanine in

186   *Salmonella* intracellular replication, we analyzed the expression level of the

187   *Salmonella panD* gene in macrophages and the impact of *panD* mutation on the

188   ability of *Salmonella* to replicate in macrophages. Quantitative real-time PCR

189   (qRT□PCR) assays revealed that the expression level of *panD* was significantly ($P <$

190   0.01) greater in RAW264.7 cells than in RPMI-1640 medium (Figure 2B). Increased

191   expression of *panD* was also observed in N-minimal medium, a widely used medium

192   that mimics the conditions inside macrophages, as revealed by qRT□PCR and

193   bioluminescent reporter assays (Figure 2C, D). These results demonstrate that *panD*

194   expression is enhanced during *Salmonella* growth inside macrophages, suggesting a

195   relationship between *panD* expression and intracellular *Salmonella* growth.

196   We then constructed the *panD* mutant strain Δ*panD* and compared the replication

197   ability of the Δ*panD* strain and the *Salmonella* Typhimurium 14028s wild type (WT)

198   strain in RAW264.7 cells. Gentamicin protection assays showed that the replication of

199   Δ*panD* in RAW264.7 cells decreased 2.4-fold at 20 h postinfection compared with

200   that of the WT strain ($P <$ 0.001), while complementation of Δ*panD* with the *panD*

201   gene restored the replication ability of the mutant strain in RAW264.7 cells (Figure

2E). Immunofluorescence analysis revealed that the number of Δ*panD* in each

infected RAW264.7 cell was comparable to that of the WT strain at the initial

infection stage (2 h), but at 20 h postinfection, the number of Δ*panD* in each infected

RAW264.7 cell was significantly ($P < 0.0001$) lower than that of the WT strain

(Figure 2F, G). These results indicate that *panD* contributes to *Salmonella* replication

in macrophages. The growth rates of Δ*panD* in LB medium and RPMI medium

resembled those of the WT (Figure 2_figure Supplement 1, Figure 2_figure

Supplement 2), indicating that the impaired intracellular replication ability of the

mutant was not due to a growth defect. Moreover, the replication defect of Δ*panD* in

RAW264.7 cells was relieved by the addition of 1 mM β-alanine to the RPMI medium

(Figure 2H). Furthermore, we examined the role of β-alanine synthesis in the

intracellular replication of *Salmonella* within another typical serovar, *Salmonella*

enterica serovar Typhi (*S*. Typhi), a serovar specific to humans and the causative

agent of typhoid fever (de Jong, Parry, van der Poll, & Wiersinga, 2012). We found

that the replication of *S*. Typhi Δ*panD* in human THP-1 macrophages was reduced by

2.6-fold compared to the the *S*. Typhi Ty2 WT strain ($P < 0.01$) (Figure 2_figure

Supplement 3), suggesting that *panD* also facilitates *S*. Typhi replication within

human macrophages.

These data collectively suggest that β-alanine synthesis is critical for *Salmonella*

replication inside macrophages.

***De novo* β-alanine synthesis is critical for systemic *Salmonella* infection in mice**

As replication in macrophages is a key determinant of systemic *Salmonella*

infection, we reasoned that β-alanine synthesis also influences *Salmonella* systemic

infection *in vivo*. To determine whether β-alanine influences systemic *Salmonella*

infection, we conducted mouse infection assays using intraperitoneal (i.p.) injection.

This method allows *Salmonella* to disseminate directly to systemic sites *via* the

lymphatic and bloodstream system, bypassing the need for intestinal invasion

(Pandeya et al., 2023; Silva et al., 2016). BALB/c mice were infected by i.p. injection

of 5,000 CFU of WT, Δ*panD*, or the complemented strain c*panD*. The survival rate,

body weight, bacterial burden in the liver and spleen, and liver histopathological

alterations of the infected mice were measured (Figure 3A). The WT-infected mice

exhibited high lethality and marked loss of body weight within 5 days, and all mice

died within 9 days of infection (Figure 3B, C). In contrast, the Δ*panD*-infected mice

displayed significantly improved survival rates and body weights, and no mice died

within the 10-day surveillance period (Figure 3B, C). Consistent with these results,

the bacterial burden in the liver and spleen of the Δ*panD*-infected mice was

significantly decreased, and the body weight was significantly increased compared

with that of the WT-infected mice on day 3 postinfection (Figure 3D).

Complementation of Δ*panD* with *panD* significantly decreased the survival rate and

body weight of infected mice but significantly increased the bacterial burden in the

liver and spleen of the infected mice (Figure 3B-D). Through immunofluorescence

staining, we examined the bacterial count in liver macrophages of mice infected with

WT, Δ*panD*, and complemented strain. The results showed that the bacterial count in

each macrophage from Δ*panD*-infected mice was significantly ($P < 0.0001$) lower

than that in WT-infected mice, on day 5 post-infection. Complementation of Δ*panD*

with *panD* restored the bacterial count in each macrophage to WT level (Figure 3E).

Furthermore, H&E staining revealed increased aggregation of inflammatory cells and

pyknosis in the livers of the WT-infected mice on day 5 postinfection, while these

histopathological alterations were obviously reduced in the livers of Δ*panD*-infected

mice (Figure 3F). Taken together, these results reveal that β-alanine synthesis is

critical for systemic *Salmonella* infection in mice.

β-Alanine is involved in the regulation of several metabolic pathways in

Salmonella

To explore the mechanism(s) associated with β-alanine-mediated promotion of

Salmonella replication in macrophages and *in vivo* virulence, we performed RNA

sequencing (RNA-seq) to reveal the differences in gene transcripts between

Salmonella WT and Δ*panD* strains grown in N-minimal medium. PCA plot of the

global transcriptomic profiles clearly demonstrated separation between the WT and

Δ*panD* strains (Figure 4A). Remarkable transcriptional changes were observed due to

the mutation of *panD*. Compared with those in the WT strain, 1379 genes were

differentially expressed in the Δ*panD* strain, with 561 upregulated genes and 618

downregulated genes (fold change ≥ 2 and *P* value < 0.05; Figure 4B). Gene Ontology

(GO) enrichment analysis revealed that the differentially expressed genes (DEGs)

were mainly involved in the metabolism and biosynthesis of several amino acids

(including arginine, leucine, histidine, and branched amino acids), carboxylic acid

metabolism, small molecule biosynthesis, and aerobic respiration (Figure 4C). Kyoto

Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis also

revealed a high frequency of terms related to metabolism, including amino acid

metabolism, lipid metabolism, carbohydrate metabolism, energy metabolism, and

nucleotide metabolism (Figure 4D). These data collectively indicate that β-alanine is

involved in the regulation of a series of metabolic pathways in *Salmonella.*

Further analysis of the downregulated DEGs (activated by PanD) revealed that

mutation of *panD* decreased the expression of genes involved in 7 pathways that are

associated with the virulence of *Salmonella* or other bacterial pathogens, including

methionine metabolism, fatty acid β-oxidation, histidine biosynthesis, and the

transport of zinc, galactose, potassium, and polyamine (Figure 4E). Zinc and

potassium uptake are associated with the virulence of *Salmonella* (zinc acquisition

promotes *Salmonella* Typhimurium virulence in mice, and potassium acquisition

promotes *Salmonella* Enteritidis virulence in chickens) (Ammendola et al., 2007;

Battistoni, Ammendola, Chiancone, & Ilari, 2017; Ilari et al., 2016; Liu et al., 2013),

while the remaining 5 pathways are involved in the pathogenicity of other pathogens

(Basavanna et al., 2013; de Paiva et al., 2016; Feldman et al., 2016; Lauriano et al.,

2004; Martínez-Guitián et al., 2019). In addition, the expression of the LysR-type

transcriptional regulator LeuO, which activates the expression of the *leuABCD*

leucine synthesis operon and numerous virulence genes in *Salmonella* Typhimurium

(Dillon et al., 2012; Guadarrama, Villaseñor, & Calva, 2014; Hernández-Lucas et al.,

2008), was also downregulated in the ΔpanD strain (Figure 4E). In line with the

decreased expression of *leuO*, the expression of *leuABCD* was downregulated in the

ΔpanD strain (Figure 4E).

We selected 16 downregulated DEGs (including the regulatory gene *leuO* and

genes from the above 7 pathways) for qRT□PCR analysis. The results showed that

the expression of all 16 genes significantly ($P < 0.05$) decreased in the ΔpanD mutant

compared with the WT strain (Figure 4F), and complementation of ΔpanD with *panD*

restored the gene expression to the WT level (Figure 4F), thus confirming the positive

regulation of these pathways and LeuO by β-alanine.

Salmonella pathogenesis largely depends on virulence genes encoded by

Salmonella pathogenicity islands (SPIs), with SPI-1 to SPI-5 being well-characterized

for their involvement in *Salmonella* virulence (Han et al., 2024). SPI-2 gene

expression is essential for *Salmonella* replication in macrophages and systemic

infection (Deng et al., 2017), yet the expression of SPI-2 genes was unaffected by

panD mutation (Figure 4_figure Supplement 1). Moreover, the gene expression of

four other virulence-associated SPIs, SPI-1, SPI-3, SPI-4, and SPI-5, is also

unaffected by *panD* mutation (Figure 4_figure Supplement 2, Figure 4_figure

Supplement 3).

Taken together, these data suggest that β-alanine might promote *Salmonella*

intracellular replication and virulence by activating virulence-associated pathway(s)

or activating the virulence-associated regulator LeuO, rather than by activating the

expression of virulence genes encoded within pathogenicity islands.

β-Alanine promotes *Salmonella* virulence *in vivo* partially by increasing the

expression of zinc transporter genes

Next, we inactivated the 7 downregulated pathways mentioned above, as well as

the regulatory gene *leuO* in *Salmonella,* to uncover the mechanism(s) by which

β-alanine promotes *Salmonella* virulence *in vivo*. Mouse infection assays revealed that

mutations in *fadAB*, *metR*, *hisABCDFGHL*, *kdpABC*, *mglABC*, and *potFGHI,* which

are associated with fatty acid β-oxidation, methionine metabolism, histidine

biosynthesis, potassium uptake, galactose uptake, and polyamine uptake, respectively,

did not influence *Salmonella* colonization in the mouse liver or spleen (Figure 5A, B)

or the body weight of infected mice (Figure 5C). Interestingly, although LeuO has

been reported to be associated with the regulation of a diverse set of virulence factors

(Dillon et al., 2012; Guadarrama et al., 2014), mutation of the regulatory gene *leuO*

did not influence *Salmonella* colonization in the mouse liver or spleen (Figure 5A, B)

or the body weight of infected mice (Figure 5C). In contrast, mutation of the zinc

transporter gene *znuA* significantly decreased *Salmonella* colonization in the mouse

liver and spleen (Figure 5D, left and middle panels); this result is consistent with

previous studies (Ammendola et al., 2007; Battistoni et al., 2017; Ilari et al., 2016).

Accordingly, the body weight of Δ*znuA*-infected mice was significantly (P < 0.001)

greater than that of the WT-infected mice (Figure 5D, right panel). These results

indicate that β-alanine might promote *Salmonella* virulence *in vivo* by promoting zinc

uptake.

To test this hypothesis, we initially evaluated the zinc content in the livers of

WT- and Δ*panD*-infected mice, on day 3 post-infection. We observed that the zinc

concentration in the macrophages of Δ*panD*-infected mouse livers was 3.2-fold higher

than in those of WT-infected mice ($P < 0.0001$; Figure 5E), suggesting that the *panD*

gene and β-alanine are crucial for *Salmonella* to obtain zinc from host cells.

Next, we constructed a double mutant, Δ*panD*Δ*znuA*, and compared colonization

of the mouse liver and spleen of the double mutant to that of the single mutant, Δ*znuA*.

The results showed that colonization of the liver and spleen of infected mice by

Δ*panD*Δ*znuA* was significantly lower than that of infected mice colonized by Δ*znuA*

(Figure 5D, left and middle panels). In agreement with these results, the body weight

of Δ*panD*Δ*znuA*-infected mice was greater than that of Δ*znuA*-infected mice (Figure

5D, right panel), suggesting that the contribution of *panD* to the virulence of

Salmonella is partially dependent on *znuA*.

Collectively, these data indicate that β-alanine promotes *in vivo* virulence of

Salmonella partially by increasing the expression of zinc transporter genes.

β-Alanine promotes *Salmonella* replication within macrophages partially by

increasing the expression of zinc transporter genes

To determine whether β-alanine influences *Salmonella* intracellular replication

by acting on zinc transporters, the zinc content in RAW 264.7 macrophages infected

with WT and Δ*panD* was also examined. We observed that the zinc concentration in

Δ*panD*-infected RAW 264.7 cells increased by 1.8-fold compared to WT-infected

cells ($P < 0.0001$; Figure 6A), further indicating the *panD* gene and β-alanine are

crucial for *Salmonella* to absorb zinc from macrophages.

We then analyzed the ability of Δ*znuA* and Δ*panD*Δ*znuA* to replicate in

RAW264.7 macrophages *via* gentamicin protection assays. The results showed that

the replication of Δ*panD*Δ*znuA* in RAW264.7 cells was significantly reduced

compared with that of the single mutant Δ*znuA* (Figure 6B), implying that the

contribution of *panD* to the intracellular replication of *Salmonella* is partially

dependent on *znuA*. The addition of 100 μM zinc to RPMI medium increased the

replication of Δ*panD* in RAW264.7 cells (Figure 6C), while the addition of 1 mM

β-alanine to RPMI medium did not increase the replication of Δ*znuA* (Figure 6D),

suggesting that the impaired replication due to the decrease in β-alanine can be

relieved by zinc supplementation.

Taken together, these data indicate that β-alanine promotes *Salmonella*

replication within macrophages by increasing the expression of zinc transporter genes.

Discussion

Replication within host macrophages is a crucial step for *Salmonella* to cause

life-threatening systemic infection in the host (Bomjan, Zhang, & Zhou, 2019;

Lathrop et al., 2015), while the crosstalk between *Salmonella* and macrophages at the

metabolic interface is critical for intracellular *Salmonella* replication (Dandekar et al.,

2014; Herrero-Fresno & Olsen, 2018; Lynch & Lesser, 2021; Rosenberg et al., 2021;

Thompson, Fulde, & Tedin, 2018). Emerging evidence suggests that several

metabolites affect the replication of *Salmonella* in macrophages. The promotion of

intracellular replication by metabolites is possibly achieved in three ways: i)

metabolites are utilized by *Salmonella* as nutrients for intracellular growth (Bowden,

Rowley, Hinton, & Thompson, 2009; Eisenreich, Dandekar, Heesemann, & Goebel,

2010; J. Wang et al., 2021); ii) *Salmonella* senses metabolites as environmental cues

to activate the expression of virulence genes (Jiang et al., 2021; X. Wang et al., 2023);

and iii) metabolites can regulate the immune responses of macrophages (Michelucci

et al., 2013; Peace & O'Neill, 2022; Yang & Cong, 2021). In this study, we

demonstrated that *Salmonella* promotes its replication inside macrophages by utilizing

both host- and bacterial-derived β-alanine (Figure 6E). We showed that β-alanine

promotes *Salmonella* intracellular replication and systemic infection partially by

increasing the expression of zinc transporter genes and therefore the uptake of zinc by

intracellular *Salmonella* (Figure 6E). Therefore, this work identified another

metabolite that can influence the replication of *Salmonella* in macrophages and

illustrated the mechanism by which β-alanine promotes intracellular *Salmonella*

replication.

We observed that *Salmonella*-infected macrophages contained lower β-alanine

levels than mock-infected macrophages, while β-alanine supplementation in the cell

medium increased the replication of *Salmonella* in macrophages, revealing that

Salmonella uptakes host-derived β-alanine to promote intracellular replication. In

addition, a deficiency in the biosynthesis of β-alanine (*via* mutation of the

rate-limiting gene *panD*) reduced *Salmonella* replication in macrophages and

systemic infection in mice, suggesting that *Salmonella* also utilizes bacterial-derived

β-alanine to promote intracellular replication and pathogenicity. It is known that

bacteria are quite stringent with their energy resources (Bergkessel, 2020; Bosdriesz,

Molenaar, Teusink, & Bruggeman, 2015), while the results of this work indicate that

either acquisition from the host or *de novo* synthesis of β-alanine is critical for

Salmonella replication inside macrophages. We speculate that *Salmonella* relies on a

large amount of β-alanine to efficiently replicate in macrophages, thereby highlighting

the importance of β-alanine for *Salmonella* intracellular growth. Nevertheless, unlike

the closely related species *E. coli*, which takes up β-alanine via the transporter protein

CycA (Schneider et al., 2004), *Salmonella* does not use CycA to uptake β-alanine.

GabP, the GABA transporter, is potentially involved in the uptake of β-alanine in *E.*

coli (Pavić et al., 2021). However, *Salmonella* does not utilize GabP to uptake

β-alanine either. Ultimately, we revealed that ArgT, the transporter of arginine, is

involved in the transport of β-alanine in *Salmonella*. ArgT has been reported to be

essential for *Salmonella* replication within macrophages and for full virulence *in vivo*

(Das et al., 2010). However, the attenuated virulence of the Δ*argT* mutant is due to a

deficiency in β-alanine or arginine requires further investigation.

Several amino acids, including lysine, proline, arginine, aspartate and asparagine,

have previously been reported to be associated with the pathogenicity of *Salmonella*

(Popp et al., 2015; Steeb et al., 2013). These amino acids are involved in the synthesis

of proteins in *Salmonella*. In contrast, β-alanine is not incorporated into proteins but

can participate in the regulation of bacterial activity through the synthesis of

pantothenate and CoA (Webb et al., 2004; White et al., 2001; Yuan et al., 2022).

Accordingly, our transcriptome data showed that a deficiency in β-alanine

biosynthesis affected the expression of *Salmonella* genes involved in a series of

important metabolic pathways. Importantly, although β-alanine does not influence the

gene expression of SPIs, it activates methionine metabolism; fatty acid β-oxidation;

histidine biosynthesis; and the transport of zinc, galactose, potassium, and polyamine,

which have been previously known to be associated with the virulence of *Salmonella*

and other bacterial pathogens. Further analysis revealed that β-alanine promotes

Salmonella intracellular replication and systemic infection partially by promoting the

uptake of zinc. As for the other virulence-related metabolic pathways activated by

β-alanine, methionine metabolism, fatty acid β-oxidation, and histidine biosynthesis

contribute to the virulence of *Streptococcus pneumoniae* (Basavanna et al., 2013),

Yersinia pestis (Feldman et al., 2016), and *Acinetobacter baumannii*

(Martínez-Guitián et al., 2019), respectively; the uptake of galactose and polyamine

influences the pathogenicity of *Francisella tularensis* (Lauriano et al., 2004) and the

avian pathogenic *Escherichia coli* (de Paiva et al., 2016)*,* respectively; and the uptake

of potassium is associated with the virulence of *Salmonella* Enteritidis in chickens

(Liu et al., 2013). However, blocking these pathways did not influence the systemic

Salmonella Typhimurium infection in mice, implying that different bacterial

pathogens adopt different virulence strategies to establish infection. Determining

other mechanism(s) by which β-alanine promotes the intracellular replication and

systemic infection of *Salmonella* require further investigation.

We observed that β-alanine also activates the expression of the LysR-type

transcriptional regulator LeuO, which is known to regulate the expression of a wide

variety of *Salmonella* genes that impact the stress response and virulence (Dillon et al.,

2012; Guadarrama et al., 2014; Hernández-Lucas et al., 2008). Typically, LeuO

activates the synthesis of the quiescent porins OmpS1 and OmpS2, which are required

for *Salmonella* virulence in mice (De la Cruz et al., 2007; Fernández-Mora, Puente, &

Calva, 2004; Rodríguez-Morales et al., 2006). Consistent with the positive regulation

of OmpS1 and OmpS2 by LeuO, lack of *leuO* in *Salmonella* also attenuated virulence

in a mouse model (Rodríguez-Morales et al., 2006). However, the attenuated

phenotypes of the *leuO* mutant in mice were not evident after i.p. injection relative to

oral infection, as a previous report showed that the competitive index for the *leuO*

mutant indicated approximately 1,000-fold reduced colonization in mouse systemic

tissues after oral infection but much less reduced colonization after i.p. injection (less

than 10-fold) (Rodríguez-Morales et al., 2006). These results imply that LeuO might

be predominantly associated with the invasion and intestinal infection of *Salmonella*

but is weakly implicated in *Salmonella* intracellular replication and systemic infection.

Therefore, it is not surprising that lack of *leuO* did not significantly affect *Salmonella*

colonization in the systemic tissues of mice after i.p. injection, as revealed by our

results.

Zinc is an essential micronutrient for all living organisms and is used as a

cofactor for various enzymes and proteins (Bock, Müller, & Blankenfeldt, 2016; Ilari

et al., 2016; Tan, Bramlett, & Lindahl, 2004). In bacteria, zinc-binding proteins

account for approximately 5% of the bacterial proteome and play crucial roles in

bacterial metabolism and virulence (Andreini, Banci, Bertini, & Rosato, 2006).

Knockout of the zinc transporter ZnuABC reduces the virulence of *Salmonella*,

Campylobacter jejuni, *Haemophilus ducreyi*, *Moraxella*, and urinary tract pathogenic

Escherichia coli (UPEC) in the host (Ilari et al., 2016). Moreover, zinc is also utilized

by *Salmonella* to subvert the antimicrobial host defense of macrophages by inhibiting

NF-κB activation and impairing NF-κB-dependent bacterial clearance (Jennings,

Esposito, Rittinger, & Thurston, 2018; Wu et al., 2017). Therefore, the efficient

acquisition of zinc may be crucial for *Salmonella*'s survival and replication within

macrophages, where zinc availability is limited (Ammendola et al., 2007; Ilari et al.,

2016). It has been reported that *Salmonella* utilizes the high-affinity ZnuABC zinc

transporter to optimize zinc availability in host cells (Ammendola et al., 2007). In this

study, we found that β-alanine can increase the expression of the zinc transporter

genes *znuABC*, which could represent an additional mechanism for the efficient zinc

uptake of *Salmonella* in macrophages. The results demonstrate a correlation between

Salmonella β-alanine utilization and zinc uptake during intracellular infection and

provide evidence that β-alanine can influence the macrophage immune response by

acting on zinc uptake.

Overall, our findings suggest a model in which *Salmonella* exploits host- and

bacterial-derived β-alanine to efficiently replicate in host macrophages and cause

systemic disease. We propose that *Salmonella* requires a large amount of β-alanine

during intracellular infection. The utilization of β-alanine promotes *Salmonella* uptake

of the essential micronutrient zinc, which was previously shown to be required for the

metabolic needs of intracellular *Salmonella* and to subvert the antimicrobial defense

of macrophages by *Salmonella*. These observations provide new insight into

Salmonella pathogenesis and the crosstalk between *Salmonella* and macrophages

during intracellular infection. Considering that the *panD* gene is present in the

genomes of all *Salmonella* strains and that mutation of *panD* markedly reduced

Salmonella replication ability in macrophages, as well as virulence in the mouse

model, this gene may be used as a potential target to control systemic *Salmonella*

infection.

Methods

Ethics statement

Six-week-old female BALB/c mice were obtained from Beijing Vital River Laboratory Animal Technology (Beijing, China). Mice were housed in barrier facilities under specific pathogen-free conditions with a 12 h light/dark cycle at a temperature of 24 ± 2 °C and a relative humidity of 50 ± 5%. Mice were fed a standard mouse chow diet, and they consumed food and water *ad libitum* throughout the experiment. All animal experiments were conducted in accordance with the policies of the Institutional Animal Care Committee of Nankai University (Tianjin, China) and performed under protocol no. 2021-SYDWLL-000029.

Cell culture

The RAW264.7 mouse macrophage-like cell line (ATCC TIB-71) was obtained from the Shanghai Institute of Biochemistry and Cell Biology of the Chinese Academy of Sciences (Shanghai, China). Cells were cultured in RPMI-1640 medium (Gibco #11879020) supplemented with 10% (v/v) fetal bovine serum (FBS, Gibco #10100147) at 37 □ with 5% CO_2. Cells were seeded in 24-well tissue culture plates at 1×10^5 cells per well 24 h before infection.

Bacterial strains, plasmids, and growth conditions

The bacterial strains and plasmids used in this study are listed in Supplementary Table 1. The *Salmonella enterica* serovar Typhimurium (STM) strain ATCC 14028s was used as the WT strain throughout this study and for the construction of the mutants. Mutant strains were generated using the λ Red recombination system with

the plasmid pSIM17 (Jiang et al., 2017). To construct the complemented strain of

Δ*panD*, the amplified DNA fragments of the *panD* ORF and its upstream promoter

were digested and inserted into the BamHI–EcoRI site of the low-copy-number

plasmid pBR322. To generate the *panD-lux* reporter fusion, the amplification

products of the *panD* promoter region were digested and cloned into the XhoI–BamHI

site of the plasmid pMS402, which carries a promoter-less *lux*CDABE reporter gene

cluster (Liang, Li, Dong, Surette, & Duan, 2008). The sequences of primers used for

the construction of the strains are listed in Supplementary Table 2. All the strains were

verified by PCR amplification and sequencing.

Bacterial strains were conventionally grown overnight in Luria–Bertani (LB)

medium (10 g/L tryptone, 5 g/L yeast extract, and 10 g/L NaCl) or in N-minimal

medium (10 μM $MgCl_2$, 110 μM KH_2PO_4, 7.5 mM $(NH_4)_2SO_4$, 0.5 mM K_2SO_4, 5 mM

KCl, 38 mM glycerol, and 0.1% [w/v] casamino acids) supplemented with appropriate

antibiotics at 37 °C with shaking at 180 rpm or on LB agar plates. All antibiotics were

used at their standard concentrations (chloramphenicol, 25 μg/mL; kanamycin, 50

μg/mL; ampicillin, 100 μg/mL; gentamicin, 10 or 100 μg/mL) unless otherwise

mentioned.

Growth curve

Bacterial strains were conventionally grown overnight in LB medium. The next

day, they were subcultured (1:100) in new LB medium and RPMI-1640 medium or

subcultured in N-minimal medium supplemented with glycerol or β-alanine as the

sole carbon source at 37 °C with shaking at 180 rpm. To measure the growth of

bacteria, 200 μL of the bacterial cultures were transferred to the microplate wells. The

absorbance (OD_{600}) of the bacteria was measured every half hour for 12 h with a

Spark multimode microplate reader (Tecan).

Bioluminescent reporter assays

STM WT carrying the *panD-luxCDABE* fusion plasmid was conventionally

grown overnight in LB medium, and the next day, the cells were subcultured (1:100)

in new LB medium or N-minimal medium for 8 h. The luminescence of the cultured

bacteria (200 μl) was measured with a Spark multimode microplate reader (Tecan).

Moreover, the cultured bacteria (100 μl) were serially diluted and plated on LB agar

plates to estimate bacterial CFUs. Bacterial CFUs were used to normalize

luminescence values.

Salmonella **infection of macrophages**

Bacterial strains were conventionally grown overnight in LB medium to the late

stationary phase, and the next day, the bacteria were diluted to 2×10^6 CFUs/mL and

opsonized in RPMI-1640 medium supplemented with 10% FBS for 15 min. The

macrophage monolayers were infected with the opsonized bacteria culture (0.5

mL/well, multiplicity of infection (MOI) = 10) and centrifuged at $800 \times g$ for 5 min to

synchronize infection. The infected cells were incubated for 30 min at 37 °C in 5%

CO_2 and then washed three times with $1 \times$ PBS. Fresh RPMI-1640 medium containing

100 μg/mL gentamicin was added to the infected cells to kill extracellular bacteria.

After 1 h, fresh RPMI-1640 medium containing 10 μg/mL gentamicin was added to

the infected cells for the remainder of the experiment. To assess the intarcelluar

growth of *Salmonella*, the infected cells were lysed with 1% Triton X-100 at 2 hpi and

20 hpi, and the abundance of the intracellular bacteria CFUs was estimated on LB

agar plates. The relative fold replication of intracellular bacterial strains was denoted

as the CFUs recovered at 20 hpi relative to those at 2 hpi. The relative fold change in

replication was normalized to the number of RAW264.7 cells. When indicated, 1 mM

β-alanine or 100 μM $ZnSO_4$ were added after 1 h of gentamicin treatment.

Targeted metabolomics analysis of amino acids in macrophages

Targeted metabolomics was conducted as previously described (Jiang et al.,

2021), with the combined metabolites of infected cells and intracellular bacteria were

extracted for analysis. The metabolites have been confirmed to be dominated by the

host metabolites (Jiang et al., 2021). RAW264.7 cells were mock-infected or infected

with STM WT for 8 h. The cells were harvested and washed with precooled PBS

solution to remove the medium. Cellular metabolites were extracted using ice-cold

extraction solvent (40:40:20 vol/vol/vol acetonitrile:methanol:water, 0.1 M formic

acid), incubated at -20 □ for 20 min, and then centrifuged for 10 min at 12,000×g and

4 □ to obtain the supernatant. Subsequently, the supernatant was transferred to an LC

–MS vial and analyzed using ultrahigh-performance liquid chromatography (Acquity;

Waters, Milford, MA, USA) coupled with mass spectrometry (Q Exactive Hybrid

Quadrupole-Orbitrap; Thermo Fisher Scientific, Waltham, MA, USA). Metabolites

were separated with a Luna NH_2 column (2 mm × 100 mm, 3 μm particle size;

Phenomenex). Mobile phase A was 20 mM ammonium acetate (pH 9.0), and mobile

phase B was acetonitrile containing 0.1% formic acid. The flow rate was 0.4 ml/min.

Xcalibur 4.0 software (Thermo Fisher) was used for data acquisition and processing.

Metabolite identification was achieved by high-resolution mass and retention time

matching to authentic standards. The absolute quantification of amino acids was

performed using the standard curve method, and the values were normalized to the

cell number. Four biological replicates of each sample were analyzed.

Immunofluorescence staining

RAW264.7 cells were infected with STM WT or Δ*panD* and the complemented

strain c*panD* as described above. After 2 and 20 h of cultivation, the infected cells

were fixed for 15 min in 4% paraformaldehyde, washed in PBS, and permeabilized

with 0.1% Triton X-100 in PBS for 15 min. The fixed samples were blocked in 5%

bovine serum albumin for 30 min, followed by staining with a FITC-conjugated

anti-*Salmonella* antibody (1:100 dilution, Abcam #ab20320) for 1 h at room

temperature in the dark. The nuclei were then stained with DAPI (Invitrogen

#D21490) for 2 min. A confocal laser scanning microscope (Zeiss LSM800) and ZEN

2.3 software (blue edition) were used to acquire and analyze the cell images

(Objective lense: 40×; The number of intracellular bacteria per infected cell was

estimated in random fields by Fiji-ImageJ).

RNA isolation

RNA was extracted from *Salmonella* strains cultured in N-minimal medium or

LB medium. To investigate the expression of the *Salmonella panD* gene inside

macrophages, we obtained RNA from intracellular bacteria in RAW264.7 cells at 8 h

postinfection and from bacteria in RPMI medium. RNA was extracted using an

EASYspinPlus bacterial RNA rapid extraction kit (Aidlab #RN0802) according to the

manufacturer's protocol. RNA quantity and purity were determined using a

NanoDrop 2000 spectrophotometer (NanoDrop Technologies). RNA samples were

stored at −80 °C before use.

Quantitative real-time PCR (qRT□PCR)

According to the manufacturer's protocols, qRT□PCR was performed using 2×

RealStar Power SYBR qPCR Mix (Genstar #A304) in a QuantStudio 5 Real Time

PCR system (Applied Biosystems). cDNA was synthesized using a StarScript III RT

Kit (Genstar #A232). Each sample was subjected to qRT□PCR in triplicate. The

expression level of the 16S *rRNA* gene was used to normalize that of the target genes.

We estimated the expression of each target gene using the $2^{-\Delta\Delta Ct}$ method.

RNA sequencing and analyses

The STM WT and Δ*panD* strains were conventionally grown overnight in LB

medium, subcultured (1:100) in N-minimal medium for 8 h and then collected by

centrifugation for RNA extraction. Sequencing libraries were generated using the

NEBNext Ultra RNA Library Prep Kit for Illumina (New England Biolabs) according

to the manufacturer's instructions, and sequencing was conducted using the Illumina

HiSeq 2000 platform at Shanghai Majorbio Bio-Pharm Technology Co., Ltd.

(Shanghai, China). The sequencing data were deposited in the NCBI Sequence Read

Archive under accession number (SRA, PRJNA1124637). The clean reads were

mapped to the STM ATCC 14028 reference genome (CP001363 and CP001362) by

using the short-sequence alignment software Bowtie 2. Gene expression was

evaluated using the fragments per kilobase of transcript per million mapped reads

(FPKM) method. DEGs in the *panD* mutant relative to the WT were determined using

the R statistical package software EdgeR. The thresholds for statistically significant

differences were set to a fold change ≥ 2 or ≤ 0.5 and a false discovery rate

(FDR) ≤ 0.05. P values were adjusted using the Benjamini–Hochberg procedure for

controlling the FDR. Enrichment analysis of DEGs was conducted using GO and

KEGG enrichment analyses.

Mouse infection

Salmonella strains were conventionally cultured overnight in LB medium, and

the next day, they were subcultured (1:100) in new LB medium and grown at 37 °C

with shaking at 200 rpm to stationary phase (OD_{600}~2). The bacteria were diluted to 5

$\times 10^4$ CFUs/mL in 0.9% NaCl. Groups of BALB/c mice were infected i.p. with 0.1

mL of the NaCl suspension. For survival assays, we recorded and monitored the

mortality and body weight of the infected mice daily for 10 days. To analyze the

bacterial burden of the mouse liver and spleen, we weighed the infected mice first on

day 3 postinfection and then harvested the liver and spleen. The liver and spleen of

infected mice were homogenized in ice-cold PBS, serially diluted, and plated on LB

plates containing the appropriate antibiotics to determine bacterial CFUs. H&E

staining of the mouse liver was performed to investigate the histopathological

alterations in the liver of infected mice. To evaluate the histopathological alterations

in the mouse liver, we harvested the liver of the infected mice on day 5 postinfection.

The mouse liver was washed with 0.9% NaCl, fixed with 10% neutral formalin for 48

h and subsequently processed for routine paraffin embedding. Paraffin-embedded

tissues were sectioned at a thickness of 5 μm and then stained with hematoxylin

(Sangon Biotech #E607318) and eosin (Sangon Biotech #E607318) for

histopathological examination. The stained sections were then examined by light

microscopy (Leica DM2500 LED).

Flow cytometry

RAW264.7 cells were infected with *Salmonella* WT for 8 h, in the absence or

presence of 1 mM β-alanine, which was added to the infected cells at 1 h

post-infection. The infected cells were fixed for 15 min in 4% paraformaldehyde,

washed in PBS, and permeabilized with 0.1% Triton X-100 in PBS for 15 min. The

fixed samples were blocked in 5% bovine serum albumin for 30 min, followed by

staining with an anti-CD86 antibody (Abcam, ab119857), an anti-CD163 antibody

(Abcam, ab182422) for 30 min, and a goat anti-rat IgG H&L (Alexa Fluor® 488)

(Abcam, ab150165), a donkey anti-rabbit IgG H&L (Alexa Fluor® 647) (Abcam,

ab150075) for 30 min in the dark. The infected cells were analyzed using a BD

FACSAria Flow Cytometer (BD Biosciences).

Assessment of intracellular ROS/ RNS levels

RAW264.7 cells were infected with *Salmonella* WT for 8 hours, either in the

absence or presence of 1 mM β-alanine. The β-alanine was added to the infected cells

at 1 h post-infection. Following the infection, the cells were washed three times with

PBS, and the levels of ROS and RNS were measured using a ROS fluorescence probe

(BBoxiProbe® O06, Bestbio, BB-46051) and an RNS fluorescence probe

(BBoxiProbe® O52, Bestbio, BB-470567), respectively, in accordance with the

manufacturer's protocol. The ROS/ RNS levels were normalized to the cell count.

Assessment of zinc levels in mouse liver and RAW 264.7 macrophages

To assess the zinc level in mouse liver, mice were i.p. infected with

approximately 5000 CFUs of *Salmonella* WT or Δ*panD* for 3 days, with three mice in

each injection group. Following infection, mouse livers were collected and then

dissociated into single cells using the gentleMACS/Mouse Liver Dissociation Kit

(Miltenyi Biotec, 130-105-807) in accordance with the manufacturer's protocol. The

zinc concentration in mouse liver was determined using a zinc fluorescence probe

(Zinquin ethyl ester, MKBio, MX4516) following the manufacturer's instructions and

normalized by the cell count.

To assess the zinc level in RAW 264.7 macrophages, RAW 264.7 cells were

infected with *Salmonella* WT or Δ*panD* for 8 hours. The infected cells were washed

three times with PBS, and zinc concentration was measured using a zinc fluorescence

probe (Zinquin ethyl ester, MKBio, MX4516) following the manufacturer's protocol.

The zinc concentration was normalized by the cell count.

Statistical analysis

The data are presented as the mean ± SD. All in vitro experiments were

conducted in duplicate and repeated at least three times (n ≥ 3). Mouse assays were

performed twice, with at least two mouse (n ≥ 2) in each injection group, and the

combined data from the two experiments was used for statistical analysis. Statistical

analyses were performed using GraphPad InStat software (version GraphPad Prism

9.5.1, San Diego, CA, USA) with two-sided Student's *t* tests, one-way ANOVA,

two-way ANOVA, log-rank Mantel–Cox tests, or Mann☐Whitney U tests according

to the test requirements (as stated in the figure legends). A *P* value < 0.05 indicated a

statistically significant difference. ns represents no statistical significance.

Data availability statement

The RNA-seq data generated in this study have been deposited in the NCBI

Sequence Read Archive (SRA) database under the accession number PRJNA1124637.

All other data associated with this study are available in the main text and

supplementary materials. Source data are provided in Supplementary Data 1.

Acknowledgements

This work was supported by the National Natural Science Foundation of China

(NSFC) Program (grant no. 32170110, 32270191), the Natural Science Foundation of

Tianjin (grant no. 22JCYBJC01060), and the Fundamental Research Funds for the

Central Universities, Nankai University (grant no. 63241588, 63243161).

Author contributions

LJ designed the research; SM, YS, XW, HG, RL, TY, CK, and JC performed the

research; BY provided technical support and insights; SM and BY analyzed the data;

and LJ and SM wrote the manuscript.

Disclosure statement

The authors report there are no competing interests to declare.

ORCID

Lingyan Jiang http://orcid.org/0000-0001-8580-606X

References

Ammendola, S., Pasquali, P., Pistoia, C., Petrucci, P., Petrarca, P., Rotilio, G., & Battistoni, A. (2007). High-affinity Zn2+ uptake system ZnuABC is required for bacterial zinc homeostasis in intracellular environments and contributes to the virulence of *Salmonella enterica*. *Infect Immun, 75*(12), 5867-5876. doi:10.1128/iai.00559-07

Andreini, C., Banci, L., Bertini, I., & Rosato, A. (2006). Zinc through the three domains of life. *J Proteome Res, 5*(11), 3173-3178. doi:10.1021/pr0603699

Basavanna, S., Chimalapati, S., Maqbool, A., Rubbo, B., Yuste, J., Wilson, R. J., . . . Brown, J. S. (2013). The effects of methionine acquisition and synthesis on *Streptococcus pneumoniae* growth and virulence. *PLoS One, 8*(1), e49638. doi:10.1371/journal.pone.0049638

Battistoni, A., Ammendola, S., Chiancone, E., & Ilari, A. (2017). A novel antimicrobial approach based on the inhibition of zinc uptake in *Salmonella enterica*. *Future Med Chem, 9*(9), 899-910. doi:10.4155/fmc-2017-0042

Begley, T. P., Kinsland, C., & Strauss, E. (2001). The biosynthesis of coenzyme A in bacteria. *Vitam Horm, 61*, 157-171. doi:10.1016/s0083-6729(01)61005-7

Bellinger, P. M. (2014). β-Alanine supplementation for athletic performance: an update. *J Strength Cond Res, 28*(6), 1751-1770. doi:10.1519/jsc.0000000000000327

Bergkessel, M. (2020). Regulation of protein biosynthetic activity during growth

arrest. *Curr Opin Microbiol, 57*, 62-69. doi:10.1016/j.mib.2020.07.010

Bock, T., Müller, R., & Blankenfeldt, W. (2016). Crystal structure of AibC, a

reductase involved in alternative de novo isovaleryl coenzyme A biosynthesis in

Myxococcus xanthus. *Acta Crystallogr F Struct Biol Commun, 72*(Pt 8), 652-658.

doi:10.1107/s2053230x16011146

Boldyrev, A. A., Aldini, G., & Derave, W. (2013). Physiology and pathophysiology of

carnosine. *Physiol Rev, 93*(4), 1803-1845. doi:10.1152/physrev.00039.2012

Bomjan, R., Zhang, M., & Zhou, D. (2019). YshB promotes intracellular replication

and is required for *salmonella* virulence. *J Bacteriol, 201*(17), e00314-19.

doi:10.1128/jb.00314-19

Bosdriesz, E., Molenaar, D., Teusink, B., & Bruggeman, F. J. (2015). How

fast-growing bacteria robustly tune their ribosome concentration to approximate

growth-rate maximization. *Febs j, 282*(10), 2029-2044. doi:10.1111/febs.13258

Bowden, S. D., Rowley, G., Hinton, J. C., & Thompson, A. (2009). Glucose and

glycolysis are required for the successful infection of macrophages and mice by

Salmonella enterica serovar Typhimurium. *Infect Immun, 77*(7), 3117-3126.

doi:10.1128/iai.00093-09

Brumell, J. H., Tang, P., Mills, S. D., & Finlay, B. B. (2001). Characterization of

Salmonella-induced filaments (Sifs) reveals a delayed interaction between

Salmonella-containing vacuoles and late endocytic compartments. *Traffic, 2*(9),

643-653. doi:10.1034/j.1600-0854.2001.20907.x

Bumann, D., & Schothorst, J. (2017). Intracellular *Salmonella* metabolism. *Cell*

Microbiol, 19(10), 10.1111/cmi.12766. doi:10.1111/cmi.12766

Castanheira, S., & García-Del Portillo, F. (2017). *Salmonella* populations inside host

cells. *Front Cell Infect Microbiol, 7*, 432. doi:10.3389/fcimb.2017.00432

Dandekar, T., Fieselmann, A., Fischer, E., Popp, J., Hensel, M., & Noster, J. (2014).

Salmonella-how a metabolic generalist adopts an intracellular lifestyle during

infection. *Front Cell Infect Microbiol, 4*, 191. doi:10.3389/fcimb.2014.00191

Das, P., Lahiri, A., Lahiri, A., Sen, M., Iyer, N., Kapoor, N., . . . Chakravortty, D.

(2010). Cationic amino acid transporters and Salmonella Typhimurium ArgT

collectively regulate arginine availability towards intracellular Salmonella

growth. *PLoS One, 5*(12), e15466. doi:10.1371/journal.pone.0015466

Davaapil, H., Tsuchiya, Y., & Gout, I. (2014). Signalling functions of coenzyme A and

its derivatives in mammalian cells. *Biochem Soc Trans, 42*(4), 1056-1062.

doi:10.1042/bst20140146

de Jong, H. K., Parry, C. M., van der Poll, T., & Wiersinga, W. J. (2012).

Host-pathogen interaction in invasive Salmonellosis. *PLoS Pathog, 8*(10),

e1002933. doi:10.1371/journal.ppat.1002933

De la Cruz, M. A., Fernández-Mora, M., Guadarrama, C., Flores-Valdez, M. A.,

Bustamante, V. H., Vázquez, A., & Calva, E. (2007). LeuO antagonizes H-NS

and StpA-dependent repression in *Salmonella enterica* ompS1. *Mol Microbiol,*

66(3), 727-743. doi:10.1111/j.1365-2958.2007.05958.x

de Paiva, J. B., da Silva, L. P., Casas, M. R., Conceição, R. A., Nakazato, G., de Pace,

F., . . . da Silveira, W. D. (2016). *In vivo* influence of in vitro up-regulated genes

in the virulence of an APEC strain associated with swollen head syndrome. *Avian*

Pathol, 45(1), 94-105. doi:10.1080/03079457.2015.1125995

Deng, W., Marshall, N. C., Rowland, J. L., McCoy, J. M., Worrall, L. J., Santos, A.

S., . . . Finlay, B. B. (2017). Assembly, structure, function and regulation of type

III secretion systems. *Nat Rev Microbiol, 15*(6), 323-337.

doi:10.1038/nrmicro.2017.20

Dillon, S. C., Espinosa, E., Hokamp, K., Ussery, D. W., Casadesús, J., & Dorman, C. J.

(2012). LeuO is a global regulator of gene expression in *Salmonella enterica*

serovar Typhimurium. *Mol Microbiol, 85*(6), 1072-1089.

doi:10.1111/j.1365-2958.2012.08162.x

Eisenreich, W., Dandekar, T., Heesemann, J., & Goebel, W. (2010). Carbon

metabolism of intracellular bacterial pathogens and possible links to virulence.

Nat Rev Microbiol, 8(6), 401-412. doi:10.1038/nrmicro2351

Fàbrega, A., & Vila, J. (2013). *Salmonella enterica* serovar Typhimurium skills to

succeed in the host: virulence and regulation. *Clin Microbiol Rev, 26*(2), 308-341.

doi:10.1128/cmr.00066-12

Feldman, M., Harbeck, M., Keller, M., Spyrou, M. A., Rott, A., Trautmann, B., . . .

Krause, J. (2016). A High-Coverage *Yersinia pestis* Genome from a

Sixth-Century Justinianic Plague Victim. *Mol Biol Evol, 33*(11), 2911-2923.

doi:10.1093/molbev/msw170

Fernández-Mora, M., Puente, J. L., & Calva, E. (2004). OmpR and LeuO positively

regulate the *Salmonella enterica* serovar Typhi ompS2 porin gene. *J Bacteriol,*

186(10), 2909-2920. doi:10.1128/jb.186.10.2909-2920.2004

Fields, P. I., Swanson, R. V., Haidaris, C. G., & Heffron, F. (1986). Mutants of *Salmonella* Typhimurium that cannot survive within the macrophage are avirulent. *Proc Natl Acad Sci U S A, 83*(14), 5189-5193. doi:10.1073/pnas.83.14.5189

Figueira, R., & Holden, D. W. (2012). Functions of the *Salmonella* pathogenicity island 2 (SPI-2) type III secretion system effectors. *Microbiology (Reading), 158*(Pt 5), 1147-1161. doi:10.1099/mic.0.058115-0

Gojković, Z., Sandrini, M. P., & Piskur, J. (2001). Eukaryotic beta-alanine synthases are functionally related but have a high degree of structural diversity. *Genetics, 158*(3), 999-1011. doi:10.1093/genetics/158.3.999

Gout, I. (2019). Coenzyme A: a protective thiol in bacterial antioxidant defence. *Biochem Soc Trans, 47*(1), 469-476. doi:10.1042/bst20180415

Guadarrama, C., Villaseñor, T., & Calva, E. (2014). The subtleties and contrasts of the LeuO regulator in *Salmonella* Typhi: implications in the immune response. *Front Immunol, 5*, 581. doi:10.3389/fimmu.2014.00581

Han, J., Aljahdali, N., Zhao, S., Tang, H., Harbottle, H., Hoffmann, M., . . . Foley, S. L. (2024). Infection biology of Salmonella enterica. *EcoSal Plus, 12*(1), eesp00012023. doi:10.1128/ecosalplus.esp-0001-2023

Hernández-Lucas, I., Gallego-Hernández, A. L., Encarnación, S., Fernández-Mora, M., Martínez-Batallar, A. G., Salgado, H., . . . Calva, E. (2008). The LysR-type transcriptional regulator LeuO controls expression of several genes in

Salmonella enterica serovar Typhi. *J Bacteriol, 190*(5), 1658-1670. doi:10.1128/jb.01649-07

Herrero-Fresno, A., & Olsen, J. E. (2018). *Salmonella* Typhimurium metabolism affects virulence in the host - A mini-review. *Food Microbiol, 71*, 98-110. doi:10.1016/j.fm.2017.04.016

Hoffman, J. R., Varanoske, A., & Stout, J. R. (2018). Effects of β-Alanine supplementation on carnosine elevation and physiological performance. *Adv Food Nutr Res, 84*, 183-206. doi:10.1016/bs.afnr.2017.12.003

Huerta Ojeda, Á., Tapia Cerda, C., Poblete Salvatierra, M. F., Barahona-Fuentes, G., & Jorquera Aguilera, C. (2020). Effects of beta-Alanine supplementation on physical performance in aerobic-anaerobic transition zones: a systematic review and meta-analysis. *Nutrients, 12*(9), 2490. doi:10.3390/nu12092490

Ilari, A., Pescatori, L., Di Santo, R., Battistoni, A., Ammendola, S., Falconi, M., . . . Chiancone, E. (2016). *Salmonella enterica* serovar Typhimurium growth is inhibited by the concomitant binding of Zn(II) and a pyrrolyl-hydroxamate to ZnuA, the soluble component of the ZnuABC transporter. *Biochim Biophys Acta, 1860*(3), 534-541. doi:10.1016/j.bbagen.2015.12.006

Jennings, E., Esposito, D., Rittinger, K., & Thurston, T. L. M. (2018). Structure-function analyses of the bacterial zinc metalloprotease effector protein GtgA uncover key residues required for deactivating NF-κB. *J Biol Chem, 293*(39), 15316-15329. doi:10.1074/jbc.RA118.004255

Jiang, L., Feng, L., Yang, B., Zhang, W., Wang, P., Jiang, X., & Wang, L. (2017).

Signal transduction pathway mediated by the novel regulator LoiA for low oxygen tension induced Salmonella Typhimurium invasion. *PLoS Pathog, 13*(6), e1006429. doi:10.1371/journal.ppat.1006429

Jiang, L., Wang, P., Song, X., Zhang, H., Ma, S., Wang, J., . . . Wang, L. (2021). *Salmonella* Typhimurium reprograms macrophage metabolism via T3SS effector SopE2 to promote intracellular replication and virulence. *Nat Commun, 12*(1), 879. doi:10.1038/s41467-021-21186-4

Kehl, A., Noster, J., & Hensel, M. (2020). Eat in or Take out? Metabolism of Intracellular Salmonella enterica. *Trends Microbiol, 28*(8), 644-654. doi:10.1016/j.tim.2020.03.005

Kim, S., Lee, K. S., Pak, G. D., Excler, J. L., Sahastrabuddhe, S., Marks, F., . . . Mogasale, V. (2019). Spatial and temporal patterns of typhoid and paratyphoid fever outbreaks: a worldwide review, 1990-2018. *Clin Infect Dis, 69*(Suppl 6), S499-s509. doi:10.1093/cid/ciz705

LaRock, D. L., Chaudhary, A., & Miller, S. I. (2015). Salmonellae interactions with host processes. *Nat Rev Microbiol, 13*(4), 191-205. doi:10.1038/nrmicro3420

Lathrop, S. K., Binder, K. A., Starr, T., Cooper, K. G., Chong, A., Carmody, A. B., & Steele-Mortimer, O. (2015). Replication of *Salmonella enterica* serovar Typhimurium in human monocyte-derived macrophages. *Infect Immun, 83*(7), 2661-2671. doi:10.1128/iai.00033-15

Lauriano, C. M., Barker, J. R., Yoon, S. S., Nano, F. E., Arulanandam, B. P., Hassett, D. J., & Klose, K. E. (2004). MglA regulates transcription of virulence factors

necessary for *Francisella tularensis* intraamoebae and intramacrophage survival.

Proc Natl Acad Sci U S A, 101(12), 4246-4249. doi:10.1073/pnas.0307690101

Leung, K. Y., & Finlay, B. B. (1991). Intracellular replication is essential for the

virulence of *Salmonella* Typhimurium. *Proc Natl Acad Sci U S A, 88*(24),

11470-11474. doi:10.1073/pnas.88.24.11470

Li, W., Ren, Q., Ni, T., Zhao, Y., Sang, Z., Luo, R., . . . Li, S. (2023). Strategies

adopted by *Salmonella* to survive in host: a review. *Arch Microbiol, 205*(12), 362.

doi:10.1007/s00203-023-03702-w

Liang, H., Li, L., Dong, Z., Surette, M. G., & Duan, K. (2008). The YebC family

protein PA0964 negatively regulates the Pseudomonas aeruginosa quinolone

signal system and pyocyanin production. *J Bacteriol, 190*(18), 6217-6227.

doi:10.1128/jb.00428-08

Liss, V., Swart, A. L., Kehl, A., Hermanns, N., Zhang, Y., Chikkaballi, D., . . . Hensel,

M. (2017). Salmonella enterica Remodels the Host Cell Endosomal System for

Efficient Intravacuolar Nutrition. *Cell Host Microbe, 21*(3), 390-402.

doi:10.1016/j.chom.2017.02.005

Liu, Y., Ho, K. K., Su, J., Gong, H., Chang, A. C., & Lu, S. (2013). Potassium

transport of *Salmonella* is important for type III secretion and pathogenesis.

Microbiology (Reading), 159(Pt 8), 1705-1719. doi:10.1099/mic.0.068700-0

Lynch, J. P., & Lesser, C. F. (2021). A host metabolite promotes *Salmonella* survival.

Science, 371(6527), 344-345. doi:10.1126/science.abf8414

Majowicz, S. E., Musto, J., Scallan, E., Angulo, F. J., Kirk, M., O'Brien, S. J., . . .

Hoekstra, R. M. (2010). The global burden of nontyphoidal *Salmonella*

gastroenteritis. *Clin Infect Dis, 50*(6), 882-889. doi:10.1086/650733

Martínez-Guitián, M., Vázquez-Ucha, J. C., Álvarez-Fraga, L., Conde-Pérez, K.,

Lasarte-Monterrubio, C., Vallejo, J. A., . . . Beceiro, A. (2019). Involvement of

HisF in the persistence of *Acinetobacter baumannii* during a pneumonia

infection. *Front Cell Infect Microbiol, 9*, 310. doi:10.3389/fcimb.2019.00310

Michelucci, A., Cordes, T., Ghelfi, J., Pailot, A., Reiling, N., Goldmann, O., . . . Hiller,

K. (2013). Immune-responsive gene 1 protein links metabolism to immunity by

catalyzing itaconic acid production. *Proc Natl Acad Sci U S A, 110*(19),

7820-7825. doi:10.1073/pnas.1218599110

Nozaki, S., Webb, M. E., & Niki, H. (2012). An activator for pyruvoyl-dependent

l-aspartate α-decarboxylase is conserved in a small group of the γ-proteobacteria

including Escherichia coli. *Microbiologyopen, 1*(3), 298-310.

doi:10.1002/mbo3.34

Ohl, M. E., & Miller, S. I. (2001). *Salmonella*: a model for bacterial pathogenesis.

Annu Rev Med, 52, 259-274. doi:10.1146/annurev.med.52.1.259

Pandeya, A., Zhang, Y., Cui, J., Yang, L., Li, J., Zhang, G., . . . Wei, Y. (2023).

Inflammasome activation and pyroptosis mediate coagulopathy and

inflammation in Salmonella systemic infection. *Microbiol Res, 275*, 127460.

doi:10.1016/j.micres.2023.127460

Pavić, A., Ji, Y., Serafini, A., Garza-Garcia, A., McPhillie, M. J., Holmes, A. O. M., . . .

Postis, V. L. G. (2021). Functional Characterization of the γ-Aminobutyric Acid

Transporter from Mycobacterium smegmatis MC(2) 155 Reveals Sodium-Driven

GABA Transport. *J Bacteriol, 203*(4), e00642-20. doi:10.1128/jb.00642-20

Peace, C. G., & O'Neill, L. A. (2022). The role of itaconate in host defense and

inflammation. *J Clin Invest, 132*(2), e148548. doi:10.1172/jci148548

Pillay, T. D., Hettiarachchi, S. U., Gan, J., Diaz-Del-Olmo, I., Yu, X. J., Muench, J.

H., . . . Pearson, J. S. (2023). Speaking the host language: how *Salmonella*

effector proteins manipulate the host. *Microbiology (Reading), 169*(6), 001342.

doi:10.1099/mic.0.001342

Popp, J., Noster, J., Busch, K., Kehl, A., Zur Hellen, G., & Hensel, M. (2015). Role of

host cell-derived amino acids in nutrition of intracellular *Salmonella enterica.*

Infect Immun, 83(12), 4466-4475. doi:10.1128/iai.00624-15

Rajashekar, R., Liebl, D., Seitz, A., & Hensel, M. (2008). Dynamic remodeling of the

endosomal system during formation of Salmonella-induced filaments by

intracellular Salmonella enterica. *Traffic, 9*(12), 2100-2116.

doi:10.1111/j.1600-0854.2008.00821.x

Röder, J., Felgner, P., & Hensel, M. (2021). Comprehensive Single Cell Analyses of

the Nutritional Environment of Intracellular Salmonella enterica. *Front Cell*

Infect Microbiol, 11, 624650. doi:10.3389/fcimb.2021.624650

Rodríguez-Morales, O., Fernández-Mora, M., Hernández-Lucas, I., Vázquez, A.,

Puente, J. L., & Calva, E. (2006). *Salmonella enterica* serovar Typhimurium

ompS1 and ompS2 mutants are attenuated for virulence in mice. *Infect Immun,*

74(2), 1398-1402. doi:10.1128/iai.74.2.1398-1402.2006

Rosenberg, G., Riquelme, S., Prince, A., & Avraham, R. (2022). Immunometabolic

crosstalk during bacterial infection. *Nat Microbiol, 7*(4), 497-507.

doi:10.1038/s41564-022-01080-5

Rosenberg, G., Yehezkel, D., Hoffman, D., Mattioli, C. C., Fremder, M., Ben-Arosh,

H., . . . Avraham, R. (2021). Host succinate is an activation signal for *Salmonella*

virulence during intracellular infection. *Science, 371*(6527), 400-405.

doi:10.1126/science.aba8026

Schmitzberger, F., Smith, A. G., Abell, C., & Blundell, T. L. (2003). Comparative

analysis of the *Escherichia coli* ketopantoate hydroxymethyltransferase crystal

structure confirms that it is a member of the (betaalpha)8

phosphoenolpyruvate/pyruvate superfamily. *J Bacteriol, 185*(14), 4163-4171.

doi:10.1128/jb.185.14.4163-4171.2003

Schneider, F., Krämer, R., & Burkovski, A. (2004). Identification and characterization

of the main beta-alanine uptake system in *Escherichia coli*. *Appl Microbiol*

Biotechnol, 65(5), 576-582. doi:10.1007/s00253-004-1636-0

Sibon, O. C., & Strauss, E. (2016). Coenzyme A: to make it or uptake it? *Nat Rev Mol*

Cell Biol, 17(10), 605-606. doi:10.1038/nrm.2016.110

Silva, A. F., Matos, M. P., Ralph, M. T., Silva, D. L., de Alencar, N. M., Ramos, M. V.,

& Lima-Filho, J. V. (2016). Comparison of immunomodulatory properties of

mannose-binding lectins from Canavalia brasiliensis and Cratylia argentea in a

mice model of Salmonella infection. *Int Immunopharmacol, 31*, 233-238.

doi:10.1016/j.intimp.2015.12.036

Song, P., Zhang, X., Wang, S., Xu, W., & Wei, F. (2023). Advances in the synthesis of

β-alanine. *Front   Bioeng   Biotechnol,   11*,   1283129.

doi:10.3389/fbioe.2023.1283129

Steeb, B., Claudi, B., Burton, N. A., Tienz, P., Schmidt, A., Farhan, H., . . . Bumann,

D. (2013). Parallel exploitation of diverse host nutrients enhances *Salmonella*

virulence. *PLoS Pathog, 9*(4), e1003301. doi:10.1371/journal.ppat.1003301

Tan, X., Bramlett, M. R., & Lindahl, P. A. (2004). Effect of Zn on acetyl coenzyme a

synthase: evidence for a conformational change in the alpha subunit during

catalysis. *J Am Chem Soc, 126*(19), 5954-5955. doi:10.1021/ja039600z

Theodoulou, F. L., Sibon, O. C., Jackowski, S., & Gout, I. (2014). Coenzyme A and

its derivatives: renaissance of a textbook classic. *Biochem Soc Trans, 42*(4),

1025-1032. doi:10.1042/bst20140176

Thompson, A., Fulde, M., & Tedin, K. (2018). The metabolic pathways utilized by

Salmonella Typhimurium during infection of host cells. *Environ Microbiol Rep,*

10(2), 140-154. doi:10.1111/1758-2229.12628

Tuli, A., & Sharma, M. (2019). How to do business with lysosomes: *Salmonella* leads

the way. *Curr Opin Microbiol, 47*, 1-7. doi:10.1016/j.mib.2018.10.003

Wang, J., Ma, S., Li, W., Wang, X., Huang, D., Jiang, L., & Feng, L. (2021).

Salmonella enterica serovar Typhi induces host metabolic reprogramming to

increase glucose availability for intracellular replication. *Int J Mol Sci, 22*(18),

10003. doi:10.3390/ijms221810003

Wang, L., Mao, Y., Wang, Z., Ma, H., & Chen, T. (2021). Advances in

biotechnological production of β-alanine. *World J Microbiol Biotechnol, 37*(5), 79. doi:10.1007/s11274-021-03042-1

Wang, X., Yang, B., Ma, S., Yan, X., Ma, S., Sun, H., . . . Jiang, L. (2023). Lactate promotes *Salmonella* intracellular replication and systemic infection via driving macrophage M2 polarization. *Microbiol Spectr, 11*(6), e0225323. doi:10.1128/spectrum.02253-23

Webb, M. E., Smith, A. G., & Abell, C. (2004). Biosynthesis of pantothenate. *Nat Prod Rep, 21*(6), 695-721. doi:10.1039/b316419p

West, T. P., Traut, T. W., Shanley, M. S., & O'Donovan, G. A. (1985). A Salmonella typhimurium strain defective in uracil catabolism and beta-alanine synthesis. *J Gen Microbiol, 131*(5), 1083-1090. doi:10.1099/00221287-131-5-1083

White, W. H., Gunyuzlu, P. L., & Toyn, J. H. (2001). *Saccharomyces cerevisiae* is capable of de *novo* pantothenic acid biosynthesis involving a novel pathway of beta-alanine production from spermine. *J Biol Chem, 276*(14), 10794-10800. doi:10.1074/jbc.M009804200

Wu, A., Tymoszuk, P., Haschka, D., Heeke, S., Dichtl, S., Petzer, V., . . . Weiss, G. (2017). *Salmonella* utilizes zinc to subvert antimicrobial host defense of macrophages via modulation of NF-κB signaling. *Infect Immun, 85*(12), e00418-17. doi:10.1128/iai.00418-17

Yang, W., & Cong, Y. (2021). Gut microbiota-derived metabolites in the regulation of host immune responses and immune-related inflammatory diseases. *Cell Mol Immunol, 18*(4), 866-877. doi:10.1038/s41423-021-00661-4

1001   Yuan, S. F., Nair, P. H., Borbon, D., Coleman, S. M., Fan, P. H., Lin, W. L., & Alper,

1002   H. S. (2022). Metabolic engineering of E. coli for β-alanine production using a

1003   multi-biosensor enabled approach. *Metab Eng, 74*, 24-35.

1004   doi:10.1016/j.ymben.2022.08.012

1005

Figure legends

Figure 1. Host-derived β-alanine promotes *Salmonella* replication inside macrophages. (A) Schematic workflow for targeted metabolomics investigation of mock- and *Salmonella*-infected (STM) mouse RAW264.7 macrophages. Picture materials were used from bioicons (https://bioicons.com/). (B) Principal component analysis (PCA) score plots of metabolic profiles in the mock- and *Salmonella*-infected (STM) groups (n = 4 biologically independent samples). (C) The concentrations of upregulated amino acids in the mock- and *Salmonella*-infected groups (n = 4 biologically independent samples). (D) The concentrations of downregulated amino acids in the mock- and *Salmonella*-infected groups (n = 4 biologically independent samples). (E) Fold intracellular replication (20 h *vs.* 2 h) of *Salmonella* WT in RAW264.7 cells in the presence of 0.5, 1, 2, 4 mM β-alanine. Data are presented as the mean ± SD, n = 3 independent experiments. (F) Growth curves of *Salmonella* WT and the *argT* mutant (Δ*argT*) in N-minimal (left) and M9 minimal (right) medium supplemented with β-alanine (1 mM) as the sole carbon source. Data are presented as mean ± SD, n = 4 independent experiments. Statistical significance was assessed using two-sided Student's *t*-test (C, D) and one-way ANOVA (E).

Figure 2. *De novo* β-alanine synthesis is critical for *Salmonella* replication inside macrophages. (A) Scheme of β-alanine and the downstream CoA biosynthesis pathway in *Salmonella*. (B) qRT□PCR analysis of the expression of the *Salmonella* *panD* gene in RAW264.7 cells (8 h postinfection) and RPMI-1640 medium. (C)

qRT☐PCR analysis of the expression of the *Salmonella panD* gene in N-minimal

medium and LB medium. (D) Expression of the *panD*-lux transcriptional fusion in

N-minimal medium and LB medium. Luminescence values were normalized to 10^5

bacterial CFUs. (E) Fold intracellular replication (20 h *vs.* 2 h) of *Salmonella*

Typhimurium 14028s wild type (WT), the *panD* mutant (Δ*panD*) and the

complemented strain (c*panD*) in RAW264.7 cells. (F) Number of intracellular

Salmonella WT, Δ*panD*, and c*panD* strains per RAW264.7 cell at 2 and 20 h

postinfection. The number of intracellular bacteria per infected cell was estimated in

random fields, n = 80 cells per group from 3 independent experiments. (G)

Representative immunofluorescence images of *Salmonella* WT, Δ*panD*, and c*panD* in

RAW264.7 cells at 20 h postinfection (green, *Salmonella*; blue, nuclei; scale bars, 50

µm). Images are representative of three independent experiments. (H) Replication of

Salmonella WT and Δ*panD* in RAW264.7 cells in the presence or absence of 1 mM

β-alanine. The data are presented as the mean ± SD, n = 3 (B–E, H) independent

experiments. Statistical significance was assessed using two-sided Student's *t*-test

(B-D) or one-way ANOVA (E, F, H). ns, not Significant.

1044

Figure 3. *De novo* β-alanine synthesis is critical for systemic *Salmonella* infection

in mice. (A) Schematic illustration of the mouse infection assays. Picture materials

were used from bioicons (https://bioicons.com/). (B, C) Survival curves (B) and body

weight dynamics (C) of mice infected i.p. with *Salmonella* WT, Δ*panD*, or c*panD*. n =

10 mice per group. (D) Liver and spleen bacterial burdens and body weights of mice

infected with *Salmonella* WT, Δ*panD*, or c*panD* on day 3 postinfection. n = 7 mice

per group. (E) Representative immunofluorescence images and intracellular bacterial

counts of *Salmonella* WT, Δ*panD*, and c*panD* in mouse liver at 5 days post-infection

(green, *Salmonella*; blue, nuclei; scale bars, 50 µm). Images are representative of

three independent experiments. The number of intracellular bacteria per infected cell

was estimated in random fields, with n = 80 cells per group from 3 independent

experiments. (F) Representative H&E-stained liver sections from mice that were left

uninfected or infected with *Salmonella* WT, Δ*panD*, or c*panD* on day 5 postinfection.

Arrows indicate severe inflammatory cell infiltration in the mouse liver. Images are

representative of three independent experiments. The data are presented as the mean ±

SD (B-E). Statistical significance was assessed using the log-rank Mantel–Cox test

(B), two-sided Student's *t*-test (C), or one-way ANOVA (D, E).

1062

Figure 4. β-Alanine is involved in the regulation of several metabolic pathways in

Salmonella. (A) Principal component analysis (PCA) score plots of transcriptomic

profiles of *Salmonella* WT and Δ*panD* (n = 3 biologically independent samples). (B)

Volcano plot of the differentially expressed genes (DEGs) in *Salmonella* WT versus

Δ*panD*. The upper right section (red dots) indicates the upregulated DEGs, and the

upper left section (green dots) indicates the downregulated DEGs. (C) Gene Ontology

(GO) enrichment analysis of DEGs. Bubble chart showing the top 20 enriched GO

terms. (D) Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment

analysis of DEGs. (E) Expression of the downregulated pathways (activated by PanD)

is shown in the Z score-transformed heatmap, with red representing higher abundance

and blue representing lower abundance. (F) qRT☐PCR analysis of the mRNA levels

of 16 selected downregulated DEGs in *Salmonella* WT, Δ*panD*, and c*panD*. The data

are presented as the mean ± SD, n = 3 independent experiments. Statistical

significance was assessed using two-way ANOVA.

1077

Figure 5. β-Alanine promotes *Salmonella* virulence *in vivo* partially by increasing

the expression of zinc transporter genes. (A, B, C) Liver (A) and spleen (B)

bacterial burdens and body weight (C) of mice infected with *Salmonella* WT, Δ*fadAB*,

Δ*metR*, Δ*hisABCDFGHL*, Δ*kdpABC*, Δ*mglABC*, Δ*potFGHI*, or Δ*leuO* on day 3

postinfection. n = 5 mice per group. (D) Liver and spleen bacterial burdens and body

weights of mice infected with *Salmonella* WT, Δ*panD,* Δ*znuA* or Δ*panD*Δ*znuA* on day

3 postinfection. n = 5 mice per group. (E) The zinc levels in the livers of mice

infection with either *Salmonella* WT or Δ*panD* for 3 days, n = 5 mice per group. The

data are presented as the mean ± SD (A–E). Statistical significance was assessed

using one-way ANOVA (A–D), two-sided Student's *t*-test (E). ns, not Significant.

1088

Figure 6. β-Alanine promotes *Salmonella* replication within macrophages

partially **by increasing the expression of zinc transporter genes.** (A) The zinc levels

in RAW264.7 cells after infection with *Salmonella* WT or Δ*panD* for 8 h. (B)

Replication of *Salmonella* WT, Δ*panD,* Δ*znuA* and Δ*panD*Δ*znuA* in RAW264.7 cells.

(C) Replication of *Salmonella* WT and Δ*panD* in RAW264.7 cells in the presence or

absence of 100 μM ZnSO$_4$. (D) Replication of *Salmonella* WT and Δ*znuA* in

RAW264.7 cells in the presence or absence of 1 mM β-alanine. The data are presented

as the mean ± SD, n = 3 independent experiments (A–D). Statistical significance was

assessed using two-sided Student's *t*-test (A), one-way ANOVA (B–D). ns, not

Significant. (E) Schematic model of β-alanine-mediated *Salmonella* replication inside

macrophages. In macrophages, *Salmonella* acquires β-alanine both via the uptake of

β-alanine from host macrophages and the *de novo* synthesis of β-alanine. β-alanine

promotes the expression of zinc transporter genes *ZnuABC,* which facilitate the uptake

of zinc by intracellular *Salmonella,* therefore promote *Salmonella* replication in

macrophages and subsequent systemic infection.

1104

1105 **Figure 1_figure Supplement 1.** The levels of ROS and RNS in RAW264.7 cells after

1106 infection with *Salmonella* WT for 8 hours, in the presence or absence of 1 mM

1107 β-alanine. Data are presented as mean ± SD, n = 3 independent experiments.

1108 Statistical significance was assessed using two-sided Student's *t*-test. ns, not

1109 Significant.

1110

Figure 1_figure Supplement 2. Flow cytometry analysis was conducted to determine

the percentage of pro-inflammatory M1 macrophages (CD86+) and anti-inflammatory

M2 macrophages (CD163+). RAW264.7 cells were infected with *Salmonella* WT for

8 hours, in the presence or absence of 1 mM β-alanine. Subsequently, the infected

cells were collected for flow cytometry analysis. Representative dot plots and

quantification of M1 (CD86+) and M2 (CD163+) macrophages are displayed in the

left and right panels, respectively. Data are presented as mean ± SD, n = 3

independent experiments. Statistical significance was assessed using two-way

ANOVA. ns, not Significant.

1120

1121 **Figure 1_figure Supplement 3.** Growth curves of *Salmonella* WT and the *cycA*

1122 mutant (Δ*cycA*) in N-minimal medium supplemented with β-alanine (1 mM) as the

1123 sole carbon source. Data are presented as mean ± SD, n = 3 independent experiments.

1124

1125 **Figure 1_figure Supplement 4.** Growth curves of *Salmonella* WT and the *cycA*

1126 mutant (Δ*cycA*) in M9 minimal medium supplemented with β-alanine (1 mM) as the

1127 sole carbon source. Data are presented as mean ± SD, n = 3 independent experiments.

1128

1129 **Figure 1_figure Supplement 5.** Fold intracellular replication (20 h *vs.* 2 h) of

1130 *Salmonella* wild-type and Δ*cycA* in RAW264.7 cells. Data are presented as mean ±

1131 SD, n = 4 independent experiments. Statistical significance was assessed using

1132 two-way ANOVA. ns, not Significant.

1133

1134 **Figure 1_figure Supplement 6.** Liver and spleen bacterial burdens, and body weight

1135 of mice infected with *Salmonella* wild-type (STM) and *cycA* mutant (Δ*cycA*), at day 3

1136 post-infection. n = 5 mice per group. Statistical significance was assessed using

1137 Mann-Whitney U test. ns, not Significant.

1138

1139 **Figure 1_figure Supplement 7.** Growth curves of *Salmonella* WT and the *gabP*

1140 mutant (Δ*gabP*) in N-minimal medium supplemented with β-alanine (1 mM) as the

1141 sole carbon source. Data are presented as mean ± SD, n = 3 independent experiments.

1142

1143 **Figure 1_figure Supplement 8.** Growth curves of *Salmonella* WT and the *gabP*

1144 mutant (Δ*gabP*) in M9 minimal medium supplemented with β-alanine (1 mM) as the

1145 sole carbon source. Data are presented as mean ± SD, n = 3 independent experiments.

1146

1147 **Figure 2_figure Supplement 1.** Growth curves of *Salmonella* WT, *panD* mutant

1148 (Δ*panD*) and the complemented strain (c*panD*) in LB medium. Data are presented as

1149 mean ± SD, n = 4 independent experiments.

1150

1151 **Figure 2_figure Supplement 2.** Growth curves of *Salmonella* WT, *panD* mutant

1152 (Δ*panD*) and the complemented strain (c*panD*) in RPMI-1640 medium (B). Data are

1153 presented as mean ± SD, n = 4 independent experiments.

1154

1155 **Figure 2_figure Supplement 3.** Fold intracellular replication (20 h *vs.* 2 h) of

1156 *Salmonella enterica* serovar Typhi Ty2 wild type (WT), the *panD* mutant (Δ*panD*) in

1157 human THP-1 monocyte like cell line (ATCC TIB-22). Data are presented as mean ±

1158 SD, n = 4 independent experiments. Statistical significance was assessed using

1159 two-sided Student's *t*-test. ns, not Significant.

1160

1161   **Figure 4_figure Supplement 1.** Expression of SPI-2 is shown in the Z-score

1162   transformed heatmap, with orange representing higher and blue representing lower

1163   abundance.

1164

1165   **Figure 4_figure Supplement 2.** Expression of SPI-1 is shown in the Z-score

1166   transformed heatmap, with orange representing higher and blue representing lower

1167   abundance.

1168

1169   **Figure 4_figure Supplement 3.** Expression of SPI-3, SPI-4, and SPI-5 genes is

1170   shown in the Z-score transformed heatmap, with orange representing higher and blue

1171   representing lower abundance.

1172

1173   **Figure 1_source data 1.** The numerical source data corresponds to Figure 1.

1174

1175   **Figure 2_source data 1.** The numerical source data corresponds to Figure 2.

1176

1177   **Figure 3_source data 1.** The numerical source data corresponds to Figure 3.

1178

1179   **Figure 4_source data 1.** The numerical source data corresponds to Figure 4.

1180

1181   **Figure 5_source data 1.** The numerical source data corresponds to Figure 5.

1182

1183 **Figure 6_source data 1.** The numerical source data corresponds to Figure 6.

1184

1185 **Figure 1_figure Supplement 1_source data 1.** The numerical source data

1186 corresponds to Figure 1_figure Supplement 1.

1187

1188 **Figure 1_figure Supplement 2_source data 1.** The numerical source data

1189 corresponds to Figure 1_figure Supplement 2.

1190

1191 **Figure 1_figure Supplement 3_source data 1.** The numerical source data

1192 corresponds to Figure 1_figure Supplement 3.

1193

1194 **Figure 1_figure Supplement 4_source data 1.** The numerical source data

1195 corresponds to Figure 1_figure Supplement 4.

1196

1197 **Figure 1_figure Supplement 5_source data 1.** The numerical source data

1198 corresponds to Figure 1_figure Supplement 5.

1199

1200 **Figure 1_figure Supplement 6_source data 1.** The numerical source data

1201 corresponds to Figure 1_figure Supplement 5.

1202

1203 **Figure 1_figure Supplement 7_source data 1.** The numerical source data

1204 corresponds to Figure 1_figure Supplement 5.

1205

1206 **Figure 1_figure Supplement 8_source data 1.** The numerical source data

1207 corresponds to Figure 1_figure Supplement 5.

1208

1209 **Figure 2_figure Supplement 1_source data 1.** The numerical source data

1210 corresponds to Figure 2_figure Supplement 1.

1211

1212 **Figure 2_figure Supplement 2_source data 1.** The numerical source data

1213 corresponds to Figure 2_figure Supplement 2.

1214

1215 **Figure 2_figure Supplement 3_source data 1.** The numerical source data

1216 corresponds to Figure 2_figure Supplement 3.

1217

1218 **Figure 4_figure Supplement 1_source data 1.** The numerical source data

1219 corresponds to Figure 4_figure Supplement 1.

1220

1221 **Figure 4_figure Supplement 2_source data 1.** The numerical source data

1222 corresponds to Figure 4_figure Supplement 2.

1223

1224 **Figure 4_figure Supplement 3_source data 1.** The numerical source data

1225 corresponds to Figure 4_figure Supplement 3.

1226 **Supplementary Table 1 Strains and plasmids used in this study**

Plasmid or strain	Genotype or description	Source
Plasmid		
pSim17	temperature-induced λ-Red recombinase system; BlaS^R	Laboratory collection
pKD3	template plasmid containing the Cm cassette for λ-Red recombination; Cm^R	Laboratory collection
pKD4	template plasmid containing the Km cassette for λ-Red recombination; Km^R	Laboratory collection
pBR322	low-copy-number expression vector; Ap^R	Laboratory collection
pMS402	Containing a promoterless *luxCDABE* reporter gene cluster; Km^R	Laboratory collection
pBR322-*panD*	pBR322 carrying the STM *panD* gene; Ap^R	This study
pMS402-*panD*	pMS402 carrying the STM *panD* promoter; Km^R	This study
strain		
wild-type (WT) STM	wild-type *Salmonella* enterica serovar Typhimurium strain ATCC 14028s	Laboratory collection
Δ*panD*	WT strain *panD*::Cm; Cm^R	This study
c*panD*	Δ*panD* containing plasmid pBR322-*panD*; Cm^R , Ap^R	This study
Δ*cycA*	WT strain cycA::Cm; Cm^R	This study
Δ*fadAB*	WT strain *fadAB*::Cm; Cm^R	This study
Δ*metR*	WT strain *metR*::Cm; Cm^R	This study
Δ*leuABCD*	WT strain *leuABCD*::Cm; Cm^R	This study
Δ*leuO*	WT strain *leuO*::Km; Km^R	This study
Δ*hisABCDFGHL*	WT strain *hisABCDFGHL*::Cm; Cm^R	This study
Δ*kdpABC*	WT strain *kdpABC*::Cm; Cm^R	This study
Δ*mglABC*	WT strain *mglABC*::Cm; Cm^R	This study
Δ*potFGHI*	WT strain *potFGHI*::Cm; Cm^R	This study
Δ*znuA*	WT strain *znuA*::Cm; Cm^R	This study
Δ*argT*	WT strain *argT*::Km; Km^R	This study
Δ*gabP*	WT strain *gabP*::Cm; Cm^R	This study

1227

1228

Supplementary Table 2 Primers involved in this study

Targets		Primer Sequences (5′-3′)
Primers used for the construction and identification of mutant strains		
Δ*cycA*	construction	F: GTATCATAGACCAAAGGCCGTAGAGCCCGCACAACACAG ACAGGTACAGGAAGAAAAACGTGTAGGCTGGAGCTGCTTC
		R: CCTCCAAATCAACGTTACGCTGTAAGCCCGGTAAGCGCC AGCGCCACCGGGCAAAACAACATATGAATATCCTCCTTAG
	identification	F: TGTGAGCTAATTCGCATTATCAAAG
		R: ACGGCATTAATGAGATGATTGATGA
Δ*panD*	construction	F: CGGATTCGCTGGAAACCATGTCGCGGCTGATCAGCAACC AGCCGCAGGGATAAGGACTAGTGTAGGCTGGAGCTGCTT
		R: TTTCCAAAGCAGGCCACAAGCGCCTGCTTAGCCAAGGTA AACGACAGGGTAAAGAAGTTCATATGAATATCCTCCTTAG
	identification	F: CGGTTATAACGCGACCGCGATCAAA
		R:GCTGCTGGAGCTGACGGAAACCAGC
Δ*fadAB*	construction	F: GGATTCGCTCAGTTGCCGCTGCGCTGCAATGCGAGTTATT TAGGGGATATTATCTTTGAGTGTAGGCTGGAGCTGCTTC
		R: AATACACACTTCGCTTCATCTGGTACGACCAGATCACTTT GTGGATTCAGGAGACTGACCATATGAATATCCTCCTTAG
	identification	F: AAACGGCAACTAAACTGTTTCCCGT
		R: AGTGATTCCATTTTTTACCCTTCTG
Δ*metR*	construction	F: TAAAGCTTGGCCAGTGGGGCTGCTGATGCCAGATGGCAA GATTAACCAGCAGTCCTGCGGTGTAGGCTGGAGCTGCTTC
		R: GAAAGTCCTTCACTTCGCCATGAACAAATTGCGCTTGAG GAATATACAGTACCTTTACACATATGAATATCCTCCTTAG
	identification	F: CCATTGCAGCGTGGTTTCCGGCAGC
		R: TGCAAGCCGAACTATTTAGAGTTTT
Δ*leuABCD*	construction	F: GAATATAATGGCAACCGGGTGATATTGCATAACCTGTAG GCCCGGTAAGGCGTCAAGGGGTGTAGGCTGGAGCTGCTTC
		R: CCGTTGCGCGGGTTTTTTTATGCCTGACGCAAGGCGCCCC TGGAGACAAGGACCACATCCATATGAATATCCTCCTTAG
	identification	F: TTTTTTATTTCGGCTGTTTCTGGAT
		R: GTTGACATTAAACGGCATATCCAGT
Δ*leuO*	construction	F: CCAGAAAAAGGGAGTTAAGCGTGACAGTGGAGTTAAATGTG TAGGCTGGAGCTGCTTC
		R: ATAAACCAGAATTTGTTTCTGATTTATTCTGCCCGGTTCATATG AATATCCTCCTTAG
	identification	F: CATTATGAATCGCAATGGTGTGAC
		R: CTCCGTCTGAATCACACCTGGT

 Supplementary Table 2 Primers involved in this study (continue)

Targets	Primer Sequences (5'-3')	
Primers used for the construction and identification of mutant strains		
Δ*hisABCDFGHL*	construction	F: GGCGTAAAAGTGGTTTAGGTTAAAAGGTATCAAATGAATAAGCATTCATCGGAATTTTTGTGTAGGCTGGAGCTGCTTC
		R: GATGGCTGGCATCAGGCCATCGGTTTTTTCCCAGTCCAGCTCGCGGCGTTGTTGCTCTGCATATGAATATCCTCCTTAG
	identification	F: CCTAACCAACCTAAACCGACAATTG
		R: AAACGCTGTTTCGTGCGTGAGAAGA
Δ*kdpABC*	construction	F: GTGGGGCGGGGGCGTTTGTTCCAACAAGCGATCCGGATCAGGACGCAAGGGTTCGTCGTGTGTAGGCTGGAGCTGCTTC
		R:CTGGTTTTCTTACTTTTAGGTTATCTGGTCTATGCCCTGATTAATGCGGAGGCGTTCTGCATATGAATATCCTCCTTAG
	identification	F: CAGAATGGTCAGCCCGTCAAG
		R: ATACTTTTTTTACACTCCGCCC
Δ*mglABC*	construction	F: CCGGCGCGCAGTTTCGCTGCGCCGGGTTATCTGATAAGCAAGGCAATAGGTCTGGATAAGTGTAGGCTGGAGCTGCTTC
		R: GTGCTGAACAGCCGGGCATTTTTTTACGCTATACCCTACATAATAAAACCGGAGCTACCCATATGAATATCCTCCTTAG
	identification	F: AAAAAGAGCACGCGTTAACACTGTA
		R: AGGCTCTAAATACGCTTCGGC
Δ*potFGHI*	construction	F: ATGTTTTAAACCACGCCTAATGGGTTCATTTGTTAACGGATTTCAGAAGGAAAGCGATGGTGTAGGCTGGAGCTGCTTC
		R: GAAAATGATGCCGCGACCCGCGCGGCAACGCATTGCCATAGTGGAAGATTTTAGTGGCTCATATGAATATCCTCCTTAG
	identification	F: AAAGGCGTTTTTTAATCTGGGCTAT
		R: TATTCATAATCATCAGCACATCGAG
Δ*znuA*	construction	F: AGATTATTAAATGCCAGGGCGACAGAGCGGGCTATCTGTTGCACGTATTCACTTCCTCGGTGTAGGCTGGAGCTGCTTC
		R: AAGTGATAGAATGTTATAATATCACATTTCACACATTCATTACGATGATTAGTCGCATTCATATGAATATCCTCCTTAG
	identification	F: AGGGAACGAATCTCGCTTTTCTC
		R: ATTCAAGCGACACGTCAGAGAGGAC
Δ*argT*	construction	F: CGTGATAGTTCCCCAGCGCGGCGCGTTATCCCCTTCCCGTGTAGGCTGGAGCTGCTTC
		R: GTAAAACATAAGAAAATGACGCCACTTGAGGGGTATGTCATATGAATATCCTCCTTAG
	identification	F: CAAAGCATTCTACCTTGTCAGCCGG
		R: CATACCTGCTATCTTCAACATCAGG

1232

1233

1234

1235

1236

 Supplementary Table 2 Primers involved in this study (continue)

Targets		Primer Sequences (5′-3′)
Primers used for the construction and identification of mutant strains		
Δ*gabP*	construction	F: CCGGTAAACGAGCGCAATAACAATAAAGAGGTTTTAGGGTGTAGGCTGGAGCTGCTTC
		R: TTACTCCTGAGAAGCTTCCCGGCTGACGCCGGGAATAACATATGAATATCCTCCTTAG
	identification	F: CTTTGATGAGGCGAAACAAGC
		R: AATGTCATTTTTAAGCCAGCGAT
Primers used for the construction of complementary strains		
pBR322-*panD*		F: CCCTTTCGTCTTCAAGAATTCGCTGCTGGAGCTGACGGAAAC
		R: TGCGTCCGGCGTAGAGGATCCTCAGGCAACCTGTACCGGAAT
pBR322		F: GACACGGAAATGTTGAATACTCATA
		R: CAAGGAATGGTGCATGCAAG
Primers used for luciferase reporter system		
pMS402-*panD*		F: CGAGGCCCTTTCGTCTTCACCTCGAGTCAAGGAAATCCTGGTCAATG
		R: ATTTTGCGGCCGCAACTAGAGGATCCCGGGTAATCGTGAGCTGC
pMS402		F: CTGTCTCTTGATCAGATCTT
		R: TTATTTCCATTACAATCAAT
Primers used for RT-qPCR analysis		
16s		F:GAAAGCGTGGGAGCAAAC
		R:ACATGCTCCACCGCTTGTG
panD		F:AAGGTTCCTGTGCCATTGAC
		R:ACCGTTCACCGAGATGATTC
fadH		F:GAAACCCACATGCCGATAAC
		R:AAACTGACCGCCAATTTCAC
fadB		F:GTTCGCTGTCTGGAAGAAGG
		R: TACTGCTGCGCCATATCAAG
fadI		F:TCTGACGCTTTTTGACATGC
		R:TTCATGCAACGTTTGGGTTA
fadD		F:AAGATCTGGCTTTCCTGCAA
		R:CCCAGCTCGATAAAAAGCAG
fadE		F:AATTGGTCGTCGTCACTTCC
		R: GCGTCCTACCGACAAACATT
metR		F:TTTGGTGATGACGTCGGATA
		R:CGGGTAAATCAACAGCGTTT
hisG		F:GTGCGTGATGATGACATTCC
		R:GGAGGTGCGGATATGAGGTA
hisF		F:TGTATTGTCGTCGGGATTGA
		R:TTTTCAACTGCGTCAGATCG
leuD		F:CCGGAATTCGTGTTGAACTT
		R:GTAGAAGATGTCGGCGAAGC
leuO		F:AGATATGGGCAAACCACAGC
		R: CCTGACGGACTGAACCAAAT

 Supplementary Table 2 Primers involved in this study (continue)

Targets	Primer Sequences (5′-3′)
Primers used for RT-qPCR analysis	
mglA	F:AGCGATTAACCACGGTTTTG
	R:ATTGGGTGTCGCTTTTCATC
mglB	F:AGAATCCGGTGTGATTCAGG
	R:CAGCCAGGCATCCATCTTAT
kdpA	F:ACTAATCTGGCGCAGATGCT
	R:ACCCTCCATGTTGACGCTAC
kdpB	F:TCGGTAAGCAGATGCTGATG
	R:CCAGCGGGATCAGAAAAATA
potG	F:ATTTACGAGCATCCGACGAC
	R:AGAGGCATCTGCATCGACTT
potF	F:TCTGATGGTCGTCTCTGTCG
	R:GCCTTCCAGCACTTCGTTAG
znuA	F:TTAAACCGCTTGGGTTCATC
	R:AACGGTTTTACATCGGCAAG

A

Liver
Spleen
Bacterial CFU determination

Immunofluorescence staining of liver slice

H&E staining of liver slice

i.p. 5,000 CFU

Euthanize

Mice were observed for 10 days for survival and body weight was documented.

B

Survival (%)

WT
ΔpanD
cpanD

P < 0.0001

P < 0.0001

Days post infection

C

Body weight (g)

WT vs ΔpanD P < 0.0001

WT
ΔpanD
cpanD

Days post infection

D

log₁₀(CFU/g liver)

P < 0.0001
P < 0.0001

WT   ΔpanD   cpanD

log₁₀(CFU/g spleen)

P < 0.0001
P < 0.0001

WT   ΔpanD   cpanD

3dpi body weight (g)

P < 0.0001
P = 0.0001

WT   ΔpanD   cpanD

E

Salmonella   Nucleus   Merge

WT

ΔpanD

cpanD

Bacteria per Macrophage (liver)

P < 0.0001

P < 0.0001

WT   ΔpanD   cpanD

F

Uninfect   WT   ΔpanD   cpanD

A Zinc level/ 10^5 macrophages (RAW264.7 cell)
P < 0.0001
WT
Δ*panD*

B Fold intracellular replication (20 h *vs.* 2 h)
P = 0.0002
P = 0.0004
P < 0.0001
P = 0.0422
WT
Δ*panD*
Δ*znuA*
Δ*panD*Δ*znuA*

C Fold intracellular replication (20 h *vs.* 2 h)
P = 0.0020
ns
WT
Δ*panD*
Δ*panD*+100μM ZnSO₄

D Fold intracellular replication (20 h *vs.* 2 h)
P = 0.0028
P = 0.0027
WT
Δ*znuA*
Δ*znuA*+1mM β-alanine

E

Macrophage

Zn²⁺

β-alanine — Uptake by *Salmonella* → SCV *Salmonella* — Promote *Salmonella* growth → SCV

Dissemination and systemic infection

L-aspartate
PanD
β-alanine → β-alanine → ZnuABC ↑
Zn²⁺ uptake

Macrophage
SCV (*Salmonella*-containing vacuole)
Salmonella
Zn²⁺
Activation
Upregulation