Title

Role of Thalamus in Human Conscious Perception Revealed by Low-Intensity Focused Ultrasound Neuromodulation

Author list

Hyunwoo Jang[1,2], Panagiotis Fotiadis[2,3], George A. Mashour[1,2,3,4,5], Anthony G. Hudetz[1,2,3,4], Zirui Huang[1,2,3,4,*]

Affiliations

[1] Neuroscience Graduate Program, University of Michigan, Ann Arbor, MI 48109, USA

[2] Center for Consciousness Science, University of Michigan Medical School, Ann Arbor, MI 48109, USA

[3] Department of Anesthesiology, University of Michigan Medical School, Ann Arbor, MI 48109, USA

[4] Michigan Psychedelic Center, University of Michigan Medical School, Ann Arbor, MI, 48109, USA

[5] Department of Pharmacology, University of Michigan Medical School, Ann Arbor, MI, 48109, USA

* Correspondence and requests for materials should be addressed to Z.H.

(Email: huangzu@med.umich.edu)

ABSTRACT

The neural basis of consciousness remains incompletely understood. While cortical mechanisms of conscious perception have been extensively investigated in humans, the role of subcortical structures, including the thalamus, remains less explored. Here, we elucidate the causal contributions of different thalamic regions to conscious perception using transcranial low-intensity focused ultrasound (LIFU) neuromodulation. We hypothesize that modulating distinct thalamic regions alters perceptual outcomes derived from Signal Detection Theory. We apply LIFU to healthy human anterior (transmodal-dominant) and posterior (unimodal-dominant) thalamic regions, further subdivided into ventral and dorsal regions, during a near-threshold visual perception task. We show that high duty cycle modulation of the ventral anterior (VA) part of thalamus enhances object recognition sensitivity. Sensitivity enhancement magnitude correlates with the core-matrix cell compositions of the stimulated thalamic region. Connectivity analysis of a large-scale functional magnetic resonance imaging dataset confirms strong transmodal connectivity of VA thalamus with frontoparietal and default-mode networks. We also demonstrate target-invariant effects of high duty cycle LIFU disrupting object categorization accuracy. These findings provide causal insight into the cytoarchitectural and functional organization of the thalamus that shapes human visual experience, especially the role of matrix cell-rich, transmodal-dominant VA thalamus.

INTRODUCTION

The neural basis of consciousness remains an active area of neuroscientific investigation[1–7]. While cortical mechanisms of conscious perception contents have been extensively studied[8–12], the role of subcortical structures in encoding, enabling, or modulating such contents remains less explored[7,8,13]. With its intricate connections to the cortex, the thalamus is a central structure that is well-positioned to orchestrate conscious experience[13,14]. Traditionally, thalamic nuclei have been categorized into specific and nonspecific types[15]. In general, specific nuclei relay modal sensory information to designated cortical areas, whereas nonspecific nuclei project more diffusely and support cortical arousal or higher-order cognition[14,16]. For instance, the lateral geniculate nucleus, considered a specific nucleus, relays retinal input to the primary visual cortex[15], while the nonspecific intralaminar thalamus is involved in controlling arousal, attention, and global states of consciousness[17–19].

Recent studies suggest that cortical areas are functionally organized along a unimodal-to-transmodal continuum rather than distinct categories[20,21]. Within the cortex, unimodal areas process sensorimotor information, whereas transmodal areas integrate information and support complex cognitive functions. Building on this functional organization of the cortex, our previous work revealed a unimodal-transmodal gradient in thalamocortical connectivity[20]. This connectivity gradient features anterior and medial thalamic regions connecting predominantly with transmodal cortices, whereas posterior thalamic regions connect primarily with unimodal cortices. A similar gradient is also reflected by the cytoarchitecture of the thalamus: diffusely-projecting calbindin-expressing matrix cells are enriched anteriorly, and narrowly-projecting parvalbumin-expressing core cells are concentrated in posterior thalamus[20,22]. Notably, both the connectivity gradient and cytoarchitectural gradient support the hypothesis that the matrix cell-rich anterior thalamus is crucial for conscious perception[18,23–26]. Disruption of this region has also been linked to loss of consciousness[20].

A promising approach to investigate the causal contribution of thalamic gradients to human consciousness, more specifically conscious sensory perception, is to selectively modulate the activity of specific thalamic subregions. Traditional methods such as deep-brain stimulation, optogenetics, and transcranial electric/magnetic stimulation are either invasive or lack sufficient spatial resolution[27,28]. In contrast, transcranial low-intensity focused ultrasound (LIFU; also referred to as tFUS) offers a non-invasive approach with higher precision for targeting deep structures[29–33]. Pioneering work has applied LIFU to human subcortical regions, including the thalamus, putamen, and basal ganglia[34–37]. Although the mechanisms underlying LIFU's effects are still under investigation, with theories ranging from thermal effects[38] to membrane pore formation[39] and mechanosensitive channel activation[40], the ability of LIFU to modulate neural activity and behavior is well-supported[29,39,41,42]. Nevertheless, LIFU's potential for modulating human conscious perception and for probing its mechanisms, particularly in the visual domain, remains largely unexplored[43,44].

LIFU has been shown to modulate neural activity in a bidirectional manner, with the directionality governed by sonication parameters such as the duty cycle (DC; the percentage of time the ultrasound is actively transmitted within each pulse cycle). Early work using neuroimaging, electrophysiological, and behavioral measures indicates that higher DC values generally elicit excitatory effects, whereas lower DC values result in inhibition[33–35,39,40,45,46]. This pattern is supported by both computational models and *in vivo* experiments, which demonstrate that 70%

DC effectively lowers neuronal activation thresholds and facilitates excitation[40,47–49]. In contrast,
5% DC is associated with diminished evoked potentials and suppressed blood oxygen level-
dependent (BOLD) responses consistent with functional inhibition[40,49–51]. These observations
align with the neuronal intramembrane cavitation excitation model, which posits that the value of
DC determines the direction of neuromodulation[39,47,49].

In this work, we aim to elucidate the causal roles of human thalamus in conscious visual
perception by applying LIFU with region- and parameter-specific precision. We hypothesize that
targeting distinct thalamic regions—each positioned along the unimodal-to-transmodal gradient—
and varying the sonication duty cycle will lead to distinct changes in perceptual outcomes. To test
this hypothesis, we sequentially modulate four gradient-defined thalamic areas[52] (ventral anterior,
ventral posterior, dorsal anterior, and dorsal posterior) while healthy human volunteers perform a
near-threshold visual task. Perceptual changes are quantified using Signal Detection Theory
(SDT) metrics. To interpret the observed effects at cytoarchitectural and network levels, we
conduct complementary analyses with core-matrix cell composition and thalamocortical
connectivity data. Our findings reveal that high duty cycle modulation of the matrix cell-rich,
transmodal-dominant ventral anterior thalamus selectively enhances object recognition sensitivity.
Additionally, we demonstrate target-invariant and duty cycle-dependent effects on the accuracy
of object categorization, observed regardless of the target region.

RESULTS

Sixty participants (age: 25.9 ± 6.3, mean ± SD; 38 females, 22 males) were randomly assigned
to one of two duty cycle (DC) groups. One group received LIFU with a high duty cycle (70%),
while the other received LIFU with a low duty cycle (5%) with equal spatial-peak temporal-average
intensity of I_{spta} = 0.72 W cm^{-2} (Fig. 1a). Six participants were excluded from analysis due to
technical issues, leaving 54 participants in the final analysis (27 in each group; Supplementary
Fig. 1).

The experiment involved key steps including transducer setup, image contrast titration, and six
blocks of the visual task (Fig. 1b-d). There were two outer LIFU-OFF baseline blocks (Baseline-1
and Baseline-2) and four inner LIFU-ON blocks. During each LIFU-ON block, LIFU was directed
toward one of four regions of the left thalamus, including the ventral anterior (VA), ventral posterior
(VP), dorsal anterior (DA), and dorsal posterior (DP) thalamus (Fig. 1e; see Supplementary Table
1 for sonication parameter summary; see Methods: Target locations for exact coordinates). The
order of target regions was pseudo-randomized and counterbalanced among participants, and
each participant received a consistent duty cycle throughout the session.

We employed a well-established near-threshold visual object recognition and categorization
task[8,9]. Four object categories were used: faces, houses, human-made objects, and animals (Fig.
1f). Before the main task, using an adaptive staircase paradigm, we individually titrated the image
contrast (Fig. 1g) to determine the threshold of conscious perception (i.e., a contrast yielding 50%
recognition rate)[8]. Here, "recognition" was operationally defined as the perception of an object
that makes sense in the real world, as opposed to meaningless noise-like patterns. During each
trial, participants responded to two questions: one regarding the category of the presented image
(i.e., "Which category does the image belong to?") and another concerning their subjective
perceptual experience (i.e., "Did you recognize the image?") (Fig. 1h).

Importantly, we also included scrambled images, which comprised 20 out of 100 trials. Therefore,
the recognition rates for real and scrambled images could be interpreted as hits and false alarms,
respectively (Fig. 1i). This enabled us to measure sensitivity (d') and decision bias (quantified by
criterion c) of object recognition within the SDT framework (see Methods: Signal detection theory
analysis). Sensitivity reflects the ability to discriminate real and scrambled images, and higher
criterion signifies a tendency to say "NO" (i.e., "I didn't recognize the image") to the Question-2,
reflecting a conservative stance. Categorization accuracy was also quantified for real and
scrambled images, independently of object recognition.

Before LIFU was administered (i.e., at Baseline-1), participants recognized 53.3 ± 16.1% (mean
± SD) of the real images (Supplementary Fig. 2a), which was not significantly different from the
intended 50% recognition rate, confirming a successful implementation of the staircase procedure.
See SI text: Perceptual outcomes at Baseline-1 and Supplementary Fig. 2 for detailed analysis of
baseline perceptual outcomes.

We examined whether LIFU induced any carry-over effects within our approximately one-hour
experimental timeframe by comparing two outer LIFU-OFF blocks (Baseline-1 and Baseline-2).
For both duty cycles, we found no significant difference in any perceptual outcomes ($p > 0.05$,
Wilcoxon signed-rank test, two-sided; Supplementary Fig. 3). These results suggest that LIFU did
not produce detectable long-lasting or cumulative effects on visual perception in our experimental
condition.

Fig. 1: Overview of the block design, experimental setup, and behavioral paradigm. a Randomized
study group assignment. Sixty subjects were randomly assigned to one of two groups, each consisting of
30 participants, corresponding to either a 70% or 5% DC. 54 participants (27 per group) were included in
the final analysis. **b** The timeline of experimental sessions, including initial preparation steps and the main
task. The main task included two outer baseline blocks and four inner LIFU-ON blocks with a pseudo-
randomized order of targeting regions. Each block comprised 100 trials. Each LIFU-ON block included 23
alternating 30-second ON and OFF epochs (total 11.5 min). Visual tasks were conducted continuously
within each block. Each ON epoch consists of 300 pulses (pulse repetition frequency = 10 Hz) at a fixed
DC and temporal-average intensity ($I_{spta.3}$ = 0.72 W cm^{-2}). The pulse-averaged intensity was higher at 5%
DC. **c** Measurement locations for head size, including width, depth, and side-to-top. **d** Illustration of the
transducer fixed by two perpendicular rubber bands on the participant's head. **e** Sagittal view for the four
target regions, including left VA, VP, DA, and DP. **f** Examples of real and scrambled images for each
category (face, house, human-made object, and animal). **g** Example of an animal image displayed at
different contrast levels. **h** Trial structure showing the sequence of events: a blank period, stimulus
presentation, delay, and two questions about categorization and recognition. **i** Illustration of the four trial
types (hit, miss, false alarm, and correct rejection). Due to copyright limitations, the actual images used in
our experiment are not shown. Copyright-free images included in this figure were obtained from

161 https://www.pexels.com. DC: duty cycle; VA: ventral anterior thalamus; VP: ventral posterior thalamus; DA:
162 dorsal anterior thalamus; DP: dorsal posterior thalamus.

Ultrasound beam analysis and targeting accuracy

For each LIFU-ON block, we assessed various beam profiles, including targeting accuracy, beam
shape, attenuation by tissue, and thermal effects (Fig. 2a). As proof of concept, we used a
pseudo-anatomical reference for beam targeting and post-hoc analysis (see Methods: Template
rescaling method for details). This was achieved by rescaling the MNI template image to match
each participant's head size measured before the main task (Fig. 1c). We validated this approach
on a separate dataset (n = 13) by comparing actual anatomical images with their rescaled pseudo-
anatomical images, finding an average deviation of 2.15 ± 0.82 mm (mean ± SD; Supplementary
Fig. 4).

In a free-field condition, our transducer produces a bullet-like beam shape with an aspect ratio ≈
9, measuring 30.3 mm in length and 3.3 mm in width (defined by -3 dB attenuation; Supplementary
Fig. 5). Given this elongated shape, the lateral deviation reported by the neuronavigation system
is a straightforward measure of the beam orientation accuracy. Here, lateral deviation is defined
as the Euclidean distance between the beam vector (originating from the transducer center) and
the target coordinate. The mean lateral deviation was 2.25 ± 0.81 mm (mean ± SD; Fig. 2b) and
did not differ significantly across regions (p = 0.2155, Friedman's test; see Source Data for
statistics), with maximal deviation not exceeding 5 mm, together confirming that the beam was
successfully oriented toward the targets.

We then simulated the beam on the segmented pseudo-anatomical image *in silico*. In line with
the literature[53–55], we observed substantial intensity attenuation of -8.78 ± 0.56 dB (mean ± SD;
Fig. 2c), equating to approximately 87% energy loss. This yielded estimated pressure of 72.6 ±
4.2 kPa for 70% DC and 258.2 ± 18.3 kPa for 5% DC (mean ± SD; Supplementary Fig. 6). This
pronounced energy loss is mainly attributable to skull[53,54,56,57]. The maximal temperature rise
simulated across skin, skull, and brain was 0.23 ± 0.04 °C (mean ± SD) and didn't exceed 0.5 °C
(Fig. 2d), complying with ITRUSST risk recommendations[58]. Additionally, the beam depth (i.e.,
the Euclidean distance from the head-transducer contact point to the location of maximal
simulated intensity) was simulated as 67.4 ± 3.4 mm (mean ± SD; Supplementary Fig. 7),
approximately 15% shorter than the free-field focal depth of 80.7 mm (Supplementary Table 1,2),
consistent with skull-induced beam shortening reported previously[55].

Next, we evaluated how LIFU beams covered each thalamic region, as visualized by the mean
relative intensity across subjects (Fig. 2e-i; see Supplementary Fig. 8-10 for individual trajectories).
Because the LIFU beam originated from the right temple (Fig. 2e) and was effectively shortened
by the skull (Supplementary Fig. 7), we observed significant bilateral coverage for all four target
regions (Fig. 2g-i). To quantify this further, we computed a coverage ratio defined by how much
of the "beam-present" volume (defined by > 50% intensity) belongs to the thalamic (or other non-
thalamic) regions (Fig. 2j-m). We found excellent beam localization in bilateral VA, VP, and DA
(Fig. 2j-l). In the case of DP targeting, the beam partially encompassed bilateral VP, presumably
because the DP regions lie lateral to VP (Fig. 2m). Overall, these post-hoc analyses confirm
successful targeting of each thalamic region.

Fig. 2: Post-hoc LIFU beam analysis and targeting accuracy. a Flowchart summarizing the beam
analysis and simulation pipeline. **b** Lateral deviation of the beam from its target, defined as the Euclidean
distance between the beam vector and the target coordinate. **c, d** Simulated **c** tissue-induced attenuation
and **d** maximal tissue temperature. In panels b-d, boxes indicate interquartile ranges, and white circles
show median values. **e** Three-dimensional rendering of the mean relative beam intensity across participants.
Colored regions highlight voxels where the mean relative intensity exceeds 10%. An arrow indicates the
LIFU beam originating from the right side. **f** Zoomed-in view of panel e, with dashed lines indicating the
locations of cross-sections. **g-i** Cross-sectional views in the **g** sagittal, **h** coronal, and **i** axial planes,
displaying mean relative intensity and outlines of bilateral thalamic regions. Higher opacity reflects greater

intensity. **j-m** Bilateral coverage percentages for beams targeting left **j** VA, **k** VP, **l** DA, and **m** DP. The
coverage percentage indicates how much of a "beam-present" volume (defined by > 50% intensity) falls
within the thalamic (or other) regions. Boxes indicate mean values, and error bars show standard error. For
panel b, VA: n = 51; VP: n = 50; DA: n = 53; DP: n = 54. For panels c-m, VA: n = 44; VP: n = 44; DA: n =
45; DP: n = 46. Color coding: VA (red), VP (green), DA (blue), and DP (yellow). VA: ventral anterior thalamus;
VP: ventral posterior thalamus; DA: dorsal anterior thalamus; DP: dorsal posterior thalamus; LH: left
hemisphere; RH: right hemisphere. Source data are provided as a Source Data file.

Effects of LIFU administration on perceptual outcomes

To determine whether LIFU-targeting of different thalamic regions produces differential effects on
visual perception, we conducted omnibus F-tests using linear mixed-effects (LME) models for
each perceptual outcome and duty cycle condition. Following the observation that the two
baseline conditions did not differ significantly (Supplementary Fig. 3), we averaged them into a
single baseline measure, yielding five targeting conditions (baseline plus four thalamic targets)
for comparison.

The omnibus analysis revealed significant regional effects for three conditions (Table 1). At 70%
DC, we observed significant omnibus effects for sensitivity d' ($F(4,90)$ = 4.6129, p_{FDR} = 0.0118)
and real-image accuracy ($F(4,107)$ = 5.4470, p_{FDR} = 0.0060). At 5% DC, criterion c showed a
significant omnibus effect ($F(4,77)$ = 4.1100, p_{FDR} = 0.0181). Other perceptual measures,
including hit rate, false alarm rate, and scrambled image accuracy, did not demonstrate significant
effects after FDR correction at either duty cycle condition. To ensure that these effects were not
artifacts of baseline averaging, we conducted control analyses using each baseline condition
separately, showing that the main findings remained robust across both individual baselines
(Supplementary Table 3). We also confirmed that none of the nuisance variables included in our
LME model showed significant effects in omnibus ANOVA (all p_{FDR} > 0.05; Supplementary Table
4; see Methods: Statistics for details).

Table 1: Omnibus ANOVA for the effect of sonication target region on behavioral outcomes.

DC	Perceptual outcome	df₁	df₂	*F*-value	*p*-value	FDR-corrected *p*-value
70%	Hit rate	4	107	1.3407	0.2596	0.5192
	FA rate	4	110	0.6525	0.6263	0.8351
	Sensitivity *d'*	4	90	4.6129	0.0020	0.0118
	Criterion *c*	4	89	0.5325	0.7122	0.8546
	Real image accuracy	4	107	5.4470	0.0005	0.0060
	Scrambled image accuracy	4	111	0.2665	0.8989	0.9553
5%	Hit rate	4	104	2.3734	0.0570	0.1710
	FA rate	4	104	2.0005	0.0999	0.2398
	Sensitivity *d'*	4	78	0.7452	0.5641	0.8351
	Criterion *c*	4	77	4.1100	0.0045	0.0181
	Real image accuracy	4	103	0.9165	0.4574	0.7841
	Scrambled image accuracy	4	106	0.1658	0.9553	0.9553

Significant values (FDR-corrected p < 0.05) are highlighted as red. DC: duty cycle; df: degree of freedom; FA: false
alarm.

For the three conditions that demonstrated significant omnibus effects, we conducted post-hoc
pairwise comparisons to identify which specific thalamic targets potentially drove the observed
regional differences (Fig. 3; see Supplementary Fig. 11 for all other perceptual outcomes).

At 70% DC, LIFU targeting of the VA nucleus specifically enhanced sensitivity d' compared to
baseline (VA vs. baseline: p_{FDR} = 0.0246; paired Cohen's d = 0.9164) and significantly
outperformed all other target regions (Fig. 3a; VA vs. VP: p_{FDR} = 0.0246, VA vs. DA: p_{FDR} = 0.0034,
VA vs. DP: p_{FDR} = 0.0119; see Supplementary Table 5 for full statistics). This pattern demonstrates
that the omnibus effect was driven primarily by VA-specific enhancement rather than a
generalized effect.

We also identified that the increase in sensitivity during VA thalamus at 70% DC sonication
became more robust as the lateral deviation of the beam decreased (Supplementary Fig. 12).
Specifically, the height of the fitted Gaussian curve was significantly positive for 70% DC (p_{FDR} =
0.0286) but not for 5% DC (p_{FDR} = 0.7679; see Source Data for full statistics). These findings
support the target-specificity of this effect and validate our targeting precision.

For categorization accuracy of real images, a consistent decrease across all four thalamic regions
relative to baseline was observed (Fig. 3b; baseline vs. all four regions: p_{FDR} < 0.05; average
paired Cohen's d = -0.4545; Supplementary Table 5), suggesting a target-invariant disruptive
effect of high-DC thalamic sonication on visual categorization tasks.

At 5% DC, the effects of LIFU on criterion c revealed a more complex pattern. We observed
elevated criterion (i.e., shift toward a conservative stance) in both VA and VP sonication compared
to baseline and to dorsal targets (DA and DP), although many of these pairwise comparisons
reached only marginal significance after FDR correction (Fig. 3c; VA vs. DP and VP vs. DP: p_{FDR}
< 0.05; Supplementary Table 5). This nuanced pattern suggests regionally heterogeneous effects
that are not uniformly robust across all comparisons.

Fig. 3: Target-specific changes in perceptual outcomes during human thalamic LIFU application.
Perceptual outcomes that demonstrated significant omnibus effects, measured at **a-b** 70% and **c** 5% DC,
including **a** sensitivity d', **b** categorization accuracy for real images, and **c** criterion c, are shown. Asterisks
and pounds denote significant differences in post-hoc pairwise comparisons (asterisk: FDR-corrected p <
0.05; pound: FDR-uncorrected p < 0.05). Boxes indicate interquartile ranges. Median values are marked

by white circles. Dashed lines are reference values for each measure: no discrimination of $d' = 0$ (a), chance
accuracy of 0.25 (b), and no bias of $c = 0$ (c). For panel a, Baseline: $n = 25$; VA: $n = 23$; VP: $n = 23$; DA: n
$= 21$; DP: $n = 20$; for panel b, Baseline: $n = 27$; VA: $n = 26$; VP: $n = 26$; DA: $n = 27$; DP: $n = 27$; for panel c,
Baseline: $n = 24$; VA: $n = 21$; VP: $n = 19$; DA: $n = 18$; DP: $n = 20$. DC: duty cycle; FA: false alarm; VA:
ventral anterior thalamus; VP: ventral posterior thalamus; DA: dorsal anterior thalamus; DP: dorsal posterior
thalamus. Statistics and source data are provided as a Source Data file.

Cytoarchitectural correlates of LIFU-induced perceptual modulation

To test whether thalamic cytoarchitecture predicts the observed perceptual effects, we conducted
correlation analyses between the core-matrix cell composition of stimulated regions and
behavioral outcomes. Core and matrix cells can be distinguished by their expression of calcium-
binding proteins, with core cells expressing higher levels of Parvalbumin (PVALB) and matrix cells
expressing higher levels of Calbindin (CALB1)[22]. We quantified the relative composition of these
cell types using the CP_T metric, as defined by the difference between CALB1 and PVALB
expression levels derived from the Allen Human Brain Atlas, which can be used as a proxy for
relative matrix cell richness[20,22,59], i.e., a higher CP_T value indicates a more matrix-cell rich area.

For each LIFU-ON block, we calculated the mean CP_T value across beam-present voxels (> 50%
relative intensity; black contours in Supplementary Fig. 8-10). We then correlated these CP_T
values with baseline-subtracted changes in the three perceptual outcomes that survived omnibus
testing. The analysis revealed a significant positive correlation between CP_T and change in
sensitivity d' at 70% DC (Spearman's $\rho = 0.3479$, $p_{FDR} = 0.0135$; Fig. 4a), indicating that regions
with higher matrix cell density produced greater enhancement in perceptual sensitivity when
sonicated by high DC. In contrast, no significant correlations emerged for real image accuracy at
70% duty cycle ($\rho = -0.1026$, $p_{FDR} = 0.3590$; Fig. 4b) or criterion c at 5% duty cycle ($\rho = 0.1748$,
$p_{FDR} = 0.2304$; Fig. 4c, see Source Data for full statistics).

Fig. 4: Correlation between matrix cell richness at beam-present regions (*x*-axis) and behavioral
outcome changes (*y*-axis) across LIFU-ON blocks. CP_T reflects the relative difference between
Calbindin (CALB1) and Parvalbumin (PVALB) expressions. Positive and negative CP_T values indicate
matrix cell-richness and core cell-richness at the beam center (i.e., voxels exceeding 50% relative intensity),
respectively. Spearman's ρ and FDR-corrected *p*-values are provided. Circle colors indicate target regions.
Solid lines and shades show linear fits and 95% confidence intervals. VA: ventral anterior thalamus; VP:

ventral posterior thalamus; DA: dorsal anterior thalamus; DP: dorsal posterior thalamus. Statistics and
source data are provided as a Source Data file.

Functional connectivity between thalamic regions and the cortex

To gain deeper insights into the network-level neural correlates of both target-specific and target-
invariant effects of thalamic LIFU, we analyzed the connectivity profiles of the four thalamic
regions by utilizing a large-scale functional magnetic resonance imaging (fMRI) dataset of the
Human Connectome Project (n = 1009)[60].

First, we assessed the connectivity profiles of the four thalamic regions at the network level (Fig.
5a,b). For each thalamic region of interest (ROI), we calculated the functional connectivity with
400 cortical ROIs and averaged their percentile rank across seven canonical networks[61]. In line
with our prior research[20], we observed a graded shift in connectivity profiles along a unimodal-
transmodal gradient. The VA thalamus displayed transmodal dominance, exhibiting higher
connectivity with frontoparietal and default-mode networks (red curve in Fig. 5b) compared to the
other three thalamic regions. This was followed by DA thalamus, with a trend toward unimodal
dominance observed in DP and VP thalamus.

Second, we examined the thalamocortical connectivity at a finer spatial scale, specifically at voxel
level. We retained the cortical voxels exhibiting the top 10% strongest connectivity with each
respective thalamic region as a seed. The connectivity profiles of the four regions showed
considerable overlap, particularly within the more primary visual cortex (cyan regions in Fig. 5c).
Region-specific connectivity patterns were also observed. The VA thalamus was predominantly
connected with the medial prefrontal cortex (mPFC) and dorsolateral prefrontal cortex (dlPFC)
(red regions in Fig. 5d). The VP thalamus exhibited unique connectivity with the somatomotor and
auditory cortices (green regions in Fig. 5d, see also high connectivity with somatomotor network
in Fig. 5a). The DP thalamus showed unique clusters within various regions in the visual cortex
(yellow regions in Fig. 5d). The connectivity profile of the DA thalamus largely overlapped with
the VA thalamus (Supplementary Fig. 13), with only a few unique clusters located in left dlPFC
and right temporal-parietal junction (blue regions in Fig. 5d).

Fig. 5: Functional connectivity profiles between four thalamic regions and the cortex. a, b Network-level analysis. **a** Percentile ranks of functional connectivity of 400 cortical regions and each thalamic region, averaged within seven canonical Schaefer-Yeo networks (visual, somatomotor, dorsal attention, ventral attention, limbic, frontoparietal, and default-mode). Each dot represents an individual (n = 1009). White circles indicate median values and error bars denote interquartile ranges. **b** Distribution of mean connectivity ranks of unimodal (visual and somatomotor) vs. transmodal (frontoparietal and default-mode) networks with each thalamic region. Curves represent kernel-smoothed histograms across the direction of maximal variance. **c, d** Voxel-based analysis. Cortical voxels exhibiting the top 10% strongest connectivity with each thalamic region as a seed, that are **c** common across four thalamic regions (i.e., region-invariant) and **d** unique to each region (i.e., region-specific). VA: ventral anterior thalamus; VP: ventral posterior thalamus; DA: dorsal anterior thalamus; DP: dorsal posterior thalamus. Source data are provided as a Source Data file.

DISCUSSION

This study aimed to causally probe how different thalamic regions contribute to conscious perception in humans. Through low-intensity focused ultrasound (LIFU) neuromodulation, we identified a specific role for the matrix cell-rich, transmodal-dominant ventral anterior (VA) thalamus in modulating the sensitivity of conscious visual perception (Fig. 6a). Sonicating the VA thalamus with a high duty cycle (DC) significantly increased object recognition sensitivity, consistent with the strong functional connectivity of the matrix cell-rich VA thalamus with transmodal cortical areas, particularly the medial and dorsolateral prefrontal cortices. Additionally, we observed target-invariant decrease in object categorization accuracy under high DC sonication

361 (Fig. 6b). Together, these findings demonstrate the causal involvement of matrix cell-rich thalamic
362 regions in conscious perception and underscore the modulatory potential of thalamocortical
363 networks in shaping visual experience.

Fig. 6: Summary of thalamic LIFU effects on human visual perception. a Sensitivity increases when
the VA thalamus is modulated at 70% duty cycle. This is associated with its higher matrix cell density and
higher functional connectivity between the VA thalamus and transmodal cortical areas compared to other
thalamic nuclei. **b** Categorization accuracy decreases at 70% duty cycle in a target-invariant manner,
potentially driven by common interactions between the thalamus and the visual network (cyan). VA: ventral
anterior thalamus; VP: ventral posterior thalamus; DA: dorsal anterior thalamus; DP: dorsal posterior
thalamus; DC: duty cycle.

Being primarily nonspecific and matrix cell-rich, the VA thalamus likely interacts extensively with
thick-tufted layer-5 (TTL5) pyramidal neurons in the cortex, potentially promoting large-scale
cortical integration essential for conscious sensory processing[1,13,23,24,62]. This also supports
theoretical frameworks such as thalamic gating mechanisms and dendritic integration theory,
which suggest that matrix cells in higher-order thalamus modulate the threshold for conscious
perception by influencing the bursting activity of TTL5 pyramidal neurons in the cortex[1,13,63–65].
These matrix cells, with their diffuse connectivity, are hypothesized to facilitate the entry of
sensory content into consciousness, potentially by lowering the activation threshold of L5
pyramidal neurons[13,26]. Our LIFU intervention at high DC may have induced excitation of the
matrix cells in the VA thalamus, possibly increasing dendritic integration in L5 pyramidal neurons
and promoting widespread cortical activation that enhances perceptual sensitivity (further
discussion on DC-dependence follows below).

The ventral anterior part of the thalamus (not to be confused with the ventral anterior thalamic
nucleus) encompasses the intralaminar nuclei, which project to various cortical and subcortical
regions[19,66]. Electrical stimulation of these nuclei via implanted electrodes has been shown to
awaken non-human primates from anesthesia[18,67] and aid in the recovery of consciousness in
patients with neuropathological conditions[68,69]. Similarly, microinjection of nicotine or potassium
channel-blocking antibodies into these nuclei has restored consciousness in rodents[70]. A previous
LIFU study targeting these nuclei demonstrated improvements in patients with disorders of
consciousness[36]. Our findings provide preliminary evidence that the intralaminar nuclei might also
influence the contents of consciousness, in addition to modulating the global states of
consciousness.

One important question is whether the increased sensitivity of conscious perception was due to
an excitation of the VA thalamus. Previous empirical studies suggest that LIFU selectively
activates excitatory vs. inhibitory neurons depending on the DC, thus exerting a general excitatory
effect at high DCs (e.g., > 50%) and an inhibitory effect at low DCs (e.g., < 20%)[39,40,48,71–74]. Based
on these observations and the subsequently established neuronal intramembrane cavitation
excitation model[48], we hypothesize that LIFU at 70% DC applied to the VA thalamus likely induced
an excitatory effect in our study. This interpretation is also supported by relevant human
neuroimaging studies. For example, Wu et al. demonstrated that higher pre-stimulus activity in
the VA thalamus predicts greater sensitivity in conscious perception[12]. This is in line with our
interpretation that 70% DC LIFU administered to the VA thalamus elevated its overall activity level,
leading to heightened perceptual sensitivity. Furthermore, given the strong bidirectional excitatory
connections[75] and functional connectivity (Fig. 5a, d), LIFU applied to the VA thalamus may
influence the activity of the mPFC. This is also supported by evidence that single-pulse electrical
stimulation of the VA activates the mPFC[76]. Higher pre-stimulus mPFC activity also predicts
greater perceptual sensitivity—a prediction intriguingly aligned with the previous findings[12].
Collectively, these observations suggest that 70% DC LIFU exerted an excitatory effect on the
VA thalamus, which in turn induced co-activation of the mPFC, which could account for the
observed increase in perceptual sensitivity.

Furthermore, we found that matrix cell richness, measured by the difference in CALB1 and PVALB
expression levels (CP_T)[22], significantly correlated with enhancements in perceptual sensitivity.
This relationship was specific to sensitivity and not observed for other outcomes, suggesting a
distinct role for matrix cell-rich regions in modulating perceptual discrimination. Unlike region-
based analyses that assume fixed anatomical targets, our approach incorporated individual beam
profiles at the voxel level, accounting for variability in sonication sites and local cell composition.
By linking behavioral effects to the actual sonication location and underlying cytoarchitecture, our
findings offer more direct evidence that matrix cell density in the thalamus plays a causal role in
conscious perception.

Unlike the increase in perceptual sensitivity with VA thalamus stimulation, categorization accuracy
was reduced with all four thalamic targets at 70% DC (Fig. 6b). This target-invariant effect did not
correlate with core-matrix cell composition, suggesting a distinct underlying mechanism. Given
the excitatory nature of high-DC LIFU and the shared thalamovisual connectivity among the
targeted regions (Fig. 5c), it is plausible that 70% DC sonication consistently co-activated the
visual cortex during the LIFU-ON blocks[44]. Such co-activation may have broadly disrupted
thalamocortical visual processing, affecting visual cortical networks that integrate inputs from

multiple thalamic regions. This interpretation aligns with previous work demonstrating a negative
correlation between the pre-stimulus activity of the visual network and categorization accuracy[12].

It is possible that the target-invariant effects were confounded by auditory artifacts[77]. Sonication
with 5% DC generated more noticeable noise from the transducer due to the higher pulse-
averaged intensity than with 70% DC (14.4 W cm^{-2} for 5% DC vs. 1.03 W cm^{-2} for 70% DC), given
a fixed temporal-average intensity. This led to a higher percentage of participants reporting
awareness of the transducer noise at 5% DC (93% vs. 52% of participants, see Source Data).
However, our results cannot be explained solely by auditory artifacts. First, when we included
transducer noise awareness as a covariate in our statistical model, it showed no significant effect
on behavioral outcomes (all $p_{FDR} > 0.05$; Supplementary Table 4). Second, if auditory artifacts
were the primary driver of the perceptual changes, one should expect target-invariant effects, with
a larger magnitude at 5% DC (at which transducer noise is more noticeable). Yet, the decrease
in categorization accuracy occurred only at 70% DC (Table 1). Nonetheless, to better isolate and
understand the effects of auditory artifacts, future research may include sham control by masking
transducer noise or physically blocking the ultrasound beam.

While previous research demonstrated long-lasting LIFU effects ranging from hours to weeks[78,79],
we did not observe persistent changes as evidenced by the comparison between two baselines
(Supplementary Fig. 3). This discrepancy may be attributed to two factors. First, the visual system,
being highly adaptable and dynamic, might respond differently to LIFU compared to states like
addiction, anxiety, or depression in which longer-term effects have been observed[74,78,80]. Second,
the intensity of our LIFU modulation might have been insufficient to trigger long-term plasticity. To
better understand the implications of LIFU for scientific study and therapeutic application, future
investigations should assess the factors that influence the duration of sonication effects.

Methodologically, LIFU offers unique advantages for causal and mechanistic investigations of the
neural substrates of consciousness in humans. Non-invasive techniques such as transcranial
magnetic stimulation have been employed to transiently disrupt the contents of consciousness,
for example, visual awareness[81–83]. However, these approaches are largely limited to superficial
cortical regions and lack the spatial resolution necessary to target deep subcortical circuits. Other
techniques such as deep brain stimulation and intracranial electrical stimulation applied to non-
human primates[11,67] or surgical patients[84–86] have provided important insights into the neural
mechanisms of conscious perception, but their invasiveness restricts clinical accessibility and
broader applicability. By enabling non-invasive, focal neuromodulation of both cortical and deep
subcortical structures with high spatiotemporal precision, LIFU provides a powerful new tool for
dissecting the causal architecture of human consciousness.

Our work represents several refinements in the field of human LIFU research. First, in contrast to
prior human studies that employed only one DC[87–89], we incorporated two duty cycles (70% vs.
5%) into our experimental design and found DC-dependent effects. Second, while regional
targeting subcortical regions has been demonstrated in rodents[40], previous human LIFU
investigations mainly targeted thalamus as a single target[35,74,87]. Our work extends these findings
by systematically targeting various thalamic subregions with sub-centimeter precision. The
average lateral deviation of the beam from the target was approximately 2 mm (Fig. 2b), smaller
than beam width of about 5 mm. This was comparable to or better than those in previous works
(3–9 mm)[87,90]. Therefore, our targeting approach suggests promising avenues for translational
scientific investigations and clinical applications. Third, of relevance to future clinical applications,

we mitigated the reliance on anatomical imaging (e.g., computed tomography or T1 MRI) by using
a template rescaling method. We verified that this method meets the desired targeting accuracy
in two ways: (1) using a separate MRI dataset, we confirmed small deviations of approximately 2
mm relative to the actual T1 image (Supplementary Fig. 4); (2) we observed that effect size
increased as the beam-target lateral deviation decreased, which is a good indicator that our
method effectively estimates the target location within an acceptable error range (Supplementary
Fig. 11a). Our approach could expedite the experimental procedures and enable LIFU application
in populations where anatomical scans are challenging, such as infants, individuals with implanted
medical devices, or those with severe claustrophobia.

Our study has limitations. First, although the template rescaling approach described above
allowed us to effectively approximate each participant's thalamic coordinates, we did not create
subject-specific acoustic models or apply skull-dependent phase corrections. The beam profiles
therefore cannot capture the focal shift, peak-pressure drop, or off-axis energy leakage introduced
by each participant's skull morphology. This lack of individualized skull modeling represents a
limitation that may have introduced unmodeled variability in beam propagation and targeting
precision, potentially affecting both the magnitude and localization of the observed
neuromodulatory effects. Likewise, while the transducer was positioned with an optical
neuronavigation system, its final offset and tilt were not iteratively refined based on native MR or
computed tomography images. This limitation further compromises targeting precision and should
be addressed in future investigations through individualized anatomical imaging or automated
transducer positioning protocols. Second, we did not perform functional imaging in the study
participants; while our findings are consistent with previously suggested DC-dependent LIFU
effects, we acknowledge that further investigation using concurrent functional imaging is
necessary to conclusively prove the existence and directionality of neural activity and connectivity
changes. Third, our simulations suggest significant energy loss (about 87%), mainly due to
skull[53,54,57,91], with potential additional attenuation by residual air pockets in hairs. Despite such
loss, previous human studies using comparable or lower intensities have still demonstrated
significant neuromodulatory effects[36,92–94]. In a recent rodent study, a temporal-average intensity
as low as 0.0075 W cm^{-2} (roughly 100-fold lower than ours; about 10-fold lower after considering
skull-induced attenuation) was sufficient to induce a detectable calcium signal[40]. Nonetheless,
advanced approaches such as direct attenuation compensation can help mitigate these issues[91].
Currently, FDA intensity guidelines (e.g., derated spatial-peak temporal-average intensity $I_{spta.3}$ =
0.72 W cm^{-2}) are based on standards for diagnostic and imaging ultrasound devices, and no
specific guidelines exist for LIFU applications in the human brain. As studies in animal models
have shown that higher intensities can yield stronger effects without causing tissue damage[40,95],
future work should advocate for updated FDA guidelines tailored to human brain LIFU.

In conclusion, our findings illuminate the causal roles of thalamic regions in conscious perception,
particularly highlighting the unique contribution of the matrix cell-rich, transmodal-dominant VA
thalamus in modulating visual discrimination sensitivity. These results provide insight into the
relationship between cytoarchitectural and functional organization of the thalamus, and human
conscious perception.

METHODS

Participants. The study protocol was approved by the Institutional Board Review of the University of Michigan Medical School. A total of 60 participants (age: 25.9 ± 6.3 years, mean ± SD; 38 females, 22 males) participated in the current experiment. All subjects provided written informed consent prior to the experiment and were compensated. All subjects were right-handed, did not have any hearing loss, were not colorblind, and had normal or corrected-to-normal vision. Although hair shaving was not required for this study, seven participants chose to shave their right temple, and they were given additional compensation in exchange. Participants were assigned to one of two equal-sized groups (duty cycle = 70% or 5%) by a random number generator, each group comprising 30 participants. Since the effect of LIFU neuromodulation is dependent on various factors, such as parameters and target regions, determining a prior effect size is challenging. However, we planned for a subject number of n = 30 per group, exceeding the mean sample size (n = 19) reported in the literature[96]. One subject who performed a different visual task was not included in data analysis. Three subjects with technical issues during targeting were excluded from data analysis. One subject did not complete the task due to technical issues in perceptual threshold determination. One subject was excluded because the hit rate during Baseline-1 was too low (< 15%). Thus, a total sample size was n = 54 (70% duty cycle: n = 27; 5% duty cycle: n = 27; see also Supplementary Fig. 1 for a comprehensive flow chart). Sex or gender analysis was not conducted because there were no sex or gender specific hypotheses regarding the influence of ultrasound neuromodulation on visual perception. Participants were blinded to the intervention conditions by providing identical procedural setups and device operations across both groups, ensuring that all participants did not know the specific duty cycle assignment.

Low-intensity focused ultrasound devices and parameters. LIFU was administered using the BrainSonix BXPulsar 1002 System (BrainSonix Corporations, Sherman Oaks, CA, USA). This system includes a single transducer with a 61.5 mm diameter. The free-field focal depth is 80.7 mm, validated by Blatek Industries, Inc. (Boalsburg, PA, USA), following ASTM E1065/E1065M Standard guidelines (see Supplementary Table 2 for test data report). This transducer is mounted in a plastic housing and sealed with a thin polyethylene membrane. Free-field beam volume including length and width was verified by BrainSonix Corporations with a representative 80-mm transducer using a bilaminar membrane hydrophone with a 0.4 mm diameter active receive aperture (Supplementary Fig. 5). Brainsight Neuronavigation System (Rogue Research, Montreal, Quebec, Canada) was used for beam targeting.

Sonication parameters were as follows: fundamental frequency: 650 kHz; pulse repetition frequency: 10 Hz; pulse duration: 100 ms; duty cycle: 70% and 5% (yielding pulse width of 70 and 5 ms, respectively); sonication duration: 30 s; inter-sonication interval: 30 s; LIFU-ON epochs per block: 12 epochs; non-derated spatial-peak temporal average intensity ($I_{spta.0}$): 1.02 W cm^{-2}; derated spatial-peak temporal average intensity ($I_{spta.3}$): 0.72 W cm^{-2}; non-derated spatial-peak pulse-average intensity ($I_{sppa.0}$): 1.03 W cm^{-2} (70% DC) and 14.4 W cm^{-2} (DC 5%); non-derated incident pressure ($P_{r.0}$): 197 kPa (70% DC) and 710 kPa (5% DC); derated incident pressure ($P_{r.3}$): 165 kPa (70% DC) and 619 kPa (5% DC). Since we used a fixed $I_{spta.3}$ for two duty cycles, we note that pulse-average intensity (I_{sppa}) was higher at DC 5% condition. Given that the Mechanical Index (MI) is the derated pressure divided by the square root of the fundamental frequency (0.65 MHz), MI values were estimated as: 0.20 (70% DC) and 0.77 (5% DC), below the ITRUSST risk recommendations[58]. The derated values were calculated using the US FDA's derating method,

which assumes a uniform tissue attenuation rate of 0.3 dB cm^{-1} MHz^{-1}. A comprehensive
parameter report summary based on the ITRUSST reporting guidelines[97] can be found in
Supplementary Table 1.

Transducer setup. Aquasonic Ultrasound Gel (Bio-Medical Instruments, Inc., Clinton Township,
MI, USA) was applied to the right temple before positioning the transducer, ensuring that all hairs
in the area were thoroughly coated with gel to enhance ultrasound transmission. The transducer
was then positioned on the right temple and secured with two adjustable fabric bands: one band
ran from the chin over the top of the head, and the other band passed from the forehead to the
back of the head. Additional self-adhering bandages were used on top of the fabric bands to fine-
tune the angle of the transducer for precise alignment. For post-hoc analysis, at the start and end
of each LIFU-ON block, lateral deviation and coordinates of the ideal beam center (i.e., 80 mm
away from the center of the transducer surface) and the head-transducer contact point were
recorded from the neuronavigation system. Beam vector was then defined as a vector originating
from the recorded head-transducer contact point, orienting toward the ideal beam center.

Target locations. Following a gradient-based thalamic parcellation scheme[52], the thalamus was
coarsely divided into four regions: ventral anterior (VA), ventral posterior (VP), dorsal anterior
(DA), dorsal posterior (DP). The center-of-mass coordinates for the four regions in the MNI space
are as follows; Left VA: (-8 mm, -11 mm, 6 mm), Left VP: (-13 mm, -23 mm, 2 mm), Left DA: (-12
mm, -23 mm, 12 mm), Left DP: (-16 mm, -31 mm, 2 mm).

Head size measurement. We used a digital outside-diameter caliper to measure head size in
three dimensions: width, depth, and side-to-top distance (Fig. 1b). Head width was defined as the
distance between the left and right supratragic notches, which are indentations in the ear cartilage
right above the tragus. Head depth was measured from the most posterior (back) point to the
most anterior (front) point of the head. The side-to-top distance was defined as the measurement
from the one supratragic notch to the highest top point of the head.

Template rescaling method. We rescaled the ICBM 2009c Nonlinear Asymmetric template of
the MNI152 linear template to match each participant's head size based on the three measured
dimensions. The head height, defined as the height of a triangle formed by the top of the head,
the supratragic notch, and the center of the head at the plane of the supratragic notch, was
calculated using the formula:

$$Height = \sqrt{(Side-to-Top\ Distance)^2 - (Width)^2} \qquad (1)$$

The default head size of the MNI template were as follows: width: 16 cm, depth: 21 cm, and height:
15 cm. We then calculated the size ratio between the MNI template and the participant's head
dimensions. The NIFTI image of the MNI template was resliced using B-spline interpolation to
adjust to the participant's head size.

We performed a post-hoc analysis with a separate MRI dataset (*n* = 13) to assess the accuracy
of our template rescaling method. For each participant, we obtained an anatomical T1 image and
measured head size (width, depth, and side-to-top distance). We then created a pseudo-
anatomical rescaled template as described above. The actual T1 and the rescaled template
images were aligned according to the anterior and posterior commissures, segmented,
normalized to MNI space, and used to generate inverse deformation fields. We applied the inverse
deformation fields to map four target MNI coordinates into each image's native space and
computed the Euclidean distances between the resulting coordinates for each thalamic region.

This distance provided a quantitative measure of how closely the rescaled template matched each
participant's true anatomy at the thalamic targets.

Simulation of ultrasound beam. Simulation of ultrasound intensity and thermal profiles was
performed using a two-step segmentation and modeling approach. We initially segmented the
pseudo-anatomical image into scalp, skull, and brain employing the Complete Head Anatomy
Reconstruction Method (CHARM)[98]. For a more accurate segmentation, we used a fat-
suppression method (https://github.com/ProteusMRIgHIFU/EnhanceTUS_Zadeh_JNE)[99]. For
cases where participant-specific ultrasound trajectories extended beyond the headspace due to
anatomical variability, the label assignment of scalp was manually extended outward by 1 cm,
and the mesh was recalculated with CHARM, ensuring that all recorded head-transducer contact
points lie within the adjusted headspace.

Subsequently, acoustic property assignments and ultrasound beam simulations were performed
using an open-source software named BabelBrain[100]. Simplified masks based on the CHARM-
generated MRI segmentations were utilized, with trabecular bone proportion set at 80%. These
masks assigned homogenized acoustic properties, including density, longitudinal and shear
speeds of sound, and attenuation, derived from established literature values[100]. Simulations were
conducted up to 100 mm below the transducer with a spatial resolution of six points-per-
wavelength.

The tissue attenuation ratio was determined by how much acoustic energy is lost when ultrasound
propagates through tissue compared to a water reference (exported as "RatioLosses" variable in
BabelBrain software). In short, the software selects the plane at which the maximum tissue
pressure occurs. It then computes the acoustic energy in this plane as the sum over all voxels of
the squared pressure amplitude. Similarly, the code processes the free-field pressure amplitude
data at the same plane. The tissue attenuation ratio is then defined as the ratio of the tissue
acoustic energy to the water acoustic energy at that specific plane.

Thermal effects resulting from ultrasound exposure were estimated using the Bio Heat Transfer
Equation (BHTE). Baseline temperature for thermal modeling was set at 37 ºC. During the
simulation, BabelBrain software calculates the temperature evolution in the tissue (i.e., skin, skull,
brain, and target point) both during ON and OFF epochs, which are then concatenated, yielding
an array of $4 \times 72,000$ size. Maximal temperature was determined by the maximal value within
this array.

The simulated relative intensity values exported from BabelBrain software were normalized to
MNI space using the forward deformation fields that MNI-normalized the corresponding pseudo-
anatomical image and then 3D-interpolated using 'interp3' function in MATLAB with the
refinement factor of 4. The individual beam trajectory images in Supplementary Fig. 8-10 were
also MNI-normalized and interpolated. The spatial coverage of each beam was determined as
the percentage of "beam-present" voxels (defined as voxels exhibiting more than 50% of the
relative intensity[101]) that fell within specific thalamic (or other non-thalamic) regions. For example,
the sum of beam coverage values across 10 regions (eight thalamic and two non-thalamic regions)
equals 100%.

Overall experimental procedure. The experiment was conducted in a single session lasting
approximately two hours, where the main task took approximately 1.5 hours (Fig. 1a). First, we
measured the participant's head size (details above), followed by task instructions. The participant
then completed a brief practice run consisting of 30 trials (a shorter version of staircase procedure;

see below) while the experimenter generated the pseudo-anatomical image. Next, the ultrasound
transducer was positioned, and participants underwent the image contrast staircase procedure
involving 60 trials. After confirming the convergence of contrast value (see below), participants
then engaged in the main task, which consisted of six blocks, each containing 100 trials and
lasting approximately 11.5 minutes (Fig. 1a). Each block featured four trials per each real image
(80 real trials in total) and five trials per each scrambled image (20 scrambled trials in total; see
below for details in visual stimuli). Participants were unaware of the fact that scrambled images
would appear.

During the inner four blocks, LIFU was administered toward one of four left thalamic areas in a
pre-assigned, pseudo-randomized order (determined by a random number generator), counter-
balanced across the participants. Within one participant, only one duty cycle (either 70% or 5%)
was applied throughout the four LIFU-ON blocks. Participants were allowed to rest between
blocks (3.8 ± 2.4 min; mean ± SD; see Source Data), during which the transducer was readjusted.
We also measured the eye-to-screen distance and readjusted the position of the laptop if
necessary (see below). After the transducer was successfully repositioned and the participants
were willing to proceed, the visual task was self-initiated by participants and LIFU administration
was manually triggered by the experimenter after the task onset.

Visual stimuli. Visual images encompassed four categories: faces, houses, human-made
objects, and animals (Fig. 1g), which were used in a previous study[8]. The images were sourced
from publicly available labeled photographs or the Psychological Image Collection at Stirling
(PICS; https://pics.stir.ac.uk). Each category included five distinct images, resulting in a total of
20 unique real images. All images were converted to grayscale and resized to 300 × 300 pixels.
The pixel intensities were normalized by subtracting the mean and dividing by the standard
deviation. Subsequently, the images were processed with a 2D Gaussian smoothing filter, using
a standard deviation of 1.5 pixels and a 7 × 7 pixel kernel. Scrambled images that maintain low-
level features were generated for each category by randomizing the phase of the 2D Fourier
transformation for one image from each category.

The contrast of an image was defined as:

$$contrast = \frac{I_{max} - I_{min}}{2b} \tag{2}$$

where I_{max} and I_{min} are the intensity values of lightest and darkest pixels in the image, respectively
(in the range of 0 to 255). To ensure a smooth transition to the background (set as gray, intensity
= 127), the edges of all images were gradually blended by applying a Gaussian window with a
standard deviation of 0.2 to the image intensity.

The stimuli were displayed on a 14-inch laptop (HP ProBook 440 G4) with a 60-Hz refresh rate
LCD screen, set to the maximum screen brightness. The viewing distance between the screen
and the participants' eyes was maintained at 65.8 ± 5.7 cm (mean ± SD). During the experiment,
participants were asked to maintain constant eye-monitor distance. We did not adopt masking
because masking is known to influence the visual information processing[9].

Image contrast staircase procedure. Before the main task, participants completed the QUEST
adaptive staircase procedure[8] to determine the contrast value that would yield a 50% recognition
rate (i.e., the rate of "YES" response to Question-2). The QUEST procedure was similar to the
main task but with shorter timing parameters and without scrambled images. The inter-trial interval
was 0.75 seconds, and the delay period between the stimulus and the first question was 2

seconds. The threshold contrast was identified through a QUEST process consisting of 60 trials.
Each trial was randomly assigned to one of three sets, and the staircase procedure was
conducted separately for each set. Task performance was deemed acceptable if the three sets
successfully converged on a specific contrast value. The staircase procedure was repeated until
the convergence of the three curves was confirmed. Rather than adjusting the contrast of each
image individually, the staircasing aimed for 50% recognition across all images.

Trial structure. Each trial began with a fixation cross displayed on a gray background for a
randomly determined duration between 3 and 6 seconds, following an exponential distribution to
prevent participants from predicting stimulus onset (Fig. 1e). The stimulus image was then shown
behind the fixation cross for 8 frames (67 ms), with the image intensity linearly increasing across
frames. After the stimulus disappeared, the fixation cross remained on the screen for an additional
2 seconds. Each trial concluded with two sequential questions about the stimulus, each displayed
for up to 3 seconds.

Question-1 asked participants to categorize the image as a face, house, object, or animal.
Participants were instructed to guess the category even if they did not consciously recognize the
object (four alternative forced-choice). Question-2 assessed their subjective experience, asking
whether they had a "meaningful visual experience" of the object stimulus ("YES" or "NO"). Before
the practice run, participants were told that a "meaningful" stimulus was defined as something
that makes sense in the real world, as opposed to random noise or meaningless shapes. They
were instructed to respond "YES" even if they recognized only part of an object. Participants
provided their answers by pressing buttons on a 4-key external USB keyboard placed on their lap.
If participants did not respond to Question-1, it was recorded as incorrect, and for Question-2, a
lack of response was recorded as "NO."

Signal detection theory analysis. We employed signal detection theory (SDT) to examine the
perceptual outcomes, focusing on sensitivity and criterion[9]. Sensitivity measures the ability to
distinguish between real images and scrambled images. We calculated parametric sensitivity
metric d' through standard SDT analysis defined as follows:

$$d' = \Phi^{-1}(H) - \Phi^{-1}(FA) \tag{3}$$

717 where Φ^{-1} is the inverse normal cumulative distribution function, H is the hit rate, and FA is the
718 false alarm rate, respectively. High d' indicates better discrimination between real and scrambled
719 images.

720 Decision bias refers to the inclination to certain response, regardless of whether the stimulus is
721 real or scrambled. We assessed decision bias using parametric metric criterion c defined below.

$$c = -\frac{\Phi^{-1}(H) + \Phi^{-1}(FA)}{2} \tag{4}$$

High c values indicate the tendency to say "NO" to Question-2, indicating a more conservative
object recognition.

Statistics. To ensure stabilization of task performance, the first ten trials of each block were
omitted from the analysis. Due to technical issues, a few participants failed to complete specific
LIFU-ON blocks, which were consequently excluded from the analysis. The numbers of affected
participants for each condition are as follows: 70% DC (VA: n = 1; VP: n = 1) and 5% DC (VA: n

= 2; VP: n = 2; DA: n = 1). For one subject, the lateral deviation during VP sonication was not
recorded. The coordinates for head-transducer contact point and ideal beam center were not
recorded for a few participants, resulting in their omission from simulation analysis. The numbers
of affected participants were as follows; VA: n = 8; VP: n = 9; DA: n = 8; DP: n = 8. Because d'
and c are ill-defined at perfect performance (i.e., $\Phi^{-1}(H) = \infty$ at H = 1)[102,103], we omitted such
blocks with from the statistical analyses of d' and c.

To address potential confounds, we included six nuisance variables in the analysis. Subject-
specific factors correspond to age, sex, and the awareness of transducer noise during LIFU-ON
blocks (coded as 0 for unaware, 0.5 for unsure, and 1 for aware). Block-specific factors such as
lateral deviation, sonication order, and tissue attenuation ratio (in dB) were also incorporated.
Missing data for noise awareness was imputed using modal category (noise awareness: 1).
Missing or baseline values of lateral deviation and attenuation ratio were imputed using group
means (lateral deviation: 2.251 mm; attenuation ratio: -8.7812 dB). Given no significant
differences between Baseline-1 and Baseline-2 across all measures (Supplementary Fig. 3), we
averaged these into a single baseline measure. No imputation was performed for the main
perceptual outcome data.

We implemented a sequential analytical approach beginning with omnibus testing using linear
mixed-effects (LME) models. This approach was chosen because LME models can directly
accommodate covariates, handle non-normal data distributions, and properly account for missing
outcome data through restricted maximum likelihood estimation. Our statistical implementation
involved fitting a single linear mixed-effects model specified as:

Outcome ~ Target + Covariates + (1|Subject)

where *Target* includes baseline and four LIFU regions (five categories total) and *Covariates*
encompass the six nuisance variables described above. For each perceptual outcome, we first
conducted an omnibus F-test to determine whether there were any differences among the five
experimental conditions using MATLAB's 'fitlme' and 'anova' functions with Satterthwaite
approximation.

Only outcomes that demonstrated significant omnibus effects after FDR correction proceeded to
post-hoc analysis. For significant outcomes, we conducted pairwise comparisons between each
LIFU condition and baseline using linear contrasts, with additional FDR correction applied within
this family of contrasts at α = 0.05. This approach ensures that each dataset undergoes a single,
unified analytical pathway while controlling family-wise error rate at the outcome level. For the
entire FDR correction steps, we used Benjamini-Hochberg procedure using MATLAB's 'mafdr'
function. Full statistics of main and supplementary analyses are included in Source Data.

Thalamic Cytoarchitecture Analysis. To examine the relationship between thalamic
cytoarchitecture and perceptual outcomes, we analyzed the core-matrix cell composition of
stimulated regions. Thalamic cell types were inferred from the mRNA expression levels of CALB1
and PVALB, as provided by the Allen Human Brain Atlas[59]. The probes
CUST_11451_PI416261804 and A_23_P17844 were used to estimate PVALB expression. For
estimating CALB1 expression, CUST_140_PI416408490, CUST_16773_PI416261804, and
A_23_P43197 were used. The CP_T metric was computed as the weighted difference between
CALB1 and PVALB expression levels[22]. Higher CP_T values reflected greater CALB1 expression
(i.e., matrix cell-rich regions), while lower values reflected greater PVALB expression (i.e., core
cell-rich regions).

Using the thalamic CP_T map provided by James M. Shine's Lab ('calb_minus_pvalb.nii' file in
https://github.com/macshine/corematrix), we calculated the mean CP_T value within the beam-
present voxels for each LIFU-ON block, defined as voxels exceeding 50% relative intensity based
on acoustic simulations. These CP_T values were then correlated with baseline-subtracted
changes in the three perceptual outcomes that survived omnibus testing.

Human Connectome Project Dataset. The dataset was obtained from the S1200 Release of the
WU-Minn Human Connectome Project (HCP) database, which has been extensively described in
prior studies[60]. The participants were healthy young adults aged 22 to 37 years. Participants who
had completed two sessions of resting-state fMRI scans (Rest1 and Rest2) were included in our
analysis, resulting in n = 1009. The data were collected using a customized Siemens 3T MR
scanner (Skyra system) with multiband EPI. Each scanning session comprised two sequences
with opposite phase encoding directions (left-to-right and right-to-left), each lasting 14 minutes
and 33 seconds. The sequences were acquired with a repetition time (TR) of 720 ms, an echo
time (TE) of 33.1 ms, and a voxel size of 2 mm isotropic. To maximize data quality and minimize
bias from phase encoding direction, the sequences from each session were combined, resulting
in a total of 29 minutes and 6 seconds. The denoised volumetric data, preprocessed through ICA-
FIX, were accessed from the online HCP database. Further details on the resting-state fMRI data
collection and preprocessing are available in previous publications[60]. We employed standard
preprocessing procedure, including resampling to a 3 × 3 × 3 mm resolution, band-pass filtering
within the 0.01–0.1 Hz frequency range, spatial smoothing with a 6-mm Full Width at Half
Maximum isotropic Gaussian kernel, and temporal normalization to attain zero mean and unit
variance.

Network-level analysis. We parcellated the cortex into 400 regions of interest (ROIs) based on
a well-established parcellation scheme, assigning each ROI to one of seven canonical cortical
networks: visual, somatomotor, dorsal attention, ventral attention, limbic, frontoparietal, and
default-mode networks[61,104]. Additionally, we used the Subcortical Atlas[52] to extract BOLD signal
time courses from 14 distinct thalamic ROIs, including bilateral ventral anterior inferior (VAi),
ventral anterior superior (VAs), ventral posterior lateral (VPl), ventral posterior medial (VPm),
dorsal anterior medial (DAm), dorsal anterior lateral (DAl), and dorsal posterior (DP) regions. For
analysis purposes, we averaged bilateral and related subdivisions into four primary thalamic
regions with equal weights: VA (left and right VAi and VAs), VP (left and right VPl and VPm), DA
(left and right DAm and DAl), and DP (left and right DP). For each resting-state session (Rest1
and Rest2), we computed the functional connectivity between each of these four thalamic regions
and the 400 cortical ROIs using Fisher-z transformed Pearson correlation coefficients.
Subsequently, for each session and thalamic region, we converted the correlation values into
percentile ranks across all 400 cortical ROIs. We then averaged these percentile ranks within
each of the seven canonical networks and across both resting-state sessions to derive network-
level thalamocortical connectivity measures. Finally, we calculated the average rank scores for
unimodal (visual and somatomotor) and transmodal (frontoparietal and default-mode) networks,
visualizing these results in a 2D plot to illustrate relative connectivity patterns.

Voxel-level analysis. As described above, mean time courses were extracted from bilateral VA,
VP, DA, and DP thalamus, serving as seeds for voxel-level functional connectivity analysis. Seed-
based maps (Fisher-z transformed Pearson correlation) were generated for each participant and
region. Group-level z-score maps were generated by standardizing individual connectivity values
(subtracting the mean and dividing by the standard deviation across participants). Within the

cortex, we retained the top 10% of cortical voxels with the highest connectivity. This analysis was
performed independently for Rest1 and Rest2 sessions, and their overlap was used to ensure
consistency. Region-specific and region-invariant clusters of high-connectivity cortical voxels
were then identified.

Software. The staircase and main task were based on a custom-modified version of the publicly-
available code for the near-threshold behavioral paradigm
(https://github.com/BiyuHeLab/NatCommun_Levinson2021/) utilizing the Psychophysics Toolbox
v3.0.19.4 (http://psychtoolbox.org/) executed on MATLAB R2023b[8]. Neuronagivation processes
including transducer targeting and coordinate recording were conducted on proprietary Brainsight
software (https://www.rogue-research.com/downloads/). Template rescaling and its post-hoc
validation were performed with custom-built code and SPM12 (https://www.fil.ion.ucl.ac.uk/spm)
both executed on MATLAB R2024a. Head segmentation with CHARM was conducted from the
SimNIBS v4.1.0 (https://simnibs.github.io/simnibs/build/html/index.html). Ultrasound beam
simulation was performed on BabelBrain v0.4.3 (https://proteusmrighifu.github.io/BabelBrain/). All
subsequent analyses involving volumetric beam intensity data (e.g., MNI normalization, average
beam intensity calculation, and CP_T calculation) were performed with custom scripts utilizing
SPM12 on MATLAB R2024b. The 3D renderings of brain and beam overlaps were performed
using MRIcroGL v1.2.20220720 (https://www.nitrc.org/projects/mricrogl). For fMRI data
preprocessing and voxel-level connectivity analysis, we employed the AFNI software suite
(linux_ubuntu_16_64; http://afni.nimh.nih.gov/). All other analyses (e.g., outcome calculation,
statistical tests, and network-level connectivity analysis) and visualizations were conducted on
MATLAB R2024b.

DATA AVAILABILITY

Source data are provided with this paper. The HCP dataset is available from online repository
(https://www.humanconnectome.org/). The thalamic CP_T map can be downloaded from Github
repository (https://github.com/macshine/corematrix).

CODE AVAILABILITY

Original code for near-threshold paradigm is available at Github
(https://github.com/BiyuHeLab/NatCommun_Levinson2021/)[8]. Custom-built code for template
rescaling is available at Github (https://github.com/janghw4/template_rescaling).

REFERENCES

1. Aru, J., Suzuki, M. & Larkum, M. E. Cellular Mechanisms of Conscious Processing. *Trends in*

Cognitive Sciences **24**, 814–825 (2020).

2. Dehaene, S., Changeux, J.-P., Naccache, L., Sackur, J. & Sergent, C. Conscious, preconscious,

and subliminal processing: a testable taxonomy. *Trends Cogn Sci* **10**, 204–211 (2006).

3. Dehaene, S. & Changeux, J.-P. Experimental and Theoretical Approaches to Conscious

Processing. *Neuron* **70**, 200–227 (2011).

4. Koch, C., Massimini, M., Boly, M. & Tononi, G. Neural correlates of consciousness: progress

and problems. *Nat Rev Neurosci* **17**, 307–321 (2016).

5. Mashour, G. A. & Hudetz, A. G. Neural Correlates of Unconsciousness in Large-Scale Brain

Networks. *Trends in Neurosciences* **41**, 150–160 (2018).

6. Storm, J. F. *et al.* An integrative, multiscale view on neural theories of consciousness. *Neuron*

S0896627324000886 (2024) doi:10.1016/j.neuron.2024.02.004.

7. Martín-Signes, M., Chica, A. B., Bartolomeo, P. & Thiebaut de Schotten, M. Streams of

conscious visual experience. *Commun Biol* **7**, 908 (2024).

8. Levinson, M., Podvalny, E., Baete, S. H. & He, B. J. Cortical and subcortical signatures of

conscious object recognition. *Nat Commun* **12**, 2930 (2021).

9. Podvalny, E., Flounders, M. W., King, L. E., Holroyd, T. & He, B. J. A dual role of prestimulus

spontaneous neural activity in visual object recognition. *Nat Commun* **10**, 3910 (2019).

10. Takahashi, N., Oertner, T. G., Hegemann, P. & Larkum, M. E. Active cortical dendrites modulate

perception. *Science* **354**, 1587–1590 (2016).

11. Van Vugt, B. *et al.* The threshold for conscious report: Signal loss and response bias in visual

and frontal cortex. *Science* **360**, 537–542 (2018).

12. Wu, Y., Podvalny, E., Levinson, M. & He, B. J. Network mechanisms of ongoing brain activity's

influence on conscious visual perception. *Nat Commun* **15**, 5720 (2024).

13. Whyte, C. J., Redinbaugh, M. J., Shine, J. M. & Saalmann, Y. B. Thalamic contributions to the

state and contents of consciousness. *Neuron* **112**, 1611–1625 (2024).

14. Shine, J. M., Lewis, L. D., Garrett, D. D. & Hwang, K. The impact of the human thalamus on

brain-wide information processing. *Nat Rev Neurosci* **24**, 416–430 (2023).

15. Halassa, M. M. *The Thalamus*. (Cambridge University Press, 2022).

16. Jones, E. G. Synchrony in the Interconnected Circuitry of the Thalamus and Cerebral Cortex.

Annals of the New York Academy of Sciences **1157**, 10–23 (2009).

17. Müller, E. J. *et al.* The non-specific matrix thalamus facilitates the cortical information processing

modes relevant for conscious awareness. *Cell Reports* **42**, 112844 (2023).

18. Redinbaugh, M. J. *et al.* Thalamus Modulates Consciousness via Layer-Specific Control of

Cortex. *Neuron* **106**, 66-75.e12 (2020).

19. Vertes, R. P., Linley, S. B. & Rojas, A. K. P. Structural and functional organization of the midline

and intralaminar nuclei of the thalamus. *Front. Behav. Neurosci.* **16**, 964644 (2022).

20. Huang, Z., Mashour, G. A. & Hudetz, A. G. Propofol disrupts the functional core-matrix

architecture of the thalamus in humans. *Nat Commun* **15**, 7496 (2024).

21. Margulies, D. S. *et al.* Situating the default-mode network along a principal gradient of

macroscale cortical organization. *Proceedings of the National Academy of Sciences* **113**,

12574–12579 (2016).

22. Müller, E. J. *et al.* Core and matrix thalamic sub-populations relate to spatio-temporal cortical

connectivity gradients. *NeuroImage* **222**, 117224 (2020).

23. Aru, J., Suzuki, M., Rutiku, R., Larkum, M. E. & Bachmann, T. Coupling the State and Contents

of Consciousness. *Front. Syst. Neurosci.* **13**, (2019).

24. Honjoh, S. *et al.* Regulation of cortical activity and arousal by the matrix cells of the ventromedial

thalamic nucleus. *Nat Commun* **9**, 2100 (2018).

25. Shepherd, G. M. G. & Yamawaki, N. Untangling the cortico-thalamo-cortical loop: cellular pieces

of a knotty circuit puzzle. *Nat Rev Neurosci* **22**, 389–406 (2021).

26. Takahashi, N. *et al.* Active dendritic currents gate descending cortical outputs in perception. *Nat*

Neurosci **23**, 1277–1285 (2020).

27. Liu, X., Qiu, F., Hou, L. & Wang, X. Review of Noninvasive or Minimally Invasive Deep Brain

Stimulation. *Front. Behav. Neurosci.* **15**, (2022).

28. Lozano, A. M. *et al.* Deep brain stimulation: current challenges and future directions. *Nat Rev*

Neurol **15**, 148–160 (2019).

29. Bystritsky, A. *et al.* A review of low-intensity focused ultrasound pulsation. *Brain Stimulation* **4**,

125–136 (2011).

30. Legon, W. *et al.* Transcranial focused ultrasound modulates the activity of primary

somatosensory cortex in humans. *Nat Neurosci* **17**, 322–329 (2014).

31. Rabut, C. *et al.* Ultrasound Technologies for Imaging and Modulating Neural Activity. *Neuron*

108, 93–110 (2020).

32. Tufail, Y., Yoshihiro, A., Pati, S., Li, M. M. & Tyler, W. J. Ultrasonic neuromodulation by brain

stimulation with transcranial ultrasound. *Nat Protoc* **6**, 1453–1470 (2011).

33. Tufail, Y. *et al.* Transcranial Pulsed Ultrasound Stimulates Intact Brain Circuits. *Neuron* **66**, 681–

694 (2010).

34. Nakajima, K. *et al.* A causal role of anterior prefrontal-putamen circuit for response inhibition

revealed by transcranial ultrasound stimulation in humans. *Cell Reports* **40**, (2022).

35. Legon, W., Ai, L., Bansal, P. & Mueller, J. K. Neuromodulation with single-element transcranial

focused ultrasound in human thalamus. *Hum Brain Mapp* **39**, 1995–2006 (2018).

36. Monti, M. M., Schnakers, C., Korb, A. S., Bystritsky, A. & Vespa, P. M. Non-Invasive Ultrasonic

Thalamic Stimulation in Disorders of Consciousness after Severe Brain Injury: A First-in-Man

Report. *Brain Stimulation* **9**, 940–941 (2016).

37. Darmani, G. *et al.* Individualized non-invasive deep brain stimulation of the basal ganglia using

transcranial ultrasound stimulation. *Nat Commun* **16**, 2693 (2025).

38. Darrow, D. P., O'Brien, P., Richner, T. J., Netoff, T. I. & Ebbini, E. S. Reversible neuroinhibition

by focused ultrasound is mediated by a thermal mechanism. *Brain Stimulation* **12**, 1439–1447

(2019).

39. Dell'Italia, J., Sanguinetti, J. L., Monti, M. M., Bystritsky, A. & Reggente, N. Current State of

Potential Mechanisms Supporting Low Intensity Focused Ultrasound for Neuromodulation.

Front. Hum. Neurosci. **16**, 872639 (2022).

40. Murphy, K. R. *et al.* Optimized ultrasound neuromodulation for non-invasive control of behavior

and physiology. *Neuron* (2024) doi:10.1016/j.neuron.2024.07.002.

41. Fomenko, A., Neudorfer, C., Dallapiazza, R. F., Kalia, S. K. & Lozano, A. M. Low-intensity

ultrasound neuromodulation: An overview of mechanisms and emerging human applications.

Brain Stimulation **11**, 1209–1217 (2018).

42. Tyler, W. J., Lani, S. W. & Hwang, G. M. Ultrasonic modulation of neural circuit activity. *Current*

Opinion in Neurobiology **50**, 222–231 (2018).

43. Mesik, L. *et al.* Transcranial Low-Intensity Focused Ultrasound Stimulation of the Visual

Thalamus Produces Long-Term Depression of Thalamocortical Synapses in the Adult Visual

Cortex. *J. Neurosci.* **44**, (2024).

44. Yu, K. & He, B. Transcranial Focused Ultrasound Modulates Visual Thalamus in a Nonhuman

Primate Model. *IEEE Trans Biomed Eng* **PP**, (2025).

45. Darmani, G. *et al.* Non-invasive transcranial ultrasound stimulation for neuromodulation. *Clinical*

Neurophysiology **135**, 51–73 (2022).

46. Kubanek, J. *et al.* Remote, brain region–specific control of choice behavior with ultrasonic

waves. *Sci. Adv.* **6**, eaaz4193 (2020).

47. Plaksin, M., Shoham, S. & Kimmel, E. Intramembrane Cavitation as a Predictive Bio-

Piezoelectric Mechanism for Ultrasonic Brain Stimulation. *Phys. Rev. X* **4**, 011004 (2014).

48. Plaksin, M., Kimmel, E. & Shoham, S. Cell-Type-Selective Effects of Intramembrane Cavitation

as a Unifying Theoretical Framework for Ultrasonic Neuromodulation. *eNeuro* **3**, ENEURO.0136-

15.2016 (2016).

49. Yoon, K. *et al.* Effects of sonication parameters on transcranial focused ultrasound brain

stimulation in an ovine model. *PLoS ONE* **14**, e0224311 (2019).

50. Kim, H. *et al.* Suppression of EEG visual-evoked potentials in rats through neuromodulatory

focused ultrasound. *NeuroReport* **26**, 211–215 (2015).

51. Cain, J. A. *et al.* Real time and delayed effects of subcortical low intensity focused ultrasound.

Sci Rep **11**, 6100 (2021).

52. Tian, Y., Margulies, D. S., Breakspear, M. & Zalesky, A. Topographic organization of the human

subcortex unveiled with functional connectivity gradients. *Nat Neurosci* **23**, 1421–1432 (2020).

53. Bernard, S. *et al.* Influence of Plastination on Ultrasound Transmission Through the Human

Skull. *Ultrasound in Medicine & Biology* **49**, 901–907 (2023).

54. Legon, W. & Strohman, A. Low-intensity focused ultrasound for human neuromodulation. *Nat*

Rev Methods Primers **4**, 1–27 (2024).

55. Lord, B. *et al.* Transcranial focused ultrasound to the posterior cingulate cortex modulates

default mode network and subjective experience: an fMRI pilot study. *Front. Hum. Neurosci.* **18**,

1392199 (2024).

56. Jing, B., Strassle Rojas, S. & Lindsey, B. D. Effect of skull porosity on ultrasound transmission

and wave mode conversion at large incidence angles. *Medical Physics* **50**, 3092–3102 (2023).

57. Mueller, J. K., Ai, L., Bansal, P. & Legon, W. Numerical evaluation of the skull for human

neuromodulation with transcranial focused ultrasound. *J. Neural Eng.* **14**, 066012 (2017).

58. Aubry, J.-F. *et al.* ITRUSST Consensus on Biophysical Safety for Transcranial Ultrasonic

Stimulation. Preprint at https://doi.org/10.48550/arXiv.2311.05359 (2024).

59. Hawrylycz, M. J. *et al.* An anatomically comprehensive atlas of the adult human brain

transcriptome. *Nature* **489**, 391–399 (2012).

60. Van Essen, D. C. *et al.* The WU-Minn Human Connectome Project: An overview. *NeuroImage*

80, 62–79 (2013).

61. Thomas Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic

functional connectivity. *Journal of Neurophysiology* **106**, 1125–1165 (2011).

62. Suzuki, M. & Larkum, M. E. General Anesthesia Decouples Cortical Pyramidal Neurons. *Cell*

180, 666-676.e13 (2020).

63. Mukherjee, A., Lam, N. H., Wimmer, R. D. & Halassa, M. M. Thalamic circuits for independent

control of prefrontal signal and noise. *Nature* **600**, 100–104 (2021).

64. Fang, Z. *et al.* Human high-order thalamic nuclei gate conscious perception through the

thalamofrontal loop. *Science* **388**, eadr3675 (2025).

65. Fang, Z. *et al.* Human high-order thalamic nuclei gate conscious perception through the

thalamofrontal loop. *Science* **388**, eadr3675 (2025).

66. Jones, E. G. The thalamic matrix and thalamocortical synchrony. *Trends in Neurosciences* **24**,

595–601 (2001).

67. Tasserie, J. *et al.* Deep brain stimulation of the thalamus restores signatures of consciousness

in a nonhuman primate model. *Sci. Adv.* **8**, eabl5547 (2022).

68. Schiff, N. D. *et al.* Behavioural improvements with thalamic stimulation after severe traumatic

brain injury. *Nature* **448**, 600–603 (2007).

69. Schiff, N. D. *et al.* Thalamic deep brain stimulation in traumatic brain injury: a phase 1,

randomized feasibility study. *Nat Med* **29**, 3162–3174 (2023).

70. Alkire, M. T., Asher, C. D., Franciscus, A. M. & Hahn, E. L. Thalamic Microinfusion of Antibody to

a Voltage-gated Potassium Channel Restores Consciousness during Anesthesia.

Anesthesiology **110**, 766–773 (2009).

71. Yu, K., Niu, X., Krook-Magnuson, E. & He, B. Intrinsic functional neuron-type selectivity of

transcranial focused ultrasound neuromodulation. *Nat Commun* **12**, 2519 (2021).

72. Fomenko, A. *et al.* Systematic examination of low-intensity ultrasound parameters on human

motor cortex excitability and behavior. *eLife* **9**, e54497 (2020).

73. Murphy, K. R. *et al.* A tool for monitoring cell type–specific focused ultrasound neuromodulation

and control of chronic epilepsy. *Proceedings of the National Academy of Sciences* **119**,

e2206828119 (2022).

74. Fan, J. M. *et al.* Thalamic transcranial ultrasound stimulation in treatment resistant depression.

Brain Stimulation **17**, 1001–1004 (2024).

75. Yang, C. *et al.* Medial prefrontal cortex and anteromedial thalamus interaction regulates goal-

directed behavior and dopaminergic neuron activity. *Nat Commun* **13**, 1386 (2022).

76. Wu, D. *et al.* Human anterior thalamic stimulation evoked cortical potentials align with intrinsic

functional connectivity. *Neuroimage* **277**, 120243 (2023).

77. Sato, T., Shapiro, M. G. & Tsao, D. Y. Ultrasonic Neuromodulation Causes Widespread Cortical

Activation via an Indirect Auditory Mechanism. *Neuron* **98**, 1031-1041.e5 (2018).

78. Mahoney, J. J. *et al.* Low-intensity focused ultrasound targeting the nucleus accumbens as a

potential treatment for substance use disorder: safety and feasibility clinical trial. *Front.*

Psychiatry **14**, (2023).

79. Yaakub, S. N. *et al.* Transcranial focused ultrasound-mediated neurochemical and functional

connectivity changes in deep cortical regions in humans. *Nat Commun* **14**, 5318 (2023).

80. Mahdavi, K. D. *et al.* A pilot study of low-intensity focused ultrasound for treatment-resistant

generalized anxiety disorder. *Journal of Psychiatric Research* **168**, 125–132 (2023).

81. Koivisto, M., Harjuniemi, I., Railo, H., Salminen-Vaparanta, N. & Revonsuo, A. Transcranial

magnetic stimulation of early visual cortex suppresses conscious representations in a

dichotomous manner without gradually decreasing their precision. *Neuroimage* **158**, 308–318

(2017).

82. Rounis, E., Maniscalco, B., Rothwell, J. C., Passingham, R. E. & Lau, H. Theta-burst

transcranial magnetic stimulation to the prefrontal cortex impairs metacognitive visual

awareness. *Cogn Neurosci* **1**, 165–175 (2010).

83. Jacobs, C., Goebel, R. & Sack, A. T. Visual awareness suppression by pre-stimulus brain

stimulation; a neural effect. *NeuroImage* **59**, 616–624 (2012).

84. Raccah, O., Block, N. & Fox, K. C. R. Does the Prefrontal Cortex Play an Essential Role in

Consciousness? Insights from Intracranial Electrical Stimulation of the Human Brain. *J.*

Neurosci. **41**, 2076–2087 (2021).

85. Blanke, O., Ortigue, S., Landis, T. & Seeck, M. Stimulating illusory own-body perceptions.

Nature **419**, 269–270 (2002).

86. Vesuna, S. *et al.* Deep posteromedial cortical rhythm in dissociation. *Nature* **586**, 87–94 (2020).

87. Badran, B. W. *et al.* Sonication of the anterior thalamus with MRI-Guided transcranial focused

ultrasound (tFUS) alters pain thresholds in healthy adults: A double-blind, sham-controlled

study. *Brain Stimulation* **13**, 1805–1812 (2020).

88. Legon, W., Bansal, P., Tyshynsky, R., Ai, L. & Mueller, J. K. Transcranial focused ultrasound

neuromodulation of the human primary motor cortex. *Sci Rep* **8**, 10007 (2018).

89. Xia, X. *et al.* Time course of the effects of low-intensity transcranial ultrasound on the excitability

of ipsilateral and contralateral human primary motor cortex. *NeuroImage* **243**, 118557 (2021).

90. Xu, L. *et al.* Characterization of the Targeting Accuracy of a Neuronavigation-Guided

Transcranial FUS System In Vitro, In Vivo, and In Silico. *IEEE Transactions on Biomedical*

Engineering **70**, 1528–1538 (2023).

91. Riis, T., Feldman, D., Mickey, B. & Kubanek, J. Controlled noninvasive modulation of deep brain

regions in humans. *Commun Eng* **3**, 1–12 (2024).

92. Sanguinetti, J. L. *et al.* Transcranial Focused Ultrasound to the Right Prefrontal Cortex Improves

Mood and Alters Functional Connectivity in Humans. *Front. Hum. Neurosci.* **14**, 52 (2020).

93. Lee, W., Weisholtz, D. S., Strangman, G. E. & Yoo, S.-S. Safety Review and Perspectives of

Transcranial Focused Ultrasound Brain Stimulation. *Brain Neurorehabil* **14**, e4 (2021).

94. Chou, T. *et al.* Transcranial focused ultrasound of the amygdala modulates fear network

activation and connectivity. *Brain Stimulation* **17**, 312–320 (2024).

95. Folloni, D. *et al.* Manipulation of Subcortical and Deep Cortical Activity in the Primate Brain

Using Transcranial Focused Ultrasound Stimulation. *Neuron* **101**, 1109-1116.e5 (2019).

96. Sarica, C. *et al.* Human Studies of Transcranial Ultrasound neuromodulation: A systematic

review of effectiveness and safety. *Brain Stimulation* **15**, 737–746 (2022).

97. Martin, E. *et al.* ITRUSST consensus on standardised reporting for transcranial ultrasound

stimulation. *Brain Stimulation* **17**, 607–615 (2024).

98. Puonti, O. *et al.* Accurate and robust whole-head segmentation from magnetic resonance

images for individualized head modeling. *NeuroImage* **219**, 117044 (2020).

99. Zadeh, A. K. *et al.* Enhancing transcranial ultrasound stimulation planning with MRI-derived skull

masks: a comparative analysis with CT-based processing. *J. Neural Eng.* **22**, 016020 (2025).

100.   Pichardo, S. BabelBrain: An Open-Source Application for Prospective Modeling of

Transcranial Focused Ultrasound for Neuromodulation Applications. *IEEE Transactions on*

Ultrasonics, Ferroelectrics, and Frequency Control **70**, 587–599 (2023).

101.   Murphy, K. R. *et al.* A practical guide to transcranial ultrasonic stimulation from the IFCN-

1068   endorsed ITRUSST consortium. *Clin Neurophysiol* **171**, 192–226 (2025).

1069   102.   Snodgrass, J. G. & Corwin, J. Pragmatics of measuring recognition memory: Applications to

1070   dementia and amnesia. *Journal of Experimental Psychology: General* **117**, 34–50 (1988).

1071   103.   Miller, J. The sampling distribution of d′. *Perception & Psychophysics* **58**, 65–72 (1996).

1072   104.   Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic

1073   Functional Connectivity MRI. *Cerebral Cortex* **28**, 3095–3114 (2018).

1074

1075

ACKNOWLEDGEMENTS

We express our sincere gratitude to our study coordinators, Amy McKinney and Aaron Ellis, for their invaluable contributions to participant recruitment, meticulous scheduling, and ensuring the smooth execution of this study. We sincerely appreciate Drs. Mark E. Schafer and Alexander Bystritsky for providing free-field beam characteristics and Dr. Kelli A. Sullivan for thoughtful comments on skull-induced energy loss. This work was supported by the National Institute of General Medical Sciences of the National Institutes of Health grants R01GM103894 (to A.G.H. and Z.H.). The content is solely the responsibility of the author and does not necessarily represent the official views of the National Institutes of Health.

AUTHOR CONTRIBUTIONS STATEMENT

H.J. and Z.H. designed the study and set up the experimental apparatus. H.J. devised the template rescaling method and developed the code. H.J., P.F., and Z.H. conducted the experiments. H.J. analyzed the data, prepared the figures, and drafted the manuscript. Z.H. co-analyzed the data. Z.H., G.A.M. and A.G.H. interpreted the data and edited the manuscript.

COMPETING INTERESTS STATEMENT

The authors declare no competing interests.